Cysteine signaling in plant pathogen response

Jannis Moormann[1], Björn Heinemann[1], Cecile Angermann[1], Anna Koprivova[1], Ute Armbruster[2], Stanislav Kopriva[1], Tatjana M. Hildebrandt[1]

[1]Institute for Plant Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), University of Cologne, Zülpicher Straße 47a, 50674 Cologne, Germany
[2]Molecular Photosynthesis, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich-Heine-University Düsseldorf, Universitätsstraße 1, D-40225, Düsseldorf, Germany

Abstract

The amino acid cysteine is the precursor for a wide range of sulfur containing functional molecules in plants including enzyme cofactors and defense compounds. Due to its redox active thiol group cysteine is highly reactive. Synthesis and degradation pathways are present in several subcellular compartments to adjust the intracellular cysteine concentration. However, stress conditions can lead to a transient increase in local cysteine levels. Here we investigate links between cysteine homeostasis and metabolic signaling in *Arabidopsis thaliana*. The systemic proteome response to cysteine feeding strongly suggests that Arabidopsis seedlings interpret accumulation of cysteine above a certain threshold as a signal for a biotic threat. Cysteine supplementation of Arabidopsis plants via the roots increases their resistance to the hemibiotrophic bacterium *Pseudomonas syringae* confirming the protective function of the cysteine induced defense pathways. Analysis of mutant plants reveals that the balance of cysteine synthesis between the cytosol and organelles is crucial during Arabidopsis immune response to *Pseudomonas syringae*. The induction profile of pathogen responsive proteins by cysteine provides insight into potential modes of action. Our results highlight the role of cysteine as a metabolic signal in the plant immune response and add evidence to the emerging concept of intracellular organelles as important players in plant stress signaling.

Keywords: amino acid metabolism, sulfur signaling, infochemicals, immunometabolism, mitochondria, proteomics

Abbreviations: ABA, abscisic acid; AOX, alternative oxidase; APK, adenosine 5'-phosphosulfate kinase; APS, adenosine-5'-phosphosulfate; BTH, benzothiadiazole; CAS-C1, cyanoalanine synthase; DAMP, damage-associated molecular pattern; DES1, L-cysteine desulfhydrase; dpi, days past infection; ETHE1, sulfur dioxygenase; GLR, Glutamate receptor-like calcium channels; GSH, reduced glutathione; GST, glutathione-S transferases; hpi, hours past infection; JA, jasmonic acid; NFS1, cysteine desulfurase; OAS, O-acetylserine; OASTL, O-acetylserine (thiol)lyase; PAMP, pathogen-associated molecular pattern; *Pst*, *Pseudomonas syringae* pv tomato DC3000; ROS, reactive oxygen species; SA, salicylic acid; SAR, systemic acquired resistance; SERAT, serine acetyltransferase; STR1, 3-mercaptopyruvate sulfur transferase; UDP, uridine diphosphate; UGT, uridine diphosphate-dependent glucosyltransferase.

Introduction

Amino acids have multiple functions in plants. In addition to their role during protein biosynthesis, they are an integral part of several biosynthetic pathways and involved in signaling processes including plant stress responses (Heinemann & Hildebrandt, 2021). Amino acids can serve as markers reflecting nutrient availability or as carbon source for alternative respiratory pathways under energy starvation (Pedrotti et al., 2018). Proline is known to function as an osmolyte during osmotic stress and as molecular chaperone preventing protein aggregation (Szabados & Savouré, 2010). In addition, amino acids serve as precursors for various molecules involved in plant immunity. Perturbations in amino acid metabolism have been reported to affect plant immune responses in various ways (Moormann et al., 2022). The phloem-mobile signaling molecule N-hydroxypipecolic acid, required for establishing systemic acquired resistance, is synthesized from lysine (Gupta & Spenser, 1969; Návarová et al., 2012). Aromatic amino acids are precursors for a broad range of specialized molecules that crucially shape the interaction between plants and microbes such as coumarins, phytoalexins and indolic glucosinolates (Glawischnig, 2007; Harun et al., 2020; Maeda & Dudareva, 2012; Pastorczyk et al., 2020). The thiazole ring of camalexin, the characteristic phytoalexin of *Arabidopsis thaliana*, originates from the cysteine residue of glutathione (Su et al., 2011).

Among amino acids, cysteine is unique as it occupies a central position in plant sulfur metabolism. It is the precursor for a wide range of sulfur containing molecules in the cell, including methionine, essential vitamins and cofactors such as thiamin, lipoic acid, biotin, Fe-S clusters and molybdenum cofactor (Droux, 2004; Giovanelli et al., 1985; van Hoewyk et al., 2008). Furthermore, the tripeptide γ-glutamyl-cysteinyl-glycine, also known as glutathione, relies on the incorporation of cysteine to function as the main determinant of cellular redox homeostasis (Foyer & Noctor, 2011). Its antioxidant property is based on the redox potential of the thiol group. Next to oxidative stress protection, glutathione also acts in detoxification of heavy metals and xenobiotics as well as in the plant defense response (Noctor et al., 2024). In proteins, cysteine residues contribute to structure, stability and function. When located in the active sites of enzymes they are essential for catalysis of many enzymatic reactions. Moreover, thiol groups can undergo oxidation to form covalent disulfide bridges, aiding protein folding and stability. Reversible oxidation/reduction of these disulfide bridges poses a mechanism for redox regulation of proteins (Buchanan & Balmer, 2005). Another layer of regulation is added by protein persulfidation whereby the thiol group of reactive cysteine residues (-SH) is converted to a persulfide (-SSH) resulting in functional changes (Moseler et al., 2024).

Cysteine is the product of the plant sulfur assimilatory pathway. Sulfate is taken up from the soil by specific transporters and activated by ATP sulfurylase. The resulting adenosine-5'-phosphosulfate (APS) is reduced in a two-step reaction via sulfite to sulfide by APS reductase and sulfite reductase. Sulfide is incorporated into O-acetylserine (OAS) by OAS-(thiol)lyase (OASTL) to produce cysteine (Takahashi, 2010; Takahashi et al., 2011). The amino acid precursor OAS is synthesized by serine acetyltransferase (SERAT) from serine and acetyl-CoA. OASTL and SERAT form a cysteine synthase complex which is required for regulating cysteine synthesis based on substrate availability (Droux, 2003; Wirtz & Hell, 2006). Both enzymes have isoforms localized in the cytosol (SERAT1;1, SERAT3;1, SERAT3;2, OASTL-A), plastids (SERAT2;1, OASTL-B) and mitochondria (SEART2;2, OASTL-C) enabling OAS and cysteine synthesis in different subcellular locations (Hell & Wirtz, 2011; Ruffet et al., 1995; Watanabe et al., 2008). Cysteine desulfurases, which transfer sulfur to Fe-S cluster scaffold

proteins, are present in plastids and mitochondria as well (Couturier et al., 2013). Cysteine levels are tightly regulated and generally kept low within the cell requiring adequate rates of cysteine degradation (Hildebrandt et al., 2015). In the cytosol, the L-cysteine desulfhydrase DES1 deaminates cysteine resulting in pyruvate, ammonia and sulfide (Alvarez et al., 2010). In mitochondria, a four-step process catalyzing the complete oxidation of cysteine to pyruvate and thiosulfate is facilitated by an unknown aminotransferase, the 3-mercaptopyruvate sulfur transferase STR1, and the sulfur dioxygenase ETHE1 (Höfler et al., 2016). Cysteine levels vary among cellular compartments and are highest in the cytosol which is known to be the main contributor of cysteine with concentrations around 300 µM (Heeg et al., 2008; Krüger et al., 2009). Accordingly, OASTL-A together with OASTL-B were found to account for 95 % of OASTL activity in *Arabidopsis* protein extracts, while OASTL-C contributed only 5% of the activity. However, all three major OASTL isoforms were shown to largely compensate for each other's absence in null mutant studies (Birke et al., 2013; Heeg et al., 2008).

Previous findings provide some insight into the relevance of compartment specific cysteine metabolic pathways. Cytochrome c oxidase, the last enzyme of the mitochondrial respiratory chain, is strongly inhibited by cyanide as well as by hydrogen sulfide (Cooper & Brown, 2008). Biosynthesis of camalexin and the gaseous phytohormone ethylene both lead to the production of cyanide in non-cyanogenic plants such as *Arabidopsis* (Böttcher et al., 2009; Peiser et al., 1984). Cyanide detoxification in the mitochondria is achieved by conversion of cysteine and cyanide to hydrogen sulfide and β-cyanoalanine catalyzed by cyanoalanine synthase (CAS-C1) (Hatzfeld et al., 2000; Watanabe et al., 2008). Hydrogen sulfide in turn is detoxified via incorporation into cysteine by mitochondrial OASTL-C, creating a cyclic detoxification pathway. OASTL-C might also be involved the regulation of sulfur homeostasis since loss or decreased activity due to a single nucleotide polymorphism in Arabidopsis accessions leads to reduced sulfate uptake (Koprivova et al., 2023). In addition, cysteine synthesis was found to play a role during stomatal closure in response to drought stress. Upon soil drying, sulfate is transported to the guard cells via the xylem and is incorporated into cysteine (Ernst et al., 2010). Subsequently, cysteine mediates abscisic acid (ABA)-dependent stomatal closure in multiple ways (Heinemann & Hildebrandt, 2021). It is a substrate of the MoCo-sulfurylase ABA3 involved in ABA synthesis. ABA induces the expression of DES1, which uses cysteine as a substrate to produce hydrogen sulfide in the cytosol. Hydrogen sulfide accumulation leads to persulfidation and activation of several proteins involved in ABA-induced stomatal closure including kinases and transcription factors. These findings illustrate the wide range of functions and different modes of action of cysteine and its related metabolism. Taken together, the versatile chemical nature of cysteine and the high degree of compartmentalization of its metabolism further highlight its potential for multiple, yet unknown, regulatory functions.

In order to identify additional potential signaling functions of cysteine in *Arabidopsis*, we analyzed the response of seedlings to an artificial increase in cysteine levels. The proteome signature of the cysteine treated seedlings indicated perception of the disturbance in cysteine homeostasis as a biotic threat. Thus, we further investigated the role of compartment specific cysteine metabolism in the interaction of *Arabidopsis* plants with the leaf pathogen *Pseudomonas syringae* pv tomato DC3000 (*Pst*) and identified cytosolic and mitochondrial cysteine synthesis as major contributors to pathogen resistance.

Results

The seedling proteome response to increased cysteine concentrations indicates biotic stress signaling

To understand the systemic response of *Arabidopsis thaliana* to an increased cysteine content we performed a feeding experiment. Six-day-old seedlings grown in liquid culture were supplemented with 1 mM L-cysteine for 24h (Fig. 1A). This treatment led to a 7.6-fold increase in the seedling cysteine content, and the glutathione content was also significantly higher (1.7 -fold) than in control seedlings (Fig. 1B). The treated seedlings were visually indistinguishable from the controls and the overall seedling composition with respect to carbohydrates, lipids and proteins remained unchanged with slightly decreased chlorophyll contents indicating no major toxic effects of the cysteine feeding (Suppl. Fig. S1). The cysteine in the medium was completely oxidized within the 24h incubation time. To test, whether cysteine oxidation led to a depletion of oxygen or had an effect on seedling respiration we analyzed the oxygen content of the medium as well as seedling oxygen consumption rates, but did not detect any significant differences between cysteine treatment and control.

Shotgun proteome analysis revealed a distinct effect of cysteine feeding on the composition of the seedling proteome (Fig. 1C). 479 of the 4613 detected protein groups were significantly increased and 670 significantly decreased in cysteine treated compared to control samples (Fig. 1D). Among the strongly downregulated proteins were the transcriptional activator Hem1, kinases associated with ABA signaling, and proteins involved in sulfur assimilation whereas glutathione-S transferases (GSTs) as well as UDP-glucosyltransferases were most drastically increased (Suppl. Dataset 1). In order to systematically identify major features of the proteome response to increased cysteine levels, we performed an enrichment analysis on functional annotations of the significantly changed proteins (Suppl. Dataset S1). Enrichment of metabolic pathways was analyzed based on a modified version of the MapMan annotation system (https://mapman.gabipd.org; Fig. 1E). The results indicated a decrease in pathways related to photosynthesis including photophosphorylation, the Calvin-Benson-Bassham cycle and also tetrapyrrole and organellar protein synthesis (Fig. 1E, blue). In contrast, pathways required for heterotrophic energy metabolism such as carbohydrate and lipid degradation as well as cellular respiration showed increased abundance after cysteine treatment (Fig. 1E, red). The enrichment of stress-related categories (glucosyltransferases, oxidoreductases, glutathione-S-transferases) among the significantly increased proteins prompted us to systematically probe our dataset for characteristic stress response profiles. To this end, we performed an additional enrichment analysis on the basis of published transcriptome profiles for diverse abiotic and biotic stress conditions as well as elicitor and hormone treatments focusing on the proteins significantly increased after cysteine feeding (Fig. 1F, Suppl. Dataset S1). Strikingly, six of the seven significantly enriched categories in cysteine treated seedlings were associated with pathogen response. Enriched categories of proteins were those induced by pathogen-associated molecular patterns (PAMPs), damage-associated molecular patterns (DAMPs), systemic acquired resistance (SAR), treatment with jasmonic acid (JA) or the salicylic acid analogon benzothiadiazole (BTH) strongly suggesting a role for cysteine during biotic stress signaling (Fig. 1F). Indeed, the concentration of the phytoalexin camalexin was significantly (4.5-fold) increased in the cysteine-treated seedlings indicating a similarity to an active pathogen response (Suppl. Fig. 1E).

Fig. 1: **Seedling proteome response to cysteine treatment**. (**A**) An *A. thaliana* seedling culture was supplemented with 1 mM L-cysteine for 24 hours before harvest. (**B**) Seedling content of L-cysteine and reduced glutathione (GSH) [µmol · g fresh weight $^{-1}$] (n = 8-9). Mean (bars) and individual (dots) values ± SD are shown. Asterisks indicate statistically significant differences compared with control seedlings (P < 0.05 > 0.01; *** P < 0,001 > 0,0001; **** P < 0,0001). (**C**) Principal component analysis of shotgun proteomics dataset with control (black) and cysteine-treated (yellow) seedlings (**D**) Volcano plot illustrating differences in the proteome of cysteine treated seedlings vs. controls. Proteins of significantly higher or lower abundance after cysteine feeding are marked in red and blue, respectively (t-test, FDR < 0.05). (**E**) Enrichment of functional categories in proteins that are significantly increased (top, red) or decreased (bottom, blue) in the presence of cysteine. (**F**) Enrichment of stress induced categories among proteins that are significantly increased in the presence of cysteine. ABA, abscisic acid; BCAA, branched-chain amino acid; BTH, benzothiadiazole, DAMP, damage-associated molecular pattern; JA, jasmonic acid; PAMP, pathogen-associated molecular pattern; SAM, S-adenosylmethionine; SAR, systemic acquired resistance. The complete proteomics dataset including the enrichment analysis is provided as Suppl. Dataset S1.

Cysteine treatment induces pathogen resistance

To test the physiological relevance of the observed induction of pathogen response pathways by cysteine feeding we watered six-week-old *A. thaliana* plants grown on soil with 10 mM cysteine 24 hours before performing a pathogen assay using the hemibiotrophic bacterium *Pseudomonas syringae* pv. tomato DC3000 (*Pst*) (Fig. 2A). Quantification of thiols confirmed that cysteine was taken up by the plants and transported to the leaves (Fig. 2B). Cysteine accumulated in the rosette leaves to a similar extent as in the seedling culture (7-fold) in treated compared to control plants. The cysteine treated plants were significantly less susceptible to *Pst* indicated by reduced bacterial growth and less chlorosis after three days of infection (Fig. 2C,D). Cysteine treatment had no significant effect on the photosynthetic performance of the leaves as determined by chlorophyll a fluorescence analysis (Suppl. Figure S2).

Fig. 2: **Cysteine treatment induces resistance to the virulent pathogen *Pseudomonas syringae*.** (**A**) Schematic representation of the workflow: *A. thaliana* plants were grown for 6 weeks under short-day conditions and incubated in 10 mM L-cysteine for 24 h before further analysis. (**B**) Content of L-cysteine and reduced glutathione (GSH) [nmol · g fresh weight^{-1}] in the rosette leaves of mock treated (white bars) and cysteine treated (yellow bars) plants (n = 10) (**C**) *Pst* infection assay: Four leaves per plant were infiltrated with *P. syringae* DC3000 solution including 5 × 10^5 colony forming units (CFU) per ml using a needleless syringe and sampled at 0 and 3 days past infection (dpi) to quantify CFU per cm leaf area (n = 11). (**D**) Phenotype of four representative leaves per treatment at three days after inoculation. Mean (bars) and individual (dots) values ± SD are shown. Asterisks indicate statistically significant differences compared with mock-treated plants (** P < 0.01 > 0.001; *** P < 0,001 > 0,0001).

Pathogen attack induces cysteine synthesis, and proteome responses to cysteine treatment and pathogen interaction overlap

To investigate the function of cysteine during plant-pathogen interaction, we quantified thiols in *Arabidopsis* leaves infected with *Pst*. Cysteine levels were significantly increased already 12 hours post infection (hpi), peaked at 24 hpi at 6.6-fold level of the mock treated controls and remained constantly high until 48 hpi (Fig. 3A). The proteome of the infected leaves showed clear differences to the mock treated samples (Fig. 3B). The pattern of changes in individual protein abundances was highly consistent between the two timepoints analyzed (24 hpi and 48 hpi; Suppl. Dataset S2; Suppl. Fig. S3). Highlighting proteins with significant changes in the cysteine-treated seedlings in the *Pst* infection dataset illustrates that there is also a strong overlap between the *Arabidopsis* proteome response to increased cysteine contents and pathogen attack (Fig. 3C; Suppl. Fig. S3). Among the consistently highly induced proteins in both responses were several glutathione-S-transferases, UDP-glucosyltransferases, and ALTERNATIVE OXIDASE 1A (Suppl. Dataset S2). Enrichment analysis of functional annotations identifies a repression of photosynthesis related pathways, coenzyme metabolism as well as organellar protein translation (Fig. 3D, blue) and an induction of lipid catabolism

(Fig. 3D, red) as common effects of both treatments. Other aspects of the proteome response to *Pst* interaction such as a decrease in terpenoid synthesis and an increase in hormone metabolism or vesicle trafficking did not become apparent during cysteine feeding and thus might be unrelated. The general effect on amino acid metabolism included an increase in the catabolic pathways mainly localized in the mitochondria and a decrease in plastidic synthesis pathways. A clear exception were the synthesis pathway of serine and the aromatic amino acids, which were strongly induced together with downstream pathways required for their conversion to specialized metabolites (Fig. 3E).

Fig. 3: **Cysteine accumulation and proteome response during pathogen interaction**. Six weeks old *A. thaliana* plants were infiltrated with *P. syringae* DC3000 (*Pst*) (2.5 × 10^6 colony forming units per ml). Six leaves per plant were sampled at the indicated time points. (**A**) Content of L-cysteine [nmol · g fresh weight^{-1}] in infected vs. mock treated leaves (n = 4-9). Mean values ± SD are shown. Asterisks indicate statistically significant differences compared with mock-treated plants (**** P < 0,0001). (**B**) Principal component analysis of the proteome of *Pst*- and mock-infected leaves (circles and squares, respectively) at 24 and 48 h after inoculation (black and grey, respectively) (**C**) Volcano plots illustrating

differences in the proteome of *Pst*- vs. mock-infected plants at 24 and 48 hpi. Proteins of significantly higher or lower abundance after treatment with 1 mM L-cysteine (Fig. 1C) are highlighted in red and blue, respectively. (**D**) Enrichment of functional categories in proteins that are significantly increased (top) or decreased (bottom) at 24 hpi with *Pst*. Red and blue fonts indicate significantly increased or decreased categories, respectively, that were also enriched after cysteine treatment (Fig. 1E). (**E**) Effect of *Pst* infection on amino acid metabolic pathways. Relative protein abundances in infected vs. mock treated leaves at 48 hours after inoculation *Pst*. Proteins significantly increasing or decreasing in abundance are indicated by red or blue squares, respectively. 2-OB, 2-oxobutyrate; 2-OG, 2-oxoglutarate; 3PG, 3-phosphoglycerate; ACoA, Acetyl-CoA; AdoMet, S-Adenosylmethionin; C1, C1-metabolism; CM, chorismate; ET, ethylene; FeS, iron-sulfur cluster; FU, fumarate; GABA, γ-aminobutyric acid; Glce, glycerate; Glco, glycolate; GX, glyoxylate; Hpyr, Hydroxypyruvate; MoCo, molybdenum cofactor; NHP, N-hydroxypipecolic acid; OA, oxaloacetic acid; Pyr, pyruvate; R5P, ribose-5-phosphate; SSA, succinic semialdehyde. The complete proteomics dataset including the enrichment analysis is provided in Suppl. Dataset S2.

Compartment specific cysteine synthesis is required for pathogen resistance

A focus on the proteins involved in cysteine metabolism illustrates that the cytosolic and the mitochondrial pathway for cysteine synthesis as well as plastidic GSH production increased in abundance after pathogen attack whereas sulfate reduction and cysteine synthesis in the chloroplasts decreased (Fig. 4A). In order to estimate the contributions of the individual compartments to cellular cysteine production we compared the total abundance of O-acetyl serine lyase (OASTL) isoforms catalyzing the last step in cysteine synthesis in the leaf proteome (Fig. 4B). The cytosolic and plastidic isoforms OASTL-A and OASTL-B were of similar abundance, but OASTL-C localized in the mitochondria represented only about 5% of the total OASTL leaf content. Next, we tested the response of knockout mutant lines for the individual OASTL isoforms during interaction with *Pst* (Fig. 4C-E). All OASTL mutant lines were more susceptible to *Pst* infection than the wild type (Fig. 4D). The strongest effect was visible in *oastl-c* (mitochondrial) mutant lines in the phenotype of the infected leaves as well as in the bacterial growth rates, which were 4.7-fold increased compared to the wild type (Fig. 4D, E). Additionally, all three OASTL mutants showed significantly lower camalexin levels than the wild type at 24 hpi (Fig. 4F). Leaf cysteine levels were lower than in the wild type already under control conditions in *oastl-a* (deficient in cytosolic cysteine synthesis) and also the cysteine increase during infection lagged behind in this line but not in the others (Fig. 4C, grey bars).

Fig. 4: **Compartmentalization of cysteine synthesis in pathogen resistance**. (**A**) Effect of *Pst* infection on cysteine metabolic pathways: Relative protein abundances in infected vs. mock treated leaves at 48 hours after inoculation with *P. syringae* DC3000 (*Pst*) (2.5 × 10⁶ colony forming units (CFU) per ml). Proteins increasing or decreasing in abundance are indicated by red or blue squares, respectively (n.d., undetected proteins). APS, adenosine 5'-phophosulfate; ATPS, ATP sulfhydrase; APR, APS reductase; SiR, sulfite reductase; SERAT, serine acetyltransferase; OAS, O-acetylserine; OAS-A, O-acetylserin(thiol)lyase; GSH, glutathione; GSH1, γ-glutamyl-cysteine synthetase; GSH2, GSH syntethase; SCS, S-sulfocysteine synthase; DES1, L-cysteine desulfhydrase; Pyr, Pyruvate; 3-MP, 3-mercaptopyruvate; STR1, 3-MP sulfurtransferase; GSSH, GSH-persulfide; ETHE1, sulfur dioxygenase; β-CA, β-cyanoalanine; CAS-C1, CAS synthase. (**B**) Total abundance of O-acetylserin(thiol)lyase isoforms in control, mock treated and *Pst* infected leaves of wild type *A. thaliana* plants calculated from quantitative iBAQ-values (n = 4). (**C**) Cysteine content [nmol · g fresh weight⁻¹] in control, mock treated, and *Pst* infected leaves of wild type *A. thaliana* plants and OASTL-deficient mutant lines (n = 5-10). (**D**) *Pst* infection assay: Four leaves per plant were infiltrated with *P. syringae* DC3000 solution including 5 × 10⁵ colony forming units (CFU) per ml using a needleless syringe and sampled at 0 and 3 days past infection (dpi) to quantify CFU per cm leaf area. (n = 8). (**E**) Phenotype of four representative leaves per treatment at three days after inoculation. (**F**) Camalexin content [nmol · g fresh weight⁻¹] in mock treated and *Pst* infected leaves of wild type *A. thaliana* plants and OASTL-deficient mutant lines (n = 4-5). Mean (bars) and individual (dots) values ± SD are shown. Asterisks indicate statistically significant differences compared with wild type plants (** P < 0.01 > 0.001; *** P < 0,001 > 0,0001)

Discussion

Links of cysteine homeostasis to plant immune signaling

The systemic proteome response to cysteine treatment strongly suggests that Arabidopsis seedlings interpret accumulation of cysteine above a certain threshold as a signal for a biotic threat. In accordance with these results, artificially increasing the intracellular cysteine concentration in mature plants renders them significantly less susceptible to infection by the hemibiotrophic bacterium *Pseudomonas syringae,* confirming the protective function of the defense pathways activated in response to cysteine treatment. Cysteine strongly accumulated within the first 24h of infection in control plants that had not been treated prior to inoculation showing that a disturbance in cysteine homeostasis is an integral feature of the interaction between plant and pathogen.

Several proteins were strongly induced by both, cysteine feeding as well as infection with *Pst* and thus might provide some insight into potential links between cysteine homeostasis and immune signaling. The functional category with the strongest enrichment in cysteine induced proteins consisted of UDP-glucosyltransferases (UGTs) of the 73B subfamily annotated as flavonoid glycosidases. Several additional UGTs responded similarly to *Pst* infection and cysteine treatment, and among them UGT71C5, UGT74F2, and UGT87A2 showed the strongest consistent induction (Suppl. Dataset S2). The multigene family of uridine diphosphate-dependent glucosyltransferases catalyzes the covalent addition of sugars from nucleotide UDP sugar donors to functional groups on a variety of compounds which can increase the solubility of lipophilic metabolites or inactivate hormones and signaling molecules (Lairson et al., 2008). A number of UGTs have already been linked to plant immunity. UGT74F2 and UGT76B1 glycosylate and thereby inactivate immune signals including salicylic acid (SA) and N-hydroxypipecolic acid. Knockout lines develop an autoimmune phenotype indicating that this function is required for balancing and containing immune responses (Bauer et al., 2021; Lim et al., 2002). UGT71C5 uses the phytohormone abscisic acid (ABA) as a substrate and is involved in regulating ABA homeostasis (Liu et al., 2015). ABA mediates abiotic stress responses and acts antagonistically to SA induced immune signaling (Torres Zabala et al., 2009; Torres-Zabala et al., 2007). The induction of UGT71C5 by cysteine will therefore amplify the plant pathogen response by suppressing antagonistic signals. A modeling approach recently suggested that UGT71C5 is activated by persulfidation of a cysteine residue at the active site, which would add an additional post-translational layer to sulfur signaling in pathogen response (M. Li et al., 2024). The strongly cysteine-responsive UGTs of the 73B subfamily also seem to have a positive regulatory role in immune signaling since *Arabidopsis* mutant plants lacking functional UGT73B3 or UGT73B5 showed increased susceptibility to *Pst avrRpm1* (Langlois-Meurinne et al., 2005). However, the relevant substrate(s) and the mechanism behind this effect have not been identified yet.

Among the group of proteins accumulating in response to cysteine treatment as well as during pathogen infection were also several glutathione-S-transferases (GSTs) of the plant-specific phi (GSTF) and tau (GSTU) classes. GSTs catalyze the conjugation of glutathione to various substrates, mostly for detoxification purposes, and several of them are induced by pathogen infection and SA (Cummins et al., 2011; Lieberherr et al., 2003; Sappl et al., 2004). However, very limited information is available on endogenous substrates or the exact metabolic functions of disease-induced GST isoenzymes. GSTF2, GSTF8, GSTF10 and GSTF11 were shown to bind to SA, though the biological implications remain unclear (Tian et al., 2012). Also, GSTF2

interacts with camalexin and was suggested to translocate plant defense compounds (Dixon et al., 2011).

Alternative oxidase (AOX) is induced by *Pst* infection and has been proposed to modulate reactive oxygen species (ROS) signaling in the mitochondria (Colombatti et al., 2014; Maxwell et al., 1999; Simons et al., 1999). The balance between AOX and manganese superoxide dismutase activities seems to be relevant for the specificity of ROS signaling by either confining the signal to the mitochondrial matrix (O_2^-) or spreading it to the rest of the cell (H_2O_2) (Cvetkoska & Vanlerberghe, 2012). Our results indicate that induction of AOX can be triggered in response to cysteine accumulation prior to or independently of a pathogen-induced ROS burst.

The general response of the Arabidopsis leaf proteome to pathogen attack revealed that microbe interaction induces profound changes in organelle metabolism. Mitochondria strongly increase their oxidative phosphorylation capacity and use amino acids as alternative respiratory substrates. In contrast, chloroplasts decrease most of their functions including photosynthesis as well as amino acid synthesis and focus on the production of precursors for specialized metabolites and immune signals. This general metabolic shift from growth to defense is also clearly visible in the proteome response of cysteine treated seedlings, which in addition had a slightly decreased chlorophyll content compared to the control. However, the macromolecular composition and respiration rate of the seedlings as well as the photosynthetic performance of the plants was unaffected 24h after cysteine treatment indicating that the systemic induction of defense pathways was able to protect the plant without causing major growth restrictions at this stage.

Interestingly, the protein most decreased in response to cysteine was HEM1, which has recently been reported to act as a translational regulator involved in the attenuation of the immune response during effector-triggered immunity (Zhou et al., 2023). The loss of HEM1 caused exaggerated cell death and restricted bacterial growth. Thus, a major aspect of cysteine signaling during pathogen response might be the repression of antagonistic signals.

Compartment specific functions of cysteine synthesis in immune signaling

While most amino acid anabolic pathways are localized in the chloroplasts, cysteine can be synthesized in different subcellular compartments (Heeg et al., 2008; Watanabe et al., 2008). The total leaf cysteine content after pathogen attack was reduced only in *oastl-A* mutant plants defective in cytosolic cysteine synthesis, but not in *oastl-B* or *oastl-C* indicating that the remaining isoforms can at least partially compensate for the defect. These finding confirm the quantitatively dominant role of the cytosol in cysteine production reported before (Heeg et al., 2008; Krüger et al., 2009). A previous study with a focus on cytosolic cysteine homeostasis had already demonstrated an increased susceptibility of *oastl-A* plants to *Botrytis cinerea* and *Pst* in contrast to a constitutive systemic immune response with high resistance to biotrophic and necrotrophic pathogens in *des1* mutants, which accumulate cysteine due to defects in the cytosolic degradation pathway (Álvarez et al., 2012). Our results are in good agreement with these finding. In addition, they clearly show that cysteine synthesis in the mitochondria is also crucial for Arabidopsis pathogen resistance. Lack of mitochondrial OASTL-C, despite constituting only about 5% of the total OASTL leaf content, had the strongest impact on pathogen response as indicated by the highest bacterial colony counts and leaf chlorosis after infection with *Pst*. Mitochondria require cysteine as a precursor for the synthesis of iron-sulfur clusters and the cofactors biotin and lipoate as well as for cyanide detoxification. It was previously reported that overexpression of the mitochondrial cysteine desulfurase NFS1, which

provides inorganic sulfur for FeS-cluster synthesis, results in constitutive upregulation of defense-related genes and increased resistance against *Pst* (Fonseca et al., 2020). The mechanism behind this response is currently unknown. However, increased cysteine desulfurase activity would potentially lead to an increase in the product sulfane sulfur, which, if produced in excess, can be reduced to hydrogen sulfide in the presence of a reductant like GSH (Frazzon et al., 2007). Hydrogen sulfide production by the mitochondrial β-cyanoalanine synthase CAS-C1 in turn has recently been shown to be crucial for stomatal closure in response to the pathogen-associated molecular pattern flagellin (Pantaleno et al., 2025). NFS1 was significantly induced by cysteine treatment and during *Pst* infection. Thus, an increase in compartment specific cysteine production as well as additional downstream metabolic processes seem to be involved in cysteine signaling during pathogen response.

Sulfate metabolism in plants branches at the level of adenosine 5'-phosphosulfate (APS), which can either be reduced to sulfide and incorporated into cysteine or phosphorylated by APS kinase (APK) and used for sulfation reactions. Blocking the sulfatation branch leads to an increased flux through the reductive branch of the pathway resulting in 4-fold increased leaf cysteine contents in *apk1xapk2* double knockout mutant plants compared to the wild type (Mugford et al., 2009). Re-evaluation of published datasets reveals that the expression profile of the *apk1xapk2* mutant strongly overlaps with the characteristic transcriptional response of *Arabidopsis* plants to bacterial pathogens, which would be in line with cysteine acting as a metabolic signal (Suppl. Figure S4). Since the cytosolic and mitochondrial but not the plastidic isoform of OASTL were induced in the APK deficient plants, compartment specific effects might again be relevant.

Potential mechanisms of cysteine immune signaling

Signaling events can be mediated by receptor proteins coupled to either kinase cascades or ion channels, allosteric protein regulation or post-translational modifications. The mechanism of cysteine sensing and signaling during pathogen response is largely unknown. However, first indications have already been found. Glutamate receptor-like calcium channels (GLRs) are involved in long-distance plant defense signaling in response to wounding (Toyota et al., 2018). GLR3.3 is activated by GSH and several amino acids including cysteine. Treatment of the leaves with GSH or cysteine suppressed *Pst* propagation in wild type but not *glr3-3* mutant plants indicating that induction of pathogen response in reaction to extracellular cysteine is associated with this signaling pathway (F. Li et al., 2013). Based on a detailed transcriptome study and subsequent experimental confirmation clade 2 GLRs have also been linked to immune signaling (Bjornson et al., 2021). Since GLRs are mostly localized in the plasma membrane, they can detect extracellular changes in cysteine and other amino acids due to cell damage or release by pathogens but are unlikely to be involved in intracellular signaling. The induction of proteins associated with a DAMP-response in the cysteine treated seedlings indicates that extracellular receptors might be relevant. Persulfidation of protein cysteine residues has been established as a post-translational modification involved in ABA signaling during drought response (Chen et al., 2020; Shen et al., 2020). In *Aspergillus fumigatus*, a human fungal pathogen causing severe pulmonary infections, persulfidation levels have been linked to both, the pathogenic potential of the fungus as well as the antifungal potency of alveolar macrophages and epithelial cells of the host (Sueiro-Olivares et al., 2021). The potential role of protein persulfidation in plant microbe interactions still has to be established. Although the mechanisms of protein persulfidation and de-persulfidation is not entirely clear yet, cysteine most likely serves as the sulfur donor either via desulfhydration producing hydrogen sulfide or via transamination to 3-mercaptopyruvate and subsequent transsulfuration

(Pedre et al., 2023; Shen et al., 2020). Thus, an increase in cysteine content might be translated to persulfidation signals. Downstream effects can include transcriptional or post-translational regulation. According to our results, the modulation of phytohormones and defense compounds as well as the alteration of immune gene translation pose potential hubs for cysteine-induced biotic stress signaling.

In conclusion, cysteine, an important precursor for defense compounds, was shown to elicit plant immune signaling and thus serve as an infochemical during plant microbe interactions. Compartmental cysteine synthesis appears to be vital for properly mounting a pathogen response. The compartment specific signaling pathways still need to be identified including potential intracellular cysteine receptors as well as connections to hormonal crosstalk during stress response.

Materials and Methods

Plant Material and growth conditions

All experiments were performed with *Arabidopsis thaliana* ecotype Col-0 as wild type control. All mutants were derived from this accession. Homozygous seeds of *oastl-A* (N572213; SALK_072213; aka *oastl-a1.1*) and *oastl-B* (N521183; SALK_021183) as well as of *oastl-C* (N500860; SALK_000860) were kindly provided by Stanislav Kopriva and Andreas Meyer, respectively. Stratified seeds were sown in pots containing soil and grown in a climate chamber under short-day conditions with 8h of light (120 µmol · m^{-2} · s^{-1}). The temperature during the day and night changed from 22 to 18 °C. Soil-grown plants were kept under these conditions for 6 weeks until experiments were performed.

For seedling cultures, seeds were incubated in 100% (v/v) ethanol for 2 min followed by two consecutive incubation steps in 6% (v/v) sodium hypochlorite (Carl Roth, Germany) for 2 min each. Afterwards, seeds were washed five times using sterile ddH$_2$O and transferred to liquid growth media containing 0.43% (w/v) Murashige & Skoog salt mixture (Merck, Germany), 3 mM MES; and 0.5% (w/v) sucrose at pH 5.8. Individual cultures consisting of approximately 3 mg seeds in 50 ml liquid growth media were cultivated under long day conditions with 16 h of light (120 µmol · m^{-2} · s^{-1}) shaking at 100 rpm. The temperature during the day and night changed from 22 to 18 °C. Seedlings cultures were kept under these conditions for 6 days until treatment.

Cysteine treatment

For seedling cultures, 1 mM L-cysteine (Merck, Germany) was supplemented after 6 days of growth. The cysteine content in the medium decreased to low µM concentrations within the 24h treatment.

For soil grown plants, pots were soaked in 1 mM MES pH 5.8 with or without 10 mM L-cysteine (Merck, Germany) for 24 hours starting at the beginning of the light period. Afterwards, plants were either harvested and flash frozen in liquid nitrogen immediately or follow-up experiments were performed.

Quantification of total lipids

The total lipid content was determined using the sulpho-phospho-vanillin method described in (Park et al., 2016). Absorbance was measured at 530 nm using a spectrophotometer (Multiscan Sykhigh, Thermo Fisher Scientific, Germany).

Quantification of total carbohydrates

The total carbohydrate content was determined using the phenol-sulphuric acid method described by. (Tamboli et al., 2020). 5 mg of lyophilised plant powder was dissolved in 1 ml of 2.5 N HCl and incubated for 3 h at 95 °C, shaking. The extracts were diluted (1:50) with demineralized water and 10 µl phenol and 1 ml concentrated sulphuric acid were added. After incubation (10 min, 95 °C, shaking) the absorbance was measured at 490 nm with a spectrophotometer (Multiskan Skyhigh, Thermo Fisher Scientific, Germany).

Extraction and quantification of total protein

5 mg lyophilized seedling powder was dissolved in 700 µl methanol (100% (v/v)) and incubated for 20 min shaking at 80 °C. After centrifugation (10 min, 4 °C, 18,800 xg) the pellet was washed twice in 1 ml ethanol (70% (v/v)) and resuspended in 400 µl NaOH (0.1 M). The solution was incubated for 1h shaking at 95 °C and centrifuged again. The protein content of the supernatant was quantified using Ready-to-use Coomassie Blue G-250 Protein Assay Reagent (Thermo Fisher Scientific, Germany) and Albumin Standard 23209 (Thermo Fisher Scientific, Germany).

Quantification of chlorophyll

The quantification of chlorophyll was carried out according to a modified version of the method described by (Lichtenthaler, 1987). 5 mg plant powder was dissolved in 700 ml methanol (100% (v/v)) and incubated for 20 min at 80 °C with shaking. After centrifugation (10 min, 4 °C, 18,800 xg), the chlorophyll content of the supernatant was quantified with a spectrophotometer (Multiskan Skyhigh, Thermo Fisher Scientific, Germany) (wavelengths: 470 nm, 653 nm and 666 nm).

Analysis of seedling respiration

Respiration rates were measured at ambient conditions using a Clark type oxygen electrode (model DW1, Hansatech Instruments Ltd, United Kingdom). Measurements were performed on control seedlings and seedlings fed 1 mM L-cysteine for 24 h in fresh, air-saturated liquid growth medium under constant stirring. After equilibration, 1 mM L-cysteine (Merck, Germany) was added. After respiration measurements, seedlings were lyophilized and oxygen consumption rates were calculated on the basis of dry weight.

Analysis of photosynthetic performance

For Chl a fluorescence analyses, the HEXAGON IMAGING-PAM (Walz, Effeltrich, Germany) was used on whole rosettes. Saturation light pulses of 0.5 s were applied after 30 min dark treatment to determine Fm and Fm' during the illumination with increasing actinic light intensities. Non-photochemical quenching (NPQ) was calculated as $(Fm − Fm')/Fm'$ and Φ_{PSII} as $(Fm' − F')/Fm$ (Baker, 2008).

Pseudomonas syringae infection assays

For bacterial growth curves and thiol quantification, *Pseudomonas syringea pv. tomato* DC3000 (*Pst*), grown on Kings-B Agar (KB) plates containing 50 µg · ml^{-1} Rifampicin, were resuspended in 10 mM MgCl$_2$ to a final concentration of 5 × 10^5 colony forming units (cfu)·ml^{-1} (OD$_{600}$ 0.001). For shot-gun proteomics analysis, 2.5 × 10^6 cfu · ml^{-1} (OD$_{600}$ 0.005) *Pst* DC3000 was prepared as described. Five to seven of the youngest, fully matured leaves per plant were hand-infiltrated with the bacterial suspension using a needless syringe. For bacterial growth curves, leaf discs were cut from infected leaves 0 days past infection (dpi) and 3 dpi and ground in 200 µl sterile MilliQ water using a tissue lyser (Mill Retsch MM400, Retsch GmbH, Haan, Germany). Dilution series were plated on KB plates containing 50 µg · ml^{-1} Rifampicin and 50 µg · ml^{-1} Cycloheximide, and colony forming units were counted after two days of growth at 28 °C. Students t-tests were performed with 8 biological replicates per time point and genotype or treatment from two independent experiments showing similar results. For thiol quantification and shot-gun proteomics, all infected leaves were harvested, flash frozen in liquid nitrogen and ground for further procedures.

Quantification of sulfur metabolites

Sulfur compounds in plant samples were derivatized using bromobimane and quantified by reverse phase high performance liquid chromatography (HPLC). Samples were solved in derivatisation buffer (1.5 mM bromobimane (Merck, Darmstadt, Germany); 32% (v/v) acetonitrile; 10.3 mM EDTA and 103 mM HEPES pH 8) and incubated at 1400 rpm for 30 min in darkness. Afterwards, 15.9 mM methanesulfonic acid was added, cell debris pelleted at 18,000 xg for 5 min and the supernatant filtered using Corning Costar Spin-X (0.22 µm) centrifuge filter tubes (Merck, Darmstadt, Germany). Samples were diluted and measured using an Agilent 1260 Infinity II HPLC (Agilent Technologies, Waldbronn, Germany) by fluorescence detection (ex. 380 nm; em. 480 nm). Peaks were evaluated and quantified using OpenLabCDS software (Agilent, Santa Clara, United States).

Quantification of camalexin

Camalexin was determined by HPLC as described in (Koprivova et al., 2019). In short, approximately 50 mg plant material were extracted in 150 µl of dimethlysulfoxide (DMSO) for 20 min with shaking and centrifuged. 20 µl of extracts were injected into a HPLC system with a Spherisorb ODS-2 column (250 mm x 4.6 mm, 5 µm) and resolved using a gradient of acetonitrile in 0.01% (v/v) formic acid. Camalexin was detected by a FLD detector with an excitation at 318 nm and emission at 368 nm as described in (Bednarek et al., 2011). For the quantification of camalexin external standards were used ranging from 1 pg to 1 ng per µl.

Protein extraction, digestion and sample preparation for proteome analysis via mass spectrometry

Protein extraction, purification and digestion was performed with an adapted single-pot solid-phase-enhanced sample preparations (SP3) protocol from (Mikulášek et al., 2021) which originates from (Hughes et al., 2019). In short, 5 mg of lyophilized plant powder were solved in 500 µl SDT buffer (4% sodium dodecyl sulfate, 0.1 M dithiothreitol, 0.1 M Tris pH 7.6) and incubated at 60 °C for 30 min. Thirty microliters of the supernatants were mixed with 7.5 µl iodoacetamide (0.1 M) and incubated for 30 min in the dark. Then 2 µl dithiothreitol (0.1 M) was added. Equal shares (v/v) of hydrophobic and hydrophilic carboxylate-modified magnetic beads (Sera-Mag: No. 441521050250, No. 241521050250, GE Healthcare) were prepared for protein binding. Here, 600 µg beads were used per sample. Washing steps with ethanol (80%) were performed on a magnetic rack as described in (Mikulášek et al., 2021). The bound

proteins were digested for 18 h at 37 °C with 0.5 µg trypsin (MS grade, Promega) per sample. The peptide-containing supernatants were collected in tubes with low peptide binding properties. The beads were rinsed in 60 µl ammonium bicarbonate (50 mM) to recover the remaining peptides. The eluates were acidified with formic acid and desalted on 50 mg Sep-Pak tC18 columns (Waters). Peptide concentrations were quantified using the Pierce Quantitative Colorimetric Peptide Assay Kit (Thermo Fisher Scientific) and adjusted to the same concentration in 0.1% formic acid.

Shotgun proteomics by ion mobility mass spectrometry (IMS-MS/MS)

For proteome profiling of the cysteine treated seedlings, 200 ng of peptides were injected via a nanoElute 1 (Bruker Daltonic, Bremen, Germany), separated on an analytical reversed-phase C18 column (Aurora Ultimate 25 cm×75 µm, 1.6 µm, 120 Å; IonOpticks) and analyzed with a timsTOF Pro 2 mass spectrometer. Using a multi-staged linear gradient (Eluent A: MS grade water containing 0.1% formic acid, Eluent B: acetonitrile containing 0.1% formic acid, gradient: 0 min, 2%; 54 min, 25%; 60 min, 37%; 62 min, 95%; 70 min, 95% eluent B), peptides were eluted and ionized by a CaptiveSpray 1 ion source with a flow of 300 nl min^{-1}. The ionized peptides were analyzed with a standard data-dependent acquisition parallel accumulation–serial fragmentation application method (DDA-PASEF) with the following settings: Ion mobility window: 0.6–1.6 V·s/cm², 10 PASEF ramps, target intensity of 20,000 (threshold 2,500), and a cycle time of ~1.1 s.

For proteome analysis of plants infected with *Pseudomonas syringae* 400 ng of peptides were injected via a nanoElute 2 UHPLC (Bruker Daltonic) and separated on the same type of analytical column with the same multi-staged gradient as above. Here, the eluting peptides were ionized by a CaptiveSpray 2 source and analyzed with a timsTOF-HT mass spectrometer. It was programmed with the following DDA-PASEF method parameters: Ion mobility window of 0.7–1.5 V·s/cm², 4 PASEF ramps, target intensity 14,500 (threshold 1,200), and a cycle time of ~0.53 s.

Data Processing and Evaluation

The ion mobility spectrometry (IMS–MS/MS) spectra from both experiments were analyzed with MaxQuant software (Cox & Mann, 2008) using default search parameters and TAIR10 (Arabidopsis.org) as database for protein identification. The calculation of label-free quantification (LFQ) values and intensity-based absolute quantification (iBAQ) values were both enabled. Data evaluation was performed using Perseus software (Tyanova et al., 2016). Proteins were excluded from further analysis if they were not detected in at least n−1 biological replicates in at least one of the sample groups. Subsequently, missing values were replaced with randomly chosen low values from a normal distribution. Significant changes were calculated from the LFQ values using Student's t-tests (p = 0.05). Fisher exact tests were performed in Perseus to identify significantly enriched or depleted metabolic pathways as well as environmental response patterns. The metabolic pathway categories were based on a modified version of MapMan (see Suppl. Datasets S1 and S2; mapman.gabipd.org). Several RNA-seq datasets were used to annotate proteins that showed a response to different stimuli, including fungal PAMPs, DAMPs, bacterial PAMPs (Bjornson et al., 2021), hypoxia (Klecker et al., 2014), ABA treatment, JA treatment (Goda et al., 2008), BTH treatment (Wang et al., 2006), or systemic acquired resistance (Gruner et al., 2013). Transcriptomics datasets for additional abiotic stress conditions (heat, cold, drought, salt, darkness) were used as summarized in (Hildebrandt, 2018).

Funding

Research in the labs of TMH, UA, and SK is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC-2048/1 – project ID 390686111. The proteomics unit in TMH's lab (timsTOF-HT, Bruker Daltonic) is funded via DFG-INST 216/1290-1 FUGG.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository (Perez-Riverol et al., 2022) with the dataset identifier PXD054723.

Supplementary Data

Supplementary Figure S1: Seedling response to cysteine treatment: protein, carbohydrate, lipid, chlorophyll and camalexin content, oxygen consumption rate.

Supplementary Figure S2: Photosynthetic performance of Arabidopsis thaliana rosette leaves after cysteine feeding via the roots.

Supplementary Figure S3: Overlap between the Arabidopsis proteome responses to pathogen interaction and cysteine treatment.

Supplementary Figure S4; Overlap between proteome and transcriptome responses to bacterial pathogens and cysteine accumulation.

Supplementary Dataset S1: Proteomics dataset of cysteine-treated Arabdiopsis seedlings

Supplementary Dataset S2: Proteomics dataset of *Pst*-infected Arabidopsis leaves

Supplementary Dataset S3: Raw data for Fig. 1, Fig. 2, Fig. 3 and Fig. 4.

Acknowledgments

We thank Christina Mack, Cosima Sies, and Dagmar Lewejohann for skillful technical assistance and Isabella Rotthäuser for thorough help with plant cultivation.

Author contributions

TMH, JM, and BH designed the research; BH, CA, and JM performed and evaluated the shotgun proteomics experiments; AK and SK quantified camalexin; UA measured photosynthetic parameters, CA analyzed seedling respiration, JM performed all other experiments, TMH, JM and BH analyzed the data; JM and TMH wrote the paper with support from all other authors; TMH agrees to serve as the author responsible for contact and ensures communication.

Conflict of interest

The authors have no conflicts of interest to declare.

References

Alvarez, C., Calo, L., Romero, L. C., Garcia, I., and Gotor, C. 2010. "An O-Acetylserine(thiol)lyase Homolog with l-Cysteine Desulfhydrase Activity Regulates Cysteine Homeostasis in Arabidopsis." *Plant Physiology*, *152*(2), 656–669. https://doi.org/10.1104/pp.109.147975

Álvarez, C., Ángeles Bermúdez, M., Romero, L. C., Gotor, C., and García, I. 2012. "Cysteine homeostasis plays an essential role in plant immunity." *New Phytologist*, *193*(1), 165–177. https://doi.org/10.1111/j.1469-8137.2011.03889.x

Baker, N. R. 2008. "Chlorophyll Fluorescence: A Probe of Photosynthesis In Vivo." *Annual Review of Plant Biology*, *59*(Volume 59, 2008), 89–113. https://doi.org/10.1146/annurev.arplant.59.032607.092759

Bauer, S., Mekonnen, D. W., Hartmann, M., Yildiz, I., Janowski, R., Lange, B., Geist, B., Zeier, J., and Schäffner, A. R. 2021. "UGT76B1, a promiscuous hub of small molecule-based immune signaling, glucosylates N-hydroxypipecolic acid, and balances plant immunity." *The Plant Cell*, *33*(3), 714–734. https://doi.org/10.1093/plcell/koaa044

Bednarek, P., Piślewska-Bednarek, M., Ver Loren Themaat, E., Maddula, R. K., Svatoš, A., and Schulze-Lefert, P. 2011. "Conservation and clade-specific diversification of pathogen-inducible tryptophan and indole glucosinolate metabolism in Arabidopsis thaliana relatives." *New Phytologist*, *192*(3), 713–726. https://doi.org/10.1111/j.1469-8137.2011.03824.x

Birke, H., Heeg, C., Wirtz, M., and Hell, R. 2013. "Successful Fertilization Requires the Presence of at Least One Major O-Acetylserine(thiol)lyase for Cysteine Synthesis in Pollen of Arabidopsis." *Plant Physiology*, *163*(2), 959–972. https://doi.org/10.1104/pp.113.221200

Bjornson, M., Pimprikar, P., Nürnberger, T., and Zipfel, C. 2021. "The transcriptional landscape of Arabidopsis thaliana pattern-triggered immunity." *Nature Plants*, *7*(5), 579–586. https://doi.org/10.1038/s41477-021-00874-5

Böttcher, C., Westphal, L., Schmotz, C., Prade, E., Scheel, D., and Glawischnig, E. 2009. "The Multifunctional Enzyme CYP71B15 (PHYTOALEXIN DEFICIENT3) Converts Cysteine-Indole-3-Acetonitrile to Camalexin in the Indole-3-Acetonitrile Metabolic Network of Arabidopsis thaliana." *The Plant Cell*, *21*(6), 1830–1845. https://doi.org/10.1105/tpc.109.066670

Buchanan, B. B., and Balmer, Y. 2005. "REDOX REGULATION: A Broadening Horizon." *Annual Review of Plant Biology*, *56*(Volume 56, 2005), 187–220. https://doi.org/10.1146/annurev.arplant.56.032604.144246

Chen, S., Jia, H., Wang, X., Shi, C., Ma, P., Wang, J., Ren, M., and Li, J. 2020. "Hydrogen Sulfide Positively Regulates Abscisic Acid Signaling through Persulfidation of SnRK2.6 in Guard Cells." *Molecular Plant*, *13*(5), 732–744. https://doi.org/10.1016/j.molp.2020.01.004

Colombatti, F., Gonzalez, D. H., and Welchen, E. 2014. "Plant mitochondria under pathogen attack: A sigh of relief or a last breath?" *Mitochondrion*, *19*, 238–244. https://doi.org/10.1016/j.mito.2014.03.006

Cooper, C. E., and Brown, G. C. 2008. "The inhibition of mitochondrial cytochrome oxidase by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and physiological significance." *Journal of Bioenergetics and Biomembranes*, *40*(5), 533–539. https://doi.org/10.1007/s10863-008-9166-6

Couturier, J., Touraine, B., Briat, J.-F., Gaymard, F., and Rouhier, N. 2013. "The iron-sulfur cluster assembly machineries in plants: current knowledge and open questions." *Frontiers*

in Plant Science, *4*. Retrieved from https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2013.00259

Cox, J., and Mann, M. 2008. "MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification." *Nature Biotechnology*, *26*(12), 1367–1372. https://doi.org/10.1038/nbt.1511

Cummins, I., Dixon, D. P., Freitag-Pohl, S., Skipsey, M., and Edwards, R. 2011. "Multiple roles for plant glutathione transferases in xenobiotic detoxification." *Drug Metabolism Reviews*, *43*(2), 266–280. https://doi.org/10.3109/03602532.2011.552910

Cvetkoska, M., and Vanlerberghe, G. C. 2012. "Coordination of a mitochondrial superoxide burst during the hypersensitive response to bacterial pathogen in Nicotiana tabacum." *Plant, Cell & Environment*, *35*(6), 1121–1136. https://doi.org/10.1111/j.1365-3040.2011.02477.x

Dixon, D. P., Sellars, J. D., and Edwards, R. 2011. "The Arabidopsis phi class glutathione transferase AtGSTF2: binding and regulation by biologically active heterocyclic ligands." *Biochemical Journal*, *438*(1), 63–70. https://doi.org/10.1042/BJ20101884

Droux, M. 2003. "Plant serine acetyltransferase: new insights for regulation of sulphur metabolism in plant cells." *Plant Physiology and Biochemistry*, *41*(6), 619–627. https://doi.org/10.1016/S0981-9428(03)00083-4

Droux, M. 2004. "Sulfur Assimilation and the Role of Sulfur in Plant Metabolism: A Survey." *Photosynthesis Research*, *79*(3), 331–348. https://doi.org/10.1023/B:PRES.0000017196.95499.11

Ernst, L., Goodger, J. Q. D., Alvarez, S., Marsh, E. L., Berla, B., Lockhart, E., Jung, J., Li, P., Bohnert, H. J., and Schachtman, D. P. 2010. "Sulphate as a xylem-borne chemical signal precedes the expression of ABA biosynthetic genes in maize roots." *Journal of Experimental Botany*, *61*(12), 3395–3405. https://doi.org/10.1093/jxb/erq160

Fonseca, J. P., Lee, H.-K., Boschiero, C., Griffiths, M., Lee, S., Zhao, P., York, L. M., and Mysore, K. S. 2020. "Iron–Sulfur Cluster Protein NITROGEN FIXATION S-LIKE1 and Its Interactor FRATAXIN Function in Plant Immunity." *Plant Physiology*, *184*(3), 1532–1548. https://doi.org/10.1104/pp.20.00950

Foyer, C. H., and Noctor, G. 2011. "Ascorbate and Glutathione: The Heart of the Redox Hub." *Plant Physiology*, *155*(1), 2–18. https://doi.org/10.1104/pp.110.167569

Frazzon, A. P. G., Ramirez, M. V., Warek, U., Balk, J., Frazzon, J., Dean, D. R., and Winkel, B. S. J. 2007. "Functional analysis of Arabidopsis genes involved in mitochondrial iron–sulfur cluster assembly." *Plant Molecular Biology*, *64*(3), 225–240. https://doi.org/10.1007/s11103-007-9147-x

Giovanelli, J., Mudd, S. H., and Datko, A. H. 1985. "Quantitative Analysis of Pathways of Methionine Metabolism and Their Regulation in Lemna." *Plant Physiology*, *78*(3), 555–560. https://doi.org/10.1104/pp.78.3.555

Glawischnig, E. 2007. "Camalexin." *Phytochemistry*, *68*(4), 401–406. https://doi.org/10.1016/j.phytochem.2006.12.005

Goda, H., Sasaki, E., Akiyama, K., Maruyama-Nakashita, A., Nakabayashi, K., Li, W., Ogawa, M., Yamauchi, Y., Preston, J., Aoki, K., Kiba, T., Takatsuto, S., Fujioka, S., Asami, T., Nakano, T., Kato, H., Mizuno, T., Sakakibara, H., Yamaguchi, S., Nambara, E., Kamiya, Y., Takahashi, H., Hirai, M. Y., Sakurai, T., Shinozaki, K., Saito, K., Yoshida, S., and Shimada, Y. 2008. "The AtGenExpress hormone and chemical treatment data set: experimental design, data evaluation, model data analysis and data access." *The Plant Journal*, *55*(3), 526–542. https://doi.org/10.1111/j.1365-313X.2008.03510.x

Gruner, K., Griebel, T., Návarová, H., Attaran, E., and Zeier, J. 2013. "Reprogramming of plants during systemic acquired resistance." *Frontiers in Plant Science*, *4*. https://doi.org/10.3389/fpls.2013.00252

Gupta, R. N., and Spenser, I. D. 1969. "Biosynthesis of the Piperidine Nucleus: The mode of incorporation of lysine into pipecolic acid and into piperidine alkaloids." *Journal of Biological Chemistry*, *244*(1), 88–94. https://doi.org/10.1016/S0021-9258(19)78195-2

Harun, S., Abdullah-Zawawi, M.-R., Goh, H.-H., and Mohamed-Hussein, Z.-A. 2020. "A Comprehensive Gene Inventory for Glucosinolate Biosynthetic Pathway in Arabidopsis thaliana." *Journal of Agricultural and Food Chemistry*, *68*(28), 7281–7297. https://doi.org/10.1021/acs.jafc.0c01916

Hatzfeld, Y., Maruyama, A., Schmidt, A., Noji, M., Ishizawa, K., and Saito, K. 2000. "β-Cyanoalanine Synthase Is a Mitochondrial Cysteine Synthase-Like Protein in Spinach and Arabidopsis1." *Plant Physiology*, *123*(3), 1163–1172. https://doi.org/10.1104/pp.123.3.1163

Heeg, C., Kruse, C., Jost, R., Gutensohn, M., Ruppert, T., Wirtz, M., and Hell, R. 2008. "Analysis of the Arabidopsis O-Acetylserine(thiol)lyase Gene Family Demonstrates Compartment-Specific Differences in the Regulation of Cysteine Synthesis." *The Plant Cell*, *20*(1), 168–185. https://doi.org/10.1105/tpc.107.056747

Heinemann, B., and Hildebrandt, T. M. 2021. "The role of amino acid metabolism in signaling and metabolic adaptation to stress-induced energy deficiency in plants." *Journal of Experimental Botany*, *72*(13), 4634–4645. https://doi.org/10.1093/jxb/erab182

Hell, R., and Wirtz, M. 2011. "Molecular Biology, Biochemistry and Cellular Physiology of Cysteine Metabolism in *Arabidopsis thaliana*." *The Arabidopsis Book*, *2011*(9). https://doi.org/10.1199/tab.0154

Hildebrandt, T. M. 2018. "Synthesis versus degradation: directions of amino acid metabolism during Arabidopsis abiotic stress response." *Plant Molecular Biology*, *98*(1), 121–135. https://doi.org/10.1007/s11103-018-0767-0

Hildebrandt, T. M., Nunes Nesi, A., Araújo, W. L., and Braun, H.-P. 2015. "Amino Acid Catabolism in Plants." *Molecular Plant*, *8*(11), 1563–1579. https://doi.org/10.1016/j.molp.2015.09.005

Höfler, S., Lorenz, C., Busch, T., Brinkkötter, M., Tohge, T., Fernie, A. R., Braun, H.-P., and Hildebrandt, T. M. 2016. "Dealing with the sulfur part of cysteine: four enzymatic steps degrade l-cysteine to pyruvate and thiosulfate in Arabidopsis mitochondria." *Physiologia Plantarum*, *157*(3), 352–366. https://doi.org/10.1111/ppl.12454

Hughes, C. S., Moggridge, S., Müller, T., Sorensen, P. H., Morin, G. B., and Krijgsveld, J. 2019. "Single-pot, solid-phase-enhanced sample preparation for proteomics experiments." *Nature Protocols*, *14*(1), 68–85. https://doi.org/10.1038/s41596-018-0082-x

Klecker, M., Gasch, P., Peisker, H., Dörmann, P., Schlicke, H., Grimm, B., and Mustroph, A. 2014. "A Shoot-Specific Hypoxic Response of Arabidopsis Sheds Light on the Role of the Phosphate-Responsive Transcription Factor PHOSPHATE STARVATION RESPONSE1." *Plant Physiology*, *165*(2), 774–790. https://doi.org/10.1104/pp.114.237990

Koprivova, A., Elkatmis, B., Gerlich, S. C., Trick, M., Harper, A. L., Bancroft, I., and Kopriva, S. 2023. "Natural Variation in OASC Gene for Mitochondrial O-Acetylserine Thiollyase Affects Sulfate Levels in Arabidopsis." *Plants*, *12*(1). https://doi.org/10.3390/plants12010035

Koprivova, A., Schuck, S., Jacoby, R. P., Klinkhammer, I., Welter, B., Leson, L., Martyn, A., Nauen, J., Grabenhorst, N., Mandelkow, J. F., Zuccaro, A., Zeier, J., and Kopriva, S. 2019. "Root-specific camalexin biosynthesis controls the plant growth-promoting effects of

multiple bacterial strains." *Proceedings of the National Academy of Sciences*, *116*(31), 15735–15744. https://doi.org/10.1073/pnas.1818604116

Krüger, S., Niehl, A., Lopez Martin, M. C., Steinhauser, D., Donath, A., Hildebrandt, T., Romero, L. C., Hoefgen, R., Gotor, C., and Hesse, H. 2009. "Analysis of cytosolic and plastidic serine acetyltransferase mutants and subcellular metabolite distributions suggests interplay of the cellular compartments for cysteine biosynthesis in Arabidopsis." *Plant, Cell & Environment*, *32*(4), 349–367. https://doi.org/10.1111/j.1365-3040.2009.01928.x

Lairson, L. L., Henrissat, B., Davies, G. J., and Withers, S. G. 2008. "Glycosyltransferases: Structures, Functions, and Mechanisms." *Annual Review of Biochemistry*, *77*(Volume 77, 2008), 521–555. https://doi.org/10.1146/annurev.biochem.76.061005.092322

Langlois-Meurinne, M., Gachon, C. M., and Saindrenan, P. 2005. "Pathogen-Responsive Expression of Glycosyltransferase Genes UGT73B3 and UGT73B5 Is Necessary for Resistance to Pseudomonas syringae pv tomato in Arabidopsis." *Plant Physiology*, *139*(4), 1890–1901. https://doi.org/10.1104/pp.105.067223

Li, F., Wang, J., Ma, C., Zhao, Y., Wang, Y., Hasi, A., and Qi, Z. 2013. "Glutamate Receptor-Like Channel3.3 Is Involved in Mediating Glutathione-Triggered Cytosolic Calcium Transients, Transcriptional Changes, and Innate Immunity Responses in Arabidopsis." *Plant Physiology*, *162*(3), 1497–1509. https://doi.org/10.1104/pp.113.217208

Li, M., Xiao, L., Sun, K., Qiu, T., Lai, S., Chen, G., Geng, L., Huang, S., and Xie, Y. 2024. "Insights from Structure-Based Simulations into the Persulfidation of Uridine Diphosphate-Glycosyltransferase71c5 Facilitating the Reversible Inactivation of Abscisic Acid." *International Journal of Molecular Sciences*, *25*(17). https://doi.org/10.3390/ijms25179679

Lichtenthaler, H. K. (1987). "[34] Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. In *Methods in Enzymology. Plant Cell Membranes* (Vol. 148, pp. 350–382). Elsevier. https://doi.org/10.1016/0076-6879(87)48036-1

Lieberherr, D., Wagner, U., Dubuis, P.-H., Métraux, J.-P., and Mauch, F. 2003. "The Rapid Induction of Glutathione S-Transferases AtGSTF2 and AtGSTF6 by Avirulent Pseudomonas syringae is the Result of Combined Salicylic Acid and Ethylene Signaling." *Plant and Cell Physiology*, *44*(7), 750–757. https://doi.org/10.1093/pcp/pcg093

Lim, E.-K., Doucet, C. J., Li, Y., Elias, L., Worrall, D., Spencer, S. P., Ross, J., and Bowles, D. J. 2002. "The Activity of ArabidopsisGlycosyltransferases toward Salicylic Acid, 4-Hydroxybenzoic Acid, and Other Benzoates" *Journal of Biological Chemistry*, *277*(1), 586–592. https://doi.org/10.1074/jbc.M109287200

Liu, Z., Yan, J.-P., Li, D.-K., Luo, Q., Yan, Q., Liu, Z.-B., Ye, L.-M., Wang, J., Li, X.-F., and Yang, Y. 2015. "UDP-Glucosyltransferase71C5, a Major Glucosyltransferase, Mediates Abscisic Acid Homeostasis in Arabidopsis." *Plant Physiology*, *167*(4), 1659–1670. https://doi.org/10.1104/pp.15.00053

Maeda, H., and Dudareva, N. 2012. "The Shikimate Pathway and Aromatic Amino Acid Biosynthesis in Plants." *Annual Review of Plant Biology*, *63*(Volume 63, 2012), 73–105. https://doi.org/10.1146/annurev-arplant-042811-105439

Maxwell, D. P., Wang, Y., and McIntosh, L. 1999. "The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells." *Proceedings of the National Academy of Sciences*, *96*(14), 8271–8276. https://doi.org/10.1073/pnas.96.14.8271

Mikulášek, K., Konečná, H., Potěšil, D., Holánková, R., Havliš, J., and Zdráhal, Z. 2021. "SP3 Protocol for Proteomic Plant Sample Preparation Prior LC-MS/MS." *Frontiers in Plant Science*, *12*. Retrieved from https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2021.635550

Moormann, J., Heinemann, B., and Hildebrandt, T. M. 2022. "News about amino acid metabolism in plant–microbe interactions." *Trends in Biochemical Sciences*, *47*(10), 839–850. https://doi.org/10.1016/j.tibs.2022.07.001

Moseler, A., Wagner, S., and Meyer, A. J. 2024. "Protein persulfidation in plants: mechanisms and functions beyond a simple stress response." *Biological Chemistry*, *405*(9-10), 547–566. https://doi.org/10.1515/hsz-2024-0038

Mugford, S. G., Yoshimoto, N., Reichelt, M., Wirtz, M., Hill, L., Mugford, S. T., Nakazato, Y., Noji, M., Takahashi, H., Kramell, R., Gigolashvili, T., Flügge, U.-I., Wasternack, C., Gershenzon, J., Hell, R., Saito, K., and Kopriva, S. 2009. "Disruption of Adenosine-5′-Phosphosulfate Kinase in Arabidopsis Reduces Levels of Sulfated Secondary Metabolites." *The Plant Cell*, *21*(3), 910–927. https://doi.org/10.1105/tpc.109.065581

Návarová, H., Bernsdorff, F., Döring, A.-C., and Zeier, J. 2012. "Pipecolic Acid, an Endogenous Mediator of Defense Amplification and Priming, Is a Critical Regulator of Inducible Plant Immunity." *The Plant Cell*, *24*(12), 5123–5141. https://doi.org/10.1105/tpc.112.103564

Noctor, G., Cohen, M., Trémulot, L., Châtel-Innocenti, G., van Breusegem, F., and Mhamdi, A. 2024. "Glutathione: a key modulator of plant defence and metabolism through multiple mechanisms." *Journal of Experimental Botany*, *75*(15), 4549–4572. https://doi.org/10.1093/jxb/erae194

Pantaleno, R., Scuffi, D., Schiel, P., Schwarzländer, M., Costa, A., and García-Mata, C. 2025. "Mitochondrial ß-Cyanoalanine Synthase Participates in flg22-Induced Stomatal Immunity." *Plant, Cell & Environment*, *48*(1), 537–552. https://doi.org/10.1111/pce.15155

Park, J., Jeong, H. J., Yoon, E. Y., and Moon, S. J. 2016. "Easy and rapid quantification of lipid contents of marine dinoflagellates using the sulpho-phospho-vanillin method." *Algae*, *31*(4), 391–401. https://doi.org/10.4490/algae.2016.31.12.7

Pastorczyk, M., Kosaka, A., Piślewska-Bednarek, M., López, G., Frerigmann, H., Kułak, K., Glawischnig, E., Molina, A., Takano, Y., and Bednarek, P. 2020. "The role of CYP71A12 monooxygenase in pathogen-triggered tryptophan metabolism and Arabidopsis immunity." *New Phytologist*, *225*(1), 400–412. https://doi.org/10.1111/nph.16118

Pedre, B., Talwar, D., Barayeu, U., Schilling, D., Luzarowski, M., Sokolowski, M., Glatt, S., and Dick, T. P. 2023. "3-Mercaptopyruvate sulfur transferase is a protein persulfidase." *Nature Chemical Biology*, *19*(4), 507–517. https://doi.org/10.1038/s41589-022-01244-8

Pedrotti, L., Weiste, C., Nägele, T., Wolf, E., Lorenzin, F., Dietrich, K., Mair, A., Weckwerth, W., Teige, M., Baena-González, E., and Dröge-Laser, W. 2018. "Snf1-RELATED KINASE1-Controlled C/S1-bZIP Signaling Activates Alternative Mitochondrial Metabolic Pathways to Ensure Plant Survival in Extended Darkness." *The Plant Cell*, *30*(2), 495–509. https://doi.org/10.1105/tpc.17.00414

Peiser, G. D., Wang, T.-T., Hoffman, N. E., Yang, S. F., Liu, H., and Walsh, C. T. 1984. "Formation of cyanide from carbon 1 of 1-aminocyclopropane-1-carboxylic acid during its conversion to ethylene." *Proceedings of the National Academy of Sciences*, *81*(10), 3059–3063. https://doi.org/10.1073/pnas.81.10.3059

Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S., Kundu, D. J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., Walzer, M., Wang, S., Brazma, A., and Vizcaíno, J. A. 2022. "The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences." *Nucleic Acids Research*, *50*(D1), D543-D552. https://doi.org/10.1093/nar/gkab1038

Ruffet, M.-L., Lebrun, M., Droux, M., and Douce, R. 1995. "Subcellular Distribution of Serine Acetyltransferase from Pisum sativum and Characterization of an Arabidopsis thaliana

Putative Cytosolic Isoform." *European Journal of Biochemistry*, *227*(1-2), 500–509. https://doi.org/10.1111/j.1432-1033.1995.tb20416.x

Sappl, P. G., Oñate-Sánchez, L., Singh, K. B., and Millar, A. H. 2004. "Proteomic Analysis of Glutathione S-Transferases of Arabidopsis thaliana Reveals Differential Salicylic Acid-Induced Expression of the Plant-Specific Phi and Tau Classes." *Plant Molecular Biology*, *54*(2), 205–219. https://doi.org/10.1023/B:PLAN.0000028786.57439.b3

Shen, J., Zhang, J., Zhou, M., Zhou, H., Cui, B., Gotor, C., Romero, L. C., Fu, L., Yang, J., Foyer, C. H., Pan, Q., Shen, W., and Xie, Y. 2020. "Persulfidation-based Modification of Cysteine Desulfhydrase and the NADPH Oxidase RBOHD Controls Guard Cell Abscisic Acid Signaling." *The Plant Cell*, *32*(4), 1000–1017. https://doi.org/10.1105/tpc.19.00826

Simons, B. H., Millenaar, F. F., Mulder, L., van Loon, L. C., and Lambers, H. 1999. "Enhanced Expression and Activation of the Alternative Oxidase during Infection of Arabidopsis withPseudomonas syringae pv tomato1." *Plant Physiology*, *120*(2), 529–538. https://doi.org/10.1104/pp.120.2.529

Su, T., Xu, J., Li, Y., Lei, L., Zhao, L., Yang, H., Feng, J., Liu, G., and Ren, D. 2011. "Glutathione-Indole-3-Acetonitrile Is Required for Camalexin Biosynthesis in Arabidopsis thaliana." *The Plant Cell*, *23*(1), 364–380. https://doi.org/10.1105/tpc.110.079145

Sueiro-Olivares, M., Scott, J., Gago, S., Petrovic, D., Kouroussis, E., Zivanovic, J., Yu, Y., Strobel, M., Cunha, C., Thomson, D., Fortune-Grant, R., Thusek, S., Bowyer, P., Beilhack, A., Carvalho, A., Bignell, E., Filipovic, M. R., and Amich, J. 2021. "Fungal and host protein persulfidation are functionally correlated and modulate both virulence and antifungal response." *PLOS Biology*, *19*(6), e3001247. https://doi.org/10.1371/journal.pbio.3001247

Szabados, L., and Savouré, A. 2010. "Proline: a multifunctional amino acid." *Trends in Plant Science*, *15*(2), 89–97. https://doi.org/10.1016/j.tplants.2009.11.009

Takahashi, H. (2010). "Chapter 4 - Regulation of Sulfate Transport and Assimilation in Plants. In K. W. Jeon (Ed.), *International Review of Cell and Molecular Biology* (Vol. 281, pp. 129–159). Academic Press. https://doi.org/10.1016/S1937-6448(10)81004-4

Takahashi, H., Kopriva, S., Giordano, M., Saito, K., and Hell, R. 2011. "Sulfur Assimilation in Photosynthetic Organisms: Molecular Functions and Regulations of Transporters and Assimilatory Enzymes." *Annual Review of Plant Biology*, *62*(Volume 62, 2011), 157–184. https://doi.org/10.1146/annurev-arplant-042110-103921

Tamboli, F. A., More, H. N., Bhandugare, S. S., Patil, A. S., Jadhav, N. R., and Killedar, S. G. 2020. "Estimation of Total Carbohydrate content by Phenol Sulphuric acid Method from Eichhornia crassipes (Mart.) Solms." *Asian Journal of Research in Chemistry*, *13*(5), 357–359. https://doi.org/10.5958/0974-4150.2020.00067.X

Tian, M., Dahl, C. C. von, Liu, P.-P., Friso, G., van Wijk, K. J., and Klessig, D. F. 2012. "The combined use of photoaffinity labeling and surface plasmon resonance-based technology identifies multiple salicylic acid-binding proteins." *The Plant Journal*, *72*(6), 1027–1038. https://doi.org/10.1111/tpj.12016

Torres Zabala, M. de, Bennett, M. H., Truman, W. H., and Grant, M. R. 2009. "Antagonism between salicylic and abscisic acid reflects early host–pathogen conflict and moulds plant defence responses." *The Plant Journal*, *59*(3), 375–386. https://doi.org/10.1111/j.1365-313X.2009.03875.x

Torres-Zabala, M. de, Truman, W., Bennett, M. H., Lafforgue, G., Mansfield, J. W., Rodriguez Egea, P., Bögre, L., and Grant, M. 2007. "*Pseudomonas syringae* pv. *tomato* hijacks the *Arabidopsis* abscisic acid signalling pathway to cause disease." *The EMBO Journal*, *26*(5), 1434-1443-1443. https://doi.org/10.1038/sj.emboj.7601575

Toyota, M., Spencer, D., Sawai-Toyota, S., Jiaqi, W., Zhang, T., Koo, A. J., Howe, G. A., and Gilroy, S. 2018. "Glutamate triggers long-distance, calcium-based plant defense signaling." *Science*, *361*(6407), 1112–1115. https://doi.org/10.1126/science.aat7744

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M. Y., Geiger, T., Mann, M., and Cox, J. 2016. "The Perseus computational platform for comprehensive analysis of (prote)omics data." *Nature Methods*, *13*(9), 731–740. https://doi.org/10.1038/nmeth.3901

Van Hoewyk, D., Pilon, M., and Pilon-Smits, E. A. 2008. "The functions of NifS-like proteins in plant sulfur and selenium metabolism." *Plant Science*, *174*(2), 117–123. https://doi.org/10.1016/j.plantsci.2007.10.004

Wang, D., Amornsiripanitch, N., and Dong, X. 2006. "A Genomic Approach to Identify Regulatory Nodes in the Transcriptional Network of Systemic Acquired Resistance in Plants." *PLOS Pathogens*, *2*(11), e123. https://doi.org/10.1371/journal.ppat.0020123

Watanabe, M., Kusano, M., Oikawa, A., Fukushima, A., Noji, M., and Saito, K. 2008. "Physiological Roles of the β-Substituted Alanine Synthase Gene Family in Arabidopsis." *Plant Physiology*, *146*(1), 310–320. https://doi.org/10.1104/pp.107.106831

Wirtz, M., and Hell, R. 2006. "Functional analysis of the cysteine synthase protein complex from plants: Structural, biochemical and regulatory properties." *Journal of Plant Physiology*, *163*(3), 273–286. https://doi.org/10.1016/j.jplph.2005.11.013

Zhou, Y., Niu, R., Tang, Z., Mou, R., Wang, Z., Zhu, S., Yang, H., Ding, P., and Xu, G. 2023. "Plant HEM1 specifies a condensation domain to control immune gene translation." *Nature Plants*, *9*(2), 289–301. https://doi.org/10.1038/s41477-023-01355-7