Agricultural and Forest Entomology

Larger and polymorphic noctuid moths tend to show less inter-annual abundance variation in the canopy of a temperate forest

Journal:	*Agricultural and Forest Entomology*
Manuscript ID	Draft
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Molleman, Freerk; Adam Mickiewicz University, Department of Systematic Zoology Wąsala, Roman; Poznan University of Life Sciences, Department of Entomology and Environmental Protection Tambay, Ahmet; Adam Mickiewicz University, Department of Systematic Zoology Davis, Robert; University of Tartu, Department of Zoology Õunap, Erki; University of Tartu, Department of Zoology Walczak, Urszula; Adam Mickiewicz University, Department of Systematic Zoology
Keywords:	population dynamics, migration, canopy, phenology, phylogeny
Abstract:	Inter-annual variation in insect abundance and seasonal phenology can be related to species traits such as body size, larval diet, overwintering stage, and colour variations. We sampled noctuid moths in the canopy of a forest in Western Poland using Malaise-interception traps during two vegetative seasons. We calculated inter-annual variation for 23 species and date of peak abundance and length of the flight season for 17 species. We found that larger moths and those with adult colour variation showed less inter-annual abundance variation, which corroborates results of previous studies. We found no indication that phenological traits are associated with the tested species traits. However, species that can feed on the dominant broad-leaved tree in which the traps were placed (Quercus petraea) tended to have a later date of peak abundance than other species. Further work may uncover whether smaller-bodied species and those lacking colour variations are more susceptible to climate change and other stressors.

Temporal abundance and species traits

Larger and polymorphic noctuid moths tend to show less inter-annual abundance variation in the canopy of a temperate forest

Freerk Molleman[1], Roman Wąsala[2], Ahmet Tambay[1], Robert B. Davis[3], Erki Õunap[3], Urszula Walczak[1]

1 Department of Systematic Zoology, Institute of Environmental Biology, Faculty of Biology, Adam Mickiewicz University in Poznań, Uniwersytetu Poznańskiego Str. 6, PL-61-614 Poznań, Poland
fremol@amu.edu.pl, 0000-0002-6551-266X
urszulaw@amu.edu.pl, 0000-0003-3698-7831
ahmtam@st.amu.edu.pl

2 Department of Entomology and Environmental Protection, Poznan University of Life Sciences, Dąbrowskiego 159, 60-594 Poznań, Poland.
roman.wasala@up.poznan.pl, 0000-0002-7825-3273

3 Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu, J. Liivi 2, 50409 Tartu, Estonia.
davis@ut.ee, 0000-0002-0104-0747
erki.ounap@ut.ee, 0000-0001-6489-0055

Abstract

Inter-annual variation in insect abundance and seasonal phenology can be related to species traits such as body size, larval diet, overwintering stage, and colour variations. We sampled noctuid moths in the canopy of a forest in Western Poland using Malaise-interception traps during two vegetative seasons. We calculated inter-annual variation for 23 species and date of peak abundance and length of the flight season for 17 species. We found that larger moths and those with adult colour variation showed less inter-annual abundance variation, which corroborates results of previous studies. We found no indication that phenological traits are associated with the tested species traits. However, species that can feed on the dominant broad-leaved tree in which the traps were placed (*Quercus petraea*) tended to have a later date of peak abundance than other species. Further work may uncover whether smaller-bodied species and those lacking colour variations are more susceptible to climate change and other stressors.

Keywords: population dynamics, migration, phenology, canopy

INTRODUCTION

Climate change and other anthropogenic factors are impacting both the population sizes and phenology of insects in temperate habitats (Wagner et al., 2021, Fox et al., 2014), but we have limited understanding on how this is affected by species traits. The abundance of adult insects at a given location varies as a result of their phenology, local population dynamics, and movements (Molleman, 2018). These processes are, on the one hand, affected by environmental variation (in particular weather), and on the other, by species traits such as diet and body size (Seifert et al., 2023, Tammaru and Haukioja, 1996, Gaston, 1988). Moreover, a species' traits can affect how it is affected by environmental variation (Végvári et al., 2015). More insight into insect temporal abundance variation and its drivers is needed to predict species' responses to the changing climate (Coulthard et al., 2019).

Moths, in particular noctuids, are often used as a model group to study these temporal dynamics, as they are often abundant, species-rich, and can be sampled with light traps at ground level. As the effectiveness of light traps is affected by ambient light (moon phase, cloud cover) and vegetation structure, and such traps attract moths from unknown distances, Malaise traps can potentially provide more reliable data (Butler et al., 1999, Busse et al., 2022, La Cava et al., 2024). Furthermore, studies including trapping in the canopy have also recovered pronounced vertical stratification so that parts of the moth community are missed by ground-level sampling (De Smedt et al., 2019, Ashton et al., 2016, Ruchin, 2023, La Cava et al., 2024, Böttger et al., 2025). Analyses of temporal patterns in the abundance of moths found in the canopy of forests are scarce, even for the temperate zone.

Variation in inter-annual abundance of adult moths results from population dynamics within the typical dispersal range of the species. Thus, the temporal patterns observed at a particular location will, in some species, represent very local population dynamics (possibly even at the level of an individual tree), and in highly dispersive or migratory species may reflect those happening over hundreds of kilometres. Thus, abundance fluctuations are not the same as population dynamics, and this may lead to wrongful interpretation of abundance fluctuations measured at a given location (Molleman, 2018). Species traits that may affect abundance fluctuations are often correlated with each other. Migratory behaviour tends to be more common in larger-bodied species and in those that are more polyphagous (Nieminen et al., 1999, Logghe et al., 2024). The effect of adverse weather on adults may also be stronger for smaller-bodied and shorter-lived species, and individuals of smaller-bodied moth species also tend to be shorter-lived (Holm et al., 2016). In particular, smaller-bodied species may more often fail to reproduce when their short life span happens to coincide with adverse weather, possibly leading to more erratic population fluctuations (Gaston, 1988, Gaston and Lawton, 1988). Consistently, larger arthropods feeding on bracken ferns were shown to have less variable abundance than smaller-bodied species (Gaston and Lawton, 1984). Overall, body size is likely correlated with the degree of inter-annual variation in population size due to such indirect relationships (Gaston and McArdle, 1994, Gaston and Reavey, 2009).

Larval diet breadth can also affect population dynamics. When these dynamics result from interactions with a small number of resources (host-plant species) and natural enemies

Temporal abundance and species traits

(parasitoids, predators, and pathogens), the overall population trends will tend to be more stochastic (MacArthur, 1955). Species that require close synchrony between their phenology and the phenology of host plants and suitable weather also tend to show a lot of inter-annual abundance variation (Price, 2003, Tammaru and Haukioja, 1996). Specifically, populations of spring feeders often appear to respond strongly to variation in synchrony between egg hatching and budburst phenology (e.g. winter moths) and also suffer more from unseasonably late episodes of frost than those that breed during the summer (Hunter and Elkinton, 2000, Varley and Gradwell, 1967). Furthermore, long-term studies on moths usually show more inter-annual abundance variation in species with more polyphagous larvae (Spitzer and Jaroš, 2009 and associated bibliography). Lastly, variable adult colouration, which is thought to reduce predation risk when moths occur at high densities, is associated with reduced inter-annual variation in the abundance of noctuid moths (Forsman et al., 2015). Overall, we can expect that body size, host-plant relationships, and adult polymorphism are correlated with inter-annual variation in moth abundance.

Moth phenology is usually controlled by temperature, photoperiod, or a combination of these factors (Valtonen et al., 2011), and is thus likely to both plastically and adaptively change as the climate is warming (Teder, 2020, O'Neill et al., 2012, Hickinbotham et al., 2024, Hällfors et al., 2021). Other species traits are also correlated with phenology, and may affect how phenology responds to climate change. This may include body size (Foerster et al., 2024). For example, smaller-bodied, spring-feeding species may be expected to complete their development faster and thus emerge earlier in the season. Across the entire season, species of geometrid moth whose larvae develop in the spring are generally larger than species developing during late summer (Seifert et al., 2023). Overwintering stage is also related to moth phenology, as species overwintering as pupae emerge earlier than those that overwinter as larvae or eggs (Hickinbotham et al., 2024). Furthermore, species feeding on woody plants tend to emerge later than those feeding on herbaceous plants (Hickinbotham et al., 2024). Moths feeding on woody plants also show larger shifts in flight periods but smaller increases in voltinism in response to climate change than those feeding on herbaceous plants (Altermatt, 2010). Moreover, phenological responses to climatic variation can be stronger for species with later overwintering life stages (e.g. overwintering as adults vs as eggs; Hickinbotham et al., 2024). Similarly, spring-flying species shift their phenology more in response to climate warming than summer- or autumn-flying species (Maurer et al., 2018). Overall, body size, timing of development, overwintering stage, and type of host plant can be expected to correlate with the phenology of moths.

To gauge how species traits are related to inter-annual variation in adult abundance and phenology of moths in the canopy of a temperate forest, we sampled noctuid moths in the canopy of a forest in Western Poland using Malaise-interception traps during two years. To our knowledge, this is the first such study performed on noctuids in the canopy of a forest. Moreover, we used traps that do not attract but only intercept ambient moths, in contrast to most moth studies that use light traps (see e.g. Butler et al., 1999 , Busse et al., 2022 for moth sampling with Malaise traps).

METHODS

Study site and insect collection

We sampled moths in Puszcza Zielonka forest, Western Poland (52.524, 17.059). Most compartments of this forest are dominated by pine (*Pinus sylvestris* L.) or sessile oak (*Quercus petraea* (Matt.) Liebl.), while some smaller areas have mainly beech (*Fagus sylvatica* L.), pedunculate oak (*Q. robur* L.), or spruce (*Picea abies* (L.) H. Karst.; Forest_Data_Bank, 2018). In some areas, there is a subcanopy of hornbeams (*Carpinus betulus* L.; Forest_Data_Bank, 2018).

We trapped insects with Malaise-interception traps (Figure 1; Gossner et al., 2014) during the vegetative seasons of 2019 (April–October) and 2020 (March–October) in the canopy of 34 sessile oak trees (the same trees as used in Molleman et al., 2022 with some additions). Focal oak trees were located in four stands near Zielonka village and seven stands near Kamińsko village in sets of two to four trees. Within sets, trees were selected to vary widely in the species identity of neighbouring trees (mainly oaks, pines, beeches, or hornbeams). Effects of tree neighbourhood and tree traits on noctuoid moth communities captured will be treated elsewhere. Traps were serviced every 3–4 weeks. Usually, not all traps could be serviced on one day, causing slight differences in collection dates among trees. Moths were preserved in the traps in polypropylene glycol and then transferred to jars with 70% ethanol, and these were kept at -20 °C. The macro moths were sorted from the samples and provisionally set (curated at AMU). Moths were identified to species using genitalia dissection when needed (Fibiger, 1990, Fibiger, 1993, Fibiger, 1997, Fibiger and Hacker, 2007, Fibiger et al., 2009, Fibiger et al., 2010, Ronkay and Ronkay, 1994, Ronkay and Ronkay, 1995, Ronkay et al., 2011, Ronkay et al., 2001).

Species traits

To obtain body size data, we measured the forewing length and thorax width of up to ten individuals of each species. We used specimens from our traps when possible, but completed the data using other specimens collected in the region. We then calculated wing loading as wing length^2/thorax width^3. To obtain information on larval diet, overwintering stage, feeding period, migratory behaviour, and polymorphism, we used authoritative websites (Wagner, 2024, Lepiforum, 2020, Jonko, 2020).

Temporal abundance and species traits

Figure 1. Malaise-interception trap suspended on a rope that passes over a high branch. Traps were pulled into the canopy using a pulley system and lowered for servicing about once a month. Insects intercepted by the see-through mesh either climbed up and ended up in the top container or fell into the bottom container. Both containers contained polypropylene glycol to preserve the insects. The bottom container was fitted with a hole covered with fine wire mesh near the top to allow excess rainwater to flow out while retaining captured insects.

Data analysis

We calculated inter-annual variation as 'one minus (the smallest number divided by the largest number)' for each species for which we caught at least ten specimens. Given that we have two years of data, this is preferable over the coefficient of variation (CV) recommended for larger data sets (Gaston and McArdle, 1994), as it is simple, little affected by the mean, and does not rely on assumptions on the data distribution.

To reconstruct the phenology of each species of moth collected from the traps, we summed the number of individuals over all the traps in a given trapping period (time between the current and previous trap servicing day). It could take multiple days to collect material from all the traps, and the servicing day was defined as the middle of this period. We subsequently calculated the number of individuals caught per trap per day. This value was then assigned to the middle of the period that the traps had been active, assuming homogenous capture probability during each trapping period. For the species of which we caught at least 50 individuals, we then plotted the abundance per trap per day against date for both years separately, and fitted a line using the loess function in R (R_Core_Team, 2024). From this line, we extracted the date of peak abundance and the first and last occurrence of the moth species (abundance > 0). The length of the flight period was then calculated as the time between the first and last occurrence of the species.

Temporal abundance and species traits

We created the timed phylogenetic tree of Noctuidae to be able to perform phylogenetic comparative analyses by relying on the recent phylogeny of Noctuidae (Nedumpally et al., in prep). We used this phylogeny as the fixed topology and incorporated the corresponding molecular data matrix. Since the full molecular dataset was too large for node age calculations, we selected the 24 longest protein-coding loci that contained data for at least 90% of the taxa, totalling 53,990 bp. First, we converted the initial tree into an ultrametric one using the makeChronosCalib and chronos functions in the R package ape (Paradis and Schliep, 2019). Because the full tree of 333 species was too large for node age calculations, we split the tree and corresponding molecular dataset into two subsets. We used the drop.tip function in ape to exclude almost all Noctuinae: Xylenini + Apameini from one subset (subset 1) and almost all Noctuinae: Noctuini + Hadenini s.l. from the other (subset 2). Both subsets retained 37 outgroup species from eight families of Macroheterocera (Drepanidae, Uraniidae, Geometridae, Lasiocampidae, Saturniidae, Sphingidae, Notodontidae, and Erebidae) along with 90 to 91 species of Noctuidae. The final data matrix included 246 species for subset 1 and 227 species for subset 2. To determine the optimal partitioning scheme based on predefined loci, we used MODELFINDER (Kalyaanamoorthy et al., 2017) within IQ-TREE 2.3.2 (Nguyen et al., 2015) on the CIPRES portal (Miller et al., 2010). We followed the results suggested by MODELFINDER when dividing the data into partitions for node age calculations, resulting in 13 partitions for subset 1 and 15 for subset 2.

We calculated node ages using BEAST v1.10.4 (Suchard et al., 2018) at the High-Performance Computing Center of the University of Tartu (Anonymous, 2025). Since test runs with more complex clock models failed to converge, we applied a strict molecular clock, linked clock and tree priors across all partitions, and implemented a birth–death tree prior with incomplete sampling (Stadler, 2009) for the final runs. To time-calibrate both ultrametric trees, we defined five calibration points for each subset based on Kawahara et al. (2019), with ages (in millions of years) as follows: root of the tree (Normal distribution, mean 93.7 ± 11.1 SD), Noctuoidea (Normal distribution, mean 77.6 ± 11.0 SD), Erebidae + Noctuidae clade (Normal distribution, 68.6 ± 10.2 SD), Chloridea + *Helicoverpa* clade (Normal distribution, 9.3 ± 5.6 SD), and *Spodoptera* (Normal distribution, mean 10.6 ± 5.7 SD). For each subset, we ran a single BEAST analysis with Bayesian MCMC for 30 million generations, sampling every 1000th generation. After inspecting the results in TRACER v1.7.2 (supplementary software to BEAST), we discarded the first 3 million generations as 'burn-in.' We used TREEANNOTATOR v1.10.4 (also supplementary software to BEAST) to construct the final trees for both subsets. Finally, we combined the two partitions to generate the full tree, making minor branch length adjustments in the basal branches to ensure ultrametricity.

To investigate relationships between species traits and inter-annual abundance variation, we conducted phylogenetic comparative analyses. For continuous predictors, such as mean forewing length, we performed PGLS analyses using the R package ape (Paradis and Schliep, 2019). When possible, we applied the Pagel model to account for phylogenetic signal. We tested residual normality using a Shapiro–Wilk test and QQ plot inspections. When necessary, we reran PGLS analyses after removing outlying data points. For categorical predictors, such as food type, we conducted a phylogenetic ANOVA using the phylANOVA function in the R

package phytools (Revell, 2012). For categorical predictors that included groups with only one species, we ran analyses both with and without those groups, as phylANOVA only provides pairwise t-values for groups with more than one species.

To explore relationships between species traits and phenology, we first made a correlogram for each study year using the R package corrplot for traits that were continuous or binary (Wei and Simko, 2021). As the number of species with at least 50 individuals captured was too small to implement phylogenetic comparative methods, we then implemented linear models for the average date of peak abundance and the average length of the flight period with as predictors the species traits in R (R_Core_Team, 2024). We then performed AICc model selection using the R package MuMln (Barton, 2022), limiting the maximum number of predictors to three to avoid over-parametrization.

RESULTS

Overview of the data: We caught 5300 individual noctuoid moths (link to Dryad data, see additional files). For 23 species we caught at least ten individuals (Appendix 1), and of 17 species at least 50 individuals (Figure 2). Most of the species had their abundance peak during summer (July–August), and a few in early autumn (October).

Temporal abundance and species traits

Figure 2. Phenology of the most abundant noctuid moths (N > 50) during 2019 (left) and 2020 (right) sampled with Malaise-interception traps in the canopy of a forest in Western Poland.

Temporal abundance and species traits

Phylogeny of the study species
The sufficiently abundant noctuoids sampled all belonged to the family Noctuidae and these species were included in the larger analyses from which we reconstructed their phylogeny (Figure 3).

Figure 3. Reconstructed phylogeny of the noctuid moths used in this study.

Inter-annual variation in abundance: Inter-annual variation in abundance varied from species that were caught only during one of the study years, to species that were quite evenly distributed across the two years (Figure 4). Inter-annual variation in abundance tended to be less pronounced in noctuid moths with longer wings (Figure 4, Table 1), and in species with pronounced adult colour variation (Figure 4; Table 1). Other species traits did not show significant relationships with inter-annual variation in abundance (Table 1).

Fig 4. Relationship between average forewing length and inter-annual variation in adult abundance of noctuid moths (1 - (abundance in lowest abundance year / abundance in highest abundance year)). Species with adult colour variation (polymorphic) are indicated with grey circles. See Table 1 for results of statistical analyses accounting for phylogeny.

Temporal abundance and species traits

Table 1. Results of phylogenetic comparative analyses of inter-annual variation in noctuid moth abundance. Where N is marked with **, outliers have been removed. The exclusion of outliers had no qualitative impact on the result of the analyses. P-values < 0.05 are in bold font. Thorax width = the average width of the thorax; Wing length = average forewing length; WA/TV = wing loading calculated as Wing length^2/Thorax width^3; Oak diet = whether the larval diet includes the dominant broad-leaved tree species (*Q. petraea*) in which traps were placed; Dietary specialization = specialist (one host plant genus), generalist (feeding on multiple genera, or grass-feeding); Plant form = whether larvae feed on trees, herbs, trees and herbs, or on grasses; Food type = leaves, roots, or stems; Food type excluding stem = same as Food type but excluding the one species feeding on stems; Overwintering stage = egg, larva, pupa, imago, or not overwintering in Poland; Feeding period = time of year when larvae are feeding (spring, summer, spring & summer, spring & autumn, summer & spring, vegetation period); Colour polymorphism = whether the species is known to have adult colour variations within the population; Migration = whether the species is known to migrate or have mass dispersal events.

Predictor	N	Test	P	Coefficient (continuous predictor), Difference in means (categorical predictor) for significant results
Mean TW	21**	PGLS	0.2284	
Mean FW	21**	PGLS	**0.0219**	-0.0304
Mean WATV	20**	PGLS	0.6785	
Oak diet	23	phylANOVA	0.3958	
Diet specialization	23	phylANOVA	0.8970	
Diet specialization (Gen v Gram)	22	phylANOVA	0.6266	
Plant form	23	phylANOVA	0.5411	
Food type	23	phylANOVA	0.9811	
Food type (no stem-feeder)	22	phylANOVA	0.6626	
Overwintering stage	23	phylANOVA	0.4415	
Feeding period simple	23	phylANOVA	0.6055	
Feeding period simple (no autumn–spring)	22	phylANOVA	0.7279	
Colour polymorphism	23	phylANOVA	**0.0093**	Yes = 0.517, No = 0.764
Migration	23	phylANOVA	0.4090	

Temporal abundance and species traits

Phenology: Species varied in the date of peak abundance and length of their flight season, with the peak abundance ranging from June to October, and the flight period tending to be long, ranging from 123 to 224 days (Figure 2). Across these species, the date of peak abundance did not differ significantly between the two study years (paired t-test: t = -1.66, df = 17, p-value = 0.12). The length of the flight season was not correlated with overall abundance (Figure 5), such as would be the case if sample size was too small to capture the actual flight period. In 2019, oak-feeding moths tended towards a shorter flight season than non-oak-feeding moths (p = 0.06, Figure 5), but no such tendency was detected in 2020. Oak-feeding species tended to have a later peak date than non-oak-feeding species (significant at p = 0.01 in 2020, Figure 5).

Figure 5. Correlations between phenology and species traits for 2019 and 2020 with p-values when they were < 0.1.

Model selection based on AICc for the average length of the flight season resulted in a model without predictors, followed by the model with forewing length. However, forewing length was not a significant predictor (R^2 = 8%, Adj-R^2 = 2%, F = 1.38 on 1 and 15 DF, P-value = 0.26). Similarly, the top model for the average date of peak abundance did not include any predictors, and the second best included only wing length, but this was not significant either (R^2 = 11%, AdjR^2 = 5%, F = 1.81 on 1 and 15 DF, P-value = 0.20).

Temporal abundance and species traits

DISCUSSION

We sampled noctuid moths in a forest in Poland during two seasons and explored relationships between some species traits that have been suggested to correlate with inter-annual variation in abundance and phenology. We used Malaise-interception traps rather than the usual light traps, avoiding some of the biases associated with the latter. Furthermore, we trapped in the canopy in contrast to other multi-year moth studies that were all performed at ground level. Even though the two-year duration is the minimum for looking into inter-annual variation, and the number of species for which we had enough data was only twenty-three, we found that larger moths and those with adult colour variation showed less abundance variation, which corroborates results of previous studies (e. g. Gaston and Lawton, 1988, Forsman et al., 2015). Notably, species traits with more than two levels never showed a significant effect, which may be due to the low number of species available in our data set. In addition, we described the phenology of seventeen species, but found no indication that phenological traits are associated with the tested species traits. However, species that were able to feed on the dominant broad-leaved tree in which the traps were placed (*Q. petraea*) tended to have a later date of peak abundance than other species.

The effect of body size on inter-annual variation in moth abundance can result from various indirect effects (see discussion in Gaston and McArdle, 1994, and Introduction). It may be related to life span, dispersal, and susceptibility to weather. Colour polymorphism is thought to allow populations to maintain higher densities, as it reduces search image formation in predators such as birds (Bond, 2007, Karpestam et al., 2016). The association between reduced inter-annual variation in abundance and colour polymorphism that we found in our data was shown before for noctuids using longer time series of more species (Forsman et al., 2015). Overall, our results suggest that small-bodied species lacking adult colour variation may vary more in abundance and may thus be more prone to (local) extinction. Further work may uncover whether this translates into greater susceptibility to climate change and other stressors.

That species that are able to feed on the dominant broad-leaved tree in which the traps were placed (*Q. petraea*) tended to have a later peak date than other species suggests that these moths eclose later in the season than other species that developed in the understory or immigrated. Perhaps the late budburst of oaks makes their leaves available for herbivores later than those of most other plant species (Molleman and Walczak, 2024, Wesołowski and Rowiński, 2006), thus causing their later emergence. Otherwise, no species traits appeared associated with their date of peak abundance, nor with the length of their flight season. The results of these analyses, however, must be taken with some caution, as we had only 17 species with at least 50 specimens, and we were not able to take into account their phylogeny.

Conclusions

We for the first time documented temporal abundance fluctuations of noctuid moths using a) Malaise-interception traps, b) in the canopy of a temperate forest, c) in Poland. Although the sample size remained small for comparative analyses, we found that species with longer forewings and with variable adult colourations had less variable abundance across the two study years.

ACKNOWLEDGEMENTS
Fieldwork was carried out with the kind permission of the Forestry Experimental Station in Murowana Goślina (Poznań University of Life Sciences). Traps were generously lent by Martin Gossner and repaired and completed with advice from Edward Baraniak. We thank Jaroslaw Kordy, Öznur Tilkioğlu, Fatmanur Selvi, and Meriç Atar for help with setting moths, Pasquale Di Pratola for help with calculating phenology from the raw data, and Mariia Sokulska for measuring the moths.

Funding
The research was funded by grant No. 2018/29/B/NZ8/00112 to Freerk Molleman from the National Science Centre (NCN, Poland) and supported by the Estonian Research Council grants PRG741 and PRG2618.

Conflict of Interests
The authors have no relevant financial or non-financial conflict of interests to disclose.

Data Availability Statement
Data and code used in this manuscript will be made available from the Dryad Digital Repository and are here uploaded for review purposes.

Author contributions
Freerk Molleman developed the initial concept of the study, participated in the fieldwork, set part of the moths, analysed the data, and wrote the first draft of the manuscript. Roman Wasala identified the moths and compiled their diet information. Ahmet Tambay entered the data and contributed to the data analysis. Robert B. Davis performed comparative analyses and Erki Õunap provided the phylogeny. Urszula Walczak participated in developing the concept, planning of the study, and in fieldwork. All authors contributed to the manuscript writing.

Literature
ALTERMATT, F. 2010. Tell me what you eat and I'll tell you when you fly: diet can predict phenological changes in response to climate change. *Ecology Letters,* 13**,** 1475-1484.

ANONYMOUS. 2025. *University of Tartu "UT Rocket."* [Online]. Available: https://share.neic.no/#/marketplace-public-offering/c8107e145e0d41f7a016b72825072287/ [Accessed].

ASHTON, L. A., NAKAMURA, A., BASSET, Y., BURWELL, C. J., CAO, M., EASTWOOD, R., ODELL, E., DE OLIVEIRA, E. G., HURLEY, K., KATABUCHI, M., MAUNSELL, S., MCBROOM, J., SCHMIDL, J., SUN, Z., TANG, Y., WHITAKER, T., LAIDLAW, M. J., MCDONALD, W. J. F. & KITCHING, R. L. 2016. Vertical stratification of moths across elevation and latitude. *Journal of Biogeography,* 43**,** 59-69.

BARTON, K. 2022. MuMln: Multi-model inference, version 1.47. 5. CRAN. https://cran. r-proje ct. org/packa ge= MuMln.

BOND, A. B. 2007. The evolution of color polymorphism: crypticity, searching images, and apostatic selection. *Annual Review of Ecology, Evolution, and Systematics,* 38**,** 489-514.

BÖTTGER, D., SINGH, R. P., FRIEDRICH, E. & BREHM, G. 2025. The moth fauna is more diverse in the understorey than in the canopy in a European forest. *Bulletin of Entomological Research,* 115**,** 1-11.

BUSSE, A., BÄSSLER, C., BRANDL, R., FRIESS, N., HACKER, H., HEIDRICH, L., HILMERS, T., MERKEL-WALLNER, G., SCHMID-EGGER, C. & SEIFERT, L. 2022. Light and Malaise traps tell different stories about the spatial variations in arthropod

biomass and method-specific insect abundance. *Insect Conservation and Diversity,* 15**,** 655-665.

BUTLER, L., KONDO, V., BARROWS, E. M. & TOWNSEND, E. C. 1999. Effects of weather conditions and trap types on sampling for richness and abundance of forest macrolepidoptera. *Environmental Entomology,* 28**,** 795-811.

COULTHARD, E., NORREY, J., SHORTALL, C. & HARRIS, W. E. 2019. Ecological traits predict population changes in moths. *Biological Conservation,* 233**,** 213-219.

DE SMEDT, P., VANGANSBEKE, P., BRACKE, R., SCHAUWVLIEGE, W., WILLEMS, L., MERTENS, J., VERHEYEN, K., DIDHAM, R. & MÜLLER, J. 2019. Vertical stratification of moth communities in a deciduous forest in Belgium. *Insect Conservation & Diversity,* 12.

FIBIGER, M. 1990. *Noctuinae I.,* Sorø, Entomological Doss.

FIBIGER, M. 1993. *Noctuinae II,* Sorø, Entomological Doss.

FIBIGER, M. 1997. *Noctuinae III,* Sorø, Entomological Doss.

FIBIGER, M. & HACKER, H. 2007. *Amphipyrinae – Xyleninae,* Sorø, Entomological Doss.

FIBIGER, M., RONKAY, L., STEINER, A. & ZILLI, A. 2009. *Pantheinae – Bryophilinae,* Sorø, Entomological Doss.

FIBIGER, M., RONKAY, L., YELA, J. L. & ZILLI, A. 2010. *Rivuliinae – Euteliinae and Micronoctuidae and supplement to volumes 1-11,* Sorø, Entomological Doss.

FOERSTER, S. I. A., CLARKE, J. T., ÕUNAP, E., TEDER, T. & TAMMARU, T. 2024. A comparative study of body size evolution in moths: evidence of correlated evolution with feeding and phenology-related traits. *Journal of Evolutionary Biology***,** voae072.

FOREST_DATA_BANK. 2018. *Forests on maps* [Online]. Available: https://www.bdl.lasy.gov.pl/portal/mapy-en [Accessed 11/01/2018].

FORSMAN, A., BETZHOLTZ, P. E. & FRANZEN, M. 2015. Variable coloration is associated with dampened population fluctuations in noctuid moths. *Proc Biol Sci,* 282**,** 20142922.

FOX, R., OLIVER, T. H., HARROWER, C., PARSONS, M. S., THOMAS, C. D. & ROY, D. B. 2014. Long-term changes to the frequency of occurrence of British moths are consistent with opposing and synergistic effects of climate and land-use changes. *Journal of Applied Ecology,* 51**,** 949-957.

GASTON, K. J. 1988. Patterns in the local and regional dynamics of moth populations. *Oikos***,** 49-57.

GASTON, K. J. & LAWTON, J. H. 1988. Patterns in body size, population dynamics, and regional distribution of bracken herbivores. *The American Naturalist,* 132**,** 662-680.

GASTON, K. J. & MCARDLE, B. H. 1994. The temporal variability of animal abundances: measures, methods and patterns. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences,* 345**,** 335-358.

GASTON, K. J. & REAVEY, D. 2009. Patterns in the life histories and feeding strategies of British macrolepidoptera. *Biological Journal of the Linnean Society,* 37**,** 367-381.

GOSSNER, M. M., WEISSER, W. W., GERSHENZON, J. & UNSICKER, S. B. 2014. Insect attraction to herbivore-induced beech volatiles under different forest management regimes. *Oecologia,* 176**,** 569-80.

HÄLLFORS, M. H., PÖYRY, J., HELIÖLÄ, J., KOHONEN, I., KUUSSAARI, M., LEINONEN, R., SCHMUCKI, R., SIHVONEN, P. & SAASTAMOINEN, M. 2021. Combining range and phenology shifts offers a winning strategy for boreal Lepidoptera. *Ecology Letters,* 24**,** 1619-1632.

HICKINBOTHAM, E. J., PATTISON, Z., FOX, R. & RUSHTON, S. P. 2024. Drivers of moth phenology in England and Wales. *Journal of Insect Conservation***,** 1-11.

HOLM, S., DAVIS, R., JAVOIŠ, J., ÕUNAP, E., KAASIK, A., MOLLEMAN, F. & TAMMARU, T. 2016. A comparative perspective on longevity: the effect of body size dominates over ecology in moths. *Journal of evolutionary biology,* 29**,** 2422-2435.

HUNTER, A. F. & ELKINTON, J. S. 2000. Effects of synchrony with host plant on populations of a spring-feeding lepidopteran. *Ecology,* 81**,** 1248-1261.

Temporal abundance and species traits

JONKO, K. 2020. *Lepidoptera Mundi* [Online]. Available: https://lepidoptera.eu/start/pl [Accessed 01/10/2024].

KALYAANAMOORTHY, S., MINH, B. Q., WONG, T. K., VON HAESELER, A. & JERMIIN, L. S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods,* 14**,** 587-589.

KARPESTAM, E., MERILAITA, S. & FORSMAN, A. 2016. Colour polymorphism protects prey individuals and populations against predation. *Scientific Reports,* 6**,** 22122.

KAWAHARA, A. Y., PLOTKIN, D., ESPELAND, M., MEUSEMANN, K., TOUSSAINT, E. F., DONATH, A., GIMNICH, F., FRANDSEN, P. B., ZWICK, A. & DOS REIS, M. 2019. Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths. *Proceedings of the National Academy of Sciences,* 116**,** 22657-22663.

LA CAVA, S., RIJLLO, G., ZUCCO, G., INNOCENTI, S., GUASTI, M., PULETTI, N., FERRARA, C. & SCALERCIO, S. 2024. Moths in space: The below-canopy structure of forest drives abundance and mobility-related traits of moth communities. *Science of The Total Environment,* 921**,** 171056.

LEPIFORUM, E. V. 2020. *Bestimmungshilfe für die in Europa nachgewiesenen Schmetterlingsarten* [Online]. Available: https://www.lepiforum.org [Accessed 1/10/2024].

LOGGHE, G., TAELMAN, C., VAN HECKE, F., BATSLEER, F., MAES, D. & BONTE, D. 2024. Unravelling arthropod movement in natural landscapes: Small-scale effects of body size and weather conditions. *Journal of Animal Ecology*.

MACARTHUR, R. 1955. Fluctuations of animal populations and a measure of community stability. *Ecology,* 36**,** 533-536.

MAURER, J. A., SHEPARD, J. H., CRABO, L. G., HAMMOND, P. C., ZACK, R. S. & PETERSON, M. A. 2018. Phenological responses of 215 moth species to interannual climate variation in the Pacific Northwest from 1895 through 2013. *PLoS One,* 13**,** e0202850.

MILLER, M. A., PFEIFFER, W. & SCHWARTZ, T. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 gateway computing environments workshop (GCE), 2010. Ieee, 1-8.

MOLLEMAN, F. 2018. Moving beyond phenology: new directions in the study of temporal dynamics of tropical insect communities. *Current Science,* 114.

MOLLEMAN, F. & WALCZAK, U. 2024. Winter moth populations are isolated on co-occurring tree species with contrasting budburst-phenology. *Ecological Entomology,* 49**,** 744-747.

MOLLEMAN, F., WALCZAK, U., MELOSIK, I., BARANIAK, E., PIOSIK, Ł. & PRINZING, A. 2022. What drives caterpillar guilds on a tree: enemy pressure, leaf or tree growth, genetic traits, or phylogenetic neighbourhood? *Insects,* 13**,** 367.

NGUYEN, L.-T., SCHMIDT, H. A., VON HAESELER, A. & MINH, B. Q. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular biology and evolution,* 32**,** 268-274.

NIEMINEN, M., RITA, H. & UUVANA, P. 1999. Body size and migration rate in moths. *Ecography,* 22**,** 697-707.

O'NEILL, B. F., BOND, K., TYNER, A., SHEPPARD, R., BRYANT, T., CHAPMAN, J., BELL, J. & DONNELLY, A. 2012. Climatic change is advancing the phenology of moth species in Ireland. *Entomologia Experimentalis et Applicata,* 143**,** 74-88.

PARADIS, E. & SCHLIEP, K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics,* 35**,** 526-528.

PRICE, P. W. 2003. *Macroevolutionary theory on macroecological patterns,* Cambridge, Cambridge University Press.

R_CORE_TEAM 2024. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

REVELL, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in ecology and evolution***,** 217-223.

RONKAY, G. & RONKAY, L. 1994. *Cuculliinae I,* Sorø, Entomological Doss.

RONKAY, G. & RONKAY, L. 1995. *Cuculliinae II,* Sorø, Entomological Doss.

RONKAY, G., RONKAY, L. & GYULAI, P. 2011. *Cuculliinae II and Psaphidinae,* Budapest, Heterocera Press.

RONKAY, L., YELA, J. L. & HREBLAY, M. 2001. *Hadeninae II,* Sorø, Entomological Doss.

RUCHIN, A. Vertical stratification and dynamics of insect communities in deciduous forests (Center of European Russia). E3S Web of Conferences, 2023. EDP Sciences, 07021.

SEIFERT, C. L., STRUTZENBERGER, P. & FIEDLER, K. 2023. How do host plant use and seasonal life cycle relate to insect body size: A case study on European geometrid moths (Lepidoptera: Geometridae). *Journal of Evolutionary Biology,* 36**,** 743-752.

SPITZER, K. & JAROŠ, J. 2009. Long-term monitoring of moth populations (Lepidoptera) associated with a natural wetland forest: synthesis after 25 years. *Terrestrial Arthropod Reviews,* 1**,** 155-163.

SUCHARD, M. A., LEMEY, P., BAELE, G., AYRES, D. L., DRUMMOND, A. J. & RAMBAUT, A. 2018. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus evolution,* 4**,** vey016.

TAMMARU, T. & HAUKIOJA, E. 1996. Capital breeders and income breeders among Lepidoptera: consequences to population dynamics. *Oikos,* 77**,** 561-564.

TEDER, T. 2020. Phenological responses to climate warming in temperate moths and butterflies: species traits predict future changes in voltinism. *Oikos,* 129**,** 1051-1060.

VALTONEN, A., AYRES, M. P., ROININEN, H., PÖYRY, J. & LEINONEN, R. 2011. Environmental controls on the phenology of moths: predicting plasticity and constraint under climate change. *Oecologia,* 165**,** 237-248.

VARLEY, G. & GRADWELL, G. 1967. Population models for the winter moth.

VÉGVÁRI, Z., JUHÁSZ, E., TÓTH, J. P., BARTA, Z., BOLDOGH, S., SZABÓ, S. & VARGA, Z. 2015. Life-history traits and climatic responsiveness in noctuid moths. *Oikos,* 124**,** 235-242.

WAGNER, D. L., FOX, R., SALCIDO, D. M. & DYER, L. A. 2021. A window to the world of global insect declines: Moth biodiversity trends are complex and heterogeneous. *Proc Natl Acad Sci U S A,* 118.

WAGNER, W. 2024. *Lepidoptera and their ecology* [Online]. Available: www.pyrgus.de [Accessed 01/10/2024].

WEI, T. & SIMKO, V. 2021. R package 'corrplot': Visualization of a Correlation Matrix (Version 0.92).

WESOŁOWSKI, T. & ROWIŃSKI, P. 2006. Timing of bud burst and tree-leaf development in a multispecies temperate forest. *Forest Ecology and Management,* 237**,** 387-393.

Phenology of the most abundant noctuid moths (N > 50) during 2019 (left) and 2020 (right) sampled with Malaise-interception traps in the canopy of a forest in Western Poland.

708x2531mm (79 x 79 DPI)