Title

A highly resolved integrated transcriptomic atlas of human breast cancers

Authors

Andrew Chen[1,2], Lina Kroehling[1,2], Christina S. Ennis[3], Gerald V. Denis[3,4,5], Stefano Monti[1,2,6]

Affiliations

1. Section of Computational Biomedicine, Boston University Chobanian and Avesidian School of Medicine, Boston, MA 02118, USA.
2. Bioinformatics Program, Center for Computing & Data Sciences, Boston University, Boston, MA 022158, USA.
3. Boston University-Boston Medical Center Cancer Center, Boston University Chobanian and Avesidian School of Medicine, Boston, MA, 02118, USA
4. Section of Hematology and Medical Oncology, Department of Medicine, Boston University Chobanian and Avesidian School of Medicine and Boston Medical Center, Boston, MA, 02118, USA
5. Department of Pharmacology and Experimental Therapeutics, Boston University Chobanian and Avesidian School of Medicine, Boston, MA, 02118, USA
6. Department of Biostatistics, Boston University School of Public Health, Boston, MA 02118, USA.
corresponding author(s): Stefano Monti (smonti@bu.edu)

Abstract

In this study, we developed an integrated single cell transcriptomic (scRNAseq) atlas of human breast cancer (BC), the largest resource of its kind, totaling > 600,000 cells across 138 patients. Rigorous integration and annotation of publicly available scRNAseq data enabled a highly resolved characterization of epithelial, immune, and stromal heterogeneity within the tumor microenvironment (TME). Within the immune compartment we were able to characterize heterogeneity of CD4, CD8 T cells and macrophage subpopulations. Within the stromal compartment, subpopulations of endothelial cells (ECs) and cancer associated fibroblasts (CAFs) were resolved. Within the cancer epithelial compartment, we characterized the functional heterogeneity of cells across the axes of stemness, epithelial-mesenchymal plasticity, and canonical cancer pathways. Across all subpopulations observed in the TME, we performed a multi-resolution survival analysis to identify epithelial cell states and immune cell types which conferred a survival advantage in both The Cancer Genome Atlas (TCGA) and METABRIC. We also identified robust associations between TME composition and clinical phenotypes such as tumor subtype and grade that were not discernible when the analysis was limited to individual datasets, highlighting the need for atlas-based analyses. This atlas represents a valuable resource for further high-resolution analyses of TME heterogeneity within BC.

Introduction

Breast cancer (BC) is the most prevalent cancer and the second most common cause of cancer
death in women [1]. It is defined as a malignancy of the epithelial duct in breast tissue, and is a
highly heterogeneous disease [2,3] for which clinical outcomes and responsiveness to treatment
depend greatly on intrinsic factors related to the cancer cell, as well as extrinsic factors related
to altered states of immune and stromal cells in the tumor microenvironment (TME) [4–6]. To
improve our understanding of breast cancer progression and treatment, an extensive
characterization of the heterogeneity in BC cancer cells and the surrounding tumor
microenvironment is required.
53
Single cell transcriptomics (scRNAseq) remains the method of choice to characterize
heterogeneity in tissue composition in terms of cell *types* – categories of cells that perform
consistent functions, and in terms of cell *states* – categories of transient functions that can be
shared across different cell types [7,8]. The technology has been extensively applied to profile
healthy breast tissue, as part of the larger human cell atlas effort. There are now highly-
resolved scRNAseq atlases that have annotated 700,000-800,000 cells across 55-126 samples
yielding new insights into the diversity of immune, stromal, and epithelial populations in
healthy breast tissue and how these are linked to clinical metadata [9,10]. However, the
application of scRNAseq to profile breast cancer (BC) samples has been limited by cohort sizes,
with most studies having fewer than 30 samples [11–20]. While these individual studies have laid
substantial groundwork in characterizing tumor heterogeneity within BC, their limited sample
size poses a challenge with reproducibility and identifying statistically significant associations
between tissue composition and clinical metadata.
67
To address this challenge, we integrated eight publicly available scRNAseq datasets of
untreated and unsorted BC samples [11–20] to create an atlas of transcriptional heterogeneity in
BC. The analysis of this combined atlas enabled 1) the reconciliation of mislabeled cell types,
2) the identification of novel cell types, and 3) the association of changes in TME heterogeneity
with clinical phenotypes such as tumor subtype and grade with increased statistical power (Fig
1a). Another study has attempted to integrate publicly-available scRNAseq BC data to create
an atlas but compared to Xu et al. our study only contains unsorted BC samples, allowing for
an unbiased characterization of inter-patient heterogeneity, and our atlas has a much a larger
sample size (> 600,000 cells), enabling more robust identification of cell types and
differentially abundant populations across phenotypes [21].
78
79 This integrated scRNAseq atlas provides an unbiased reference landscape of transcriptional
80 heterogeneity in BC, comprised of more than 600,000 cells across 138 patients, the largest of
81 its kind, and represents a valuable resource for hypothesis-driven analysis of TME
82 heterogeneity. The data and methods described herein provide a robust framework for
83 updating this BC atlas as new datasets are made publicly available [22].
84

Results

Constructing the integrated core atlas

The BC atlas was constructed by integrating eight publicly available scRNAseq datasets of
untreated primary BC tumors. Across the eight datasets, samples were sequenced at varying
throughput levels, and had different distributions of clinical and patient metadata including
subtype, grade, and age (Fig. 1b,c,d). Before integration, batch effects were evident as cells
separated by dataset of origin, rather than by biological factors (Supp Fig. 1). Six different
integration approaches were applied and the best-performing method, as determined by atlas
integration batch correction metrics [23] was Seurat's reciprocal PCA method (RPCA) (Fig 2a).

Relabeling of broad cell types

To annotate cells in the integrated atlas, we first defined three compartments: epithelial, immune, and stromal, by cross-referencing annotations from SingleR, CellTypist, and author annotations with broad clustering (Louvain clustering with resolution 0.1) (Supp Fig. 2). The resulting compartments showed distinct expression of canonical marker genes (Epithelial: *EPCAM, KRT8, KRT18*; Immune: *PTPRC, CD3D, CD3E, CD68*; Stromal: *COL1A1, COL1A2, ENG*) and diverse representation across all eight source datasets (Fig. 2b,c,d). More than 300,000 cells that previously had no available cell type annotation were annotated using the cluster-based annotations from the atlas (Fig. 2e).

These annotations of broad cell types based on integrated data also resulted in a relabeling of broad cell types for over 9172 cells (around 1% of the atlas) across 5 datasets (e.g., relabeling a cell as stromal that was originally labeled immune).

Cancer Epithelial Diversity

Intra-tumor heterogeneity within cancer cells stems from multiple factors including but not limited to genomic instability, epigenetic alterations, and clonal evolution which are all implicated in the treatment of cancer [24]. In this BC atlas, we sought to characterize intra and inter-tumor transcriptional heterogeneity by scoring each cancer cell with respect to potency/stemness [25], epithelial-mesenchymal plasticity (EMP) [26], malignancy [27], 'activity' of hallmark pathways [28], expression of epithelial cell type markers from HBCA [10], as well as clinical metadata related to age, tumor grade, and subtype (Fig. 3a). Further, we sought to find the driving factors of variation by leveraging a recursive partitioning tree-based method *K2 Taxonomer* [29] to identify meta-clusters and hierarchical relationships between them (Fig. 3b), and also trained random forests to extract the most discriminative features distinguishing each cluster and meta-cluster (Fig. 3c).

To identify functional groups of cancer cells shared across patients in the atlas, we performed unsupervised clustering (Louvain clustering with resolution 0.1) of cancer epithelial cells and identified 23 clusters that largely stratified by cell typist annotation based on the healthy Human Breast Cell Atlas (HBCA), author subtype, and Pam50 inferred subtype (Supp Fig. 3a,b,c,d). The differentially expressed genes for each epithelial cluster were identified using MAST (Supp Table 1). Several resolution parameters were evaluated (0.1-1.0) with even the coarsest resolution of 0.1 yielding multiple clusters (7,9,10,11,12,19,21) mostly consisting of cells from individual patients, as also revealed by their lowest entropy of donor proportions (Supp Fig. 3e,f). This suggests that the variation captured within these clusters was mostly driven by tumor-specific clonal expansion and, given our goal of defining shared programs of cancer epithelial heterogeneity, they were excluded from further downstream characterization.

<u>Multi-resolution characterization of cancer epithelial cells</u>
Among the remaining 16 epithelial clusters we identified the driving factors of variation within our cancer epithelial compartment by training random forest based classifiers at three separate levels of taxonomic resolution (Fig. 3b,c). In previous integrated single cell studies of breast cancer, cancer epithelial heterogeneity was characterized with respect to pam50 and clinical subtypes[11,12], as well as functional annotations by, e.g., Hallmark signatures [21], but has not yet been done in a hierarchical manner to identify the driving factors of variation.

This multi-level characterization revealed segregation of cancer epithelial cells at the highest level (Level 1) by stemness, EMP, and basal phenotypes, with clusters 13 through 4 (13~4) on the right side of the dendrogram exhibiting more stemness, more EMP (more mesenchymal),

and being more basal-like compared to clusters on the left side (clusters 0~16). At the next
level of the dendrogram (Level 2), cancer cells on both sides segregated by age, with the left
branch of the tree (clusters 0~16) segregating by older donors (clusters 0~20, 8, and 15~16)
and younger donors (clusters 1~5), and the right branch of the tree (clusters 13~4) segregating
by older (cluster 4) and younger (clusters 13~2) patients. At the next level of characterization
(Level 3), finer groups of cancer epithelial heterogeneity emerge with clusters exhibiting
decreased mitotic spindle activity (cluster 0), decreased TNFA signaling (cluster 20), increased
EMT activity (cluster 13), as well as subtype-specific clustering (Luminal A: 1~17, 18; Luminal
B: 6, 5, 15, 16; HER2: 3, 14), and grade specific clustering (high grade tumors: 8; low grade
tumors: 6, 5, 18).

Immune and Stromal Cell Diversity in the TME

156   Across the immune and stromal compartments of the atlas, 20 immune and 14 stromal
157   subpopulations were characterized (Fig. 4a,5a). These subpopulations were identified by
158   integrating data from automatic annotation methods: SingleR [30] and CellTypist [31] (Supp Fig.
159   4,5), hierarchical tree-based methods (K2 Taxonomer) [29] (Fig. 4b, 5b), as well as marker genes
160   obtained from differential expression analysis (MAST) [32] (Supp Table 2,3).
161
Reconciliation of immune cell subtypes to pan-cancer subtypes
Macrophages formed three clusters, which were all found to have analogs in previous pan-
cancer and breast cancer specific characterizations of myeloid cells [10,33]. These included a
cluster of lipid-associated macrophages *Mac. Lipo* defined by high expression of genes
associated with lipid transport and metabolism (*LIPA, LPL, APOE, APOC1*); a cluster
characterized by high expression of *MIK67* and *TOP2A* indicating a more proliferative state
(*Mac Prolif)*; and a cluster of macrophages *Mac. Col* defined by strong expression of collagen
genes (*COL1A1, COL1A2, COL3A1*), a subtype that has been previously been connected to ECM
remodeling [34] (Fig. 4a,b,c).
171
T cells formed nine clusters (Fig. 4a), which were also found to have analogs in previous pan-
cancer studies [13,35,36]. These included CD4 and CD8 effector memory T cells (*CD4/CD8 Tem*),
which are characterized by expression of *LMNA, FOS,* and *GZMK, EOMES* respectively; CD4
Tregs defined by expression of *FOXP3, IL2RA,* and *TNFRSF4*; an interferon stimulated group of
CD8 T cells (*CD8 ISG*) defined by strong expression of interferon signaling genes *IFIT1, IFIT2,*
IFIT3; naïve CD4 cells characterized by expression of *CCR7, PASK, LEF1*; exhausted CD4 T cells
(*CD4 Ex*) defined by expression of canonical exhaustion markers such as *LAG3, HAVCR2, PDCD1*
and *TIGIT*; CD4 T follicular helper cells (*CD4 Tfh*) expressing transcription factors *TOX, TOX2*
and chemokine *CXCL13*; CD4 T helper cells (*CD4 Th*) which in comparison to *CD4 Tfh* cells
expressed high levels of interleukin genes *IL7R*, as well as the chemokine receptor *CCR6*.
Similar to the macrophages, a proliferative T cell cluster (*T. Prolif*) was identified exhibiting
strong expression of *MIK67* and *STMN1* (Fig. 4,a,b,d).
184
NK cells formed one cluster (Fig. 4a,d), which was defined by expression of Killer Cell Lectin
receptors (*KLRD1, KLRF1*) and growth factor *FGFBP2*. A previous single cell breast cancer atlas
identified 6 NK subtypes, although their atlas only contained 3720 NK cells across all studies,
whereas our study identified one cluster with no discernable functional differences across a
population of 23508 NK cells. This discrepancy may be attributed to the difference in sample
size and clustering strategies between the two atlases [37].
191
192   Mast cells formed one cluster (Fig. 4a,c), defined by expression of the enzymes *TPSB2, CPA3,*
193   and *TPSAB1*.
194

195 B Cells also formed a single cluster defined by canonical B cell markers such as *CD19*, *MS4A1*,
196 *CD74* as well as expression of human leukocyte antigen genes such as *HLA-DRA* and *HLA-DPB1*
197 (Fig. 4a,d).
198
Identification of dendritic subtypes
Dendritic cells formed four clusters (Fig. 4a,c), which include two conventional dendritic cell
populations, one characterized by expression of *CLEC9A* (*cDC1*), and another characterized by
expression of *CD1C* and *CLEC10A* (*cDC2*); a mature dendritic cell population (*mDC*) defined by
CCL22, *CCR7*, and *CCL19* expression; and a plasmacytoid dendritic cell population (*pDC*)
defined by expression of *GZMB*, *JCHAIN*, and *PTGDS*.
205
Reconciliation of stromal cell subtypes to pan-cancer subtypes
Fibroblasts formed six clusters (Fig. 5a,b,c), which were also reconciled to previously
characterized pan-cancer stromal subtypes [38]. This includes five clusters of cancer-associated
fibroblasts (CAFs) consisting of a population of *COL11A1+ CAFs* (*COL11A1*, *COL8A2*) implicated
in collagen metabolism [38], a population of *LAMP5 CAFs* defined by expression of *LAMP5* and
cystatins *CST1* and *CST2* implicated in promoting EMT [38], a population of PI16+ CAFs defined
by expression of *PI16* and *PCOLCE2*, a population of DPT+ CAFs defined by expression of *DPT*
and *CAPN6*, as well as population of CA12+ CAFs defined by expression of *CA12* and *SLC2A1*
which has been implicated in glycolysis metabolism and hypoxia [38]. Apart from these CAFs
subtypes, myofibroblasts also formed one cluster by expression of *RGS5* and *CDH6*.
216
Endothelial cells (ECs) surround blood vessels in the TME which are responsible for
angiogenesis and tumor growth and formed a total of five clusters in the atlas (Fig. 6a). Four
clusters emerged based on the function of these endothelial cells in supporting vasculature:
endothelial vein, arterial, capillary, and lymphatic cells segregated based on expression of
different marker genes (Fig 5a,b,c). A further cluster of immature endothelial cells (*Endo Imm*),
defined by expression of *PLVAP, VWA1*, and *CA4*, was identified and has been implicated in
poor tumor prognosis [38].
224
225 Other stromal subtypes captured in the atlas include vascular smooth muscle cells (*VSMC*)
226 defined by expression of *RERGL*, and *MYH11*, extracellular matrix related pericytes (*ECM PCs*)
227 defined by expression of collagen genes *COL4A1*, and *COL4A2*, as well as a set of proliferating
228 stromal cells (*Prolif. Stromal)* characterized by high expression of proliferative markers
229 including *TOP2A, MKI67*.
230
231 **Taxonomic characterization of immune and stromal subtypes**
232 We utilized K2Taxonomer to organize the clusters in the immune and stromal compartments
233 into hierarchical taxonomies. Annotating groups of clusters, or clades, mitigates the
234 uncertainty in annotation due to over-clustering. Taxonomic characterization also captures
235 groups of cells sharing similar functions and cell states rather than just cell types.
236
In the immune compartment, cells largely segregated by lymphoid and myeloid lineages. The
exceptions being a population of *pDCs*, which clustered with plasma cells in a clade due to
their shared expression of *MZB1* [39], and Mast cells, which clustered within the lymphoid clade
due to the lack of expression of macrophage and dendritic markers (*CD68*, *CD1C*). These
exceptions suggest that immune cells in the TME may share similar transcriptional states
despite having different lineages (Fig. 4b).
243
244 In the stromal compartment, cells partitioned by endothelial and non-endothelial at the first
245 level, and within the non-endothelial clade, further separated between CAFs and mural cells,

246 and lastly between smooth muscle cells, pericytes, and myofibroblasts, indicating strong
247 transcriptome-based partitioning of these stromal subtypes (Fig. 5b).
248

Cell type diversity of tumors changes across phenotypes

250 Using these improved annotations within the atlas, we used an entropy-based cell type
251 diversity score (CTDS) [40] to assess how the overall diversity of the TME changes across patient
252 phenotypes (Fig. 6a). The CTDS score yields the highest value when all cell types (or states)
253 have equal representation in a sample, and the lowest value when a single cell type (or state)
254 is represented in a sample.
255

Across clinical subtypes, the CTDS of observed epithelial cell states and stromal cell types
observed was lowest in TNBC compared to ER+ and HER2+, whereas the average CTDS of
immune cell types was highest in TNBC. Across age groups, we observed a consistent decrease
in CTDS across all three compartments in older tumors compared to younger tumors. This
expands upon previous studies that identified decreased immune cell type diversity in old
subjects compared to young subjects, but only within PBMCs [40]. As tumor grade increased, we
also observed a consistent decrease in cell type diversity within the epithelial and stromal
compartment. The cell type diversity scores provide a global summary of changes in tumor
heterogeneity that can complement the more granular differential abundance analysis of
individual cell types within the tumor.

Improved estimates of differential immune cell type abundance across phenotypes

268 Using these improved annotations within the atlas, we used *sccomp* [41] to estimate differential
269 immune cell type abundance across phenotypes such as tumor grade and subtype (Methods).
270

271 HER2+ tumors are typically considered to be 'hot' tumors with increased immune infiltration
272 compared to other subtypes of tumors [42], but the details of their cell type composition vis-à-
273 vis other tumor subtypes is unclear. In our analysis, we observed that one specific subset of T
274 cells, *CD4 Tregs*, increased in abundance in HER2+ tumors relative to other tumor types,
275 whereas one subset of macrophages, *Mac Col.*, decreased in relative abundance (Fig. 6b).
276

277 The immune microenvironment of ER+ tumors have been characterized as macrophage-
278 driven, but the exact subtypes of macrophages driving this microenvironment remain
279 unknown [43]. In our analysis, we observed that one subset of macrophages, *Mac Col.*, was
280 enriched in ER+ tumors compared to other tumors (Fig. 6b).
281

TNBC tumors are known to have a heterogeneous immune microenvironment with tumors
exhibiting both increased tumor infiltrating lymphocytes (TILS) [44], and increased tumor
associated macrophages [45]. However, the exact nature of which immune cell types are
enriched has not yet been characterized in studies with adequate statistical power. In our
analysis we found that *Mac Lipo.* were enriched in TNBC tumors, as well as a subpopulation of
CLEC10A conventional dendritic cells (*cDC2*). Previous studies have found that pDCs are
expanded [46] and that mast cells are be depleted in TNBC tumors [47]. However, although these
specific populations were also found to be expanded and depleted, respectively, in the atlas,
their changes were not statistically significant (fdr > 0.05) (Fig. 6b).
291

Comparison of the atlas-based estimates of differential cell type abundances across tumor
grades with the same estimates obtained on individual datasets further highlighted the utility
of the integrated atlas to uncover unique associations (Fig. 6c). We found that significant

estimates of differential cell type abundance in the atlas were not significant or in discordant
directions when tested in individual datasets. For instance, *CD4 Tem* cells were observed to
increase with grade in two studies (Wang 2024 and Wu Natgen 2021) but after including all
the datasets in the atlas, they did not reach atlas-wide significance. Similarly, *Mast* cells
yielded no significant changes in two of the datasets tested (Pal 2021 and Wang 2024),
although when all datasets were combined in the atlas it was the cell type most significantly
decreasing with grade. Only three datasets were included in this comparison since datasets
need to have both high grade (grade 3) and low-grade tumors (grade 1 & 2) to be included in
this differential abundance analysis, which further highlights the need for integration so that
diverse patient and clinical metadata can be pooled in mega-analysis-based approaches.
305

Multi-resolution cell type associations with patient survival

To assess the clinical relevance of the subpopulations identified in the atlas, we performed
survival analyses by projecting bulk RNA expression data from both TCGA and METABRIC onto
the gene signatures that defined each subgroup within our taxonomic analyses (Fig. 7). To
correct for the confounding effects of patient age, proliferation, and inflammation with
respect to patient survival [29,48], the Cox proportional hazard model also included these factors
as covariates (Methods).
313

Two cell populations were significantly associated with survival in both TCGA and METABRIC
and both had a positive survival association (hazard ratio less than 1). These include a
subpopulation of epithelial cells (Cluster 5) which is characterized by low stemness (more
differentiated) and low grade (Fig 3a,c), and a cluster of *CD8 Tem* immune cells (Cluster 1).
Both CD8 effector memory T cell activity and cytotoxicity as well as low grade phenotype
within tumors have been previously associated with favorable prognosis in BC [29,49,50]. Though
no single cell type/state was associated with decreased survival in both datasets separately,
when combining the results by fisher's method, increased proliferative stromal cell (*Strom
Prolif.*) activity was collectively associated with decreased survival.

Discussion

In this study we integrated scRNAseq data from 138 primary BC tumors collected across eight
studies to create the largest transcriptomic atlas of human BC to date. The large number of
patients in our integrated atlas, along with the rigorous annotation of both discrete cell types
and continuous cell states, will provide a valuable resource for researchers seeking to explore
associations of gene expression and or cell type composition with clinical and molecular
phenotypes with increased statistical power.
330

Our characterization of the cancer epithelial compartment of the atlas revealed substantial
heterogeneity between cancer cells along the axes of potency/stemness (differentiation
continuum), EMP (epithelial-mesenchymal continuum), expression of hallmark and HBCA
pathways, as well as clinical phenotypes. By leveraging K2Taxonomer and random forest
models, we performed a multi-resolution characterization of the cancer epithelial
compartment and identified the most discriminative factors differentiating epithelial clusters
at different levels of the taxonomy of BC epithelial cells. At the first level, we found that BC
tumor cells primarily segregated due to EMP, stemness, and basal phenotypes. At the second
level, tumor cells primarily segregated by age, suggesting that aging processes can be a major
driver of tumor heterogeneity [51]. At the third level, tumor cells separate through more granular
hallmark pathways, tumor subtype and grade. A multi-resolution characterization of cancer
cells has several advantages over typical unsupervised clustering approaches including
mitigating the risks of over-clustering and thus over-annotation, as well as being able to
identify the major factors driving heterogeneity in a dataset.

For the immune and stromal compartments, our analysis reconciled cell type information to
established pan-cancer cell types previously characterized, and identified cell states unique to
our BC atlas: a collagen expressing Macrophage subtype (*Mac Col.*) and exhausted CD4 cells
(*CD4 Ex*), a T cell state associated with the progressive loss of cytokine production and effector
function [52].

Using these annotations of immune cell type and state, we performed a cell type diversity and
differential abundance analysis to capture global and more granular changes in cell type
composition within the TME with respect to clinical phenotypes such as tumor subtype (ER+,
HER2+, TNBC). Through the cell type diversity analysis, we uncovered associations between
increasing tumor grade and decreasing global diversity within the TME, as well as age-
associated changes across epithelial, stromal, and immune cells. Through the differential
abundance analysis, we resolved differences in immune cell type composition unique to each
subtype that provide insight into effective cancer therapies. The expansion of *Mac Lipo* in
TNBC is of interest and may help explain recent findings suggesting that TNBC is uniquely
susceptible to treatment with lipid-acting drugs [53]. The expansion of *CD4 Tregs* in HER2+
tumors, may explain the effectiveness of targeted therapies such as Traztuzumab that are
known to modulate *Treg* activity [54], and highlights the need to identify additional therapies
targeting *Treg* activity given that *CD4 Tregs* are known to promote an immunosuppressive TME
that can lead to tumor escape [55]. The enrichment of conventional dendritic cells *cDC* and *Mac*
Col. in TNBC and ER+, respectively, also represent promising targets for future therapies given
their known role in antigen presentation to activate anti-tumor T cells and in remodeling the
extracellular matrix, respectively [56,57]. Collectively, these subtype specific differences in the
immune TME may help explain the effectiveness of existing cancer therapies and prioritize
future targets. Further, by repeating this analysis in subsets of the atlas limited to the original
datasets, we found that estimates of differential cell type abundance vary considerably
between datasets, highlighting the need to pool data to arrive at more robust estimates and
to increase the statistical significance of associations found between cell type composition and
clinical phenotypes.

Finally, our atlas-wide survival analysis of cell type subpopulations based on bulk data from
both TCGA and METABRIC (Supp Table 4) revealed both known associations, such as decreased
mortality risk associated with increased CD8 memory T cell activity and low-grade cancer
epithelial cells, as well as novel associations, such as increased mortality risk associated with
increased proliferative stromal cells (*Strom Prolif.*), which could be prognostic in BC.

One limitation of this atlas is the absence of adipocytes, despite their known importance in
cancer progression and aggressiveness [58], due to the technical limitations of scRNAseq. Other
technologies such as single-nucleus RNA sequencing, and spatial transcriptomics are needed
to characterize the heterogeneity in adipocytes across patients. Another limitation is that the
atlas represents a static snapshot of the transcriptome and does not capture dynamic changes
in response to treatments like radiation or chemotherapy, which could significantly alter the
cellular composition and gene expression profiles. Future studies incorporating temporal
analyses of cellular responses to various therapies could enhance the atlas' utility in guiding
personalized treatment strategies and understanding mechanisms of treatment resistance.

In summary, this integrated scRNAseq BC atlas, the largest of its kind, fills a crucial need in our
understanding of tumor heterogeneity by providing a 'reference' landscape of cell populations
which not only enables statistically powered analyses of cell type diversity, composition, and
survival, but also provides a framework for the construction of further atlases to interrogate
tumor heterogeneity.

Methods

Data Acquisition

To create an unbiased scRNAseq atlas of human BC, we integrated publicly available datasets that contained untreated samples (samples from patients who had not undergone chemotherapy or other treatment before sequencing) and samples that were not sorted prior to sequencing (not biased towards a particular cell type). Given these criteria, we identified eight datasets to include in this atlas, totaling 138 patients and 621,200 cells.

Dataset	# Patients	# Cells	System
Pal et al. 2021	33	224,823	10x 3'
Bassez et al. 2021	31	78,549	10x 5'
Wu et al. 2021	26	99,876	10x 3' v2
Wang et al. 2024	14	48,553	10x 3' v2
Qian et al. 2020	14	40,665	10x 3' v2 & 10x 5' v2
Tietscher et al. 2023	12	97,301	10x 3' v3
Liu et al. 2023	4	27,490	10x 3' v3
Gao et al. 2021	4	3,943	10x 3' v2 & 10x 5' v2
Total	**138**	**621,200**	

Table 1: Summary of Datasets used in Atlas

Data Preprocessing

Each scRNAseq dataset was first pre-processed using Seurat [59] to remove low-quality cells, doublets, and normalize data before integration. We adopted current scRNAseq pre-processing best-practices by using adaptive thresholding based on median absolute deviation, instead of manual cutoffs, to filter out outlier cells based on three criteria: mitochondrial genes expression, number of unique genes detected per cell, and total number of genes detected per cell [60,61]. For studies that only provided pre-filtered data, we retained all cells for downstream analysis. Following filtering, doublet scores were obtained using scDblFinder [62] and expression profiles were normalized by total expression and log-transformed using Seurat's *LogNormalize* function which has been demonstrated to work well in prior benchmarking [63].

Integration of Single Cell Data

Integration of filtered scRNAseq data was performed using various methods: Seurat's V5 reciprocal principal component analysis (RPCA) [59], fastMNN [64], Harmony [65], scVI [66], and scANVI [67]. To facilitate fair comparison between the integration methods, the same number of variable features (5000), and PC dimensions (200) were used for integration when possible. For scVI and scANVI, integration is performed at the level of raw counts instead of embeddings, so only the variable feature selection process was included. To avoid biasing the integration towards individual sources of annotations e.g. author annotations or SingleR/CellTypist annotations, only the batch correction metrics: silhouette score, k-nearest neighbor batch effect test (kBET), and integration local inverse Simpson's index (iLISI), were used to assess integration performance. The best performing integration according to these metrics – Seurat V5's RPCA, was chosen for downstream analysis (Fig 2a).

Inference of PAM50 Subtype

To infer molecular subtypes of each sample using the PAM50 method, we first calculated a 'pseudo-bulk' profile for each tumor using Seurat's *AggregateExpression* function, and then

431   applied *molecular.subtyping* from the *genefu* R package to each pseudo-bulk profile with
432   default parameters [68].

Discrete annotation of cell types

434   Annotation of cell types was performed using a combination of reference based annotation
435   methods: singleR [30], CellTypist [31,69], unsupervised recursive partitioning: K2Taxonomer [29], and
436   unsupervised clustering followed by differential expression analysis: MAST [32].
437
438   Reference-based annotation of cell type
Both singleR and CellTypist require a labelled scRNAseq reference dataset to compare query
cells with. Since all of the samples in this study originate from breast tissue, we leveraged
labeled scRNAseq data from the healthy breast cell atlas (HBCA) as reference [10]. For singleR,
the wilcox method *de.method = "Wilcox"* was used as this is more appropriate than the default
methods when applying singleR to single cell data. For CellTypist, we used a pre-trained breast
tissue model that used labeled data from HBCA [69], and used majority voting to determine the
final cell type label *majority_voting = True, mode = 'best match'*.
446
447   Unsupervised clustering-based annotation
Both K2Taxonomer and MAST-based annotation rely on unsupervised clustering which was
performed using the Louvain algorithm for three subsets of the atlas separately: epithelial,
immune, and stromal, based on expression of canonical markers (Epithelial: *EPCAM, KRT8, KRT
18*; Immune*: PTPRC, CD3D, CD3E, CD68*; Stromal: *COL1A1, COL1A2, ENG*). Performing
clustering and annotation within each compartment separately enables the detailed
characterization of heterogeneity within each compartment.
454
455   To identify differentially expressed genes (DEGs) between clusters, we used Seurat's
456   *FindAllMarkers* with the parameters *test.use = 'MAST'*, and *latent.vars = 'batch'* to account for
457   the negative-binomial distribution of single cell data and to control for dataset-specific effects
458   respectively.
459
Whilst methods like MAST can quantify differences at the cluster level, the number of clusters,
and the resulting DEGs obtained from downstream analysis, relies heavily on arbitrary values
of the 'resolution' parameter in typical unsupervised clustering algorithms. To mitigate these
limitations, we leveraged K2Taxonomer to learn hierarchical relationships between the
clusters identified in each compartment, enabling the annotation of clades of clusters instead
of just individual clusters. To accomplish this, K2Taxonomer was applied to the integrated
embeddings obtained from RPCA, with the following parameters (*featMetric="F", nBoots=400,
clustFunc=cKmeansDownsampleSqrt*) which are recommended for large scRNAseq datasets [29].
To estimate the differentially expressed genes (DEGs) between nodes in the dendrogram, we
used the normalized gene expression matrix and K2Taxonomer's built in function *runDGEmods*
with batch variable as a covariate to control for dataset-specific effects.
471
472   Random Forest-based annotation of discriminative factors
The *randomForest* package was used to train random forests to classify cluster membership,
or meta-cluster membership (as determined by K2 Taxonomer), of single cell profiles based
on expression of continuous scores related to cell stemness, EMP, HBCA cell type, malignancy,
and clinical metadata. The model was run with the parameter *importance = TRUE* to extract
the most important features (as determined by gini-based importances), that differentiated
epithelial clusters from each other.

Continuous annotation of cell states

Classifying discrete cell types is just one facet of heterogeneity, and there are many transient functional programs or cell states that require a more continuous scoring [70]. To characterize heterogeneity of functional states within the cancer epithelial compartment of our BC atlas we scored each cell in terms of its malignancy with inferCNV [27], stemness with Cytotrace2 [25], Epithelial-Mesenchymal plasticity [26], expression of hallmark pathways [28,71], as well as expression of epithelial subtypes identified in HBCA [10].

Inference of copy number variations

InferCNV was used to estimate copy number alterations for tumor epithelial cells from each dataset independently with the following parameters: *cutoff = 0.1*, *cluster_by_groups = TRUE*, *denoise = TRUE* and *HMM = FALSE* that are recommended for 10X scRNAseq data. Immune and stromal cells from each dataset were used as normal references, and from these normal reference sets, 100 immune and stromal cells were sampled to act as negative controls to validate inferCNV output.

For each cell, inferCNV estimates a score for each gene that is above one if there is inferred copy number gain, and below one if there is inferred copy number loss. We summarized these scores to estimate a malignancy score per cell that represents the mean absolute CNV score.

$$Malignancy\ Score = \frac{\sum |CNV\ score|}{\#Genes}$$

To classify each cell as malignant or non-malignant, for each dataset we found the 90% interval of malignancy scores of the negative control cells (non-epithelial cells) and labeled all epithelial cells outside of this interval as malignant (Supp Fig. 6).

Scoring of gene set activity

To characterize the heterogeneity of functional states present in the atlas, we leveraged publicly available transcriptional signatures (Supp Table 4) and used a rank-based method (AUCell) [72], with default parameters, to score enrichment of signatures in each cell. For the immune compartment, we scored cells using signatures from pan-cancer studies of myeloid and lymphoid cell states [33,35], as well as signatures of immune cell types from HBCA [10]. For the stromal compartment, we scored cells using signatures from a pan-cancer study of stromal cell heterogeneity [38]. For the epithelial compartment, we scored cells using canonical hallmark signatures [28], as well as signatures of epithelial cell types from HBCA [10].

Differential Diversity and Abundance Analysis

Cell Type Diversity Analysis

We used an entropy-based cell type diversity metric (CTDS) [40] to assess how the overall diversity in a patient's TME changes across subtype, age, and tumor grade. The CTDS metric accounts for the number of cells per sample and the compositional nature of proportional data which enables comparison across samples.

Differential Abundance Analysis

To test differences in cell type proportions across different phenotype groups e.g. tumor grade and subtype, we used the R package *sccomp* with *bimodal_mean_variability_association = TRUE* which is recommended for scRNAseq data [41]. To isolate the effects of tumor grade on TME composition, we controlled for tumor subtype, age, and batch with the following formula:

$$celltype_{count} \sim grade + pam50 + age + (1|batch)$$

Similarly, to isolate the effects of tumor subtype on TME composition, we controlled for grade, age, and batch.

$$celltype_{count} \sim (LumA/LumB/Her2/Normal\_like) + grade + age + (1|batch)$$

Multi-resolution survival analysis of cell type populations

To identify the clinical relevance of cell type annotations identified in the atlas, we extracted differentially expressed genes defining each subpopulation found in our taxonomic analysis of each compartment (K2Taxonomer) [29], and used the R package *brcasurv* [73] to model the association between the expression of subpopulation specific genes and patient survival in both TCGA [3] and METABRIC [74].

To address possible confounders in the survival analysis, we included patients age, as well as patient-level gene set projection scores of inflammation and proliferation that have previously been associated with poor prognosis and demonstrated to be necessary for correction in survival modelling [29,48,75]. The patient-level gene set scores were obtained by using the R package *GSVA* [76] to project inflammation [75], proliferation [48], as well as the cell type specific gene sets onto transcriptional profiles from TCGA and METABRIC. The Cox proportional hazards model underlying the survival analysis controlled for these confounders with the following formula:

$$Surv(patient) \sim celltype_{gsva} + age + inflam_{gsva} + prolif_{gsva}$$

To control for multiple testing, we corrected p-values in each cox model using the FDR method [77] in the *p.adjust* R package, and to reconcile p-values between analyses performed in TCGA and METABRIC, we used the Fisher's method for combining p-values [78].

Code Availability

The atlas is deposited at cellxgene at: https://cellxgene.cziscience.com/collections/9432ae97-4803-4b9f-8f64-2b41e42ad3cb.

All code is available at: https://github.com/montilab/brca_atlas

Author contributions

A.C. performed all analyses and wrote the manuscript.
L.K., and S.M. provided computational guidance and feedback.
C.S.E., L.K., and G.V.D, provided biological guidance.
S.M. oversaw the project.
The manuscript was edited by all authors.

Acknowledgments

We thank Gat Rauner, and Yuhan Qiu for feedback regarding the construction and annotation of the atlas. We also thank Corinn Small for providing thorough feedback on making the data accessible on cellxgene. This work was supported by grants from the National Institute of Health (NIH): U01CA243004 (G.V.D. and S.M.), the National Institute of General Medical Sciences of the NIH under award number T32GM100842, and by a gift from Find The Cause Breast Cancer Foundation [79]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Competing interests

There are no conflicts to be declared.

Figure 1

Figure 1. Construction and composition of the integrated breast cancer atlas. A) Overview of the construction of the integrated atlas and downstream analysis. B) Distribution of patients and number of cells per patient sequenced across each of the eight datasets. C) Distribution of tumor subtype and grade information. D) Distribution of ages for all patient donors.

Figure 2

Figure 2. Compartment specific analysis of atlas. A) Batch correction metrics for each integration method. B) UMAP of embeddings from chosen integration method (RPCA) and the separation by broad cell type. C) Expression of canonical markers for each of the broad cell types separated from the atlas. D) Distribution of cells from each dataset in each of the three broad compartments. E) Comparison of broad cell type label between author annotations and atlas-based annotations. F) Expression of canonical markers for the relabeled cells if categorized by the atlas labels. G) Expression of canonical markers for the relabeled cells if categorized by their original labels.

Figure 3

Figure 3. Multi-resolution characterization of Cancer Epithelial Compartment. A) Heatmap of patient metadata, malignancy data (InferCNV), annotation data (HBCA), cell potency data (Cytotrace), plasticity data (EMP), and expression of hallmark pathways (Hallmark). B) Dendrogram of epithelial clusters learned from K2 Taxonomer. C) Top 5 important features, as determined by gini-based importance, from random forest models trained to differentiate different clades within each level of the dendrogram. Red cells indicate that scores related to that feature were higher than the mean in that subset of the dataset.

Figure 4

Figure 4. Diversity of immune cell types in the TME. A) Sankey diagram of all immune subtypes characterized in the atlas. B) Dendrogram of immune annotations learned from K2 Taxonomer. C) Dot plot of markers for the lymphoid subtypes identified in the atlas. D) Dot plot of markers for the myeloid subtypes identified in the atlas.

Figure 5

Figure 5. Diversity of stromal cell types in the TME. A) Sankey diagram of all stromal subtypes characterized in the atlas. B) Dendrogram of stromal annotations learned from K2 Taxonomer. C) Dot plot of markers for the stromal subtypes identified in the atlas.

Figure 6

Figure 6. Differential cell type diversity and abundance analysis. A) Cell Type Diversity Score (CTDS) across tumor subtype, patient age, and tumor grade. B) Differential cell type estimates within the immune compartment between HER2+, ER+ and TNBC tumors. B) Differential cell type estimates within the immune compartment between high and low grade tumors.

Figure 7

Figure 7. Survival associations of subpopulations found within taxonomic analysis of epithelial, stromal, and immune compartments with respect to TCGA and METABRIC. Green branches indicate a clade that confers a survival advantage (hazard ratio < 1), and red branches indicate a survival disadvantage (hazard ratio > 1) in the respective dataset.

Supplementary Figure 1. UMAP of unintegrated data colored by dataset, subtype, grade, celltypist, singleR, and author annotation.

Supplementary Figure 2. UMAP of RPCA integrated data colored by clustering, and broad cell type annotations from singleR, CellTypist, and author provided metadata.

Supplementary Figure 3. Cancer Epithelial Compartment Diversity. A) UMAP of cancer epithelial cells colored by clustering performed at 0.2 resolution. B) UMAP of cancer epithelial cells colored by HBCA annotation from CellTypist. C) UMAP of cancer epithelial cells colored by tumor subtype. D) UMAP of cancer epithelial cells 625 colored by PAM 50 subtype. E) Distribution of patients across each of the clusters. F) Entropy of donor proportions across each of the clusters.

Supplementary Figure 4. UMAP of immune compartment colored by clustering, and cell type annotations from singleR, CellTypist, and author provided metadata.

S5

Supplementary Figure 5. UMAP of stromal compartment colored by clustering, and cell type annotations from singleR, CellTypist, and author provided metadata.

Supplementary Figure 6. Distribution of inferCNV scores for each of the datasets included in the atlas. Immune and stromal cells sampled from each donor are in yellow whereas epithelial cells are in orange. The empirical 90% interval of malignancy scores for normal cells is highlighted for each dataset and is the interval used to classify epithelial cells as malignant or non-malignant.

References

1. Siegel, R. L., Giaquinto, A. N. & Jemal, A. Cancer statistics, 2024. *CA Cancer J. Clin.* **74**, 12–49 (2024).

2. Perou, C. M. *et al.* Molecular portraits of human breast tumours. *Nature* **406**, 747–752 (2000).

3. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61–70 (2012).

4. Polyak, K. Breast cancer: origins and evolution. *J. Clin. Invest.* **117**, 3155–3163 (2007).

5. Lüönd, F., Tiede, S. & Christofori, G. Breast cancer as an example of tumour heterogeneity and tumour cell plasticity during malignant progression. *Br. J. Cancer* **125**, 164–175 (2021).

6. Place, A. E., Jin Huh, S. & Polyak, K. The microenvironment in breast cancer progression: biology and implications for treatment. *Breast Cancer Res.* **13**, 227 (2011).

7. Kolodziejczyk, A. A., Kim, J. K., Svensson, V., Marioni, J. C. & Teichmann, S. A. The technology and biology of single-cell RNA sequencing. *Mol. Cell* **58**, 610–620 (2015).

8. Fleck, J. S., Camp, J. G. & Treutlein, B. What is a cell type? *Science* **381**, 733–734 (2023).

9. Reed, A. D. *et al.* A single-cell atlas enables mapping of homeostatic cellular shifts in the adult human breast. *Nat. Genet.* **56**, 652–662 (2024).

10. Kumar, T. *et al.* A spatially resolved single-cell genomic atlas of the adult human breast. *Nature* **620**, 181–191 (2023).

11. Pal, B. *et al.* A single-cell RNA expression atlas of normal, preneoplastic and tumorigenic states in the human breast. *EMBO J.* **40**, e107333 (2021).

12. Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat. Genet.* **53**, 1334–1347 (2021).

13. Tietscher, S. *et al.* A comprehensive single-cell map of T cell exhaustion-associated

665   immune environments in human breast cancer. *Nat. Commun.* **14**, 98 (2023).

666  14. Bassez, A. *et al.* A single-cell map of intratumoral changes during anti-PD1 treatment of

667    patients with breast cancer. *Nat. Med.* **27**, 820–832 (2021).

668  15. Wang, L. *et al.* PD-L1-expressing tumor-associated macrophages are immunostimulatory

669    and associate with good clinical outcome in human breast cancer. *Cell Rep. Med.* **5**,

670    101420 (2024).

671  16. Qian, J. *et al.* A pan-cancer blueprint of the heterogeneous tumor microenvironment

672    revealed by single-cell profiling. *Cell Res.* **30**, 745–762 (2020).

673  17. Liu, Y.-M. *et al.* Combined single-cell and spatial transcriptomics reveal the metabolic

674    evolvement of breast cancer during early dissemination. *Adv. Sci. (Weinh.)* **10**, e2205395

675    (2023).

676  18. Gao, R. *et al.* Delineating copy number and clonal substructure in human tumors from

677    single-cell transcriptomes. *Nat. Biotechnol.* **39**, 599–608 (2021).

678  19. Wu, S. Z. *et al.* Stromal cell diversity associated with immune evasion in human triple-

679    negative breast cancer. *EMBO J.* **39**, e104063 (2020).

680  20. Wu, S. Z. *et al.* Cryopreservation of human cancers conserves tumour heterogeneity for

681    single-cell multi-omics analysis. *Genome Med.* **13**, 81 (2021).

682  21. Xu, L. *et al.* A comprehensive single-cell breast tumor atlas defines epithelial and immune

683    heterogeneity and interactions predicting anti-PD-1 therapy response. *Cell Rep Med*

684    101511 (2024).

685  22. Johnston, K. G., Grieco, S. F., Nie, Q., Theis, F. J. & Xu, X. Small data methods in omics: the

686    power of one. *Nat. Methods* (2024) doi:10.1038/s41592-024-02390-8.

687  23. Luecken, M. D. *et al.* Benchmarking atlas-level data integration in single-cell genomics.

688    *Nat. Methods* **19**, 41–50 (2022).

689  24. Sun, X.-X. & Yu, Q. Intra-tumor heterogeneity of cancer cells and its implications for

cancer treatment. *Acta Pharmacol. Sin.* **36**, 1219–1227 (2015).

25. Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of developmental

potential. *Science* **367**, 405–411 (2020).

26. Winkler, J. *et al.* Single-cell analysis of breast cancer metastasis reveals epithelial-

mesenchymal plasticity signatures associated with poor outcomes. *J. Clin. Invest.* **134**,

(2024).

27. Tickle, T., Tirosh, I., Georgescu, C., Brown, M. & Haas, B. inferCNV of the Trinity CTAT

Project. Preprint at https://github.com/broadinstitute/inferCNV (2019).

28. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set

collection. *Cell Syst* **1**, 417–425 (2015).

29. Reed, E. R. & Monti, S. Multi-resolution characterization of molecular taxonomies in bulk

and single-cell transcriptomics data. *Nucleic Acids Res.* **49**, e98 (2021).

30. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a

transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

31. Domínguez Conde, C. *et al.* Cross-tissue immune cell analysis reveals tissue-specific

features in humans. *Science* **376**, eabl5197 (2022).

32. Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional changes

and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol.* **16**,

278 (2015).

33. Guimarães, G. R. *et al.* Single-cell resolution characterization of myeloid-derived cell

states with implication in cancer outcome. *Nat. Commun.* **15**, 5694 (2024).

34. Strack, E. *et al.* Identification of tumor-associated macrophage subsets that are

associated with breast cancer prognosis. *Clin. Transl. Med.* **10**, e239 (2020).

35. Chu, Y. *et al.* Pan-cancer T cell atlas links a cellular stress response state to

immunotherapy resistance. *Nat Med* **29**, 1550–1562 (2023).

36. Zheng, L. *et al.* Pan-cancer single-cell landscape of tumor-infiltrating T cells. *Science* **374**,

abe6474 (2021).

37. Kiselev, V. Y., Andrews, T. S. & Hemberg, M. Challenges in unsupervised clustering of

single-cell RNA-seq data. *Nat. Rev. Genet.* **20**, 273–282 (2019).

38. Du, Y. *et al.* Integration of Pan-Cancer Single-Cell and Spatial Transcriptomics Reveals

Stromal Cell Features and Therapeutic Targets in Tumor Microenvironment. *Cancer Res.*

84, 192–210 (2024).

39. Kapoor, T., Corrado, M., Pearce, E. L., Pearce, E. J. & Grosschedl, R. MZB1 enables efficient

interferon α secretion in stimulated plasmacytoid dendritic cells. *Sci. Rep.* **10**, 21626

(2020).

40. Karagiannis, T. T., Monti, S. & Sebastiani, P. Cell Type Diversity Statistic: An Entropy-Based

Metric to Compare Overall Cell Type Composition Across Samples. *Front. Genet.* **13**,

855076 (2022).

41. Mangiola, S. *et al.* sccomp: Robust differential composition and variability analysis for

single-cell data. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2203828120 (2023).

42. Perrone, M., Talarico, G., Chiodoni, C. & Sangaletti, S. Impact of immune cell

heterogeneity on HER2+ breast cancer prognosis and response to therapy. *Cancers*

(Basel) **13**, 6352 (2021).

43. Onkar, S. *et al.* Immune landscape in invasive ductal and lobular breast cancer reveals a

divergent macrophage-driven microenvironment. *Nat. Cancer* **4**, 516–534 (2023).

44. Fan, Y. & He, S. The characteristics of tumor microenvironment in triple negative breast

cancer. *Cancer Manag. Res.* **14**, 1–17 (2022).

45. Yuan, Z.-Y., Luo, R.-Z., Peng, R.-J., Wang, S.-S. & Xue, C. High infiltration of tumor-

associated macrophages in triple-negative breast cancer is associated with a higher risk

of distant metastasis. *Onco. Targets. Ther.* **7**, 1475–1480 (2014).

46. Oshi, M. *et al.* Plasmacytoid dendritic cell (pDC) infiltration correlate with tumor

infiltrating lymphocytes, cancer immunity, and better survival in triple negative breast

cancer (TNBC) more strongly than conventional dendritic cell (cDC). *Cancers (Basel)* **12**,

3342 (2020).

47. Majorini, M. T., Colombo, M. P. & Lecis, D. Few, but efficient: The role of mast cells in

breast cancer and other solid tumors. *Cancer Res.* **82**, 1439–1447 (2022).

48. Venet, D., Dumont, J. E. & Detours, V. Most random gene expression signatures are

significantly associated with breast cancer outcome. *PLoS Comput. Biol.* **7**, e1002240

(2011).

49. Jin, Y. *et al.* Prognostic impact of memory CD8(+) T cells on immunotherapy in human

cancers: A systematic review and meta-analysis. *Front. Oncol.* **11**, 698076 (2021).

50. Henson, D. E., Ries, L., Freedman, L. S. & Carriaga, M. Relationship among outcome, stage

of disease, and histologic grade for 22,616 cases of breast cancer. The basis for a

prognostic index. *Cancer* **68**, 2142–2149 (1991).

51. Chatsirisupachai, K., Lagger, C. & de Magalhães, J. P. Age-associated differences in the

cancer molecular landscape. *Trends Cancer* **8**, 962–971 (2022).

52. Miggelbrink, A. M. *et al.* CD4 T-cell exhaustion: Does it exist and what are its roles in

cancer? *Clin. Cancer Res.* **27**, 5742–5752 (2021).

53. Scully, T., Kase, N., Gallagher, E. J. & LeRoith, D. Regulation of low-density lipoprotein

receptor expression in triple negative breast cancer by EGFR-MAPK signaling. *Sci. Rep.* **11**,

17927 (2021).

54. Horlock, C. *et al.* The effects of trastuzumab on the CD4+CD25+FoxP3+ and CD4+IL17A+

T-cell axis in patients with breast cancer. *Br. J. Cancer* **100**, 1061–1067 (2009).

55. Plitas, G. *et al.* Regulatory T cells exhibit distinct features in human breast cancer.

Immunity **45**, 1122–1134 (2016).

56. Sebastião, A. I. *et al.* Dendritic cells in triple-negative breast cancer: From

pathophysiology to therapeutic applications. *Cancer Treat. Rev.* **133**, 102884 (2025).

57. Deligne, C. & Midwood, K. S. Macrophages and extracellular matrix in breast cancer:

Partners in crime or protective allies? *Front. Oncol.* **11**, 620773 (2021).

58. Gao, Y. *et al.* Adipocytes promote breast tumorigenesis through TAZ-dependent secretion

of Resistin. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 33295–33304 (2020).

59. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

60. Heumos, L. *et al.* Best practices for single-cell analysis across modalities. *Nat. Rev. Genet.*

24, 550–572 (2023).

61. Amezquita, R. A. *et al.* Orchestrating single-cell analysis with Bioconductor. *Nat. Methods*

17, 137–145 (2020).

62. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet

identification in single-cell sequencing data using scDblFinder. *F1000Res.* **10**, 979 (2021).

63. Ahlmann-Eltze, C. & Huber, W. Comparison of transformations for single-cell RNA-seq

data. *Nat. Methods* **20**, 665–672 (2023).

64. Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell RNA-

sequencing data are corrected by matching mutual nearest neighbors. *Nat. Biotechnol.*

36, 421–427 (2018).

65. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

66. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for

single-cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).

67. Xu, C. *et al.* Probabilistic harmonization and annotation of single-cell transcriptomics data

with deep generative models. *Mol. Syst. Biol.* **17**, e9620 (2021).

68. Gendoo, D. M. A. *et al.* Genefu: an R/Bioconductor package for computation of gene

expression-based signatures in breast cancer. *Bioinformatics* **32**, 1097–1099 (2016).

69. Xu, C. *et al.* Automatic cell-type harmonization and integration across Human Cell Atlas

datasets. *Cell* **186**, 5876-5891.e20 (2023).

70. Dance, A. What is a cell type, really? The quest to categorize life's myriad forms. *Nature*

633, 754–756 (2024).

71. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–

15550 (2005).

72. Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat.*

Methods **14**, 1083–1086 (2017).

73. Chen, A. *Brcasurv*. (2025).

74. Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours

reveals novel subgroups. *Nature* **486**, 346–352 (2012).

75. Winslow, S., Leandersson, K., Edsjö, A. & Larsson, C. Prognostic stromal gene signatures

in breast cancer. *Breast Cancer Res.* **17**, 23 (2015).

76. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray

and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).

77. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful

approach to multiple testing. *Journal of the royal statistical society series b-*

methodological **57**, 289–300 (1995).

78. Fisher, R. A. *Statistical Methods for Research Workers*. vol. no. V (Oliver and Boyd,

Edinburgh, 1938).

79. Find The Cause Breast Cancer Foundation. https://findthecausebcf.org/ (2016).