1 **Independent mechanisms of benzimidazole resistance across *Caenorhabditis* nematodes**
2

3 Amanda O. Shaver[1]*, Ryan McKeown[2]*, Joyce M. Reyes Otero[1], and Erik C. Andersen[1]
4
5 [1] Dept. of Biology, Johns Hopkins University, Baltimore, Maryland, USA
6 [2] Dept. of Molecular Biosciences, Northwestern University, Evanston, Illinois, USA
7 * These authors contributed equally.
8
Corresponding author:
Erik C. Andersen
Johns Hopkins University
Department of Biology
3400 N. Charles St.
Bascom UTL 383
Baltimore, MD 21218
410-516-1282
erik.andersen@gmail.com
18
ORCIDs and emails:
Amanda O. Shaver: 0000-0002-2910-1505, amanda.shaver@jhu.edu,
amandaoshaver@gmail.com
Ryan McKeown: 0000-0003-3570-2494, ryanmckeown2021@u.northwestern.edu
Joyce M. Reyes Otero: 0009-0002-5039-4434, joyce.reyes2@upr.edu
Erik C. Andersen: 0000-0003-0229-9651, erik.andersen@gmail.com

Keywords: Repeated evolution, *C. elegans*, *C. briggsae*, *C. tropicalis*, beta-tubulin,
benzimidazoles

ABBREVIATIONS

ABZ — Albendazole

BZs — Benzimidazoles

Cbr-ben-1 — *ben-1* in *Caenorhabditis briggsae*

Cbr-BEN-1 — BEN-1 in *Caenorhabditis briggsae*

Cbr-tbb-1 — *tbb-1* in *Caenorhabditis briggsae*

Cbr-TBB-1 — TBB-1 in *Caenorhabditis briggsae*

Cbr-tbb-2 — *tbb-2* in *Caenorhabditis briggsae*

Cbr-TBB-2 — TBB-2 in *Caenorhabditis briggsae*

Cel-ben-1 — *ben-1* in *Caenorhabditis elegans*

Cel-BEN-1 — BEN-1 in *Caenorhabditis elegans*

Cel-tbb-1 — *tbb-1* in *Caenorhabditis elegans*

Cel-TBB-1 — TBB-1 in *Caenorhabditis elegans*

Cel-tbb-2 — *tbb-2* in *Caenorhabditis elegans*

Cel-TBB-2 — TBB-2 in *Caenorhabditis elegans*

Ctr-ben-1 — *ben-1* in *Caenorhabditis tropicalis*

Ctr-BEN-1 — BEN-1 in *Caenorhabditis tropicalis*

Ctr-tbb-1 — *tbb-1* in *Caenorhabditis tropicalis*

Ctr-TBB-1 — TBB-1 in *Caenorhabditis tropicalis*

Ctr-tbb-2 — *tbb-2* in *Caenorhabditis tropicalis*

Ctr-TBB-2 — TBB-2 in *Caenorhabditis tropicalis*

HSB — Horvitz Super Broth

HTA — High-throughput larval development assay

LoF — Loss-of-function

NGMA — Nematode growth media agar

SNV — Single nucleotide variants

SV   Structural variants

<u>ABSTRACT</u>

Benzimidazoles (BZs), a widely used class of anthelmintic drugs, target beta-tubulin proteins,

disrupt microtubule formation, and cause nematode death. In parasitic nematode species,

mutations in beta-tubulin genes (*e.g.*, isotype-1 beta-tubulin) are predicted to inhibit BZ binding

and are associated with BZ resistance. Similarly, in the free-living nematode *Caenorhabditis*

elegans, mutations in an isotype-1 beta-tubulin ortholog, *ben-1*, are the primary drivers of BZ

resistance. The recurrent association of BZ resistance with beta-tubulins suggests that BZ

resistance is repeatedly caused by mutations in beta-tubulin genes, an example of repeated

evolution of drug resistance across nematode species. To evaluate the hypothesis of repeated

evolution of BZ resistance mediated by beta-tubulin, we identified predicted resistance alleles in

beta-tubulin genes across wild strains from three *Caenorhabditis* species: *C. elegans*,

Caenorhabditis briggsae, and *Caenorhabditis tropicalis*. We hypothesized that, if these species

experienced similar selective pressures, they would evolve resistance to BZs by mutations in any

of three beta-tubulin genes (*ben-1*, *tbb-1*, and *tbb-2*). Using high-throughput development assays,

we tested the association of predicted beta-tubulin alleles with BZ resistance. We found that a

heterogeneous set of variants identified in *C. elegans ben-1* were associated with BZ resistance.

In *C. briggsae*, only two variants in *ben-1*, predicted to encode a premature stop codon (W21stop)

and a missense substitution (Q134H), were associated with BZ resistance. In *C. tropicalis*, two

missense variants were identified in *ben-1*, but neither was associated with BZ resistance. *C.*

briggsae and *C. tropicalis* might have evolved BZ resistance by mutations in other beta-tubulin

genes, but we found that variants in *tbb-1* or *tbb-2* in these species were not associated with BZ

resistance. Our findings reveal a lack of repeated evolution of BZ resistance across the three

Caenorhabditis species and highlight the importance of defining BZ resistance mechanisms

outside of beta-tubulins.

1. INTRODUCTION

Global control of parasitic nematode infections relies on the efficacy of a small arsenal of anthelmintic drugs, including benzimidazoles (BZs) (A. C. Kotze et al. 2020). BZs are a widely used class of anthelmintic drugs that target beta-tubulin proteins (Lubega and Prichard 1990), disrupt microtubule formation (Neff et al. 1983; Laclette, Guerra, and Zetina 1980; Ireland et al. 1979), and lead to nematode death. Although BZs are essential in human and veterinary medicine, resistance is prominent in parasitic nematode populations (Theodorides, Scott, and Lademan 1970; Roos, Kwa, and Grant 1995). To effectively screen for and manage the emergence of BZ resistance, it is necessary to understand the underlying genetics that contribute to the evolution of BZ resistance in nematode species.

Historically, BZ resistance in parasite populations (*i.e.*, *Haemonchus contortus*, *Teladorsagia circumcincta*, and *Trichostrongylus colubriformis*) has been associated with three canonical missense variants (F167Y, E198A, and F200Y) in orthologs of the *H. contortus isotype-1* (Silvestre and Cabaret 2002; Ghisi, Kaminsky, and Mäser 2007; de Lourdes Mottier and Prichard 2008; Kwa, Veenstra, and Roos 1994) and *isotype-2 beta-tubulin* genes (A. C. Kotze and Prichard 2016; Avramenko et al. 2019). These three missense variants have been used to track and manage BZ resistance across nematode species globally (Kwa, Veenstra, and Roos 1994; Silvestre and Cabaret 2002; Ghisi, Kaminsky, and Mäser 2007). However, these three missense variants do not explain all of the BZ resistance observed in parasitic nematode populations (Andrew C. Kotze et al. 2014; Krücken et al. 2017). Recently, additional novel missense variants have been associated with BZ resistance (*e.g.*, E198I, E198K, E198T, E198stop, and Q134H) (Venkatesan et al. 2023; Mohammedsalih et al. 2020). The three canonical missense variants, along with novel missense variants of these parasitic nematode beta-tubulin alleles, conferred BZ resistance when introduced in the free-living nematode *Caenorhabditis elegans* (Dilks et al. 2020, 2021). Unlike parasitic species, *C. elegans* wild strains

have a heterogeneous set of variants in one of the *isotype-1 beta-tubulin* orthologs, *ben-1*,

responsible for much of BZ resistance in the species (Hahnel et al. 2018). The recurrent

association of BZ resistance with alleles predicted to impact beta-tubulin function across

nematode species suggests that BZ resistance repeatedly evolves by standing variation or

recurrent mutations in beta-tubulin genes and provides compelling evidence to predict the

emergence of BZ resistance across nematode species.

Repeated evolution is the development of a similar phenotype (*e.g.*, BZ resistance) and

genotype (*e.g.*, the same beta-tubulin alleles confer BZ resistance) in response to similar

environmental pressures (*e.g.*, BZ exposure) (Cerca 2023). Because parasitic and free-living

nematodes have evolved BZ resistance by variants in conserved beta-tubulin genes, we

hypothesized that nematodes acquire BZ resistance by repeated variants or mutations in beta-

tubulin genes. However, this hypothesis is difficult to test directly in parasitic nematode species

because of their host-dependent life cycles, poorly annotated reference genomes, and limited

molecular and genetic tools (Hahnel et al. 2020; Mariene and Wasmuth 2025). By contrast, the

availability of high-quality genomic data for hundreds of wild strains (Crombie et al. 2023) and the

laboratory tractability of the free-living *Caenorhabditis* nematode species (*C. elegans*,

Caenorhabditis briggsae, and *Caenorhabditis tropicalis*) provide an opportunity to test for the

repeated evolution of BZ resistance in the *Caenorhabditis* genus.

Using the global natural diversity of *C. elegans*, *C. briggsae*, and *C. tropicalis*, we

assessed variation in conserved beta-tubulin genes to identify predicted BZ resistance alleles.

We identified single nucleotide variants (SNVs), small insertions or deletions (INDELs), and

structural variants (SVs) predicted to impact function in conserved beta-tubulin genes, herein

called "high-impact" variants. Because high-impact variants in the *Cel-ben-1* gene are known to

account for much of the BZ resistance in *C. elegans*, we first used an established high-throughput

larval development assay (HTA) to expose *C. elegans* strains with novel *ben-1* variants to the

prominently used BZ, albendazole (ABZ), to identify new resistance alleles. We found eight novel

variants in *Cel-ben-1*, of which six were associated with ABZ resistance in the HTAs, a result that

further confirms the role *Cel-ben-1* plays in ABZ resistance. Next, we used the HTA to expose

C. briggsae and *C. tropicalis* strains with novel *ben-1* variants to ABZ to identify if these two

species confer ABZ resistance by the same mechanism as in *C. elegans*. In *C. briggsae*, two of

the eleven BEN-1 variants (W21stop and Q134H) were correlated with ABZ resistance. In *C.*

tropicalis, neither of the two BEN-1 variants (P80T and R121Q) were associated with ABZ

resistance. Because we found a lack of repeated evolution of BZ resistance by variants in *ben-1*,

we tested whether *C. briggsae* and *C. tropicalis* evolved BZ resistance by variants in the beta-

tubulin genes *tbb-1* and *tbb-2*, but found no high-impact variants in these genes associated with

ABZ resistance. Overall, our results indicate a lack of repeated evolution of ABZ resistance across

Caenorhabditis species, suggesting that beta-tubulin cannot underlie BZ resistance across all

nematode species.

2. MATERIALS AND METHODS

2.1 Identification of beta-tubulin loci

Amino acid sequences for all six *C. elegans* beta-tubulin proteins were obtained from

WormBase (WS283) (Sternberg et al. 2024) and used as queries in a BLASTp search (Version

2.12.0) (Camacho et al. 2009). The search was performed against protein sequence databases

constructed using gene models for *C. briggsae* (Moya et al. 2023) and *C. tropicalis* (Noble et al.

2021). To construct the protein sequence database, we extracted gene model transcript features

from the gene feature file with *gffread* (Version 0.9.11) (Pertea and Pertea 2020) and processed

them using the *makeblastdb* function from BLAST (Version 2.12.0). From the BLASTp search, we

identified *C. briggsae* and *C. tropicalis* protein sequences with the highest percent identity (PID)

to each *C. elegans* beta-tubulin protein. Only protein sequences with the highest PID in both

searches were considered orthologs (**S1 Table**). For some *C. elegans* beta-tubulin orthologs,

multiple *C. briggsae* or *C. tropicalis* gene models contained multiple splice isoforms. All gene

models for all beta-tubulin transcripts were manually inspected, and isoforms that were not fully

supported by short-read RNA sequencing data were removed.

2.2 Single nucleotide variant (SNV) and indel calling and annotation

To identify single nucleotide variants (SNVs) or indels (insertions and deletions) in the

beta-tubulin genes across the selfing *Caenorhabditis* species, we used the Variant Annotation

Tool from the *Caenorhabditis* Natural Diversity Resource (CaeNDR) (Release IDs: *C. elegans* -

20231213, *C. briggsae* - 20240129, *C. tropicalis* - 20231201) (Crombie et al. 2023). The identified

SNVs and indels (**S2, S3, and S4 Tables**) included small insertions and deletions, frameshifts,

altered stop and start codons, nonsynonymous changes, and splice variants.

2.3 Structural variant (SV) calling and annotation

Structural variant (SV) calling was performed using *DELLY* (Version 0.8.3), a SV caller

optimized to detect large insertions, deletions, and other complex structural variants such as

inversions, translocations, and duplications in paired-end short-read alignments (Rausch et al.

2012) and shown to perform well on *C. elegans* short-read sequence data (Lesack et al. 2022).

SVs that overlapped with beta-tubulin genes were extracted using *bcftools* (Version 1.10.1)

(Danecek et al. 2021). Insertions, deletions, inversions, and duplications that passed the *DELLY*

(Version 0.8.3) default quality threshold (greater than three supporting read pairs with a median

MAPQ > 20), filtered to high-quality genotypes (genotype quality > 15), and had at least one

alternative allele were retained. For complex variants (inversions and duplications), the

identification of at least one split-read pair was required (variants flagged as a precise SV by

DELLY). To validate SVs that passed quality filtering, each SV was manually inspected for

breakpoints in the raw-read alignments (*Wally*, Version 0.5.8) and for impacts on the beta-tubulin

coding sequence (CaeNDR Genome Browser) (Crombie et al. 2023) (**S5 Table**). We retained

SVs where raw read alignments suggested that the SV impacted the beta-tubulin coding

sequence. We compared the *Cel-ben-1* SVs called by *DELLY* to those SVs identified previously

(Hahnel et al. 2018). *DELLY* successfully recalled structural variants in several strains, including

deletions in JU751, JU830, JU1395, JU2582, JU2587, JU2593, JU2829, and QX1233, as well as

an inversion in MY518. However, *DELLY* did not detect a previously reported transposon insertion

in strain JU3125. To assess if other SVs could have been missed by *DELLY*, we manually

inspected the read alignments for all strains that had not been previously phenotyped to check if

any other SVs were not detected by *DELLY*. We confirmed the presence of novel *Cel-ben-1* SVs

in multiple strains, including deletions in ECA706 and NIC1832, and a duplication in NIC1107.

Additionally we identified a previously undetected deletion in JU4287.

2.4 Association of low *Cel-ben-1* expression with ABZ response

Two previous assays measured developmental responses of wild *C. elegans* strains after

ABZ exposure (Hahnel et al. 2018; Shaver et al. 2023). For 180 of these wild *C. elegans* strains,

expression levels (transcripts per million estimates [TPM]) were collected at the young-adult stage

(Zhang et al. 2022). Of the 180 wild *C. elegans* strains, 105 strains were shared between both

assays. A linear model was built using the *lm* function in R to account for assay effects.

Subsequently, the residuals of the linear model were used to normalize previous measures of

ABZ response. We evaluated the linear fit between each strain's expression of *ben-1*, *tbb-1*, or

tbb-2 and the developmental delay following ABZ exposure.

2.5 Phylogenetic analysis

We characterized the relatedness of isotype reference strains (genetically unique strains)

with beta-tubulin variants using species trees downloaded from CaeNDR and generated by the

'post-gatk-nf' pipeline (https://github.com/AndersenLab/post-gatk-nf) (Release IDs: *C. elegans* -

20231213, *C. briggsae* - 20240129, *C. tropicalis* - 20231201) (Crombie et al. 2023). Briefly, the

trees were generated with high-quality SNVs in isotype reference strains retained in the hard-

filtered variant call format (VCF) file. *vcf2phylip* (Ortiz 2019) and the *bioconvert* (Caro et al. 2023)

function *phylip2stockholm* were used to prepare inputs for *quicktree*, which was used to construct

a tree using a neighbor-joining algorithm (Saitou and Nei 1987). All versions of these software

can   be   accessed   from   the   'post-gatk'   docker   container

(https://hub.docker.com/r/andersenlab/tree), used by the 'post-gatk-nf' pipeline. We visualized the

trees for each species using the *ggtree* function from the *ggtree* (v3.6.2) R package (Yu 2020).

2.6 Strain selection and maintenance

Fifteen *C. elegans* strains, 38 *C. briggsae* strains, and seven *C. tropicalis* strains from the

CaeNDR (Crombie et al. 2023) were used in this study (**S2, S3, and S4 Tables)**. Isolation details

for each strain are included in CaeNDR. For each species, we selected strains that had unique

high-impact consequences (SNV or SV) in *ben-1*, *tbb-1*, or *tbb-2* that had not been previously

phenotyped. Strains with high-impact consequences in a beta-tubulin gene are herein referred to

as "predicted resistant" strains. Strains with no high-impact consequences in beta-tubulin genes

that were closely related to predicted resistant strains were also included for *C. briggsae* and *C.*

tropicalis and herein classified as "predicted susceptible" strains (**S3 and S4 Figures**) (**Table S6**).

The reference strain(s) for all three species was included. Finally, for *C. elegans*, a strain with a

loss of *ben-1* (ECA882) was also included.

Before measuring ABZ responses, *C. elegans* and *C. briggsae* animals were maintained

at 20ºC and *C. tropicalis* animals were maintained at 25ºC. All animals were maintained on 6 cm

plates with modified nematode growth medium (NGMA), which contains 1% agar and 0.7%

agarose to prevent animals from burrowing (Andersen et al. 2014). The NGMA plates were

seeded with the *Escherichia coli* strain OP50 as a nematode food source. All strains were grown

for three generations without starvation on NGMA plates before anthelmintic exposure to reduce

the transgenerational effects of starvation stress (Andersen et al. 2015).

2.7 Nematode food preparation for NGMA 6 cm plates

A batch of OP50 *E. coli* was grown and used as a nematode food source for NGMA plates. A frozen stock of OP50 *E. coli* was streaked onto a 10 cm Luria-Bertani (LB) agar plate and incubated overnight at 37ºC. The following day, a single bacterial colony was transferred into two culture tubes that contained 5 mL of 1x LB. The starter cultures and two negative controls (1X LB without *E. coli*) were incubated for 18 hours at 37ºC shaking at 210 rpm. The OD_{600} value of the starter cultures were measured using a spectrophotometer (BioRad, SmartSpec Plus) to calculate how much starter culture was needed to inoculate a 1 L culture at an OD_{600} value of 0.005. For each assay, one culture containing one liter of pre-warmed 1X LB inoculated with the starter culture grew for approximately 4 - 4.5 hours at 37ºC at 210 rpm to an OD_{600} value between 0.45 and 0.6. Cultures were transferred to 4ºC to slow growth. OP50 was spotted on NGMA test plates (two per culture) and grown at 37ºC overnight to ensure no contamination.

2.8 Nematode food preparation for HTAs

One batch of HB101 *E. coli* was used as a nematode food source for all HTAs in this study. A frozen stock of HB101 *E. coli* was streaked onto a 10 cm LB agar plate and incubated overnight at 37ºC. The following day, a single bacterial colony was transferred into three culture tubes that contained 5 mL of 1x Horvitz Super Broth (HSB). The starter cultures and two negative controls (1X HSB without *E. coli*) were incubated for 18 hours at 37ºC shaking at 180 rpm. The OD_{600} value of the starter cultures were measured using a spectrophotometer (BioRad, SmartSpec Plus) to calculate how much starter culture was needed to inoculate a 1 L culture at an OD_{600} value of 0.001. A total of four cultures each containing 1 L of pre-warmed 1X HSB inoculated with the starter culture grew for 15 hours at 37ºC while shaking at 180 rpm. After 15 hours, flasks were removed from the incubator and transferred to 4ºC to slow growth. The 1X HSB was removed from the cultures by performing three rounds of centrifugation, where the

supernatant was removed, and the bacterial cells were pelleted. Bacterial cells were washed with

K medium, resuspended in K medium, pooled, and transferred to a 2 L glass beaker. The OD_{600}

value of the bacterial suspension was measured and diluted to a final concentration of OD_{600}100

with K medium, aliquoted to 15 mL conicals, and stored at -80ºC for use in the HTAs.

2.9 ABZ dose-response assays for *C. briggsae* and *C. tropicalis*

Because ABZ response has been minimally characterized in *C. briggsae* (Zamanian et al.

2018) and has not yet been described in *C. tropicalis*, we first measured dose-response curves

for both species after exposure to ABZ to assess developmental delay. Before performing HTAs,

ABZ (Sigma-Aldrich, Catalog # A4673-10G) stock solutions were prepared in dimethyl sulfoxide

(DMSO) (Fisher Scientific, Catalog # D1281), aliquoted, and stored at -20ºC for use in the assays.

For the dose-response assays, animals were exposed to ABZ at the following concentrations

(μM): 0 (1% DMSO), 0.12, 0.23, 0.47, 0.94, 1.88, 3.75, 7.5, 15, 30, 60, and 120. Animals were

allowed to develop in the presence of ABZ as described in *HTAs to assess nematode*

development.

Dose-response model estimation and statistics were performed as described previously

(Widmayer et al. 2022; Shaver et al. 2023). Briefly, a four-parameter log-logistic dose-response

curve was fit independently for a genetically diverse set of 11 *C. briggsae* strains (**S5 Figure**) and

seven *C. tropicalis* strains (**S6 Figure**), where normalized median animal length was used as a

metric for phenotypic response (see *Methods*). For each strain-specific dose-response model,

slope (*b*) and concentration (*e*) were estimated with strain as a covariate. We calculated EC_{10} as

we have previously found EC_{10} response to be more heritable than half maximal effective

concentration (EC_{50}) estimates and were therefore used in our analysis (Shaver et al. 2023;

Widmayer et al. 2022). A dosage of 30 μM ABZ was closest to the EC_{10} for *C. briggsae* and *C.*

tropicalis, consistent with ABZ concentrations used in past *C. elegans* assays (Dilks et al. 2020,

2021; Shaver et al. 2024) and in all HTAs in this study.

2.10 HTAs to assess nematode development

Populations of each strain were amplified and bleach-synchronized in three independent assays. Independent bleach synchronizations controlled for variation in embryo survival and subsequent effects on developmental rates. After bleach synchronization, approximately 30 embryos were dispensed into the wells of a 96-well microplate in 50 µL of K medium. Forty-eight wells were prepared per bleach for each strain. Each 96-well microplate was prepared, labeled, and sealed using gas-permeable sealing films (Fisher Scientific, Catalog # 14-222-043). Plates were placed in humidity chambers to incubate for 24 hours at 20°C while shaking at 170 rpm (INFORS HT Multitron shaker). After 24 hours, every plate was inspected to ensure that all embryos hatched and animals were developmentally arrested at the first larval (L1) stage so all strains started each assay at the same developmental stage. Next, food was prepared to feed the developmentally arrested L1 animals using the required number of OD_{600}100 HB101 aliquots (see *Nematode food preparation for HTAs*). The HB101 aliquots were thawed at room temperature, combined into a single conical tube, and diluted to an OD_{600}30 with K medium. To inhibit further bacterial growth and prevent contamination, 150 µL of kanamycin was added to the HB101. An aliquot of 100 µM ABZ stock solution was thawed at room temperature and added to an aliquot of OD_{600}30 K medium at a 3% volume/volume ratio. Next, 25 µL of the food and ABZ mixture was transferred into the appropriate wells of the 96-well microplates to feed the arrested L1 animals at a final HB101 concentration of OD_{600}10 and expose L1 animals to ABZ. Immediately afterward, the 96-well microplates were sealed using a new gas permeable sealing film, returned to the humidity chambers, and incubated for 48 hours at 20°C (*C. elegans* and *C. briggsae*) or 25°C (*C. tropicalis*) while shaking at 170 rpm. After 48 hours (*C. elegans* and *C. briggsae*) or 42 hours (*C. tropicalis*) of incubation and shaking in the presence of food and either DMSO or ABZ, the 96-well microplates were removed from the incubator and treated with 50 mM sodium azide in M9 for 10 minutes to paralyze and straighten nematodes. After 10 minutes, images of nematodes in

the microplates were immediately captured using Molecular Devices ImageXpress Nano

microscope (Molecular Devices, San Jose, CA) using a 2X objective. The ImageXpress Nano

microscope acquires brightfield images using a 4.7 megapixel CMOS camera and stores images

in a 16-bit TIFF format. The images were used to quantify the development of nematodes in the

presence of DMSO or ABZ as described below (see *High-throughputput imager assays [HTA]*

data collection and cleaning). A full step-by-step protocol for the HTA has been deposited in

protocols.io.

2.11 HTA data collection and data cleaning

CellProfiler (Version 24.10.1) was used to characterize and quantify biological data from

the image-based assays. Custom software packages designed to extract animal measurements

from images collected on the Molecular Devices ImageXpress Nano microscope were previously

described (Nyaanga et al. 2021). *CellProfiler* modules and *Worm Toolbox* were developed to

extract morphological features of individual animals from images from the HTA (Wählby et al.

2012). Worm model estimations and custom *CellProfiler* pipelines were written using the

WormToolbox in the GUI-based instance of *CellProfiler* (Widmayer et al. 2022). Next, a Nextflow

pipeline (Version 24) was written to run command-line instances of *CellProfiler* in parallel on the

Rockfish High-Performance Computing Cluster (Johns Hopkins University). The *CellProfiler*

workflow can be found at https://github.com/AndersenLab/cellprofiler-nf. The custom *CellProfiler*

pipeline generates animal measurements by using four worm models: three worm models tailored

to capture animals at the L4 larval stage, in the L2 and L3 larval stages, and the L1 larval stage,

as well as a "multi-drug high dose" (MDHD) model, to capture animals with more abnormal body

sizes caused by extreme anthelmintic responses. These measurements comprised our raw

dataset. Two *C. briggsae* strains (NIC1052 and VX34) were not fully paralyzed and straightened

at the time of imaging, which created some misclassification of animal measurements. Thus, the

animal lengths for strains NIC1052 and VX34 measured by CellProfiler are shorter than the actual

animal lengths. However, the difference discrepancy in animal lengths does not affect the

classification of a strain as resistant or sensitive to ABZ. Data cleaning and analysis steps were

performed using a custom R package, *easyXpress* (Version 2.0) (Nyaanga et al. 2021) and

followed methods previously reported (Shaver et al. 2024). All analyses were performed using

the R statistical environment (Version 4.2.1) unless stated otherwise.

3. RESULTS

3.1 In three free-living *Caenorhabditis* species, strains with high-impact variants in beta-

tubulin genes or low levels of beta-tubulin expression are predicted to confer ABZ

resistance

To investigate if the mechanisms of ABZ resistance are repeated across the three selfing

Caenorhabditis species, we identified "predicted resistant" strains that have high-impact variants

(*i.e.*, SNVs, INDELs, or SVs) (Crombie et al. 2023) likely to affect the function of the five conserved

beta-tubulin genes (*i.e.*, *ben-1*, *tbb-1*, *tbb-2*, *tbb-4*, and *mec-7*) (see *Methods*). Although ABZ

response has been highly characterized and associated with mutations in *ben-1* across *C.*

elegans wild strains (Hahnel et al. 2018; Shaver et al. 2024), less is known about how *C. briggsae*

(Zamanian et al. 2018) and *C. tropicalis* respond to ABZ exposure. To minimize the amount of

genetic variation outside of the beta-tubulin that could affect the BZ response, we also selected

C. briggsae and *C. tropicalis* strains that lacked high-impact variants in beta-tubulin genes but

were closely related to predicted resistant strains (**S3 and S4 Figures**). The strains with no high-

impact variants in any beta-tubulin gene were classified as "predicted susceptible".

In a set of 611 *C. elegans* strains (Crombie et al. 2023), we identified 65 strains with 33

unique high-impact variants in *Cel-ben-1* (**Table S2**). Of the 33 unique high-impact variants, 24

were previously phenotyped and 23 were associated with ABZ resistance (**S1A Figure**) (Hahnel

et al. 2018; Shaver et al. 2024). Here, we report eight novel high-impact variants predicted to

impact *Cel*-BEN-1 (E3stop, Y50C, P80S, VDN113N, frameshift 319, frameshift 368, stop445S,

and a duplication) that have not previously associated with ABZ resistance in nematodes (**Table**

S2). The missense variants predicted to encode Y50C and P80S *Cel*-BEN-1 substitutions have

been found in fungus and human beta-tubulins, respectively (Qiu et al. 2011; Dingerdissen et al.

2018). We found no strains with high-impact variants predicted to impact *Cel*-TBB-1 or *Cel*-TBB-

2.

Previously, low *ben-1* expression was correlated with ABZ resistance in *C. elegans* strains

(Zhang et al. 2022), providing another parameter to investigate BZ resistance across the species.

We evaluated if beta-tubulin expression levels were predictive of ABZ responses in *C. elegans.*

We used a linear regression model to assess the relationship between the expression of *Cel-ben-*

1, *Cel-tbb-1*, or *Cel-tbb-2* (Zhang et al. 2022) and ABZ responses in 180 wild strains (Hahnel et

al. 2018; Shaver et al. 2024). We found a significant negative relationship between *Cel-ben-1*

expression (>=3.75 TPM) and ABZ resistance (*p*-value = 5.16e-16, r^2= 0.344) (**S1 Figure**).

Importantly, 19 of the 23 strains with low *ben-1* expression also had high-impact variants in *ben-*

1 (SVs, frameshifts, or stop/start altering variants). Next, we identified strains with no high-impact

variants in *ben-1* but had low *ben-1* expression that we predicted to cause ABZ resistance. Using

these parameters, we found one strain, JU1581, with low *Cel-ben-1* expression to test for ABZ

resistance and to further define the role *Cel-ben-1* plays in ABZ resistance. No significant

relationship was found between *Cel-tbb*-2 and *Cel-tbb-1* expression and ABZ response (**S2**

Figure).

In a set of 641 *C. briggsae* strains, we identified 22 strains with eight unique high-impact

variants in *Cbr-ben-1* (**Table S3**). The eight unique high-impact variants in *Cbr*-BEN-1 included

seven amino acid substitutions (V91I, Q94K, D128E, Q134H, S218L, M299V, and R359H) and

one premature stop codon (W21stop). Outside of *Cbr*-BEN-1, we identified missense variants in

Cbr-TBB-1 (T35A, A275T, L377I, and V64I) and *Cbr*-TBB-2 (E441A) predicted to disrupt function.

All *C. briggsae* predicted resistant strains, predicted susceptible strains, and the reference strains

(AF16 and QX1410) (Stevens et al. 2022; Moya et al. 2023) were phenotyped for ABZ response

(**S1 Figure**).

In a set of 518 *C. tropicalis* strains, we identified two strains with unique high-impact

variants in *Ctr*-BEN-1 (**Table S4**) (P80T and R121Q). Outside of *Ctr*-BEN-1, we identified one

missense variant in *Ctr*-TBB-2 (N89S) and no high-impact variants in *Ctr*-TBB-1. All *C. tropicalis*

predicted resistant strains, three predicted susceptible strains, and the reference strain (NIC58)

(Luke et al. 2021) were phenotyped for ABZ response.

3.2 Natural allelic variation in *ben-1* is associated with ABZ resistance in *C. elegans*

We performed image-based HTAs across all *C. elegans* wild strains with novel high-

impact variants in *ben-1* (*i.e.*, predicted resistant strains) to measure nematode length (a proxy

for larval development) in drug (ABZ) and control (DMSO) conditions (see *Methods*). The

canonical laboratory strain N2 was included as an ABZ-susceptible control (Dilks et al. 2020,

2021; Shaver et al. 2024) and a strain with the loss of *ben-1* in the N2 strain background (ECA882)

was included as an ABZ-resistant control. Previously, two high-impact variants (frameshift at

position 299 and a deletion) were associated with ABZ resistance (Hahnel et al. 2018; Dilks et al.

2020). With expanded sampling, we identified two novel strains with the same frameshift at

position 299 (JR4305) and the same deletion (NIC1832) in *Cel-ben-1*. Therefore, we included

JR4305 and NIC1832 to validate the role that a frameshift at position 299 and a deletion play in

ABZ resistance. The assay included 48 replicates per strain with 5 to 30 animals per replicate in

ABZ or DMSO conditions. The reported nematode length of each strain is the delta between

animal lengths in DMSO (**S7 Figure**) and ABZ to obtain normalized animal length and assess

drug effects. We classified a strain as resistant to ABZ if its nematode length after ABZ exposure

was 75% greater than that of the reference strain (see *Methods*). A longer median animal length

(*i.e.*, larger animals) indicated resistance to ABZ, and a shorter median animal length (*i.e.*, smaller

animals) indicated susceptibility to ABZ.

As expected, when exposed to ABZ, strains ECA882 (loss of *ben-1*), JR4305 (frameshift

at position 299), and NIC1832 (deletion) all recapitulated a resistance phenotype (nematode

length ≥ 75% of the reference strain N2). Of the eight predicted resistant *C.elegans* strains, six

displayed a resistance phenotype (**Fig. 1**). All six wild strains showed minimal developmental

delays after exposure to ABZ, a phenotype similar to loss of *ben-1* (**Fig. 1**). Of all the assayed

predicted resistant strains with variants in *Cel*-BEN-1, three (P80S, stop445S, and low *ben-1*

expression) were not classified as resistant to ABZ. The P80S variant might partially alter *ben-1*

function, causing a moderate resistance phenotype. In stop445S, the normal stop codon was

replaced with a serine, likely allowing translation to continue beyond the termination point. This

variant likely does not affect *ben-1* function because position 445 is at the end of the BEN-1

protein. Finally, the strain with low *ben-1* expression (JU1581), which did not have any high-

impact variants in *ben-1*, was sensitive to ABZ, indicating that the selected threshold of *ben-1*

expression (≥3.75 TPM) still retained strains with adequate *ben-1* function.

Figure 1. High-throughput assays for each *C. elegans* strain with a high-impact variant in BEN-1 in the presence of albendazole

The regressed median animal length values for populations of nematodes grown in 30 μM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status. The gray dashed line marks the resistance threshold (nematode length ≥ 75% of the reference strain N2).

3.3 Rare missense and nonsense variants in *Cbr-ben-1* are associated with ABZ resistance,

whereas high-impact variants in *Ctr-ben-1* are not

We performed HTAs on 11 predicted resistant *C. briggsae* strains with eight unique

variants in *Cbr-ben-1* and on 13 predicted susceptible strains in DMSO (**S8 Figure**) and ABZ

conditions. We found that in *C. briggsae*, only two variants in BEN-1 (Q134H and W21stop)

conferred ABZ resistance (nematode length ≥ 75% of the reference strain AF16) (**Fig. 2**). The

Q134H amino acid change has been associated with ABZ resistance in *Ancylostoma caninum*

and validated in *C. elegans* (Venkatesan et al. 2023). An early stop gain at position 21 is predicted

to cause the premature termination of protein synthesis and *Cbr-ben-1* loss-of-function (LoF). Of

all the assayed strains with predicted resistance variants in *Cbr*-BEN-1, six (V91I, Q94K, D128E,

S218L, M299V, and R359H) were not resistant to ABZ. Overall, because only one of the seven

missense variants was associated with ABZ resistance, we could not reliably predict how a strain

responded to ABZ based on the presence of a missense variant alone. These results indicate that

ABZ resistance associated with *Cbr-ben-1* variants is uncommon and that accurately predicting

ABZ resistance in *C. briggsae* requires additional factors beyond *ben-1* variation.

Figure 2. High-throughput assays for each *C. briggsae* strain with a high-impact variant in BEN-1 and paired predicted susceptible strains in the presence of albendazole

The regressed median animal length values for populations of nematodes grown in 30 µM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status. The gray dashed line marks the resistance threshold (nematode length ≥ 75% of the reference strain AF16).

We performed HTAs on two *C. tropicalis* predicted resistant strains with high-impact

variants in *Ctr*-BEN-1 (P80T and R121Q) and two predicted susceptible strains in DMSO (**S9**

Figure) and ABZ conditions. We found that both predicted resistant strains had a similar ABZ

response as the predicted susceptible strains, where no strains met the resistance threshold

(nematode length ≥ 75% of the reference strain NIC58), indicating that P80T and R121Q are not

associated with ABZ resistance (**Fig. 3**). These results indicate that *Ctr*-BEN-1 variants do not

confer ABZ resistance and that predicting ABZ resistance in *C. tropicalis* requires considering

factors beyond *ben-1* variation. Additionally, to evaluate if the predicted resistant variants in *Cbr*-

BEN-1 or *Ctr*-BEN-1 were less deleterious compared to *Cel*-BEN-1, we assessed the BLOSUM

and Grantham scores of each high-impact variant (Crombie et al. 2023). The BLOSUM and

Grantham scores measure the evolutionary likelihood of observing a particular missense

substitution, and we found that these scores do not accurately predict ABZ resistance (**Table S7**).

We performed a linear regression analysis of strain ABZ responses by the BLOSUM or Grantham

scores of the beta-tubulin variant in each strain. We found no significant relationship between

either BLOSUM or Grantham scores and ABZ response (**S10, S11, S12 Figures**). For example,

the only missense substitution to cause ABZ resistance in Cbr-BEN-1, Q134H, had substantially

less radical BLOSUM and Grantham scores (BLOSUM: 0, Grantham: 24) compared to other

missense substitutions, such as S218L (BLOSUM: -2, Grantham: 145).

Figure 3. High-throughput assays for each *C. tropicalis* strain with a high-impact variant in BEN-1 or TBB-2 and paired predicted susceptible strains in the presence of albendazole

The regressed median animal length values for populations of nematodes grown in 30 μM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status. The gray dashed line marks the resistance threshold (nematode length ≥ 75% of the reference strain NIC58).

3.4 Natural allelic variants in *tbb-1* and *tbb-2* are not associated with ABZ resistance across three free-living *Caenorhabditis* species

To identify if high-impact variants in other beta-tubulin genes beyond *ben-1* play a role in ABZ resistance in the three free-living *Caenorhabditis* species, we categorized variants in the two genes most highly expressed in *C. elegans* (*tbb-1* and *tbb-2*). Because the loss of *tbb-1* or *tbb-2* is deleterious in *C. elegans* (Collins et al. 2024), LoF mutations in either gene would likely be purged by purifying selection from natural populations. Therefore, as expected, we found no variants in TBB-1 or TBB-2 in *C. elegans* wild strains. Unlike *C. elegans*, *C. briggsae* had four missense variants in TBB-1 (T35A, V64I, A275T, and L377I) and one missense variant in TBB-2 (E441A). In *C. tropicalis*, we identified one rare missense variant in TBB-2 (N89S). Performing the same HTA to assess nematode development as described previously, we found that none of the missense variants in TBB-1 or TBB-2 in *C. briggsae* (**Fig. 4**) or *C. tropicalis* conferred ABZ resistance (**Fig. 3**). These results suggest that missense variants in TBB-1 and TBB-2 are not associated with ABZ resistance in *Caenorhabditis* species.

Figure 4. High-throughput assays for each *C. briggsae* strain with a high-impact variant in TBB-1 or TBB-2 in the presence of albendazole

The regressed median animal length values for populations of nematodes grown in 30 μM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status. The gray dashed line marks the resistance threshold (nematode length ≥ 75% of the reference strain AF16.)

3.5 High-impact variants in beta-tubulin genes are rare and not enriched for geography or

substrate

To better understand the evolution of BZ resistance alleles in three *Caenorhabditis*

species, we assessed the population-wide frequencies of each beta-tubulin consequence, along

with the geographic location and substrate of where each strain was isolated from nature. First,

to determine the prevalence of high-impact variants in the five conserved beta-tubulin genes (*tbb-*

1, *tbb-2*, *mec-7*, *tbb-4*, and *ben-1*) across *Caenorhabditis* species, we quantified the frequency of

each consequence (deletion, duplication, frameshift, in-frame deletion, inversion, missense,

splice donor, and start/stop altering) in each species. Despite extensive gobal sampling of the

three *Caenorhabditis* species (*C. elegans*: 611 strains, *C. briggsae*: 641 strains, *C. tropicalis*: 518

strains), we found that all consequences in all three *Caenorhabditis* species were rare (< 0.05%

of all strains in a species) (**Fig. 5**). The most variation in consequences was found in *Cel*-BEN-1,

where deletions, frameshifts, in-frame deletions, inversions, missense, stop/start altering variants,

and a duplication were found. However, we found one missense variant in *Cel*-TBB-4 and several

missense variants and a splice donor in *Cel*-MEC-7. *Cel-tbb-6* is unique to *C. elegans* and was

therefore not assessed. Overall, *C. elegans* has acquired the most diverse set of consequences

in BEN-1 and has conferred BZ resistance primarily by variation in *Cel-ben-1*. In *C. briggsae*, we

found 21 missense amino-acid substitutions and a single strain with a start/stop altering

consequence in *Cbr*-BEN-1. In both *Cbr*-TBB-1 and *Cbr*-TBB-2, we identified rare missense

consequences. Finally, we found nine strains with splice variants in *Cbr*-TBB-4 and one strain

with a missense variant. In *C. tropicalis*, we found missense consequences in all beta-tubulin

genes except *tbb-1*. These findings highlight that *C. elegans* has the most diverse set of beta-

tubulin variants, particularly in *ben-1*, reinforcing its role in BZ resistance.

Figure 5. The frequency of predicted high-impact consequences in the five beta-tubulin
genes present in natural populations of *C. elegans*, *C. briggsae*, and *C. tropicalis*
The frequency of SNVs present in natural populations of *C. elegans* (n = 611), *C. briggsae* (n =
641), and *C. tropicalis* (n = 518) (y-axis) are shown by their predicted consequence in each beta-
tubulin gene (x-axis). The total number of isotype reference strains with a given predicted
consequence are displayed on top of each bar plot.
553

Second, we examined the geographic distribution of strains carrying high-impact variants

in BEN-1, TBB-1, and TBB-2 to identify if beta-tubulin variants were associated with geography.

Variants in BEN-1 were found globally across the three *Caenorhabditis* species (**Fig. 6A**), with no

discernible geographic pattern. In *C. elegans*, variants in BEN-1 were found in more recent clades

(*i.e.*, strains that experienced recent selective sweeps) (Andersen et al. 2012; Zhang, Mostad,

and Andersen 2021) (**Fig. 6B**). *Cel*-BEN-1 variants in swept clades suggest that these mutations

arose as relatively recent evolutionary events and might be maintained in response to BZ-like

compounds in the natural niche. By contrast, in *C. briggsae*, variants in BEN-1 were distributed

throughout the species tree and found on more ancestral branches (*i.e.*, earlier diverged lineages

in the species) (**Fig. 6C**). However, it is unlikely that variants in *Cbr*-BEN-1 emerged from

ancestral evolutionary events, because none of the variants identified were shared across strains.

For *C. tropicalis*, the limited number of variants in BEN-1 precludes any definitive conclusions

regarding their evolutionary patterns (**Fig. 6D**). Finally, because few variants are found in TBB-1

and TBB-2 in *C. briggsae* and *C. tropicalis*, we cannot identify the evolutionary patterns of BZ

resistance in these genes (**S13 and S14 Figures**).

Finally, as substrates harbor distinct microbial communities that can influence the

evolution of BZ resistance alleles, we determined if strains carrying a high-impact variant in a

beta-tubulin gene were associated with specific substrates. We performed substrate enrichment

analysis to assess enrichment between 12 substrates and all strains in the three *Caenorhabditis*

species (**Table S8**). However, no significant enrichment was observed between a high-impact

variant in a beta-tubulin gene and any given substrate (Fisher's Exact Test, *p*=1) (**S15 Figure**).

Because no geographic or substrate enrichment was observed, evolutionary pressures driving

beta-tubulin variation are likely not strongly tied to substrate. Instead, it is likely that species-

specific evolutionary trajectories (*e.g.*, selective pressures or gene family redundancy) shape

differential susceptibility to BZs.

579
Figure 6. The global distribution of *Caenorhabditis* strains that contain predicted high-
impact variation in BEN-1
(A) Each point corresponds to the sampling location of an individual *C. elegans* (orange),
C. briggsae (green), or *C. tropicalis* (blue) isotype reference strain with a predicted high-impact
consequence in BEN-1. A genome-wide phylogeny of **(B)** 611 *C. elegans*, **(C)** 641 *C. briggsae*,
and **(D)** 518 *C. tropicalis* isotype reference strains where each point denotes an isotype reference
strain with a predicted high-impact consequence in BEN-1 is shown.

4. DISCUSSION

4.1 A lack of repeated evolution of beta-tubulin mediated BZ resistance across

***Caenorhabditis* nematodes**

This study provides new insights into the mechanisms of ABZ resistance across three

Caenorhabditis species. No comprehensive survey of variation in the beta-tubulin gene family or

BZ response had been conducted across the three *Caenorhabditis* species to date. First, we

found that each *Caenorhabditis* species had unique predicted high-impact beta-tubulin alleles that

were not shared among the three species. Next, with an expanded sampling of *C. elegans*, we

confirmed that a heterogeneous set of rare high-impact alleles in *ben-1* were associated with ABZ

resistance (Hahnel et al. 2018; Dilks et al. 2021). Additionally, we identified a heterogenous set

of rare high-impact alleles in *Cbr-ben-1*, but only two (W21stop and Q134H) were associated with

ABZ resistance. The lack of ABZ resistance associated with *Cbr-ben-1* alleles suggested that

other beta-tubulin genes might play a role in BZ resistance. Unlike in *C. elegans*, we identified

high-impact alleles in *Cbr-tbb-1* and *Cbr-tbb-2*, but alleles in these genes were not associated

with ABZ resistance. Finally, although we identified high-impact alleles in *ben-1*, *tbb-1*, and *tbb-2*

in *C. tropicalis*, we did not observe ABZ resistance. The absence of repeated evolution of ABZ

resistance linked to variation in beta-tubulin genes across the three *Caenorhabditis* species

suggests that each species either employs distinct strategies to overcome ABZ exposure,

possibly arising from distinct ecological niches with unique selection pressures, or they are not

exposed to BZ compounds in the natural niche.

4.2 What drives the evolution of BZ resistance across *Caenorhabditis* species?

The lack of shared resistance alleles among the three species highlights the complexity

of BZ resistance. Across clade V nematodes, BZ resistance alleles come in two types (Collins et

al. 2024). First, when redundant beta-tubulin genes are present, LoF alleles of a beta-tubulin gene

can cause resistance. Second, without redundant beta-tubulin genes, LoF alleles are

hypothesized to reduce fitness or even cause lethality, so only alleles that alter BZ binding cause

resistance. Across *Caenorhabditis* nematode species, either type could be present so additional

criteria must be assessed to understand the evolution of BZ resistance. We must clearly

determine (1) the specific role of each beta-tubulin in ABZ resistance, (2) the tissue-specific

expression patterns of beta-tubulins, (3) the unique selective pressures acting on each

Caenorhabditis species, and (4) non-beta-tubulin genes involved in the ABZ response.

In *C. elegans*, the contribution of each beta-tubulin gene to BZ response has been

characterized in a controlled genetic background (*i.e.*, LoF beta-tubulin alleles in the N2 strain

background) (Collins et al. 2024). Similarly, we must first definitively identify the role each beta-

tubulin gene plays in BZ resistance in *C. briggsae* and *C. tropicalis*. CRISPR-Cas9 genome

editing should be used to delete each beta-tubulin gene in the reference strain background of

each species to determine the gene's contribution to resistance. Because the reference strains

for *C. briggsae* (AF16 and QX1410) and *C. tropicalis* (NIC58) are sensitive to ABZ, a resistance

phenotype following gene deletion would confirm the role of the gene in ABZ resistance.

Additionally, it is possible that disrupting the function of *tbb-1* and *tbb-2* in *C. briggsae* and

C. tropicalis could be highly detrimental to animal development, as seen in *C. elegans* (Collins et

al. 2024). Altogether, the creation of LoF alleles will allow us to define the role each gene plays

on nematode fitness and BZ response, defining the likelihood that BZ resistance will evolve

through naturally occurring LoF alleles.

In *C. elegans*, significant BZ susceptibility was identified in animals that express *ben-1* in

cholinergic neurons, suggesting that *ben-1* function in this cell type underlies susceptibility to BZ

(Gibson, Ness-Cohn, and Andersen 2022). The identification of beta-tubulin specific expression

improves our understanding of the cellular mode of action of BZs in *C. elegans*. Similarly, we must

characterize the temporal and tissue-specific expression patterns of beta-tubulins in *C. briggsae*

and *C. tropicalis*. Tissue-specific differences in beta-tubulin expression among the three

Caenorhabditis species could influence susceptibility and the evolution of BZ resistance. To

compare the expression patterns of *ben-1*, *tbb-1*, and *tbb-2*, we analyzed a publicly available

dataset that estimated the conservation of expression between *C. elegans* and *C. briggsae*

orthologs in homologous cell types (Large et al. 2024). The expression patterns of *ben-1* (Jensen-

Shannon Distance of expression [JSDgene] = 0.21), *tbb-1* (JSDgene = 0.24), and *tbb-2* (JSDgene

= 0.27) were highly conserved between *C. elegans* and *C. briggsae* (Large et al. 2024), indicating

that these genes maintain similar transcriptional profiles across species. Lower JSD values

indicate higher conservation, suggesting that beta-tubulin expression is largely preserved in

homologous cells, which might imply that the beta-tubulins have a conserved functional role in BZ

response. However, the homologous cells are not cholinergic neurons, so conserved activities in

this specific neuronal cell type might be divergent.

However, some neuronal cell-types (*e.g.*, PVQ, AVD, AVJ) expressed *Cel-ben-1* at higher

levels than in the *Cbr-ben-1*, whereas two cell types, ADL and interior arcade cells, showed higher

ben-1 expression in *C. briggsae* than in *C. elegans*. (Large et al. 2024). Therefore, differences in

cell-specific expression patterns of beta-tubulins might influence BZ susceptibility in each species

and should be further investigated. One approach involves introducing *ben-1* into a *C. briggsae*

ben-1 knockout strain background using transgenesis, where multi-copy arrays express *ben-1* in

specific tissues, as previously done in *C. elegans* (Gibson, Ness-Cohn, and Andersen 2022).

Plasmids containing the coding sequence of *ben-1* are fused to tissue-specific promoters to form

extrachromosomal arrays in transgenic animals, allowing precise tissue-specific expression.

Investigating beta-tubulin expression patterns in specific tissues will clarify the role of each beta-

tubulin and identify the tissues that BZs target in *C. briggsae*. The same approach can be applied

to other beta-tubulin genes in *C. briggsae* and *C. tropicalis*. Additionally, understanding the tissue-

specific roles of each beta-tubulin across *Caenorhabditis* species can inform strategies for

managing resistance in parasitic nematodes by identifying key sites of BZ action. For instance, if

ben-1 function in specific tissues is critical for BZ sensitivity, it might be possible to identify similar

tissues in parasitic species where beta-tubulins play a pivotal role in BZ resistance. This

information could lead to targeted approaches for enhancing drug efficacy, such as developing

treatments that specifically affect these tissues or identifying novel drug targets within them to

combat resistance.

Next, the lack of shared resistance variants across *C. elegans, C. briggsae*, and

C. tropicalis suggests that each species experiences distinct selective pressures in the evolution

of BZ resistance. Although *Caenorhabditis* species share overlapping niches, differences in

microbial communities that produce natural BZs and proximity to synthetic BZs in agriculture could

lead to differences in BZ response. Despite the fact that our substrate enrichment tests indicate

no significant association between the substrate where a strain was collected and the probability

of having a predicted resistance variant in a beta-tubulin gene, our broad substrate categories

and small number of BZ resistant *C. briggsae* and *C. tropicalis* strains might obscure finer-scale

ecological patterns. For instance, the microbial community composition on each substrate, such

as the presence or absence of BZ producing bacterial species, could shape the evolution of BZ

resistance. Future studies characterizing microbial communities associated with each substrate

might clarify the selective pressures on *Caenorhabditis* nematodes.

Finally, although beta-tubulins are the primary targets of BZs in Clade V nematodes, other

mechanisms, such as variation in detoxification pathways or drug efflux transporters, could drive

the evolution of BZ resistance outside the beta-tubulin gene family. In *C. elegans*, variation in BZ

responses among wild strains has been leveraged to identify alleles in a cytochrome P450 that

cause BZ resistance independent of *Cel-ben-1* (Collins et al. 2025). Similarly, variation in ABZ

response among *C. briggsae* and *C. tropicalis* strains can be investigated with quantitative

genetic mapping techniques to discover sources of variation in ABZ response among wild strains.

Expanding genomic resources to better characterize additional sources of variation in ABZ

response among wild strains (*e.g.,* SVs, transcriptomic variation) is critical to disentangle the lack

of repeated evolution in BZ resistance.

DATA AVAILABILITY STATEMENT

All code and data used to replicate the data analysis and figures are available on GitHub

at: https://github.com/AndersenLab/ce_cb_ct_betatubulin. Table S1 contains all beta-tubulin

transcript IDs in the three *Caenorhabditis* species. Tables S2, S3, and S4 contain the list of *C.*

elegans, *C. briggsae*, and *C. tropicalis* isotype strains, collection location, substrate type, and their

beta-tubulin variant status, respectively. Table S5 contains the manual curation of SVs found in

beta-tubulin genes. Table S6 contains all *C. briggsae* and *C. tropicalis* isotype strains with beta-

tubulin variants (*i.e.*, predicted resistant strains) and their corresponding predicted susceptible

strains. Table S7 contains the BLOSUM and Granthum scores for amino acid changes in the

beta-tubulin genes in the three *Caenorhabditis* species. Table S8 contains results from the

substrate enrichment analysis.

FUNDING SOURCES

Amanda O. Shaver was funded by the National Institutes of Health grant F32AI181342. Ryan

McKeown was supported by the National Institutes of Health Training Grant (T32 GM008449)

through Northwestern University's Biotechnology Training Program. This work was supported by

the National Institutes of Health grant R01AI153088 to ECA.

CRediT AUTHOR STATEMENT

Conceptualization: ECA

Data curation: AOS, RM

Formal analysis: AOS, RM

Funding acquisition: ECA

Investigation: AOS, JMRO

Methodology: AOS, RM

Project administration: ECA

Resources: ECA

Software: AOS, RM

Supervision: ECA

Validation: AOS

Visualization: AOS, RM

Writing - original draft: AOS

Writing - reviewing & editing: AOS, RM, ECA

DECLARATION OF COMPETING INTERESTS

The authors have declared that no competing interests exist.

ACKNOWLEDGEMENTS

We would like to thank members of the Andersen laboratory for their feedback and helpful

comments on this manuscript. We thank the *Caenorhabditis* Natural Diversity Resource (NSF

Capacity grant 2224885) for providing us with strains for this study.

REFERENCES

Andersen, Erik C., Joshua S. Bloom, Justin P. Gerke, and Leonid Kruglyak. 2014. "A Variant in
the Neuropeptide Receptor Npr-1 Is a Major Determinant of Caenorhabditis Elegans
Growth and Physiology." *PLoS Genetics* 10 (2): e1004156.
Andersen, Erik C., Justin P. Gerke, Joshua A. Shapiro, Jonathan R. Crissman, Rajarshi Ghosh,
Joshua S. Bloom, Marie-Anne Félix, and Leonid Kruglyak. 2012. "Chromosome-Scale
Selective Sweeps Shape Caenorhabditis Elegans Genomic Diversity." *Nature Genetics* 44
(3): 285–90.
Andersen, Erik C., Tyler C. Shimko, Jonathan R. Crissman, Rajarshi Ghosh, Joshua S. Bloom,
Hannah S. Seidel, Justin P. Gerke, and Leonid Kruglyak. 2015. "A Powerful New
Quantitative Genetics Platform, Combining Caenorhabditis Elegans High-Throughput
Fitness Assays with a Large Collection of Recombinant Strains." *G3* 5 (5): 911–20.
Avramenko, Russell W., Elizabeth M. Redman, Lynsey Melville, Yvonne Bartley, Janneke Wit,
Camila Queiroz, Dave J. Bartley, and John S. Gilleard. 2019. "Deep Amplicon Sequencing
as a Powerful New Tool to Screen for Sequence Polymorphisms Associated with
Anthelmintic Resistance in Parasitic Nematode Populations." *International Journal for
Parasitology* 49 (1): 13–26.
Camacho, Christiam, George Coulouris, Vahram Avagyan, Ning Ma, Jason Papadopoulos,
Kevin Bealer, and Thomas L. Madden. 2009. "BLAST+: Architecture and Applications."
BMC Bioinformatics 10 (1): 421.
Caro, Hugo, Sulyvan Dollin, Anne Biton, Bryan Brancotte, Dimitri Desvillechabrol, Yoann
Dufresne, Blaise Li, et al. 2023. "BioConvert: A Comprehensive Format Converter for Life
Sciences." *NAR Genomics and Bioinformatics* 5 (3): lqad074.
Cerca, José. 2023. "Understanding Natural Selection and Similarity: Convergent, Parallel and
Repeated Evolution." *Molecular Ecology* 32 (20): 5451–62.
Collins, J. B., Clayton M. Dilks, Steffen R. Hahnel, Briana Rodriguez, Bennett W. Fox, Elizabeth
Redman, Jingfang Yu, et al. 2025. "Naturally Occurring Variation in a Cytochrome P450
Modifies Thiabendazole Responses Independently of Beta-Tubulin." *PLoS Pathogens* 21
(1): e1012602.
Collins, J. B., Skyler A. Stone, Emily J. Koury, Anna G. Paredes, Fiona Shao, Crystal Lovato,
Michael Chen, et al. 2024. "Quantitative Tests of Albendazole Resistance in Caenorhabditis
Elegans Beta-Tubulin Mutants." *International Journal for Parasitology, Drugs and Drug
Resistance* 25 (August):100556.
Crombie, Timothy A., Ryan McKeown, Nicolas D. Moya, Kathryn S. Evans, Samuel J.
Widmayer, Vincent LaGrassa, Natalie Roman, et al. 2023. "CaeNDR, the Caenorhabditis
Natural Diversity Resource." *Nucleic Acids Research*, October.
https://doi.org/10.1093/nar/gkad887.
Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O.
Pollard, Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools."
GigaScience 10 (2). https://doi.org/10.1093/gigascience/giab008.
Dilks, Clayton M., Steffen R. Hahnel, Qicong Sheng, Lijiang Long, Patrick T. McGrath, and Erik
C. Andersen. 2020. "Quantitative Benzimidazole Resistance and Fitness Effects of
Parasitic Nematode Beta-Tubulin Alleles." *International Journal for Parasitology, Drugs and
Drug Resistance* 14 (December):28–36.
Dilks, Clayton M., Emily J. Koury, Claire M. Buchanan, and Erik C. Andersen. 2021. "Newly
Identified Parasitic Nematode Beta-Tubulin Alleles Confer Resistance to Benzimidazoles."
International Journal for Parasitology, Drugs and Drug Resistance 17 (December):168–75.
Dingerdissen, Hayley M., John Torcivia-Rodriguez, Yu Hu, Ting-Chia Chang, Raja Mazumder,
and Robel Kahsay. 2018. "BioMuta and BioXpress: Mutation and Expression
Knowledgebases for Cancer Biomarker Discovery." *Nucleic Acids Research* 46 (D1):

D1128–36.
Ghisi, Marc, Ronald Kaminsky, and Pascal Mäser. 2007. "Phenotyping and Genotyping of
Haemonchus Contortus Isolates Reveals a New Putative Candidate Mutation for
Benzimidazole Resistance in Nematodes." *Veterinary Parasitology* 144 (3-4): 313–20.
Gibson, Sophia B., Elan Ness-Cohn, and Erik C. Andersen. 2022. "Benzimidazoles Cause
Lethality by Inhibiting the Function of Caenorhabditis Elegans Neuronal Beta-Tubulin."
International Journal for Parasitology, Drugs and Drug Resistance 20 (December):89–96.
Hahnel, Dilks, Heisler, and Andersen. 2020. "Caenorhabditis Elegans in Anthelmintic research–
Old Model, New Perspectives." *International Journal for Educational and Vocational
Guidance*. https://www.sciencedirect.com/science/article/pii/S2211320720300312.
Hahnel, Steffen R., Stefan Zdraljevic, Briana C. Rodriguez, Yuehui Zhao, Patrick T. McGrath,
and Erik C. Andersen. 2018. "Extreme Allelic Heterogeneity at a Caenorhabditis Elegans
Beta-Tubulin Locus Explains Natural Resistance to Benzimidazoles." *PLoS Pathogens* 14
(10): e1007226.
Ireland, C. M., K. Gull, W. E. Gutteridge, and C. I. Pogson. 1979. "The Interaction of
Benzimidazole Carbamates with Mammalian Microtobule Protein." *Biochemical
Pharmacology* 28 (17): 2680–82.
Kotze, A. C., J. S. Gilleard, S. R. Doyle, and R. K. Prichard. 2020. "Challenges and
Opportunities for the Adoption of Molecular Diagnostics for Anthelmintic Resistance."
International Journal of Forecasting.
https://www.sciencedirect.com/science/article/pii/S2211320720300476.
Kotze, A. C., and R. K. Prichard. 2016. "Anthelmintic Resistance in Haemonchus Contortus:
History, Mechanisms and Diagnosis." *Advances in Parasitology* 93 (March):397–428.
Kotze, Andrew C., Peter W. Hunt, Philip Skuce, Georg von Samson-Himmelstjerna, Richard J.
Martin, Heinz Sager, Jürgen Krücken, et al. 2014. "Recent Advances in Candidate-Gene
and Whole-Genome Approaches to the Discovery of Anthelmintic Resistance Markers and
the Description of Drug/receptor Interactions." *International Journal for Parasitology, Drugs
and Drug Resistance* 4 (3): 164–84.
Krücken, Jürgen, Kira Fraundorfer, Jean Claude Mugisha, Sabrina Ramünke, Kevin C. Sifft,
Dominik Geus, Felix Habarugira, et al. 2017. "Reduced Efficacy of Albendazole against
Ascaris Lumbricoides in Rwandan Schoolchildren." *International Journal for Parasitology,
Drugs and Drug Resistance* 7 (3): 262–71.
Kwa, M. S., J. G. Veenstra, and M. H. Roos. 1994. "Benzimidazole Resistance in Haemonchus
Contortus Is Correlated with a Conserved Mutation at Amino Acid 200 in Beta-Tubulin
Isotype 1." *Molecular and Biochemical Parasitology* 63 (2): 299–303.
Laclette, J. P., G. Guerra, and C. Zetina. 1980. "Inhibition of Tubulin Polymerization by
Mebendazole." *Biochemical and Biophysical Research Communications* 92 (2): 417–23.
Large, Christopher R. L., Rupa Khanal, Ladeana Hillier, Chau Huynh, Connor Kubo, Junhyong
Kim, Robert H. Waterston, and John I. Murray. 2024. "Lineage-Resolved Analysis of
Embryonic Gene Expression Evolution in C. Elegans and C. Briggsae." *bioRxiv*.
https://doi.org/10.1101/2024.02.03.578695.
Lesack, Kyle, Grace M. Mariene, Erik C. Andersen, and James D. Wasmuth. 2022. "Different
Structural Variant Prediction Tools Yield Considerably Different Results in Caenorhabditis
Elegans." *PloS One* 17 (12): e0278424.
Lourdes Mottier, Maria de, and Roger K. Prichard. 2008. "Genetic Analysis of a Relationship
between Macrocyclic Lactone and Benzimidazole Anthelmintic Selection on Haemonchus
Contortus." *Pharmacogenetics and Genomics* 18 (2): 129–40.
Lubega, G. W., and R. K. Prichard. 1990. "Specific Interaction of Benzimidazole Anthelmintics
with Tubulin: High-Affinity Binding and Benzimidazole Resistance in Haemonchus
Contortus." *Molecular and Biochemical Parasitology* 38 (2): 221–32.
Luke, M., John Noble, Lewis Yuen, Nicolas Stevens, and Riaad Moya. 2021. "Selfing Is the

Safest Sex for Caenorhabditis Tropicalis." *eLife* 10.

Mariene, Grace M., and James D. Wasmuth. 2025. "Genome Assembly Variation and Its
Implications for Gene Discovery in Nematodes." *International Journal for Parasitology*,
January. https://doi.org/10.1016/j.ijpara.2025.01.004.

Mohammedsalih, Khalid M., Jürgen Krücken, Amna Khalafalla, Ahmed Bashar, Fathel-Rahman
Juma, Adam Abakar, Abdalhakaim A. H. Abdalmalaik, Gerald Coles, and Georg von
Samson-Himmelstjerna. 2020. "New Codon 198 β-Tubulin Polymorphisms in Highly
Benzimidazole Resistant Haemonchus Contortus from Goats in Three Different States in
Sudan." *Parasites & Vectors* 13 (1): 114.

Moya, Nicolas D., Lewis Stevens, Isabella R. Miller, Chloe E. Sokol, Joseph L. Galindo,
Alexandra D. Bardas, Edward S. H. Koh, et al. 2023. "Novel and Improved Caenorhabditis
Briggsae Gene Models Generated by Community Curation." *BMC Genomics* 24 (1): 486.

Neff, N. F., J. H. Thomas, P. Grisafi, and D. Botstein. 1983. "Isolation of the Beta-Tubulin Gene
from Yeast and Demonstration of Its Essential Function in Vivo." *Cell* 33 (1): 211–19.

Noble, Luke M., John Yuen, Lewis Stevens, Nicolas Moya, Riaad Persaud, Marc Moscatelli,
Jacqueline L. Jackson, et al. 2021. "Selfing Is the Safest Sex for Caenorhabditis Tropicalis."
eLife. https://doi.org/10.7554/elife.62587.

Nyaanga, Joy, Timothy A. Crombie, Samuel J. Widmayer, and Erik C. Andersen. 2021.
"easyXpress: An R Package to Analyze and Visualize High-Throughput C. Elegans
Microscopy Data Generated Using CellProfiler." *PloS One* 16 (8): e0252000.

Ortiz, Edgardo M. 2019. *vcf2phylip v2.0: Convert a VCF Matrix into Several Matrix Formats for
Phylogenetic Analysis*. Zenodo. https://doi.org/10.5281/ZENODO.2540861.

Pertea, Geo, and Mihaela Pertea. 2020. "GFF Utilities: GffRead and GffCompare."
F1000Research 9 (April):304.

Qiu, Jianbo, Jianqiang Xu, Junjie Yu, Chaowei Bi, Changjun Chen, and Mingguo Zhou. 2011.
"Localisation of the Benzimidazole Fungicide Binding Site of Gibberella Zeae β2-Tubulin
Studied by Site-Directed Mutagenesis." *Pest Management Science* 67 (2): 191–98.

Rausch, Tobias, Thomas Zichner, Andreas Schlattl, Adrian M. Stütz, Vladimir Benes, and Jan
O. Korbel. 2012. "DELLY: Structural Variant Discovery by Integrated Paired-End and Split-
Read Analysis." *Bioinformatics* 28 (18): i333–39.

Roos, M. H., M. S. G. Kwa, and W. N. Grant. 1995. "New Genetic and Practical Implications of
Selection for Anthelmintic Resistance in Parasitic Nematodes." *Parasitology Today
(Personal Ed.)* 11 (4): 148–50.

Saitou, N., and M. Nei. 1987. "The Neighbor-Joining Method: A New Method for Reconstructing
Phylogenetic Trees." *Molecular Biology and Evolution* 4 (4): 406–25.

Shaver, Amanda O., Isabella R. Miller, Etta S. Schaye, Nicolas D. Moya, J. B. Collins, Janneke
Wit, Alyssa H. Blanco, et al. 2024. "Quantifying the Fitness Effects of Resistance Alleles
with and without Anthelmintic Selection Pressure Using Caenorhabditis Elegans." *PLoS
Pathogens* 20 (5): e1012245.

Shaver, Amanda O., Janneke Wit, Clayton M. Dilks, Timothy A. Crombie, Hanchen Li, Raffi V.
Aroian, and Erik C. Andersen. 2023. "Variation in Anthelmintic Responses Are Driven by
Genetic Differences among Diverse C. Elegans Wild Strains." *PLoS Pathogens* 19 (4):
e1011285.

Silvestre, Anne, and Jacques Cabaret. 2002. "Mutation in Position 167 of Isotype 1 Beta-Tubulin
Gene of Trichostrongylid Nematodes: Role in Benzimidazole Resistance?" *Molecular and
Biochemical Parasitology* 120 (2): 297–300.

Sternberg, Paul W., Kimberly Van Auken, Qinghua Wang, Adam Wright, Karen Yook,
Magdalena Zarowiecki, Valerio Arnaboldi, et al. 2024. "WormBase 2024: Status and
Transitioning to Alliance Infrastructure." *Genetics* 227 (1).
https://doi.org/10.1093/genetics/iyae050.

Stevens, Lewis, Nicolas D. Moya, Robyn E. Tanny, Sophia B. Gibson, Alan Tracey, Huimin Na,

Rojin Chitrakar, et al. 2022. "Chromosome-Level Reference Genomes for Two Strains of
Caenorhabditis Briggsae: An Improved Platform for Comparative Genomics." *Genome*
Biology and Evolution 14 (4). https://doi.org/10.1093/gbe/evac042.
Theodorides, V. J., G. C. Scott, and M. S. Lademan. 1970. "Strains of Haemonchus Contortus
Resistant against Benzimidazole Anthelmintics." *American Journal of Veterinary Research*
31 (5): 859–63.
Venkatesan, Abhinaya, Pablo D. Jimenez Castro, Arianna Morosetti, Hannah Horvath, Rebecca
Chen, Elizabeth Redman, Kayla Dunn, et al. 2023. "Molecular Evidence of Widespread
Benzimidazole Drug Resistance in Ancylostoma Caninum from Domestic Dogs throughout
the USA and Discovery of a Novel β-Tubulin Benzimidazole Resistance Mutation." *PLoS*
Pathogens 19 (3): e1011146.
Wählby, Carolina, Lee Kamentsky, Zihan H. Liu, Tammy Riklin-Raviv, Annie L. Conery, Eyleen
J. O'Rourke, Katherine L. Sokolnicki, et al. 2012. "An Image Analysis Toolbox for High-
Throughput C. Elegans Assays." *Nature Methods* 9 (7): 714–16.
Widmayer, Samuel J., Timothy A. Crombie, Joy N. Nyaanga, Kathryn S. Evans, and Erik C.
Andersen. 2022. "C. Elegans Toxicant Responses Vary among Genetically Diverse
Individuals." *Toxicology*, August, 153292.
Yu, Guangchuang. 2020. "Using Ggtree to Visualize Data on Tree-Like Structures." *Et Al*
[Current Protocols in Bioinformatics] 69 (1): e96.
Zamanian, Mostafa, Daniel E. Cook, Stefan Zdraljevic, Shannon C. Brady, Daehan Lee, Junho
Lee, and Erik C. Andersen. 2018. "Discovery of Genomic Intervals That Underlie Nematode
Responses to Benzimidazoles." *PLoS Neglected Tropical Diseases* 12 (3): e0006368.
Zhang, Gaotian, Jake D. Mostad, and Erik C. Andersen. 2021. "Natural Variation in Fecundity Is
Correlated with Species-Wide Levels of Divergence in Caenorhabditis Elegans." *G3* 11 (8).
https://doi.org/10.1093/g3journal/jkab168.
Zhang, Gaotian, Nicole M. Roberto, Daehan Lee, Steffen R. Hahnel, and Erik C. Andersen.
2022. "The Impact of Species-Wide Gene Expression Variation on Caenorhabditis Elegans
Complex Traits." *Nature Communications* 13 (1): 3462.

<u>SUPPORTING INFORMATION</u>

Supplemental Table 1. Beta-tubulin transcript IDs
Supplemental Table 2. *C. elegans* isotype variant table
Supplemental Table 3. *C. briggsae* isotype variant table
Supplemental Table 4. *C. tropicalis* isotype variant table
Supplemental Table 5. Manual curation of SVs
Supplemental Table 6. *C. briggsae* and *C. tropicalis* strain pairs
Supplemental Table 7. BLOSUM and Grantham scores for amino acid changes in beta-tubulin
genes in the three *Caenorhabditis* species
Supplemental Table 8. Location and substrate where each isotype reference strain was
collected

 SUPPLEMENTAL FIGURES

**Supplemental Figure 1. The relationship between *ben-1* expression levels and albendazole
response in *C. elegans* strains**
(A) Scatterplot of the relationship between *ben-1* expression levels and normalized albendazole
(ABZ) response across *C. elegans* wild strains. Each point represents a strain phenotyped for
ABZ response in previous publications (Hahnel *et al.*, 2018; Shaver *et al.*, 2024) with *ben-1*
expression data (Zhang et al. 2022)The *ben-1* expression level measured in transcripts per million
(TPM) is displayed on the x-axis. The normalized ABZ response values adjusted for assay-
specific effects are displayed on the y-axis. The gray line represents the linear regression fit
between *ben-1* expression and normalized response (R^2 = 0.34, *p*-value = 5.16e-18), with the

linear model's coefficient of determination (R^2). Data points are colored based on the predicted
functional consequence of their *ben-1* allele (*e.g.*, large structural variant (SV), Frame Altering,
Missense substitution, Disrupted Start/Stop sequence, No high-impact variant, Low *ben-1*
expression). A horizontal red line represents a resistance threshold calculated from the
normalized phenotypes of all strains from previous publications set at 75% of the reference strain
N2. Strains with normalized responses above this threshold are considered resistant to ABZ. **(B)**
Boxplots of *ben-1* expression levels among individuals grouped by the predicted functional
consequences of their *ben-1* alleles. Each point represents the *ben-1* expression level of an
individual within each group. We tested for statistically significant differences in the expression
between each consequence type and strains without a high-impact *ben-1* allele with an unpaired
Wilcoxon test. Significance levels are indicated by symbols: '*' ($p < 0.05$), '**' ($p < 0.01$), '***' ($p <$
0.001), '****' ($p < 0.0001$).

Supplemental Figure 2. The relationship between *C. elegans* beta-tubulin expression and albendazole response

Scatterplot of the relationship between **(A)** *tbb-1* and **(B)** *tbb-2* expression levels and normalized albendazole (ABZ) response across *C. elegans* wild strains. Each point represents a strain phenotyped for ABZ response in previous publications (Hahnel *et al.*, 2018; Shaver *et al.*, 2024) with *tbb-1* and *tbb-2* expression data (Zhang et al. 2022)The *tbb-1* and *tbb-2* expression levels measured in transcripts per million (TPM) are displayed on the x-axis. The normalized ABZ response values adjusted for assay-specific effects are displayed on the y-axis. The gray line represents the linear regression fit between beta-tubulin gene expression and normalized response, with the linear model's coefficient of determination (R^2). Data points are colored based

on the predicted functional consequence of their *ben-1* alleles (*e.g.*, large structural variant (SV),
Frame Altering, Missense substitution, Disrupted Start/Stop sequence, No high-impact variant,
Low *ben-1* expression). A horizontal red line represents a resistance threshold calculated from
the normalized phenotypes of all strains from previous publications set at 75% of the reference
strain N2. Strains with normalized responses above this threshold are considered resistant to
ABZ.

Beta-tubulin variant ○ False ● True

0.02

Supplemental Figure 3. *C. briggsae* **species tree highlighting isotype reference strains**
tested for ABZ resistance. *C. briggsae* strains with predicted high-impact variants are colored
red, and strains with no predicted variants in beta-tubulin genes are colored gray.

Supplemental Figure 4. ***C. tropicalis* species tree highlighting isotype reference strains**
tested for ABZ resistance. *C. tropicalis* strains with predicted high-impact variants are colored
red, and strains with no predicted variants in beta-tubulin genes are colored gray.

964
Supplemental Figure 5. Dose-response curves for *C. briggsae* strains in albendazole
Normalized animal lengths (y-axis) are plotted for each strain as a function of the dose of
albendazole (ABZ) in the high-throughput development assay (x-axis). Strains are denoted by
color. Lines extending vertically from points represent the standard deviation from the mean
response. Statistical normalization of animal lengths is described in *Methods*. *C. elegans* strains
N2 and ECA882 were added for ABZ-susceptible and ABZ-resistant controls, respectively.
971

972
973
974 **Supplemental Figure 6. Dose-response curves for *C. tropicalis* strains in albendazole**
975 Normalized animal lengths (y-axis) are plotted for each strain as a function of the dose of
976 albendazole (ABZ) in the high-throughput development assay (x-axis). Strains are denoted by
977 color. Lines extending vertically from points represent the standard deviation from the mean
978 response. Statistical normalization of animal lengths is described in *Methods*.

Supplemental Figure 7. High-throughput assays for each *C. elegans* strain in control conditions

Median animal length values from populations of nematodes grown in DMSO are shown on the y-axis. Each point represents the median animal length from a well containing approximately 5-30 animals. Data are shown as Tukey box plots with the median as a solid horizontal line, the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within a 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile.

Supplemental Figure 8. High-throughput assays for each *C. briggsae* strain in control conditions.

Median animal length values from populations of nematodes grown in DMSO are shown on the y-axis. Each point represents the median animal length from a well containing approximately 5-30 animals. Data are shown as Tukey box plots with the median as a solid horizontal line, the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within a 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status.

999
Supplemental Figure 9. High-throughput assays for each *C. tropicalis* strain in control conditions
Median animal length values from populations of nematodes grown in DMSO are shown on the y-axis. Each point represents the median animal length from a well containing approximately 5-30 animals. Data are shown as Tukey box plots with the median as a solid horizontal line, the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within a 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. Strains are colored by beta-tubulin variant status.

Supplemental Figure 10. Relationship between missense substitutions in BEN-1 and
albendazole response in *C. elegans* strains. Scatterplots show the relationship between
normalized median animal length (y-axis) and three amino acid substitution scoring metrics (x-
axis): BLOSUM (R^2 = 0.97, *p*-value = 0.11), Grantham (R^2 = 0.39, *p*-value = 0.57), and percent
protein (R^2 = 0.1, *p*-value = 0.8). Each point represents a *C. elegans* strain with a missense
substitution in BEN-1 that was **(A)** phenotyped for ABZ response inprevious assays (Hahnel et
al. 2018; Shaver et al. 2024) or **(B)** in the high-throughput assays performed for this manuscript
are reported. Gray lines indicate the linear regression fit for these models.

Supplemental Figure 11. Relationship between missense substitutions in BEN-1, TBB-1,
TBB-2 and albendazole response in *C. briggsae* strains. Scatterplots show the relationship
between normalized median animal length (y-axis) and three amino acid substitution scoring
metrics (x-axis): BLOSUM (R^2 = 0.0, *p*-value = 0.85), Grantham (R^2 = 0.08, *p*-value = 0.28),
percent protein (R^2 = 0.01, *p*-value = 0.79). Each point represents a *C. briggsae* strain with a
missense substitution in either BEN-1, TBB-1, or TBB-2. Gray lines indicate the linear regression
fit, and the R^2 and *p*-values of these models are displayed in each panel

Supplemental Figure 12. Relationship between missense substitutions in BEN-1 or TBB-2
and albendazole response in *C. tropicalis* strains. Scatterplots show the relationship between
normalized median animal length (y-axis) and three amino acid substitution scoring metrics (x-
axis): BLOSUM (R^2 = 0.04, *p*-value = 0.87), Grantham (R^2 = 0.03, *p*-value = 0.89), percent protein
(R^2 = 0.77, *p*-value = 0.32). Each point represents a *C. tropicalis* strain with a missense
substitution in BEN-1 or TBB-2. Gray lines indicate the linear regression fit of these models.

C. briggsae   *C. tropicalis*

Supplemental Figure 13. The global distribution of *Caenorhabditis* strains that contain
predicted high-impact variation in *tbb-2*
(A) Each point corresponds to the sampling location of an individual *C. briggsae* or *C. tropicalis*
isotype reference strain with a predicted high-impact consequence in the gene *tbb-2*. A genome-
wide phylogeny of **(B)** 641 *C. briggsae* and **(C)** 518 *C. tropicalis* isotype reference strains where
each point denotes an isotype reference strain with a predicted high-impact consequence in *tbb-*
2.

C. briggsae

1037
Supplemental Figure 14. The global distribution of *Caenorhabditis* strains that contain
predicted high-impact variation in *tbb-1*
(A) Each point corresponds to the sampling location of an individual *C. briggsae* or *C. tropicalis*
isotype reference strain with a predicted high-impact consequence in the gene *tbb-1*. **(B)** A
genome-wide phylogeny of 641 *C. briggsae* isotype reference strains where each point denotes
an isotype reference strain with a predicted high-impact consequence in *tbb-1*. One isotype,
XZ1213 has a high-impact *tbb-1* variant, but sampling coordinates were not recorded. **(C)** A
genome-wide phylogeny of 518 *C. tropicalis* isotype reference strains where each point denotes
an isotype reference strain with a predicted high-impact consequence in *tbb-1*.

1047
Supplemental Figure 15. The proportion of strains with high-impact resistant variants in beta-tubulin genes and the substrates where those strains were found
The proportion of strains (y-axis) found in a given substrate (x-axis) are displayed. Strains with a high-impact variant in a beta-tubulin gene are colored salmon. Strains with no variants in a beta-tubulin gene are colored teal. The total number of strains isolated from a given substrate is displayed above each column. Moss and rotting wood were not included in the substrate enrichment analysis due to the small sample size. No significant relationship between beta-tubulin gene variant status and substrate were identified (Fisher's Exact Test, p=1).