Evaluating beta-tubulin variants as predictors of benzimidazole resistance across *Caenorhabditis* nematodes

Amanda O. Shaver[1]*, Ryan McKeown[2]*, Joyce M. Reyes Otero[3], J.B. Collins[1], Daniel W. Hogan[4], James S. Fraser[4], Stephen M. Dreyer[5], Erik J. Ragsdale[5], and Erik C. Andersen[1]

[1] Dept. of Biology, Johns Hopkins University, Mudd Hall 144, 3400 N. Charles Street, Baltimore, Maryland 21218
[2] Dept. of Molecular Biosciences, Northwestern University, 2205 Tech Dr, Evanston, Illinois 60208
[3] Dept. of Biology, University of Puerto Rico, CX33+QM7, San Juan, 00925, Puerto Rico t
[4] Dept. of Bioengineering and Therapeutic Sciences, University of California San Francisco, 600 16th St., Genentech Hall Room S472E, San Francisco, CA 94143
[5] Dept. of Biology, Indiana University, Biology Building, Rm. 142, 1001 East Third St., Bloomington, Indiana, 47405
* These authors contributed equally.

Corresponding author:
Erik C. Andersen
Johns Hopkins University
Department of Biology
3400 N. Charles St.
Bascom UTL 383
Baltimore, MD 21218
410-516-1282
erik.andersen@gmail.com

ORCIDs and emails:
Amanda O. Shaver: 0000-0002-2910-1505, amanda.shaver@jhu.edu, amandaoshaver@gmail.com
Ryan McKeown: 0000-0003-3570-2494, ryanmckeown2021@u.northwestern.edu
Joyce M. Reyes Otero: 0009-0002-5039-4434, joyce.reyes2@upr.edu
J.B. Collins: 0000-0003-0808-5216, jcolli95@jh.edu
Daniel W. Hogan: 0000-0003-3375-408X, Daniel.Hogan@ucsf.edu
James S. Fraser: 0000-0002-5080-2859, jfraser@fraserlab.com
Stephen Dreyer: 0000-0003-3900-6031, stdreyer@iu.edu
Erik J. Ragsdale: 0000-0002-9367-8552, ragsdale@iu.edu
Erik C. Andersen: 0000-0003-0229-9651, erik.andersen@gmail.com

Short title: Beta-tubulin mediated benzimidazole resistance

Keywords: *C. elegans*, *C. briggsae*, *C. tropicalis*, beta-tubulin, benzimidazole resistance

ABSTRACT

Benzimidazoles, a widely used class of anthelmintic drugs, target beta-tubulin, disrupt microtubule formation,

and delay nematode development. In parasitic nematodes, mutations in beta-tubulin genes are predicted to

inhibit benzimidazole binding and are associated with resistance. In the free-living nematode *Caenorhabditis*

elegans, loss-of-function mutations in the beta-tubulin gene *ben-1* cause benzimidazole resistance. Although

several beta-tubulin mutations serve as established markers of resistance, the prediction of the effects of novel

variants in different nematode species remains challenging. Here, we identified novel beta-tubulin variants

predicted to confer benzimidazole resistance across wild strains in three *Caenorhabditis* species: *C. elegans*,

Caenorhabditis briggsae, and *Caenorhabditis tropicalis*. The three *Caenorhabditis* species are experimentally

tractable, have characterized beta-tubulin gene complements, and defined natural niches, which allowed us to

identify variants in beta-tubulin genes and test which variants are associated with resistance. We hypothesized

that, if these species experienced similar selective pressures, they would evolve resistance to benzimidazoles

by mutations in a beta-tubulin gene (*tbb-1*, *tbb-2*, *mec-7*, *tbb-4*, and *ben-1*). In the three *Caenorhabditis* species,

we tested all strains harboring variants in the five conserved beta-tubulin genes for benzimidazole resistance. In

C. elegans, we found that a heterogeneous set of variants in *ben-1* were associated with resistance. By contrast,

only two variants in *C. briggsae ben-1* (W21stop and Q134H) were associated with resistance. *C. tropicalis* was

distinct from the other two species, where no strains with variants in any beta-tubulin gene were resistant. We

generated deletions of *ben-1* in *C. briggsae* and *C. tropicalis* and confirmed that loss of *ben-1* confers resistance

in both species. Our findings reveal species-specific patterns of beta-tubulin-mediated benzimidazole resistance

and emphasize that prediction of variants in beta-tubulin genes alone is not sufficient to predict resistance,

especially across diverse nematode species.

AUTHOR SUMMARY

Mutations in beta-tubulin genes have been associated with benzimidazole resistance across nematode species, yet predicting novel resistance variants remains challenging. Using wild strains from three *Caenorhabditis* species, we identified strains with variants in beta-tubulin genes and tested each strain for benzimidazole resistance. In *C. elegans,* a diverse set of loss-of-function variants in *ben-1* were associated with resistance. Whereas in *C. briggsae*, only two *ben-1* alleles were associated with resistance, suggesting selection acts differently in this species despite a similar niche as *C. elegans*. *C. tropicalis* had no strains with beta-tubulin variants that were resistant. Our results highlight species-specific patterns of benzimidazole resistance.

INTRODUCTION

Global control of parasitic nematode infections relies on the efficacy of a small arsenal of anthelmintic drugs, including benzimidazoles (BZs) [1]. BZs are a widely used class of anthelmintic drugs that inhibit the polymerization of microtubules [2–4] and delay nematode development [5]. Although BZs are essential to human and veterinary health, resistance is prominent in parasitic nematode populations [6,7]. In clade V nematodes (*e.g.*, *Ancylostoma caninum*, *Ancylostoma duodenale*, *Caenorhabditis elegans*, *Haemonchus contortus*, *Necator americanus*, *Teladorsagia circumcincta*, and *Trichostrongylus colubriformis*), BZ resistance is associated with mutations in beta-tubulin genes [8–16].

Among parasitic nematodes, BZ resistance has been best characterized in the small ruminant parasite *H. contortus* [17]. The *H. contortus* genome contains four genes that encode beta-tubulins (*tbb-isotype-1*, *tbb-isotype-2*, *tbb-isotype-3*, and *tbb-isotype-4*), where each encoded protein has a phenylalanine at position 200, which is thought to confer binding to BZs [18]. In field populations of *H. contortus*, BZ resistance has been associated historically with three canonical missense variants (F167Y, E198A, and F200Y) in *tbb-isotype-1* [19–22]. Recently, additional novel missense variants in *tbb-isotype-1* have been associated with *H. contortus* BZ resistance (*e.g.*, E198I, E198K, E198T, and E198stop) [23] These newly described missense variants in *tbb-isotype-1* are hypothesized to disrupt BZ binding, without causing loss-of-function (LoF). Given the high expression of *tbb-isotype-1*, it is likely an essential beta-tubulin in *H. contortus [18]*. By contrast, *tbb-isotype-2* LoF alleles have been identified in highly resistant *H. contortus* populations [24], indicating that it is not an essential gene and does not confer resistance on its own because variants are not observed in resistant *H. contortus* populations [18]. The other two beta-tubulin genes are expressed at low levels and have not been associated with BZ resistance [18]. The beta-tubulin genes and alleles involved in BZ resistance suggest that specific variants that prevent or reduce BZ binding can be tolerated, whereas complete loss of essential beta-tubulin genes cannot.

Unlike *H. contortus*, the free-living model nematode species *C. elegans* has six beta-tubulin genes (*tbb-1*, *tbb-2*, *mec-7*, *tbb-4*, *ben-1*, and *tbb-6*) [25]. LoF mutations in *ben-1*, an ortholog of *H. contortus tbb-isotype-1* and *tbb-isotype-2* [18], confer BZ resistance [26]. Additionally, a heterogeneous set of LoF variants in *ben-1* identified in *C. elegans* wild strains cause natural BZ resistance in this species [14]. The ability to maintain wild-

type growth despite the loss of *ben-1* is likely explained by functional redundancy among beta-tubulin genes

[27]. Previous studies have demonstrated that *Cel-tbb-1* and *Cel-tbb-2* act redundantly for viability [28–30] and

for movement, body morphology, and growth [31]. In *C. elegans*, *tbb-1* and *tbb-2* are the most broadly and highly

expressed beta-tubulin genes, whereas *ben-1* expression is restricted largely to neurons, specifically cholinergic

and glutamatergic neurons [25,32]. Consistent with this pattern, cell-specific rescue experiments demonstrated

that *ben-1* acts in cholinergic and GABAergic neurons to confer BZ susceptibility, suggesting that these neurons

are key sites of BZ action [33]. Of the six *C. elegans* beta-tubulins, only MEC-7, TBB-4, and BEN-1 contain a

phenylalanine at position 200, the residue associated with BZ binding in *H. contortus*, whereas TBB-1 and TBB-

2 contain tyrosine at this position and are predicted not to bind BZs. Together, these findings illustrate how loss

of a *C. elegans* beta-tubulin that can bind BZs causes resistance because other beta-tubulin genes are

functionally redundant for viability and do not bind BZs.

The recurrent association of beta-tubulin variants with BZ resistance in clade V parasitic and free-living

nematodes suggests the hypothesis that resistance might be predictable across species. However, the reliability

of beta-tubulin variants as predictors of BZ resistance across nematode species is unknown. This hypothesis is

difficult to test directly in parasitic nematode species because of their host-dependent life cycles, poorly

annotated reference genomes, and limited molecular and genetic tools [34,35]. By contrast, the availability of

high-quality genomic data for hundreds of wild strains [36] and the laboratory tractability of the free-living

Caenorhabditis nematode species, *C. elegans*, *Caenorhabditis briggsae*, and *Caenorhabditis tropicalis*, provide

an opportunity to test predictions of beta-tubulin mediated BZ resistance in the *Caenorhabditis* genus.

Establishing and interrogating patterns of repeated evolution in these three free-living species would provide a

mechanistic framework to anticipate resistance-associated variants in beta-tubulin genes across other clade V

nematode species, including parasitic nematodes where direct experimental validation of resistance

mechanisms remains a challenge.

Using the global natural diversity of *C. elegans*, *C. briggsae*, and *C. tropicalis*, we assessed variation in

the beta-tubulin genes *tbb-1*, *tbb-2*, *mec-7*, *tbb-4*, and *ben-1* and identified high-impact variants (*i.e.*, single

nucleotide variants (SNVs), small insertions or deletions (INDELs), and structural variants (SVs) predicted to

disrupt beta-tubulin function and confer BZ resistance. Because *ben-1* is the primary driver of BZ resistance in

C. elegans, we used an established high-throughput larval development assay (HTLDA) to expose strains with novel *ben-1* variants to the highly used BZ, albendazole (ABZ). In *C. elegans*, strains harboring seven of nine novel *Cel-ben-1* variants were resistant to ABZ, adding more alleles hypothesized to confer resistance in this species. By contrast, strains harboring only two of the eight unique *Cbr-ben-1* variants (W21stop and Q134H) were resistant to ABZ. *C. tropicalis* was distinct from the other two species, where no strains with variants in *ben-1* were resistant to ABZ. To validate the roles of *Cbr-ben-1* and *Ctr-ben-1* in BZ resistance, we generated deletion alleles and confirmed that loss of *ben-1* confers resistance in both *C. briggsae* and *C. tropicalis*. Fecundity assays showed that a loss of *Cbr-ben-1* did not affect *C. briggsae* fitness, whereas a deletion of *Ctr-ben-1* significantly reduced fecundity in *C. tropicalis*. Because *ben-1* variants could not always predict ABZ resistance, we tested whether strains harboring high-impact variants in the other four conserved beta-tubulin genes were resistant to ABZ. Strains with variants in *tbb-1*, *tbb-2*, *mec-7*, and *tbb-4* were not resistant to ABZ in any of the three *Caenorhabditis* species, indicating that *ben-1* is the primary beta-tubulin that confers resistance. The species-specific patterns of beta-tubulin-mediated BZ resistance might reflect species-specific selection pressures, such as exposure to natural BZs, and can shape the evolution of beta-tubulin-mediated resistance. The identification of alleles associated with BZ resistance in experimentally tractable species establishes a framework to predict beta-tubulin variants associated with resistance in other clade V nematodes.

RESULTS

Strains with variants or expression differences in *ben-1* were predicted to be ABZ resistant

To predict ABZ resistant strains in the three *Caenorhabditis* species, we identified high-impact variants (SNVs, INDELs, or SVs) in *ben-1* (see *Materials and Methods*). In a set of 611 *C. elegans* wild strains [36], we identified 65 strains with 33 unique high-impact variants in *ben-1* (**S1 Table**). Of the 33 variants, 24 were previously phenotyped and 20 were associated with ABZ resistance (**S1A Fig.**) [14,37]. In 641 *C. briggsae* wild strains, 22 strains with eight unique high-impact variants in *ben-1* were identified (**S2 Table**). In a set of 518 *C. tropicalis* wild strains, only two strains with unique high-impact variants in *ben-1* were identified (**S3 Table**).

Next, because low *ben-1* expression was previously correlated with ABZ resistance in *C. elegans* wild strains [38], we evaluated whether *ben-1* expression levels were predictive of ABZ responses in *C. elegans.* We

assessed the relationship between the expression of *ben-1* [38] and ABZ responses in 180 *C. elegans* wild strains [14,37] (*p*-value = 5.16e-16, r^2= 0.344). We hypothesized that low *ben-1* expression could contribute to ABZ resistance in *C. briggsae* and *C. tropicalis*, but expression data from wild strains in these species have yet to be collected.

Altogether, we identified nine novel variants in *Cel-ben-1* (E3stop, Y50C, P80S, VDN113N, Q131L, frameshift 319, frameshift 368, stop445S, and a duplication) (**S1 Table**), one strain with low *Cel-ben-1* expression, eight variants in *Cbr-ben-1* (W21stop, V91I, Q94K, D128E, Q134H, S218L, M299V, and R359H) (**S2 Table**), and two variants in *Ctr-ben-1* (P80T and R121Q) (**S3 Table**) to test for ABZ resistance. In addition to these selected strains, we also selected *C. briggsae* and *C. tropicalis* strains that lacked high-impact variants in beta-tubulin genes but were closely related to strains with variants (**S2 Fig., S3 Fig., and S4 Table**). The strains with no high-impact variants in any beta-tubulin gene might control for differences in wild strain genetic backgrounds.

High-throughput larval development assays (HTLDAs) reveal species-specific associations between *ben-1* variation and ABZ resistance

Nematodes grow longer as they progress through development, and BZs slow this progression [14–16,27,37]. A longer animal length (*i.e.*, larger animals) corresponds to increased ABZ resistance, and a shorter animal length (*i.e.*, smaller animals) corresponds to increased ABZ sensitivity. To evaluate the effects of ABZ on animal length (a proxy for development), we used image-based HTLDAs to expose all *C. elegans*, *C. briggsae*, and *C. tropicalis* wild strains with novel high-impact variants in *ben-1* or low *ben-1* expression (*i.e.*, predicted resistant strains) to control (DMSO) (**S4 Fig., S5 Fig., and S6 Fig.**) and drug (ABZ) conditions (**S7 Fig., S8 Fig., and S9 Fig.**) (see *Materials and Methods*). The assay included 48 replicates per strain with five to 30 animals per replicate in DMSO or ABZ conditions. The reported nematode length of each strain is the difference between animal lengths in DMSO and ABZ.

To define the role of *ben-1* in BZ response and classify wild strains as BZ resistant, we first obtained or created *ben-1* deletions in the reference strains for each of the three *Caenorhabditis* species. In *C. elegans*, the strain with a *ben-1* LoF variant in the N2 strain background (ECA882) has been used as an ABZ-resistant control

previously [14,27,37]. For *C. briggsae* and *C. tropicalis*, we used CRISPR-Cas9 genome editing to generate two

independent *ben-1* deletion alleles per species (see *Materials and Methods*). In *C. briggsae*, deletions of *Cbr-*

ben-1 were created in the AF16 reference strain background (ECA3953 and ECA3954) (**S10 Fig. and S5 Table**).

In *C. tropicalis*, deletions of *Ctr-ben-1* were created in the NIC58 reference strain background (ECA4247 and

ECA4248) (**S11 Fig. and S5 Table**). All deletions of *ben-1* in each of the three *Caenorhabditis* species conferred

high levels of ABZ resistance (**S12 Fig. and S13 Fig.**). Wild strains were then classified as resistant using a

threshold where the median animal lengths after ABZ exposure were no more than two SD below the median

animal length of the species-specific *ben-1* deletion strain.

In *C. elegans*, seven of the nine strains with *ben-1* variants showed minimal developmental delays after

exposure to ABZ, a phenotype similar to loss of *ben-1*, and were classified as resistant (**Fig. 1**). The two

remaining strains with *ben-1* variants (P80S, stop445S) and the strain with low *ben-1* expression were not

strongly resistant to ABZ. The P80S variant might partially alter *ben-1* function, causing a moderate resistance

phenotype. In stop445S, the normal stop codon was replaced with a serine. This variant likely does not affect

ben-1 function because position 445 is at the end of the BEN-1 protein, and only four amino acid residues (NRKL)

are added beyond the wild-type stop codon. Finally, the strain with low *ben-1* expression and no high-impact

variants in *ben-1* (JU1581) was sensitive to ABZ, indicating that the selected threshold of *ben-1* expression

(≥3.75 TPM) still retained strains with adequate *ben-1* function. Overall, natural allelic variation in *Cel-ben-1* is

associated with BZ resistance, recapitulating previous findings that BZ resistance is associated with a diverse

set of LoF variants in *Cel-ben-1* [14,27,31,37].

Fig 1. High-throughput larval development assays in the presence of albendazole for *C. elegans* strains with unique high-impact variants in BEN-1.

The regressed median animal length values for populations of nematodes grown in 30 µM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. The gray dashed line marks the *C. elegans* resistance threshold, defined as two standard deviations below the mean of the *ben-1* deletion strain in the N2 reference strain background. Results for **(A)** the N2 reference strain (orange) and a strain with a *ben-1* deletion in the N2 background (red), and **(B)** all wild *C. elegans* strains with unique high-impact variants in *ben-1* are sorted by their relative resistance to ABZ based on median animal length. Wild *C. elegans* strains are colored by beta-tubulin variant status.

In *C. briggsae*, we performed HTLDAs on the AF16 reference strain (ABZ sensitive), two *ben-1* deletion strains (ECA3953 and ECA3954) (ABZ resistant), 11 strains with eight unique variants in *ben-1*, and 13 strains genetically related to strains with *ben-1* variants in DMSO (control) (**S5 Fig.**) and ABZ conditions (**Fig. 2, S8 Fig., and S12 Fig.**). Of the 11 strains with *Cbr-ben-1* variants, only two strains with unique variants in BEN-1 (Q134H and W21stop) were ABZ resistant (**Fig. 2**). The Q134H amino acid change has been associated with ABZ

resistance in *A. caninum* and validated in *C. elegans* [11]. An early stop gain at position 21 is predicted to cause

the premature termination of protein synthesis and LoF. Of all the assayed strains with variants in BEN-1, six

(V91I, Q94K, D128E, S218L, M299V, and R359H) were not resistant to ABZ. Additionally, eleven *C. briggsae*

strains were exposed to a maximum concentration of 120 μM ABZ, where only two strains with the variants

Q134H and W21stop were resistant at these higher concentrations (**S14 Fig.**). Overall, because only one of the

seven missense variants was associated with ABZ resistance, we could not reliably predict *C. briggsae* ABZ

resistance based on the presence of a missense variant alone. To evaluate the potential fitness consequences

of *ben-1* LoF alleles, fecundity assays were performed between the AF16 reference strain and the two strains

with a loss of *ben-1* in the AF16 background (ECA3953 and ECA3954) (**S15A Fig. and S6 Table**). We found no

significant differences in fecundity between the three strains, indicating that a loss of *ben-1* does not affect fitness

in this species. These results indicate that, although a loss of *Cbr-ben-1* confers ABZ resistance with no fitness

detriment, ABZ resistance associated with *Cbr-ben-1* variants is uncommon.

233
234 **Fig 2. High-throughput larval development assays in the presence of albendazole for *C. briggsae* strains**
235 **with unique high-impact variants in BEN-1.**

The regressed median animal length values for populations of nematodes grown in 30 µM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. The gray dashed line marks the *C. briggsae* resistance threshold, defined as two standard deviations below the mean of the *ben-1* deletion strain in the AF16 reference strain background. Results for **(A)** the AF16 reference strain (green) and a strain with a *ben-1* deletion in the AF16 background (red), and **(B)** all wild *C. briggsae* strains with unique high-impact variants in *ben-1* are sorted by their relative resistance to ABZ based on median animal length. No variant (N. V.) strains (gray) paired with strains that have a high-impact variant in *ben-1* that pass the resistance threshold are shown alongside each corresponding strain with a high-impact variant in *ben-1*. Wild *C. briggsae* strains are colored by beta-tubulin variant status.

In *C. tropicalis*, we performed HTLDAs on the reference strain NIC58 (ABZ sensitive), two independent deletions of *Ctr-ben-1* (ECA4247 and ECA4248) (ABZ resistant), two *C. tropicalis* strains with variants in *ben-1* (P80T and R121Q), and two strains genetically related to *ben-1* variant strains in DMSO (**S6 Fig.**) and ABZ conditions (**Fig. 3 and S9 Fig.**). We found that all *C. tropicalis* wild strains displayed ABZ sensitivity, indicating that P80T and R121Q are not associated with ABZ resistance (**Fig. 3**). Additionally, all tested wild strains were exposed to a maximum concentration of 120 µM ABZ, and none of the strains displayed resistance at these higher concentrations (**S16 Fig.**). Finally, to evaluate the potential fitness consequences caused by a loss of *ben-1*, fecundity assays were performed between the NIC58 reference strain and the two strains with a loss of *ben-1* in the NIC58 background (ECA4247 and ECA4248) (**S15B Fig. and S7 Table**). We found that a deletion of *ben-1* caused a significant reduction in fecundity in *C. tropicalis*. The fitness defect caused by a loss of *ben-1* could explain the absence of naturally occurring *ben-1* variants in wild *C. tropicalis* strains, suggesting that *C. tropicalis* is unlikely to acquire natural *ben-1* variants that confer ABZ resistance without strong selection. If ABZ resistance is present in *C. tropicalis* wild strains, its genetic basis remains unknown and involves factors distinct from *ben-1*.

Fig 3. High-throughput larval development assays in the presence of albendazole for *C. tropicalis* strains with unique high-impact variants in BEN-1.
The regressed median animal length values for populations of nematodes grown in 30 μM albendazole (ABZ) are shown on the y-axis. Each point represents the normalized median animal length value of a well containing approximately 5-30 animals. Strains are sorted by their relative resistance to ABZ based on median animal length. Data are shown as Tukey box plots with the median as a solid horizontal line, and the top and bottom of the box representing the 75th and 25th quartiles, respectively. The top whisker is extended to the maximum point that is within the 1.5 interquartile range from the 75th quartile. The bottom whisker is extended to the minimum point that is within the 1.5 interquartile range from the 25th quartile. The gray dashed line marks the *C. tropicalis* resistance threshold, defined as two standard deviations below the mean of the *ben-1* deletion strain in the NIC58 reference strain background. Results for **(A)** the NIC58 reference strain (blue) and a strain with a *ben-1* deletion in the NIC58 background (red), and **(B)** all wild *C. tropicalis* strains with unique high-impact variants in *ben-1* are sorted by their relative resistance to ABZ based on median animal length. Wild *C. tropicalis* strains are colored by beta-tubulin variant status.

The predicted molecular consequence and severity of *ben-1* variants do not predict BZ resistance

To evaluate whether the predicted molecular consequence and severity of *ben-1* variants were predictive of BZ resistance, we characterized the identities and predicted functional impacts of each amino acid altering variant. We distinguished between alleles that disrupt multiple amino acids (*i.e.,* frameshifts, stop/start altering variants, and SVs) and alleles that affect a single amino acid (*i.e.,* missense alleles). Multi-site alleles were excluded from further analysis because they are predicted to severely disrupt or terminate BEN-1 function,

whereas missense alleles might preserve an intact protein and allow interpretation of residue-level effects on BZ

interaction.

In *C. elegans*, seven unique missense substitutions in BEN-1 were identified, of which five (Y50C, Q131L,

S145F, A185P, M275I) were associated with BZ resistance (**S17A Fig.**) [14]. In *C. briggsae*, seven unique

missense substitutions in *Cbr*-BEN-1 were identified, but only Q134H was associated with BZ resistance (**S17B**

Fig.). In *C. tropicalis*, both missense substitutions (P80T and R121Q) were not associated with BZ resistance

(**S17C Fig.**). To evaluate if missense variants with larger predicted functional impacts were associated with BZ

resistance, we quantified the severity of each substitution using BLOSUM62 and Grantham scores [39,40].

BLOSUM scores measure the evolutionary likelihood of observing a specific missense substitution, and

Grantham scores measure the physicochemical severity of a substitution. Because *C. elegans* had the most

missense variants associated with BZ resistance, we performed a regression analysis of each strain's response

to ABZ by the BLOSUM or Grantham scores of the strain's beta-tubulin variant. For *C. elegans*, neither BLOSUM

or Grantham scores showed a significant correlation with ABZ response (**S18 Fig.**). By contrast, too few

missense variants were associated with BZ resistance among *C. briggsae* and *C. tropicalis* strains to

meaningfully interpret BLOSUM or Grantham scores for these species (**S8 Table**). The limited number of

missense substitutions among wild strains and the even smaller subset associated with BZ resistance

constrained statistical power to evaluate structure-function relationships between naturally occurring beta-tubulin

variants and BZ response.

Next, we mapped missense variants onto an AlphaFold-predicted BEN-1 protein structure that showed

that the five *C. elegans* missense substitutions associated with resistance are located within the core, whereas

the two substitutions not associated with resistance (P80S and D404N) are on the protein surface (**S17A Fig.**).

By contrast, all *C. briggsae* (except Q134H) and all *C. tropicalis* missense substitutions are on the protein surface

and are not associated with resistance (**S17B Fig.**). For structural comparison, we modeled *H. contortus tbb-*

isotype-1 beta-tubulin bound to ABZ, and highlighted the canonical missense variants at positions 167, 198, and

200 (**S17D Fig.**), all of which are located within the core of the structure, near the predicted BZ-binding pocket

and are associated with resistance (**S17D Fig.**) [18–22,41]. These results suggest that BZ resistance is

associated with missense substitutions occurring within the protein core likely affecting the theoretical BZ-binding

pocket.

Natural variants in *tbb-1*, *tbb-2*, *mec-7*, and *tbb-4* are not associated with ABZ resistance across the three

free-living *Caenorhabditis* species

Although it has been established that *ben-1* is the primary gene involved in ABZ resistance in *C. elegans*

[14,27,37], we know less about the role each beta-tubulin gene plays in *C. briggsae* and *C. tropicalis* ABZ

resistance. Therefore, we identified variants in the other four conserved beta-tubulin genes across the three

Caenorhabditis species. We found no variants in *Cel-tbb-1* or *Cel-tbb-2* in any *C. elegans* wild strains (CaeNDR

Release ID: *C. elegans* - 20231213) (**Fig. 4**). However, we did identify strains with one missense variant (A9T)

in *Cel-mec-7*, a splice donor variant in *Cel-mec-7*, or one missense variant in *Cel-tbb-4* (Q8H) (**S1 Table**). The

strain with a splice donor variant in *Cel-mec-7* also carried a high-impact variant (*445S) in *Cel-ben-1*. We also

assessed the relationship between the expression of *Cel-tbb-1*, *Cel-tbb-2*, *Cel-mec-7*, or *Cel-tbb-4* [38] and ABZ

responses in 180 wild strains (**S19 Fig.**) [14,37]. In *C. briggsae*, we identified strains with missense variants in

Cbr-tbb-1 (T35A, V64I, A275T, and L377I), *Cbr-tbb-2* (E441A), *Cbr-tbb-4* (A271V), and *Cbr-mec-7* (T136I,

N165S, and S338C) (CaeNDR Release ID: *C. briggsae* - 20240129) (**S2 Table**). In *C. tropicalis*, we identified

no high-impact variants in *Ctr-tbb-1*, one missense variant in *Ctr-tbb-2* (N89S), two missense variants in *Ctr-*

mec-7 (P80F, D433E), and one missense variant in *Ctr-tbb-4* (A78V) (CaeNDR Release ID: *C. tropicalis* -

20231201) (**S3 Table**). Because expression data from *C. briggsae* and *C. tropicalis* wild strains have yet to be

collected, we could not test correlations of beta-tubulin gene expression with resistance. We performed HTLDAs

under control (DMSO) (**S20 Fig., S21 Fig., and S22 Fig.**) and ABZ conditions (**S23 Fig., S24 Fig., and S25**

Fig.) on strains in the three *Caenorhabditis* species carrying variants in the conserved beta-tubulin genes *tbb-1*,

tbb-2, *mec-7*, and *tbb-4*, along with strains genetically related to strains with variants in those beta-tubulin genes.

No variants outside of *ben-1* conferred ABZ resistance in any species, indicating that natural missense variants

in *tbb-1*, *tbb-2*, *mec-7*, and *tbb-4* are not associated with BZ resistance in the three selfing *Caenorhabditis*

species.

339
340
Fig 4. The frequency of predicted high-impact consequences in the five conserved beta-tubulin genes
present in natural populations of *C. elegans*, *C. briggsae*, and *C. tropicalis*.
The frequency of single nucleotide variants (SNVs) and structural variants (SVs) present in natural populations
of *C. elegans* (n = 611), *C. briggsae* (n = 641), and *C. tropicalis* (n = 518) (y-axis) are shown by their predicted

345 consequence in each beta-tubulin gene (x-axis). The total number of isotype reference strains with a given
346 predicted consequence are displayed on top of each bar plot.
347

Differences in tissue- and cell-specific expression patterns of beta-tubulins might influence BZ

susceptibility

Tissue-specific differences in beta-tubulin expression among the three *Caenorhabditis* species could

influence BZ resistance. To compare beta-tubulin expression patterns, we analyzed two whole-animal single-

cell transcriptomic datasets that quantify gene expression patterns across *C. elegans*, *C. briggsae*, and

C. tropicalis [42,43]. A comparison of embryonic gene expression levels in *C. elegans* and *C. briggsae* revealed

that beta-tubulin gene expression during embryogenesis is highly conserved (**S9 Table and S10 Table**). On

average, the five beta-tubulin gene distances, which quantify expression conservation across homologous cell-

types, were small (mean beta-tubulin JSDgene = 0.29), which indicated conserved embryonic expression

patterns between *C. elegans* and *C. briggsae* (JSDgene < 0.45) (**S9 Table**) [43]. Expression breadth across cell-

types (*i.e.*, cell-type specificity) was also conserved. The broad expression of *tbb-1* and *tbb-2*, and the cell-

specific expression of *ben-1*, *tbb-4*, and *mec-7* are conserved between *C. elegans* and *C. briggsae* during

embryogenesis (**S10 Table**). Among beta-tubulin genes, *tbb-2* displayed the greatest divergence in embryonic

gene expression breadth, the largest difference across species in cell-type specificity (**S10 Table**). Overall,

embryonic beta-tubulin expression patterns are conserved between *C. elegans* and *C. briggsae*, although *tbb-2*

expression breadth is an exception (**S10 Table**). Next, because neuronal *ben-1* expression restores BZ

susceptibility in *C. elegans* [33]we focused on the divergence of beta-tubulin expression across homologous

neuronal cell classes at the L2 larval stage [43]. For each gene, we quantified neuron-class-specific expression

divergence across the three *Caenorhabditis* species using the proportion of neuronal classes where gene

expression differs among species (**S11 Table**) [43]. At least one species expressed *ben-1* in 98 of 118 neuronal

cell classes. However, all three *Caenorhabditis* species expressed *ben-1* in only 35 of these classes (Jaccard

distance = 0.64) (**S11 Table**). By contrast, *tbb-1* and *tbb-2* had highly conserved expression across the three

Caenorhabditis species (98 of 118 neuronal cell classes expressed *tbb-1* or *tbb-2* in all three species) (Jaccard

distance = 0) (**S11 Table**) [43]. The broad and conserved neuronal expression of *tbb-1* and *tbb-2* contrasts with

ben-1, which showed narrower and species-specific expression patterns. Next, because *ben-1* expression in

cholinergic neurons restores BZ susceptibility in *C. elegans [33]*, we compared cholinergic neuronal expression

and identified several cell classes with species-specific patterns. Generally, beta-tubulin cell-specificity is

conserved across the three *Caenorhabditis* species (**S9 Table**, **S10 Table**, **S11 Table**). However, examples of

expression divergence among *C. elegans*, *C. briggsae*, and *C. tropicalis* were identified for *ben-1* and *tbb-2.* The

divergence in *ben-1* expression across cholinergic cell classes indicates that the neuronal sites where *ben-1*

function causes BZ susceptibility could differ among species. Future work should test this hypothesis using

transgenic strains that express *ben-1* in different tissues and neuronal cells.

The most diverse high-impact variants are found in *Cel-ben-1*

To better understand the evolution of predicted BZ resistance alleles in the three *Caenorhabditis* species,

we assessed the population-wide frequencies of each beta-tubulin variant. First, to determine the prevalence of

high-impact variants in the five conserved beta-tubulin genes (*tbb-1*, *tbb-2*, *mec-7*, *tbb-4*, and *ben-1*) across

Caenorhabditis species, we quantified the frequency of each variant (deletion, duplication, frameshift, in-frame

deletion, inversion, missense, splice donor, and start/stop altering) in each species. With global sampling of the

three *Caenorhabditis* species (*C. elegans*: 611 strains, *C. briggsae*: 641 strains, *C. tropicalis*: 518 strains), we

found that *C. elegans* variants predicted to cause deleterious functional effects were present in 1% of strains. By

contrast, variants predicted to cause deleterious functional effects in *C. briggsae* and *C. tropicalis* were rare (<

0.05% of all strains in either species) (**Fig. 4**). The most variation in beta-tubulin genes was identified in *Cel-ben-*

1, where deletions, frameshifts, in-frame deletions, inversions, missense, stop/start altering variants, and a

duplication were found. Next, we found one missense variant in *Cel-tbb-4* and several missense variants and a

splice donor in *Cel-mec-7*. Overall, *C. elegans* has acquired the most diverse set of high-impact variants in and

predicted functional effects on *ben-1*. In *C. briggsae*, we found 21 missense amino acid substitutions and a single

start/stop altering consequence in *Cbr-ben-1*. In both *Cbr-tbb-1* and *Cbr-tbb-2*, we identified rare missense

consequences. Additionally, we found nine strains with splice variants in *Cbr-tbb-4* and one strain with a

missense variant. In *C. tropicalis*, we found missense consequences in all beta-tubulin genes, except *tbb-1*.

Additionally, we found six strains with splice variants in *Ctr-tbb-4*. These findings highlight that *Cel-ben-1* has the

most diverse set of variants, reinforcing its role in BZ resistance.

Next, we examined the geographic distribution of strains carrying high-impact variants in *tbb-1*, *tbb-2*,

mec-7, *tbb-4*, and *ben-1* to determine if beta-tubulin variants were associated with natural sampling location. In

C. elegans, variants in *ben-1* were found globally with no discernible geographic pattern but were concentrated

in clades that have experienced recent selective sweeps [44,45] (**Fig. 5**). *Cel-ben-1* variants in swept clades

suggest that these mutations arose as relatively recent evolutionary events in response to BZ-like compounds

in the natural niche. By contrast, in *C. briggsae*, variants in *ben-1* were distributed throughout the species tree

and found on more ancestral branches (*i.e.*, earlier diverged lineages in the species) (**Fig. 5C**). For *C. tropicalis*,

the limited number of variants in *ben-1* precludes any definitive conclusions regarding their evolutionary patterns

(**Fig. 5D**). Because few variants are found in *tbb-1* (**S26 Fig.**), *tbb-2* (**S27 Fig.**), *mec-7* (**S28 Fig.**), and *tbb-4* (**S29**

Fig.), we cannot identify the evolutionary patterns of BZ resistance in these genes for any of the three species.

Fig 5. The global distribution of *Caenorhabditis* strains that contain predicted high-impact variation in BEN-1.
(A) Each point corresponds to the sampling location of an individual *C. elegans* (orange), *C. briggsae* (green), or *C. tropicalis* (blue) isotype reference strain with a predicted high-impact consequence in BEN-1. A genome-wide phylogeny of **(B)** 611 *C. elegans*, **(C)** 641 *C. briggsae*, and **(D)** 518 *C. tropicalis* isotype reference strains, where each point denotes an isotype reference strain with a predicted high-impact consequence in BEN-1 is shown.

Finally, because substrates harbor distinct microbial communities that can influence the evolution of BZ

resistance alleles, we determined if strains carrying a high-impact variant in a beta-tubulin gene were associated

with specific substrates. We performed substrate enrichment analysis to assess correlations between 12

substrates and all strains in the three *Caenorhabditis* species (**S12 Table**). However, no significant enrichment

was observed between a high-impact variant in a beta-tubulin gene and any given substrate (Fisher's Exact

Test, p=1) (**S30 Fig.**). Because no geographic or substrate enrichment was observed, evolutionary pressures

driving beta-tubulin variation are likely not strongly tied to substrate.

DISCUSSION

Beta-tubulin-mediated BZ resistance varies across natural populations of *Caenorhabditis* nematodes

This study provides new insights into beta-tubulin-mediated ABZ resistance across three *Caenorhabditis*

species. Each *Caenorhabditis* species harbored a unique set of predicted high-impact beta-tubulin alleles, but

only variants in *ben-1* conferred resistance. With additional wild strains since our first study, we identified

additional *Cel-ben-1* LoF alleles associated with BZ resistance, which confirmed that a diverse collection of

predicted LoF variants in *ben-1* are associated with ABZ resistance in *C. elegans* [14,16]. In *C. briggsae*, strains

harboring only two of the eight unique *Cbr-ben-1* variants were resistant to ABZ. One *Cbr-ben-1* variant

(W21stop) causes early protein termination, and the other (Q134H) alters a residue within the *Cbr*-BEN-1 protein

that likely affects ABZ binding and is associated with resistance in *A. caninum [11]*. A CRISPR-Cas9-generated

deletion of *Cbr-ben-1* conferred resistance similar to that displayed by wild strains with these alleles and did not

impact fitness. To date, resistance has yet to be identified in *C. tropicalis* wild strains. However, a CRISPR-Cas9-

generated deletion of *Ctr-ben-1* conferred resistance and caused a significant reduction in fitness (fecundity),

which likely explains why just two high-impact variants were identified among wild strains and neither caused

loss of *ben-1* function. Overall, our results highlight the complexity of BZ resistance. Accurate prediction of BZ

resistance across nematode species requires a clear understanding of the contributions of both beta-tubulin

dependent and beta-tubulin independent mechanisms.

Nematode survival under BZ exposure depends on beta-tubulin dosage and drug-binding ability

BZs bind to beta-tubulins and inhibit the polymerization of microtubules [2–4]. Therefore, despite the

presence of beta-tubulin independent resistance mechanisms [46,47], beta-tubulins have a large impact on BZ

efficacy as an anthelmintic treatment. Two factors determine how beta-tubulin impacts susceptibility to BZs: (1)

the beta-tubulin's ability to bind to BZs and (2) the dosage of the beta-tubulin protein, which can be modified by

tissue-specific expression. Nematode species have distinct beta-tubulin gene complements with divergence in

number of genes, BZ binding affinities, expression levels and cell types, and redundancy. Ultimately, this

divergence in beta-tubulin complement shapes the type(s) of resistance-associated alleles that arise in each

nematode species. In species where the primary BZ-binding beta-tubulin is redundant with other isoforms, as in

selfing *Caenorhabditis* species, loss of a single copy can confer resistance without disrupting essential

microtubule functions. In species without redundancy, such as *H. contortus*, beta-tubulin variants that alter the

BZ binding affinity are the only available beta-tubulin dependent route to BZ resistance.

In *C. elegans*, *ben-1* is the primary target of BZs but is redundant with *tbb-1* and *tbb-2* [28–30]. LoF alleles

in *ben-1* reduce the total amount of BZ-binding protein while preserving essential microtubule functions, which

permits nematode development during BZ exposure. A similar pattern likely occurs in *C. briggsae*, where rare

high-impact variants in *ben-1* confer resistance, and other beta-tubulin genes maintain essential microtubule

functions. By contrast, *H. contortus* relies on the essential beta-tubulin *tbb-isotype-1* and cannot tolerate loss

without severe fitness consequences [18]. The differences between the *Caenorhabditis* species and *H. contortus*

demonstrate that the prediction of beta-tubulin dependent resistance requires the identification of the total

number and expression levels of beta-tubulin genes and determination of which encoded beta-tubulins can bind

to BZ. Altogether, BZ resistance depends not on the copy number of beta-tubulins alone but on the availability

of functional beta-tubulins capable of interacting with BZs.

How can we accurately predict BZ resistance across nematode species?

To accurately predict BZ resistance across nematode species, we need (1) tractable nematode models

with beta-tubulin gene complements that resemble that of parasite species, (2) improved parasitic nematode

genomes that enable the comprehensive identification of beta-tubulin genes, (3) functional tests that define how

parasite beta-tubulin alleles contribute to BZ response, and (4) to define the contribution of non-beta-tubulin

genes to overall BZ response. To date, no experimentally tractable model nematode species exist with beta-

tubulin genes that have both limited redundancy and BZ binding properties similar to those of parasitic nematodes. *Pristionchus pacificus*, a free-living clade V nematode has three beta-tubulin genes, comprising two *ben-1* orthologs (*Ppa-ben-1.1* and *Ppa-ben-1.2*) and an ortholog of *Cel-mec-7* (*Ppa-mec-7*) [48]. All three beta-tubulin genes in *Pristionchus pacificus* are predicted to have high ABZ binding affinity (**S31 Fig.**). Using CRISPR-Cas9 genome editing, we found that homozygous LoF alleles for either *Ppa-ben-1.1* or *Ppa-ben-1.2* could not be recovered in *P. pacificus*, suggesting that both genes are essential **(S13 Table)**. The fact that *P. pacificus* has fewer beta-tubulin genes than *C. elegans* likely contributes to the lack of redundancy among beta-tubulins. The number, essentiality, and BZ binding affinity of the *P. pacificus* beta-tubulins more closely resembles that of *H. contortus*, which positions *P. pacificus* as a valuable free-living model to study essential beta-tubulin function, beta-tubulin dosage effects, and BZ resistance relevant to parasitic nematode species.

Second, to understand the role beta-tubulins play in BZ resistance across parasitic nematode species, we must significantly improve genomes and gene models. Most parasitic nematode genomes remain incomplete or are poorly annotated, which obscures beta-tubulin copy number and gene identity. Recent efforts have produced higher quality genomes for some parasitic nematodes [49–51], but we must still define beta-tubulin copy number across diverse nematode species. For example, improved reference genomes and gene models for *C. briggsae* [52,53] and *C. tropicalis* [54] enabled accurate beta-tubulin gene identification for these species. Until genome assemblies, technologies, and analytical techniques improve, we will be unable to accurately predict the number of beta-tubulin genes in a given species and their respective BZ binding affinities.

Third, functional validation of parasitic resistance alleles presents additional challenges. Although tools such as RNAi can work in parasitic nematodes [55], delivery challenges and difficulty isolating edited individuals limit its use. An alternative strategy is to introduce predicted parasite resistance alleles into *C. elegans* to test effects on fitness and BZ response, as shown previously [14–16]. Finally, some parasitic nematodes, such as ascarid species, exhibit BZ resistance without known beta-tubulin resistance alleles [56–58]. This pattern suggests that resistance arises independently of detected beta-tubulin sequence changes or reflects incomplete identification of beta-tubulin genes caused by poor genome assemblies. To date, the underlying cause of BZ resistance in ascarid species remains unclear. Reduced functional beta-tubulin dosage, altered BZ binding, or both could contribute to BZ resistance in ascarids [59]. Improved genome assemblies and annotation are

required to identify all beta-tubulin genes and drug-binding sites to define the mechanisms of BZ resistance in ascarid species. In the future, the introduction of newly identified ascarid alleles into free-living nematode models can directly test resistance and fitness, and clarify the mechanisms that drive BZ resistance in clade III nematodes. Additionally, we must define how redundancy, essentiality, beta-tubulin dosage, and drug-binding properties shape the evolution of resistance to improve the prediction of BZ resistance across the huge diversity of parasitic nematode species.

MATERIALS AND METHODS

Identification of beta-tubulin loci

Amino acid sequences for all six *C. elegans* beta-tubulin proteins were obtained from WormBase (WS283) [60] and used as queries in a BLASTp search (Version 2.12.0) [61] against protein sequence databases constructed using gene models for *C. briggsae* [53] and *C. tropicalis* [62]. To construct the protein sequence database, we extracted gene model transcript features from the gene feature file with *gffread* (Version 0.9.11) [63] and processed them using the *makeblastdb* function from BLAST (Version 2.12.0). From the BLASTp search, we identified *C. briggsae* and *C. tropicalis* protein sequences with the highest percent identity (PID) to each *C. elegans* beta-tubulin protein. Only protein sequences with the highest PID in both searches were considered orthologs (**S14 Table**). For some *C. elegans* beta-tubulin orthologs, multiple *C. briggsae* or *C. tropicalis* gene models contained multiple splice isoforms. All gene models for all beta-tubulin transcripts were manually inspected, and isoforms that were not fully supported by short-read RNA sequencing data were removed.

Single nucleotide variant (SNV) and indel calling and annotation

To identify single nucleotide variants (SNVs) or indels (insertions and deletions) in the beta-tubulin genes across the selfing *Caenorhabditis* species, we used the Variant Annotation Tool from the *Caenorhabditis* Natural Diversity Resource (CaeNDR) (Release IDs: *C. elegans* - 20231213, *C. briggsae* - 20240129, *C. tropicalis* - 20231201) [36]. The identified SNVs and indels included small insertions and deletions, frameshifts, altered stop and start codons, nonsynonymous changes, and splice variants (**S1 Table, S2 Table, and S3 Table**).

Structural variant (SV) calling and annotation

Structural variant (SV) calling was performed using *DELLY* (Version 0.8.3), a SV caller optimized to detect

large insertions, deletions, and other complex structural variants such as inversions, translocations, and

duplications in paired-end short-read alignments [64] and shown to perform well on *C. elegans* short-read

sequence data [65]. SVs that overlapped with beta-tubulin genes were extracted using *bcftools* (Version 1.10.1)

[66]. Insertions, deletions, inversions, and duplications that passed the *DELLY* (Version 0.8.3) default quality

threshold (greater than three supporting read pairs with a median MAPQ > 20), filtered to high-quality genotypes

(genotype quality > 15), and had at least one alternative allele were retained. For complex variants (inversions

and duplications), the identification of at least one split-read pair was required (variants flagged as a precise SV

by *DELLY*). To validate SVs that passed quality filtering, each SV was manually inspected for breakpoints in the

raw-read alignments (*Wally*, Version 0.5.8) and for impacts on the beta-tubulin coding sequence (CaeNDR

Genome Browser) [36] (**S15 Table**). We retained SVs where raw read alignments suggested that the SV

impacted the beta-tubulin coding sequence. We compared the *Cel-ben-1* SVs called by *DELLY* to those SVs

identified previously [14]. *DELLY* successfully recalled structural variants in several strains, including deletions

in JU751, JU830, JU1395, JU2582, JU2587, JU2593, JU2829, and QX1233, as well as an inversion in MY518.

However, *DELLY* did not detect a previously reported transposon insertion in strain JU3125. To assess if other

SVs could have been missed by *DELLY*, we manually inspected the read alignments for all strains that had not

been previously phenotyped to check if any other SVs were not detected by *DELLY*. We confirmed the presence

of novel *Cel-ben-1* SVs in multiple strains, including putative deletions in ECA706 and NIC1832, and a duplication

in NIC1107. Additionally, we identified a previously undetected putative deletion in JU4287. We also examined

the amino acids at position 200 in TBB-1, TBB-2, MEC-7, TBB-4, and BEN-1 orthologs and found that all MEC-

7, TBB-4, and BEN-1 orthologs contained phenylalanine, and that all TBB-1 and TBB-2 orthologs contained

tyrosine at position 200 (**S31 Fig.**).

Association of *Cel-ben-1* expression with ABZ response

Two previous assays measured developmental responses of wild *C. elegans* strains after ABZ exposure [14,67]. For 180 of these wild *C. elegans* strains, whole-animal expression levels (transcripts per million estimates [TPM]) were collected from untreated young-adult animals [38]. We identified strains with low *Cel-ben-1* expression by selecting strains with TPM values more than one standard deviation (SD) below the mean expression level across all 207 wild strains with expression data. Of the 180 wild *C. elegans* strains, 105 strains were measured for both ABZ response and gene expression. A linear model was built using the *lm* function in R to account for assay effects. Subsequently, the residuals of the linear model were used to normalize previous measures of ABZ response. We evaluated the linear fit between each strain's expression of *ben-1* (**S1 Fig.**), *tbb-1*, *tbb-2*, *mec-7*, or *tbb-4* (**S19 Fig.**) and the developmental delay following ABZ exposure.

Phylogenetic analysis

We characterized the relatedness of isotype reference strains (genetically unique strains) with beta-tubulin variants using species trees downloaded from CaeNDR and generated by the 'post-gatk-nf' pipeline (https://github.com/AndersenLab/post-gatk-nf) (Release IDs: *C. elegans* - 20231213, *C. briggsae* - 20240129, *C. tropicalis* - 20231201) [36]. Briefly, the trees were generated using high-quality SNVs in isotype reference strains retained in the hard-filtered variant call format (VCF) file. *vcf2phylip* [68] and the *bioconvert* [69] function *phylip2stockholm* were used to prepare inputs for *quicktree*, which was used to construct a tree using a neighbor-joining algorithm [70]. All versions of these software can be accessed from the 'post-gatk' docker container (https://hub.docker.com/r/andersenlab/tree), used by the 'post-gatk-nf' pipeline. We visualized the trees for each species using the *ggtree* function from the *ggtree* (v3.6.2) R package [71].

Strain selection and maintenance

Eighteen *C. elegans* strains, 45 *C. briggsae* strains, and 15 *C. tropicalis* strains from the CaeNDR [36] were used in this study (**S1 Table, S2 Table, and S3 Table)**. Isolation details for each strain are included in CaeNDR. For each species, we selected strains that had variants (SNV or SV) with unique high-impact consequences in *tbb-1*, *tbb-2*, *mec-7*, *tbb-4*, or *ben-1* that had not been previously phenotyped. High-impact consequences included changes to amino acids, start and stop codon positions, or splice variants [36]. Strains

with high-impact consequences in a beta-tubulin gene are herein referred to as "predicted resistant" strains. Strains that were closely related to predicted resistant strains with no high-impact consequences in beta-tubulin genes were also included for *C. briggsae* and *C. tropicalis* and herein classified as "predicted susceptible" strains (**S2 Fig., S3 Fig., and S4 Table**). The reference strains for all three species were included.

Before measuring ABZ responses, *C. elegans* and *C. briggsae* animals were maintained at 20ºC and *C. tropicalis* animals were maintained at 25ºC. All animals were maintained on 6 cm plates with modified nematode growth medium (NGMA), which contains 1% agar and 0.7% agarose to prevent animals from burrowing [72]. The NGMA plates were seeded with the *Escherichia coli* strain OP50 as a nematode food source. All strains were grown for three generations without starvation on NGMA plates before anthelmintic exposure to reduce the transgenerational effects of starvation stress [73].

CRISPR-Cas9 genome editing

To validate the role that *ben-1* plays in ABZ resistance in the three *Caenorhabditis* species, we used CRISPR-Cas9 to create *ben-1* deletions. For *C. elegans*, we used a previously generated strain (ECA882) with a *ben-1* deletion in the N2 background [14,15,27,37]. The *ben-1* deletions were generated in the AF16 background for *C. briggsae* and the NIC58 background for *C. tropicalis*. Injections were performed by InVivo Biosystems (Eugene, OR), and deletions of *Cbr-ben-1* and *Ctr-ben-1* were confirmed using PCR (**S10 Fig. and S11 Fig.**). Briefly, two primer pairs were designed for the deletion alleles for each species, with each pair designed to bind to a region external or internal to both of the deletions. Confirmation of deletion was performed by performing two amplification reactions for each sample: (1) the use of both external primers, and (2) the use of an internal and external pair (**S5 Table**). The parental strain was used as a control in each PCR. Deletions were confirmed by a reduction in the size of the external-external amplicon in the edited strains compared to the unedited parental control strain. Homozygosity was confirmed by the loss of a band amplified from the external-internal primer pair. Edited strains underwent two generations of PCR confirmation for homozygosity. Two independent edits of each allele in each species were generated to control for any potential off-target effects caused by CRISPR-Cas9 genome editing (**S12 Fig., S13 Fig., and S5 Table**).

Nematode food preparation for NGMA 6 cm plates

The OP50 *E. coli* strain was used as a nematode food source for NGMA plates. A frozen stock of OP50 was streaked onto a 10 cm Luria-Bertani (LB) agar plate and incubated overnight at 37ºC. The following day, a single bacterial colony was transferred into each of two culture tubes that contained 5 mL of 1x LB. The starter cultures and two negative controls (1X LB without *E. coli*) were incubated for 18 hours at 37ºC shaking at 210 rpm. The OD_{600} value of the starter cultures were measured using a spectrophotometer (BioRad, SmartSpec Plus) to calculate how much starter culture was needed to inoculate a 1 L culture at an OD_{600} value of 0.005. For each assay, one culture containing 1 L of pre-warmed 1X LB inoculated with the starter culture grew for approximately 4 - 4.5 hours at 37ºC at 210 rpm to an OD_{600} value between 0.45 and 0.6. Cultures were transferred to 4ºC to slow growth. OP50 was spotted on NGMA test plates (two per culture) and grown at 37ºC overnight to assay for contamination.

Nematode food preparation for high-throughput larval development assays (HTLDAs)

One batch of HB101 *E. coli* was used as a nematode food source for all HTLDAs in this study. A frozen stock of HB101 *E. coli* was streaked onto a 10 cm LB agar plate and incubated overnight at 37ºC. The following day, a single bacterial colony was transferred into three culture tubes that contained 5 mL of 1x Horvitz Super Broth (HSB). The starter cultures and two negative controls (1X HSB without *E. coli*) were incubated for 18 hours at 37ºC shaking at 180 rpm. The OD_{600} value of the starter cultures were measured using a spectrophotometer (BioRad, SmartSpec Plus) to calculate how much starter culture was needed to inoculate a 1 L culture at an OD_{600} value of 0.001. A total of four cultures each containing 1 L of pre-warmed 1X HSB inoculated with the starter culture grew for 15 hours at 37ºC while shaking at 180 rpm. After 15 hours, flasks were removed from the incubator and transferred to 4ºC to slow growth. The 1X HSB was removed from the cultures by performing three rounds of centrifugation, where the supernatant was removed, and the bacterial cells were pelleted. Bacterial cells were washed with K medium, resuspended in K medium, pooled, and transferred to a 2 L glass beaker. The OD_{600} value of the bacterial suspension was measured and diluted to a final concentration of OD_{600}100 with K medium, aliquoted to 15 mL conicals, and stored at -80ºC for use in the HTLDAs.

ABZ dose-response assays for *C. briggsae* and *C. tropicalis*

Because ABZ response had been minimally characterized in *C. briggsae* [74] and has not yet been described in *C. tropicalis*, we first measured dose-response curves for both species after exposure to ABZ to assess developmental delay. Before performing HTLDAs, ABZ (Sigma-Aldrich, Catalog # A4673-10G) stock solutions were prepared in dimethyl sulfoxide (DMSO) (Fisher Scientific, Catalog # D1281), aliquoted, and stored at -20ºC for use in the assays. For the dose-response assays, animals were exposed to ABZ at the following concentrations (µM): 0 (1% DMSO), 0.12, 0.23, 0.47, 0.94, 1.88, 3.75, 7.5, 15, 30, 60, and 120. Animals developed in the presence of ABZ as described in *HTLDAs to assess nematode development*.

Dose-response model estimation and statistics were performed as described previously [67,75]. Briefly, a four-parameter log-logistic dose-response curve was fit independently for a genetically diverse set of 11 *C. briggsae* strains (**S14 Fig.**) and seven *C. tropicalis* strains (**S16 Fig.**), where normalized median animal length was used as a metric for phenotypic response (see *HTLDA data collection and data cleaning*). For each strain-specific dose-response model, slope (*b*) and concentration (*e*) were estimated with strain as a covariate. We calculated EC_{10} as we have previously found EC_{10} response to be more heritable than half maximal effective concentration (EC_{50}) estimates and were used in our analysis [67,75]. A dosage of 30 µM ABZ was closest to the EC_{10} for *C. briggsae* and *C. tropicalis*, consistent with ABZ concentrations used in past *C. elegans* assays [15,16,37] and in all HTLDAs in this study.

HTLDAs to assess nematode development

Populations of each strain were amplified and bleach-synchronized in three independent assays. Independent bleach synchronizations controlled for variation in embryo survival and subsequent effects on developmental rates. After bleach synchronization, approximately 30 embryos were dispensed into each well of a 96-well microplate in 50 µL of K medium. Each strain had sixteen wells per condition (DMSO or ABZ) in each assay. Three independent assays yielded a total of forty-eight wells per condition per strain. Each 96-well microplate was prepared, labeled, and sealed using gas-permeable sealing films (Fisher Scientific, Catalog # 14-222-043). Plates were placed in humidity chambers to incubate for 24 hours at 20°C for *C. elegans* and *C. briggsae*, and 25°C for *C. tropicalis* while shaking at 170 rpm (INFORS HT Multitron shaker). After 24 hours,

every plate was inspected to ensure that all embryos hatched and animals were developmentally arrested at the

first larval (L1) stage so all strains started each assay at the same developmental stage. Next, food was prepared

to feed the developmentally arrested L1 animals using the required number of $OD_{600}100$ HB101 aliquots (see

Nematode food preparation for HTLDAs). The HB101 aliquots were thawed at room temperature, combined into

a single conical tube, and diluted to an $OD_{600}30$ with K medium. To inhibit further bacterial growth and prevent

contamination, 150 μL of kanamycin was added to the HB101. An aliquot of 100 μM ABZ stock solution was

thawed at room temperature and added to an aliquot of $OD_{600}30$ K medium at a 3% volume/volume ratio. Next,

25 μL of the food and ABZ mixture was transferred into the appropriate wells of the 96-well microplates to feed

the arrested L1 animals at a final HB101 concentration of $OD_{600}10$ and expose L1 animals to ABZ. Immediately

afterward, the 96-well microplates were sealed using a new gas permeable sealing film, returned to the humidity

chambers, and incubated for 48 hours at 20°C (*C. elegans* and *C. briggsae*) or 42 hours at 25°C (*C. tropicalis*)

while shaking at 170 rpm. After 48 hours (*C. elegans* and *C. briggsae*) or 42 hours (*C. tropicalis*) of incubation

and shaking in the presence of food and either DMSO or ABZ, the 96-well microplates were removed from the

incubator and treated with 50 mM sodium azide in M9 for 10 minutes to paralyze and straighten nematodes.

After 10 minutes, images of nematodes in the microplates were immediately captured using Molecular Devices

ImageXpress Nano microscope (Molecular Devices, San Jose, CA) using a 2X objective. The ImageXpress

Nano microscope acquires brightfield images using a 4.7 megapixel CMOS camera and stores images in a 16-

bit TIFF format. The images were used to quantify the development of nematodes in the presence of DMSO or

ABZ as described below (see *High-throughputput imager assays [HTLDA] data collection and cleaning*). A full

step-by-step protocol for the HTLDA has been deposited on protocols.io [76].

HTLDA data collection and data cleaning

CellProfiler (Version 24.10.1) was used to characterize and quantify biological data from the image-based

assays. Custom software packages designed to extract animal measurements from images collected on the

Molecular Devices ImageXpress Nano microscope were previously described [77]. *CellProfiler* modules and

WormToolbox were developed to extract morphological features of individual animals from images from the

HTLDA [78]. Worm model estimations and custom *CellProfiler* pipelines were written using the *WormToolbox* in

the GUI-based instance of *CellProfiler* [75]. Next, a Nextflow pipeline (Version 24) was written to run command-

line instances of *CellProfiler* in parallel on the Rockfish High-Performance Computing Cluster (Johns Hopkins

University). The *CellProfiler* workflow can be found at https://github.com/AndersenLab/cellprofiler-nf. The custom

CellProfiler pipeline generates animal measurements by using four worm models: three worm models tailored to

capture animals at the L4 larval stage, in the L2 and L3 larval stages, and the L1 larval stage, as well as a "multi-

drug high dose" (MDHD) model, to capture animals with more abnormal body sizes caused by extreme

anthelmintic responses. These measurements comprised our raw dataset. Two *C. briggsae* strains (NIC1052

and VX34) were not fully paralyzed and straightened at the time of imaging, which created some misclassification

of animal measurements. Thus, the animal lengths for strains NIC1052 and VX34 measured by *CellProfiler* are

shorter than the actual animal lengths. However, the difference in animal lengths does not affect the classification

of a strain as resistant or sensitive to ABZ. Data cleaning and analysis steps were performed using a custom R

package, *easyXpress* (Version 2.0) [77] and followed methods previously reported [37]. All analyses were

performed using the R statistical environment (Version 4.2.1) unless stated otherwise.

C. briggsae and C. tropicalis fecundity assays

To define the fitness costs associated with a loss of *ben-1* in *C. briggsae* and *C. tropicalis*, we performed

fecundity assays. For *C. briggsae*, we used the two strains with independent edits of *ben-1* in the AF16

background (ECA3953 and ECA3954) and the AF16 reference strain. For *C. tropicalis*, we used the two strains

with independent edits of *ben-1* in the NIC58 background (ECA4247 and ECA4248) and the NIC58 reference

strain. To perform fecundity assays, we placed a single L4 larval stage hermaphrodite from each strain onto a 6

cm NGMA plate that was spotted with *E. coli* OP50. *C. briggsae* assay plates were maintained at 20°C, and

C. tropicalis assay plates were maintained at 25°C. For each assay plate, the original hermaphrodite parent was

transferred to a fresh 6 cm NGMA plate every 24 hours for 192 hours. Ten technical replicates were prepared

for each strain. The Basic Imaging Platform from Tau Scientific was used to collect images for each of the assay

plates (0, 24, 48, 72, 96, and 120-hours) at either 48 or 72 hours for *C. briggsae* and *C. tropicalis* animals,

respectively, after the removal of the parent from each NGMA plate. The total offspring was counted from each

image by visual inspection using the Multi-point tool in ImageJ (Version 1.54g). The original hermaphrodite

parents were excluded from the counts. Replicates where the original hermaphrodite parent died were excluded

from the analysis. Only biological replicates with data from at least six assay plates were used to calculate total

fecundity (**S15 Fig., S6 Table, and S7 Table**).

Tissue-specific beta-tubulin gene expression conservation in *C. elegans*, *C. briggsae,* and *C. tropicalis*

To assess beta-tubulin expression divergence across neuronal cell classes, we used whole-animal

single-cell transcriptomes of *C. elegans*, *C. briggsae*, and *C. tropicalis* [43]. We quantified neuronal cell

expression divergence across species using precomputed Jaccard distances calculated from the neuronal cell

classes in at least one species and the number of neuron classes in which a gene is expressed in all three

species [43]. We also downloaded gene expression summary data for each homologous cell class from

CaenoGen and analyzed expression patterns in cell classes that used acetylcholine (ACh) as a neurotransmitter

in *C. elegans [79]*. We identified cell classes in which *ben-1* orthologs showed species-specific presence or

absence of expression.

We also analyzed embryonic single-cell transcriptomes from the *C. briggsae* and *C. elegans* reference

strains to examine the conservation of beta-tubulin gene expression in homologous cell types [42]. We used pre-

computed summary statistics to quantify two components of expression divergence among beta-tubulin genes:

gene distance (reported as Jensen-Shannon Distances for each gene) and expression breadth (reported as Tau

metrics for each species). Beta-tubulin gene distances and expression breadth metrics are available in the gene

data summary table file (**S9 Table and S10 Table**). Estimates for three cell populations are distinguished by cell-

type assignment: progenitor, terminal, and joint (combined progenitor and terminal cells).

Protein structure and visualization of *ben-1* and *tbb-isotype-1* variants

For *C. elegans*, the BEN-1 amino acid sequence was obtained from WormBase (WS283) [60]. For

C. briggsae and *C. tropicalis*, BEN-1 amino acid sequences were the best hits to the *C. elegans* BEN-1 query

from the reciprocal BLASTp search used to identify beta-tubulin orthologs (*see Identification of beta-tubulin loci*)

(**S16 Table**). The *H. contortus tbb-isotype-1* amino acid sequence was obtained from WormBase Parasite

(Version: WBPS19) [41]. Protein structures were predicted using AlphaFold3 [80]. All BEN-1 variant data and

associated benzimidazole-response phenotypes were compiled from this study and previously published data [14,37]. A custom Python script was used to generate a PyMOL script (Version 3.1.6.1) for structural visualization. Each predicted beta-tubulin structure was aligned to *Cel*-BEN-1 using the PyMOL `align` command.

Mutation of *ben-1* orthologs in *Pristionchus pacificus*

To test the function of *ben-1* in *Pristionchus pacificus*, we used CRISPR-Cas9 genome editing to delete the two orthologs of *Cel-ben-1*. Designed crRNAs and tracrRNA were synthesized by Integrated DNA Technologies (IDT). For the creation of the guide RNAs, 3 µL of 100 µM tracrRNA (IDT) was combined with 3 µL of 100 µM crRNA (IDT) and incubated at 95ºC for five minutes, followed by five minutes at room temperature for annealing. The ribonucleoprotein (RNP) complex was then created by combining 0.61 µL of the guide-RNA mixture with 0.25 µL of Cas9 protein (IDT), followed by incubation at 37ºC for 10 minutes. The final microinjection mix was then created by combining the 0.86 µL RNP complex with 9.14 µL of a TE buffer mixture containing a (55 ng/µL) plasmid containing a *P. pacificus* codon-optimized *egl-20p::TurboRFP::rpl-23UTR* construct [81] used as a co-injection marker for visual identification of successful injections by the presence of fluorescent F1 individuals (**S13 Table)**. All F1 individuals displaying fluorescence were then isolated and allowed to self-fertilize. After successful F2 embryo hatching, F1 individuals were genotyped for mutations using a heteroduplex mobility assay (HMA). Individuals from F2 broods that were determined to contain heterozygous mutations in F1 mothers were then isolated, allowed to self-fertilize, and genotyped by HMA after F3 embryos had hatched to identify homozygous mutant lines, followed by Sanger sequencing to determine types of mutations induced. Additionally, for mutations that were homozygous lethal, the F3 from heterozygous F2 individuals were likewise genotyped by HMA to quantify survivability based on deviations from expected Mendelian ratios.

DATA AVAILABILITY STATEMENT

All code and data used to replicate the data analysis and figures are available on GitHub at: https://github.com/AndersenLab/ce_cb_ct_betatubulin. S1 Table contains the list of *C. elegans* isotype reference strains, any beta-tubulin variants, sample collection locations, and substrate types. S2 Table contains the list of

C. briggsae isotype reference strains, any beta-tubulin variants, sample collection locations, and substrate types.

S3 Table contains the list of *C. tropicalis* isotype reference strains, any beta-tubulin variants, sample collection

locations, and substrate types. S4 Table contains all *C. briggsae* and *C. tropicalis* isotype reference strains with

beta-tubulin variants and genetically related strains. S5 Table contains details about the CRISPR-Cas9 genome-

edited strains, CRISPR-Cas9 reagents, and oligonucleotide sequences for the deletion of *ben-1* in *C. briggsae*

and *C. tropicalis*. S6 Table contains the results from the fecundity assays of the *C. briggsae* reference strain

(AF16) and the *Cbr-ben-1* deletion strains.Table S7 contains the results from the fecundity assays of the *C.*

tropicalis strain NIC58 and the *Ctr-ben-1* deletion strains. S8 Table contains the BLOSUM62 and Granthum

scores for amino acid changes in the beta-tubulin genes in the three *Caenorhabditis* species. S9 Table contains

expression breadth (Tau) estimates for beta-tubulin genes calculated on *C. elegans* and *C. briggsae* embryonic

cells by Large *et al.* 2024. S10 Table contains conservation estimates of embryonic expression patterns between

C. elegans and *C. briggsae* with Jensen-Shannon gene distances estimated in Large *et al.* 2024 for beta-tubulin

genes. S11 Table contains the neuronal cell-class expression divergence of beta-tubulin orthologs across *C.*

elegans, *C. briggsae*, and *C. tropicalis*, reported as Jaccard distances obtained from Toker *et al.* 2025. S12

Table contains results from the substrate enrichment analysis. S13 Table contains details about the CRISPR-

Cas9 genome edited strains, CRISPR-Cas9 reagents, and oligonucleotide sequences for the deletion of *ben-1*

in *P. pacificus*. S14 Table contains all beta-tubulin transcript IDs in the three *Caenorhabditis* species. S15 Table

contains the manual curation of SVs found in beta-tubulin genes. S16 Table contains the amino acid sequences

for the BEN-1 proteins from the three *Caenorhabditis* species.

FUNDING

Amanda O. Shaver was funded by the National Institutes of Health grant F32AI181342. Ryan McKeown was

supported by Northwestern University's Biotechnology Training Program (NIH T32 GM008449). This work was

supported by the National Institutes of Health grant R01AI153088 to E.C.A.

CRediT AUTHOR STATEMENT

Conceptualization: ECA

Data curation: AOS, RM

Formal analysis: AOS, RM

Funding acquisition: ECA

Investigation: AOS, JMRO, JBC, DWH, JSF, SD, EJR

Methodology: AOS, RM

Project administration: ECA

Resources: ECA

Software: AOS, RM

Supervision: ECA

Validation: AOS, RM

Visualization: AOS, RM

Writing - original draft: AOS

Writing - reviewing & editing: AOS, RM, ECA

DECLARATION OF COMPETING INTERESTS

The authors have declared that no competing interests exist.

ACKNOWLEDGEMENTS

We would like to thank members of the Andersen laboratory for their feedback and helpful comments on this

manuscript. We thank members of the *C. elegans* community for collecting the diverse set *Caenorhabditis* strains

included in this study and the *Caenorhabditis* Natural Diversity Resource (NSF Capacity grant 2224885) for

providing the strains for this study. We thank WormBase for providing the amino acid sequences for all six *C.*

elegans beta-tubulin proteins [60]. We thank WormBase Parasite for providing the amino acid sequences for

TBB-ISOTYPE-1 beta-tubulin protein in *H. contortus* [41].

REFERENCES

1. Kotze AC, Gilleard JS, Doyle SR, Prichard RK. Challenges and opportunities for the adoption of molecular diagnostics for anthelmintic resistance. Int J Forecast. 2020. Available: https://www.sciencedirect.com/science/article/pii/S2211320720300476

2. Neff NF, Thomas JH, Grisafi P, Botstein D. Isolation of the beta-tubulin gene from yeast and demonstration of its essential function in vivo. Cell. 1983;33: 211–219.

3. Laclette JP, Guerra G, Zetina C. Inhibition of tubulin polymerization by mebendazole. Biochem Biophys Res Commun. 1980;92: 417–423.

4. Ireland CM, Gull K, Gutteridge WE, Pogson CI. The interaction of benzimidazole carbamates with mammalian microtobule protein. Biochem Pharmacol. 1979;28: 2680–2682.

5. Chalfie M, Thomson JN. Structural and functional diversity in the neuronal microtubules of Caenorhabditis elegans. J Cell Biol. 1982;93: 15–23.

6. Theodorides VJ, Scott GC, Lademan MS. Strains of Haemonchus contortus resistant against benzimidazole anthelmintics. Am J Vet Res. 1970;31: 859–863.

7. Roos MH, Kwa MSG, Grant WN. New genetic and practical implications of selection for anthelmintic resistance in parasitic nematodes. Parasitol Today. 1995;11: 148–150.

8. Beech RN, Prichard RK, Scott ME. Genetic variability of the beta-tubulin genes in benzimidazole-susceptible and -resistant strains of Haemonchus contortus. Genetics. 1994;138: 103–110.

9. Sangster NC, Prichard RK, Lacey E. Tubulin and benzimidazole-resistance in Trichostrongylus colubriformis (Nematoda). J Parasitol. 1985;71: 645–651.

10. Martínez-Valladares M, Valderas-García E, Gandasegui J, Skuce P, Morrison A, Castilla Gómez de Agüero V, et al. Teladorsagia circumcincta beta tubulin: the presence of the E198L polymorphism on its own is associated with benzimidazole resistance. Parasit Vectors. 2020;13: 453.

11. Venkatesan A, Jimenez Castro PD, Morosetti A, Horvath H, Chen R, Redman E, et al. Molecular evidence of widespread benzimidazole drug resistance in Ancylostoma caninum from domestic dogs throughout the USA and discovery of a novel β-tubulin benzimidazole resistance mutation. PLoS Pathog. 2023;19: e1011146.

12. Furtado LFV, Dos Santos TR, de Oliveira VNGM, Rabelo ÉML. Genotypic profile of benzimidazole resistance associated with SNP F167Y in the beta-tubulin gene of Necator americanus helminths obtained from Brazilian populations. Infect Genet Evol. 2020;86: 104594.

13. Schwenkenbecher JM, Albonico M, Bickle Q, Kaplan RM. Characterization of beta-tubulin genes in hookworms and investigation of resistance-associated mutations using real-time PCR. Mol Biochem Parasitol. 2007;156: 167–174.

14. Hahnel SR, Zdraljevic S, Rodriguez BC, Zhao Y, McGrath PT, Andersen EC. Extreme allelic heterogeneity at a Caenorhabditis elegans beta-tubulin locus explains natural resistance to benzimidazoles. PLoS Pathog. 2018;14: e1007226.

15. Dilks CM, Hahnel SR, Sheng Q, Long L, McGrath PT, Andersen EC. Quantitative benzimidazole resistance and fitness effects of parasitic nematode beta-tubulin alleles. Int J Parasitol Drugs Drug Resist. 2020;14: 28–36.

16. Dilks CM, Koury EJ, Buchanan CM, Andersen EC. Newly identified parasitic nematode beta-tubulin alleles

confer resistance to benzimidazoles. Int J Parasitol Drugs Drug Resist. 2021;17: 168–175.

17. Wit J, Dilks CM, Andersen EC. Complementary Approaches with Free-living and Parasitic Nematodes to
Understanding Anthelmintic Resistance. Trends Parasitol. 2021;37: 240–250.

18. Saunders GI, Wasmuth JD, Beech R, Laing R, Hunt M, Naghra H, et al. Characterization and comparative
analysis of the complete Haemonchus contortus beta-tubulin gene family and implications for
benzimidazole resistance in strongylid nematodes. Int J Parasitol. 2013;43: 465–475.

19. Silvestre A, Cabaret J. Mutation in position 167 of isotype 1 beta-tubulin gene of Trichostrongylid
nematodes: role in benzimidazole resistance? Mol Biochem Parasitol. 2002;120: 297–300.

20. Ghisi M, Kaminsky R, Mäser P. Phenotyping and genotyping of Haemonchus contortus isolates reveals a
new putative candidate mutation for benzimidazole resistance in nematodes. Vet Parasitol. 2007;144:
313–320.

21. de Lourdes Mottier M, Prichard RK. Genetic analysis of a relationship between macrocyclic lactone and
benzimidazole anthelmintic selection on Haemonchus contortus. Pharmacogenet Genomics. 2008;18:
129–140.

22. Kwa MS, Veenstra JG, Roos MH. Benzimidazole resistance in Haemonchus contortus is correlated with a
conserved mutation at amino acid 200 in beta-tubulin isotype 1. Mol Biochem Parasitol. 1994;63: 299–
303.

23. Mohammedsalih KM, Krücken J, Khalafalla A, Bashar A, Juma F-R, Abakar A, et al. New codon 198 β-
tubulin polymorphisms in highly benzimidazole resistant Haemonchus contortus from goats in three
different states in Sudan. Parasit Vectors. 2020;13: 114.

24. Kwa MS, Kooyman FN, Boersema JH, Roos MH. Effect of selection for benzimidazole resistance in
Haemonchus contortus on beta-tubulin isotype 1 and isotype 2 genes. Biochem Biophys Res Commun.
1993;191: 413–419.

25. Hurd DD. Tubulins in C. elegans. WormBook. 2018;2018: 1–32.

26. Driscoll M, Dean E, Reilly E, Bergholz E, Chalfie M. Genetic and molecular analysis of a Caenorhabditis
elegans beta-tubulin that conveys benzimidazole sensitivity. J Cell Biol. 1989;109: 2993–3003.

27. Collins JB, Stone SA, Koury EJ, Paredes AG, Shao F, Lovato C, et al. Quantitative tests of albendazole
resistance in Caenorhabditis elegans beta-tubulin mutants. Int J Parasitol Drugs Drug Resist. 2024;25:
100556.

28. Wright AJ, Hunter CP. Mutations in a beta-tubulin disrupt spindle orientation and microtubule dynamics in
the early Caenorhabditis elegans embryo. Mol Biol Cell. 2003;14: 4512–4525.

29. Ellis GC, Phillips JB, O'Rourke S, Lyczak R, Bowerman B. Maternally expressed and partially redundant
beta-tubulins in Caenorhabditis elegans are autoregulated. J Cell Sci. 2004;117: 457–464.

30. Lu C, Srayko M, Mains PE. The Caenorhabditis elegans microtubule-severing complex MEI-1/MEI-2
katanin interacts differently with two superficially redundant beta-tubulin isotypes. Mol Biol Cell. 2004;15:
142–150.

31. Pallotto LM, Dilks CM, Park Y-J, Smit RB, Lu BT, Gopalakrishnan C, et al. Interactions of Caenorhabditis
elegans β-tubulins with the microtubule inhibitor and anthelmintic drug albendazole. Genetics. 2022;221.
doi:10.1093/genetics/iyac093

32. Boulin T, Etchberger JF, Hobert O. Reporter gene fusions. WormBook. 2006; 1–23.

33. Gibson SB, Ness-Cohn E, Andersen EC. Benzimidazoles cause lethality by inhibiting the function of
Caenorhabditis elegans neuronal beta-tubulin. Int J Parasitol Drugs Drug Resist. 2022;20: 89–96.

34. Hahnel, Dilks, Heisler, Andersen. Caenorhabditis elegans in anthelmintic research–Old model, new
perspectives. Int J Educ Vocat Guid. 2020. Available:
https://www.sciencedirect.com/science/article/pii/S2211320720300312

35. Mariene GM, Wasmuth JD. Genome assembly variation and its implications for gene discovery in
nematodes. Int J Parasitol. 2025. doi:10.1016/j.ijpara.2025.01.004

36. Crombie TA, McKeown R, Moya ND, Evans KS, Widmayer SJ, LaGrassa V, et al. CaeNDR, the
Caenorhabditis Natural Diversity Resource. Nucleic Acids Res. 2023. doi:10.1093/nar/gkad887

37. Shaver AO, Miller IR, Schaye ES, Moya ND, Collins JB, Wit J, et al. Quantifying the fitness effects of
resistance alleles with and without anthelmintic selection pressure using Caenorhabditis elegans. PLoS
Pathog. 2024;20: e1012245.

38. Zhang G, Roberto NM, Lee D, Hahnel SR, Andersen EC. The impact of species-wide gene expression
variation on Caenorhabditis elegans complex traits. Nat Commun. 2022;13: 3462.

39. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. Proc Natl Acad Sci U S A.
1992;89: 10915–10919.

40. Grantham R. Amino acid difference formula to help explain protein evolution. Science. 1974;185: 862–
864.

41. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite - a comprehensive resource for
helminth genomics. Mol Biochem Parasitol. 2017;215: 2–10.

42. Large CRL, Khanal R, Hillier L, Huynh C, Kubo C, Kim J, et al. Lineage-resolved analysis of embryonic
gene expression evolution in *C. elegans* and *C. briggsae*. Science. 2025;388: eadu8249.

43. Toker IA, Ripoll-Sánchez L, Geiger LT, Sussfeld A, Saini KS, Beets I, et al. Divergence in neuronal
signaling pathways despite conserved neuronal identity among Caenorhabditis species. Curr Biol.
2025;35: 2927–2945.e7.

44. Andersen EC, Gerke JP, Shapiro JA, Crissman JR, Ghosh R, Bloom JS, et al. Chromosome-scale
selective sweeps shape Caenorhabditis elegans genomic diversity. Nat Genet. 2012;44: 285–290.

45. Zhang G, Mostad JD, Andersen EC. Natural variation in fecundity is correlated with species-wide levels of
divergence in Caenorhabditis elegans. G3 . 2021;11. doi:10.1093/g3journal/jkab168

46. Laing ST, Ivens A, Laing R, Ravikumar S, Butler V, Woods DJ, et al. Characterization of the xenobiotic
response of Caenorhabditis elegans to the anthelmintic drug albendazole and the identification of novel
drug glucoside metabolites. Biochem J. 2010;432: 505–514.

47. Collins JB, Dilks CM, Hahnel SR, Rodriguez B, Fox BW, Redman E, et al. Naturally occurring variation in
a cytochrome P450 modifies thiabendazole responses independently of beta-tubulin. PLoS Pathog.
2025;21: e1012602.

48. Athanasouli M, Witte H, Weiler C, Loschko T, Eberhardt G, Sommer RJ, et al. Comparative genomics and
community curation further improve gene annotations in the nematode Pristionchus pacificus. BMC
Genomics. 2020;21: 708.

49. Doyle SR, Tracey A, Laing R, Holroyd N, Bartley D, Bazant W, et al. Genomic and transcriptomic variation
defines the chromosome-scale assembly of Haemonchus contortus, a model gastrointestinal worm.

Commun Biol. 2020;3: 656.

50. Doyle SR. Improving helminth genome resources in the post-genomic era. Trends Parasitol. 2022 [cited 4
Jul 2022]. Available: https://pubag.nal.usda.gov/catalog/7806127

51. Montarry J, Mimee B, Danchin EGJ, Koutsovoulos GD, Ste-Croix DT, Grenier E. Recent Advances in
Population Genomics of Plant-Parasitic Nematodes. Phytopathology. 2021;111: 40–48.

52. Stevens L, Moya ND, Tanny RE, Gibson SB, Tracey A, Na H, et al. Chromosome-level reference
genomes for two strains of Caenorhabditis briggsae: An improved platform for comparative genomics.
Genome Biol Evol. 2022;14. doi:10.1093/gbe/evac042

53. Moya ND, Stevens L, Miller IR, Sokol CE, Galindo JL, Bardas AD, et al. Novel and improved
Caenorhabditis briggsae gene models generated by community curation. BMC Genomics. 2023;24: 486.

54. Luke M, Noble J, Yuen L, Stevens N, Moya R. Selfing is the safest sex for Caenorhabditis tropicalis. eLife.
2021;10.

55. Geldhof P, Murray L, Couthier A, Gilleard JS, McLauchlan G, Knox DP, et al. Testing the efficacy of RNA
interference in Haemonchus contortus. Int J Parasitol. 2006;36: 801–810.

56. Roose S, Avramenko RW, Pollo SMJ, Wasmuth JD, Ame S, Ayana M, et al. Characterization of the β-
tubulin gene family in Ascaris lumbricoides and Ascaris suum and its implication for the molecular
detection of benzimidazole resistance. PLoS Negl Trop Dis. 2021;15: e0009777.

57. Krücken J, Fraundorfer K, Mugisha JC, Ramünke S, Sifft KC, Geus D, et al. Reduced efficacy of
albendazole against Ascaris lumbricoides in Rwandan schoolchildren. Int J Parasitol Drugs Drug Resist.
2017;7: 262–271.

58. Palma A, Matamoros G, Escobar D, Sánchez AL, Fontecha G. Absence of mutations associated with
resistance to benzimidazole in the beta-tubulin gene of Ascaris suum. Rev Soc Bras Med Trop. 2020;53:
e20190155.

59. Collins JB, Andersen EC. The turkey ascarid, Ascaridia dissimilis, as a model genetic system. Int J
Parasitol. 2023;53: 405–409.

60. Sternberg PW, Van Auken K, Wang Q, Wright A, Yook K, Zarowiecki M, et al. WormBase 2024: status
and transitioning to Alliance infrastructure. Genetics. 2024;227. doi:10.1093/genetics/iyae050

61. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and
applications. BMC Bioinformatics. 2009;10: 421.

62. Noble LM, Yuen J, Stevens L, Moya N, Persaud R, Moscatelli M, et al. Selfing is the safest sex for
Caenorhabditis tropicalis. eLife. 2021. doi:10.7554/elife.62587

63. Pertea G, Pertea M. GFF utilities: GffRead and GffCompare. F1000Res. 2020;9: 304.

64. Rausch T, Zichner T, Schlattl A, Stütz AM, Benes V, Korbel JO. DELLY: structural variant discovery by
integrated paired-end and split-read analysis. Bioinformatics. 2012;28: i333–i339.

65. Lesack K, Mariene GM, Andersen EC, Wasmuth JD. Different structural variant prediction tools yield
considerably different results in Caenorhabditis elegans. PLoS One. 2022;17: e0278424.

66. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and
BCFtools. Gigascience. 2021;10. doi:10.1093/gigascience/giab008

67. Shaver AO, Wit J, Dilks CM, Crombie TA, Li H, Aroian RV, et al. Variation in anthelmintic responses are

driven by genetic differences among diverse C. elegans wild strains. PLoS Pathog. 2023;19: e1011285.

68. Ortiz EM. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for phylogenetic analysis.
Zenodo; 2019. doi:10.5281/ZENODO.2540861

69. Caro H, Dollin S, Biton A, Brancotte B, Desvillechabrol D, Dufresne Y, et al. BioConvert: a comprehensive
format converter for life sciences. NAR Genom Bioinform. 2023;5: lqad074.

70. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol
Biol Evol. 1987;4: 406–425.

71. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc Bioinformatics. 2020;69: e96.

72. Andersen EC, Bloom JS, Gerke JP, Kruglyak L. A variant in the neuropeptide receptor npr-1 is a major
determinant of Caenorhabditis elegans growth and physiology. PLoS Genet. 2014;10: e1004156.

73. Andersen EC, Shimko TC, Crissman JR, Ghosh R, Bloom JS, Seidel HS, et al. A Powerful New
Quantitative Genetics Platform, Combining Caenorhabditis elegans High-Throughput Fitness Assays with
a Large Collection of Recombinant Strains. G3 . 2015;5: 911–920.

74. Zamanian M, Cook DE, Zdraljevic S, Brady SC, Lee D, Lee J, et al. Discovery of genomic intervals that
underlie nematode responses to benzimidazoles. PLoS Negl Trop Dis. 2018;12: e0006368.

75. Widmayer SJ, Crombie TA, Nyaanga JN, Evans KS, Andersen EC. C. elegans toxicant responses vary
among genetically diverse individuals. Toxicology. 2022; 153292.

76. Shaver A, Andersen E. High-throughput larval development assay (HTLDA) v1. 2025.
doi:10.17504/protocols.io.n92ldrqwng5b/v1

77. Nyaanga J, Crombie TA, Widmayer SJ, Andersen EC. easyXpress: An R package to analyze and
visualize high-throughput C. elegans microscopy data generated using CellProfiler. PLoS One. 2021;16:
e0252000.

78. Wählby C, Kamentsky L, Liu ZH, Riklin-Raviv T, Conery AL, O'Rourke EJ, et al. An image analysis toolbox
for high-throughput C. elegans assays. Nat Methods. 2012;9: 714–716.

79. Wang C, Vidal B, Sural S, Loer C, Aguilar GR, Merritt DM, et al. A neurotransmitter atlas of C. elegans
males and hermaphrodites. Elife. 2024;13. doi:10.7554/elife.95402.3

80. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature. 2024;630: 493–500.

81. Han Z, Lo W-S, Lightfoot JW, Witte H, Sun S, Sommer RJ. Improving transgenesis efficiency and
CRISPR-associated tools through Codon optimization and native intron addition in Pristionchus
nematodes. Genetics. 2020;216: 947–956.