Lipid packing frustration in the outer leaflet of the plasma membrane prevents scission of caveolae

Larsson, E.[1], Kabedev, A.[2], Pace H.[3]., Lindwall J.[1] Fouzia, B.[3], Parton, R.G.[4,5], Bergström, C.A.S.[2], Parmryd, I.[6], Bally, M.[3], Lundmark, R.[1],*.

Affiliations

1. Department of Medical and Translational Biology and Laboratory for Molecular Infection Medicine Sweden, Umeå Centre for Microbial Research, SciLifeLab, Umeå University, 901 87, Umeå, Sweden.

2. Department of Pharmacy, Uppsala University, Uppsala Biomedical Center, 751 23 Uppsala, Sweden.

3. Department of Clinical Microbiology, Wallenberg Centre for Molecular Medicine & Umeå Centre for Microbial Research, Umeå University, 901 85, Umeå, Sweden.

4. Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland, Australia.

5. Centre for Microscopy and Microanalysis, The University of Queensland, Brisbane, Queensland, Australia.

6. Department of Medical Biochemistry and Cell Biology, University of Gothenburg, 405 30 Göteborg, Sweden.

*Corresponding author: Richard Lundmark, Department of Medical and translational biology, Umeå University, 901 87 Umeå, Sweden, phone: +46 706202464, email: richard.lundmark@umu.se.

Running title: Lipid packing affects caveola dynamics

Keywords: Caveolae, lipid packing, membrane curvature, membrane scission, lipid model membrane, molecular dynamics simulations, Dyngo-4a.

Abstract

Lipid packing is a fundamental characteristic of bilayer membranes. It affects all membrane-associated processes ranging from curvature generation to membrane fission. Yet, we lack detailed, mechanistic understanding of how lipid packing directly affects these processes in cellular membranes. Here, we address this by focusing on caveolae, small Ω-shaped invaginations of the plasma membrane which serve as key regulators of cellular lipid sorting and mechano-responses. In addition to caveolae coat proteins, the lipid membrane is a core component of caveolae that critically impacts both the biogenesis, morphology and stability of such membrane invaginations. We show that the small compound Dyngo-4a inserts into the membrane, resulting in a dramatic dynamin-independent inhibition of caveola scission. Analysis of model membranes in combination with molecular dynamics simulations revealed that a substantial amount of Dyngo-4a was inserted and positioned at the level of cholesterol in the bilayer. Dyngo-4a-treatment resulted in decreased lipid packing in the outer leaflet of the plasma membrane preventing scission without affecting caveola morphology, caveolae-associated proteins, or the overall membrane stiffness. Artificially increasing plasma membrane cholesterol levels was found to counteract the block in caveola scission caused by Dyngo-4a-mediated lipid packing frustration. Therefore, we propose that the packing of cholesterol in the outer leaflet of the plasma membrane critically controls the confinement of caveolae to the plasma membrane.

Significance statement

Larsson et al., demonstrate that lipid packing critically impacts the stability of small Ω-shaped plasma membrane cavities termed caveolae. The scission of caveolae is dynamin independent and halted by outer leaflet incorporation of the small compound Dyngo-4a, which leads to lipid packing frustration as characterized by the authors. The block in caveola scission can be counteracted by increased PM levels of cholesterol which is enriched in caveolae and increases the lipid packing. This proof-of-principle study show that lipid packing control scission which can be used by cells to generate membrane vesicles that remain stably associated with the donor membrane for an extended period of time.

Introduction

The plasma membrane (PM) of eukaryotic cells has a unique assymetric lipid composition with up to 50% cholesterol (chol). The PM lipid composition and asymmetry has a major role in lipid packing and thus also membrane remodelling processes such as the budding and scission of membrane vesicles (1, 2). During scission, the lipids in the membrane neck of budding vesicles reorganise, which results in membrane fission and release of the vesicle. The process is mediated by proteins such as dynamin (Dyn2), but the lipid packing of the membrane is also critical (1). Classically, most attention has been directed towards vesicles that rapidly undergo scission. However, some vesicles, remain associated with the donor membrane for a more extended period of time thereby forming Ω-shaped cavities in the PM (3, 4). Little is known about how lipid packing contributes to stalling scission of such membrane structures. Examples of more stable membrane cavities are caveolae, which play important roles in lipid trafficking and monitoring of the PM composition and tension. Such membrane structures assemble by cooperative interaction of the integral membrane proteins caveolins 1-3 and peripherally attached cavins 1-4, which form the caveola coat (5, 6). The Ω-shaped structure is stabilized by the ATPase EH domain containing 2 (EHD2), which multimerizes at the caveola (singular) neck and prevents scission from the PM (3). Assembly of the caveola coat is highly dependent on distinct lipid species. Caveolin 1 (Cav1) assembly is driven by high chol content, which results in the enrichment of sphingomyelin (SM) and glycosphingolipids in the outer lipid leaflet (7-9). Modelling of the lipid packing around the Cav1 oligomer shows that the inner leaflet lipids are displaced by the Cav1 oligomer, and that chol molecules together with sphingolipids form a disordered monolayer in the outer leaflet facing the oligomer (9). The inner lipid leaflet is also enriched in phosphatidylserine (PS) and phosphatidylinositol 4,5 bisphosphate (PIP2) due to their interactions with cavins and EHD2 (7). This generates a unique membrane domain with distinct lipid packing and a very high curvature.

Although caveolae are relatively stable PM structures, they exhibit dynamics in terms of lateral mobility and rounds of scission from and fusion with the membrane. The scission of caveolae

from the PM is controlled by the lipid composition (10). Caveola-enrichment of chol and glycosphingolipids has been found to promote scission, while sphingomyelin prevents it, suggesting that lipid packing might be important for this process (10). Recently, it was shown that the scission process is independent of dynamin (11), a GTPase which mediates scission of clathrin-coated pits from the PM (12). In fact, dynamin2 acted to prevent scission of caveolae (11).

Intriguingly, caveola internalisation has been shown to be inhibited by small pyrimidyl-based compounds such as dynasore and Dyngo-4a (11, 13). These compounds inhibit dynamin-mediated scission and the uptake of, for example, the transferrin receptor. Yet, cell viability and endocytosis via clathrin independent carriers is unaffected (14-16). Interestingly, Dyngo-4a constrains caveolae dynamics also in cells depleted of Dyn2 (11), suggesting that the drug acts via another mechanism. Furthermore, both Dynasore and Dyngo-4a have been shown to have wider effects on cells, for example affecting chol, lipid rafts, lamellipodia formation and cell migration (17, 18). Dynasore was shown to act protectively against chol-dependent pore forming toxins by disrupting PM lipid rafts (19). Dynamin depletion or inhibition using a dynamin inhibitor peptide showed that the effect was dynamin independent (19). Dyngo-4a-treatment has been shown to block membrane ruffling and inhibit fluid phase endocytosis in dynamin triple knockout (TKO) cells (20). These results clearly show that these compounds affect cellular processes at the plasma membrane that are dynamin independent and suggests that their actions are multifaceted. These insights have clouded the view on how this frequently used drug should be utilized in experiments addressing endocytosis and raise questions on how the resulting data should be interpreted.

Since Dyngo-4a-treatment dramatically affected caveola scission in Dyn2-depleted HeLa cells, we aimed to use this compound to reveal principles that control caveola dynamics. To get an understanding of the underlying mechanisms, we used interdisciplinary experimental setups and show that Dyngo-4a binds and inserts beneath the headgroups of the plasma membrane affecting the lipid order and membrane remodeling.

Results

The drug Dyngo-4a confines caveolae at the plasma membrane independently of dynamin

The small inhibitory molecule Dyngo-4a (Fig. S1A) has been demonstrated to increase the number of caveolae per PM surface area (13), and to have a dramatic stabilizing effect on caveola dynamics even in cells depleted of Dyn2 (11). This suggested that the drug acts via an unknown mechanism, which, if clarified, might advance our understanding of the principles that control caveola dynamics. To assure that the effect of Dyngo-4a was totally independent of all dynamin isoforms we utilized the well-established dynamin TKO cell line where the addition of tamoxifen can be used to silence the three different isoforms of *dnm* (20). This abolishes dynamin protein expression compared to the untreated control (ctrl) cells (Fig. 1A) (20). Analysis of the of the caveolae core proteins Cav1, cavin1 and EHD2 by fluorescence microscopy showed that, similarly to control cells, typical punctate caveolae structures covered the plasma membrane of TKO cells (Fig. 1B and Fig. S1B-C). Quantification of the amount of caveola revealed no significant difference between the wild type (wt) and knock out cells (Fig. 1C), in agreement with previous results (21).

To monitor the dynamics of caveolae, Cav1-GFP was transiently expressed in Tamoxifen-treated TKO cells. Cells were treated for 30 min with either Dyngo-4a or DMSO and then imaged using total internal reflection fluorescence (TIRF) microscopy to visualize Cav1-GFP at, or in close proximity to the basal PM (Video 1 and 2). Using single particle tracking, where Cav1-GFP punctae of a specific size and fluorescent intensity were identified as caveolae, we could determine their lateral mobility (displacement length) and axial movements in and out of the TIRF field (duration time) (Fig. 1D-F and Fig. S1D) (10, 11, 22, 23). Tracking of a large number of caveolae from both ctrl and Dyngo-4a-treated cells, showed that the average PM duration time was significantly increased in the Dyngo-4a-treated cells, indicative of stalled scission (Fig. 1G). Previous work has shown that, in addition to caveola scission, membrane undulations interfering with the TIRF field contribute to axial mobility, resulting in an overrepresentation of short duration tracks (10, 11). However, in Dyngo-4a-treated cells, the majority of the caveolae present at time zero were tracked persistently throughout the movie (Fig. 1G). Almost no short duration tracks were detected in agreement with that both caveola scission and membrane undulations are blocked. Disassembly of the caveola complex would result in progressive loss of caveolae from the plasma membrane and thereby shorter track duration times. We further found that the average displacement length was greatly reduced in the Dyngo-4a-treated cells as compared to ctrl-treated cells (Fig. 1E), which is in agreement with fewer scissoned and mobile caveolae as well as less membrane undulations. Detailed analysis of the lateral mobility of caveolae showed that both directed transport and Brownian

motion decreased even in tracks with long duration time, following Dyngo-4a treatment (Fig. 1F). This demonstrated that Dyngo-4a-treatment prevents scission of caveolae from the PM and reduce their lateral movement even in cells completely devoid of dynamins.

Dyngo-4a adsorbs and inserts into the lipid bilayer

Despite the striking effect on caveola dynamics, Dyngo-4a has no apparent effect on the recruitment and assembly of the individual caveola coat proteins EHD2 and cavin1 which remain colocalized with caveolae after 30 minutes of Dyngo-4a treatment (11). Because Dyngo-4a is an amphiphilic compound (Fig. S1A) we reasoned that it might interact with the lipid bilayer and influence the properties of the membrane. To decipher if Dyngo-4a inserts into membranes, we created a lipid monolayer of 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC) lipids suspended at an air-water interphase and monitored the changes in the lateral pressure after injection of the compound using Langmuir-Blodgett trough tensiometry (Fig. 2A, illustration)(24). Injection of Dyngo-4a under the equilibrated lipid monolayer resulted in a steep rise of the surface pressure (Fig. 2A). These results show that Dyngo-4a indeed inserts between the lipids of the planar monolayer. Injection of a control solution containing the same DMSO concentration as the Dyngo-4a solution, had a minor effect on the surface pressure (Fig. 2A), in agreement with the known interaction between DMSO lipid bilayers (25).

To further confirm the surface interaction between Dyngo-4a and the membrane, we used quartz crystal microbalance with dissipation monitoring (QCM-D). This technique uses an acoustic sensor to track adsorption and desorption to the sensor surface, which is expressed as a function of frequency shift (Δf) (Fig. 2B, illustration)(26). Supported lipid bilayers (SLBs) of POPC were generated on the sensor surface via vesicle fusion. The system was first equilibrated in buffer containing DMSO. Upon addition of Dyngo-4a in the same buffer the frequency decreased substantially, indicating that Dyngo-4a adsorbed to the SLB (Fig. 2B). Rinsing the SLB with buffer lead to a desorption of Dyngo-4a, showing that binding was at least partially reversible in DMSO-containing buffer (Fig. 2B). After returning to DMSO-free buffer, the system was of low enough softness (Fig. S2), in order to use the Sauerbrey equation to convert the frequency change into adsorbed mass. A comparison of the calculated mass of the SLB alone with the calculated mass of the Dyngo-4a-containing-SLB allowed us to estimate that the amount of Dyngo-4a retained after washing (0.9 ± 0.3 Hz), accounted for 6 ±

2 mol% (Fig. 2B). Taking this value as a reference, the amount of Dyngo-4a incorporated into the bilayer before rinsing, which resulted in a frequency shift of 3 Hz, was extrapolated to be about 15.6 mol%.

Dyngo-4a inserts just beneath the head group region of the lipid bilayer at a similar position as cholesterol

To gain detailed information about how Dyngo-4a is positioned in the membrane, we performed molecular dynamic (MD) simulations where Dyngo-4a was added to a POPC bilayer. The free sampling simulations were performed for 200 ns and showed that Dyngo-4a was incorporated into the bilayer underneath the headgroups (Fig. 3A). This is in agreement with the hydrophobic properties of the compound. To gain information about the energy landscape with barriers and preferred regions of positioning of Dyngo-4a within the membrane, Dyngo-4a was sampled through the POPC membrane using simulations with a convolved adaptive biasing potential based on the Accelerated Weight Histogram (AWH) method. The potential of mean force (PMF) profile showed that Dyngo-4a preferred the area of the POPC membrane, around 1.1 to 1.2 nm from the bilayer center (Fig. 3B). Interestingly, an energy difference of approximately 40 kJ/mol between the lowest PMF point and the center of the membrane (Figure 3B) suggests that Dyngo-4a is unlikely to cross the POPC bilayer spontaneously.

As chol is highly abundant within the PM, we hypothesized that Dyngo-4a might compete with chol for the space within the lipid tail region. AWH-enhanced simulations of a chol molecule interacting with the POPC membrane showed that the center of mass located in regions (0.5 to 1.0 nm) overlapping with the preferred region of Dyngo-4a (Fig. 3C and B). To test if addition of chol to the membrane would alter the position of Dyngo-4a, MD simulations were performd on a membrane consisting of 70% POPC and 30% chol. This showed that the chol molecules localized to the POPC tail region as expected. In this setup Dyngo-4a was still incorporated under the headgroups (Fig. 3D). However, simulation employing AWH showed that with chol in the membrane, the preferred position of Dyngo-4a was closer to the head groups compared to when chol was absent (Fig. E). Furthermore, the mid-membrane energy barrier that Dyngo-4a would need to overcome to pass the membrane was increased to approximately 60 kJ/mol (Fig. 3E). These results further support that Dyngo-4a insert into the lipid bilayer affecting the lipid packing. Notably, when more than one Dyngo-4a molecule was added to the limited membrane area of the simulation, single molecules inserted in the bilayer, but most of the

molecules aggregated and formed stacks that were not able to entirely penetrate through the headgroup region of the membrane (Fig. S3).

Membrane insertion of Dyngo-4a decreases lipid packing in the outer leaflet of the plasma membrane

Next, we aimed to address how membrane incorporation of Dyngo-4a affected the properties of the PM. Interestingly, we found that Dyngo-4a majorly impaired the typical undulations of plasma membrane protrusion and retraction in a concentration-dependent manner (Fig.4A-B and S4A). This resulted in a "frozen" cell membrane with no membrane ruffling and lamellipodia formation which has been reported previously as a dynamin-independent effect of Dyngo-4a (20). Similarly, we observed a steady and continuous concentration-dependent decrease in lateral mobility of caveolae as concentrations of Dyngo-4a were increased (Fig. S4B). Quantification of the effect on lateral movement over time revealed a fast initial decrease in response to Dyngo-4a and that the maximum level of inhibition was reached after 20 to 25 minutes (Fig. S4C). This is comparable to effects on clathrin-coated vesicle (CCV) endocytosis as measured by the internalization of transferrin (Tfn), although the inhibition of this process was slightly faster and completely blocked already at 10 µM of Dyngo-4a (Fig. S4C). Taken together, these data showed that Dyngo-4a affects several active processes that involve remodelling of the PM. To make sure that highly curved regions of the PM such as caveolae did not flatten in response to the membrane freezing effect, we analyzed the PM using electron microscopy. In agreement with previous observations (16), the typical Ω-shaped caveolae could be detected also in Dyngo-4a treated cells (Fig. 4C).

To determine if Dyngo-4a affected the mechanical properties of the membrane, we quantified Young's modulus (a measure of the elastic properties of materials) using an atomic force microscopy (AFM) -based nanomechanical setup (Fig. S4A). Young's moduli for ctrl cells and cells treated with Dyngo-4a were obtained by fitting force-indentation curves with the Hertz model (Fig. S4D). Quantification revealed that Dyngo-4a treatment had no significant effect on the elastic modulus, showing that the drug does not affect the overall cell stiffness (Fig. 4D). It is likely that membrane insertion of Dyngo-4a might influence lipid packing due to its hydrophobic nature. To analyze if Dyngo-4a affected this property of the plasma membrane bilayer, we used the fluorescent probe 6-dodecanoyl-2-[*N*-methyl-*N*-(carboxymethyl)amino] naphthalene (C-Laurdan), a probe commonly used to measure the degree of lipid packing in

the outer leaflet of the PM (27). The fluorochrome of the probe inserts into the interphase between the lipid headgroups and the acyl chains of the lipid bilayer, and will emit at different wavelengths depending on the level of water penetration in the membrane. First, we confirmed that Dyngo-4a had a similar effect on caveolae duration in the presence and absence of C-Laurdan (Fig. S4E). C-Laurdan was added to cells treated with Dyngo-4a and z-stacks were acquired using fluorescent microscopy. Using the intensities from the different emission channels, the relative membrane order was established by calculating the generalized polarization (GP) value of regions of the PM (Fig. 4E, left panel). Quantification of the GP-values showed that the PM of Dyngo-4a treated cells displayed significantly lower mean GP-value than ctrl-treated cells (Fig. 4F), showing that the compound decreases the lipid packing. Based on this, we propose that membrane insertion of Dyngo-4a increases the hydrated hydrophilic area in the outer lipid leaflet.

Increased PM levels of cholesterol counteracts the inhibition in caveola scission caused by Dyngo-4a-induced lipid packing frustration

Contrary to Dyngo-4a, cholesterol is known to increase lipid packing (28-31). Since our simulation data showed that Dyngo-4a is competing with cholesterol, we hypothesized that Dyngo-4a displaces chol in the outer leaflet of the PM leading to decreased lipid packing which might be the cause of halted caveolae scission. To test if increased levels of chol would restore the ability of caveolae to scission, we used fusogenic liposomes to incorporate chol in the PM (10). However, Dyngo-4a treatment prevented fusion of the fusogenic liposomes (Fig S4F). This suggests that the Dyngo-4a-mediated lipid packing frustration which increases the hydrated hydrophilic area of the outer leaflet also affects membrane fusion. The PM-levels of chol can also be increased by the addition of chol prebound to methyl-β-cyclodexrin (MβCD) as previously reported (32). Using this approach and fluorescently labelled cholesterol, we could visualize efficient incorporation of chol using TIRF microscopy (Fig S4G). This methodology allowed us to analyse and compare caveola duration time in cells treated with Dyngo-4a alone, or followed by the addition of MβCD-chol to control cells. Strikingly, increasing the chol levels following Dyngo-4a treatment resulted in a caveolae track duration similar to control cells not treated with Dyngo-4a (Fig 4G). This rescue effect was not observed in cells where Dyngo-4 treatment was followed by incubation in media alone. These data showed that the block in caveolae scission induced by Dyngo-4a could be counteracted by increased levels of chol in the PM.

Taken together, our data show that membrane incorporation of Dyngo-4a results in lipid packing frustration and halted caveola scission which can be overcome by increasing the PM levels of chol. Based on this we conclude that outer leaflet lipid packing is a critical determinant in the balance between scission and stable PM association of membrane vesicles.

Discussion

Lipid composition and membrane properties are known to be critical determinants of both caveolae biogenesis and endocytosis (33). Yet, determining which distinct bilayer property dictates caveola dynamics has proven difficult. Here, we show that the small pyramidyl-based compound, Dyngo-4a, inserts into membranes and this has a dramatic, dynamin-independent, effect on caveola scission. This enabled the use of this compound to identify lipid packing as a driving force behind confinement of such highly curved invaginations to the PM. Detailed characterization of the membrane interaction of Dyngo-4a using model membranes showed that the compound was inserted into a lipid monolayer. Simulations employing AWH showed that Dyngo-4a was positioned just beneath the headgroup region of the lipids. Interestingly this coincides with the preferred localization of chol and indeed the positioning of Dyngo-4a was influenced by the presence of chol. Using SLBs and the QCM-D system we could determine that a substantial amount of Dyngo-4a adsorbed to the bilayer. We estimated that this would be equivalent to 15.6 mol% of the membrane lipids following incubation with 30 µM Dyngo-4a without washes. Simulation data suggests that Dyngo-4a is restricted to the outer lipid bilayer, which would double the effective concentration in the leaflet. Yet, the simulations also suggested that Dyngo-4a might form complexes that only associate with the lipid head groups without inserting. Therefore, it is difficult to precisely determine the amount of Dyngo-4a inserted into cellular membranes.

Analysis of the PM of cells treated with Dyngo-4a using C-Laurdan revealed an increased water accessibility of the outer leaflet of the membrane as compared to control cells. This showed that incorporation of the compound caused lipid packing frustration increasing the hydrated hydrophilic area, which will influence the properties of the membrane. We could not detect any significant effect on the overall stiffness of the PM implying that the altered packing does not influence the general deformability of the PM. Furthermore, highly curved invaginated caveolae were still present in cells, and previous results show that clathrin

independent endocytic carriers are actively formed in Dyngo-4a-treated cells (16). This suggests that Dyngo-4a incorporation does not have a generic impact on the elasticity of the membrane. However, membrane insertion of Dyngo-4a prevented the axial mobility of caveolae, increasing their average duration time detected at the PM. Dyngo-4a-treated cells were almost completely devoid of short tracks and no scission events could be detected. This also showed that the short tracks typically derived from membrane undulations interfering with the TIRF field were absent. This is in agreement with the finding that PM movements were stalled in Dyngo-4a-treated cells. Therefore, we conclude that the lipid packing frustration caused by Dyngo-4a prevents both the scission of caveolae as well as axial movements caused by membrane undulations. The effect that lipid packing has on scission is in line with the model suggesting that the caveola coat proteins together with specific lipids drives scission as a continuum of membrane curvature generation (10). Lipid packing around the caveolin 8S complex was recently proposed to drive membrane curvature by inducing elastic stresses of tilt and splay in the monolayer proximal to the caveolin complex (34). Furthermore, recent simulations of the lipid packing around the caveolin 8S complex show that the lipids are displaced by the Cav1 oligomer in the inner leaflet. In the outer leaflet facing the complex, 40-70 chol molecules and sphingolipids form a disordered monolayer (9). This provides molecular insight into how lipids are sorted by the 8S complex and how lipid packing might control the curvature and scission of caveolae. Our data suggests that Dyngo-4a is restricted to the outer lipid bilayer in a similar position as chol. Thus, incorporation of Dyngo-4a might be expected to interfere with lipid packing of the disordered monolayer facing the Cav1 oligomer preventing increased curvature leading to membrane scission. Indeed, our data show that artificially increased levels of chol in the PM can counteract the block in caveola scission induced by Dyngo-4a. Chol is enriched in caveolae and known to increase lipid packing whereby these data further support that outer leaflet lipid packing is a critical determinant of membrane scission.

Based on previous literature and our data, it is clear that Dyngo-4a efficiently blocks scission and hence endocytosis of both CCVs and caveolae. We and others have also shown that some clathrin-independent endocytic pathways are still operative under conditions in which Dyngo-4a blocks clathrin and caveolar endocytosis (15, 16). This demonstrates that, despite the general effects on membrane properties reported here, there is considerable specificity in terms of regulation of specific cellular processes. Therefore, the numerous studies that have used this drug to block endocytosis of caveola and CCVs in relation to other endocytic pathways are still

valid. Yet, according to our data, the inhibition of caveolae scission is independent of dynamin and instead due to the insertion of Dyngo-4 in the outer lipid bilayer and thereby decreasing the lipid packing. Currently, we cannot address if this effect is specific to caveolae, or general to the different on- and off-targets effects that has been ascribed to this drug. Dyngo-4a has not been shown to pass membranes and enter cells and our data show that the energy barrier for Dyngo-4a to pass from the outer to the inner leaflet is very high. This would imply that in order for Dyngo-4a to directly interact with dynamin in the cytosol, membrane passage would have to be mediated by, for example, a transporter. Yet, it is difficult to definitively disentangle if Dyngo-4a acts via one or several inhibitory mechanisms. Our comparison of the effects on CCV endocytosis and caveola mobility suggests that CCV endocytosis is more sensitive to Dyngo-4a in terms of time and concentration. However, these differences could either be due to Dyngo-4a acting via more than one mechanism or could be due to the same mechanism of action having quantitatively different impacts on CCVs and caveolae due to their different sensitivities to perturbations of specific properties of the membrane.

Our results show that caution should be taken when conclusions are drawn based on the use of small inhibitory compounds. The drug Pitstop 2, developed to inhibit clathrin assembly and thus clathrin mediated endocytosis has also been reported to freeze the membrane (35). Currently, very few small compounds used for chemical screening are tested for their ability to insert into membranes. Yet, we also show Dyngo-4a can be used to efficiently block endocytosis of both CCV and caveola, and that such compounds can be very useful when addressing how membrane properties affect cellular processes. Based on this work, we propose that lipid packing is instrumental for the balancing of lateral movement and scission of caveolae.

Acknowledgment

We acknowledge the Biochemical Imaging Center (BICU) at Umeå University within the National Microscopy Infrastructure, NMI (VR-RFI 2016-00968) and the Microscopy Australia Research Facility at the Center for Microscopy and Microanalysis at the University of Queensland for providing support and assistance with microscopy. We especially thank Irene Martinez at BICU for assistance and expertise with image analysis and data visualization. We acknowledge the Biochemical Macromolecular Characterization Umeå at Umeå university (BMCU). The computations were enabled by resources provided by the National Academic

Infrastructure for Supercomputing in Sweden (NAISS), partially funded by the Swedish Research Council through grant agreement no. 2022-06725. We thank Jeremy Adler for support with the script for C-Laurdan software analysis and James Rae for assistance with electron microscopy. The work was supported by grants to RL from the Swedish Research Council (dnr 2021-05117) the Swedish Cancer Society (CAN 23 3004 Pj 01 H) and grants to IP from Magnus Bergvall's Foundation. RGP was supported by an Australian Research Council (ARC) Laureate Fellowship (FL210100107). HP and FB were supported by the Knut and Alice Wallenberg foundation.

Author contributions

Elin Larsson and Richard Lundmark designed the research and Elin Larsson performed and analysed the molecular biology, live cell imaging and tracking experiments. Aleksei Kabedev performed modelling experiments, Jakob Lindwall tracking experiments, Hudson Pace QCM-D experiments and analysis, Fouzia Bano AFM experiments and analysis, Ingela Parmryd enabled the C-Laurdan measurements and performed their analysis. Robert Parton enabled the electron microscopy experiments and performed their analysis. Christel A. Bergström and Marta Bally provided financial contribution. Elin Larsson and Richard Lundmark conceived the work and wrote the manuscript, and all authors contributed with text, figures and editing.

Conflict of interest

The authors declare no competing interests.

Materials and methods

Reagents

Dyngo-4a (ab120689, Abcam) was dissolved in DMSO to make a stock solution of 30 mM. For experiments the working concentration was 30 µM unless otherwise stated and Ctrl samples were treated with equivalent DMSO concentration (0.1%) unless otherwise stated. 4-hydroxy-tamoxifen (Sigma) was dissolved in ethanol (99%) to make a stock solution of 10 mM. C-Laurdan was a generous gift from Dr. Bong Rae Cho, Korea University. Human Transferrin Alexa Fluor 647-conjugate (T23366) was bought from Invitrogen. POPC (1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine) was purchased from Avanti Polar Lipids Inc (Alabaster). DMSO (34943), chloroform ($CHCl_3$), and methanol (MeOH) were purchased from Sigma-Aldrich.

Cell lines and constructs

The dynamin triple knockout cells were a kind gift from professor De Camilli (20). The cells were maintained in DMEM supplemented with 10% (vol/vol) FBS and 1x penicillin-streptomycin. The HeLa Flp-In T-REx Caveolin1-mCherry cells were previously generated by us (10). Cells were maintained in DMEM supplemented with 10% (vol/vol) FBS, 100 μg/ml hygromycin B (Thermo Fisher Scientific), and 5 μg/ml blasticidin S HCl (Thermo Fisher Scientific) for plasmid selection. Expression at near endogenous levels was induced by incubation with 0.5 ng/ml doxycycline hyclate (Dox; Sigma-Aldrich) for 16 to 24 h prior to imaging. All cell lines were cultivated at 37°C, 5% CO_2 in a humidified incubator and tested negative for mycoplasma.

Cell treatments

Ablation of *dynamin 1- 3* from Dynamin TKO cells was achieved by culturing the cells with 2 μM 4-hydroxy-tamoxifen for two days. Cells were split to prevent overcrowding and cultured with 300 nM 4-hydroxy-tamoxifen for an additional three days prior to experiment. Protein levels were analyzed by SDS-PAGE and immunoblotting to Amersham Hybond P 0.45 polyvinylidene difluoride membrane (MERCK) using rabbit anti-Dyn2 (PA1-661; Thermo Fisher Scientific), and mouse anti-β-actin (3700; Cell Signaling). Following HRP-conjugated secondary antibodies were used: goat anti-rabbit (AS09 602; Agrisera) and goat anti-mouse (A9917; MERCK). Dynamin TKO cells were transfected with Lipofectamine 2000 (Thermo Fisher Scientific) using Opti-MEM I reduced serum medium (Thermo Fisher Scientific) for transient protein expression of Cav1-GFP according to manufacturer's instructions. For Dyngo-4a treatment, cells were serum-starved 1 hour prior to addition of 30 μM Dyngo-4a in live cell medium for 30 min prior to experiment or as indicated in Fig. 4 B and D-E. DMSO (0.1%) was used as control. For C-Laurdan experiments, Cav1-mCherry HeLa FlpIn TRex cells were incubated in 3μM C-Laurdan in imaging buffer (phenol-red free DMEM) containing catalase and glucose oxidase (7000 units/ml and 16 units/ml, respectively) at 37°C, 15 min prior to imaging. Fusogenic liposomes were prepared from a lipid mixture of DOPE, DOTAP, and Bodipy-tagged cholesterol at a ratio of 47.5:47.5:5. Lipid blends were in MeOH:CHCl$_3$ (1:3, vol/vol). Following the generation of a thin film using a stream of nitrogen gas, the vesicles were formed by addition of 20 mM Hepes (VWR, pH 7.5, final lipid concentration 2.8 μmol/ml) and incubated for 1.5 h at room temperature. Glass beads were

added to facilitate rehydration. The liposome dispersion was sonicated for 30 min (Transsonic T 310, Elma). Ctrl or Dyngo-4a treated cells were incubated with fusogenic liposomes (7nmol/ml) and incorporation of the fluorescent lipid in the PM was imaged using TIRF microscopy. For chol: MβCD -complex treatment, MβCD was dissolved in serum free media (phenol-red free DMEM) supplemented with HEPES (25mM) to give a concentration of 5 mM. Chol in $CHCl_3$ was added to a glass vial and a thin film of chol was created using a stream of nitrogen gas while rotating the vial on an angle. Thereafter, nitrogen gas was flowed into the vial for an hour to evaoporate remaining solvent. The chol was resuspended by vortexing in freshly made MβCD solution (5mM) for a final chol concentration of 480 mg/ml. The chol: MβCD solution was sonicated for 10 x 5 sec and incubated at 37°C ON with constant stirring. The chol:MβCD mixture was diluted 1:1 and cells were incubated with the solution and imaged using TIRF microscopy. For visualization of the incorporation, chol and Bodipy-chol at a ratio of 90:10 was complexed to MβCD and used as described above.

Immunostaining and transferrin uptake

Tamoxifen-treated and non-treated (ctrl) dynamin TKO cells were seeded on high precision coverslips (No. 1.5H, Paul Marienfeld GmbH and Co. KG) in 24-well plates at 3×10^4 cells/well and incubated overnight (37°C, 5% CO_2). Cells were fixed with 3% PFA in PBS (Electron Microscopy Sciences) for 15 min at room temperature (RT) and subsequent permeabilization and blocking was carried out simultaneously using PBS containing 5% goat serum and 0.05% saponin. Cells were then immunostained with rabbit anti-Caveolin1, rabbit anti-PTRF (RRID:AB_88224) both from Abcam, rabbit anti-EHD2, RRID:AB_2833022 (36), followed by goat anti-rabbit IgG secondary antibody coupled to Alexa Fluor 488, (Thermo Fisher Scientific) as previously described (37). Confocal images were acquired using a Zeiss Spinning Disk Confocal microscope controlled by ZEN interface (RRID:SCR_013672) with an Axio Observer.Z1 inverted microscope equipped with an EMCCD camera iXonUltra from ANDOR. For Transferrin uptake, Cav1-mCherry HeLa FlpIn TRex cells were seeded on precision coverslips in 24-well plates at 6×10^4 cells/well and induced with Dox and incubated at 37°C, 5% CO_2 16-18 hours prior to uptake assay. Cells were pretreated with Dyngo-4a or DMSO as specified in Fig. 4 D-E and then incubated 10 min at 37°C, 5% CO_2 with Alexa flour-647 conjugated transferrin (5 µg/ml). Cells were washed with ice-cold PBS and kept on ice for a total of 15 min and immediately fixed with 3% PFA for 15 min at RT. Images were aquired using a Leica DMI8 inverted Thunder widefield microscope equipped with a LEICA DFC 9000

GTC camera controlled by Leica Application Suite X 3.9.1.28433. Micrographs were prepared using Fiji, RRID:SCR_002285 (38) and Adobe Photoshop v25.6.

Live cell fluorescence microscopy imaging

Dynamin TKO cells were treated with 4-hydroxy-tamoxifen for five days in total and the day before experiments the cells were seeded on glass coverslips (CS-25R17, Warner instruments) in 6-well plates at 1.5×10^5 cells/well and incubated at 37°C, 5% CO_2. T-REx cells were induced with 0.5 ng/ml Dox and seeded on glass coverslips (CS-25R17) in 6-well plates at 3×10^5 cells/well and incubated at 37°C, 5% CO_2. Before imaging, the media was replaced with live-cell media (DMEM high glucose, no phenol red (Gibco), supplemented with 1 mM sodium pyruvate (Gibco) and imaged at 37°C with 5% CO_2. For TIRF-microscopy, images were acquired for 5 min at 3 s intervals (every third image with definitive focus) using a Zeiss Axio Observer.Z1 inverted microscope that was equipped with an EMCCD camera iXon Ultra from ANDOR and an alpha Plan-Apochromat TIRF 100×/1.46 oil objective controlled by ZEN software. Only the basal membrane was analyzed as the laser penetration depth can not be adjusted to image the apical and lateral plasma membrane. All micrographs and videos were prepared with Fiji (38) and Adobe Photoshop CS6. C-Laurdan imaging was performed on a wide-field fluorescence Nikon Ti-U microscope (Nikon, Tokyo, Japan) equipped with a Cascade 1 k camera and a dual viewer (Photometrics, Tucson, AZ). A 60X water objective (NA 1.27) and refractive index-matched immersion oil (Zeiss, Oberkochen, Germany) were used. Excitation was with a LED at 380 nm (CoolLED, Andover, UK). A FF01-387/485/559/649 excitation filter (Semrock, Lake Forrest, IL) and a z365/577rpc dichroic (Chroma, Rockingham, VT) were used in the microscope and in the dual viewer a ms-470LDX dichroic (Chroma) and emissions filters 442/46 and FF01–483/32 from Semrock were used. Stacks of eleven images with a z-step size of 200 nm were acquired close to the basal membrane of the cells.

Analysis of caveola dynamics

Imaris software 10.0.1 (Bitplane) was used for tracking analysis of Cav1-GFP or Cav1-mCh positive structures segmented as spots as previously described (10). Structures with a diameter of 0.4 μm were tracked and the applied algorithm was based on Brownian motion with max distance traveled of 0.8 μm and a max gap size of 3. Statistics for track duration and track displacement length from three independent experiments were extracted and analyzed. Tracks in figures were displayed using the spectrum colormap for track duration showing the caveolae

heterogeneity within a cell where red tracks represent stable caveolae and blue/purple highly dynamic caveolae. The number of quantified cells is specified in the respective figure legend. Unpaired two-sample t test was performed on track duration (s) and track displacement (μm) using Prism 10.0.2, and data are shown as fold change.

Analysis of PM dynamics

Masks of the total area of the basal membranes of the first and last image of 5 minute TIRF movies were established based on the membrane epiflouroscense using ImageJ. To quantify the PM motion, the area change was calculated by subtracting the end mask from the start mask (1) and by subtracting the start mask from the end mask (2) using the Image J image calculator. The total area change was calculated by subtracting the result from (1) and (2) from the area of the start image. The number of quantified cell areas are specified in respective figure legend. Unpaired two-sample t test was performed and data are shown as fold change.

Image deconvolution and GP analysis of C-Laurdan images

The images were deconvolved to improve the resolution (39). The image stacks were first cropped around single cells. Image deconvolution was performed using Huygens Professional (SVI, Netherlands) with default settings and automatic background detection. A theoretical point spread function (PSF) was used and the maximal number of iterations was set to 50. The upper and lower channels (CH483 and CH442)) were deconvolved concurrently with a signal to noise ratio of 1. The concurrently deconvolved images were saved using a linked scale intended to retain their relative intensities. The regions of interest (ROIs) were created using the ImageJ macro Find_plasma_membrane (40), where the center images from the deconvolved z-stacks from the two channels were aligned using the ImageJ plugin *StackReg* (41). Only plasma membrane stretches without adjacent stained intracellular membranes were selected for the analysis. First, a segmented line ROI was created by manually selecting a few points close to the plasma membrane section to be analyzed. Then, the pixels with the highest intensity along lines perpendicular to each selected point were identified, additional points interpolated between them and their positions optimized in the same manner. The background was subtracted based on the mean intensities in an area outside the cells. Generalized polarization (GP) values were calculated for all the pixels in the plasma membrane ROIs according to the formula:

$$GP = \frac{I(419-465)-I(470-499)}{I(419-465)+I(470-499)} \quad (1)$$

Generation of SLBs and analysis by QCM-D

POPC vesicles for forming SLBs were prepared by diluting POPC in chloroform:methanol (3:1 volume/volume), drying it under a stream of nitrogen and rehydrating in 20 mM HEPES buffer, 150 mM NaCl (pH 7.4) (HBS buffer) at a final concentration of 1 mg/ml. Liposomes were sonicated using a bath sonicator and then extruded 11 times (Mini Extruder, Avanti) through a polycarbonate filter (Nuclepore Track-Etched Membranes, Whatman) with 100-nm pore size. An X4 unit (AWSensors) equipped with four flow chambers was used to conduct the QCM-D measurements. Wrapped 14-mm (5 MHz, Cr/Au–SiO$_2$, polished) sensors were used for all experiments. Each sensor was stored in 2% SDS overnight and treated with UV/ozone (BioForce Nanosciences) for 30 min before use. The frequency and dissipation changes for overtones 1, 3, 5, 7, 9, and 11 were all recorded, but only the third overtone is reported herein. POPC vesicles (100 µL, 0.1 mg/mL) in HBS buffer were injected in a continuous flow and SLB formation was monitored. After SLB formation, the chambers were rinsed with HBS buffer. After reaching a stable baseline, DMSO (0.2%) was injected into the chamber. The flow was paused once the solution had filled the sensor chamber, and the system was allowed to reach equilibrium before 30 µM Dyngo-4a in DMSO (0.2%) or DMSO (0.2%) alone as control was injected into the chamber. The flow was paused again once the solution had filled the sensor chamber, and the system was allowed to reach equilibrium before rinsing with the buffer.

Estimation of Dyngo-4a retatined in SLBs

The adsorbed mass of Dyngo-4a was calculated from the frequency shift using the Sauerbrey Equation (42) using the AWSensors software (AWS suite). The system followed the necessary criteria for valid calculations: 1) thin adsorbed layer relative to the sensor, 2) homogenous adsorbed layer and 3) ridged adsorbed layer (43). The POPC SLBs were very thin (~4 nm) and homogenous displaying 2D fluidity. The SLB rigidity, before DMSO addition and after DMSO removal, was established to $\Delta D/\Delta F < 6.6 \times 10^{-8}$ Hz^{-1} as measured by the AWSensors using the third overtone of a 5MHz sensor (Fig. S2). The clean sensor surface before SLB formation was used as the zero reference. The mole percentage of Dyngo-4a retained in the SLB after rinsing was calculated by using the molecular weights of POPC (759.578 g/mol) and Dyngo-4a (338.3 g/mol).

Lipid Monolayer Experiments

Lipid monolayer experiments were performed with a Microtrough G1 System (Ø 53 × 4 mm, Kibron). The trough was covered to prevent temperature and humidity loss. POPC lipid was prepared at a total lipid concentration of 1 mM in chloroform:methanol (3:1 vol/vol). A microbalance equipped with a Wilhelmy plate was used to measure the surface pressure (π) and calibrated before each measurement. Lipid solutions were deposited onto the surface of the subphase (20 mM HEPES, 150 mM NaCl, pH 7.4) to obtain the initial surface pressure (π_0). The subphase was continuously stirred by a magnetic stirrer. After the solvent had been allowed to evaporate for 15 min and a stable monolayer was formed, Dyngo-4a in DMSO or DMSO was injected under the lipid film directly into the subphase using a thin syringe needle (final concentration 30 µM Dyngo-4a, 0.1% DMSO). Experiments were run in triplicates and analysis of the adsorption curves was performed with GraphPad Prism.

MD simulations

The CHARMM force field (44-46) was used to run MD simulations. The Dyngo-4a molecule was built using ligand modeler (47) and validated by octanol-water partitioning compared to cLogP value provided by the supplier (2.5 against 2.6). All simulation series were run in triplicates and presented as average and standard deviation. Initial equilibration simulations were conducted using the Nose-Hoover thermostat (48) ($\tau_T = 1, T_{ref} = 300$) and semi-isotropic Parrinello-Rahman (49) barostat ($\tau_P = 5, P_{ref} = 1\ atm, compressibility = 4.5e - 5\ bar^{-1}$) were employed to control temperature and pressure in the simulations. For AWH simulations, a convolved adaptive biasing potential (50) was applied with a force constant of 12800kJ mol^{-1} nm^{-2} defining the curvature of the biasing potential along the reaction coordinate. Verlet cut-off (51) scheme was applied with a cut-off radius of 1.2 nm set to both van der Waals and electrostatic interactions. LINCS algorithm (52) was used to constraint all bonds with hydrogen atoms.

AFM experiments

HeLa cells were seeded on a 35 mm glass bottom Petri dish (Cellvis, Cat. # D35-10-1.5-N) and starved as described above. The cell media was exchanged to serum-free media with 20 mM HEPES containing either Dyngo (30 µM) or DMSO (0.1%) and set for 30 minutes before performing AFM experiments. All AFM experiments were conducted by NanoWizard 4 XP bioscience system (Bruker/JPK, Berlin, Germany) at ambient temperature. Commercial colloid probes (Si$_3$N$_4$ AFM probes, 0.03 N/m cantilever; 4.5 µm polystyrene particle from Novascan

Technologies, Ames, IA USA) were used to collect force maps on Cav1 FlpIn T-REx HeLa cells, with and without the treatment of Dyngo after calibrating the cantilever. The spring constant for each cantilever was calculated based on the thermal noise method. A glass dish with cells was mounted on an inverted Nikon Eclipse Ti-E2 microscope (Nikon Corp., Japan) with a Prime 95B sCMOS camera (Photometrics Teledyne, USA) to localize cells using a 40x objective. The cantilever was carefully centred on cells. The whole duration of experiments was kept below 2 hours. For both conditions, force maps on 8 or 9 individual cells were collected. An approach curve with a length of 5 µm at a speed of 2 µm/s and a retract curve with a length of 7 µm and a speed of 4 µm/s at the maximum force load of 1 nN was used for mapping 8x8 matrix (or 64 points) over 10 µm² scan size on individual cells and glass (i.e., a region of a glass dish with no cells). All experiments were carried out independently at least twice, with separately but identically prepared cells and AFM tips.

AFM data analysis

Individual force curves from force maps were analyzed using the JPK data processing software. Before fitting the force curves, data was smoothed with a Gaussian filter with a width of 1. After smoothing and setting the measured force close to the contact point and offset to zero of the extend curve, the Young's modulus (*E*) was determined by fitting with the built-in Hertz/Sneddon model (Eq. 1). The major assumption of this model is that the indentation depth (~0.5 to 2 µm) is very small or nearly negligible as compared to the thickness of the cells (10 to 12 µm). This model also assumes that the sample is elastic and homogenous. As this is not the case for cells, we fitted the data from 0 to 500 nm of indentation (*δ*) from a contact point for both conditions.

$$F = \frac{4}{3}\frac{E}{1-\mu^2}\sqrt{R\delta^3} \qquad\qquad (1)$$

where *F* is force, R represents the radius of the spherical probe (2.25 µm) and μ is the Poisson ratio. The ratio varies from 0 to 0.5 depending on the material properties. For simplicity, we set it to 0.5. Curves with the root mean square (RMS) value below 15 pN were discarded from further analysis.

Statistical analysis

Two-sample *t* test was used to compare values in the different experiments using the GraphPad prism program version 10.0.2. P < 0.05 was considered a statistically significant change. *P <

0.05; **P < 0.01; ***P < 0.001; NS, not significant. All the values were presented as mean ± standard deviation (SD) as specified in the figure legends.

Data sharing plans

All data needed to evaluate the conclusions in this article are included in the article and/or SI appendix.

Online supplementary material

Fig. S1 show the molecular structures of Dyngo-4a and cholesterol, immunofluorescent stainings of cavin1 and EHD2 in Dynamin TKO cells and an illustration of caveola mobility and TIRF imaging. Fig. S2 show the softness plot from QCM-D measurements during SLB formation. Fig. S3 provides additional MD simulation data of ten Dyngo-4A molecules in POPC bilayer or in POPC:Chol bilayer. Fig. S4 shows the analysis of the effect of Dyngo-4a on PM mobility, caveolae displacement and Tfn uptake as well as AFM force-indentation curves, and incorporation of Chol using fusogenic liposomes and MβCD. Video 1 show TIRF microscopy timelapse of a Dynamin TKO cell treated with DMSO (0.1%) for 30 min. Video 2 show TIRF microscopy timelapse of a Dynamin TKO cell treated with Dyngo-4a (30μM) for 30 min.

Figure legends

Figure 1. The drug Dyngo-4a constrains both lateral mobility and scission of caveolae independently of dynamin. (A) Immunoblots of dynamin triple knock out cells showing dynamin 2 expression after 5 days of Tamoxifen-treatment (KO) or no treatment (Ctrl) as indicated. **(B)** Representative immunofluorescent staining of Caveolin1 in dyn TKO cells treated as in (A) as indicated, Scale bar, 5 μm. **(C)** Quantification of the number of caveolae per square micrometer counted in the basal PM in immunofluorescent labeled dyn TKO cells treated as in (A) as indicated. Mean ± SD from at least 20 cells per condition. Significance was assessed using *t* test, p =0.7421. **(D)** Color-coded trajectories showing caveolae movements in the PM. Cav1-GFP was transiently expressed in dyn TKO cells after 5 days of Tamoxifen treatment. Cells were serum-starved 1 h prior to 30 min treatment with DMSO (Ctrl) or 30 μM Dyngo-4a before imaged on TIRF over 5 min. Red trajectories represent caveolae with long duration times and purple represents caveolae with short duration times as indicated. Scale bar, 10 μm. **(E-F)** Quantification of Cav1-GFP track duration time (E) and track displacement

length (F). Numbers were related to ctrl-treated cells, and track mean from at least 23 cells per condition are shown ± SD (E) and track length mean from at least 15 cells per condition are shown ± SD (F). Significance was assessed using t test, ****p ≤ 0.0001. **(G)** Kymographic visualization of the track duration of color-coded trajectories in (D) that were present at time zero, representing axial stability. **(H)** Visualization of the track displacement of color-coded trajectories in (D) by alignment of the starting position of all tracks, representing lateral mobility.

Figure 2. Dyngo-4a binds and inserts into membranes. (A) Top, Illustration of Langmuir-Blodget experiment. Bottom, Dyngo-4a adsorption to POPC monolayer. Dyngo-4a (30 μM) (black line) or DMSO (gray line) was injected underneath the lipid film and the surface pressure shift (Δπ) was recorded over time. **(B)** Top, illustration of the QCM-D set up. Bottom, QCM-D measurements of SLB formation and Dyngo-4a (30 μM) (black line) or DMSO (gray line) adsorption to a POPC SLB, followed by rinsing. Arrows depict time of injection of DMSO (equivalent to the concentration that Dyngo-4a is dissolved in), sample injection (Dyngo-4a or DMSO), DMSO injection and saline buffer wash (150 mM NaCl, pH 7.4) injection respectively.

Figure 3. Dyngo-4a inserts underneath the phospholipid headgroups at a similar position as cholesterol. (A) Molecular dynamic simulation of Dyngo-4a in a membrane. Snapshot of a single Dyngo-4A molecule (cyan stick and red volume) in POPC bilayer (phospholipid headgroups in dark gray and lipid chains as light gray sticks, dark blue represents the aqueous phase) **(B-C)** AWH simulations of Dyngo-4a (B) or Chol (C) in POPC membrane. PMF profile show the energy required to move a Dyngo-4a molecule or Chol molecule from the membrane center to the aqueous phase, respectively. **(D)** Molecular dynamic simulation of Dyngo-4a in a membrane. Snapshot of a single Dyngo-4A molecule (cyan stick and red volume) in POPC:Chol (70:30%) bilayer (phospholipid headgroups in dark gray and lipid chains as light gray sticks, chol as orange sticks, dark blue represents water). **(E)** AWH simulations of Dyngo-4a in POPC:Chol (70:30%) membrane. The PMF profile shows the energy required to move a Dyngo-4a molecule from the membrane center to the aqueous phase. In the PMF profiles the lines represent the mean and the shaded regions represent the SD from triplicate simulations.

Figure 4. Dyngo-4a decreases the lipid packing in the outer leaflet of the PM which is counteracted by increased levels of cholesterol. (A) Visualization of the PM mobility of a ctrl-treated cell (top) and a Dyngo-4a-treated cell (bottom) from 5 minute TIRF movies. The outline of the PM at the start (0 s, red), middle (60, 120, 180 and 240 s, gray) and end (300 s, black) are depicted. Insets to the right show magnifications of the indicated areas. Scale bar, 10 μm. **(B)** Quantification of the PM mobility of cells following treatment with DMSO (ctrl) or Dyngo-4a at different concentrations as indicated. The means is shown from five cells per condition ± SD. **(C)** Representative electron micrographs of caveolae in DMSO treated (ctrl) and Dyngo-4a treated cells. Scale bar, 100 nm. **(D)** Young moduli calculated from force-indentation curves from the representative curves in (Fig S4) over 31 and 33 cells treated with DMSO and Dyngo-4a, respectively. Each data point is an average value calculated from at least 40 individual force curves on each cell. Statistical analysis: parametric student t-test with Welch's correction. ns = non significant. **(E)** Representative deconvolved fluorescent micrographs of HeLa cells stained with C-Laurdan captured at 442 nm (left panel) and 483 nm (middle panel). Yellow arrows depict analyzed stretch of PM with no neighboring fluorescent internal membrane as seen in the right panel. Red area corresponds to membrane with high membrane order and purple corresponds to membrane with low membrane order. Scale bar, 5 μm. **(F)** Quantification of the mean GP-values for DMSO (ctrl) or Dyngo-4a treated cells as indicated. All data points from at least 26 cells per condition are shown, n = 3, mean ± SD. Significance was assessed using t test, ** $p \leq 0.0065$. **(G)** Quantification of Cav1-mCh track duration time following chol:MβCD treatment as indicated. Numbers were related to ctrl-treated cells. All data points from at least 26 cells per condition are shown, n = 3, mean ± SD. Significance was assessed using t test, ****$p \leq 0.0001$.

References

1.   L. V. Chernomordik, M. M. Kozlov, Protein-lipid interplay in fusion and fission of biological membranes. *Annu Rev Biochem* **72**, 175-207 (2003).
2.   J. H. Lorent *et al.*, Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nat Chem Biol* **16**, 644-652 (2020).
3.   M. Hubert, E. Larsson, R. Lundmark, Keeping in touch with the membrane; protein- and lipid-mediated confinement of caveolae to the cell surface. *Biochem Soc Trans* **48**, 155-163 (2020).

4. T. Laurent *et al.*, Architecture of the chikungunya virus replication organelle. *Elife* **11** (2022).
5. R. G. Parton, Caveolae: Structure, Function, and Relationship to Disease. *Annu Rev Cell Dev Biol* **34**, 111-136 (2018).
6. A. Echarri, M. A. Del Pozo, Caveolae. *Curr Biol* **22**, R114-116 (2012).
7. A. K. Kenworthy, B. Han, N. Ariotti, R. G. Parton, The Role of Membrane Lipids in the Formation and Function of Caveolae. *Cold Spring Harb Perspect Biol* **15** (2023).
8. J. C. Porta *et al.*, Molecular architecture of the human caveolin-1 complex. *Sci Adv* **8**, eabn7232 (2022).
9. M. Doktorova *et al.*, Caveolin assemblies displace one bilayer leaflet to organize and bend membranes. *bioRxiv* 10.1101/2024.08.28.610209 (2024).
10. M. Hubert *et al.*, Lipid accumulation controls the balance between surface connection and scission of caveolae. *Elife* **9** (2020).
11. E. Larsson, B. Moren, K. A. McMahon, R. G. Parton, R. Lundmark, Dynamin2 functions as an accessory protein to reduce the rate of caveola internalization. *J Cell Biol* **222** (2023).
12. M. Kaksonen, A. Roux, Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **19**, 313-326 (2018).
13. B. Sinha *et al.*, Cells respond to mechanical stress by rapid disassembly of caveolae. *Cell* **144**, 402-413 (2011).
14. E. Macia *et al.*, Dynasore, a cell-permeable inhibitor of dynamin. *Dev Cell* **10**, 839-850 (2006).
15. A. McCluskey *et al.*, Building a better dynasore: the dyngo compounds potently inhibit dynamin and endocytosis. *Traffic* **14**, 1272-1289 (2013).
16. M. T. Howes *et al.*, Clathrin-independent carriers form a high capacity endocytic sorting system at the leading edge of migrating cells. *J Cell Biol* **190**, 675-691 (2010).
17. E. Girard *et al.*, The dynamin chemical inhibitor dynasore impairs cholesterol trafficking and sterol-sensitive genes transcription in human HeLa cells and macrophages. *PLoS One* **6**, e29042 (2011).
18. H. Yamada *et al.*, Dynasore, a dynamin inhibitor, suppresses lamellipodia formation and cancer cell invasion by destabilizing actin filaments. *Biochem Biophys Res Commun* **390**, 1142-1148 (2009).
19. G. Preta, V. Lotti, J. G. Cronin, I. M. Sheldon, Protective role of the dynamin inhibitor Dynasore against the cholesterol-dependent cytolysin of Trueperella pyogenes. *FASEB J* **29**, 1516-1528 (2015).
20. R. J. Park *et al.*, Dynamin triple knockout cells reveal off target effects of commonly used dynamin inhibitors. *J Cell Sci* **126**, 5305-5312 (2013).
21. C. Matthaeus *et al.*, The molecular organization of differentially curved caveolae indicates bendable structural units at the plasma membrane. *Nat Commun* **13**, 7234 (2022).
22. M. Hoernke *et al.*, EHD2 restrains dynamics of caveolae by an ATP-dependent, membrane-bound, open conformation. *Proc Natl Acad Sci U S A* **114**, E4360-E4369 (2017).
23. J. Mohan, B. Moren, E. Larsson, M. R. Holst, R. Lundmark, Cavin3 interacts with cavin1 and caveolin1 to increase surface dynamics of caveolae. *J Cell Sci* **128**, 979-991 (2015).

24. A. Blume, A. Kerth, Peptide and protein binding to lipid monolayers studied by FT-IRRA spectroscopy. *Biochim Biophys Acta* **1828**, 2294-2305 (2013).

25. V. I. Gordeliy, M. A. Kiselev, P. Lesieur, A. V. Pole, J. Teixeira, Lipid membrane structure and interactions in dimethyl sulfoxide/water mixtures. *Biophys J* **75**, 2343-2351 (1998).

26. N. J. Cho, C. W. Frank, B. Kasemo, F. Hook, Quartz crystal microbalance with dissipation monitoring of supported lipid bilayers on various substrates. *Nat Protoc* **5**, 1096-1106 (2010).

27. H. M. Kim *et al.*, A two-photon fluorescent probe for lipid raft imaging: C-laurdan. *Chembiochem* **8**, 553-559 (2007).

28. O. G. Mouritsen, M. J. Zuckermann, What's so special about cholesterol? *Lipids* **39**, 1101-1113 (2004).

29. K. Simons, E. Ikonen, How cells handle cholesterol. *Science* **290**, 1721-1726 (2000).

30. C. M. MacDermaid *et al.*, Molecular dynamics simulations of cholesterol-rich membranes using a coarse-grained force field for cyclic alkanes. *J Chem Phys* **143**, 243144 (2015).

31. C. Yuan, L. J. Johnston, Phase evolution in cholesterol/DPPC monolayers: atomic force microscopy and near field scanning optical microscopy studies. *J Microsc* **205**, 136-146 (2002).

32. S. Mahammad, J. Dinic, J. Adler, I. Parmryd, Limited cholesterol depletion causes aggregation of plasma membrane lipid rafts inducing T cell activation. *Biochim Biophys Acta* **1801**, 625-634 (2010).

33. R. Lundmark, E. Larsson, L. I. A. Pulkkinen, The adaptable caveola coat generates a plasma membrane sensory system. *Curr Opin Cell Biol* **88**, 102371 (2024).

34. A. Barnoy, R. G. Parton, M. M. Kozlov, Mechanism of membrane curvature generation and caveola formation by flat disc-like complexes of caveolin. *bioRxiv* 10.1101/2024.07.09.602763, 2024.2007.2009.602763 (2024).

35. D. Dutta, C. D. Williamson, N. B. Cole, J. G. Donaldson, Pitstop 2 is a potent inhibitor of clathrin-independent endocytosis. *PLoS One* **7**, e45799 (2012).

36. B. Moren *et al.*, EHD2 regulates caveolar dynamics via ATP-driven targeting and oligomerization. *Mol Biol Cell* **23**, 1316-1329 (2012).

37. R. Lundmark *et al.*, The GTPase-activating protein GRAF1 regulates the CLIC/GEEC endocytic pathway. *Curr Biol* **18**, 1802-1808 (2008).

38. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).

39. A. Huang, J. Adler, I. Parmryd, Optimised generalized polarisation analysis of C-laurdan reveals clear order differences between T cell membrane compartments. *Biochim Biophys Acta Biomembr* **1865**, 184094 (2023).

40. J. Adler, A. Huang, I. Parmryd, Find_plasma_membrane and measure_plasma_membrane: ImageJ macros for efficient identification of and measurements at and around the plasma membrane. *SoftwareX* **24**, 101570 (2023).

41. P. Thevenaz, U. E. Ruttimann, M. Unser, A pyramid approach to subpixel registration based on intensity. *IEEE Trans Image Process* **7**, 27-41 (1998).

42. G. Sauerbrey, Verwendung von Schwingquarzen zur Wägung dünner Schichten und zur Mikrowägung. *Zeitschrift für Physik* **155**, 206-222 (1959).

43.   I. Reviakine, D. Johannsmann, R. P. Richter, Hearing what you cannot see and visualizing what you hear: interpreting quartz crystal microbalance data from solvated interfaces. *Anal Chem* **83**, 8838-8848 (2011).

44.   S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: a web-based graphical user interface for CHARMM. *J Comput Chem* **29**, 1859-1865 (2008).

45.   J. Lee *et al.*, CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J Chem Theory Comput* **12**, 405-413 (2016).

46.   E. L. Wu *et al.*, CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J Comput Chem* **35**, 1997-2004 (2014).

47.   H. Guterres, S. J. Park, Y. Cao, W. Im, CHARMM-GUI Ligand Designer for Template-Based Virtual Ligand Design in a Binding Site. *J Chem Inf Model* **61**, 5336-5342 (2021).

48.   D. J. Evans, B. L. Holian, The Nose-Hoover Thermostat. *J Chem Phys* **83**, 4069-4074 (1985).

49.   M. Parrinello, A. Rahman, Polymorphic Transitions in Single-Crystals - a New Molecular-Dynamics Method. *J Appl Phys* **52**, 7182-7190 (1981).

50.   J. Lidmar, Improving the efficiency of extended ensemble simulations: The accelerated weight histogram method. *Phys Rev E* **85** (2012).

51.   S. Páll, B. Hess, A flexible algorithm for calculating pair interactions on SIMD architectures. *Comput Phys Commun* **184**, 2641-2650 (2013).

52.   B. Hess, H. Bekker, H. J. C. Berendsen, J. G. E. M. Fraaije, LINCS: A linear constraint solver for molecular simulations. *Journal of Computational Chemistry* **18**, 1463-1472 (1997).

Figure 1

A

130-
95-   a-Dyn2

55-
       a-actin

B Ctrl / Cav1   TKO / Cav1

C
Caveola/µm²
ns
Ctrl   TKO

D Ctrl   Dyngo-4a

9   Track duration time (s)   300

E
Relative track duration time
****
Ctrl   Dyngo-4a

F
Relative track displacement length
****
Ctrl   Dyngo-4a

G Axial stability
Ctrl   9 / 300 Time
Dyngo-4a   9 300 / 300 Time

H Lateral mobility
Ctrl   Dyngo-4a
9   Track time (s)   300

Figure 2

A

Lipid monolayer →
Sample injection →

Air
Aqueous

Injection point

$\Delta\pi$ (mN m^{-1})

-1

30   60   90   120

Time (min)

— Dyngo
— Ctrl

B

Sample injection →
SLB →
Sensor surface →

Aqueous

ΔF (Hz)

2
0
-2
-4

DMSO   Dyngo-4a   DMSO   Saline
                              Buffer

0   1000   2000   3000

Time (s)

— Dyngo
— Ctrl

Figure 3

A 1 Dyngo-4a in POPC

Aqueous

Membrane bilayer

Aqueous

B Single Dyngo-4a entering POPC

C Single Chol entering POPC

D 1 Dyngo-4a in POPC:Chol

Aqueous

Membrane bilayer

Aqueous

E Single Dyngo-4a entering POPC:Chol

Figure 4

A Cell outline over time: start, middle, end

Ctrl

Dyngo-4a

B PM mobility

Relative area difference (start and end) vs Dyngo-4a conc (μM)

C

Ctrl

Dyngo-4a

D Elasticity

ns

Young's modulus (kPa) — Ctrl, Dyngo-4a

E

442 nm   483 nm   GP value

Ctrl

Dyngo-4a

0.206

GP value

-0.19

F Lipid packing

**

GP value — Ctrl, Dyngo-4a

G Caveola duration

ns
****
****
***

Relative track duration time

Ctrl   Dyngo-4a   +Chol   +MOCK