The conserved nature of nitrogen recycling within the bacterial pouch of *Tetraponera* ants

Mingjie Ma[1], Biru Zhu[1], Dayong Zhang[1], Piotr Łukasik[2], Yi Hu[1]

1. Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing 100875, China
2. Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Kraków, Poland

Abstract

Microbial symbioses are fundamental to the nutrition of many animal groups, yet the mechanisms of nutrient recycling have only been characterized in a few social insect systems. Here, we investigate the functional significance of a specialized and unusual symbiotic organ—the bacterial pouch—in four ant species within the *Tetraponera nigra*-group, focusing on the symbionts' roles in nitrogen recycling and colony fitness. This unique organ houses a microbial community consistently dominated by the co-diversifying bacterial symbiont *Tokpelaia*, occasionally accompanied by other stable bacterial associates. Metagenomic sequencing and targeted ^{15}N-labeled urea feeding experiments demonstrate that these symbionts assimilate nitrogen from urea (likely derived from uric acid) and transfer it to adult workers and developing larvae, incorporating recycled nitrogen into host tissues. Disruption of this symbiosis severely impairs larval growth and overall colony fitness. Overall, our study highlights the critical role of the bacterial pouch in sustaining colony fitness in nitrogen-limited environments, providing new insights into the co-evolutionary dynamics between insect hosts and their microbial partners.

Keywords:

29   microbiome, microbiota, bacteriome, amino acid biosynthesis, Formicidae

Introduction

The symbiotic relationships have significantly influenced ecological dynamics and

evolutionary trajectories of interacting multicellular organisms and their microbiota.

Notably, the associations formed between specific microbial partners and dedicated

host organs play a critical role in shaping these interactions. Classic examples include

plant root nodules, which harbor nitrogen-fixing rhizobia[1, 2], and the light organs of

bobtail squid[3, 4], home to bioluminescent bacteria. Similarly, mycangia in ambrosia

beetles[5] serve as reservoirs for fungal symbionts, while specialized gut organs in bean

bugs[6] and bacteriomes in many Hemiptera insects[7, 8] provide a habitat for nutrient-

provisioning bacteria. Despite the importance of symbiont-hosting organs, our

understanding of their diversity and distribution across various phylogenetic lineages

and the specificity and ecological significance of host-microbe associations remains

notably incomplete.

Ants (Hymenoptera: Formicidae) are among the most ecologically successful and

significant groups of insects on Earth. Their mutualistic interactions with microbes

have been identified as an important factor driving their long-term evolution and

ecological dominance[9, 10, 11, 12]. Recent studies have shown that several ant groups

harbor diverse microbial communities that influence multiple life history traits of their

ant hosts[10, 13]. In particular, microbes' abilities to provide essential nutrients such as

amino acids, vitamins, and other metabolic byproducts that ants cannot synthesize

themselves are considered a key driver that enables ants to thrive in various nutrient-

imbalanced niches, contributing to their widespread ecological dominance[11, 14, 15, 16, 17].

A remarkable pattern in ant evolution is multiple shifts from nitrogen-rich

carnivory and omnivory to nitrogen-poor herbivory. Herbivorous ants typically

scavenge for foods such as honeydew from tended sap-feeding insects, extrafloral

nectar, pollen, fungi, and plant wound secretions[18, 19, 20]. The availability of accessible

nitrogen and essential amino acids in these food sources is typically limited. The

relatively high occurrence of specialized bacteria in herbivorous ant taxa indicates that they may often rely on microbial partners to mitigate nitrogen limitation and nutritional imbalance[10, 11, 12, 13]. Recent studies using high-throughput sequencing and functional assays have shown a trend of functional convergence in nitrogen recycling capabilities among symbionts of unrelated herbivorous ants that feed on N-poor or N-inaccessible foods. Such nitrogen provisioning by symbionts has been demonstrated in carpenter ants (genus *Camponotus*), whose intracellular symbiont *Blochmannia* synthesizes essential amino acids from recycled nitrogen for their hosts[21, 22]. In the herbivorous ant tribe Cephalotini, analogous roles fulfilled by abundant, conserved multi-species bacterial communities present within the gut lumen were both predicted from metagenomic data and demonstrated *in vivo* and *in vitro*[17, 23]. In the herbivorous genus *Dolichoderus*, the same functions were predicted based on metagenomic data but not demonstrated experimentally[16].

Due to the intricate social structure of ant colonies, the nutritional demands of their colony members differ significantly. Larvae require substantial protein intake for their rapid growth, and adult workers, with their higher carbohydrate needs for energy[24], have distinct dietary needs. This dietary divergence proposes a greater challenge for ant larvae in obtaining sufficient nitrogen than adult workers. The critical role of symbiotic bacteria in the overall nutrition of ant colonies has been well studied in *Camponotus* ants, which harbor intracellular symbionts *Blochmannia* at all developmental stages. It was shown that antibiotic-treated colonies suffered from a decline in the number of larvae developing into pupae, smaller body size of newly eclosed workers, and reduced overall colony sizes[25], highlighting the essential role that *Blochmannia* plays in supporting colony development through its impact on nitrogen balance and brood nutrition. In contrast, extracellular symbionts associated with ants are typically restricted to specific life stages. Their role in nutrition supplementation across the entire colony and their broader impact on colony-wide nutrition and development remains an area that requires further investigation.

The arboreal ant genus *Tetraponera,* restricted to the Palaeotropics and Australia, encompasses approximately 110 described species divided into six distinct species

groups[26, 27] that were shown to vary in their nutritional biology and anatomy[28, 29, 30]. The

T. nigra-group, comprising 20 species, is uniquely characterized by an unusual bacterial

pouch located at the junction between the midgut and hindgut. This pouch is connected

to Malpighian tubules and the tracheal system and is densely populated with bacteria[31].

This structure bears a resemblance to the symbiotic organs observed in other insects

such as leaf beetles[32], bean bugs[33], or ambrosia beetles[5], but also with root nodules of

legumes[1], suggesting that these ants engage in a specialized form of nutritional

symbiosis with microbial partners[34]. Stable isotopic analysis revealed that *T. nigra*-

group ants are at a relatively low trophic level (Fig. 1B). Indeed, three species in the

group, *T. tucurua*[35], *T. binghami*[30, 36], and *T. punctulata*[35], have been documented

feeding on honeydew produced by scale insects, and one species *T. rotula*[37] feeding on

extrafloral nectar – foods considered nutritionally imbalanced. We hypothesized that

the nutritional symbiosis with specialized microorganisms housed within gut pouches

may have enabled these ants to occupy ecological niches where essential nutrients are

otherwise scarce, providing a potential explanation for the success of this group in such

habitats. Hence, in a broad sample spanning multiple colonies of several species, we

anticipated finding evidence for the symbiosis stability, the conservation and

expression of symbiont functions related to nitrogen recycling and nutrient biosynthesis,

and a correlation between symbiosis disruption and colony health.

The goal of this study was to comprehensively characterize the diversity and

functional roles of bacterial symbionts and their significance in the nutritional biology

of the *T. nigra*-group ants. We first asked about the general organization of the gut

structure in connection with the bacterial pouch across ant life stages. Second, we

characterized the diversity of microbial communities within bacterial pouches across

colonies of four species and the phylogenetic position and genomic characteristics of

the dominant bacteria. Third, we assessed the roles of these symbiotic bacteria in

nutrient metabolism, with an emphasis on nitrogen recycling and provisioning. Lastly,

we explored how they contribute to nutritional supplementation for the entire colony,

particularly for larvae. We addressed these questions by combining microscopic

analysis, 16S rRNA gene amplicon sequencing, metagenomics, and *in vivo* colony
feeding experiments. Through this multi-pronged investigation, we demonstrated how
specialized symbiotic relationships underpin the nutritional ecology of these
herbivorous ants.

Results

T. nigra-group ant workers have evolved an unusual organ – bacterial pouch

Previous studies have revealed the presence of a pouch-like structure in the gut of
adult workers from three *T. nigra*-group ant species[31]. Our microscopy analyses of the
four ant species from the *T. nigra*-group, including two previously characterized[31] (*T.*
attenuata, *T. binghami)* and two newly identified and so far undescribed (*T.* sp1 and *T.*
sp2 - Fig. 1A and Supplementary Fig. 1), confirmed that this specialized gut structure
is consistently present within the group. In adult workers of four ant species, this pouch-
like structure is located at the junction between the midgut and hindgut and is connected
to several Malpighian tubules (Fig. 1C, I-L). The inner lining of the pouch lumen
consists of microvillar cells containing numerous mitochondria, and a dense population
of rod-shaped bacteria enclosed by a double membrane are housed inside the pouch
(Fig. 1F and G). The whole-mount fluorescence in situ hybridization (FISH) reveals
that bacteria are concentrated within the pouch-like structure in all four ant species (Fig.
1I-L), confirming its role as a symbiotic organ, referred to as the bacterial pouch in the
T. nigra-group ants.

When comparing across life stages of *T. nigra*-group ants, scanning electron
microscopy observations revealed that larvae lack the specialized bacterial pouch (Fig.
1E), which only develops during the pupal stage (Fig. 1D). The symbiotic organ in the
pupal stage is underdeveloped compared to that of adults: smaller in size and not yet
fully swollen. Furthermore, the results from the transmission electron microscopy

showed a notable absence of bacteria in the symbiotic organ during the pupal stage (Fig. 1H). This indicates that the establishment of symbiotic relationships occurs after worker eclosion.

Fig. 1 Origins of specimens and gut structure of the _T. nigra_-group ants. **(a)** Sampling locales for four ant species used in this study (image credit: Mingjie Ma). **(b)** Nitrogen isotope ratios utilized to assess the trophic levels of four species. Shown are the mean and standard deviation of nitrogen isotope ratios for four _Tetraponera_ ant species and herbivorous (purple box) and predatory (blue box) insects from the same site. The bottom side displays a phylogeny of the four ant species based on 1329 single-copy orthologs. **(c-e)** Scanning electron microscope (SEM) images showing the gut structure of _T. attenuata_ at different developmental stages: worker, pupa, and larva. **(f-h)** TEM images of ultrathin sections illustrating the bacteria within the bacterial pouch of _T. attenuata_ at worker (with different magnifications) and pupal stage, respectively. **(i-l)** Fluorescence in situ hybridization (FISH) detection of symbiotic bacteria within the bacterial pouch of four _Tetraponera_ ant species. DNA and bacteria are denoted by blue and magenta, respectively. CP, crop; MG, midgut; MT, Malpighian tubules; BP, bacterial pouch; HG, hindgut; B, bacteria; mv, microvillar differentiation; mt, mitochondria.

Microbiota of _T. nigra_-group ants are consistently dominated by pouch-associated Hyphomicrobiales bacteria

16S rRNA amplicon sequencing for dissected gasters of replicate workers from four to six colonies from each of the four _T. nigra_-group species revealed striking consistency in bacterial communities (Fig. 2, Supplementary Data 4). In all species, the microbiota was predominantly composed of a single OTU assigned to the order Hyphomicrobiales (formerly Rhizobiales), representing 99.4–100% of all reads in _T. attenuata_, 98.7–99.9% in _T._ sp1, 61.1–96.9% in _T. binghami_, 77.1–99.9% in _T._ sp2. Each ant species harbored a single distinct Hyphomicrobiales zOTU. In addition, we identified an OTU representing unclassified members of the Alphaproteobacteria class in three species. One of these zOTUs was present in all individuals of _T._ sp1 and _T._ sp2, and the other was found in all individuals of _T. binghami._ Despite consistent presence, their relative abundance was relatively low (e.g., 0.06–3.4% in _T._ sp2, 0.1–1.1% in _T._ sp1, 0.3–11.5% in _T. binghami_). Furthermore, we found that all _T. binghami_ individuals hosted an additional zOTU from the order Xanthomonadales, with relative abundances ranging from 1.4% to 27.7%. All other bacterial zOTUs combined comprised an average of 0.13% of reads in a library (ranging from 0 to 2.55%).

We used fluorescence in situ hybridization, with probes specific to each Hyphomicrobiales zOTU to localize them in ant guts. The signal overlapped with that for broad-spectrum bacterial probes – confirming that Hyphomicrobiales fill up the bacterial pouch (Fig. 2).

Fig. 2 Symbiont diversity and the localization of dominant Hyphomicrobiales bacteria within the gut of workers from four *T. nigra*-group species. The bar graphs show the relative abundance of bacterial zOTUs across individual workers from replicate colonies of four ant species. zOTUs with a relative abundance of less than 1% in any sample and the prevalence of less than 80% occurrence in any ant species are summed up as "others". ZOTUs were clustered into operational taxonomic units (OTUs) with 97% similarity. Above the bar graphs, fluorescence in situ hybridization (FISH) results illustrate the localization of Hyphomicrobiales bacteria within the worker ants' guts. Blue, DAPI; Red, Hyphomicrobiales-specific probe.

Comparative genomics reveals conserved symbiont lineages in the *T. nigra*-group

After sequencing metagenomes for pooled worker guts from the four *T. nigra*-group species, we performed metagenomic binning to generate draft symbiont genomes, all with more than 90% completeness and less than 1% contamination from the gut microbiomes of four ant species (Supplementary Table 3). The draft genome classification through comparison with the Genome Taxonomy Database aligned with the 16S rRNA amplicon-based findings (Fig. 3A). Specifically, all *Tetraponera* species hosted a dominant and unique Hyphomicrobiales strain. Additionally, *T. binghami* and *T.* sp1 were found to carry an unclassified Alphaproteobacteria strain, while *T. binghami* uniquely possessed a distinct Xanthomonadales strain.

The draft genomes of Hyphomicrobiales strains from the four ant species ranged

in size between 1.18 and 1.26 Mb, with an average GC content from 47.4% to 49.5%.

The maximum likelihood phylogeny based on a concatenated alignment of 385 amino

acid sequences of single-copy orthologous genes confirmed that the four *Tetraponera*

symbionts grouped together and formed a distinct, well-supported monophyletic clade,

along with other ant-associated Hyphomicrobiales symbionts with sequenced genomes

(Fig. 3B). This ant-associated bacterial clade forms a sister group to the *Bartonella*

clade, which includes *B. apis* of honeybees and pathogenic *Bartonella* species. This

result is also supported by the phylogenetic tree constructed based on 16S rRNA

sequences (Supplementary Fig. 2). Intriguingly, the phylogenomic relationships among

bacterial strains did not fully agree with the phylogeny of their ant hosts: strains from

T. attenuata and *T.* sp1 were not identified as sister clades, unlike their hosts. Further,

genome organization comparisons revealed that despite multiple rearrangements, there

are large areas of synteny among *T. nigra*-group symbiont genomes – much more so

than relative to *Tokpelaia* genomes from other ants (Supplementary Fig. 5). The 16S

rRNA gene identities and the average amino acid identities in all pairwise comparisons

between all ant-associated Hyphomicrobiales symbionts were 94.61%-97.91% and

62.33%-97.23% (Supplementary Table 4), respectively, suggesting that they should be

considered as members of the same genus *Tokpelaia*. Thus, we proposed to consider

the *Tetraponera nigra* group symbionts as strains of a single species, *candidatus*

Tokpelaia tetraponerae.

Our 16S rRNA gene phylogeny (Supplementary Fig. 2) indicated that the *T. nigra*-

group associated with unclassified Alphaproteobacteria and Xanthomonadales

symbionts were each grouped into a well-supported *Tetraponera*-specific clade,

respectively. Metagenomic binning further revealed the genomic characteristics of

these symbiont bacteria. The two unclassified Alphaproteobacteria strains had genomes

of approximately 1.13 Mb with a GC content of 27.4% (Supplementary Table 3).

Phylogenetic analysis placed them within the order Holosporales (Supplementary Fig.

3), a group known to include obligate intracellular or intranuclear symbionts of

protists[38, 39]. In addition, the Xanthomonadales bacterium, found exclusively in *T.*

binghami, had a genome size of 2.36 Mb and a GC content of 60.99% (Supplementary

Table 3). Phylogenetic analysis revealed its close relationship with 100% bootstrap support to *Frateuria aurantia* isolated from the lily plant *Lilium auratum.* However, taxonomic inconsistencies within the *Frateuria* genus preclude precise genus-level assignment, and thus, we provisionally classify it within the Rhodanobacteraceae family. While this last symbiont has the largest genome among the symbionts of *T. nigra*-group ants, it is still significantly reduced compared to the closely related bacteria (Supplementary Fig. 3).

Fig. 3 Taxon-annotated GC-coverage plots for four *Tetraponera* metagenomes and phylogenetic relationships of dominant Hyphomicrobiales bacteria of *T. nigra*-group ants and other hosts. (a) Assembled scaffolds in each metagenome are plotted based on their %GC content (x-axis) and their depth of sequencing coverage (y-axis, log scale). The colors of the blob plots indicate the taxa to which the scaffolds were assigned. Circle size represents the length of each scaffold. Scaffolds assigned to unclassified organisms are not shown. **(b)** Phylogenomic tree generated by maximum likelihood analysis based on 385 homologous single-copy genes. Red branches represent the Hyphomicrobiales genomes obtained from the metagenome of four *T. nigra*-group ant species. The purple bars represent the genome size, and the blue bars represent the GC content.

Nitrogen recycling and provisioning facilitated by the *T. nigra*-group ant symbionts

Given that the diets of these ants are deficient in nitrogen, yet their symbionts reside in a bacterial pouch enriched with nitrogenous waste such as uric acid delivered through the Malpighian tubules, we proposed that these symbionts likely convert nitrogenous waste into forms usable for synthesizing amino acids required by the ant host. Analysis of *Tokpelaia* genomes supports this hypothesis, as they retained genes encoding enzymes involved in nitrogen cycling from nitrogenous waste and biosynthetic pathways for the production of most essential amino acids that are needed by the ant hosts (Fig. 4).

All the *Tokpelaia* genomes harbored the *uacT* gene, which encoded a transporter for uric acid—the primary metabolic waste transported by the host's Malpighian tubules into the bacterial pouch. Further, all *Tokpelaia* strains contain gene clusters encoding enzymes involved in all the uric acid degradation pathway steps except for the last (Fig. 4). Interestingly, the missing gene, *alc*, which encodes allantoicase that converts allantoate to urea, is ancestrally lost in ant-associated *Tokpelaia* strains (Supplementary Fig. 10). This gene is present on Hymenoptera-assigned scaffolds in metagenomes of the *T. nigra*-group ants except *T.* sp1, suggesting that this reaction is likely carried out by the ant hosts. Following uric acid degradation, the product urea could be further processed into ammonia through the urease, encoded by three core genes, *ureC*, *ureB*, and *ureA* (Supplementary Data 7), present in all sampled metagenomes. Further, all *Tokpelaia* strains possess the genes *gdhA* and *glnA*, encoding enzymes glutamate dehydrogenase (GDH) and glutamine synthetase (GS), respectively (Fig. 4). These enzymes assimilate ammonia incorporated from urease-driven urea degradation into glutamate and glutamine, which serve as the nitrogen donors for the biosynthesis of amino acids.

The *Tokpelaia* genomes retain complete pathways for the biosynthesis of essential amino acids, including phenylalanine, tryptophan, tyrosine, arginine, lysine, threonine,

and methionine (Fig. 4, Supplementary Fig. 8). The isoleucine, leucine, and valine pathways are nearly complete, although *ilvH* gene encoding the small subunit of acetolactate synthase is missing. It is possible that these reactions are catalyzed by the large subunit (*ilvB*) of acetolactate synthase alone in *Tokpelaia,* as demonstrated in some Enterobacteria intact[40] enzymes.

Despite their specialized role in nitrogen metabolism, the four *Tokpelaia* strains exhibit striking metabolic constraints. While retaining genes for core cellular processes, including translation, replication, and transcription, their metabolic gene repertoire is notably limited (Supplementary Figs. 6-9). The symbionts' genomes lacked key genes involved in glycolysis, the biosynthesis of most B vitamins, certain non-essential amino acids, nucleotides, and KDO_2-lipid A. However, the metabolic pathways of these symbionts remain intact for the TCA cycle, oxidative phosphorylation, the biosynthesis of peptidoglycan, and fatty acids. Notably, the primary metabolic functions contributed by these symbionts are nitrogen recycling and provisioning, which account for approximately 12% of all protein-coding genes (Supplementary Fig. 7, Supplementary Data 7).

Analysis of the Xanthomonadales genome uniquely found in *T. binghami* revealed its metabolic redundancy, as it harbored genes encoding enzymes involved in ammonia assimilation and the biosynthesis of most amino acids (Fig. 4). Particularly, while genes *ilvH* and *hisB* were missing from across all examined *Tokpelaia* genomes, both were present in the Xanthomonadales genomes. In contrast, Holosporales strains lack biosynthetic pathways for almost all amino acids (Fig. 4, Supplementary Fig. 8), but encode transporters for oligopeptides and several amino acids. In addition, they also harbor glucose PTS transporters encoded by ptsG, enabling glucose to serve as the substrate fueling their TCA cycle (Supplementary Data 7). Like other Holosporales bacteria reported previously[41, 42], these bacteria are unlikely to be involved in ant nutrition, despite their strikingly consistent presence.

305 **Fig. 4 Amino acid biosynthetic and nitrogen (N) cycle utilization genes and pathways in the**
306 **genomes of *Tetraponera* ants-related symbionts.** The distinct colors of the arrows represent the
307 genomes of different symbiotic bacteria or hosts. Please note that this figure does not annotate the
308 entire metabolic symbiosis of the host; only the steps crucial to the nitrogen cycle pathway are
309 highlighted

Microbiota disruption negatively affects colony-wide fitness

Given the metagenomic evidence for nutritional provisioning by pouch-associated

symbionts, we assessed the symbiont impact on host survivorship and development. We

first generated symbiont-free workers by removing pupae from the original colonies to

sterile dishes. Those isolated pupae kept under sterile conditions emerged as adult

workers with no bacteria in the bacterial pouch, confirmed by universal bacterial PCRs,

as well as *Tokpelaia*-targeted 16S rRNA gene PCRs and whole-mount FISH staining

(Supplementary Fig. 11).

We then compared fitness parameters between the two groups of larvae with similar

sizes: one reared by normal workers (Sym-workers) and the other reared by symbiont-

free, aposymbiotic workers (Apo-workers), in two ant species, *T. attenuata* and *T.* sp1.

Through detailed daily observation of the larval development status over a 30-day

period, we observed that larvae tended by symbiont-free workers in both species

exhibited significantly lower survivorship (Fig. 5A; Wald statistic = 192.6, $p < 2e\text{-}16$

for *T.* sp1; Wald statistic = 180.6, $p < 2e\text{-}16$ for *T. attenuata*) and slower development,

as indicated by reduced body length (Fig. 5B; $\chi^2 = 897.982$, $p < 2.2e\text{-}16$ for *T.* sp1;

$\chi^2 = 721.6$, $p < 2.2e\text{-}16$ for *T. attenuata*; GLMM) compared to larvae reared by normal

workers.

In another long-term experiment lasting 90 days, we monitored the continuous

development of the larvae until they emerged as adults. After 90 days, the number of

missing broods was higher in the group tended by symbiont-free workers compared to

the one reared by normal workers only in *T. attenuata* (Fig. 5C, Supplementary Table

8; p = 0.0848; t-test). In both ant species, we observed that larvae raised by symbiont-

free workers exhibited significantly lower pupation rates (Fig. 5C, Supplementary

Table 8; *T.* sp1, p = 0.0096; *T. attenuata*, p = 0.2326; T-test), reduced adult emergence

rates from pupae to adults (Fig. 5C, Supplementary Table 8; *T.* sp1, p = 0.1999,

Wilcoxon-Mann-Whitney test; *T. attenuata*, p = 0.2271, t-test), as well as smaller head

widths at the time of emergence (Fig. 5C, Supplementary Table 8; *T.* sp1, p = 0.1818,

Wilcoxon-Mann-Whitney test; *T. attenuata*, p = 0.0453, t-test), compared to those

reared by normal workers. We also noted that among the adult ants tending the larvae,

symbiont-free workers exhibited significantly lower survivorship compared to their

symbiont-bearing counterparts (Supplementary Fig. 13; Wald statistic = 94.22, p $<$ 2e-

16 for *T.* sp1; Wald statistic = 3.44, p = 0.06 for *T. attenuata*).

Larvae by Sym-worker ⎯⎯ Larvae by Apo-worker ⋯⋯⋯ *T. attenuata* ⎯⎯ *T.* sp1

Fig. 5 Comparison of fitness parameters between two groups of larvae with similar sizes, one reared by normal workers (Sym-workers) and the other reared by symbiont-free workers (Apo-workers), in two ant species, *T. attenuata* and *T.* sp1. (a) The survival rate of larvae cared for by Apo-workers or Sym-workers over a 30-day experimental period. **(b)** Changes in the relative body length of larvae reared by Sym-workers and Apo-workers. Body length data from day 2 to day 30 were normalized to the average body length of larvae on day 1 for each treatment group. **(c)** Comparison of fitness parameters at day 90, including the number of missing larvae, pupation rate, eclosion rate, and head width of newly emerged workers as an indicator of body size. The light green color represents the control group of worker ants with symbiotic bacteria, while the purple indicates the experimental group of worker ants without symbiotic bacteria. In the line graphs, the dashed lines represent *T. attenuata*, and the solid lines represent *T.* sp1.

Symbionts help ants recycle nitrogen from urea

To experimentally test whether genetic signatures reflect the abilities of symbionts to degrade nitrogenous wastes and synthesize amino acids, we conducted feeding experiments with ^{15}N-labelled urea (Supplementary Fig 12) to two species from the *T. nigra*-group. Within each ant colony, individuals were assigned to one of three experimental groups: (1) symbiotic workers fed ^{15}N-labelled urea; (2) symbiont-free workers fed ^{15}N-labelled urea; and (3) symbiont workers fed ^{14}N-labelled urea as a control.

In the first experiment, we focused exclusively on workers. We examined whether recycled nitrogen from ^{15}N-labelled urea could be incorporated into the free amino acids present in the hemolymph of adult ants. Among symbiotic workers fed with ^{15}N-urea, we detected a significantly higher proportion of ^{15}N in 14 of 19 measurable amino acids compared to the control group fed diets containing only standard urea (significant p-value range: 1.04E−05 to 0.076, Fig. 6A, Supplementary Table 9). These enriched amino acids included seven essential amino acids (arginine, histidine, isoleucine, phenylalanine, tryptophan, tyrosine, and valine), as well as seven non-essential amino acids (alanine, asparagine, glutamate, glutamine, glycine, proline, and serine). Crucially, this heavy isotopic enrichment was dependent on symbiont presence. Symbiont-free workers fed ^{15}N-labelled urea showed no such enrichment in the same amino acids relative to the ^{14}N-labelled urea fed control ants. These results highlighted the pivotal role of bacterial pouch symbionts in recycling nitrogen, particularly in facilitating amino acid synthesis essential to the host.

In the second experiment, we investigated how symbiotic bacteria within the bacterial pouch of workers mediate nitrogen allocation to larvae, given the severe negative effects of worker microbiota disruption on larval growth and development (Fig. 5). By tracking ^{15}N incorporation from ^{15}N-urea into the amino acids of larval body tissue, we found that larvae reared by symbiotic workers fed ^{15}N-urea exhibited ^{15}N enrichment in 11 amino acids of 18 measurable amino acids (significant p-value range: 0.001 to 0.0861, Fig. 6B, Supplementary Table 10). In contrast, larvae tended by either symbiont-free workers (^{15}N-urea fed) or symbiotic workers (^{14}N-urea fed) showed no such enrichment. Among these 11 enriched amino acids, there are nine essential amino acids (arginine, histidine, isoleucine, leucine, lysine, phenylalanine, tryptophan, tyrosine, and valine) and two non-essential amino acids (glycine and serine). This highlights the worker symbionts' role in bridging nutritional needs critical for larval growth and development.

To evaluate how symbiont-processed nitrogen is allocated among members within the whole colony, the relative incorporation of ^{15}N into worker and larval tissues was quantified. We observed that δ^{15}N values in the body tissues of larvae attended by

symbiotic workers for 30 days were significantly higher compared to those reared by aposymbiotic workers in two ant species (*T. attenuata*, p=0.0373; *T.* sp1, p=0.0399; post hoc Dunn's test; Supplementary Fig. 14, Supplementary Table 11). Similarly, $\delta^{15}N$ values in the body tissues of symbiotic workers were higher than those of aposymbiotic workers (*T. attenuata*, p = 0.1062; *T.* sp1, p = 0.1052; post hoc Dunn's test; Supplementary Fig. 14, Supplementary Table 11). Notably, comparative analysis revealed that larvae exhibited 2.2-fold and 2.3-fold higher ^{15}N enrichment compared to symbiotic workers tending them in *T. attenuata* and *T.* sp1 respectively (*T. attenuata*, p = 0.0022; *T.* sp1, p = 0.0005; t-test). These findings reflected this larval-biased nitrogen allocation as an adaptive investment strategy, where colonies prioritize developmental stages to maximum fitness under N-limiting conditions.

Fig. 6 Depriving symbiotic bacteria reduces the proportion of ^{15}N-labeled amino acids in the worker ants and larvae that have consumed ^{15}N-labeled urea. (a) The proportion of ^{15}N-labeled amino acids in the hemolymph of *T. attenuata* workers was measured under three conditions: symbiotic workers fed ^{15}N-labeled urea, symbiont-free workers fed ^{15}N-labeled urea, and symbiotic workers fed ^{14}N-labeled urea as a control. Detailed information on sample sizes for each treatment can be found in Supplementary Data 10. The statistical results are described in Supplementary Methods and Supplementary Table 9. **(b)** The proportions of ^{15}N-labeled amino acids in the body tissues of larvae cared for by workers of *T. attenuata* and *T.* sp1 under the same three conditions. Due to limited sample size per species (n=3 under one condition) and non-significant interspecific effect on the proportion of ^{15}N amino acids (Supplementary Table 11), data from two species were

415 pooled across group for a combined analysis using one-way ANOVA (for normally distributed data)
416 or Kruskal Wallis test (for non-normal data).

Discussion

Our study has demonstrated the unique location of symbiotic bacteria within a gut

pouch (Fig. 1H-1K), a special structure consistently present across the *T. nigra*-group.

We also demonstrated for the first time the ants' ancestral symbioses with a co-

diversifying *Tokpelaia* symbiont, while also maintaining consistent associations with

some other bacteria (Figs. 2 and 3). Through the characterization of the symbionts'

genomic characteristics, we have elucidated the mechanisms underlying this nutritional

symbiosis (Fig. 4). Additionally, we have demonstrated experimentally the effects of

symbiosis disruption on colony development, particularly highlighting the impact on

larval growth (Fig. 5). Finally, we have experimentally established links between

symbiont genomic characteristics and ant nutrition, explaining the observed fitness

effects (Fig. 6). Considering the diversity, global distribution, and ecological

importance of social insects, it is surprising how few clades and species have

comparably detailed data on microbial associations. Through this study, the *T. nigra*-

group ants are added to the short list of social insect clades whose symbioses are

considered well-understood.

The *T. nigra*-group is distinguished by a unique bacterial pouch—a structure not

previously reported in other hymenopteran clades—highlighting an unconventional

method of symbiont localization. In contrast, other hymenopteran insects exhibit

diverse anatomical niches for hosting symbiotic bacteria. For instance, *Blochmannia* is

found in bacteriocytes situated between midgut cells and in female ovaries[43, 44]; species

such as honeybees[45, 46], bumblebees[47], turtle ants[17, 48], and army ants[49] harbor multi-

partite bacteria within their guts; and fungus-farming ants maintain bacterial

communities in specialized exoskeletal crypts essential for the health of their fungal

gardens[50]. However, several other groups of insects, including stinkbugs[51] and leaf

beetles[32, 52], host extracellular bacteria in a symbiotic organ consisting of pouch-like gut

invagination. These symbiotic organs ensure a stable supply of resources for the

symbiotic bacteria, which promoting their growth while limiting competition with other

microbial species[53]. Meanwhile, these structures restrict the entry of symbionts into

other host tissues, minimizing disruption to the hosts' other physiological processes[53].

We observed that the larvae of *T. nigra*-group ants lack a bacterial pouch (Fig. 1E).

Although the pouch first develops during the pupal stage (Fig. 1D), it remains

uncolonized (Fig. 1H) until the ants eclose into adults (Fig. 1C). This indicated that this

symbiotic organ develops independently of symbiont colonization. A similar pattern of

organ development, proceeding without symbiont triggering, has been observed in

tortoise beetles, reflecting an ancient evolutionary trait in which the formation of such

organs proceeds autonomously[54]. Moreover, pupae isolated from their colony and

allowed to eclose in isolation showed no bacterial colonization of their pouch

(Supplementary Fig. 15), suggesting that symbiotic bacteria are likely transmitted to

newly emerged adults through trophallactic interactions within the colony. Such

trophallaxis behaviors may play a key role in maintaining partner fidelity, not only in

these groups of ants but also across several other social insects. For instance, symbionts

are transmitted between termite siblings via oral-anal trophallaxis[55], and a similar

transmission mode has been observed in *Cephalotes* and *Procryptocerus* ants[56]. In

honeybees, gut symbiotic bacteria are transmitted through a combination of trophallaxis

or contact with nest materials within the hive[57]. Future investigation of transmission

mechanisms of symbiotic bacteria in the bacterial pouch can shed light on key drivers

of co-diversification between the *T. nigra*-group ants and their symbiotic bacteria.

High-throughput sequencing revealed an ant-specific, *Tokpelaia*-dominated

bacterial community in the *T. nigra*-group ants' pouch, suggesting a specialized and

long-standing association between the *T. nigra*-group ants and their symbiotic bacteria.

Our conclusions are further strengthened by the phylogenetic similarity between the

symbionts characterized here and those previously generated from other *T. nigra*-group

species[58, 59]. Our findings suggest that the *T. nigra*-group ants belong to a growing

number of insect clades whose extracellular microbiota are dominated by one or a few

specialized symbionts—a pattern similarly observed in other insect groups with

specialized symbiotic organs, such as stinkbugs[7] and leaf beetles[32, 60, 61]. One possible

explanation is that these symbiotic organs function to filter microbial strains, allowing

only specific symbionts to access the host and maintain a beneficial partnership. This

contrasts with the complex multi-partite gut microbiota observed in termites[62, 63],

Cephalotes ants[17], and corbiculate bees[46], or with occasional and non-specialized

microbial association seen in diverse other ants and other insects[64].

The critical role of symbiotic bacteria in the whole ant colony fitness is evident in

two key aspects. Symbiont workers fed diets where urea was the only source of nitrogen

exhibited lower survival rates, and the larvae they cared for showed impaired

development, reduced survival, lower pupation and emergence rates, and small adult

body sizes upon eclosion (Fig. 4). These results suggest the importance of symbiont-

mediated nitrogen metabolism, which supports the overall fitness of the ant colony. We

indeed observe that, in the presence of symbionts within the bacterial pouch, a greater

proportion of nitrogen derived from urea was assimilated into the body tissues of both

adults and larvae, with larvae exhibiting a higher rate of nitrogen assimilation

(Supplementary Fig. 14). Genomic analysis and heavy nitrogen urea labeling

experiments revealed that symbionts in the bacterial pouch recycle nitrogenous waste,

such as urate or urea, and convert symbiont-recycled nitrogen in the form of essential

and non-essential amino acids utilized by ant hosts (Figs. 5 and 6). Taken together, the

conserved N-recycling and N-provisioning roles of symbionts in the bacterial pouch of

T. nigra-group ants support the hypothesis that this nutritional symbiosis plays an

important role in the exploitation of an N-limited dietary niche. Similar functions are

observed in other herbivorous ants, such as the endosymbiont *Blochmannia* in

Camponotini[25] and the multi-partite gut bacterial communities in *Cephalotes* ants[17],

where symbionts improve nitrogen economies by recycling nitrogen and further fueling

the synthesis of essential amino acids. Despite differences in the symbiotic microbes

involved, these herbivorous ants with nitrogen-poor diets exhibit a functional

convergence in the adaptive role of symbiosis for nitrogen acquisition. This reflects the

generality of such functional convergence across herbivorous ant species inhabiting N-limited environments.

The specialized structure connecting the bacterial pouch to the Malpighian tubules reinforces the proposed biological role of *Tokpelaia*. The nitrogen recycling activity by *Tokpelaia* is likely mediated through the delivery of host-derived waste nitrogen via the Malpighian tubules into the bacterial pouch, where *Tokpelaia* resides. Additionally, this structure's connection to the gut facilitates the export of amino acids synthesized from assimilated recycled nitrogen by *Tokpelaia* cells, which are then transported to the host gut. The high abundance of nitrogen-recycling symbionts at the site of insect nitrogen waste delivery via the Malpighian tubules is also observed in the *Cephalotes* system. Positioned at the site of host nitrogen waste delivery[17, 48], *Ischyrobacter davidsoniae* plays a central role in a collaborative nitrogen-recycling symbiosis[23]. These findings highlight the critical role of specialized symbiotic structures in facilitating nitrogen recycling in insects, where the spatial arrangement of symbionts near nitrogen waste delivery sites enables efficient nutrient cycling and contributes to the overall ecological success of the host.

During the metamorphosis of holometabolous insects, significant morphological and physiological changes provide opportunities for symbiotic mutualisms to become dissociated. A notable example is the bacterium *Acetobacter thailandicus*, which stably colonizes the gut of Drosophila adults. While it does not enhance adult fitness, it significantly increases larval fitness[65]. This pattern may reflect a broader trend in holometabolous insects, where adults carry and transmit symbiotic bacteria that primarily benefit larvae, even without direct advantages to the adults. In social insects, such cross-stage symbiont benefits are more pronounced due to the close physical association between adults and larvae and the capacity for trophallactic feeding of symbiont-derived metabolites produced by adults. In the *T. nigra*-group ants, specialized *Tokpelaia* symbionts present exclusively in adults, enhance colony-wide nitrogen economies. Their contributions to larval nitrogen budgets promote larval growth and development. This adult-specific symbiosis is likely sustained through the

531 trophallactic transfer of extracellular symbionts during the adult stage.

532 Our findings suggest that this sophisticated structural design enables a highly

533 efficient nitrogen economy within the holobiont of *T. nigra*-group ants. The strict

534 compartmentalization of these structures stabilizes the symbiotic relationship by

535 allowing the ant hosts to deliver precursors of the uricolytic pathway directly to the

536 bacterial pouch. The symbionts, which reside within this structure, import uric acid

537 and produce metabolites that are beneficial to the ants, functioning as "nutrient

538 factories". These symbiont-derived metabolites are then distributed to colony members,

539 particularly larvae that lack these symbionts, via trophallaxis, ensuring that the entire

540 colony benefits from the nutritional symbiosis. This study enhances our understanding

541 of insect-microbe symbioses, particularly in social insects, by adding the *T. nigra*-group

542 ants to the list of species with well-characterized, specialized microbial associations.

543 The findings highlight the importance of spatial organization in symbiotic systems,

544 where specialized organs facilitate nutrient recycling and support host fitness. This

545 work also provides new insights into nitrogen acquisition strategies in other

546 herbivorous insects, like *Cephalotes* and *Camponotus* ants, where convergent symbiotic

547 mechanisms may help overcome nitrogen limitations[17, 25]. Ultimately, this research

548 opens avenues for further exploration of microbial co-diversification and the functional

549 roles of symbioses in host evolution.

Methods

Ant collection and specimen identification

552 We collected 53 *Tetraponera* colonies from southern provinces of China, including

553 Yunnan, Guangdong, and Jiangxi, between 2019 and 2024 (Supplementary Data 1). For

554 molecular analysis, multiple workers were preserved in anhydrous ethanol and stored

555 at -20°C before DNA extraction. Colonies designated for dietary manipulation

556 experiments or microscopic observation were kept alive for laboratory rearing.

557 For identification, one individual from each colony was photographed and

classified based on morphology. Additionally, we selected another individual per colony for DNA barcoding of the mitochondrial cytochrome oxidase I gene (COI) to refine species identification, with COI sequences >600 bp unambiguously matched to *Tetraponera* species in GenBank and BOLD database[66] included as a taxonomic reference. Using the COI sequence of *Pseudomyrmex gracilis* (FJ436821.1), a sister genus of *Tetraponera*, as an outgroup, we aligned all sequences with MAFFT v7.520[67] and removed divergent or ambiguously aligned regions using Gblocks v0.91b[68]. Phylogenetic analyses were conducted using RAxML v8.2.12[69] with 1000 bootstrap replicates under the GTRGAMMA model, and the resulting phylogeny was visualized using iTOL[70]. Species delimitation was further assessed using the Automatic Barcode Gap Discovery (ABGD) method[71], which detects barcode gaps by assuming non-overlapping intra- and interspecific genetic distances. Pairwise K2P distance matrices were calculated from the COI alignment (excluding the outgroup) using MEGA v11.0.13[72]. These matrices were used for ABGD analysis performed online, with a relative gap width (X) of 1.5, and results recorded for intraspecific divergence (P) ranging from 0.001 to 0.1 (Supplementary Fig. 1).

Trophic levels of *Tetraponera* ants determined by nitrogen isotope ratios

While collecting *Tetraponera* ants, we also gathered local plants, herbivores, and predatory arthropods (Supplementary Data 2). To prevent contamination from residual food in arthropod guts, we removed ant abdominal segments (gasters) before isotopic analysis. Plant leaves and arthropod samples were dried using a vacuum-freeze dryer, then ground for $\delta^{15}N$ analysis using a Delta A Advantage isotope ratio mass spectrometer coupled with an EA-HT elemental analyzer (Thermo Fisher Scientific Inc., Bremen, Germany). The nitrogen isotope ratios expressed as $\delta^{15}N$ value was calculated according to the equation: $\delta(‰) = [(Rsample - Rstandard)/Rstandard]*1000$, where Rsample is the $^{15}N/^{14}N$ ratio of the sample and Rstandard is the $^{15}N/^{14}N$ ratio of atmospheric N_2 (Rstandard = 0.003676). To conduct comparisons across different

587 locations, we computed the average δ^{15}N values for plants (δ^{15}N plant) examined at

588 each specific site. The relative trophic level of ants and arthropods was then determined

589 by subtracting δ^{15}N plant from δ^{15}N ant/arthropod at each location. Finally, we adjusted

590 for local variation by subtracting the median δ^{15}N value of plants at each site from the

591 overall median value across all locations.

Microscopic analysis of bacterial pouch structure and symbiont localization in *T. nigra*-group ants

To explore the organization of the bacterial pouch across different life stages in *T.*

nigra-group ants, we examined the gut structure of workers, pupae, and larvae of *T.*

attenuata using scanning electron microscopy (SEM). Guts were dissected and fixed

overnight at 4°C in 2.5% glutaraldehyde, followed by dehydration through a graded

ethanol series and drying using a critical point dryer (Leica CPD300). After sputter-

coating with gold (Hitachi E-1045), the samples were observed under an SEM (Hitachi

SU8010, Tokyo, Japan).

Fluorescence in situ hybridization (FISH) was then used to localize the symbiotic

bacteria in workers from four *T. nigra*-group species, following a previously published

protocol[49] Łukasik et al. (2017). Guts were dissected, fixed in 4% formaldehyde in PBS

at room temperature for 2 hours, and treated with 80% ethanol and 6% hydrogen

peroxide for 1–4 weeks to reduce autofluorescence. The samples were then rehydrated

by washing with PBSTx (PBS with 0.3% TritonX-100) for 10 minutes three times,

followed by washing with hybridization buffer (30% formamide, 0.01% SDS, 0.9 M

NaCl, and 0.02 M Tris-HCl, pH 8.0) for 10 minutes three times. Finally, samples were

incubated overnight at room temperature in a hybridization buffer containing 100 nM

EUB338 universal bacterial probe and 1% DAPI. After washing the specimens three

times for 10 minutes each with PBSTx, specimens were imaged using an inverted

fluorescence microscope (Observer Z1, Zeiss, Germany). To further identify specific

symbionts, a second round of FISH was performed using species-specific probes

targeting   Hyphomicrobiales,   including   RHIZ1244[73]   (Cy5-

TCGCTGCCCACTGTCACC) for *T. attenuata* and *T.* sp2, TBIRHI (Cy5-

TCGCTACCCACTGTCACC) for *T. binghami*, and TSP1RHI (Cy5-

TCGCTACCCATTGTCACC) for *T.* sp1.

To obtain detailed structure information on the bacterial pouch and associated

bacteria, we used transmission electron microscopy (TEM). Guts of *T. attenuata* were

dissected and fixed overnight in 2.5% glutaraldehyde in PBS at 4°C. After washing with

PBS four times, the samples were postfixed with 1% (w/v) osmium tetroxide in 0.1 M

PBS for 2h at room temperature. Then, the samples were dehydrated in a graded ethanol

series and then in pure acetone for 10 minutes twice. Then, the samples were infiltrated

in a graded mixture (3:1, 1:1, 1:3) of acetone and SPI-PON812 resin (16.2 g SPI-

PON812, 10 g DDSA and 8.9 g NMA) and then infiltrated in pure resin. Finally, tissues

were embedded in pure resin with 1.5% BDMA and polymerized at 45°C for 12h and

then at 60°C for 48 h. The ultrathin sections (70nm thick) were sectioned

with microtome (Leica EM UC6), double-stained by uranyl acetate and lead citrate, and

examined under a transmission electron microscope (Tecnai Spirit120kV, FEI, USA).

Bacterial 16S rRNA amplicon sequencing

We investigated the microbial community diversity of *T. nigra*-group ants through

the V4 region of the 16S rRNA amplicon sequencing. For each species, samples were

collected from 4-6 colonies, with 3-5 workers sampled per colony. The gaster of a single

worker ant was dissected using sterile forceps, and total genomic DNA was extracted

from each gaster using the DNeasy Blood and Tissue Kit (Qiagen, Germany), following

the manufacturer's protocol for Gram-positive bacteria. DNA was submitted to Beijing

Novogene Bioinformatics Technology Co., Ltd for amplicon library preparation. Along

with insect DNA samples, blank samples in our DNA extraction batches were included

to monitor potential contamination during sample preparation and sequencing. Pooled

libraries were sequenced on the Illumina NovaSeq6000 SP platform, generating

2x250bp reads (Supplementary Data 3).

The raw reads were demultiplexed, trimmed barcodes and primers, and merged

reads using FLASH v1.2.7[74]. Quality filtering was done with fastp v0.23.4[75], and

downstream analysis was performed using a modified version of the

"modified_LSD.py" script[76] from

https://github.com/catesval/army_ant_myrmecophiles. We made changes to the

database in the script to suit our requirements. Briefly, low-quality sequences were

removed using vsearch-2.15.2[77], followed by chimera removal and denoising with

usearch11.0.667[78], which produced zOTU abundance tables. zOTUs were clustered

into OTUs based on 97% similarity using the -cluster_otus and -uparseout parameters

in usearch, and the resulting clusters were checked for chimeras. Species classification

was performed by comparing sequences against the RDP database

(http://www.drive5.com/sintax/rdp_16s_v18.fa.gz) using vsearch.

Bacterial DNA present in reagents and the environment can significantly impact

microbial community profiles, especially in samples with low template concentrations[79].

To eliminate potential contamination, we implemented several steps following

established criteria[49] . First, we performed BLASTn searches against the NCBI database

to assign taxonomy. Sequences identified as mitochondria, chloroplasts, Archaea,

Eukarya, or chimera were then removed from further analysis. Next, sequences

obtained from cloning amplification experiments (Accession numbers PQ549644-

PQ549668), specifically designed to identify bacterial species in the ant gut, were used

as reference sequences. zOTUs that perfectly matched these reference sequences were

initially added to the final zOTU list. The remaining zOTUs were then screened for

potential contamination. Contaminants were identified by comparing their relative

abundance in experimental libraries to that in blank libraries. Specifically, zOTUs with

a maximum relative abundance at least ten times greater in any experimental library

than in any blank library were considered non-contaminants. Those non-contaminant

zOTUs were added to the final zOTU list for further analysis (Supplementary Data 4).

Metagenomics

670　　　For each ant species, we dissected the gut tissues of 10 workers from the same

671　colony using sterile forceps and pooled them into a single sample. The extracted DNA

was then sent to Beijing Novogene Bioinformatics Technology Co., Ltd for library preparation, followed by metagenomic sequencing on the Illumina NovaSeq 6000 platform. Quality trimming of raw reads was done with Trimmomatic-0.39[80], followed by read quality checks using FastQC v0.11.9(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). The trimmed reads were assembled with IDBA-UD 1.1.3[81] using k values of 20, 40, 60, 80, and 100, and assembly statistics were calculated with QUAST-5.0.2[82] (Supplementary Table 2). Scaffolds were uploaded to the Integrated Microbial Genomes with Microbiome Samples Expert Review (IMG/M-ER) website for taxonomic classification and gene annotation[83]. We focused on N metabolism by utilizing KEGG as references to manually construct the pathways involved in the degradation of nitrogenous wastes and the biosynthesis of amino acids (Supplementary Data 7). The bacterial community composition of all metagenomes in this study was visualized by the Taxon-Annotated GC Coverage plots, which were generated by mapping raw reads to classified scaffolds using BWA 0.7.1[84] and employing custom scripts based on a previously published method[85].

Metagenomic binning and genome-based phylogenetic analysis of symbiotic Bacteria

In order to obtain the draft genomes of individual symbiont strains, we used the metaWRAP pipeline[86] using CONCOCT, MaxBin, and metaBAT binning modules. The draft genomes were refined with the bin_refinement module using ≥80% completeness and ≤10% contamination. Few scaffolds belonging to these draft genomes are not classified or not predicted to be present in the *T. nigra*-group ants. To improve accuracy, scaffolds from seven draft genomes were compared against the NCBI database using BLASTn and BLASTx with an e-value threshold of 10^{-6} and 70% identity. Scaffolds with unexpected classification or their read coverage significantly deviates from that of most scaffolds within their respective draft genomes were manually removed from the binning datasets (Supplementary Data 5). Bin completeness and contamination were

assessed with CheckM[87] and the taxonomic assignment was performed using the

Genome Taxonomy Database with gtdbtk-2.1.1[88] (Supplementary Table 3).

We further performed genome-based phylogenetic analysis of symbiotic bacteria

in *T. nigra*-group ants using draft genomic data. OrthoFinder v2.5.5[89] was used to

identify single-copy orthologous genes. Amino acid sequences of these genes were

aligned with MAFFT v7.520, and divergent/ambiguous regions were removed using

Gblocks v0.91b. Alignments were concatenated for phylogenetic inference. Prottest

v3.4.2[90] was used to select the optimal amino acid substitution model. Phylogenetic

analysis was then performed with RAxML v8.2.12 using the best model and 1000

bootstrap replicates. Results were visualized and annotated on iTOL (Supplementary

Data 6).

Symbiont manipulation and ^{15}N-labeled urea feeding experiments

In order to perform symbiont manipulation experiments, we first obtained large

numbers of symbiont-free workers without the use of antibiotics. *T. attenuata* and *T.*

sp1 colonies were collected from Yunnan Province (Supplementary Data 1) and reared

on a diet of 30% honey water and mealworms at 25°C with a 14:10 light-dark cycle. A

fresh diet was provided every two days. To generate symbiont-free workers, pupae with

pigmented eyes were removed from the original cages and placed in sterile dishes.

Those pupae kept under sterile conditions will emerge as adult workers with no bacteria

in the bacterial pouch, confirmed by PCR amplification using both universal bacterial

and *Tokpelaia*-specific 16S rRNA gene primers, as well as whole-mount FISH staining

(Supplementary Fig. 11).

Controlled lab experiments were conducted on two ant species, *T. attenuata* and *T.*

sp1, to quantify the symbionts' contributions to the colony-wide nitrogen economy

through N recycling of dietary N waste (Supplementary Fig. 12). For each species, three

or four colonies were selected as biological replicates. Within each colony, individuals

were randomly assigned to one of the three treatments: (1) workers harboring symbiotic

bacteria in the bacterial pouch (Sym-worker), fed 30% sucrose water containing 1%

(w/v) ^{15}N-labeled urea; (2) workers without symbiotic bacteria in the bacterial pouch

(Apo-worker), fed the same ^{15}N-labeled urea solution; (3) workers with symbiotic

bacteria in the bacterial pouch, fed 30% sucrose water containing 1% (w/v) ^{14}N-labeled

urea as a control.

We initially conducted the ^{15}N-labeled urea feeding experiments exclusively with

worker ants for 3 weeks. At the end of the experiment, hemolymph was collected from

the surviving workers. Based on the number of remaining workers, we aimed to ensure

2–3 replicates for each treatment group within each colony (Supplementary Data 10).

Hemolymph from each replicate was pooled to measure the enrichment of ^{15}N in amino

acids within ant hemolymph at the Scale Biomedicine Technology Co., LTD (Beijing,

China). Chromatographic separation of each hemolymph-water mixture was performed

on Ultra-High Performance Liquid Chromatography (UHPLC)–High Resolution

Tandem Mass Spectrometry (HRMS/MS).

Next, we extended the ^{15}N-labeled urea feeding experiments to include larvae

tended by adult workers, with experimental durations of 30 days and 90 days. In the

30-day experiment, each treatment group was assigned 10 workers and 30 larvae. We

recorded worker and larval survival throughout the whole period. In addition, we

monitored larval growth and development by measuring larval body length daily. At the

end of the experiment, the ^{15}N abundance in larval and worker body tissues, excluding

the gut to avoid potential food contamination, was measured using a MAT253 stable

isotope mass spectrometer at the Beijing Academy of Agriculture and Forestry Sciences

(BAAFS) to determine whether the nitrogen assimilated from urea by symbiotic

bacteria was utilized by the host and incorporated into its body tissues. At the same time,

3-5 larvae from each group were sent to Scale Biomedicine Technology Co., LTD

(Beijing, China), where the relative abundance of ^{15}N-labeled amino acids was analyzed

using UHPLC-HRMS/MS. In addition, we performed the 90-day experiment, with each

group assigned 15 workers and 50 larvae, to record larval pupation and eclosion rates

and measure the head width of newly emerged workers as an indicator of the worker

body size. The detailed methods and statistical analysis can be found in the

Supplementary Methods.

Acknowledgements

This study was supported by the National Natural Science Foundation of China 32370448 to Y. H., Beijing Advanced Innovation Program for Land Surface Science and the "111" Program of Introducing Talents of Discipline to Universities (B13008).

Data Availability

All data generated in this study are publicly available. The 16S amplicon sequencing data have been deposited in the GenBank Short Read Archive under BioProject PRJNA1159735, while the 16S rRNA sequences obtained via Sanger sequencing are accessible in GenBank under accession numbers PQ549644-PQ549668. Additionally, raw data from the stable isotope mass spectrometer and UHPLC-HRMS/MS are provided in the Supplementary Tables, and details on the PCR primers and FISH probes can be found in the Supplementary Tables and Methods.

References

1. Markmann, K. & Parniske, M. Evolution of root endosymbiosis with bacteria: how novel are nodules? *Trends Plant Sci.* **14**, 77-86 (2009).

2. Desbrosses, G. J. & Stougaard, J. Root nodulation: a paradigm for how plant-microbe symbiosis influences host developmental pathways. *Cell Host Microbe* **10**, 348-358 (2011).

3. McFall-Ngai, M. J. The importance of microbes in animal development: lessons from the squid-vibrio symbiosis. *Annu. Rev. Microbiol.* **68**, 177-194 (2014).

4. Belcaid, M., et al. Symbiotic organs shaped by distinct modes of genome evolution in cephalopods. *Proc. Natl. Acad. Sci. USA* **116**, 3030-3035 (2019).

5. Skelton, J., Johnson, A. J., Jusino, M. A., Bateman, C. C., Li, Y. & Hulcr, J. A selective fungal transport organ (mycangium) maintains coarse phylogenetic congruence between fungus-farming ambrosia beetles and their symbionts. *Proc. R. Soc. B* **286**, 20182127 (2019).

6. Goodchild, A. J. P. Studies on the functional anatomy of the intestines of Heteroptera. *Proceedings of the Zoological Society of London* **141**, 851-910 (1963).

7. Kuechler, S. M., Fukatsu, T. & Matsuura, Y. Repeated evolution of bacteriocytes in lygaeoid stinkbugs. *Environ. Microbiol.* **21**, 4378-4394 (2019).

8. Oishi, S., Moriyama, M., Koga, R. & Fukatsu, T. Morphogenesis and development of midgut symbiotic organ of the stinkbug *Plautia stali* (Hemiptera: Pentatomidae). *Zool. Lett.* **5**, 16-29 (2019).

9. Jackson, R., et al. Convergent evolution of a labile nutritional symbiosis in ants. *ISME J.* **16**, 2114-2122 (2022).

10. Russell, J. A., Sanders, J. G. & Moreau, C. S. Hotspots for symbiosis: function, evolution, and
specificity of ant-microbe associations from trunk to tips of the ant phylogeny (Hymenoptera:
Formicidae). *Myrmecological News* **24**, 43-69 (2017).

11. Russell, J. A., Moreau, C. S., Goldman-Huertas, B., Fujiwara, M. & Lohman, D. J. Bacterial gut
symbionts are tightly linked with the evolution of herbivory in ants. *Proc. Natl. Acad. Sci. USA* **106**,
21236-21241 (2009).

12. Pringle, E. G. Convergence, constraint and the potential for mutualism between ants and gut microbes.
Mol. Ecol. **28**, 699-702 (2019).

13. Sanders, J. G., et al. Dramatic differences in gut bacterial densities correlate with diet and habitat in
rainforest ants. *Integr. Comp. Biol.* **57**, 705-722 (2017).

14. Klein, A., et al. A novel intracellular mutualistic bacterium in the invasive ant *Cardiocondyla*
obscurior. *ISME J.* **10**, 376-388 (2016).

15. Neuvonen, M. M., et al. The genome of Rhizobiales bacteria in predatory ants reveals urease gene
functions but no genes for nitrogen fixation. *Scientific Reports* **6**, 39197 (2016).

16. Bisch, G., et al. Genome evolution of Bartonellaceae symbionts of ants at the opposite ends of the
trophic scale. *Genome Biol. Evol.* **10**, 1687-1704 (2018).

17. Hu, Y., et al. Herbivorous turtle ants obtain essential nutrients from a conserved nitrogen-recycling
gut microbiome. *Nat. Commun.* **9**, 964-978 (2018).

18. Blüthgen, N., Gebauer, G. & Fiedler, K. Disentangling a rainforest food web using stable isotopes:
dietary diversity in a species-rich ant community. *Oecologia* **137**, 426-435 (2003).

19. Davidson, D. W., Cook, S. C., Snelling, R. R. & Chua, T. H. Explaining the abundance of ants in
lowland tropical rainforest canopies. *Science* **300**, 969-972 (2003).

20. Davidson, D. W. Ecological stoichiometry of ants in a New World rain forest. *Oecologia* **142**, 221-
231 (2004).

21. Gil, R., et al. The genome sequence of *Blochmannia floridanus*: comparative analysis of reduced
genomes. *Proc. Natl. Acad. Sci. USA* **100**, 9388-9393 (2003).

22. de Souza, D. J., Bézier, A., Depoix, D., Drezen, J. M. & Lenoir, A. *Blochmannia* endosymbionts
improve colony growth and immune defence in the ant *Camponotus fellah*. *BMC Microbiol.* **9**, 29-37
(2009).

23. Béchade, B., et al. Physiological and evolutionary contexts of a new symbiotic species from the
nitrogen-recycling gut community of turtle ants. *ISME J.* **17**, 1751-1764 (2023).

24. Csata, E. & Dussutour, A. Nutrient regulation in ants (Hymenoptera: Formicidae): a review.
Myrmecological News **29**, 111-124 (2019).

25. Feldhaar, H., et al. Nutritional upgrading for omnivorous carpenter ants by the endosymbiont
Blochmannia. *BMC Biol.* **5**, 48-59 (2007).

26. Ward, P. S. *Taxonomy, phylogeny and biogeography of the ant genus Tetraponera*
(Hymenoptera:Formicidae) in the Oriental and Australian regions. Legare Street Press (2001).

27. Ward, P. S. The ant genus *Tetraponera* in the Afrotropical region: synopsis of species groups and
revision of the *T. ambigua*-group (Hymenoptera: Formicidae). In: *Myrmecologische Nachrichten*).
antbase.org (2006).

28. Chong, K. F. & Lee, C. Y. Inter- and intraspecific aggression in the invasive longlegged ant
(Hymenoptera: Formicidae). *J. Econ. Entomol.* **103**, 1775-1783 (2010).

29. Wheeler, W. M. & Bailey, I. W. The feeding habits of Pseudomyrmine and other ants. *Transactions*
of the American Philosophical Society **22**, 235-279 (1920).

30. Klein, R. W., Kovac, D., Schellerich, A. & Maschwitz, U. Mealybug-carrying by swarming queens
of a southeast asian bamboo-inhabiting ant. *Naturwissenschaften* **79**, 422-423 (1992).
31. Billen, J. & Buschinger, A. Morphology and ultrastructure of a specialized bacterial pouch in the
digestive tract of *Tetraponera* ants (Formicidae, Pseudomyrmecinae). *Arthropod Struct. Dev.* **29**, 259-
266 (2000).
32. Salem, H., et al. Drastic genome reduction in an herbivore's pectinolytic symbiont. *Cell* **171**, 1520-
1531 (2017).
33. Hosokawa, T., Imanishi, M., Koga, R. & Fukatsu, T. Diversity and evolution of bacterial symbionts
in the gut symbiotic organ of jewel stinkbugs (Hemiptera: Scutelleridae). *Appl. Entomol. Zool.* **54**,
359-367 (2019).
34. Cook, S. C. & Davidson, D. W. Nutritional and functional biology of exudate-feeding ants. *Entomol.*
Exp. Appl. **118**, 1-10 (2006).
35. Ward, P. S. Phylogenetic analysis of Pseudomyrmecine ants associated with domatia-bearing plants.
In: *Ant-Plant Interactions* (eds Huxley CR, Cutler DF). Oxford University Press (1991).
36. Buschinger, A., Klein, R. W. & Maschwitz, U. Colony structure of a bamboo-dwelling *Tetraponera*
sp. (Hymenoptera: Formicidae: Pseudomyrmecinae) from Malaysia. *Ins. Soc.* **41**, 29-41 (1994).
37. Braekman, J. C., et al. *Tetraponerine*-8, an alkaloidal contact poison in a Neoguinean
Pseudomyrmecine ant, *Tetraponera* sp. *Zeitschrift für Naturforschung C* **42**, 627-630 (1987).
38. Midha, S., Rigden, D. J., Siozios, S., Hurst, G. D. D. & Jackson, A. P. *Bodo saltans* (Kinetoplastida)
is dependent on a novel *Paracaedibacter*-like endosymbiont that possesses multiple putative toxin-
antitoxin systems. *ISME J.* **15**, 1680-1694 (2021).
39. George, E. E., et al. Highly reduced genomes of protist endosymbionts show evolutionary
convergence. *Curr. Biol.* **30**, 925-933 (2020).
40. Weinstock, O., Weinstock, O., Sella, C., Chipman, D. M. & Barak, Z. Properties of subcloned subunits
of bacterial acetohydroxy acid synthases. *J Bacteriol.* **174**, 5560-5566 (1992).
41. Dittmer, J., Bredon, M., Moumen, B., Raimond, M., Grève, P. & Bouchon, D. The terrestrial isopod
symbiont 'Candidatus Hepatincola porcellionum' is a potential nutrient scavenger related to
Holosporales symbionts of protists. *ISME Commun.* **3**, 18-30 (2023).
42. Giovannini, M., Petroni, G. & Castelli, M. Novel evolutionary insights on the interactions of the
Holosporales (Alphaproteobacteria) with eukaryotic hosts from comparative genomics. *Environ.*
Microbiol. **26**, e16562 (2024).
43. Sauer, C., Dudaczek, D., Hölldobler, B. & Gross, R. Tissue localization of the endosymbiotic
bacterium "*Candidatus* Blochmannia floridanus" in adults and larvae of the carpenter ant *Camponotus*
floridanus. *Appl. Environ. Microbiol.* **68**, 4187-4193 (2002).
44. Ramalho, M. O., Vieira, A. S., Pereira, M. C., Moreau, C. S. & Bueno, O. C. Transovarian
transmission of *Blochmannia* and *Wolbachia* endosymbionts in the Neotropical weaver ant
Camponotus textor (Hymenoptera, Formicidae). *Curr. Microbiol.* **75**, 866-873 (2018).
45. Kwong, W. K. & Moran, N. A. Gut microbial communities of social bees. *Nat. Rev. Microbiol.* **14**,
374-384 (2016).
46. Moran, N. A., Hansen, A. K., Powell, J. E. & Sabree, Z. L. Distinctive gut microbiota of honey bees
assessed using deep sampling from individual worker bees. *PLoS One* **7**, e36393 (2012).
47. Martinson, V. G., Danforth, B. N., Minckley, R. L., Rueppell, O., Tingek, S. & Moran, N. A. A simple
and distinctive microbiota associated with honey bees and bumble bees. *Mol. Ecol.* **20**, 619-628
(2011).

48. Flynn, P. J., D'Amelio, C. L., Sanders, J. G., Russell, J. A. & Moreau, C. S. Localization of bacterial
communities within gut compartments across *Cephalotes* turtle ants. *Appl. Environ. Microbiol.* **87**,
e02803-02820 (2021).

49. Łukasik, P., et al. The structured diversity of specialized gut symbionts of the New World army ants.
Mol. Ecol. **26**, 3808-3825 (2017).

50. Li, H., et al. Convergent evolution of complex structures for ant-bacterial defensive symbiosis in
fungus-farming ants. *Proc. Natl. Acad. Sci. USA* **115**, 10720-10725 (2018).

51. Oishi, S., Moriyama, M., Mizutani, M., Futahashi, R. & Fukatsu, T. Regulation and remodeling of
microbial symbiosis in insect metamorphosis. *Proc. Natl. Acad. Sci. USA* **120**, e2304879120 (2023).

52. Oguchi, K., Harumoto, T., Katsuno, T., Matsuura, Y., Chiyoda, S. & Fukatsu, T. Intracellularity,
extracellularity, and squeezing in the symbiotic organ underpin nurturing and functioning of bacterial
symbiont in leaf beetles. *iScience* **27**, 109731-109744 (2024).

53. Fronk, D. C. & Sachs, J. L. Symbiotic organs: the nexus of host-microbe evolution. *Trends Ecol. Evol.*
37, 599-610 (2022).

54. González Porras, M. Á., et al. Extracellular symbiont colonizes insect during embryo development.
ISME Commun. **4**, ycae005 (2024).

55. Nalepa, C. A. Origin of termite eusociality: trophallaxis integrates the social, nutritional, and
microbial environments. *Ecol. Entomol.* **40**, 323-335 (2015).

56. Lanan, M. C., Rodrigues, P. A. R., Agellon, A., Jansma, P. & Wheeler, D. E. A bacterial filter protects
and structures the gut microbiome of an insect. *ISME J.* **10**, 1866-1876 (2016).

57. Powell, J. E., Martinson, V. G., Urban-Mead, K., Moran, N. A. & Goodrich-Blair, H. Routes of
acquisition of the gut microbiota of the honey bee *Apis mellifera*. *Appl. Environ. Microbiol.* **80**, 7378-
7387 (2014).

58. Van Borm, S., Buschinger, A., Boomsma, J. J. & Billen, J. *Tetraponera* ants have gut symbionts
related to nitrogen-fixing root-nodule bacteria. *Proc. R. Soc. Lond. B.* **269**, 2023-2027 (2002).

59. Stoll, S., Gadau, J., Gross, R. & Feldhaar, H. Bacterial microbiota associated with ants of the genus
Tetraponera. *Biol. J. Linn. Soc.*, 399–412 (2007).

60. Salem, H. & Kaltenpoth, M. Beetle–bacterial symbioses: endless forms most functional. *Annu. Rev.*
Entomol. **67**, 201-219 (2022).

61. Salem, H., et al. Symbiont digestive range reflects host plant breadth in herbivorous beetles. *Curr.*
Biol. **30**, 2875-2886 (2020).

62. Brune, A. & Dietrich, C. The gut microbiota of termites: digesting the diversity in the light of ecology
and evolution. *Annu. Rev. Microbiol.* **69**, 145-166 (2015).

63. Arora, J., et al. The functional evolution of termite gut microbiota. *Microbiome* **10**, 78-100 (2022).

64. Engel, P. & Moran, N. A. The gut microbiota of insects - diversity in structure and function. *FEMS*
Microbiol. Rev. **37**, 699-735 (2013).

65. Pais, I. S., Valente, R. S., Sporniak, M. & Teixeira, L. *Drosophila melanogaster* establishes a species-
specific mutualistic interaction with stable gut-colonizing bacteria. *PLoS Biol.* **16**, e2005710 (2018).

66. Ratnasingham, S. & Hebert, P. D. N. BOLD : the barcode of life data system. *Mol. Ecol. Notes* **7**,
355-364 (2007).

67. Rozewicki, J., Li, S., Amada, K. M., Standley, D. M. & Katoh, K. MAFFT-DASH: integrated protein
sequence and structural alignment. *Nucleic Acids Res.* **47**, W5-W10 (2019).

68. Talavera, G. & Castresana, J. Improvement of phylogenies after removing divergent and ambiguously
aligned blocks from protein sequence alignments. *Syst. Biol.* **56**, 564-577 (2007).

69. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

70. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display
and annotation. *Nucleic Acids Res.* **49**, W293-W296 (2021).

71. Puillandre, N., Lambert, A., Brouillet, S. & Achaz, G. ABGD, Automatic Barcode Gap Discovery for
primary species delimitation. *Mol. Ecol.* **21**, 1864-1877 (2012).

72. Tamura, K., Stecher, G., Kumar, S. & Battistuzzi, F. U. MEGA11: molecular evolutionary genetics
analysis version 11. *Mol. Biol. Evol.* **38**, 3022-3027 (2021).

73. Thayanukul, P., Zang, K., Janhom, T., Kurisu, F., Kasuga, I. & Furumai, H. Concentration-dependent
response of estrone-degrading bacterial community in activated sludge analyzed by
microautoradiography-fluorescence *in situ* hybridization. *Water Res.* **44**, 4878-4887 (2010).

74. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics* **27**, 2957-2963 (2011).

75. Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp.
iMeta **2**, e107 (2023).

76. Valdivia, C., et al. Microbial symbionts are shared between ants and their associated beetles. *Environ.*
Microbiol. **25**, 3466-3483 (2023).

77. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool
for metagenomics. *PeerJ* **4**, e2584 (2016).

78. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460-
2461 (2010).

79. Salter, S. J., et al. Reagent and laboratory contamination can critically impact sequence-based
microbiome analyses. *BMC Biol.* **12**, 87-99 (2014).

80. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**, 2114-2120 (2014).

81. Peng, Y., Leung, H. C., Yiu, S. M. & Chin, F. Y. IDBA-UD: a de novo assembler for single-cell and
metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**, 1420-1428 (2012).

82. Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D. & Gurevich, A. Versatile genome assembly
evaluation with QUAST-LG. *Bioinformatics* **34**, i142-i150 (2018).

83. Chen, I. M. A., et al. The IMG/M data management and analysis system v.7: content updates and new
features. *Nucleic Acids Res.* **51**, D723-D732 (2023).

84. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754-1760 (2009).

85. Kumar, S., Jones, M., Koutsovoulos, G., Clarke, M. & Blaxter, M. Blobology: exploring raw genome
data for contaminants, symbionts and parasites using taxon-annotated GC-coverage plots. *Front*
Genet **4**, 237-249 (2013).

86. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-resolved
metagenomic data analysis. *Microbiome* **6**, 158-171 (2018).

87. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.*
25, 1043-1055 (2015).

88. Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify genomes
with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925-1927 (2019).

89. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome comparisons

968 dramatically improves orthogroup inference accuracy. *Genome Biology* **16**, 157-171 (2015).

969 90. Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. ProtTest 3: fast selection of best-fit models of

970 protein evolution. *Bioinformatics* **27**, 1164-1165 (2011).