Title: Premature vision drives aberrant development of
response properties in primary visual cortex

Abbreviated Title: Premature vision drives aberrant development

Authors: Sophie V. Griswold (0000-0003-2340-9602)[1,2], Stephen D. Van Hooser
(0000-0002-1112-5832)[1,2,3]
1: Department of Biology
2: Volen Center for Complex Systems
3: Sloan-Swartz Center for Theoretical Neurobiology
Brandeis University, Waltham, MA 02454 USA
11
12 **Corresponding author:**
13 Stephen D. Van Hooser, vanhooser@brandeis.edu
14 Brandeis University, 415 South St. MS008, Waltham, MA 02454 USA

16 **Conflicts of interest:** The authors declare no competing financial interests.
17
18 **Acknowledgements**: This work was funded by NIH EY022122 (SDV). We thank Jiwon
19 Sun for help with histology and some experiments and members of the Van Hooser lab
20 for comments.
21
22 **Contributions:** SVG performed experiments. SVG and SDV designed the
23 experiments, analyzed the data, and wrote the paper.
24
25 **Keywords:** motion, temporal frequency, direction selectivity, striate cortex, area 17

ABSTRACT:

Development of the mammalian visual system is thought to proceed in two stages. In the first stage, before birth in primates and before eye opening in altricial mammals, spontaneous activity generated by the retina and cortex shapes visual brain circuits in an activity-dependent but experience-independent manner. In the second stage, visual activity generated by sensory experience refines receptive fields. Here we investigated the consequences of altering this sequence of events by prematurely opening one or both eyes of ferrets and examining visual receptive fields in monocular cortex after the closure of the critical period for ocular dominance plasticity. We observed that many cells in animals with prematurely-opened eyes exhibited low-pass temporal frequency tuning and increased temporal frequency bandwidths, and these cells showed slightly increased orientation and direction selectivity index values. Spontaneous activity was greatly elevated in both hemispheres following the premature opening of one or both eyes, suggesting a global change in circuit excitability that was not restricted to cells that viewed the world through the prematurely opened eye. No major changes were noted in spatial frequency tuning. These results suggest that premature visual experience alters circuit excitability and visual receptive fields, in particular with respect to temporal processing. We speculate that closed lids in altricial mammals serve to prevent visual experience until circuits are initially established and are ready to be refined by visual experience.

INTRODUCTION

The proper development of visual cortical circuits depends on both experience-independent (Katz and Shatz 1996, Wong 1999) and experience-dependent (Huberman et al 2008, Wang et al., 2010) processes. Classically, these mechanisms have been divided into two sequential stages: an early stage, before the onset of visual experience, in which molecular cues and spontaneous activity in retina and cortex guide initial circuit formation (Katz and Shatz 1996, Wong 1999), followed by a period in which activity driven by visual experience refines circuit function (Huberman et al 2008, Li et al,. 2006, Wang et al., 2010). The sequential nature of these two phases presumably ensures a developmental trajectory in which circuits achieve a state of maturation that provides proper support for the onset of experience-driven mechanisms that finalize the construction process.

Given the sequential nature of visual development, it is interesting to ask if there are any positive or negative consequences of violating this sequence by introducing patterned visual stimulation at earlier time points. Are there significant changes that cannot be reversed even with subsequent visual experience? These questions are especially important to address in light of the rising population of human infants born very prematurely (Crump et al., 2019), who receive visual stimulation during what is typically a presensory period (Colonnese et al. 2010).

To address these questions, we have examined the impact of premature visual experience on receptive fields in visual cortex of the ferret. Ferrets undergo a long postnatal developmental period and have a history of foundational developmental studies in vision. Ferrets are born with their eyes closed – a condition that is maintained for 30-35 days (Issa et al., 1999) – making it possible to introduce inappropriately early visual experience by opening the lids with forceps,

without introducing factors related to immature birth. We opened zero, one, or both eyes early,

and evaluated receptive fields in primary visual cortex electrophysiologically after the closure of

the critical periods for direction selectivity and ocular dominance. We studied monocular

receptive fields only, so that we could be sure of the developmental condition of the eye,

necessarily the contralateral eye, that was driving the receptive field properties we observed.

We observed substantial changes in temporal frequency tuning in monocular cells that

observed the world through an early-opened eye. These cells, on average, exhibited strikingly

broad tuning for temporal frequency with markedly enhanced responses at low temporal

frequencies. These major alterations in visual selectivity were not present in all receptive field

properties, as orientation and direction tuning in these cells were only slightly altered. By

contrast, we observed substantial changes in spontaneous firing rates in both hemispheres after

either or both eyes were opened early, suggesting that premature vision has a more global impact

on the establishment of a firing rate set point in cortex (Turrigiano and Nelson, 2004; Hengen et

al. 2013).

In all, these results suggest that premature visual experience causes lasting changes in

temporal processing and in the baseline tone of cortical circuits. The changes in temporal

frequency tuning are likely to have an impact on motion perception. To perceive a moving

object, an animal must have knowledge of the object's direction and speed. The ratio of the

temporal frequency of a stimulus to its spatial frequency is its speed, so temporal frequency

tuning provides critical information about stimulus speed. Human babies that are born very

prematurely are a heterogenous group with a wide variety of visual deficits (Kozeis 2010, Dutton

2013, Sakki et al., 2018), but even when one excludes cases of known brain or retinal damage,

very premature babies exhibit increased thresholds for motion perception later in life (Taylor et

al., 2009, Hou et al., 2011). Our results in ferret raise the possibility that inappropriately early

visual experience could be an important factor that contributes to lasting motion processing

deficits in humans born very prematurely.

Part of this work appeared as the PhD thesis of SVG (Griswold, 2024).

METHODS

All experimental procedures were approved by the Brandeis University's Institutional Animal

Care and Use Committee (IACUC) and performed in compliance with National Institutes of

Health guidelines.

Animal source and housing: Ferrets (*Mustelo putorius furo*) were obtained from Marshall Bio-

Resources. Litters of 4 or more kits arrived with a jill between postnatal days (P) 12-21. Animals

were housed in a room with timed lights (12 h on, 12 h off) in a custom stainless-steel cage (60

cm × 60 cm × 35 cm) with a hammock and small toys. For the entire study, a total of 27 female

ferrets were used and all experimental procedures were carried out between postnatal days P55–

68. Female ferrets were used because housing mature male ferrets in the same room with mature

female ferrets causes stress to the female ferrets.

Facilitating premature visual experience: We divided animals into three experimental groups.

The first group of animals had one of their eyes gently opened with forceps at P25, while the

other eye was left closed to open naturally, and time of natural eye opening was noted. We used

the label early Eye Opening One (henceforth EO1) to indicate neurons in the monocular visual

cortex that viewed the world through the eye that was opened early, and early Eye Opening

Ipsilateral (henceforth EOIPS) to indicate neurons in the monocular visual cortex that viewed the

world through the eye that opened on time. In the second group of animals, both eyes were

opened at P25, and neurons from the monocular visual cortex that viewed the world through

either of these eyes were labeled early Eye Opening 2 (henceforth EO2). Finally, the third group

of animals were two eye controls, and were allowed to open both eyes normally, with time of

natural eye opening noted. Ferrets frequently open one eye up to a day before the other, and any

disparity in the time of opening between the eyes was noted. Ferret kits are actively kept in their

nests by their mothers prior to eye opening, so in order to ensure that animals with early-opened

eyes actually had patterned visual experience, animals were brought to the lab for 2 hours a day

for 4 consecutive days beginning at P25. These ferret kits were placed in a rat cage atop a

heating pad, and gently manipulated to maintain wakefulness during this natural, unguided

viewing. We studied 17 animals that had one eye prematurely opened, 4 animals that had both

eyes were opened, and 11 control.

Surgical preparation for terminal physiology experiments P55-68: The ferret was sedated with

ketamine (20 mg kg−1 im). Atropine (0.16–0.8 mg kg−1 im) and dexamethasone (0.5 mg kg−1

im) were administered to reduce bronchial and salivary secretion and to reduce inflammation,

respectively. The animal was next anesthetized with a mixture of isoflurane, oxygen, and nitrous

oxide through a mask and a tracheostomy was performed. The animal was then ventilated with

1.5–3% isoflurane in a 2:1 mixture of nitrous oxide and oxygen. A cannula was inserted into the

intraperitoneal (ip) cavity for delivery of neuromuscular blockers and Ringer solution (3 ml kg−1

h−1), and the animal was inserted in a custom stereotaxic frame that did not obstruct vision. All

wound margins were infused with bupivacaine. A suture was made in the scalp and the skin

resected over the posterior half of the skull. A small opening in the cranium was drilled over V1

with a dental drill (Medidenta). The dura covering the brain at the craniotomy site was removed

over a 1 mm × 1 mm area to allow for electrode placement. Craniotomies were made over V1 of

each hemisphere. The eyelids were sutured open, and contact lenses were placed on the eyes to

prevent corneal damage. Before visual stimulation commenced, the ferret was paralyzed with the

neuromuscular blocker gallamine triethiodide (10–30 mg kg−1 h−1) through the ip cannula to

suppress spontaneous eye movements, and the nitrous oxide oxygen mixture was adjusted to 1:1.

The animal's ECG was continuously monitored to ensure adequate anesthesia, and the

percentage of isoflurane was increased if the ECG indicated any distress. Body temperature was

maintained at 37 °C. At the conclusion of the experiment, the animal was killed with an overdose

of sodium pentobarbitol (200 mg/kg, IP) and transcardially perfused to retrieve the brain for

histology.

Electrophysiological recordings: 32 channel electrodes (Plexon S probes, 50um inter-tetrode

spacing; 50um intra-tetrode spacing) were used for all recordings. The signal was amplified

using the RHD2000 amplifying/ digitizing chip and USB interface board (Intan Technologies).

Stimulus timing information was acquired using a Micro1401 acquisition board and Spike2

software (Cambridge Electronic Design). Spike sorting was performed off-line using JRClust

running in Matlab (Jun et al., 2017). An electrode was inserted into the brain using a Sutter

Instruments MP-285 manipulator. To reduce sampling bias, we recorded from any site that had a

signal-to- noise ratio sufficient for isolation and had a response that appeared to be modulated by

the presentation of drifting gratings. Data are reported from all units that are responsive enough

to be included in analysis (see below). After finishing the recording at one site, the electrode was

lowered at least 640 microns before attempting to identify a suitable subsequent recording site.

The experiment was concluded when successful recordings were made from one or both

hemispheres.

Locating Monocular Neurons: After placing the electrode at a minimum depth of 200 μm in the

brain, entering the brain at an angle of 30-45 degrees, the search for monocular neurons

commenced. The electrode was driven into the brain until monocular neurons were discovered.

A population of neurons was deemed monocular if their receptive fields were in the periphery of

the eye contralateral to the recording hemisphere. Monocularity was ensured by determining

whether movement in the central visual and ipsilateral visual fields evoked a response; any such

responses indicated binocularity. If no purely monocular cells were found in a given penetration,

the electrode was removed and placed and another penetration was made elsewhere in the brain.

Locating the Optic Disk: An indirect ophthalmoscope (Heine Omega 600) and handheld 78 or

90D lens (Volk) were used to locate the optic disk and determine the orientation of the eye. This

was necessary as the orientation of the eye is necessary for determining the true eccentricity of

receptive fields. When one investigator had the optic disk in the center of view of the indirect

ophthalmoscope, another investigator introduced a wooden rod into the line of sight and moved

it until the end of the rod was centered on the image of the optic disk (also the center of view).

The position of the end of the rod was then measured in X, Y, and Z relative to the animal's eye

to make the optic disk direction.

Visual Stimuli: Visual stimuli were created in MATLAB (MathWorks) using the Psychophysics

Toolbox (Brainard 1997, Pelli 1997) and displayed on a 21-inch flat face CRT monitor (GDM-

520, Sony) with a resolution of 800×600 and a refresh rate of 100 Hz. We manually mapped

receptive fields by displaying circular patches of drifting sinusoidal gratings at different positions

and moving the monitor to accommodate different eccentricities while listening to the responses

on a loudspeaker.

Drifting grating stimuli were full-field, drifting sinusoidal gratings (4 s duration; 3.5 s

interstimulus interval) presented pseudorandomly, with direction of motion (in steps of 45°) in

either of the two directions orthogonal to the axis of orientation. We ran two sets of stimuli

where multiple grating parameters were covaried in order to sample a wide variety of receptive

field properties in our multi-neuron recordings. In the first set, orientation (angles 0°/horizontal,

45°, 90°/vertical, 135°) was co-varied with stimuli consisting of drifting sinusoidal gratings at 8

different spatial frequencies (0.04, 0.08, 0.16, 0.24, 0.32, 0.64, 0.90, and 1.25), and 6 different

contrasts (0.04, 0.08, 0.16, 0.32, 0.64, and 1), while temporal frequency was held constant at 4

Hz. In this first set of stimuli, gratings drifted back and forth every 4 cycles; for example,

horizontal stimuli drifted upward and then downward. In a second set of stimuli, direction (0°/up,

45°, 90°/right, 135°, 180°/down, 225°, 270°/left, 315°) was co-varied with stimuli of 7 different

temporal frequencies (0.5, 1, 2, 4, 8, 16, and 32 Hz) at 100% contrast at 100% contrast and a

fixed spatial frequency of 0.1 cycles/° visual angle, In this second set of stimuli, gratings drifted

in a single direction and did not move back and forth. Each stimulus set had a blank, control

stimulus that was the same duration as the other stimuli where the screen did not change. The

first set of stimuli was repeated 5 times, and the second set 7 times.

Data analysis: Responses were determined by examining either the mean response during the

entire stimulus or by taking the F1 component of the response. The same analysis (mean, or F1 at

the temporal frequency of the stimulus of interest) was performed for the closest blank (control)

stimulus and the response to the control stimulus was subtracted to yield the stimulus-driven

response. For each tuning curve for orientation, direction, and spatial frequency analyses, we

examined whether the mean or the F1 response was higher, and used those responses for analysis

and fits. For temporal frequency analysis, we only used mean responses because we found a

profound response suppression in many cells.

Orientation selectivity was examined in the first set of stimuli at the preferred spatial

frequency for each cell. Circular variance was calculated in orientation space (Ringach et al.,

2002, Mazurek et al., 2014) with the equation $V = 1 - |R|$, where R is the resultant calculated as:

$$R = \frac{\sum_k r_k e^{i2\theta_k}}{\sum_k r_k}$$

Where rk was the mean spike rate in response to a grating drifting with angle θk.

Tuning curves from cells that showed significant variation across orientation stimuli and

blank (ANOVA test, (Mazurek et al., 2014)) were fit with a double gaussian to obtain the angle

preference.

Direction selectivity was examined in the second set of stimuli for the preferred temporal

frequency of each cell (unless noted). Circular variance was calculated in direction space with

the following equation:

$$1 - DirCirVar = \left| \frac{\sum_k R(\theta_k)\exp(i\theta_k)}{\sum_k R(\theta_k)} \right|$$

Preferred spatial frequency was determined by responses to the first stimulus battery at

the preferred orientation at 100% contrast. Preferred temporal frequency was determined by

responses to the second stimulus battery at the preferred direction. Each was fitted with a model

due to Movshon et al. 2005: $R(f) = k * \exp\left(-\left(\frac{f}{fc}\right)^2\right)/[1 + (\frac{fh}{f})^\beta]$.

Low frequency cut-offs were determined by following the fit from its peak to lower frequencies

until the first frequency where the response dropped to half its maximum; high frequency cut-

offs were determined by the same procedure, except that the fit was followed from the peak

towards higher frequencies (Moore et al. 2005, Heimel et al., 2005). If the response never

dropped below half its maximum, then the low frequency cut off was said to be 0 or the high

frequency cut off was said to be infinity. Temporal frequency tuning bandwidth was defined as

log2(H50/L50).

Contrast responses were fit with Naka-Rushton functions as described in (Peirce 2007)

and sensitivity determined by the reciprocal of the contrast for which the response exceeds 5

standard deviations of the response to a blank screen. The Naka-Rushton function is defined as:

$$R = R_{max} \frac{c^n}{c^n_{50} + c^n} + B$$

Where Rmax is the maximum response of the neuron, c50 is the contrast at which the

response is halfway between baseline and maximum, c is the contrast of the stimulus presented

to the neuron, and B is the baseline response of the neuron. n is a factor simply referred to as the

exponent.

Statistics: Index values were examined with linear mixed effects models in Matlab (function:

fitlme), where treatment condition (control, EO1, EO2, EOIPS) were fixed effects and animal

identities were random effects.

Immunohistochemistry: Upon completion of experiments, an electrode coated in the fluorescent

dye DiI (DiCarlo et al., 1996) was inserted at the location and depth and left in place for 20

minutes. Animals were then transcardially perfused and the brain was placed in 4%

paraformaldehyde in 0.1M PBS at 4°C for 24 hours and then moved to 10% sucrose in PBS for

24-48 h. This was followed by placement in 30% sucrose in PBS at 4°C until sectioning. The

brain was sectioned sagittally into 100 μm sections using a sliding microtome (Leica SM2010R).

We washed sections in 0.1M PBS 3×5 min and permeabilized in 0.3% Triton-X 100 diluted in

PBS for 2 hours at room temperature on a shaker. Then slices were incubated in fluorophore-

conjugated anti-NeuN antibody (Alexa Fluor 488 Rabbit anti NeuN, Millipore ABN78A4) at a

1:300 dilution overnight (>12 hours) at room temperature on a shaker. Sections were then

washed 3×5 min in PBS and mounted on slides and allowed to air dry. Slides were then cover-

slipped with Fluoromount-G media (Electron Microscopy Sciences, Ft. Washington, PA) and

edges were sealed using nail polish. Histological sections were viewed using a fluorescent

microscope (Keyence BX-Z 710) and electrode tracks were reconstructed using DiI dye traces.

RESULTS

Our primary goal was to explore the influence of premature vision on the parameters of

receptive fields in V1. We reared ferrets under three conditions in order to examine the impact of

premature vision through one or both eyes in comparison with controls (**Figure 1**): in one group,

we opened one eye early; in a second group, we opened both eyes early; and a third, control

group were allowed to open their eyes naturally. Animals in both of the premature eye opening

groups experienced early vision beginning on postnatal day 25, a developmental timepoint when

retinal waves are still occurring (Meister et al., 1991, Wong et al., 1993), yet V1 is also

responsive to visual input through the closed lids (Krug et al., 2001, Akerman et al., 2002). This

ensured interaction between spontaneous, endogenously generated activity and patterned visual

input. We hypothesized that this interaction may drive aberrant development of cortical receptive

fields, as it is known that visual input is capable of initiating retinal waves through the closed lids

(Tiriac et al., 2018). Following premature eye opening, animals were brought to the lab for 2

hours a day of unguided visual exposure over 4 days, for a total of 8 hours of visual exposure.

Visual exposure consisted of gentle handling to prevent the young kits from sleeping thus

ensuring a sufficient amount of visual experience through the prematurely opened eye(s).

Animals received no other intervention between the end of visual stimulus training on P28 and

terminal electrophysiology experiments which occurred between P55-68.

We made multichannel recordings at a developmental timepoint when past studies have

shown that V1 receptive fields are largely mature. By postnatal day 42, contrast sensitivity (Li et

al., 2006), orientation (Chapman and Stryker 1993), direction, temporal frequency, and spatial

frequency (Li et al., 2006) tuning have reached adult-like levels and ocular dominance plasticity

is greatly reduced (Issa 1999). Thus, if premature vision causes permanent changes to receptive

field properties, we ought to be able to see them at P55-68.

We recorded responses to long batteries of visual stimulation with sinusoidal gratings

where multiple parameters were co-varied in order to assess a wide variety of possible receptive

field differences between control animals and animals that had their eyes opened prematurely. In

this study, all recordings were performed in the monocular visual cortex in order to simplify the

interpretation of these changes without needing to consider interactions between the two eyes.

We labeled cells as E01 (monocular neurons contralateral to a single early-opened eye), EOIPS

(monocular neurons ipsilateral to a single early-opened eye), E02 (monocular neurons

contralateral to an early opened eye where both eyes were opened early), and control (monocular

neurons in control animals). E01 and E02 cells observed the world through early-opened eyes,

while EOIPS neurons observed the world through eyes that opened on time but were ipsilateral

to early-opened eyes.

Orientation and direction tuning index values are slightly increased by premature

experience

Cells in all animals exhibited some degree of orientation and direction selectivity.

Orientation selectivity was assessed for stimuli that moved back and forth in two opposite

directions at each cell's preferred spatial frequency. Direction selectivity was assessed at the

neuron's optimal temporal frequency and a spatial frequency of 0.1. Example responses for each

group are shown in Fig. 2A-H. Control animals exhibited a baseline average orientation index

value (1-CV) of 0.42, and this value was not significantly elevated in EOIPS neurons (0.45,

$p < 0.54$, Linear Mixed Effects Model). However, orientation selectivity was slightly elevated in

E01 neurons (0.52, $p < 0.024$) and E02 neurons (0.54, $p < 0.016$, LMEM). Direction selectivity as

assessed by 1-DCV was slightly reduced in EOIPS neurons (0.24, $p < 0.027$, LMEM) as

compared to control values (0.29), and 1-DCV values were not significantly different in E01

neurons (0.29, p< 0.296) or E02 neurons (0.33, p<0.169, LMEM) . The overall distribution of

direction preferences in 45° bins did not significantly differ across the groups (chi-square test, p

= 0.11), nor did the distribution of orientation angle preferences in 45° bins (chi-square test, 5.3

x 10-2). Therefore, we observed only minor long-term changes in orientation and direction

tuning between animals with premature vision and control animals.

Temporal frequency tuning is aberrant in animals with premature visual experience

We were struck by the unusual temporal frequency tuning curves that we observed in

animals whose eyes were opened prematurely. Many EO1 cells exhibited strong responses at the

lowest temporal frequency tested (Fig. 3A), and we observed a trend for this in EO2 cells as well

(Fig. 3B). Further, several cells also showed evidence of a suppression of the response below

baseline. Tuning curves from E0IPS cells (Fig. 3C) superficially looked more like curves from

control animals (Fig. 3D), although quantitative analyses uncovered small differences. Most cell

types exhibited similar peak temporal frequency values (Fig. 3E), except that EOIPS cells

exhibited slightly higher temporal frequency preferences (p<0. 021, LMEM). To quantify the

response to the lowest temporal frequency tested, we calculated a low pass index, defined as the

ratio of the response to the lowest temporal frequency tested to the largest temporal frequency

response observed (Fig. 3F). Temporal frequency low pass index values of control cells were

0.20 on average, while EO1 cells showed low pass index values that averaged 0.35 (p< 0.039,

LMEM) and EO2 cells showed empirically elevated values of 0.33 that were not significantly

different from control (p< 0.142, LMEM). High pass index values, determined by calculating the

response at the highest temporal frequency tested divided by the maximum response, did not

differ across these animals (Fig. 3G).

In order to assess tuning bandwidth, we calculated low- and high-frequency cut off values

by identifying the low and high frequency values where the response dropped below half the

maximum response value. If this never occurred, the cell was considered to be low-pass (if this

never occurred on the low frequency side) or high-pass (if this never occurred on the high

frequency side). We then calculated the bandwidth between this low and high value in octaves. If

a cell was low-pass or high-pass or both, the cell was said to have infinite bandwidth. In Fig. 3H,

we show bandwidth computed from rectified responses, and saw a empirically elevated

bandwidths in E01 cells (2.33 octaves, $p< 0.07$) and E02 cells (2.33 octaves, $p<0.08$) compared

to control cells (1.95 octaves) or EOIPS cells (1.91 octaves, $p<.937$). Many cells exhibited

suppression, where the response fell below the response to a control stimulus, and we took the

absolute value of the response in calculating the absolute bandwidth so that both positive and

negative responses were considered. The median absolute bandwidth (Fig. 3I) of control cells

(4.1 octaves) was narrower than any of the cells in animals whose eyes were opened early,

including EOIPS cells (4.7 octaves, $p<0.0013$), EO1 cells (4.8 octaves, $p<0.002$), and EO2 cells

(5.1 octaves, $p<0.00015$, LMEM).

Spatial frequency tuning is relatively unaffected by premature visual experience

Spatial frequency tuning was hardly influenced by early eye opening. We assessed spatial

frequency at a cell's preferred orientation, as determined by responses to drifting gratings that

moved back and forth in opposite directions. Example tuning curves are shown in Fig 4A-D.

In linear mixed effect modeling, no significant differences were observed in peak spatial frequency preferences across these groups (control: 0.031 cpd; EOIPS: 0.045 cpd, p<0.19; EO1: 0.031 cpd, p<0.949; EO2: 0.042 cpd, p<0.362, LMEM). Similarly, median spatial frequency low pass index values were very similar, with a slightly significant change noted for EOIPS animals (control: 0.86; EOIPS: 0.76, p<0.0499; EO1: 0.85, p<0.928; EO2: 0.79, p<0.390, LMEM). We observed slightly higher spatial frequency high pass index values in EO2 cells, but effects were modest: (control: 0.02; EOIPS: 0.02, p<0.857; EO1: 0.04, p<0.090; EO2: 0.06, p<0.0323, LMEM).

We observed no significant effects with respect to spatial frequency bandwidth, either when measured with responses that were greater than 0 or measuring the absolute response (that is, interpreting responses below baseline as positive responses). Bandwidths of individual cells were often infinite, given the large preponderance of low-pass cells, and so we analyzed the data in rank order instead of the raw values. Median values for both positive-only responses and absolute responses were always above the ranks of cells with finite bandwidths, indicating that the typical cell in all groups had infinite bandwidth. Differences from control were not significant for positive-only responses (EOIPS: p<0.310, EO1: 0.622, EO2: p<0.898, LMEM) or absolute responses (EOIPS: p<0.06, EO1: p<0.597, EO2: p<0.385, LMEM).

In short, we found very little evidence of substantial changes to spatial frequency tuning in these monocularly-driven cells.

Spontaneous firing rates and response suppression are increased in animals that have had premature visual experience

Hebbian and homeostatic mechanisms are active early in development (Turrigiano and

Nelson, 2004), and may be critical for setting response gains and the background excitability of

neural circuits (Roy et al., 2018; Roy et al. 2020a). To examine whether response gains or

excitability were impacted by premature vision, we quantified maximum firing rates in response

to visual stimuli, response suppression, and background firing rates.

The peak mean responses of neurons to visual stimuli (Fig. 5A) did not differ among the

experimental groups. Control neurons exhibited average peak responses of 7.0 Hz; EOIPS

neurons did not differ significantly (6.4 Hz, $P<0.600$), nor did EO1 neurons (10.1 Hz, $p<0.518$)

or EO2 neurons (10.5 Hz, $p<0.295$).

In contrast to peak responses, response suppression below background levels was

pronounced in animals with prematurely opened eyes (Fig. 5B). We calculated all responses by

subtracting the background rate measured during control stimuli (gray screen), and took response

suppression for each cell to be the maximum amount of suppression below background rates for

the average response to any visual stimulus. Suppression was rarely observed in control cells

(average: 2.1 Hz), but was substantially elevated in EOIPS cells (7.0 Hz, $p<1.59e-9$) and slightly

elevated in EO1 cells (3.4 Hz, $p<0.022$) and EO2 cells (2.8 Hz, $p<0.035$).

Finally, background firing rates were significantly higher in animals whose eyes were

opened early (Fig. 5C). Spontaneous firing rates were low in control animals (2.6 Hz), and were

markedly increased in EOIPS hemispheres (11.1 Hz, $p<1.09e-4$), EO1 hemispheres (6.6 Hz,

$p<0.0132$) and EO2 hemispheres (6.6 Hz, $p<0.0248$). Therefore, early opening produced a long-

lasting effect on spontaneous firing rates after the critical period.

In all, early eye opening induced changes in response suppression and background firing

rates that were still evident at the time of our measurements several weeks later. We observed

these changes in both hemispheres, which indicates that the differences in direct visual drive to

visual receptive fields are not underlying these differences but instead are the result of some

more global process. These results are consistent with the idea that early eye opening induces

long-term changes to circuit-level excitability that are not corrected during the remainder of

development.

DISCUSSION

We explored the impact of premature vision on the development of a number of V1 RF

properties in ferret. We found small increases in direction and orientation tuning in cells that

viewed the world through prematurely-opened eyes. We observed profound changes in temporal

frequency tuning: cells recorded from animals that had their eyes opened prematurely exhibited

more low-pass temporal frequency tuning curves and broader temporal frequency bandwidths

compared to controls. Spatial frequency preferences were not substantially altered. Finally, cells

in either hemisphere in animals with prematurely-opened eyes had higher spontaneous firing

rates and exhibited marked suppression below this spontaneous firing rate in response to some

visual stimuli.

A novel temporal frequency tuning deficit with possible implications for motion processing

We found that ferrets with premature visual experience exhibited a pronounced

broadening of temporal frequency tuning curves and increases in the fraction of low- pass

neurons. The alterations in temporal frequency tuning identified in ferrets following premature

eye opening are strongly suggestive of a deficit in motion processing. The broadening of

temporal frequency tuning curves and increase in the proportion of low pass cells in V1 could

translate to a decrease in the animal's ability to distinguish the temporal frequency (and perhaps

the speed) of a moving visual stimulus.

Altered rearing has been shown to influence the subsequent development of motion

processing in various ways. Dark-reared cats and ferrets exhibit poor direction selectivity (Imbert

and Buisseret 1975, Li et al., 2006), but dark-reared ferrets do not show strong deficits in

temporal frequency processing. Overall responses after dark-rearing are reduced, but temporal

frequency tuning is only slightly narrowed (Li et al., 2006). Cats raised in a strobe-light

environment (mostly after eye opening) exhibited strong changes in subsequent direction

selectivity (Kennedy and Orban 1983, Humphrey and Saul 1998), but temporal frequency tuning

of these animals has not been reported in detail. Kennedy and Orban (1983) report a reduction in

high-pass velocity-tuned cells, but not an increase in low-pass cells; Humphrey and Saul (1998)

do not report temporal frequency tuning across their population but their example cell does

appear to be broadly tuned and low-pass. Temporal frequency tuning is not subject to rapid

modification shortly after eye opening, as Ritter and colleagues demonstrated that providing

ferrets that had just opened their eyes naturally with 6-9 hours of visual stimulation of a single

TF did not elicit a shift in TF tuning (Ritter et al., 2017). The same stimulation causes a dramatic

increase in direction selectivity (Ritter et al., 2017) that normally develops in the first week to 10

days after eye opening and requires visual experience (Li et al., 2006, Li et al., 2008, Van Hooser

et al., 2012). The only manipulation we know of in the literature that has been shown to modify

temporal frequency properties of developing animals is prey capture: mice that learned to hunt

crickets in the critical period for ocular dominance plasticity exhibit lower low-pass index values

(Bissen et al. 2025).

In sum, the deficits in temporal frequency tuning we observed are unlike those reported

previously. Premature vision produces permanent alteration of temporal frequency tuning curves,

while dark-rearing does not and comparable results from strobe rearing are not available.

Mechanisms underlying altered receptive fields

One way that premature patterned vision could alter the development of the brain is by

altering the character of retinal waves. Low resolution visual input before natural eye opening is

capable of driving activity in the visual circuit (Huttenlocher 1967, Chapman and Stryker 1993,

Krug et al., 2001, Akerman et al., 2002, Akerman et al., 2004), and evidence shows that this

early form of low resolution visual experience plays a role in the development of ON-OFF

segregation in dLGN (Akerman et al., 2002) and the initiation of retinal waves (Tiriac et al.,

2018). Propagation of high luminance, high contrast, and high spatial frequency input through

the developing retina may alter or disrupt the typical retinal wave formation and propagation,

and thus may have deleterious effects on developmental processes driven by endogenous

activity. Alternatively, high contrast signals may not interrupt retinal waves but may contribute

independent activity that, through experience-dependent plasticity mechanisms, produces the

alterations in receptive field properties that we observed.

Several other basic mechanistic questions remain unanswered. It is unclear where in the

visual circuit cascade these deficits first arise. Does the lateral geniculate nucleus or retina

exhibit altered temporal frequency tuning? Is the influence of the patterned visual stimulation

instructive, so that if one provided premature stimulation with only certain temporal frequencies,

one would see selectivity for those temporal frequencies, or would tuning always be broad?

Future studies will be needed to address these questions.

The broad tuning in temporal frequency and elevated spontaneous firing rates that we

observed could reflect a deficit in cortical inhibition in animals that experienced premature

vision. Yet selectivity was not entirely destroyed, as neurons still exhibited selectivity for

orientation and direction. Selectivity can be achieved with a combination of amplification of

appropriate excitatory signals and inhibitory suppression that is either selective or broad (Ben-

Yishai et al., 1995, Rubin et al., 2015, Somers et al., 1995, Troyer et al., 1998), so it is unclear

how the circuit has changed. The amount and timing of inhibition, whether feed-forward or

feedback, could be altered.

Alterations in ipsilateral receptive fields

One of our most surprising findings was a difference in the receptive fields and firing

properties of cells that were ipsilateral to the early opened eye. It is straightforward to imagine

how Hebbian mechanisms might influence monocular cells that are contralateral to the early-

opened eye, but one would have to imagine different and possibly more global mechanisms to

explain changes in monocular cells that view the world through the eye that opened naturally.

Homeostatic mechanisms such as synaptic scaling (Desai et al., 2008, Turrigiano et al.,

1998), and regulation of intrinsic excitability (Desai et al., 1998, Lambo and Turrigiano, 2013)

operate during the critical period for ocular dominance plasticity and serve to keep ongoing

firing rates within a cell-defined range (Hengen et al., 2013, Hengen et al., 2016). It is possible

that these homeostatic mechanisms are not yet operating in the presensory period, and that the

extra activity causes an increase in excitatory forces (whether synaptic or intrinsic) relative to

inhibitory forces.

Alternatively, opening a single eye early may still influence feed-forward activity in the

ipsilateral hemisphere. Work in ferrets and rats suggests a role for retinal waves in the

development of interhemispheric connectivity; bilateral enucleation of ferrets induces anomalies

in the distribution of callosal cells in primary visual cortex (Bock et al., 2012), and results in

callosal projections connecting topographically mismatched loci in VC in P4-6 rats (Olavarria

and Hiroi 2003). Further, retinal waves demonstrate coordinated activity between both

hemispheres in mice and ferrets (Weliky and Katz 1999, Ackman et al., 2012). Recent findings

describe a transient retina to retina connection in mice and ferret, which exists during the period

of retinal waves (Murcia-Belmonte et al., 2019). This connection provides a plausible

mechanism for the theorized inter- retinal coordination of waves thought to be necessary to drive

development of bilateral topographic maps (Adams and Horton 2003, Murcia-Belmonte et al.,

2019). Given that light is capable of initiating retinal waves (Tiriac et al., 2018), it is feasible that

exposure to light through a single, prematurely opened eye would initiate retinal waves capable

of propagating to the closed eye. Given the qualitative differences between vision through a

closed lid vs open eye, initiation and propagation of retinal waves would be both asymmetric and

mutually influential between retinas, potentially accounting for the differences in RFs observed

between EOIPS and control hemispheres.

Comparison with prior studies

There have been only a few animal studies that have sought to introduce inappropriately

early vision. Lichliter and colleagues (2018) prematurely opened the shells of bobwhite quail

eggs and provided visual experience to the chick, and found a subsequent loss of visual

selectivity to bobwhite hens vs. hens of another species (Sleigh and Lickliter 1995). Further, they

demonstrated multi-modal impacts on auditory processing after premature visual experience, in

that animals failed to eventually become responsive to cross-species hen calls. In behavioral

studies of rats whose eyes were prematurely opened 7 days early, investigators found that

opening a single eye (but not opening both eyes early) caused a deficit in pivoting locomotor

reactions on the side of the early-opened eye; that is, rats pivoted more frequently towards

stimuli viewed through the eye that opened on time (Foreman and Altaha 1991). To our

knowledge, the present work is the first to study cortical receptive fields after premature sensory

experience. All of these studies, including our own, have noted deficits after premature

experience.

Premature vision as a risk factor for visual disfunction

Very premature human babies exhibit a variety of visual deficits later in life that are not

explained by the condition of the eye, broadly termed Cortical/Cerebral Visual Impairment

(CVI) (Kozeis 2010, Dutton 2013, Sakki et al., 2018). These very premature babies are a

heterogenous population. Some of these infants have suffered clear brain injuries (Slidsborg et

al., 2012, Dutton 2013) of varies types, such as hypoxia, brain bleeding, or periventricular

leukomalacia (Gallo and Lennerstrand 1991, Hokken and Krabbendam 2023, Kozeis 2010), and

these injuries clearly have a direct impact on subsequent visual deficits that go beyond any

influence of early experience. But, studies that have excluded individuals with known brain

damage or retinal problems (or analyzed these populations separately) have also found that

extreme prematurity itself has an impact on later vision, particularly on motion perception

(Atkinson and Braddick, 2007, Benassi et al., 2018, Guzzetta et al., 2009, Hou et al., 2011,

MacKay et al., 2005) and sometimes lowered visual acuity (Jain et al. 2022). Previous

investigators have speculated that inappropriately early visual stimulation may contribute to

these deficits, as in Taylor et al (2009): "Why the dorsal stream should be particularly vulnerable

in this population is not completely understood. One possibility is that the unusually early visual

stimulation that very premature infants are exposed to may, in itself, have a differential effect on

the functional development of the dorsal and ventral streams". Our data are consistent with the

idea that inappropriately early visual stimulation contributes to deficits in motion processing. A

moving object has both a speed and direction, and temporal frequency is critical to determine an

object's speed. The major temporal frequency tuning deficits we have observed in receptive

fields are consistent with a motion deficit.

Premature vision as a cause of visual plasticity

Regardless of whether premature patterned vision has a clinically relevant role for human

vision, the experiments here demonstrate clearly that premature patterned vision has a long-

lasting impact on the development of receptive fields in the visual cortex. The results suggest

that the evolved late eye opening of many rodents and carnivores, and the prolonged prenatal

period of primates, may help to protect the visual system from exposure to high luminance, high

contrast, and high spatial frequency stimulation until the visual system is ready for this type of

stimulation. This shielding of the developing visual system is not found in all vertebrates, as

tadpoles and zebrafish have transparent eyelids that permit patterned vision from very early

stages (Avitan et al., 2017, Demas et al., 2012). By altering premature patterned vision, it may be

possible to uncover the plasticity rules that are at play in the developing mammalian visual

system before the natural onset of vision.

Figure 1: Recording schematics, experimental timeline, and electrode tract reconstructions.

A: Recording schematics: EO1 cells were those recorded in monocular cortex contralateral to a

single prematurely-opened eye. EOIPS cells were those recorded in monocular cortex ipsilateral

to a single prematurely-opened eye. B: EO2 cells were those recorded in monocular cortex after

both eyes were opened prematurely. C: Control cells were those recorded in monocular cortex

after both eyes were allowed to open naturally. D: Experimental timeline. Animals that

experienced premature vision had one eye or both eyes opened on P25. All animals experienced a

period outside of the nest in the form of daily 2 hour handling sessions beginning on P25 and

ending on P28, allowing unguided viewing of the laboratory environment through prematurely

opened eyes or low resolution experience through closed lids. Electrophysiological recording took

place between P55-68, at a time when the critical period for ocular dominance was closing or

closed (Issa et al. 1999). E: Electrode track reconstructions showing penetrations along the

posterior cortical surface, in monocular V1 (Law et al., 1988; White et al. 1999). Following

recording, the recording electrode was removed and replaced with a marking electrode coated in

DiI.

Figure 2: Early eye opening produced only minor changes in orientation and direction

selectivity index values: A: Example direction tuning curves of individual EO1 cells measured

after the close of the critical period. Direction tuning curves are shown for stimulation at the

optimal temporal frequency. Direction tuning curves look coarsely normal. Horizontal scale bar

represents 10 spikes per second unless otherwise indicated. Dashed line shows mean response to

blank / control stimuli. Number indicates 1-DCV value. B: Same, for EO2 cells C: Same, for

EOIPS cells: D: Same, for control cells E: Orientation tuning curves of individual EO1 cells,

measured in different epochs and in some different animals than the direction tuning curves.

Temporal frequency was 2 Hz, and the spatial frequency optimal for each cell. Numbers indicate

orientation selectivity index values 1-CV. F: EO2 cells G: EOIPS cells H: control cells I: Results

of a linear mixed effects analysis of orientation selectivity index values (1-CV) for all cells and

conditions in the study. Each small column shows the cells for an individual animal. Purple lines

indicate random effect values for each animal. Gray lines indicate condition means and standard

errors of the mean from the linear mixed effect model. Animals within a group have been sorted

by mean index value. Condition coefficients that differ significantly from 0 are indicated with *

($p<0.05$). See text for P values. J: Same for direction selectivity index values, as assessed by (1-

DCV). E01 and E02 cells exhibited slightly higher orientation selectivity index values than

control animals, while EOIPS cells exhibited slightly lower direction selectivity values than

other conditions.

(Figure 3)

Figure 3: Premature eye opening altered temporal frequency tuning after maturity:

Representative temporal frequency tuning curves recorded from single cells: A: EO1: cells contralateral to a single prematurely opened eye. Several cells exhibited strong responses at the lowest temporal frequency tested (that is, several exhibited low-pass tuning). Temporal frequency reported at each cell's overall preferred direction (assessed over all temporal frequencies). Number indicates bandwidth in octaves. $-\infty$ is low pass. B: EO2: cells recorded in an animal with both eyes prematurely opened. Again, many cells exhibited low-pass responses. C: EOIPS: cells ipsilateral to a single prematurely opened eye. D: Control: cells recorded in animals that opened both eyes naturally. E: Linear mixed effects model plot of median TF preference; linear mixed effects plotted as in **Figure 2**. Median TF preference was slightly higher in EOIPS cells than other cells. F: TF low pass index values were significantly elevated in EO1 cells, indicating stronger responses to the lowest temporal frequency tested. G: No differences were observed in high pass index values. H. Bandwidth of the rectified response. I: Bandwidth of the absolute value of responses. Cells in animals that experienced early eye opening exhibited wider bandwidths (less selectivity) than control animals. See text for numbers and p-values.

(Figure 4)

Figure 4: Premature eye opening had very small impacts on spatial frequency tuning. A:
Example spatial frequency tuning curves in EO1 cells. B: Same for EO2 cells. C: Same for
EOIPS cells. D: Same for control cells. E: Spatial frequency peak preferences for all cells,
animals, and experimental groups (same linear mixed effect model plot layout as Fig. 2). F:
Spatial frequency low pass index values. EOIPS cells exhibit a slightly significant (p<0.499)
decrease in low pass index but differences across groups were small. G: High pass index values
for spatial frequency; differences were modest across groups although cells with strong high-pass
responses were more common in E02 cells. H: Spatial frequency bandwidth, in octaves. Most
cells exhibited low-pass profiles and had infinite bandwidth, and no differences were noted
across the groups. I: Spatial frequency bandwidth for absolute response.

Figure 5: Premature eye opening altered response suppression below baseline and

spontaneous activity of V1 neurons: A: Maximum firing rates of cells to visual stimulation

were unchanged across experimental conditions. B: Response suppression rates, defined as the

largest reduction in responses to visual stimulation below background firing rates, were

substantially bigger in EOIPS, EO1, and EO2 cells, indicating that responses below baseline

were features of the stimulus response in many of these cells. C: Background firing rates.

Responses of cells during presentation of control stimuli, where the screen remained gray for the

same duration as our typical visual stimuli. EOIPS, EO1, and EO2 cells showed greatly and

significantly elevated firing rates compared to controls. See text for numbers.

Citations:

Ackman, J. B., T. J. Burbridge and M. C. Crair (2012). "Retinal waves coordinate patterned activity throughout the developing visual system." Nature **490**(7419): 219-225.

Ackman, J. B. and M. C. Crair (2014). "Role of emergent neural activity in visual map development." Curr Opin Neurobiol **24**(1): 166-175.

Adams, D. L. and J. C. Horton (2003). "Capricious expression of cortical columns in the primate brain." Nat Neurosci **6**(2): 113-114.

Akerman, C. J., D. Smyth and I. D. Thompson (2002). "Visual experience before eye-opening and the development of the retinogeniculate pathway." Neuron **36**(5): 869-879.

Akerman, C. J., M. S. Grubb and I. D. Thompson (2004). "Spatial and temporal properties of visual responses in the thalamus of the developing ferret." J Neurosci **24**(1): 170-182.

Atkinson, J., & Braddick, O. (2007). Visual and visuocognitive development in children born very prematurely. *Prog Brain Res*, *164*, 123-149.

Avitan, L., Pujic, Z., Molter, J., Van De Poll, M., Sun, B., Teng, H., Amor, R., Scott, E. K., & Goodhill, G. J. (2017). Spontaneous Activity in the Zebrafish Tectum Reorganizes over Development and Is Influenced by Visual Experience. *Curr Biol*, *27*(16), 2407-2419

Ben-Yishai, R., Bar-Or, R. L., & Sompolinsky, H. (1995). Theory of orientation tuning in visual cortex. *Proc Natl Acad Sci U S A*, *92*(9), 3844-3848.

Benassi, M., Bolzani, R., Forsman, L., Aden, U., Jacobson, L., Giovagnoli, S., & Hellgren, K. (2018). Motion Perception and Form Discrimination in Extremely Preterm School-Aged Children. *Child Dev*, *89*(6), e494-e506.

Bissen, D., Cary, B. A., Zhang, A., Sailor, K. A., Van Hooser, S. D., & Turrigiano, G. G. (2025). Prey capture learning drives critical period-specific plasticity in mouse binocular visual cortex. *bioRxiv*. https://doi.org/10.1101/2025.01.28.635373

Bock, A. S., C. D. Kroenke, E. N. Taber and J. F. Olavarria (2012). "Retinal input influences the size and corticocortical connectivity of visual cortex during postnatal development in the ferret." J Comp Neurol **520**(5): 914-932.

Brainard, D. H. (1997). "The Psychophysics Toolbox." Spat Vis **10**(4): 433-436.

Cang, J., R. C. Renteria, M. Kaneko, X. Liu, D. R. Copenhagen and M. P. Stryker (2005). "Development of precise maps in visual cortex requires patterned spontaneous activity in the retina." Neuron **48**(5): 797-809.

Chang, J. T., D. Whitney and D. Fitzpatrick (2020). "Experience-Dependent Reorganization Drives Development of a Binocularly Unified Cortical Representation of Orientation." Neuron **107**(2): 338-350 e335.

Chapman, B. and M. P. Stryker (1993). "Development of orientation selectivity in ferret visual cortex and effects of deprivation." J Neurosci **13**(12): 5251-5262.

Chapman, B., M. P. Stryker and T. Bonhoeffer (1996). "Development of orientation preference maps in ferret primary visual cortex." J Neurosci **16**(20): 6443-6453.

Colonnese, M. T., Kaminska, A., Minlebaev, M., Milh, M., Bloem, B., Lescure, S., Moriette, G., Chiron, C., Ben-Ari, Y., & Khazipov, R. (2010). A conserved switch in sensory processing prepares developing neocortex for vision. *Neuron*, *67*(3), 480–498. https://doi.org/10.1016/j.neuron.2010.07.015

Crump, C., Winkleby, M. A., Sundquist, J., & Sundquist, K. (2019). Prevalence of Survival Without Major Comorbidities Among Adults Born Prematurely. *JAMA*, *322*(16), 1580-1588

Demas, J. A., Payne, H., & Cline, H. T. (2012). Vision drives correlated activity without patterned spontaneous activity in developing Xenopus retina. *Dev Neurobiol*, *72*(4), 537-546.

Desai, N. S., Rutherford, L. C., & Turrigiano, G. G. (1999). Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nat Neurosci*, *2*(6), 515-520.

Dumas, T. C. (2004). "Early eyelid opening enhances spontaneous alternation and accelerates the development of perforant path synaptic strength in the hippocampus of juvenile rats." Dev Psychobiol **45**(1): 1-9.

Dutton, G. N. (2013). "The spectrum of cerebral visual impairment as a sequel to premature birth: an overview." Documenta Ophthalmologica **127**(1): 69-78.

Foreman, N. and M. Altaha (1991). "The development of exploration and spontaneous alternation in hooded rat pups: effects of unusually early eyelid opening." Dev Psychobiol **24**(7): 521-537.

Gallo, J. E., & Lennerstrand, G. (1991). A population-based study of ocular abnormalities in premature children aged 5 to 10 years. *Am J Ophthalmol*, *111*(5), 539-547

Griswold, S.V. (2024) Premature vision drives aberrant development of response properties in primary visual cortex. PhD Thesis, Brandeis University. https://doi.org/10.48617/etd.1185

Grubb, M. S., F. M. Rossi, J. P. Changeux and I. D. Thompson (2003). "Abnormal functional organization in the dorsal lateral geniculate nucleus of mice lacking the beta 2 subunit of the nicotinic acetylcholine receptor." Neuron **40**(6): 1161-1172.

Guzzetta, A., Tinelli, F., Del Viva, M. M., Bancale, A., Arrighi, R., Pascale, R. R., & Cioni, G.
(2009). Motion perception in preterm children: role of prematurity and brain damage.
Neuroreport, *20*(15), 1339-1343

Hatta, S., T. Kumagami, J. Qian, M. Thornton, E. L. Smith, 3rd and Y. M. Chino (1998).
"Nasotemporal directional bias of V1 neurons in young infant monkeys." Invest Ophthalmol Vis
Sci **39**(12): 2259-2267.

Heimel, J. A., S. D. Van Hooser and S. B. Nelson (2005). "Laminar organization of response
properties in primary visual cortex of the gray squirrel (Sciurus carolinensis)." J Neurophysiol
94(5): 3538-3554.

Hengen, K. B., Lambo, M. E., Van Hooser, S. D., Katz, D. B., & Turrigiano, G. G. (2013).
Firing rate homeostasis in visual cortex of freely behaving rodents. *Neuron*, *80*(2), 335-342.

Hensch, T. K. and E. M. Quinlan (2018). "Critical periods in amblyopia." Vis Neurosci **35**:
E014. Hou, C., A. M. Norcia, A. Madan, S. Tith, R. Agarwal and W. V. Good (2011). "Visual
cortical function in very low birth weight infants without retinal or cerebral pathology." Invest
Ophthalmol Vis Sci **52**(12): 9091-9098.

Hou, C., Norcia, A. M., Madan, A., Tith, S., Agarwal, R., & Good, W. V. (2011). Visual cortical
function in very low birth weight infants without retinal or cerebral pathology. *Invest*
Ophthalmol Vis Sci, *52*(12), 9091-9098

Hubel, D. H. and T. N. Wiesel (1970). "The period of susceptibility to the physiological effects
of unilateral eye closure in kittens." J Physiol **206**(2): 419-436.

Huberman, A. D., Feller, M. B., & Chapman, B. (2008). Mechanisms underlying development of
visual maps and receptive fields. *Annu Rev Neurosci*, *31*, 479-509

Hubel, D. H., T. N. Wiesel and S. LeVay (1977). "Plasticity of ocular dominance columns in

Humphrey, A. L. and A. B. Saul (1998). "Strobe rearing reduces direction selectivity in area 17
by altering spatiotemporal receptive-field structure." J Neurophysiol **80**(6): 2991-3004.

Huttenlocher, P. R. (1967). "Development of cortical neuronal activity in the neonatal cat." Exp
Neurol **17**(3): 247-262.

Imbert, M. and P. Buisseret (1975). "Receptive field characteristics and plastic properties of
visual cortical cells in kittens reared with or without visual experience." Exp Brain Res **22**(1):
25- 36.

Issa, N. P., J. T. Trachtenberg, B. Chapman, K. R. Zahs and M. P. Stryker (1999). "The critical
period for ocular dominance plasticity in the Ferret's visual cortex." J Neurosci **19**(16): 6965-
6978.

Jain, S., Sim, P. Y., Beckmann, J., Ni, Y., Uddin, N., Unwin, B., & Marlow, N. (2022).
Functional Ophthalmic Factors Associated With Extreme Prematurity in Young Adults. *JAMA*
network open, *5*(1), e2145702. https://doi.org/10.1001/jamanetworkopen.2021.45702

Jun, J. J., C. Mitelut, C. Lai, S. L. Gratiy, C. A. Anastassiou and T. D. Harris (2017). "Real-time
spike sorting platform for high-density extracellular probes with ground-truth validation and drift
correction." bioRxiv: 101030.

Katz, L. C., & Shatz, C. J. (1996). Synaptic activity and the construction of cortical circuits.
Science, *274*(5290), 1133-1138.

Kennedy, H. and G. A. Orban (1983). "Response properties of visual cortical neurons in cats
reared in stroboscopic illumination." J Neurophysiol **49**(3): 686-704.

Kenny, P. A. and G. Turkewitz (1986). "Effects of unusually early visual stimulation on the
development of homing behavior in the rat pup." Dev Psychobiol **19**(1): 57-66.

Kozeis, N. (2010). "Brain visual impairment in childhood: mini review." Hippokratia **14**(4): 249-
251.

Krug, K., C. J. Akerman and I. D. Thompson (2001). "Responses of neurons in neonatal cortex
and thalamus to patterned visual stimulation through the naturally closed lids." J Neurophysiol
85(4): 1436-1443.

Law, M. I., Zahs, K. R., & Stryker, M. P. (1988). Organization of primary visual cortex (area 17)
in the ferret. *The Journal of comparative neurology*, *278*(2), 157–180.
https://doi.org/10.1002/cne.902780202

Lehmann, K. and S. Lowel (2008). "Age-dependent ocular dominance plasticity in adult mice."
PLoS One **3**(9): e3120.

LeVay, S., T. N. Wiesel and D. H. Hubel (1980). "The development of ocular dominance
columns in normal and visually deprived monkeys." J Comp Neurol **191**(1): 1-51.

Li, Y., D. Fitzpatrick and L. E. White (2006). "The development of direction selectivity in ferret
visual cortex requires early visual experience." Nat Neurosci **9**(5): 676-681.

Li, Y., S. D. Van Hooser, M. Mazurek, L. E. White and D. Fitzpatrick (2008). "Experience with
moving visual stimuli drives the early development of cortical direction selectivity." Nature
456(7224): 952-956.

Lickliter, R. (2000). "Atypical perinatal sensory stimulation and early perceptual development:
insights from developmental psychobiology." J Perinatol **20**(8 Pt 2): S45-54.

Lickliter, R (2018). The influence of prenatal experience on behavioral and social development:
The

benefits and limitations of an animal model. Dev Psychopathol 30, 871-880, doi:10.1017/
S0954579418000640.

MacKay, T. L., L. S. Jakobson, D. Ellemberg, T. L. Lewis, D. Maurer and O. Casiro (2005).
"Deficits in the processing of local and global motion in very low birthweight children."
Neuropsychologia **43**(12): 1738-1748.

Mazurek, M., M. Kager and S. D. Van Hooser (2014). "Robust quantification of orientation
selectivity and direction selectivity." Front Neural Circuits **8**: 92.

Meister, M., R. O. Wong, D. A. Baylor and C. J. Shatz (1991). "Synchronous bursts of action
potentials in ganglion cells of the developing mammalian retina." Science **252**(5008): 939-943.

Moore, B. D., 4th, Alitto, H. J., & Usrey, W. M. (2005). Orientation tuning, but not direction
selectivity, is invariant to temporal frequency in primary visual cortex. *Journal of*
neurophysiology, *94*(2), 1336–1345. https://doi.org/10.1152/jn.01224.2004.

Movshon, J. A., Kiorpes, L., Hawken, M. J., & Cavanaugh, J. R. (2005). Functional maturation
of the macaque's lateral geniculate nucleus. *The Journal of Neuroscience*, *25*(10), 2712–2722.
https://doi.org/10.1523/JNEUROSCI.2356-04.2005.

Muir-Robinson, G., B. J. Hwang and M. B. Feller (2002). "Retinogeniculate axons undergo eye-
specific segregation in the absence of eye-specific layers." J Neurosci **22**(13): 5259-5264.

Murata, Y. and M. T. Colonnese (2016). "An excitatory cortical feedback loop gates retinal wave
transmission in rodent thalamus. Elife Oct 11:5:e18816. doi: 10.7554/eLife.18816.

Murcia-Belmonte, V., Y. Coca, C. Vegar, S. Negueruela, C. de Juan Romero, A. J. Valino, S.
Sala, R. DaSilva, A. Kania, V. Borrell, L. M. Martinez, L. Erskine and E. Herrera (2019). "A
Retino-retinal Projection Guided by Unc5c Emerged in Species with Retinal Waves." Curr Biol
29(7): 1149-1160 e1144.

Olavarria, J. F. and R. Hiroi (2003). "Retinal influences specify cortico-cortical maps by
postnatal day six in rats and mice." J Comp Neurol **459**(2): 156-172.

Ortibus, E., L. Lagae, I. Casteels, P. Demaerel and P. Stiers (2009). "Assessment of cerebral
visual impairment with the L94 visual perceptual battery: clinical value and correlation with
MRI findings." Dev Med Child Neurol **51**(3): 209-217.

Osaki, H., T. Naito, O. Sadakane, M. Okamoto and H. Sato (2011). "Surround suppression by
high spatial frequency stimuli in the cat primary visual cortex." Eur J Neurosci **33**(5): 923-932.

Pamir, Z., C. M. Bauer, E. S. Bailin, P. J. Bex, D. C. Somers and L. B. Merabet (2021). "Neural
correlates associated with impaired global motion perception in cerebral visual impairment
(CVI)." Neuroimage Clin **32**: 102821.

Peirce, J. W. (2007). "The potential importance of saturating and supersaturating contrast
response functions in visual cortex." J Vis **7**(6): 13.

Pelli, D. G. (1997). "The VideoToolbox software for visual psychophysics: transforming
numbers into movies." Spat Vis **10**(4): 437-442.

Ringach, D. L., R. M. Shapley and M. J. Hawken (2002). "Orientation selectivity in macaque
V1: diversity and laminar dependence." J Neurosci **22**(13): 5639-5651.

Ritter, N. J., N. M. Anderson and S. D. Van Hooser (2017). "Visual Stimulus Speed Does Not
Influence the Rapid Emergence of Direction Selectivity in Ferret Visual Cortex." J Neurosci
37(6): 1557-1567.

Rossi, F. M., T. Pizzorusso, V. Porciatti, L. M. Marubio, L. Maffei and J. P. Changeux (2001).
"Requirement of the nicotinic acetylcholine receptor beta 2 subunit for the anatomical and
functional development of the visual system." Proc Natl Acad Sci U S A **98**(11): 6453-6458.

Roy, A., Christie, I. K., Escobar, G. M., Osik, J. J., Popovic, M., Ritter, N. J., Stacy, A. K.,
Wang, S., Fiser, J., Miller, P., & Van Hooser, S. D. (2018). Does experience provide a
permissive or instructive influence on the development of direction selectivity in visual cortex?
Neural Dev, *13*(1), 16.

Roy, A., Osik, J. J., Meschede-Krasa, B., Alford, W. T., Leman, D. P., & Van Hooser, S. D.
(2020a). Synaptic and intrinsic mechanisms underlying development of cortical direction
selectivity. *eLife*, 9, e58509. https://doi.org/10.7554/eLife.58509.

Roy, A., Wang, S., Meschede-Krasa, B., Breffle, J., & Van Hooser, S. D. (2020b). An early
phase of instructive plasticity before the typical onset of sensory experience. *Nature*
communications, *11*(1), 11. https://doi.org/10.1038/s41467-019-13872-1.

Rubin, D. B., Van Hooser, S. D., & Miller, K. D. (2015). The stabilized supralinear network: a
unifying circuit motif underlying multi-input integration in sensory cortex. *Neuron*, *85*(2), 402-
417.

Sakki, H. E. A., N. J. Dale, J. Sargent, T. Perez-Roche and R. Bowman (2018). "Is there
consensus in defining childhood cerebral visual impairment? A systematic review of terminology
and definitions." Br J Ophthalmol **102**(4): 424-432.

Sleigh, M. J. and M. B. Casey (2014). "Prenatal sensory experience affects hatching behavior in
domestic chicks (Gallus gallus) and Japanese quail chicks (Coturnix coturnix japonica)." Dev
Psychobiol **56**(5): 935-942.

Sleigh, M. J. & Lickliter, R (1995). Augmented prenatal visual stimulation alters postnatal
auditory and
visual responsiveness in bobwhite quail chicks. Dev Psychobiol 28, 353-366, doi:10.1002/
dev.420280702

Slidsborg, C., R. Bangsgaard, H. C. Fledelius, H. Jensen, G. Greisen and M. la Cour (2012).
"Cerebral damage may be the primary risk factor for visual impairment in preschool children
born extremely premature." Arch Ophthalmol **130**(11): 1410-1417.

Smith, G. B., B. Hein, D. E. Whitney, D. Fitzpatrick and M. Kaschube (2018). "Distributed
network interactions and their emergence in developing neocortex." Nat Neurosci **21**(11): 1600-
1608.

Somers, D. C., Nelson, S. B., & Sur, M. (1995). An emergent model of orientation selectivity in
cat visual cortical simple cells. *J Neurosci*, *15*(8), 5448-5465.

Taylor, N. M., Jakobson, L. S., Maurer, D., & Lewis, T. L. (2009). Differential vulnerability of
global motion, global form, and biological motion processing in full-term and preterm
children. *Neuropsychologia*, *47*(13), 2766–2778.
https://doi.org/10.1016/j.neuropsychologia.2009.06.001

Tiriac, A., B. E. Smith and M. B. Feller (2018). "Light Prior to Eye Opening Promotes Retinal
Waves and Eye-Specific Segregation." Neuron **100**(5): 1059-1065 e1054.

Turrigiano, G. G., Leslie, K. R., Desai, N. S., Rutherford, L. C., & Nelson, S. B. (1998).
Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature*, *391*(6670), 892-
896

Turrigiano, G. G., & Nelson, S. B. (2004). Homeostatic plasticity in the developing nervous
system. *Nature reviews. Neuroscience*, *5*(2), 97–107. https://doi.org/10.1038/nrn1327

Van Hooser, S. D., Y. Li, M. Christensson, G. B. Smith, L. E. White and D. Fitzpatrick (2012).
"Initial neighborhood biases and the quality of motion stimulation jointly influence the rapid
emergence of direction preference in visual cortex." J Neurosci **32**(21): 7258-7266.

Wang, B. S., R. Sarnaik and J. Cang (2010). "Critical period plasticity matches binocular
orientation preference in the visual cortex." Neuron **65**(2): 246-256.

Weliky, M. and L. C. Katz (1999). "Correlational structure of spontaneous neuronal activity in
the developing lateral geniculate nucleus in vivo." Science **285**(5427): 599-604.

White, L. E., Bosking, W. H., Williams, S. M., & Fitzpatrick, D. (1999). Maps of central visual
space in ferret V1 and V2 lack matching inputs from the two eyes. *The Journal of*
neuroscience, *19*(16), 7089–7099. https://doi.org/10.1523/JNEUROSCI.19-16-07089.1999

Wong, R. O., M. Meister and C. J. Shatz (1993). "Transient period of correlated bursting activity
during development of the mammalian retina." Neuron **11**(5): 923-938.

Wong, R. O. (1999). Retinal waves and visual system development. *Annu Rev Neurosci*, *22*, 29-
47.