Impacts of climate change on fonio millet: seed germination and suitability modelling of an important indigenous West African crop

George P. Burton[1,2], Hillary Mireku Botey[3], Paolo Ceci[4], Caspar Chater[1], Rafal M. Gutaker[1], Amy C. Jackson[1], Philippa Ryan[1], Charlotte E. Seal[5], Colin G. N. Turnbull[2], Maria S. Vorontsova[1], Efisio Mattana[5] and Tiziana Ulian[5, 6].

[1] Royal Botanic Gardens, Kew, Richmond, Kew, TW9 3AB, United Kingdom

[2] Department of Life Sciences, Imperial College London, London, SW7 2AZ, United Kingdom

[3] CSIR-Crops Research Institute, Kumasi, Ghana

[4] Food and Agriculture Organisation of the United Nations, Sub-Regional Office for West Africa (FAO-SFW), Senegal

[5] Royal Botanic Gardens, Kew, Wellcome Trust Millennium Building, Wakehurst, Ardingly, UK

[6] Department of Life Sciences and Systems Biology (DBIOS), University of Torino, Torino, Italy

Abstract

Seed germination is highly temperature sensitive. Climate change factors such as increasing temperatures are likely to have a harmful effect on agriculture, particularly after crop sowing. Better utilisation of indigenous, arid-resilient crops like fonio (*Digitaria exilis*) are a commonly proposed solution to improving food security. This study develops knowledge of fonio germination requirements and how these correspond to future climate conditions across West Africa. We use a combined approach; integrating seed germination experiments under a range of temperatures, and niche suitability modelling to investigate how cultivation of fonio will be impacted by climate change. We find that from 37 seed accessions collected across Guinea, Togo, Mali, and Burkina Faso, the ceiling temperature for germination is around 42°C, with an optimum temperature of 30-35°C – also noted from phenotypic observations. Drought trials show successful germination to beyond -1MPa. There is no obvious difference in response by accessions originating from either hotter or cooler climates. By comparing these temperature thresholds with future climate predictions, alongside a suitability modelling approach, we predict an average decline of 10% in the suitable area for fonio cultivation, especially affecting Senegal, Mali, and Burkina Faso. Newly suitable area is predicted to increase in Guinea, Ghana, Cote d'Ivoire, and Nigeria by around 5%. These findings provide valuable insight for developing future dryland agriculture policies and prioritisation of resilient crops.

Keywords

Fonio, food security, heat stress, agriculture, climate change, *Digitaria exilis*, niche modelling, seed germination

Introduction

Climate change and heat-resilient crops

In past decades there has been a significant increase in both rising global temperatures
and unpredictable, destructive weather patterns which affect both the quality and
availability of food crops, and overall human health (Sylla et al., 2018; Fatima et al., 2020;
Ragatoa et al., 2024). Climate models suggest that an increase of 1-2°C could further
destabilise agricultural systems (Sylla et al., 2018), yet overall global temperatures are
predicted to surpass this, with increases of up to 2-3°C by 2050 (Fatima et al., 2020),
which will be particularly harmful to agriculture in vulnerable arid regions of West Africa.
This is concerning for regions that often prioritise major introduced crops such as Asian
rice (*Oryza sativa* L.) which requires irrigation and is less resilient to the local high heat
and drought than native locally-adapted crops like African rice (*Oryza glaberrima* Steud.,
Linares, 2002). Other indigenous crops, such as cereal millet fonio (*Digitaria exilis*
(Kippist) Stapf.), exhibit high tolerance to heat and drought, making them critical to food
systems for future food security in West Africa (Cruz et al., 2016).

Countries close to the Sahel in West Africa have recently experienced record weather
conditions, including a heatwave in March-April 2024, with prolonged temperatures
above 45°C in Senegal, Mali, and Burkina Faso (World Weather Attribution, April 2024) -
this had severe effects on human health causing hospitalisations and deaths. These
record temperatures were coupled with extreme rainfall and flooding in the Niger and
Lake Chad basins later in Autumn, again causing damage and loss of life (World Weather
Attribution, October 2024). Increases in consistent average temperature and frequency
of extreme weather events are predicted across the Sahel (Ragatoa et al., 2024).

Seed germination is highly sensitive to ambient temperature, and warming climates are
likely to have an adverse effect on both plant germination success and post-germination
seedling development. Millets, however, have a higher-than-average heat tolerance to
protect against these stresses (Wahid et al., 2007; Tadele et al., 2016), utilising various
adaptation strategies to grow in challenging climates (Pirnajmedin et al., 2024; Heckman
et al., 2024). Climate change impacts on food security are directly related to seed
physiology and vigour: the capacity of a crop species to regulate successful germination
and survival under heat and drought stress (Reed et al., 2022). As the climate changes, it
will be essential to identify the specific heat and drought limits of germination for
indigenous crops, to select the most climate-resilient crops appropriate for each region.

Seed germination

Seed germination studies have a longstanding history in botanical research. Systematic
methods for assessing germination rates became commonplace in the 1980's, using the
proportion of germinated seeds under a set of conditions to predict germination under
untested conditions. This involved the concepts of cardinal temperatures: base

temperature (Tb), optimum temperature (To), and ceiling temperature (Tc) – to delimit the
thresholds within a seed population will germinate under a gradient of different
temperatures, to mathematically predict the effects of temperature on seed germination
rate (Garcia-Huidobro et al., 1982; Ellis et al., 1986; Gummerson, 1986; Covell et al.,
1986). These studies introduced the use of a 'thermal-time' model, and later a
'hydrothermal-time' model, the latter to predict the effects of water potential and drought
on germination. These concepts and models are widely applied (Watt & Bloomberg,
2012; Bradford, 2002; Trudgill et al., 2000), including the use of polyethylene glycol (PEG)
to create solutions of different water potential to limit the movement of water into a seed,
thereby simulating drought. Though there have been concerns for how PEG may interact
with toxicity stress, temperature, and substrates like filter paper (Michel, 1983;
Hardegree and Emmerich, 1990), they nonetheless provide an accessible and efficient
method of testing the effects of temperature and/or simulated drought (a decrease in
water potential) on seed germination through time. More recently, dose-response curves
(which rely on a time-to-event model) have been presented as a mathematically accurate
method of calculating germination rates which are inherently non-linear, including using
non-parametric maximum likelihood estimator (NPMLE) models (Onofri et al., 2018;
Catara et al., 2016; Onofri et al., 2022).

Fonio climate resilience

Fonio is a West African millet known to tolerate extreme drought and high temperatures
(Tadele et al., 2016; Cruz et al., 2016), as well as grow in unfertilised, un-irrigated soils,
and has an important cultural role in religious and social ceremonies (Burton et al., 2023).
It is therefore an important crop in supporting food security of rural communities in West
Africa (Cruz et al., 2016). Despite its important role in sustainable, subsistence
agriculture and potential for improving livelihoods in rural communities, fonio has been
recognised as a Neglected and Underutilised Species (NUS, Ulian et al., 2020): a
traditional crop which has experienced a low degree of research and commercial
interest, due to its smaller seeds and lower yields, compared to major economic crops.
One of the only germination-based growth tests for fonio was undertaken by Portères
(1955), who observed the developmental stages of germinating fonio seedlings at a range
of temperatures up to ~30°C, observing an optimum temperature of around ~30°C, and
predicting a ceiling temperature of between 40-45°C. Portères commented that these
temperatures were perfectly suited to the climate averages in West Africa, during the
1950s. For fonio millets more recently there have been few field or laboratory
experiments which examine the specific constraints and germination limits under
different climate change scenarios. Senegalese field trials by Gueye et al. (2015)
throughout 2010 and 2011 evaluated white fonio (*D. exilis*) sowing and harvesting times
versus climate conditions, and Petri-dish germination trials by Iwuala et al. (2021)
evaluated the drought response of the close relative and sister crop black fonio (*Digitaria*
iburua Stapf.) on three genotypes from the National Cereals Research Institute in Nigeria.

More recently, Pudasaini et al. (2025) conducted an extensive lab-based experiment
comparing the difference in morphological response between one white fonio genotype
from Mali (arid) and one from Guinea (wet), presenting responses in inflorescence, root
and leaf development during irrigated versus drought conditions.

Modelling future climates and crop suitability

As well as identifying the environmental limits that constrain fonio seed germination, it is
useful to compare these with predictions of the temperatures across West Africa in the
present and future. Common methodology for predicting what future environments will
look like use 'Shared Socio-economic Pathways' (SSPs) - a set of 5 possible economic,
political, and social development factors (Van Vuuren et al., 2017; Kriegler et al., 2012).
These are categorised as: 1, an almost total transition to green energy, and commitment
to global development and reducing inequality; 2, little shift from current behaviours and
trends; 3, international division which prioritises focus on local regions, with some
inequality; 4, increased inequality both within and between regions; and 5, almost total
reliance on fossil-fuels and geo-engineering. The SSPs are commonly combined with
Representative Concentration Pathways (RCPs) which predict future greenhouse gas
emissions, ranging from RCP2.6 with low-emissions, to RCP8.5 - high emissions
(Meinshausen et al. 2011). The combination of these two pathway strategies (for example
SSP1 and RCP2.6 = SSP126) are then used to predict effects on the environment under
different future scenarios: SSP126, SSP245, SSP370, and SSP585.

Every species has its own set of biological limits for living and reproducing, including
cereal crops such as fonio. These observed thresholds can be inferred from current
geographic occurrences and matched with current climate variables, and future suitable
distributions predicted using future climate modelling – which is especially relevant in
the scope of exploring how agricultural crops will be affected by climate change.
Although this method is entirely theoretical, it can provide a valuable foundation for
further interpretation (Pearson and Dawson, 2003), especially when compared to a
species' observed physiological limitations. There are several popular analytical
methods which utilise a machine-learning approach to model and predict these species
suitability niches in the present and future, including MaxEnt (Phillips and Dudík, 2008;
Merow et al., 2013), random forest (Evans et al., 2010), and CLIMEX (Sutherst et al.,
1999). Producing an accurate set of results using these methods is dependent on careful
selection of data inputs, including valid species distribution data and a set of bioclimatic
variables which are relevant both to the species suitability limits, and the geographic
region itself (Phillips and Dudík, 2008; Merow et al, 2013).

Fonio was included in a similar study of African crop suitability predictions under future
climate change, by Chemura et al. (2024), which uses a combination of climate variables
alongside known crop traits, to predict environmental niche suitability under different
climate scenarios. Similarly, a plant growth study by Gueye et al. (2013) did not use

climate modelling against the raw physiological experiments directly for white fonio
varieties in Senegal, but the region-specific scope was useful for predicting the best
sowing times in different areas, throughout the year. A combination of these approaches
is likely to be effective: seed growth experiments and comparison of thresholds against
future temperature predictions, and future environmental niche modelling based on
current occurrences.

Aims and Scope

For this study, as our interest is in determining the thresholds of heat and water potential
resilience of fonio for comparison with climates in the future, it is important to test a wide
range of accessions from different geographic areas, using a wide range of experimental
conditions to detect specific tolerances. We combine two main methods to answer this
question: germination trials to determine temperature and water potential thresholds,
compared to current and future predicted temperatures in West Africa; and
environmental niche analysis using known fonio distribution data (from GBIF) and
associated bioclimatic variables, to model future suitability under different SSP
scenarios. We also hypothesise that accessions may have different regional tolerances
due to its large geographical range.

We investigated the germination thresholds of fonio, for temperature and PEG-induced
drought, compare them in the current and future climates, and applied predictive
modelling and projections into the near future to assess how suitable fonio cultivation
systems are across different areas of West Africa. Here, we report germination profiles of
seed accessions from across West Africa, at a range of constant temperatures, and niche
suitability models for 2024-2060 under three different SSP scenarios.

Methods

Germplasm accessions

Nineteen seed accessions of *Digitaria exilis* for conducting germination trials were accessed from the collections of the Research Institute for Development (IRD) in Montpellier, France, collected up to 50 years ago by partners across Mali, Guinea, Burkina Faso and Togo, and used in the genomic study of Abrouk et al. (2020). Eighteen accessions from across the Fouta Djallon region of Guinea were also accessed from the Herbier National De Guinée (HNG), collected during a study by Burton et al. (2024). Details of all collections can be seen in **Figure** 1 and **Table** 1 – which range from an altitude of 28-1062m, maximum average temperature of 30-43°C, and minimum average temperature of 17-29°C. Accessions 1-19 (Table 1) of white fonio from 4 countries provided an 'international' range from two typically separate climates (high maximum average temperature and arid climate in Mali and Burkina Faso, and a lower maximum average temperature and sub-tropical climate in Guinea and Togo; we refer to these as 'hot' and 'cool' climates). These samples were chosen as a good representation across the wider geographical cultivation region for white fonio, and among genetic populations (the three genetic clusters first identified by Adoukonou-Sagbadja et al., 2007). Accessions 18-37 from the HNG, collected by Burton et al. (2024), were used to provide a wider range of fonio germplasm, to determine average and maximum germination indices.

Figure 1 – Germplasm collections used in seed germination experiments. Separate collections donated by the IRD (France) and HNG (Guinea) seedbanks are shown, displayed as triangles and circles respectively, and numbered according to accession number given in Table 1. Colours show recent (1970-2000) average maximum temperature across the region in July, separated for above and below 30°C. Accessions collected in Guinea and Togo are classified to be from 'cool' vs. 'hot' climates in Mali and Burkina Faso.

Accession	Catalogue Code	Collection	Year	Country	Region	Village	Latitude	Longitude	Altitude	Landrace Name	Analysis
1	CM06496	IRD	1981	Burkina Faso	Djibo	Baraboule	14.20	-1.85	306	Sereme	IND, RM
2	CM06501	IRD	1981	Burkina Faso	Tougan	Dagale	13.37	-3.08	272	Fin	IND, RM
3	CM07890	IRD	1981	Burkina Faso	Sideradougou	Degue	10.58	-4.85	290	Foni Ba	IND, RM
4	CM05742	IRD	1978	Mali	Kayes	Takoutala	14.72	-12.07	28	Fonio	IND, RM
5	CM05772	IRD	1978	Mali	Kita	Kokofata	12.88	-9.95	331	Founi Ba	IND, RM
6	CM05796	IRD	1978	Mali	Bougouni	Dossola	11.53	-8.08	370	Fini Le	IND
7	CM05836	IRD	1978	Mali	Douentza	Konsa	15.05	-3.88	263	Sereme	IND, RM
8	CM05858	IRD	1978	Mali	Bandiagara	Dialo	14.18	-3.88	422	Mpongon	IND, DR
9	CM05863	IRD	1978	Mali	Segou	Togou	13.58	-6.00	288	Fini Ba	IND, RM
10	CM03431	IRD	1977	Togo	Badou	Benali	7.58	0.73	544	Owa	IND, RM, DR
11	CM03438	IRD	1977	Togo	Amlame	Lele	9.42	0.92	328	Gbokoui	IND, RM, DR, DRM
12	CM03439	IRD	1977	Togo	Kapalime	Kati	6.57	0.85	118	Gbokoui	IND, RM
13	CM07244	IRD	1982	Guinea	Dabola	N Dema	10.63	-10.97	496	Fognie Dani	IND, RM
14	CM07249	IRD	1982	Guinea	Dabola	Kambaya	10.70	-11.50	868	Fognie Ba Gbe	IND, RM
15	CM07268	IRD	1982	Guinea	Labe	Tarambali	11.38	-12.13	858	Fognie Hotia	IND, RM
16	CM07327	IRD	1982	Guinea	Mamou	Dar Es Salam	10.23	-11.97	719	Fognie Konso	IND, RM
17	CM07329	IRD	1982	Guinea	Mamou	Bantamaya	10.13	-11.63	393	Fognie Konso	IND, RM, DR
18	CM07354	IRD	1982	Guinea	Kerouane	Konsankoro	9.03	-9.00	537	Founi Ba	IND, RM
19	CM08537	IRD	1982	Guinea	Dalaba	Dittin	10.88	-12.18	697	Sira Rane	IND, RM
20	GPB01	HNG	2022	Guinea	Mamou	Kounkoure	10.31	-12.23	586	Liti	IND
21	GPB03	HNG	2022	Guinea	Mamou	Kigna	10.59	-11.78	643	Bode	IND
22	GPB04	HNG	2022	Guinea	Mamou	Kigna	10.59	-11.78	643	Rane	IND
23	GPB05	HNG	2022	Guinea	Mamou	Bagata	10.71	-11.66	707	Bode	IND
24	GPB06	HNG	2022	Guinea	Mamou	Bagata	10.71	-11.66	707	Bale	IND
25	GPB08	HNG	2022	Guinea	Dalaba	Fogoumba	10.87	-12.11	775	Hotia	IND, DR
26	GPB10	HNG	2022	Guinea	Labé	Popdara	11.43	-12.41	935	Koine	IND
27	GPB11	HNG	2022	Guinea	Labé	Popdara	11.43	-12.41	935	Konso	IND
28	GPB13	HNG	2022	Guinea	Labé	Sannoun	11.43	-12.11	815	Hotia	IND
29	GPB14	HNG	2022	Guinea	Labé	Sannoun	11.43	-12.11	815	Bambougou	IND
30	GPB15	HNG	2022	Guinea	Labé	Sannoun	11.43	-12.11	815	Rane	IND
31	GPB16	HNG	2022	Guinea	Labé	Pellel Koura	11.30	-12.50	970	Bode	IND, DR, DRM
32	GPB17	HNG	2022	Guinea	Labé	Pellel Koura	11.30	-12.50	970	Rane	IND, DR, DRM
33	GPB18	HNG	2022	Guinea	Mali	Fougou	11.91	-12.30	1062	Kensere	IND
34	GPB19	HNG	2022	Guinea	Mali	Fougou	11.91	-12.30	1062	Nyougou	IND
35	GPB20	HNG	2022	Guinea	Mali	Sigon	11.79	-12.21	841	Konso	IND, DR
36	GPB21	HNG	2022	Guinea	Pita	Kambako	10.89	-12.49	960	Bagala	IND
37	GPB22	HNG	2022	Guinea	Pita	Kambako	10.89	-12.49	960	Konso Sokili	IND

Table 1 – Fonio germplasm accessions used in seed germination experiments as part of this study. Details provided include seed accession numbers used in text (1-37) used in text, catalogue code, and seedbank collection (IRD is the Institut de Recherche pour le Développement, France, and HNG is the Herbier National de Guinée). Analysis column shows which accessions are used for each analysis: germination indices (IND), rate modelling (RM), drought indices (DR) and drought modelling (DRM).

Germination trials: temperature and drought simulation

Germination experiments were conducted at the Millenium Seed Bank, Wakehurst, UK, in 2023 and 2024. To test the effect of different temperatures on germination, 20 seeds from each of the 37 accessions were sown in seven 90mm diameter Petri dishes containing water-agar (30ml of 1% Agar, 80-100 Mesh, Fisher Scientific, UK), and placed in incubators (LMS LTD, UK) between 15-45°C at 5°C intervals (constant temperatures at 15°C, 20°C, 25°C, 30°C, 35°C, 40°C, and 45°C), with 3 replicates of each dish. Two photoperiods were applied, 12 hours each of light (LED 55 µmol/m²/s) and darkness. 420 seeds and 21 Petri dishes were tested for each accession, at a total of 777 Petri dishes, and 15,540 seeds. Seed germination was scored every 24hrs for 7 days. A germination event was categorised as a radicle over 2mm in length. Trials were conducted in 4 batches, containing a random order of 8-10 accessions in each (between 168-210 Petri dishes per batch). Ungerminated seeds were dissected and classed as viable or empty determined by presence of an embryo, and germination data adjusted accordingly.

To test the effect of drought (changes in water potential) on seed germination, drought
conditions were simulated using 6 different concentrations of PEG8000 combined with
water, to emulate water potentials at of 0MPa, -0.8MPa, -1MPa, -1.2MPa, -1.4MPa and -
1.6MPa. The solutions were prepared using ratios suggested by Michel (1983) and used
in similar germination drought experiments by Jadav et al. (2016) and Violita and Azhari
(2021): 1ml water containing 0g, 0.25g, 0.3g, 0.32g, 0.34g, and 0.38g, to achieve the
respective water potentials. Seeds were germinated on filter paper (Whatman filter
paper, 90mm diameter, 200µm thickness) inside a 90mm Petri dish, covered with 3ml of
the water:PEG8000 solution. Eight different accessions were tested (8, 10, 11, 17, 25, 31,
32, and 35) which were randomly chosen, but restricted by availability of seed,
particularly accessions from Mali and Burkina Faso. Twenty seeds were sown in each
Petri dish, across the 8 different water potentials, with 3 replicates for each. Seeds were
germinated in incubators at the sub-optimal (20°C) and optimal (30°C) temperatures
(determined by temperature trials), and scored every 48 hours for 10 days. Experiments
were split into 2 batches, with accessions randomly ordered in each.

Germination analysis

Germination events for all experiments were recorded per Petri-dish (each given a unique
identifying code). This included accession number, repeat, temperature, water potential,
viable seed number, number of germination events per time interval (7 time points for
heat tests, and 5 for drought tests), and final germination percentage. Final datasets for
heat and drought tests were transformed into 'wide' and 'long' formats (sorted by scoring
intervals in the first, and by cumulative germination in the second), following the methods
in Onofri et al. (2022). Analysis of germination thresholds, proportion, and rate was
conducted in R environment version 4.4.0 (R Core Team, 2024). The package GerminaR
(Lozano-Isla et al., 2019) was used to produce germination indices including germination
percentage (grp), total number of germinated seeds (grs), mean germination time (mgt),
and germination synchronicity (syn)) for each individual petri-dish. These germination
indices were analysed using GLM (generalised linear mixed), and one and two-way
ANOVA linear models, and conducted separately on each international and regional
dataset, following Gianinetti (2020), to examine the effects of temperature, accession,
seed age, and collection group. This included data from all 37 accessions. Accessions
used in each specific analysis are shown in **Table** 1. T-tests and Bartlett tests were used
to analyse the difference between germination of the two climatic groups of accessions.
The package DRCTE (Onofri et al., 2022) was used to model germination thresholds,
germination rate, and cardinal temperatures for each of the 19 international accessions
from the IRD collection, following Onofri et al. (2022) and Catara (2016). A DRCTE (dose-
responsive curve time-to-event) model object was produced using the NPLME method,
and germination rate as a response to temperature across time was calculated using an
exponential model (GRT.Ex) in the drcSeedGerm package (Onofri et al., 2018). Thermal
thresholds were modelled at time taken for a Petri-dish to reach 50% germination (t50).

The predict() function in R was used to calculate optimum temperature (To) for each
accession using data from the GRT.Ex model. Seven accessions where models would not
converge above t40 were excluded from this analysis: 6, 8, 20-24, and 37.

Similar methods were used to analyse data from drought experiments. Data from trials at
20°C and 30°C were separated, and germination indices calculated for all 8 accessions
tested (details in **Figure** 1). Germination rate curves were calculated using the water
potential model GRPsiPol (rather than GRT.Ex) in drcSeedGerm to explore the
relationship between water potential and germination rate. t50 was used for model
convergence at 30°C, and t15 at 20°C; only 3 accessions (11, 31 and 32) could be
modelled to t50 at 30°C, and none at 20°C. At t15 and 20°C only 3 accessions (10, 31, and
35) could be modelled.

Phenotypic data

After seed germination experiments were complete (at the end of 7 days for heat and 10
days for drought), the weight of total leaf matter from 5 individual seedlings from 3 Petri-
dishes tested at 20°C, 30°C, and 40°C was measured in milligrams (mg) using a
microscale (leaves from a total of 45 seedlings per accession across treatments), for 17
accessions (1, 3, 8, 10, 11, 13, 16, 17, 21, 24, 25, 27, 28, 31, 32, 35, and 37). Seedlings
grown as part of drought trials were also weighed at 0MPa and -1MPa, accessions 10, 25,
32, and 35 grown at 30°C.

Seedlings from accession 1 were photographed periodically over 7 days to capture the
development of the seedling during heat and drought experiments to record qualitive
differences across the gradient of conditions.

Climate Data and Niche Modelling

Climate data was accessed and downloaded from WorldClim ver. 2.1
(https://www.worldclim.org/), all at 10min resolution. For present/historic climate data
we used average maximum temperature (°C) (tmax) from the years 1970-2000, for the
months 5, 6, 7, and 8 (May, June, July, August) to represent the temperatures experienced
by seeds germinating during the sowing period (Burton et al., 2024). This was compared
to the maximum temperature of the same months in future climate predictions, for the
2040-2060 period, under the SSP126, SSP245, SSP370, and SSP585 scenarios, using the
CMCC-ESM2 (Euro-Mediterranean Center on Climate Change Earth System Model)
general circulation model (GCM). Climate values were attached to xy coordinates and
averaged across the sowing period, to predict a count of occurrences occurring within
different temperature ranges shown to be suitable for successful germination (<35°C).

For modelling crop suitability under future climates, occurrence data for *Digitaria exilis*
was downloaded from GBIF (https://www.gbif.org/species/5289953) as representative of
the present occurrence range of the species. GBIF records did not include any
occurrence data from Senegal or Nigeria, which are known to be key cultivation regions
for white fonio (Cruz et al. 2016), and verified records with co-ordinates from these

regions could not be accessed. These absent occurrences were instead used to assess the validity of the suitability models (i.e. see whether the model still predicted these areas to be suitable, despite missing data). Bioclimatic variables for the present (1970-2000) and future (2040-2060) were downloaded from WorldClim (https://www.worldclim.org/data/index.html), under the SSP126, SSP245, SSP370, and SSP585 scenarios, using the CMCC-ESM2 general circulation model (GCM). A full range of all bioclimatic variables were input to niche modelling analysis first, and the variables with the lowest model contributions removed systematically. The remaining bioclimatic variables BIO3 (Isothermality), BIO4 (Temperature Seasonality), BIO5 (Max Temperature of Warmest Month), BIO9 (Mean Temperature of Driest Quarter), BIO10 (Mean Temperature of Warmest Quarter), BIO14 (Precipitation of Wettest Month), BIO15 (Precipitation Seasonality), and BIO17 (Precipitation of Driest Quarter) were chosen as the most useful climatic variables which are likely to be significant indicators of fonio suitability.

Raster layers and maps were processed using ArcGIS Pro 3.4.0. The maps containing current and future variables were cropped to the extent of West Africa, and raster layers converted into ascii files. The current variables and species distribution data were input to Maxent 3.4.3 (Phillips and Dudík, 2004, available from https://biodiversityinformatics.amnh.org/open_source/maxent/), and run to produce an initial map of current species suitability, using the Cloglog model. This was rerun with random test percentage at 30%, which removed a random 30% of the occurrences, to test likelihood with a smaller dataset for the relaxed minimum training presence and strict 10th percentile training presence. The full model was then rerun again with the future climatic variables as projection layers, which was repeated for each future scenario. These results were overlaid in ArcGIS Pro, and maps of regions suitable in the present versus future produced using the Raster Calculator tool and binarized suitable niche maps. Percentage of suitable area was calculated using raw count data provided in ArcGIS.

Results

Response to temperature

Germination synchronicity between accessions (syn), which measures the consistent overlap of germination within groups, is found to be insignificantly affected by temperature and accession ($p>0.05$). Neither date of seed collection nor collection group (IRD vs. HNG) have a significant effect on seed germination time, germination proportion, or synchronicity ($p>0.05$; **Figure** S1).

Temperature had a significant effect on seed germination percentage across all accessions ($p<0.005$) when analysed using a binomial GLM including accession and temperature as fixed effect dependent variables, and grp as a response. A set of graphs demonstrating the relationship between heat and water potential, and seed germination

indices and leaf weight, are given in **Figure** 2. The temperature with the mean highest
successful germination was 30°C (87.2%), followed by 25°C (86.8%), and 35°C (86%). No
germination was observed at 45°C, confirming that this temperature exceeds fonio's
possible germination range. A significant relationship between mean germination time
(mgt) and temperature was also observed (p<0.005), where the temperature with the
fastest germination was 35°C (1.59), followed by 40°C (1.79), and 30°C (1.85), measured
in average time taken for germination.

Figure 2 – Response of seed germination proportion (row A), mean germination time (row B), and leaf weight (row C), to temperature (boxplots in red) and water potential (boxplots in blue). Graph captions give sample size (n) for each boxplot, and number of seed accessions used. MGT measurement is given as average time interval taken for successful germination.

Cardinal temperatures were modelled for 19 accessions using t50 (time taken to achieve
50% successful germination per Petri-dish). Average Tb was found to be 14.52°C
(SD=1.7°C, maximum 17.99°C, minimum 11.72°C), To was 36°C (SD=2.15°C, maximum
38°C, minimum 30°C), Tc was 42.62°C (SD=1.68°C, maximum 45.36°C, minimum
39.99°C) and mean thermal window (Tc-Tb) of possible germination found to be 28°C
(SD=2.65°C, maximum 32.72°C, minimum 23.73°C), and all were significant at p<0.001.

Estimated values for cardinal temperatures are shown in **Table** 2, and curve for
germination rate and cardinal temperatures for accession GB31 in **Figure** 3. A full set of
results for both temperature and water potential experiments are given in **Figure** S2, and
germination rates and proportions for all seed accessions under heat conditions are
shown in **Figure** S3.

Accession	Country	Tb	To	Tc
1	Burkina Faso	11.91	36	41.96
2	Burkina Faso	14.89	30	45.36
3	Burkina Faso	14.61	31	40.04
4	Mali	11.72	36	44.44
5	Mali	13.96	38	41.95
7	Mali	14.13	36	44.98
9	Mali	12.83	37	44.61
10	Togo	14.22	35	45.01
11	Togo	12.97	38	41.45
12	Togo	13.24	37	42.21
13	Guinea	12.41	37	41.11
14	Guinea	17.65	38	43.03
15	Guinea	13.49	38	42.01
16	Guinea	13.09	36	45.08
17	Guinea	13.96	36	43.51
18	Guinea	14.70	36	43.68
19	Guinea	14.15	38	41.75
Average		14	36	43

Table 2 – Cardinal temperatures of seed accessions tested as part of germination trials, including base (Tb), optimum (To), and ceiling (Tc) temperatures calculated at t50 (time taken to achieve 50% germination), given in °C. Accessions shown are from the IRD seedbank collection, included as part of an international group also used for statistical analysis against climatic variables from locality of origin.

Figure 3 – Modelled germination rate for accession 31 from Guinea, under different temperature conditions, at t50. Rate is given as a log-adjusted value based on average time taken for all seeds in a Petri-dish to achieve 50% germination. Dashed red lines show viable germination limits at base (Tb) and ceiling (Tc) temperatures, and blue line shows predicted optimum temperature (To).

Correlations between average climate data associated with accession collection localities (maximum temperature, minimum temperature, and average precipitation) and the climate groupings assigned to each accession (hot and arid vs. wet and sub-tropical) are significant, and the relationship between climate (arid Burkina Faso and Mali grouped together versus Guinea and Togo) and seed germination percentage, at the temperatures of 30°C and 40°C were all found to be significant (p=0.008 and p=0.0005, respectively). However, there is significantly unbalanced variance within groups (Bartlett test p<0.005, F=0.298, F=0.254, and F=0.149. This is shown in **Figure** S4, with a full table of statistical correlation between all variables in **Figure** S5. There was no correlation between Tb or Tc with country or climate groups. There is a similarly minor significant relationship between To and both country and climate group, (p=0.0009284 and p=2.088e-09, respectively), though again there is significantly unbalanced variance between groups (Bartlett test p<0.005, F=0.132).

Response to drought

Germination synchronicity was significantly affected by water potential (p<0.001). Water potential (WP) had a significant effect on seed germination percentage (p<0.001), using a one-way ANOVA linear model. The mean highest successful germination percentage at 30°C was at 0MPa (77.1%), followed by -0.8MPa (74.8%), -1MPa (38.4%), -1.2MPa (10.5%), and -1.4MPa (1.25%). Mean germination time was also significantly affected by WP (p<0.001), where again the fastest mean time to germination was found at 0MPa (1.09), followed by -0.8MPa (1.33), -1MPa (1.80), -1.2MPa (2.12), and -1.4MPa (3.16). At 20°C the mean seed germination was highest again at 0MPa (75.6%), followed by -0.8MPa (5.05%). There was no germination at -1MPa to -1.4MPa. Germination time was also lowest at 0MPa (1.71) versus at -0.8MPa (3.64).

The minimum water potential required for germination (Ψb), by rate, could only be modelled for 3 accessions to t50 at 30°C. Average Ψb was -1.09MPa, with a minimum of -1.22MPa (accession 11 from Togo), and a maximum of -1.01MPa (accession 32 from Guinea). Germination rate curves for water potential experiments are shown in **Figure** S6. Germination at 20°C could only be modelled using t15, at which the mean Ψb is -0.8MPa, from three accessions.

Seedling phenotypes of stress

Temperature had a significant effect on seedling weight (p<0.001), the highest mean weight was found at 30°C (2.81mg), followed by 20°C (1.41mg), and 40°C (1.11mg). This was independent of accession (p<0.001). Water potential also had a significant effect on seedling weight (p<0.001), with the highest mean weight of 1.87mg at 0MPa and 0.4mg at -1MPa, also unaffected by accession (p<0.001). Mean weights of leaves are shown in **Figure** 2.

Figure 4 – Seedling growth phenotypes of seed accession 1 from Burkina Faso, 7 days after sowing, germinated under different heat (A) and water potential (B) conditions. Germinated seed can be seen at -1.4MPa in B, second seed from the right in second row from the top.

386 Observable phenotypes of temperature conditions at 7 days after sowing include:
387 undeveloped roots and shoots with no clear coleoptile expansion at 15°C; coiled, non-
388 branching roots with a single leaf at 20-25°C; roots with multiple branchings and multiple
389 leaves at 30-35°C; and long unbranched roots, elongated and pallid leaves at 40°C.
390 Phenotypes of temperature and drought stress shown in **Figure** 4.

Figure 5 – Current and future average maximum temperatures across West Africa, in the fonio sowing months from May to August. Future predictions are shown for 4 future SSP scenarios. Black circles show fonio occurrences recorded on GBIF (https://www.gbif.org/species/5289953). Areas are coloured by maximum temperature range from yellow to dark red – areas in red and dark red (Temperatures >35°C are above the average optimum fonio germination temperature.

Germination temperature thresholds vs. future temperatures

Historic (1970-2000) and future (2024-2060) projected maximum temperature values across West Africa for the main fonio sowing period (May-August) are shown in **Figure** 5. These include future projections under 4 different scenarios, and temperature categorised to match seed germination testing thresholds. The threshold of above 35°C, corresponding to average To, is chosen as unsuitable for consistent successful germination, based on seed germination results and seedling phenotypes described above. Large parts of Senegal, Burkina Faso, and Mali experience average maximum temperature of between 35-45°C in May and June in current climates – in future climates this range increases southward. In July, only fonio cultivated to the northern most extent of range (in Mali) experience temperatures of 35-45°C in the present, but this range extends to cover large parts of cultivation in Mali and Burkina Faso in the future, especially under SSP585. August remains within the window of likely germinable temperatures in current and all future scenarios. When occurrence points are matched to maximum temperatures under each current and future scenario, out of 1609 data points, and averaged across all 4 months in current conditions, 88.3% of occurrences are within a suitable temperature range. This decreases to 79.86% to 79.08% under SSP126, SSP245, and SSP370, and down to 71.05% for SSP585. This represents a decrease of 10.45% between current and future (SSP126-370) scenarios - shown in **Figure** 6, and separated by month in **Figure** S7.

Figure 6 – Suitability of fonio cultivation in current and future climate scenarios, calculated by two methods: A. occurrences in geographic areas below the optimal temperature threshold of 35°C; and B. niche suitability modelling. Arrows and labels show the decrease in mean suitable occurrences and area between current and future scenarios.

Figure 7 – Suitable area for fonio cultivation under current and future projections, calculated using environmental niche modelling. Fonio occurrences from GBIF are shown as circles in green. Area in blue shows unsuitable area in present, purple shows suitable area both in the current and future, red shows suitable area missing in future projections, and pink shows newly suitable area.

Niche suitability modelling

Niche suitability climate maps were produced under current and 3 future (SSP126, SSP245, SSP370) scenarios. Based on 30% test data and a minimum training threshold, the omission rate was 0 (binomial test p<0.050. AUC based on 30% removal of test data = 0.905. The minimum training threshold is 0.041, and the 10th percentile training presence is 0.272, equal training sensitivity and specificity is 0.389. Fractional predicted area at the 10th percentile training presence is 0.239 (24%). All of these are significant to p<0.0001. The highest percentage contribution of bioclimatic values to the model was temperature isothermality (BIO4, 58%), followed by precipitation seasonality (BIO15, 13.3%), precipitation of driest month (BIO14, 9.4%) and mean temperature of driest quarter (BIO9, 7.2%). The lowest contributor was mean temperature of warmest quarter (BIO10, 1.2%).

Suitability maps for fonio are shown in **Figure** 7. In current and recent historic conditions, 25.9% of the whole region shown is predicted to be suitable for fonio cultivation. This area covers nearly all fonio occurrences on the map, including Senegal and Nigeria, which are known to cultivate fonio, but no available occurrence points could be used for this study. This suggests an accurate modelling of crop suitability in the current climate. Under predictions in SSP126 the percentage of suitable area decreases to 21.3%, to 19.4% at SSP245, and to 18.8% at SSP370. This represents a range of between 6.3% to 9.1% reduction between current to future, and an average of 7.9%, shown in **Figure** 6. Based on the 10min resolution raster files used, this represents around 400,000km^2 (average of 1190 squares in raster layer missing in future) of land. There is also an increase of some areas newly suitable for fonio, especially in Sierra Leone, Cote D'Ivoire, Ghana, and Nigeria, as the southern regions of West Africa experience higher average temperatures – a total average increase in new areas of 5.5% (around 250,000km^2).

Discussion

Germination responses to climate change

The germinable temperature thresholds of *Digitaria exilis* were found to be between an average Tb of 14.52°C (observed down to 11.72°C) up to an average To of 36°C. Germination rate models estimated the highest Tc to be 45.36°C, though no seed tested in germination trials was able to germinate successfully at 45°C. This corroborates the results of Portères (1955), whose germination experiments suggested that fonio has a Tc of below 45°C, and a To of above 30°C.

The effects of seed age (collected in1977, 1978, 1981 and 1982 vs. 2022), and country and climate of origin (Mali and Burkina Faso vs. Guinea and Togo), appeared to be negligible. In traditional storage, the dried grains are reported to be germinable for several years after harvest (Cruz et al. 2016), but it appears that there is little to no difference in

germination success between seeds harvested, dried and stored in the 1970s, and those stored and dried 50 years later, and no observable dormancy. Many orthodox plant seeds remain germinable after long periods, although it is uncommon to not see a significant decrease in germination success after such a long period of time (Walters et al., 2005; Kameswara Rao et al., 2017; De Vitis et al., 2020). This relationship between seed age and germinability is obviously very dependent on storage methods, and highlights the positive role of seed-banking in preserving valuable indigenous crop seeds for growth in the future.

Although country and climate groupings had a minor statistically significant relationship with seed germination proportion (grp) and optimal temperature (To), there was disproportionate variation between groups, where seeds from arid environments (Mali and Burkina Faso) had a more varied germination response to higher temperatures, and actually fail to outperform seeds from cooler climates found in Guinea and Togo. There appears to be no definite difference between germination responses between the two groups that would suggest strong adaptations to local climates. Generally, there is little variation in germination response to high heat among varieties of fonio, even those from vastly different geographic areas. This may be due to our experimental conditions and climate variables not taking into account the specific microclimates that older generations of seed experienced, or that sample size was not large enough to capture any strong, consistent difference between groups.

It is also known that fonio varieties are often transported to different regions through trade to neighbouring villages and to different habitats (upland slopes, lowland grasslands; Burton et al., 2024; Cruz et al., 2016). This would suggest a crop that is frequently moved between different nearby microclimates, possibly explaining why a significant difference in origin climate-specific heat tolerance was detected in this study. However, it is more likely that fonio is hitting an absolute limit with germination temperature, where adaptation is no longer feasible due to physiological constraints. In a recent study by Pudasaini et al. (2025), two fonio genotypes, one each from Mali and Guinea, were compared for differences in drought response in whole mature plants. While they found that there was strong difference in root architecture and yield (the variety from Mali outperforming the variety from Guinea), they make no comment on any differences in the overall germination success or early seedling development of the two varieties. This again supports that germination vigour under heat stress is generally high across white fonio varieties, but also reaching a set physiological limit of around 42°C. This high resilience is possibly a trait conserved from the likely origins of *Digitaria* alongside the subtribe Anthephorinae and other Panicoid C4 grasses in arid central and East African grasslands (Gallaher et al., 2022; Bouchenak-Khelladi et al., 2010).

Suitability of fonio in future climates

From the germination thresholds and phenotypes described above, it is sensible to predict that fonio being sown in any areas that experience consistent average maximum temperatures above 35°C for prolonged periods is likely to experience a decrease in plant health (at 40°C) and germination success and survival (above 40°C), resulting in decreased grain yield and food availability.

Analysis of seed germination results against future climate predictions above utilises a comparison of realistic thresholds for successful fonio seed germination (below 35°C) compared to current and future maximum temperatures across West Africa. In the current climate, very few occurrences appear within the regions which experience >35°C temperatures in July and August, apart from 3 occurrences in the area on the border between the Mopti and Tombouctou regions in Mali, and Burkina Faso. In May and June large areas of the upper Sahel are unsuitable, but most areas of Guinea are suitable throughout the whole season. This supports why farmers in Guinea described how early varieties of fonio may be sown as early as May in a study by Burton et al. (2024), and there has recently been a shift to sowing later seasons, and abandonment of early varieties. Meanwhile, Mali, Senegal, and Burkina Faso appear suitable for these earlier varieties. A growth study by Gueye et al. (2015) with fonio in south-eastern Senegal, suggests that early July had the best sowing dates for plant growth and grain yield, as it avoided the intense heat of May and June, and avoided rains in August. This fully corroborates with our results shown here, and suggests that in the future, farmers across West Africa, especially in arid areas, are likely to have to shift their sowing and harvest calendars to better suit the heat tolerance of their crop seeds to more suitable later months.

Between the current and future climate scenarios, using results from seed germination trials against future climates, around 10% of suitable land for fonio cultivation is likely be lost. The months May and June experience far more severe reductions than July and August, although in some scenarios July is also affected. Key cultivation regions in Senegal, Mali, and Burkina Faso appear to be affected the most, where some areas experience temperatures above 35°C more frequently. These results support those from the niche suitability model predictions: reduction in suitable niche area for fonio cultivation under these models predict a reduction of around 8% across the region. The area in Mali that appears to be currently unsuitable with the 3 occurrences in the first method, are also unsuitable in the current climate under these predictions. Even under the most optimistic scenario SSP126, there is a significant reduction in unsuitable niche area of around 6%, where the model entirely removes Senegal, and large parts of Mali and Burkina Faso from the suitable land area. This only degrades further as the SSP scenarios become decreasingly optimistic. Crop suitability models produced by Chemura et al. (2024) similarly found that fonio by 2050 will experience a 2-7% reduction in suitable area under the different SSP scenarios – though this was calculated to the scale of the whole African continent, their predictions also suggest that the upper Sahel

countries will experience the most severe reductions. A review study on the effect of
climate change on crop yields in West Africa found a median yield loss of ~11% in the
future, which was far more impactful in northern West Africa (including Mali and Burkina
Faso) vs. southern regions of Guinea and Ghana (Roudier et al., 2011). The results in this
study for fonio then of around 10% decrease in the future is similar to other studies,
tested through multiple different methods. This reduction of suitable area for fonio,
coupled with low specific local adaptation for germinating in extreme arid climates, is
worrying, considering how viable germination already appears to have reached a
consistent limit of around 42°C in seed accessions across the cultivation range.

Three key areas that appear to be affected in both methods are south-eastern Senegal,
the south-eastern Mali:Burkina Faso border, and north-eastern Benin, areas shown in
Figure S8. These areas are historically very important for the cultivation of fonio, with
records of fonio in this area of Benin as far back as 900-1400AD, and possibly having been
domesticated near the Mopti region in Mali before 500BC (Champion and Fuller, 2018).
In a study by Porcuna-Ferrer et al. (2024) on the south-eastern fonio cultivation by Bassari
farmers in Senegal, they reported that a mix of socio-political issues and policies
promoting non-indigenous crop species over fonio and Bambara groundnut, causing a
decline in fonio production of up to 40%. Around 9% of interviewees cited environmental
factors as the reason for abandonment, suggesting that cultivation in currently climate-
suitable regions across West Africa will and do now experience abandonment – the
effects of climate change are also coupled with and intensified by social attitudes of
governments and local communities.

The climate change between present and future scenarios are also beneficial in some
areas: in Sierra Leone, Cote D'Ivoire, Ghana, and Nigeria, suitable land for fonio is
predicted to expand by around 5.5%. A study by Wimalasiri et al. (2023) investigated a
similar response for proso millet (*Panicum miliaceum* L.), using the 2024-2069
predictions from the C3MP/CMIP5, with increased yield under future rainfall increases,
alongside a 2°C temperature increase in the dry zone of Sri Lanka. Chemura et al. (2024)
while predicting an overall decrease in suitable land for fonio, predicted a positive
increase in suitable area for similar millet species sorghum and pearl millet, and also
Bambara groundnut, by up to 5% by 2050.

Conclusions

This study provides critical insights into the climatic resilience of fonio. We have
established that the physiological growth limit of fonio is up to a likely absolute maximum
of 43°C, with an average optimum temperature of around 35°C based on extensive
germination experiments involving 37 seed accessions collected from across multiple
habitats and climates in West Africa. There appears to be no straightforward relationship
between heat resilience in germination trials and the seed accessions' local climate,

suggesting a low selection for heat-resilient varieties, or local adaptation to the climate,
and instead a more general high tolerance for heat for the species.

These results highlight both opportunities and risk for fonio as an important indigenous
crop. When comparing temperature suitability thresholds to current and future climates
using two different methods, we find that over the next 40 years, fonio cultivation in
regions close to the Sahel in Senegal, Benin, Mali, and Burkina Faso, are very likely to be
adversely impacted by climate change. These are some of the most important current
and historical areas for fonio cultivation, both socially and economically. However, many
other areas will remain or transition into the optimal climate window for cultivation,
including Sierra Leone, Cote D'Ivoire, and Ghana. While fonio remains an essential
component of food security in West Africa, the increasing threat of climate change
necessitates improved adaptive agronomic strategies, such as shifting sowing times to
avoid extreme heat and weather events, accelerated selection and breeding for more
heat-tolerant varieties, and expanding cultivation into newly suitable areas.

From this study, future research and policy may be implemented to target conservation
for the most under-threat areas, and promote fonio in regions where it is still vitally
suitable and sustainable, where other crops are not. Policies that promote the
conservation and commercialisation of neglected crops like fonio will be crucial in
supporting food security in climate-vulnerable regions of the Sahel. These include policy
initiatives like the Comprehensive Africa Agriculture Development Programme (CAADP,
2025) administered by the African Union, which aims to end hunger (Malabo
Commitment 3), and enhance resilience of livelihoods and production systems to
climate variability (MC6). Our research provides a further opportunity to conduct similar
studies on other dryland NUS crops (neglected and underutilised, Ulian et al. 2020) in
order to inform agricultural development policies, programmes, and projects on the
basis of comparative advantages and disadvantages of climate-resilient crops.

Supplementary Data

Figure S1. Effect of date collected and seedbank collection group on germination
proportion.

Figure S2. Cardinal temperatures (optimum temperature, ceiling temperature, base
temperature, and thermal window at t50) and base water potentials (Psib) at 30°C (t50)
and 20°C (t15) for fonio seed accessions.

Figure S3. Modelled germination rate curves and proportions (boxplots) for all
accessions used in temperature experiments.

Figure S4. Effect of seed accession origin climate (Guinea and Togo vs. Mali and Burkina
Faso) on germination proportion at 30°C, 35°C, and 40°C.

Figure S5. Correlation of variables between seed germination proportions, cardinal
temperatures, and climatic variables. Significant p values are in bold.

Figure S6. Modelled germination rate of three fonio accessions under a gradient of water
potential conditions, at threshold t50.

Figure S7. Percentage of fonio occurrence points mapped to maximum temperatures
above 35°C, in the current and under future climate SSP scenarios, grouped by month.

Figure S8. Key areas predicted to experience significant reduction in suitable land for
fonio cultivation.

Acknowledgements

We are extremely grateful to the staff and students at the Herbier National de Guinée
including director Sékou Magassouba, and Kew Science liaisons Charlotte Couch and
Martin Cheek for helping with fieldwork to collect fonio seed material. We thank Adeline
Barnaud at the IRD and Claire Billot at CIRAD (both Montpellier, France), for their valuable
collaboration and for providing seed material from the IRD for our study. We also thank
the curation, research and laboratory teams at the Millenium Seed Bank (UK), including
Pablo Gomez Barreiro, Silvia Bacci, and Bendetta Gori for assisting with germination
experiments. Thank you to Owen Corbett for guidance on producing suitability models in
ArcGIS. This study is a part of G. Burton's PhD project, and we are thankful for funding and
administrative help provided by Christiane Morgan at the Grantham Institute (Imperial
College London), and Felix Forest at Kew Science.

Author Contributions

GPB-TU: conceptualisation; GPB, HMB, CES, MSV, EM, and TU: methodology; GPB, CES,
EM: data analysis; GPB-TU: writing; PC, RMG, PR, MSV, EM, and TU: supervision; GPB,
MSV, and TU: funding acquisition.

Conflicts of Interest

We declare no conflicts of interest were found for any of the above authors.

Funding Statement

This work was supported by United Kingdom Research and Innovation (UKRI) and the
Natural Environment Research Council (NERC), administered through the Science and
Solutions for a Changing Planet DTP at the Grantham Institute and Life Sciences
department (South Kensington) at Imperial College London, and at Kew Gardens,
Richmond.

Data Availability

Data from this study will be deposited to Dryad online repository after review.

References

Abrouk M, Ahmed HI, Cubry P, Šimoníková D, Cauet S, Pailles Y, Bettgenhaeuser J, Gapa L, Scarcelli N, Couderc M, et al. 2020. Fonio millet genome unlocks African orphan crop diversity for agriculture in a changing climate. *Nature Communications* **11**: 4488.

Adoukonou-Sagbadja H, Wagner C, Dansi A, Ahlemeyer J, Daïnou O, Akpagana K, Ordon F, Friedt W. 2007. Genetic diversity and population differentiation of traditional fonio millet (Digitaria spp.) landraces from different agro-ecological zones of West Africa. *Theoretical and Applied Genetics* **115**: 917–931.

Bouchenak-Khelladi Y, Verboom GA, Savolainen V, Hodkinson TR. 2010. Biogeography of the grasses (Poaceae): a phylogenetic approach to reveal evolutionary history in geographical space and geological time: ANCESTRAL BIOGEOGRAPHY AND ECOLOGY OF GRASSES. *Botanical Journal of the Linnean Society* **162**: 543–557.

Bradford KJ. 2002. Applications of hydrothermal time to quantifying and modeling seed germination and dormancy. *Weed Science* **50**: 248–260.

Burton G, Gori B, Camara S, Ceci P, Conde N, Couch C, Magassouba S, Vorontsova MS, Ulian T, Ryan P. 2024. Landrace diversity and heritage of the indigenous millet crop fonio (*DIGITARIA EXILIS*): Socio-cultural and climatic drivers of change in the Fouta Djallon region of Guinea. *PLANTS, PEOPLE, PLANET*: ppp3.10490.

CAADP. Comprehensive Africa Agricultural Development Programme. 2025. https://caadp.org/. Accessed on 09/03/25.

Catara S, Cristaudo A, Gualtieri A, Galesi R, Impelluso C, Onofri A. 2016. Threshold temperatures for seed germination in nine species of *Verbascum* (Scrophulariaceae). *Seed Science Research* **26**: 30–46.

Champion L, Fuller DQ. 2018. New Evidence on the Development of Millet and Rice Economies in the Niger River Basin: Archaeobotanical Results from Benin. In: Mercuri AM, D'Andrea AC, Fornaciari R, Höhn A, eds. Plants and People in the African Past. Cham: Springer International Publishing, 529–547.

Chemura A, Gleixner S, Gornott C. 2024. Dataset of the suitability of major food crops in Africa under climate change. *Scientific Data* **11**: 294.

Covell S, Ellis RH, Roberts EH, Summerfield RJ. 1986. The Influence of Temperature on Seed Germination Rate in Grain Legumes: I. A COMPARISON OF CHICKPEA, LENTIL, SOYABEAN AND COWPEA AT CONSTANT TEMPERATURES. *Journal of Experimental Botany* **37**: 705–715.

Cruz J-F, Béavogui F, Dramé D, Diallo TA. **2016**. *Fonio, an African cereal*. Montpellier] [Conakry (Guinée): Cirad [UMR Qualisud] IRAG, Institut de recherche agronomique de Guinée.

De Vitis M, Hay FR, Dickie JB, Trivedi C, Choi J, Fiegener R. **2020**. Seed storage: maintaining seed viability and vigor for restoration use. *Restoration Ecology* **28**.

Ellis RH, Covell S, Roberts EH, Summerfield RJ. **1986**. The Influence of Temperature on Seed Germination Rate in Grain Legumes: II. INTRASPECIFIC VARIATION IN CHICKPEA (*Cicer arietinum* L.) AT CONSTANT TEMPERATURES. *Journal of Experimental Botany* **37**: 1503–1515.

Evans JS, Murphy MA, Holden ZA, Cushman SA. **2010**. Modeling Species Distribution and Change Using Random Forest. In: Drew CA, Wiersma YF, Huettmann F, eds. Predictive Species and Habitat Modeling in Landscape Ecology. New York, NY: Springer New York, 139–159.

Fatima Z, Ahmed M, Hussain M, Abbas G, Ul-Allah S, Ahmad S, Ahmed N, Ali MA, Sarwar G, Haque EU, *et al.* **2020**. The fingerprints of climate warming on cereal crops phenology and adaptation options. *Scientific Reports* **10**: 18013.

Fuller DQ, Champion L, Cobo Castillo C, Den Hollander A. **2024**. Cotton and post-Neolithic investment agriculture in tropical Asia and Africa, with two routes to West Africa. *Journal of Archaeological Science: Reports* **57**: 104649.

Fuller DQ, Stevens CJ. **2018**. Sorghum Domestication and Diversification: A Current Archaeobotanical Perspective | SpringerLink.

Gallaher TJ, Peterson PM, Soreng RJ, Zuloaga FO, Li D, Clark LG, Tyrrell CD, Welker CAD, Kellogg EA, Teisher JK. **2022**. Grasses through space and time: An overview of the biogeographical and macroevolutionary history of Poaceae. *Journal of Systematics and Evolution* **60**: 522–569.

Garcia-Huidobro J, Monteith JL, Squire GR. **1982**. Time, Temperature and Germination of Pearl Millet (*Pennisetum typhoides* S. & H.): I. CONSTANT TEMPERATURE. *Journal of Experimental Botany* **33**: 288–296.

Gianinetti A. **2020**. Basic Features of the Analysis of Germination Data with Generalized Linear Mixed Models. *Data* **5**: 6.

Gueye M, Kanfany G, Fofana A, Gueye M, Noba K, Grove J. **2015**. Effect of planting date on growth and grain yield of fonio millet (*Digitaria exilis* Stapf) in the Southeast of Senegal. *International Journal of Biological and Chemical Sciences* **9**: 581.

Gummerson RJ. **1986**. The Effect of Constant Temperatures and Osmotic Potentials on the Germination of Sugar Beet. *Journal of Experimental Botany* **37**: 729–741.

Hardegree SP, Emmerich WE. **1990**. Effect of Polyethylene Glycol Exclusion on the Water Potential of Solution-Saturated Filter Paper. *Plant Physiology* **92**: 462–466.

Heckman RW, Pereira CG, Aspinwall MJ, Juenger TE. **2024**. Physiological Responses of C4 Perennial Bioenergy Grasses to Climate Change: Causes, Consequences, and Constraints. *Annual Review of Plant Biology* **75**: 737–769.

Iwuala E, Unung, O., Abiodun, I. and Aneke, T. 2021. Responses of Digitaria iburu Stapf. (Black Fonio) Seedlings to Simulated Drought Condition.

Jadav SK, Bardhan K, Chaudhary K, Narwade AV. **2016**. Comparative Effect of PEG-8000 and NaCl Induced Stress on Tomato at Germination.

Kameswara Rao N, Dulloo ME, Engels JMM. **2017**. A review of factors that influence the production of quality seed for long-term conservation in genebanks. *Genetic Resources and Crop Evolution* **64**: 1061–1074.

Kriegler E, O'Neill BC, Hallegatte S, Kram T, Lempert RJ, Moss RH, Wilbanks T. **2012**. The need for and use of socio-economic scenarios for climate change analysis: A new approach based on shared socio-economic pathways. *Global Environmental Change* **22**: 807–822.

Linares OF. **2002**. African rice (*Oryza glaberrima*): History and future potential. *Proceedings of the National Academy of Sciences* **99**: 16360–16365.

Lozano-Isla F, Benites-Alfaro OE, Pompelli MF. **2019**. GerminaR: An R package for germination analysis with the interactive web application "GerminaQuant for R". *Ecological Research* **34**: 339–346.

Meinshausen M, Smith SJ, Calvin K, Daniel JS, Kainuma MLT, Lamarque J-F, Matsumoto K, Montzka SA, Raper SCB, Riahi K, *et al*. 2011. The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climatic Change* **109**: 213–241.

Merow C, Smith MJ, Silander JA. **2013**. A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* **36**: 1058–1069.

Michel BE. **1983**. Evaluation of the Water Potentials of Solutions of Polyethylene Glycol 8000 Both in the Absence and Presence of Other Solutes. *Plant Physiology* **72**: 66–70.

Onofri A, Benincasa P, Mesgaran MB, Ritz C. **2018**. Hydrothermal-time-to-event models for seed germination. *European Journal of Agronomy* **101**: 129–139.

Onofri A, Mesgaran MB, Ritz C. **2022**. A unified framework for the analysis of germination, emergence, and other time-to-event data in weed science. *Weed Science* **70**: 259–271.

Pearson RG, Dawson TP. **2003**. Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global Ecology and Biogeography* **12**: 361–371.

Phillips SJ, Dudík M. **2008**. Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* **31**: 161–175.

Pirnajmedin F, Majidi MM, Jaškūnė K. **2024**. Adaptive strategies to drought stress in
grasses of the poaceae family under climate change: Physiological, genetic and
molecular perspectives: A review. *Plant Physiology and Biochemistry* **213**:
108814.

Porcuna-Ferrer A, Calvet-Mir L, Faye NF, Klappoth B, Reyes-García V, Labeyrie V.
2024. Drought-tolerant indigenous crops decline in the face of climate change: A
political agroecology account from south-eastern Senegal. *Journal of Rural
Studies* **105**: 103163.

Portères R. **1955**. Les Céréales mineures du genre Digitaria en Afrique et en Europe.
Journal d'agriculture tropicale et de botanique appliquée **2**: 349–386.

Pudasaini R, Amyotte J, Brettingham DJL, Raizada MN. **2025**. Physiological,
metabolomic, morphological and root system architecture acclimation
responses to drought in the African orphan millet white fonio (Digitaria exilis).
Plant and Soil.

**Ragatoa DS, Amekudzi LK, Fink AH, Maranan M, Klutse NAB, Edjame KS, Ogunjobi
KO**. **2024**. Droughts and heatwaves in the West African monsoon system.
International Journal of Climatology **44**: 3681–3705.

Reed RC, Bradford KJ, Khanday I. **2022**. Seed germination and vigor: ensuring crop
sustainability in a changing climate. *Heredity* **128**: 450–459.

Roudier P, Sultan B, Quirion P, Berg A. **2011**. The impact of future climate change on
West African crop yields: What does the recent literature say? *Global
Environmental Change* **21**: 1073–1083.

Sutherst RW, Maywald, G. F., Yonow, T., and Stevens, P. M. 1999. Predicting the
Effects of Climate on Plants and Animals. CD-ROM andUser Guide.

Sylla MB, Pal JS, Faye A, Dimobe K, Kunstmann H. **2018**. Climate change to severely
impact West African basin scale irrigation in 2 °C and 1.5 °C global warming
scenarios. *Scientific Reports* **8**: 14395.

Tadele Z. **2016**. Drought Adaptation in Millets. In: Shanker AK, Shanker C, eds. Abiotic
and Biotic Stress in Plants - Recent Advances and Future Perspectives. InTech.

Trudgill DL, Squire GR, Thompson K. **2000**. A thermal time basis for comparing the
germination requirements of some British herbaceous plants. *New Phytologist*
145: 107–114.

**Ulian T, Diazgranados M, Pironon S, Padulosi S, Liu U, Davies L, Howes MR, Borrell
JS, Ondo I, Pérez-Escobar OA, *et al***. **2020**. Unlocking plant resources to
support food security and promote sustainable agriculture. *PLANTS, PEOPLE,
PLANET* **2**: 421–445.

**Van Vuuren DP, Riahi K, Calvin K, Dellink R, Emmerling J, Fujimori S, Kc S, Kriegler
E, O'Neill B**. **2017**. The Shared Socio-economic Pathways: Trajectories for
human development and global environmental change. *Global Environmental
Change* **42**: 148–152.

Violita V, Azhari S. **2021**. Effect of PEG-8000 imposed drought stress on rice varieties
germination. *Journal of Physics: Conference Series* **1940**: 012071.

Wahid A, Gelani S, Ashraf M, Foolad M. **2007**. Heat tolerance in plants: An overview.
Environmental and Experimental Botany **61**: 199–223.

Walters C, Wheeler LM, Grotenhuis JM. **2005**. Longevity of seeds stored in a
genebank: species characteristics. *Seed Science Research* **15**: 1–20.

Watt MS, Bloomberg M. **2012**. Key features of the seed germination response to high
temperatures. *New Phytologist* **196**: 332–336.

Wimalasiri EM, Ashfold MJ, Jahanshiri E, Walker S, Azam-Ali SN, Karunaratne AS.
2023. Agro-climatic sensitivity analysis for sustainable crop diversification; the
case of Proso millet (Panicum miliaceum L.) (MM Hamed, Ed.). *PLOS ONE* **18**:
e0283298.

World Weather Attribution. 2024, April. Extreme Sahel heatwave that hit highly
vulnerable population at the end of Ramadan would not have occurred without
climate change – World Weather Attribution.
https://www.worldweatherattribution.org/extreme-sahel-heatwave-that-hit-
highly-vulnerable-population-at-the-end-of-ramadan-would-not-have-occurred-
without-climate-change/
World Weather Attribution. 2024, October. Conflict, poverty and water management
issues exposing vulnerable communities in Africa to extreme floods that are now
common events because of climate change – World Weather Attribution.
https://www.worldweatherattribution.org/conflict-poverty-and-water-
management-issues-exposing-vulnerable-communities-in-africa-to-extreme-
floods-that-are-now-common-events-because-of-climate-change/

820