HaploCharmer: a Snakemake workflow for read-scale haplotype calling adapted to polyploids

Simon Rio [1,2], Sophie Abdallah[1,2], Théo Durand[1,2], Angélique D'Hont[1,2], and Olivier Garsmeur[1,2]

Abstract

The advent of next-generation sequencing (NGS) has revolutionized the study of single nucleotide polymorphisms (SNPs), making it increasingly cost-effective. Haplotypes, which combine alleles from adjacent variants, offer several advantages over bi-allelic SNPs, including enhanced information content, reduced dimensionality, and improved statistical power in genomic studies. These benefits are particularly significant for polyploid species, where distinguishing all homologous copies using SNP markers alone can be challenging. This article introduces HaploCharmer, a flexible workflow designed for read-scale haplotype calling from NGS data. HaploCharmer identifies haplotypes within preconfigured genomic regions smaller than a sequencing read, ensuring direct comparability across individuals. It integrates a series of processing steps including mapping, haplotype identification, filtration, and reporting of haplotype sequences, as presence-absence, in the panel of accessions analyzed. The performance of HaploCharmer was validated using whole-genome sequencing data from a highly polyploid sugarcane cultivar (R570) and its self-progeny. The workflow successfully identified a large number of high-quality haplotypes, with less than 1% of false positives. Single-dose haplotypes were used to construct a genetic map that accurately included known chromosomal rearrangements in the R570 cultivar, demonstrating its effectiveness in studying large chromosome structural variations. HaploCharmer provides a robust method for diversity, genetic mapping, and quantitative genetics studies in both diploid and polyploid species.

Keywords: Haplotype, Next-generation sequencing, Polyploid, Sugarcane

[1]CIRAD, UMR AGAP Institut, F-34398 Montpellier, France, [2]UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

Correspondence
simon.rio@cirad.fr

Introduction

Because of their convenience in subsequent analyses, most variant call methods from next-generation sequencing (NGS) data aim to characterize bi-allelic single nucleotide polymorphisms (SNPs). Alternatively, alleles from adjacent variants, including SNPs and other polymorphisms such as indels, can be combined into haplotypes. The advantages of haplotypes over bi-allelic SNPs include an increased information content, reduced dimensionality, better capture of linkage disequilibrium, higher statistical power in genome-wide association studies, higher genomic prediction accuracy, better genetic mapping resolution and improved population structure inference (Bhat et al., 2021; Gattepaille and Jakobsson, 2012). The multi-allelic nature of haplotypic markers is expected to be particularly useful in polyploid species, where all homologous copies cannot be distinguished by a single bi-allelic variant, unlike diploid species (Voorrips and Tumino, 2022).

Using NGS data, alleles from adjacent variants can be combined into haplotypes when they are observed on the same sequencing read. This strategy is implemented in HAPLOSWEEP (Clevenger et al., 2018), but is limited to phasing pairs of variants for polyploid species. In addition, haplotypes can be extended by assembling reads on the basis of their shared allele content, which supports an origin from the same homologous copy. This task is more complex in polyploids for which the solution space increases dramatically with the number of variants and the ploidy level (He et al., 2018). Several haplotype assembling methods have been proposed in the literature, some of them specifically dedicated to polyploids, such as HapCompass (Aguiar and Istrail, 2012, 2013), Poly-Harsh (He et al., 2018), or Ranbow (Moeinzadeh et al., 2020). However, these approaches can lead to considerable heterogeneity in haplotype length between homologous copies of the same individual or between different individuals, depending on the degree of information available to guide assembly. As a consequence, the resulting haplotypes may not be easily transformed into multi-allelic markers to be used in genomic and genetic studies.

In this article, we present HaploCharmer, a workflow for read-scale haplotype calling implemented in Snakemake (Köster and Rahmann, 2012; Mölder et al., 2021). Haplotypes are called within preconfigured genomic regions of a size smaller than a sequencing read, and within which all possible variants are considered: bi- and multi-allelic SNPs as well as small indels. Like in HAPLOSWEEP, alleles are considered as phased only if they are jointly observed on the same sequencing read. The use of preconfigured genomic regions, within which haplotypes are characterized, ensures that the resulting haplotypes are directly comparable across individuals. For whole-genome sequencing data, genomic regions can be randomly defined in non-repetitive sequences or in sequences of putative interest (e.g. exons or cis-regulatory elements). For genotyping-by-sequencing approaches, genomic regions can be defined in read-enriched regions.

HaplotyCharmer's effectiveness in identifying haplotypes in a highly polyploid and heterozygous genome was assessed by applying it to available resequencing data for a sugarcane cultivar (R570, $\approx$12x) and its self-progeny.

Materials and methods

HaploCharmer workflow

HaploCharmer takes as input single or paired-end short-read sequencing data (FASTQ) for a set of samples, which have passed quality controls such as adapter removal, read filtering and trimming of low-quality bases.

First, reads are mapped to a reference sequence genome (FASTA) using BWA-MEM (Li, 2013). Standard processing steps are applied to BAM files, such as removal of read duplicates using Picard MarkDuplicates (https://broadinstitute.github.io/picard/), local realignment around indels using GATK3 (DePristo et al., 2011), and update of the MD flag using SAMtools (Danecek et al., 2021). Haplotypes are then identified and counted within pre-configured genomic co-ordinates (BED) using a python script (BAM_to_gVCF.py) based on the pysam module (https://github.com/pysam-developers/pysam/). Only sequencing reads that perfectly overlap pre-defined genomic regions - also called phase sets (PSs) - are considered (Figure 1A). For each of the samples analyzed, all positions that fall into a PS are reported in a genomic VCF file (gVCF), regardless of whether a variant (SNP or indels) was detected at that site or not. In the gVCF file, haplotypes are reported as groups of alleles phased within each PS, which is specified following the convention of using the pipe "|" in the genotype (GT) field.

Individual sample files are merged into a global VCF file using BCFtools (Danecek et al., 2021), and filtered for haplotypes likely resulting from sequencing errors using a python script (VCF_filter.py). For this filtering step, the general approach is to discard haplotypes that are never observed with a high degree of certainty in at least one sample, i.e. with sufficient read depth and frequency to support the existence of the haplotype (see next subsection "HaploCharmer parameters"). Then, a non-standard file characterizing haplotype presence-absence (HPA) is generated using a python script (VCF_to_HapPresAbs.py) based on the pyfaidx module (Shirley et al., 2015). In this HPA file, each line corresponds to a haplotype and includes its coordinates on the reference sequence, its complete sequence, and sample-specific columns indicating its presence/absence genotyping along with read depth information (Figure 1B). In addition, the nature and position of variants (SNPs and indels) distinguishing the haplotypes of a same PS are reported in a text file (INFO). This file is made up of a concatenation of tables in VCF format, one for each PS, where haplotypes are represented in columns. Each sub-table corresponding to a specific PS can easily be accessed using a "grep" command with the name of the PS as the argument. Finally, a report (HTML) is generated presenting various statistics, such as read mapping, read duplicates, phase set read depths, and number of haplotypes per phase set. The workflow's key steps are summarized in Figure 1C.

HaploCharmer parameters

The haplotype calling relies on a limited set of parameters. Regarding the mapping, the removing of read duplicates is optional, although recommended, and a minimum mapping quality (i.e. MAPQ provided by BWA-MEM) must be specified. For the VCF filtering step, a genotyping data point must be supported by a read depth (DP) comprised between "min_gt_depth" (minimum read depth) and "max_gt_depth" (maximum read depth) otherwise the genotyping datapoint is set to a missing value. Then, each haplotype must be supported by a read depth (AD) greater than or equal to "min_hap_depth" and a frequency (AD/DP) greater than or equal to "min_hap_freq" in

Figure 1 – (A) Schematic representation of three aligned reads considered for haplotyping (green) that overlap the entire region (PS) defined by the START and END positions on the reference, unlike the other reads (grey). (B) Illustration of the haplotype representation in the VCF, HPA and INFO files for one sample genotyped for a haplotypic region "1_2_8", i.e. a phase set spanning chromosome 1 from position 2 to 8. For the HPA and INFO files, the haplotype "1_2_8_hap4" corresponds to the reference sequence haplotype. Only the genotype (GT) format field is represented here although it is normally associated with read depth information in real files. (C) Summary of the HaploCharmer (i) inputs including the reference sequence (FASTA), sequencing reads (FASTQ), PS coordinates on the reference sequence (BED), (ii) intermediate files including alignment files (BAM) and genomic VCF files (gVCF), (iii) and outputs including VCF, HPA and INFO files. Key workflow steps to obtain output from input files are described in orange rectangles.

at least one sample. Otherwise, this haplotype is considered as likely resulting from sequencing
errors and discarded from the set of haplotypes in the population.
Although a haplotype may be supported at a population level, it may be the result of sequenc-
ing errors for a particular sample. At the sample level, the haplotypes reported in the HPA file are
first re-inspected to verify they are supported by a read depth (AD) and frequency (AD/DP) su-
perior or equal to "min_pres_depth" and "min_pres_freq", respectively. Then, too rare haplotypes

with a frequency below "max_abs_freq", are considered absent in the sample. Finally, haplotypes

that do not meet any of these criteria (e.g. intermediate frequency: "max_abs_freq" $\leq$ AD/DP $<$

"min_pres_freq") are set as missing values.

The workflow parameters are explained in further details and illustrated with a toy exam-

ple in the README.md file of the Gitlab repository (see "Data, script, code, and supplementary

information availability" section).

Evaluation of HaploCharmer on polyploid sugarcane

The performance of HaploCharmer was assessed by calling haplotypes from paired-end 150

bp Illumina whole-genome sequencing data from sugarcane cultivar R570 ($\approx$12x) and 96 acces-

sions of its self-progeny (Grivet et al., 1994; Grivet et al., 1996). The identified haplotypes were

then characterized to select segregating single-dose haplotypes, which were used to construct

a genetic map.

Haplotyping of R570 and its self-progeny. The recent polyploid genome assembly of R570 (Healey

et al., 2024) was used as a sugarcane reference sequence. From this polyploid assembly, with

several hom(e)ologous copies assembled for each basic chromosome (i.e. hom(e)ology group), a

monoploid reference sequence was derived by choosing the longest scaffolds from each hom(e)ology

group, but excluding those representing rearranged chromosomes. The scaffolds considered

were the following: Chr1A, Chr2A, Chr3A, Chr4A, Chr5A, Chr6A, Chr7A, Chr8A, Chr9A, and

Chr10A.

Three sets of probes were used to define haplotypic regions: 60,000 120 bp probes defined

by Yang et al. (2018), 1,964 120 bp probes (non-redundant with the previous set) from the set

of 19,436 probes defined in Dijoux et al. (2024), and 39,867 additional 80 bp probes defined in

sugarcane gene models. All probes were mapped onto the monoploid reference genome using

BWA-MEM (Li, 2013). Different filters were applied, excluding probes that i) were not uniquely

mapped, ii) mapped on an interval below 60 bp or above 140 bp, and iii) overlapped the map-

ping coordinates of another adjacent probe. A total of 86,681 80 bp probes distributed along

the R570 monoploid genome assembly were retained and the corresponding regions were con-

sidered as PSs for the haplotype calling. The coordinates of these PSs were reported in a BED

file format.

To reduce the computational time required for aligning high-depth WGS data from cultivar

R570 and the 96 progenies, the reads corresponding to all regions were recovered through in-

silico capture using probe sequences as targets with Mirabait (Chevreux et al., 2004).

The HaploCharmer workflow was launched by specifying the following parameters in the

Snakemake YAML configuration file:

• Remove_duplicates: True

• Mapping_quality: 20

• VCF_filter: "–min_gt_depth 30 –max_gt_depth 1000 –min_hap_depth 4 –min_hap_freq

0.04"

• VCF_to_HapPresAbs: "–max_abs_freq 0.01 –min_pres_depth 3 –min_pres_freq 0.04"

Filtering of progenies. From the raw haplotypes identified and reported in the HPA output file,

progenies' genotypes (present, absent, or missing) were compared to that of R570 to potentially

identify illegitimate progeny (Figure 2). No progeny was found to have a substantial proportion

of haplotypes absent from R570, suggesting no illegitimate cross. However, for a self-progeny,

a substantial proportion of haplotypes is expected to be present in the parent but absent in
the progeny, notably resulting from the segregation of single-dose haplotypes (see subsection
"Characterization of single-dose markers"). Few progenies were discarded from the study as they
presented a genotyping profile too similar to R570, suggesting that there were clones of the par-
ent: AF411, AF413, AF439, AF454, AF491, AF499, and AF568. Two additional progenies were
discarded due to a significant percentage of missing values: AF450 and AF488. After filtering, a
total of 87 progenies were conserved for downstream analyses.

Figure 2 – Number of haplotypes called (y-axis) in each progeny (x-axis). Colors corre-
spond to distinct categories describing the presence, absence or missing information
(NA) in comparison with R570, with R570 on the left, and the progeny on the right. For
instance, "presence - absence" (in orange) corresponds to a haplotype present in R570
and absent in the progeny.

Filtering of haplotypes. A total of 247,267 haplotypes distributed in 73,100 phase sets were ob-
tained from the raw presence-absence HPA output file. Given the ploidy level of R570 ($\approx$12x
with limited aneuploidy), and the self-progeny nature of the population, a maximum of around
12 haplotypes can theoretically be expected from each PS. As a consequence, PSs with more
than 14 haplotypes were discarded as they likely resulted from reads originating from duplicated
regions. In addition, PSs revealing only one haplotype were discarded from the dataset, as well
as haplotypes with more than 25% missing values. A total of 215,704 haplotypes distributed
in 67,917 phase sets were conserved for the 87 progenies. The number of haplotypes per PS
varied from 2 to 14, with an average of 3.17.

SNPs and indels contributing to haplotypes. The total 215,704 haplotypes resulted from the com-
bination of 292,518 variants (biallelic SNPs, multiallelic SNPs, indels) reported in the VCF output
file. The presence/absence genotyping of each allele of these variants was re-inspected using the
same parameters used to generate the HPA file (see "HaploCharmer parameters" sub-section).
To this end, allele read depths that are initially reported specifically for each haplotype were ag-
gregated per allele. After filtering out alleles with more than 25% missing values 504,558 alleles
were conserved: 352,240 alleles from bi-allelic SNPs (69.81%), 130,745 alleles from bi-allelic
indels (25.91%), 6,700 alleles from multi-allelic SNPs (1.33%), and 14,873 other more complex
variants (2.95%).

Characterization of single-dose markers. Single-dose markers (a single copy is present at the lo-
cus in the parent, unlike multi-dose haplotypes) have a 50% chance of being transmitted to
the gametes, resulting in an expected presence:absence ratio of 3:1 in self-progeny. The prob-
ability of transmission of multi-dose markers depends on the number of doses, the ploidy and
the pairing of homologous chromosomes during meiosis. For instance, assuming a ploidy of 12
as in sugarcane cultivars and random bivalent chromosome pairing at meiosis (i.e. polysomic
inheritance), double-dose markers have 77% chance of being transmitted to the gametes. In
presence of strict preferential bivalent pairing of chromosomes (i.e. disomic inheritance), double-
dose markers have a 75% or 100% chance of being transmitted to gametes if the two doses
are located on unpaired or paired chromosomes, respectively. As a consequence, the expected
presence:absence ratio of multi-dose markers is always higher than 3:1 in the self-progeny of a
polyploid species like sugarcane.

To classify M markers (haplotypes or alleles) based on their segregation in the self-progeny
population, a mixture model was proposed in which the latent type Z_m of marker m is drawn
from a Multinomial distribution:

$$Z_m \sim \mathcal{M}\left(1, \pi_1, ..., \pi_K\right) \text{ iid,}$$

171 where π_k is the proportion of markers from type k (false positives, single-dose or multi-dose) and
172 iid stands for independent and identically distributed. The number of progenies X_m genotyped
173 as present for the marker is drawn from a Binomial distribution conditional on the marker type:

$$X_m | Z_m = k \sim \mathcal{B}\left(N_m; \delta_k\right) \text{ independent,}$$

174 where N_m is the number of progenies genotyped for the marker (excluding missing values), and
175 δ_k is the expected proportion of progenies genotyped as present. The inference of model param-
176 eters is done using an expectation-maximization (EM) algorithm (Dempster et al., 1977). In the
177 expectation step, the conditional probability $\mathbb{P}\left(Z_m = k | X_m\right) = \tau_{hk}$ is calculated:

$$\tau_{hk} = \frac{\pi_k \, \delta_k^{X_m}\left(1 - \delta_k\right)^{N_m - X_m}}{\sum_{k'=1} \pi_{k'} \, \delta_{k'}^{X_m}\left(1 - \delta_{k'}\right)^{N_m - X_m}}$$

178   In the maximization step, the values of parameters π_k and δ_k are updated as following:

$$\widehat{\pi}_k = \frac{\sum_{m=1}^{M} \tau_{mk}}{M}$$

$$\widehat{\delta}_k = \frac{\sum_{m=1}^{M} \tau_{mk} X_m}{\sum_{m=1}^{M} \tau_{mk} N_m}$$

179   Parameters were initialized as 1/3 for all π_k, and δ_k as 0.1, 0.75, and 0.9 for putative false
180 positives, single-dose and multi-dose markers categories, respectively. The EM algorithm iterates
181 until convergence, i.e. a difference of less than 0.001 between two iterations, of the marginal
182 log-likelihood:

$$\mathcal{L} = \sum_{m=1}^{M} \log \sum_{k=1} \pi_k \binom{N_m}{X_m} \delta_k^{X_m}\left(1 - \delta_k\right)^{N_m - X_m}$$

Haplotypes were considered single-dose in R570 when the probability of being a single dose
marker (obtained from τ_{hk}) was at least 0.99.

Genetic map. Single-dose haplotypes were grouped into non-redundant bins used as effective
markers for genetic mapping using the R-package "onemap" (MARGARIDO et al., 2007). After
estimating the recombination fraction between all bin pairs, haplotype bins were clustered into
linkage groups considering a LOD score of 7. Each linkage group was assigned to a basic chro-
mosome (hom(e)ology group) based on the mapping position of the majority of haplotypes from
the group. Note that known rearranged chromosomes in R570 (i.e. chromosomes 8/5, 5/9, 6/9,
7/10 and 8/10) were also taken into account as they displayed large numbers of haplotypes lo-
cated on two chromosomes of the monoploid reference (Healey et al., 2024). Haplotypes whose
position on the reference were discordant with the chromosome class or the recombination frac-
tion with neighboring haplotypes on the chromosome were filtered out.

Single-dose markers corresponding to SNPs and indels contributing to haplotypes were also
grouped into non-redundant bins to compare the number of effective markers obtained.

Results and discussion

Among the 215,704 haplotypes obtained from the R570 progeny, the estimated proportion
of false positives was below 1% (Table 1), demonstrating the high quality of the haplotyping ob-
tained with HaploCharmer. A haplotype was qualified as false positive if its expected proportion
in the progenies genotyped was low (δ_k = 2.25%), which is a very unlikely segregation for true
haplotypes. These false positives likely resulted from sequencing and mapping errors. Similar
proportions were found for SNPs and indels contributing to haplotypes.

Table 1 – Percentage π_k of markers (haplotypes or alleles) from each category (false pos-
itive, single-dose or multi-dose) and expected percentage δ_k of progeny genotyped as
present in the category, according to the type of marker: haplotypes obtained from Hap-
loCharmer or each type of variant constituting the haplotypes (bi- and multi-allelic SNPs,
bi-allelic indels and other more complex variants).

	False positive		Single-dose		Multi-dose	
	π_k	δ_k	π_k	δ_k	π_k	δ_k
Haplotypes	0.82%	2.25%	39.00%	72.86%	60.18%	99.31%
Bi-allelic SNPs	0.43%	2.24%	23.10%	73.72%	73.47%	99.74%
Bi-allelic indels	0.30%	1.80%	23.84%	73.84%	75.86%	99.75%
Multi-allelic SNPs	0.61%	2.20%	30.91%	72.76%	68.48%	99.52%
Others	0.73%	2.84%	30.05%	72.83%	69.22%	99.39%

The estimated proportion of single-dose haplotypes was 39.00%, which was higher than the
proportion obtained with bi-allelic SNPs (23.10%), bi-allelic indels (23.84%), multi-allelic SNPs
(30.91%) or other variants (30.05%). This result can be explained by multi-dose SNPs or indels
which, when combined, can reveal single-dose haplotypes. While the theoretical proportion of
single-dose markers present in a self-progeny is 75%, a lower proportion ($\delta_k \approx 73\%$) was ob-
served for haplotypes, SNPs and indels. This lower proportion could result from a slight segre-
gation bias due to meiosis abnormalities or insufficient sequencing depth leading to erroneous
genotyping of haplotypes or variants from which they were defined.

Multi-dose haplotypes were genotyped as present in nearly all progenies (δ_k = 99.31%), and
the same was observed for SNPs and indels contributing to haplotypes. This category aggre-
gated different dosage configurations (e.g. double-dose, triple-dose, ...). It was not realistic to

differentiate them due to (i) the limited size of our progeny sample and (ii) the conditioning of
the expected proportion of multidose markers genotyped as present (δ_k) on the pairing configu-
ration at meiosis. This means that two double-dose haplotypes may not have the same δ_k value
if they are located on chromosomes with contrasting pairing affinities at meiosis, although this
value is close to 100% in both cases.

The 81,857 single-dose haplotypes could be grouped into 25,094 non-redundant bins used
as effective markers for genetic mapping. In comparison, 21,789 and 24,435 non-redundant bins
could be obtained by considering single-dose bi-allelic SNPs or all variants included in haplotypes,
respectively. This higher number of effective markers obtained from haplotypes directly results
from the phenomenon where single-dose haplotypes can emerge from variants whose alleles
are all multi-dose.

Using haplotypes, linkage groups were created and those corresponding to copies of chromo-
some 1 are represented according to their physical position along the monoploid reference se-
quence used for read mapping (Figure 3). In addition, the consistency of these linkage groups was
validated using haplotype pairwise recombination fractions (Figure S1). Although 12 chromosome-
scale linkage groups are expected for R570 for chromosome 1 (Piperidis and D'Hont, 2020), the
linkage groups obtained did not sum to 12 copies at each position along chromosome 1. This
partial representation of the genome obtained from the linkage map was expected since large
chromosome segments are identical by descent due to inbreeding in the genealogy of cultivar
R570 (Healey et al., 2024), resulting in regions of the genome where haplotypes are multi-dose
(i.e. no single dose were available for mapping).

Figure 3 – Linkage groups obtained from single-dose haplotypes for the chromosome 1 hom(e)ology class. Each haplotype is represented by a horizontal black segment. Only linkage groups of at least 20 haplotypes are displayed.

Modern sugarcane cultivars such as R570 resulted from hybridization between two species
with distinct basic chromosome numbers, *Saccharum officinarum* (x=10) and *S. spontaneum* (x=8).
In *S. spontaneum*, two sets of three chromosomes have each been rearranged into two chro-
mosomes, in comparison with the ancestral chromosome structure observed in *S. officinarum*
(**Garsmeur2018**; Piperidis and D'Hont, 2020). Accordingly, some linkage groups corresponded
to known rearranged chromosomes. To illustrate, two linkage groups involving chromosomes 5,
6 and 9, which have been rearranged into two in *S. spontaneum*, are presented in Figure 4. The
two linkage groups and the positioning of haplotypes along the monoploid reference assembly

(ancestral genome structure) on Figure 4 well reflect this known rearrangement, with one part
of chromosome 9 merged with chromosome 5 and the other part merged with chromosome 6.
This attests of the quality of the genetic map obtained from haplotypes genotyped using Hap-
loCharmer.

Figure 4 – Two linkage groups corresponding to *S. spontaneum* rearranged chromosomes 5/9 and 6/9, with haplotypes positioned on the R570 reference (Chr5A, Chr6A and Chr9A) along with heatmaps of pairwise recombination fractions (rf) estimated between haplotypes. The positioning of haplotypes along the reference corresponds to the ancestral genome structure.

Fundings

This work was supported by the International Consortium for Sugarcane Biotechnology (Project
ICSB37). Whole genome sequencing data were produced by the Joint Genome Institute (`https:`
`//ror.org/04xm1d337`, proposal: `https://doi.org/10.46936/10.25585/60001084`), a USA De-
partment of Energy (DOE) Office of Science User Facility and the DOE Joint BioEnergy Institute,
which are supported by the Office of Science of the DOE operated under Contract No. DE-AC02-
05CH11231.

Conflict of interest disclosure

256 The authors declare that they have no conflicts of interest in relation to the content of the
257 article.

Data, script, code, and supplementary information availability

HaploCharmer source code is available at `https://gitlab.cirad.fr/agap/seg/haplocharmer`, along with documentation and examples. It can easily be installed on any Linux system. Dependencies are managed automatically via a conda installation or using a docker image.

Whole-genome sequencing data was available from the Joint Genome Institute portal: `https://genome.jgi.doe.gov/portal/Undpolgesequence/Undpolgesequence.info.html`. The genome assembly of R570 was available from the Phytozome portal: `https://phytozome-next.jgi.doe.gov/info/SofficinarumxspontaneumR570_v2_1`.

Figure S1: Linkage groups corresponding to chromosome 1, with haplotypes positioned on the monoploid reference (Chr1A) along with heatmaps of pairwise recombination fractions (rf) estimated between haplotypes. The positioning of haplotypes along the reference corresponds to the ancestral genome structure.

References

Aguiar D, Istrail S (2012). *HapCompass: A Fast Cycle Basis Algorithm for Accurate Haplotype Assembly of Sequence Data*. Journal of Computational Biology **19**, 577–590. `https://doi.org/10.1089/cmb.2012.0084`.

Aguiar D, Istrail S (2013). *Haplotype assembly in polyploid genomes and identical by descent shared tracts*. Bioinformatics **29**, i352–i360. `https://doi.org/10.1093/bioinformatics/btt213`.

Bhat JA, Yu D, Bohra A, Ganie SA, Varshney RK (2021). *Features and applications of haplotypes in crop breeding*. Communications Biology **4**. `https://doi.org/10.1038/s42003-021-02782-y`.

Chevreux B, Pfisterer T, Drescher B, Driesel AJ, Müller WE, Wetter T, Suhai S (2004). *Using the miraEST Assembler for Reliable and Automated mRNA Transcript Assembly and SNP Detection in Sequenced ESTs*. Genome Research **14**, 1147–1159. `https://doi.org/10.1101/gr.1917404`.

Clevenger JP, Korani W, Ozias-Akins P, Jackson S (2018). *Haplotype-Based Genotyping in Polyploids*. Frontiers in Plant Science **9**. `https://doi.org/10.3389/fpls.2018.00564`.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM, Li H (2021). *Twelve years of SAMtools and BCFtools*. GigaScience **10**. `https://doi.org/10.1093/gigascience/giab008`.

Dempster AP, Laird NM, Rubin DB (1977). *Maximum Likelihood from Incomplete Data Via the EM Algorithm*. Journal of the Royal Statistical Society Series B: Statistical Methodology **39**, 1–22. `https://doi.org/10.1111/j.2517-6161.1977.tb01600.x`.

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ (2011). *A framework for variation discovery and genotyping using next-generation DNA sequencing data*. Nature Genetics **43**, 491–498. `https://doi.org/10.1038/ng.806`.

Dijoux J, Rio S, Hervouet C, Garsmeur O, Barau L, Dumont T, Rott P, D'Hont A, Hoarau JY (2024). *Unveiling the predominance of Saccharum spontaneum alleles for resistance to orange rust in sugarcane using genome-wide association*. Theoretical and Applied Genetics **137**. `https://doi.org/10.1007/s00122-024-04583-3`.

Gattepaille LM, Jakobsson M (2012). *Combining Markers into Haplotypes Can Improve Population Structure Inference*. Genetics **190**, 159–174. `https://doi.org/10.1534/genetics.111.131136`.

Grivet L, D'Hont A, Dufour P, Hamon P, Roques D, Glaszmann JC (1994). *Comparative genome mapping of sugar cane with other species within the Andropogoneae tribe. Heredity* **73**, 500–508. https://doi.org/10.1038/hdy.1994.148.

Grivet L, D'Hont A, Roques D, Feldmann P, Lanaud C, Glaszmann JC (1996). *RFLP Mapping in Cultivated Sugarcane (Saccharum spp.): Genome Organization in a Highly Polyploid and Aneuploid Interspecific Hybrid. Genetics* **142**, 987–1000. https://doi.org/10.1093/genetics/142.3.987.

He D, Saha S, Finkers R, Parida L (2018). *Efficient algorithms for polyploid haplotype phasing. BMC Genomics* **19**. https://doi.org/10.1186/s12864-018-4464-9.

Healey AL, Garsmeur O, Lovell JT, Shengquiang S, Sreedasyam A, Jenkins J, Plott CB, Piperidis N, Pompidor N, Llaca V, Metcalfe CJ, Doležel J, Cápal P, Carlson JW, Hoarau JY, Hervouet C, Zini C, Dievart A, Lipzen A, Williams M, et al. (2024). *The complex polyploid genome architecture of sugarcane. Nature* **628**, 804–810. https://doi.org/10.1038/s41586-024-07231-4.

Köster J, Rahmann S (2012). *Snakemake—a scalable bioinformatics workflow engine. Bioinformatics* **28**, 2520–2522. https://doi.org/10.1093/bioinformatics/bts480.

Li H (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.* https://doi.org/10.48550/ARXIV.1303.3997.

MARGARIDO GRA, SOUZA AP, GARCIA AAF (2007). *OneMap: software for genetic mapping in outcrossing species: OneMap. Hereditas* **144**, 78–79. https://doi.org/10.1111/j.2007.0018-0661.02000.x.

Moeinzadeh MH, Yang J, Muzychenko E, Gallone G, Heller D, Reinert K, Haas S, Vingron M (2020). *Ranbow: A fast and accurate method for polyploid haplotype reconstruction. PLOS Computational Biology* **16**. Ed. by Alexander Schönhuth, e1007843. https://doi.org/10.1371/journal.pcbi.1007843.

Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, Forster J, Lee S, Twardziok SO, Kanitz A, Wilm A, Holtgrewe M, Rahmann S, Nahnsen S, Köster J (2021). *Sustainable data analysis with Snakemake. F1000Research* **10**, 33. https://doi.org/10.12688/f1000research.29032.2.

Piperidis N, D'Hont A (2020). *Sugarcane genome architecture decrypted with chromosome-specific oligo probes. The Plant Journal* **103**, 2039–2051. https://doi.org/10.1111/tpj.14881.

Shirley MD, Ma Z, Pedersen BS, Wheelan SJ (2015). *Efficient "pythonic" access to FASTA files using pyfaidx.* https://doi.org/10.7287/peerj.preprints.970v1.

Voorrips RE, Tumino G (2022). *PolyHaplotyper: haplotyping in polyploids based on bi-allelic marker dosage data. BMC Bioinformatics* **23**. https://doi.org/10.1186/s12859-022-04989-0.

Yang X, Song J, Todd J, Peng Z, Paudel D, Luo Z, Ma X, You Q, Hanson E, Zhao Z, Zhao Y, Zhang J, Ming R, Wang J (2018). *Target enrichment sequencing of 307 germplasm accessions identified ancestry of ancient and modern hybrids and signatures of adaptation and selection in sugarcane (Saccharum spp.), a 'sweet' crop with 'bitter' genomes. Plant Biotechnology Journal* **17**, 488–498. https://doi.org/10.1111/pbi.12992.