A SHERLOCK toolbox for eco-epidemiological surveillance of African trypanosomes in domestic pigs from Western Africa.

Roger-Junior Eloiflin*[1], Elena Pérez-Antón*[2], Aïssata Camara*[3], Annick Dujeancourt-Henry[2], Salimatou Boiro[3], Martial N'Djetchi[4], Mélika Taoré[4], Mathurin Koffi[4], Dramane Kaba[5], , Yann Le Pennec[6], Bakary Doukouré[6], Abdoulaye Dansy Camara[7], Moïse Kagbadouno[7], Pascal Campagne[8], Mamadou Camara[7], Vincent Jamonneau[1], Sophie Thevenon[1,9], Jean-Mathieu Bart[1,7], Lucy Glover[2,#], and Brice Rotureau[3,10,#]

[1] INTERTRYP, Université de Montpellier, Cirad, IRD, Montpellier, France

[2] Trypanosome Molecular Biology Unit, Institut Pasteur, Université de Paris, INSERM U1347, Paris, France

[3] Parasitology Unit, Institut Pasteur of Guinea, Conakry, Guinea

[4] Unité de Formation et de Recherche Environnement, Université Jean Lorougnon Guédé, Daloa, Côte d'Ivoire

[5] Unité de Recherche Trypanosomoses, Institut Pierre Richet, Bouaké, Côte d'Ivoire

[6] Virology Unit, Institut Pasteur of Guinea, Conakry, Guinea

[7] Programme National de Lutte contre la Trypanosomiase Humaine Africaine, Ministère de la Santé, Conakry, Guinea

[8] Bioinformatics and biostatistics hub, Institut Pasteur, Paris, France

[9] Cirad, UMR INTERTRYP, Montpellier, France

[10] Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France

* Equal contributions

[#] Corresponding authors: Brice Rotureau, rotureau@pasteur.fr; Lucy Glover, lucy.glover@pasteur.fr

 Short title

 A SHERLOCK toolbox for AAT surveillance

Abstract

Animal African trypanosomosis (AAT), caused by protist parasites of the genus
Trypanosoma, puts upward of a million head of livestock at risk across 37 countries in
Africa. The economic impact of AAT and the presence of human-infectious
trypanosomes in animals place a clear importance on improving diagnostics for animal
trypanosomes to map the distribution of the veterinary parasites and identify reservoirs
of human-infectious trypanosomes. We have adapted the CRISPR-based detection
toolkit SHERLOCK (Specific High-sensitivity Enzymatic Reporter unLOCKing) for
trypanosomatid parasites responsible for AAT (SHERLOCK4AAT) including Pan-
trypanosomatid, *Trypanozoon*, *T. vivax*, *T. congolense*, *T. theileri*, *T. simiae* and *T. suis*
assays. To test the applicability of this technique in the field, we analysed dried blood
spots collected from 200 farm and 224 free-ranging pigs in endemic and historical
human African trypanosomiasis foci in Guinea and Côte d'Ivoire, respectively. The
results revealed that SHERLOCK4AAT can detect and discriminate between
trypanosome species involved in multiple infections with a high sensitivity. 62.7 %
[58.1, 67.3] of pigs were found infected with at least one trypanosome species. *T.
brucei gambiense*, a human-infectious trypanosome, was found in one animal at both
sites, highlighting the risk that these animals may act as persistent reservoirs. These
data suggest that, due to their proximity to humans and their attractiveness to tsetse
flies, pigs could act as sentinels to monitor *T. b. gambiense* circulation using the
SHERLOCK4AAT toolbox.

Introduction
Through a combination of vector control, reactive screening, passive diagnosis, new
therapeutic drugs and the support of national control programmes, donors and
stakeholders, interrupting the transmission of gambiense human African
trypanosomiasis (HAT) at pan-African level is in reach (***Barrett et al., 2024***). Reported
annual cases have been maintained below 1,000 since 2018, making the elimination
(zero transmission) of gambiense HAT (gHAT) by 2030, one of the major objectives of
the World Health Organisation (WHO), achievable (***Franco et al., 2024***). Nevertheless,
animal African trypanosomosis (AAT), also known as *nagana*, remains a significant

burden to sustainable agricultural development, and represents a threat to livestock
and small holder economies (**Boulangé et al., 2022**), with a total estimated loss of
4.75 billion USD per year in terms of agricultural gross domestic product in tsetse-
infested lands (**Budd, 2001**). Unlike HAT, that is caused by either *Trypanosoma brucei*
(T. b.) gambiense in West Africa, or *T. b. rhodesiense* in East Africa, AAT
encompasses a wider range of trypanosomatid parasites. In sub-Saharan Africa,
animal infecting trypanosomes including *T. vivax*, *T. congolense*, *T. brucei*, *T. theileri*,
T. simiae and *T. suis* are transmitted by blood-sucking flies, such as Glossinidae and
Tabanidae, to a wide range of wild and domestic mammals. Socio-environmental
factors such as land cover, humidity, temperature and livestock migration impact
trypanosome transmission (**Karshima et al., 2016**). This is particularly evident in rural
areas of sub-Saharan Africa, where high incidences of *T. brucei*, *T. congolense* and *T.*
vivax are observed, due to conditions favourable to the proliferation of flies (**Okello et**
al., 2022b).

Symptoms of AAT are host and pathogen-dependent, with an infection potentially
leading to spontaneous abortions, reduction in milk and meat production, delays in
sexual maturity, low calving rates, loss of draught power, and ultimately to high
mortality, all of which compound poverty in many Sub-Saharan countries (**Kristjanson**
et al., 1999). Indirect control strategies including the combination of vector control,
breeding of trypanotolerant animals, and avoiding grazing in infested areas have
proved to be efficient in reducing the number of infections (**Meyer et al., 2016**).
Chemotherapy and chemoprophylaxis are the main means for controlling AAT,
although inadequate usage, presence of fake drugs and emergent drug resistance are
hampering trypanocide efficacy (**Giordani et al., 2016; Richards et al., 2021**).
Reliable and adapted diagnostic tests are crucial for parasite detection to better target
infected animals for treatment, as well as for parasite identification to understand the
distribution of trypanosomes in endemic areas. Microscopic, serological and molecular
methods are generally used to diagnose AAT. Microscopy based methods are the most
commonly used diagnostic methods, but lack sensitivity with a limit of detection (LoD)
ranging from 10^4 to 10^5 parasites/mL when using fresh blood, to 250 to 5,000
parasites/mL for methods combining centrifugation (**Desquesnes et al., 2022a**).
Serological methods, such as the card agglutination test (CATT) and ELISA, can be
used to identify anti-*Trypanosoma* antibodies, but their reliability is often compromised

by the poor antigen characterization in animal infecting trypanosomes or by the detection of antibodies from previous infections. In contrast, molecular diagnostic techniques, including PCR and Loop-mediated isothermal amplification (LAMP), offer increased sensitivity (73-100 %) and LoD (1-1000 parasites/mL) by targeting specific regions of the parasites DNA. For the main AAT-related *Trypanosoma* species, pan-species primers (targeting ITS1 for instance) and genus/species/type-specific primers are available and currently used to detect the DNA of parasites from the subgenus *Nannomonas* (*T. congolense*), *Duttonella* (*T. vivax*) and *Trypanozoon* (*T.b.brucei, T. evansi* and *T. equiperdum*) (***Álvarez-Rodríguez et al., 2022; Desquesnes et al., 2022a***).

AAT-related trypanosome species circulate co-endemically with *T. b. gambiense* in West and Central Africa, where domestic animals such as pigs, dogs, cattle, sheep and goats are suspected reservoirs (***Magang et al., 2023; Traoré et al., 2021***). In these regions, Suids (warthogs, domestic and bush pigs) and bovids (bushbucks and buffalos) are the preferred hosts for most tsetse fly species (***Serem et al., 2024***). In a historical HAT focus in Côte d'Ivoire, free-range pigs sharing the same environment as humans were identified as reservoirs for *T. b. gambiense* (***Traoré et al., 2021***). The diagnostic techniques employed in this study, involving single round (***Radwanska et al., 2002***) or nested PCR (***Morrison et al., 2008***), however tended to overestimate the prevalence of the human infective parasites in pigs (***Traoré et al., 2021***). Therefore, an active and accurate surveillance of the infection status of domestic animals using a more reliable diagnostic tool will be key to reach and sustain gHAT elimination (***Simo and Rayaisse, 2015; Vourchakbé et al., 2020***). In addition, a comprehensive map of the animal infective trypanosomes circulating in tsetse-infested areas would enable an improved control strategy and prevent or limit further economic losses.

In this context, developing new diagnostic tools that are species-specific and highly sensitive is essential to accurately diagnose infected individuals and define the circulation of trypanosomes in domestic animals. For this purpose, we have adapted the Specific High-sensitivity Enzymatic Reporter unLOCKing (SHERLOCK) detection that offers unprecedented levels of diagnostic specificity and sensitivity (***Gootenberg et al., 2017***), and has been adapted for the detection of HAT (***Sima et al., 2022***). SHERLOCK combines an isothermal amplification by recombinase polymerase (RPA) with Cas13a detection of the target RNA (***Gootenberg et al., 2017; Kellner et al.,***

2019; Piepenburg et al., 2006). The RPA step, in combination with a reverse transcriptase, allows the use of total nucleic acid (TNA) as an input, increasing the flexibility of the assay. Here, we have developed a broad panel of SHERLOCK assays for the detection of the six most prevalent AAT-related species. We have applied the SHERLOCK4AAT panel to two populations of domestic pigs from two eco-epidemiological regions of West Africa to determine their importance as reservoirs of both animal and human-infective trypanosomes. A total of 424 blood samples were collected (1) from free-ranging pigs in the last hypo-endemic foci of gHAT in Côte d'Ivoire, and (2) from farm pigs bred in the historical gHAT focus of forested Guinea where unconfirmed seropositive cases of gHAT are still sporadically detected (*Courtin et al., 2019*). These samples were screened using the complete SHERLOCK4AAT panel to provide an in-depth profile of the trypanosome species circulating among these domestic animals.

Materials and Methods

Ethics statement

The only reliable method available to maintain and amplify field-isolated animal trypanosomes is to experimentally infect laboratory animals, here mice. For this study, experimental infections were carried out at UMR INTERTRYP and were approved by the regional ethics committee for animal experimentation as part of the project APAFIS #34149-2020091118136527v5 and authorized by the French Ministry for Higher Education and Research. In Guinea, ethical approval was obtained from the *Comité National d'Ethique pour la Recherche en Santé* (CNERS) under agreement number 102/CNERS/19. In Côte d'Ivoire, sample collection was conducted within the framework of epidemiological surveillance activities supervised by the HAT National Elimination Program (HAT-NEP). No ethical statement is required by local authorities for domestic animal sampling. Any veterinarian may carry out blood sampling on domestic animals, with the authorization of the owner, as it is performed during prophylaxis or diagnostic campaign. Breeders gave their consent for animal sampling after being informed of the objectives of the study. For pig care, venous sampling was performed by a veterinary of the Laboratoire National d'Appui au Développement Rural (Ministry of Agriculture).

Mice infection and parasite purification

24 hours prior to infection with trypanosome parasites, 8-week-old female BALB/c mice (Charles River Laboratory) were immunosuppressed by intraperitoneal injection of 0.1 mL Endoxan 500 mg (Baxter). *T. congolense* Savannah subgroup IL1180 strain (Tanzania, Serengeti 1971, STIB212), *T. vivax* Magnan strain (Burkina Faso, Magnan 2011) and *T. theileri* (Nederland, Utrecht 2003, UNUT) parasites (INTERTRYP unit's cryobank) were inoculated by intraperitoneal injection of 0.15 mL of the cryopreserved blood containing parasites. Two mice were infected per strain and parasitaemia was monitored daily by tail blood sampling. Parasite yields were estimated by counting in a fast read 102 counting cells (Kova). Whole blood was collected by cardiac puncture under deep isoflurane anaesthesia when parasitaemia reached 1×10^8 cells per mL. Infected blood was passed over a diethylaminoethyl (DEAE)-cellulose column and the column washed using phosphate buffer saline (PBS) supplemented with 1 % glucose (PBSG). Parasites were collected in PBSG and concentrated by centrifugation (2000 rpm, 10 min, RT).

Cell culture

T. b. brucei Lister 427 bloodstream form and *T. b. gambiense* ELIANE cells were cultured in HMI-11 medium (ThermoFisher scientific) supplemented with 10 % fetal bovine serum (Sigma) at 37°C with 5 % CO_2. Human embryonic kidney (HEK) 293T cells were cultured in complemented dMEM (Merck) at 37°C with 5 % CO_2.

Epidemiological surveys

Pig blood samples were collected from the external jugular vein in heparinized tubes during two independent surveys in Côte d'Ivoire and Guinea. In Côte d'Ivoire, 224 free-ranging pigs were sampled in May, June and July 2021 in the Bonon and Sinfra hypoendemic HAT foci, the Bouaflé and Vavoua historical HAT foci and the Brobo HAT-free area {Kaba, 2023 #8655}. In Guinea, 200 farm pigs were sampled in March 2023 in the N'zérékoré prefecture, an ancient gHAT focus. Four drops of blood were dried on 4th grade Whatman filter paper (Cytivia Life Sciences) and conserved in sealed plastic bags with silica gel at ambient temperature for several months until total nucleic acid (TNA) extraction.

Nucleic acid extraction

RNA was extracted from bloodstream form parasites (*T. b. brucei* Lister 427, *T. b. gambiense* ELIANE, *T. congolense* Savannah subgroup IL1180, *T. vivax* Magnan and

T. theileri), HEK 293T cells and whole blood from French cattle, pig, sheep and goat (Bergerie de la Combe aux Loups). Bloodstream form parasites from infected mice, as well as cultured HEK 293T cells, were harvested in TRIzol (Life Technologies) and Chloroform:Isoamyl alcohol (24:1, Sigma-Aldrich) was added, followed by centrifugation at 13,000 rpm at 4°C for 15 min. The upper aqueous phase was transferred to a fresh tube containing 0.5 mL of 75 % ethanol. After incubation for 10 min at room temperature, RNA was extracted using a NucleoSpin RNA column (Macherey Nagel) or a RNAeasy kit (Qiagen) according to the manufacturer's instructions. 500 µL of blood sampled from French cattle, pig, sheep and goats was treated with 500 µL lysis buffer supplemented by 50 µL Proteinase K and incubated for 30 min at 56°C. The resulting mixture was transferred into a QIAshredder column (Qiagen) and centrifuged at 13,000 rpm for 2 min. The flow-through was then used for RNA extraction using a RNeasy Mini kit (Qiagen) according to the manufacturer recommendations. After extraction, qualities and quantities of RNA were evaluated using the NanoDrop™ One (Ozyme).

For extraction of TNA from dry blood spots (DBS), three 6 mm-punches from one spot of each animal DBS were resuspended in 300 µL of lysis buffer supplemented by 30 µL of Proteinase K and incubated for 30 min at 56°C. The punches and liquid were transferred into a QIAshredder column (Qiagen) and centrifuged at 13,000 rpm for 2 min. TNA was extracted from the flow-through using the Maxwell RSC Buccal Swab Kit (AS1640) with the Maxwell RSC Instrument (AS4500) according to the manufacturer's instructions. For each sample, all SHERLOCK assays were performed from a single TNA extraction.

Target selection, RPA primers and crRNAs design

Alignments of the *18S rRNA*, *IFX* and *GAPDH* genes were performed on sequences from different species (Supplementary Table 1) using Clustal Omega (**Sievers and Higgins, 2018**). The distance matrix representing the percentage identity between each sequence was represented using GraphPad Prism software (V.10).

BLAST analysis (**McGinnis and Madden, 2004**) of the *18S rRNA* and *GAPDH* genes selected target sequences was performed in the NCBI standard database of using the following selection of organisms: Microsporidia (taxid:6029), Ciliophora (taxid:5878), Haemosporida (taxid:5819), Diplomonadida (taxid:5738), Symbiontida (taxid:797087), Diplonemea (taxid:191814), Euglenida (taxid:3035), Kinetoplastea (taxid:5653),

Metakinetosplastina (taxid:2704647), Trypanosomatida (taxid:2704949), Trypanosomatidae (taxid:5654) and Euglenozoa (taxid:33682). The program selection was optimized for somewhat similar sequences (blastn) and in algorithm parameters, the maximum number of aligned sequences to display was set as 5000.

The presence of single nucleotide polymorphism (SNPs) in the target regions was assessed by aligning sequence variants of the parasite species using Clustal Omega (**Sievers and Higgins, 2018**). Sequence variants used are listed in Supplementary Table 1. SNPs identified in the TriTrypDB (https://tritrypdb.org) database were also taken into account in the design: NGS_SNP.Tb927_03: v5.1.921431, v5.1.921465, v5.1.921571, v5.1.921612, v5.1.921937, v5.1.922234, v5.1.922298, v5.1.922511, v5.1.922637, v5.1.922803, v5.1.923180, v5.1.959274; NGS_SNP.Tb927_07: v5.1.1951525, v5.1.1952283, v5.1.1951291; NGS_SNP.Tb927_02: v5.1.371291, v5.1.371584 and v5.1.370058.

Primers for Recombinase Polymerase Amplification (RPA) of targeted regions were designed in Primer-BLAST (**Ye et al., 2012**) using the customised parameters described by (**Kellner et al., 2019**). Oligonucleotides prone to generate self-dimers or cross primer dimers detected by Multiple Primer Analyzer (ThermoFisher) were discarded. Primers selected for each SHERLOCK assays are listed and highlighted in Supplementary Table 2. The T7 promoter sequence (5' GAAATTAATACGACTCACTATAGGG) was added at each 5' end of the selected forward primers to enable the *in vitro* transcription necessary for Cas13a detection (**Kellner et al., 2019**).

Production of crRNAs and positive controls

crRNA production was carried out as previously described (**Kellner et al., 2019; Sima et al., 2022**). Briefly, each crRNA DNA template, containing the spacer (specific targeted region) followed by the direct repeat (5' GATTTAGACTACCCCAAAAACGAAGGGGACTAAAAC) and the T7 promoter sequence, were *in vitro* transcribed using the HiScribe™ T7 Quick High Yield RNA Synthesis Kit (NEB) and purified using magnetic beads (Agencourt RNAClean XP). DNA template sequences used for the synthesis of each crRNA are detailed in Supplementary Table 2.

For the production of the synthetic positive controls, *in vitro* transcription of the targets
was performed as previously described (***Sima et al., 2022***) using RNA from the target
species for each specific assay. The amount of template used to produce the positive
control for each assay were as follows: 200 ng of RNA from *T. b. brucei* Lister 427 for
18S pan-trypanosomatid and 18S pan-*Trypanozoon* assays, 120 ng of RNA from *T. b.*
gambiense ELIANE strain for TgSGP assay, 120 ng of RNA from *T. vivax* Magnan
strain for IFX assay, 230 ng of RNA from *T. congolense* Savannah subgroup IL1180
strain for 18S *T. congolense*-specific assay and 5 ng of RNA from *T. theileri* for 18S *T.*
theileri-specific assay. Due to a lack of *T. simiae* and *T. suis* genetic material, the
positive controls were synthesized at Eurogentec using the *in-silico* predicted
sequence of the target region. Primers and target regions used to produce positive
controls are listed in the Supplementary Table 3.

<u>SHERLOCK assays</u>
SHERLOCK assays were performed as previously described (***Kellner et al., 2019;***
Sima et al., 2022). RPA was performed using the TwistAmp Basic kit (TwistDx)
following manufacturer's instructions with slight modifications. To enable reverse
transcription of the RPA (RT-RPA), 2.2 U of transcriptor reverse transcriptase (Roche)
were added to the reaction. Specific RPA primer pairs were used for each target
amplification at a concentration of 480 nM per primer, and the amplification reaction
was carried out in a heating block at 42°C for 45 minutes (***Sima et al., 2022***).
Leptotrichia wadeii Cas13a (*Lw*Cas13a) enzyme was produced as previously
described (***Kellner et al., 2019; Sima et al., 2022***). Briefly, pC013-Twinstrep-SUMO-
huLwCas13a plasmid (Addgene plasmid #90097) was inserted into *Escherichia coli*
RosettaTM 2(DE3) pLysS competent cells for *Lw*Cas13a expression and purification
(Gootenberg et al. 2017). Purified *Lw*Cas13a was conserved in Storage buffer (SB:
Tris-HCl 1M, NaCl 5M, DTT 1M and glycerol) (***Kellner et al., 2019***) in single-use
aliquots at -80°C to avoid freeze-thaw cycles.

The Cas13a-crRNA mediated target detection reaction was performed simultaneously
with the *in vitro* transcription of the target amplified by RT-RPA. *In vitro* transcription of
the amplified target was performed using T7 RNA polymerase (Biosearch technology).
Each Cas13a detection reaction was run in triplicate on a TECAN plate reader
(INFINITE F200 PRO M PLEX) and fluorescence readouts were collected every 10
minutes for 3 hours. All reactions were carried out with the inclusion of a negative

control template (NCT) adding nuclease-free water as input, and a positive control template (PCT), here the *in vitro* produced positive control for each specific SHERLOCK. For screening samples from epidemiological surveys, the same protocol as described above was used, except a TECAN plate reader (INFINITE F200 PRO Option Infinite F Nano +) was used and the assay was performed without replicates. Optimizations were carried out either at Institut Pasteur in Paris (France) or at INTERTRYP unit in Montpellier (France), while analysis of samples from epidemiological surveys was carried out at Institut Pasteur of Guinea.

Data representation and statistical analyses

Fold-change over background fluorescence was used to represent fluorescence readouts for all SHERLOCK assays. Background fluorescence is given by the NCT evaluation, where water is used as input material for the RPA reaction. For each test, sample fluorescence obtained after 3 hours of reading was divided by NCT fluorescence at the same time.

$$FC = \frac{Fsample\ t = 3h}{Fnct\ mean\ t = 3h}$$

Where *FC* represents Fold-Change over background fluorescence, *Fsample t=3h* represents sample fluorescence obtained after 3 hours and *Fnct mean t=3h* represents NCT fluorescence after 3 hours of reactions.

Analytical characteristics including threshold, sensitivity and specificity of each SHERLOCK assay were evaluated using Receiver Operating Characteristic (ROC) curves calculation comparing NCT versus PCT at controlled dilutions of parasites or hosts genetic materials. Results obtained from SHERLOCK specific assays on targeted parasites were gathered in a positive sample column, while results obtained from NCT, hosts and non-targeted parasites analysed with the assay, were placed in a negative sample column for ROC curve calculation and threshold evaluation. ROC curves were calculated using the Wilson/Brown method with a 95 % confidence interval and threshold values were selected based on the Youden's index (sensitivity + specificity – 1), that defines an optimal threshold as the one that maximizes the index of classification accuracy (**Brown et al., 2001; Schisterman et al., 2005**). The threshold of positivity for each SHERLOCK assay was selected from ROC analyses as the value corresponding to the maximum specificity (100 %). The limit of detection

(LoD) of each SHERLOCK reaction was assessed using specific tests on serial

dilutions of target parasite RNA. Dilutions ranged from 5.10^3 pg/μL (equivalent to 10^7

parasites/mL) to 5.10^{-5} pg/μL (equivalent to 0.1 parasites/mL). All plots and statistical

analyses were performed with GraphPad Prism software (V.10). The Venn diagrams

were made based on the Venny model by Oliveros J.C. (BioinfoGP, CNB-CSIC).

Maps were generated with the QGIS software version 3.42.0 with backgrounds

retrieved from data.gouv.fr (OSM2IGEO - Ivory Coast - SHP - EPSG:4326 for Côte

d'Ivoire, and OSM2IGEO - Guinea - SHP - EPSG:4326 for Guinea).

In the absence of reliable infection status in the field data from pigs, the posterior

distributions of performance estimates, such as sensitivity (Se), were jointly estimated

with a Bayesian latent class model (***Joseph et al., 1995***). The model takes boolean

outcomes (either positive or negative) of several tests as input. This Monte Carlo

method operates based on a Gibbs sampling process. The model iteratively updates

both the test performance parameters (sensitivity, specificity, and frequency of positive

individuals in sample collection) and the latent infection status. Flat prior was used.

After visually controlling that stationarity of the chain was quickly reached, we

discarded the first 150 burn-in iterations and calculated posterior densities based on >

10,000 iterations. 95 % confidence intervals were calculated for each proportion to

quantify the uncertainty of the estimates, providing a scientific range within which the

population proportions are likely to be accurate (***Brown et al., 2001***).

Results

Selection of targets for broad and species-specific SHERLOCK assays targeting AAT

species (SHERLOCK4AAT).

Target gene selection was based on two requirements: to develop (a) a broad

SHERLOCK assay for the detection of all trypanosomatid parasites and (b) for species

discrimination. To adapt SHERLOCK for the detection of AAT-related species, we

followed the criteria as described in (***Sima et al., 2022***). Briefly, candidate genes must

(1) be expressed in the bloodstream form of the parasites, (2) be conserved between

strains, (3) have few or no single nucleotide polymorphisms (SNPs), or (4) show

sequence divergence that would allow for species discrimination. All RPA primers and

crRNAs designed for each SHERLOCK assays are listed in Supplementary Table 2.

Pan-trypanosomatid assay. SHERLOCK has the capacity to discriminate between two sequences differing by a single nucleotide (***Gootenberg et al., 2017***). This study therefore focused on regions of the *18S rRNA* gene that show limited sequence homology between species. In addition to being a multicopy gene, the *18S rRNA* gene shares over 70 % identity between Kinetoplastea members (Supplementary Figure 1). Using BLAST analysis, the region between 1535-1672 bp (XR_002989632.1) was found to be conserved between all trypanosomatid species and was selected as the pan-trypanosomatid SHERLOCK target. RPA primers were designed to amplify this 138 bp region using Primer-Blast (***Ye et al., 2012***) (Supplementary Figure 2A and Supplementary Table 2). The level of identity of the target region was assessed between the different sequences described in the Trypanosomatidae family, as well as in the order, subclass and class that contain this family (Supplementary Figure 3). Examination of the Euglenozoa phylum, excluding the Kinetoplastea class, showed 98.4 % identity within the Trypanosomatidae family and 97.3 % identity within the Kinetoplastea class (Supplementary Figure 3). This was reduced to 82.9 % for the Phylum Euglenozoa, excluding the Kinetoplastea sequences (Supplementary Figure 3). Three distinct crRNAs were designed to this region, which can be used individually or multiplexed to increase the sensitivity (Supplementary Figure 4A and Supplementary Table 2). The sequences described in each group were 100 % conserved to the crRNA. RPA primers and crRNA targets of the pan-trypanosomatid SHERLOCK assay target regions (Kinetoplastea) were then compared to the homologous sequences from other classes within the Phylum Euglenozoa (Euglenida, Diplonemea, Symbiontida and Kinetoplastea) (Supplementary Figure 2A and 3). In addition, sequences from related protozoan parasite classes (Diplomonadida, Haemosporida and Ciliophora) and from the genetically close but phylogenetically distinct group Microsporidia were also included in the analysis (Supplementary Figure 2A).

Pan-*Trypanozoon* assay. A pan-*Trypanozoon* SHERLOCK assay was designed for increased selectivity to especially detect *T. brucei s.l.* (*sensu lato*). The regions within the 18S rRNA that were highly conserved among the species of the *Trypanozoon* subgenus, while showing significant divergence from other trypanosome species not belonging to the *Trypanozoon* subgenus, were analysed (Supplementary Figure 2B). A region ranging from 1318 to 1433 bp (XR_002989632.1) (Figure 1A and

Supplementary Table 2) was selected as it showed high conservation among the *Trypanozoon* subgenus (99.12 - 100 % percentage of identity, query coverage >90 %), except for three *T. evansi* sequences (91.96 - 96.49 %, OL869590.1, KT844944.1, MF142286.1). The crRNA target region selected was also conserved in the subgenus *Trypanozoon* and divergent in other parasites. Specific alignments with trypanosome species that could share the same hosts (*T. congolense*, *T. vivax*, *Leishmania spp.*, etc) were performed to assess target amplicon specificity (Supplementary Figure 2B). The pan-*Trypanozoon* SHERLOCK assay can discriminate between the species belonging to the *Trypanozoon* subgenus (*T. b. spp.*, *T. evansi* types A and B, and *T. equiperdum*) from other non-*Trypanozoon* trypanosomes or other co-endemic parasites.

Species-specific assays. For species discrimination within a single sample, several SHERLOCK assays were developed to detect the most prevalent species causing AAT. These include *T. congolense*, *T. vivax*, *T. theileri*, *T. simiae* and *T. suis*.

T. congolense: Regions conserved in the *18S rRNA* gene for all described *T. congolense* sequences, including the Savannah, Forest, Kilifi and Dzanga-Sangha groups, were analysed. The target region selected from position 116 to 246 bp of the *18S rRNA* gene (AJ009146.1) shows 100 % identity between the described *T. congolense* sequences. BLAST analysis of the *T. congolense 18S rRNA* gene sequences, excluding the target species (*T. congolense*, (taxid: 5692)) was also performed to determine the percentage of identity of closely related species. *T. simiae* (86.72 to 90.62 %), *T. sapaensis* (88.28 %), *T. godfreyi* (88.28 %), and *T. otospermophili* (87.02 %) had the highest percentage identity observed (Supplementary Figure 5). Species with common hosts and of veterinary relevance such as the two analysed *T. vivax* strains showed 82.81% and 87.5% identity. In addition, none of these sequences were fully conserved for the selected crRNA (Supplementary Figure 5), ensuring the specificity of the test and the absence of cross-detection with other species.

T. vivax: The *invariant flagellum antigen* (*IFX*) gene has no apparent homologue in any other *Trypanosoma* species or other related parasites, and was therefore selected (**Autheman et al., 2021; Romero-Ramirez et al., 2022**). BLAST analysis, using *T. vivax* Y486 (HE573024.1) as a reference genome, resulted in no matches (Figure 1A and Supplementary Table 2).

T. theileri: A conserved region between the three *T. theileri* lineages (TthI, TthII and TthIII) (**Brotánková et al., 2022**) was selected. The target region, ranging from 1014 to 1158 bp of the *18S rRNA* gene (AJ009163.1), has 100 % identity between the described sequences of the *T. theileri* lineages, and the most closely related sequences were from *T. cruzi*, *T. rangeli* and *T. dionisii*, with a 97.89 % identity (query coverage =65 %). The segment 1052-1081 bp corresponding to the crRNA sequence, is highly conserved between *T. theileri* variants (Figure 1A and Supplementary Table 2), however it contains a SNP (1060 C/T) in a number of the sequences analysed. Hence, one RPA primer pair and two multiplexed crRNAs were designed to target both version of the SNP for the *T. theileri* SHERLOCK assay (Supplementary Figure 4E).

T. simiae: The region from 1246-1346 bp of the *18S rRNA* gene (AJ404608.1) was selected as the targeted region for RPA amplification of the *T. simiae*-specific SHERLOCK assay. The segment 1273-1299 bp that corresponds to the crRNA sequence, has 100 % identity between the described sequences of *T. simiae* (Figure 1A and Supplementary Table 2).

T. suis: The multicopy *GAPDH* gene was used, as it can be phylogenetically separated into *T. suis* and *T. suis*-like strains from other African trypanosomes (**Rodrigues et al., 2020**). The region from 389 to 528 bp of the *GAPDH* gene (MK962702.1) has been selected as the targeted region for RPA amplification of the *T. suis*-specific SHERLOCK assay (Figure 1A and Supplementary Table 2). This region is 100 % conserved between the *T. suis* species described, and outside this clade, the most closely related sequences were found in *T. congolense*, with a 90 % of identity (query coverage >90 %). The segment 410-437 bp of the *gGAPDH* gene (MK962702.1), that corresponds to the crRNA target region, is highly conserved between *T. suis* species, with 100 % identity, and the BLAST analysis showed that this target fragment can only align to *T. suis* sequences but not to that of other Kinetoplastea species (taxid:5653).

<u>SHERLOCK can discriminate between closely related AAT-causing trypanosomatid species.</u>

All RPA primers and crRNAs for each targeted species were screened to select the optimal combinations, with the highest specificity and selectivity. For the pan-trypanosomatid assay, RPA primer pair A was used to screen three crRNAs (A:crRNA1, A:crRNA2 and A:crRNA3; Supplementary Figure 4A). From this, crRNA 3 was first selected for the simple pan-trypanosomatid assay. For the pan-*Trypanozoon*

assay, three RPA primer pairs (A, B and C) were screened with two crRNA candidates each (A:crRNA1, A:crRNA2, B:crRNA1, B:crRNA2, C:crRNA1, C:crRNA2, Supplementary Figure 4B). The A:crRNA1 combination was selected for the pan-*Trypanozoon* assay, being the one that obtained a higher fold change for the positive control and a lower fold change for the negative control. For the IFX *T. vivax* species-specific SHERLOCK assay, two RPA primer pairs were assessed (A and B) in combination with six crRNA candidates (A:crRNA1, A:crRNA2, A:crRNA3, B:crRNA4, B:crRNA5, B:crRNA6). RPA primer pair B and crRNA 6 showed the lowest variability between replicates and without any cross-reaction with other trypanosome species (Supplementary Figure 4C). For the *T. congolense*-specific assay, one pair of RPA primer (A) was combined with ten crRNA candidates (A:crRNA1, A:crRNA2, A:crRNA3, A:crRNA4, A:crRNA5, A:crRNA6, A:crRNA7, A:crRNA8, A:crRNA9, A:crRNA10). The crRNA1 that exhibited the lowest inter-replicate deviation and no cross-reactivity when exposed to the genetic material of non-target trypanosomatid parasites was selected (Supplementary Figure 4D). For the *T. theileri*-specific SHERLOCK assay, the combination of the RPA primer pair A with the two designed crRNAs (1 and 2) was used to cover the specific detection of all the *T. theileri* described sequences (Supplementary Figure 4E). For *T. simiae* and *T. suis*, single combinations of RPA primer and crRNA were tested. The specificity of each of these combinations was also verified by comparing the specific recognition of each RNA target to RNA from human embryonic kidney cell (HEK) (for pan-trypanosomatid and pan-*Trypanozoon* tests) or RNA from *T. brucei* (for *T. vivax*, *T. congolense* and *T. theileri* tests) (Supplementary Figure 4). In total, one RPA primer-crRNA combination was selected for each assay, based on the highest fold change over background fluorescence value, the highest specificity, and the minimum standard deviation (Figure 1A).

The specificity of the RPA primer-crRNA combinations selected for each assay was then cross-assessed against closely related parasites (Figure 1B-C). The single pan-trypanosomatid assay detected RNA from all trypanosomatid species tested (*T. b. brucei*, *T. b. gambiense*, *T. b. rhodesiense*, *T. evansi* (type A and B), *T. equiperdum*, *T. congolense*, *T. vivax*, *T. theileri*, *T. cruzi* and *Leishmania major*) but not from HEK cells, synthetic controls for *T. simiae* and *T. suis* or co-endemic parasites such as *Plasmodium falciparum* (Figure 1B). The pan-*Trypanozoon* assay detected only RNA

from members of the *Trypanozoon* subgenus (*T. b. brucei*, *T. b. gambiense*, *T. b. rhodesiense*, *T. evansi* (type A and B) and *T. equiperdum*) but not from HEK cells, synthetic controls for *T. simiae* and *T. suis* or co-endemic parasites such as *T. congolense*, *T. vivax*, *T. theileri* and *Plasmodium falciparum* (Figure 1B). The species-specific *T. congolense*, *T. vivax*, *T. theileri*, *T. simiae* and *T. suis* assays only detected their respective targeted parasite RNA or synthetic controls with no cross reactivity with any other RNAs (Figure 1C). Additionally, no unspecific cross-reaction was observed with the host-derived genetic material, as assessed by testing RNA from pig blood (SHERLOCK single pan-trypanosomatid and pan-*Trypanozoon* assays in Supplementary Figure 7).

The analytical sensitivity of each assay was quantified using serial dilutions of positive control RNAs from known numbers of parasites. For the single pan-trypanosomatid assay, the LoD was between 10 to 10^3 parasites per mL. The pan-*Trypanozoon* and *T. theileri* assays tested 10 parasites per mL. For the *T. congolense* assay, the LoD was found between 10 and 10^3 parasites per mL, and between 10^4 and 10^5 parasites per mL for the *T. vivax* test (Figure 2A). The sensitivity of *T. simiae* and *T. suis* tests were evaluated on synthetic positive controls, as no genetic materials were available, and both showed a LoD between 10 and 100 fM (Figure 2B).

Then, to improve the sensitivity of the simple pan-trypanosomatid assay, crRNA1 was tested in combination with crRNA3 for target recognition, as both crRNAs fall within the same RPA amplicon A (Figure 3A). Comparative analyses between the two versions were performed and multiplexing was shown to increase the performance of the assay, especially for the lower dilutions evaluated (100, 500 and 10^3 p/mL) by increasing the fold-change readouts of these positive controls by 1, 2.5 and 3-fold-change units, respectively (Supplementary Figure 6). Using results obtained on the blood samples from pigs in Côte d'Ivoire (see next section), we compared the two versions of the pan-trypanosomatid SHERLOCK assays to determine the posterior distribution of sensitivity estimates. We applied a Monte Carlo method (Gibbs sampling) proceeding in an iterative chain using the infection frequency, specificity, and sensitivity of the two tests to update the infection status. We determined the posterior sensitivity distribution of the simple versus multiplex pan-trypanosomatid SHERLOCK assays. Multiplexing crRNAs increased the median sensitivity from 66.1 % to 74.7 %, while keeping specificity above 80 % (Figure 3B). Furthermore, a higher concordance was observed

between the results obtained on the same samples with the pan-*Trypanozoon* SHERLOCK assay and the multiplexed pan-trypanosomatid SHERLOCK assay (25.7-% of double positives), as compared to the simple version (7.8 % of double positives) (Figure 3C). Given its improved detection efficiency, the multiplex version of the 18S rRNA pan-trypanosomatid assay was subsequently used exclusively and referred to as the multiplex pan-trypanosomatid assay.

To determine a positivity threshold value (fold-change cut-off) for each SHERLOCK assay, ROC curve analyses were performed and use to evaluate each following analytical sensitivity and specificity. The positivity threshold values were 6.050 for the pan-trypanosomatid test, 11.70 for the pan-*Trypanozoon* test, 1.923 for the *T. congolense* test, 1.831 for the *T. vivax* test, 1.805 for the *T. theileri* test, 1.546 for the *T. simiae* test, and 2.609 for the *T. suis* test (Supplementary Figure 8). Subsequently, for a given test, any sample with a fold change value above the threshold was considered positive for this test. Unsurprisingly, assays designed against multiple copy genes (*18S rRNA* and *GAPDH*) showed a greater sensitivity than those targeting single copy genes (*IFX*). Taken together, all the validated tests were able to specifically detect the RNA targets for which they were designed, making it possible to distinguish between closely related trypanosomatid parasite species. We then applied this novel SHERLOCK4AAT toolbox was applied to the analysis of field samples in a molecular eco-epidemiological survey.

Application of the SHERLOCK4AAT toolbox to eco-epidemiological surveys in pigs.

In transmission foci of western Africa, domestic pigs can host multiple trypanosome species. To investigate this, the SHERLOCK4AAT toolbox was used to survey trypanosomes circulating in domestic pigs from hypo-endemic and historical foci in Côte d'Ivoire and Guinea. In Côte d'Ivoire, blood samples were collected from 224 free-ranging pigs across the rural districts of Bonon, Bouaflé, Sinfra, Vavoua, and Brobo (Figure 4A). In the N'zérékoré prefecture of Forested Guinea, 200 farm pigs were sampled, with a focus on organized farming communities (Figure 5A). In addition to the SHERLOCK4AAT toolbox assays, all samples were also screened for *T. b. gambiense* using the previously described TgsGP SHERLOCK assay (**Sima et al., 2022**).

In total, 74.6 % [68.9, 80.3] (167/224) of the samples tested positive in at least one SHERLOCK assay in Côte d'Ivoire (Figure 4B), and 49.5 % [42.6, 56.9] (99/200) in

Guinea (Figure 5B) representing a high apparent prevalence of trypanosome infections

in these two populations of pigs. In parallel, 47.3 % [40.8, 53.8] (106/224) of the pigs

from Côte d'Ivoire tested positive using at least one parasitological method (buffy coat

examination, mAECT and /or isolation in mice; results detailed in a separate study,

manuscript in preparation). No systematic parasitological data was available for

Guinean samples. The difference in positivity rates between SHERLOCK detection

and parasitological approaches on Ivoirian samples was likely due to the higher

sensitivity of SHERLOCK as compared to classical methods. Indeed, we compared

three SHERLOCK assays (multiplex pan-trypanosomatid, pan-*Trypanozoon* and *T.*

congolense-specific SHERLOCK assays) to the three parasitological tests (buffy coat

examination, mAECT and /or isolation in mice) to determine the posterior distribution

of sensitivity estimates. A Monte Carlo method (Gibbs sampling) was used by

proceeding in an iterative chain using the infection frequency, specificity, and

sensitivity of the tests to update the infection status. This analysis suggested that, in

terms of sensitivity, the combination of SHERLOCK4AAT assays may compete with

combined parasitological methods (i.e., median sensitivity 91.4 % vs 84.3 %) while

controlling median specificity above 75 % (Supplementary Figure 9). Importantly, in

contrast to parasitological tests, SHERLOCK4AAT allowed for the identification of the

trypanosome species involved in more than half of these infections.

Using the multiplex pan-trypanosomatid assay, 41.1 % [34.6, 47.6] of the samples from

Côte d'Ivoire (90/219) and 29.5 % [23.2, 35.8] from Guinea (59/200) were positive

(Figure 4C and 5C). The pan-*Trypanozoon* assay identified 38.0 % [31.6, 44.4]

(85/224) and 39.0 % [32.2, 45.8] (78/200) of samples in each country respectively

(Figure 4C and 5C). As there is no specific test for *T. b. brucei*, samples that were

positive in the pan-*Trypanozoon* assay but negative in the TgsGP assay were

presumed to be *T. b. brucei* infections. Only one sample tested positive in the *T. b.*

gambiense-specific assay in each country (Figure 4B and 5B). The TgsGP-positive

sample from the district of Bonon, Côte d'Ivoire was negative for all other SHERLOCK

assays, as well as in all parasitological tests. In contrast, the TgsGP-positive sample

from Guinea was positive in the multiplex pan-trypanosomatid SHERLOCK assay, as

well as in the CATT and mAECT tests, but it was negative in the pan-*Trypanozoon*

SHERLOCK assay. To further explore trypanosome diversity, species-specific

SHERLOCK assays for AAT were applied to the samples. In total, 23.2 % [17.7, 28.7]

(52/224) of the samples in Côte d'Ivoire and 7.5 % [3.8, 11.2] (15/200) in Guinea were positive for *T. congolense*. Additionally, a single sample from Côte d'Ivoire tested positive in the *T. suis*-specific SHERLOCK assay. Furthermore, no RNA from *T. vivax* or *T. theileri* was detected with the IFX SHERLOCK assay.

Interestingly, a substantial number of co-infections were observed. In total, 24/219 individuals (11.0 ± 4.1 %) in Côte d'Ivoire and 13/200 in Guinea (6.5 ± 3.4 %) tested positive in the multiplex pan-trypanosomatid, the pan-*Trypanozoon* and the *T. congolense*-specific assays. These triple positive results indicate co-infections involving both a *Trypanozoon* species (likely *T. b. brucei*) and *T. congolense* (Figure 4D and 5D). In addition, 9/219 samples (4.1 ± 2.6 %) from Côte d'Ivoire tested positive in the pan-*Trypanozoon* and the *T. congolense* assays, which also represented co-infections with *T. brucei s.l.* and *T. congolense*. Overall, 15.2 % [10.5, 19.9] of the samples from Côte d'Ivoire showed co-infections with a *Trypanozoon* species and *T. congolense*. Additionally, 15 samples from Côte d'Ivoire tested positive in both the multiplex pan-trypanosomatid assay and the *T. congolense*-specific assay or were positive only for *T. congolense*. In Guinea, two samples tested positive in both the *T. congolense*-specific and multiplex pan-trypanosomatid SHERLOCK assays, indicating *T. congolense* mono-infections (Figure 4D and 5D).

Furthermore, 56 samples (25.6 ± 5.8 %) in Côte d'Ivoire, and 38 samples (19.0 ± 5.4 %) in Guinea tested positive in both the multiplex pan-trypanosomatid and the pan-*Trypanozoon* SHERLOCK assays. In contrast, 17 samples from Côte d'Ivoire and 40 from Guinea tested positive in the pan-*Trypanozoon* assay but negative in the multiplex pan-trypanosomatid assay, likely due to the lower sensitivity of the latter. Lastly, 19 samples (8.7 ± 3.7 %) in Côte d'Ivoire and 19 samples (9.5 ± 4.1 %) in Guinea tested positive in the multiplex pan-trypanosomatid assay only (Figure 4D and 5D). The absence of positivity in the *T. congolense*, *T. simiae* and *T. suis*-specific assays may be due to their lower sensitivity, or to infections with other non-targeted trypanosome species.

Discussion

The SHERLOCK4AAT toolbox, a new panel of highly sensitive and specific high-throughput assays for detecting animal trypanosomes.

Species identification is essential for assessing both the distribution of prominent parasites impacting veterinary health and spotting reservoirs for human-infective trypanosomes. In a context where symptoms are often mild and unspecific, especially in hosts like pigs, accurate species-specific detection methods are essential. Ideally, diagnostic assays should be able to analyse both fresh and stored (ambient temperature) samples to determine parasites circulating within a specific area.

In this study, conserved and species-specific targets have been used to adapt SHERLOCK for the detection of most members of the Trypanosomatidae family (Kinetoplastea class) (Supplementary Figure 3) of veterinary importance (pan-trypanosomatid), and species-specific assays to discriminate between the six most common animal trypanosomes species in the study regions: *Trypanozoon* (*T. b. brucei*, *T. evansi* and *T. equiperdum*), *T. congolense* (Savanah, Forest Kilifi and Dzanga-sangha), *T. vivax*, *T. theileri* (TthI, TthII and TthIII), *T. simiae* and *T. suis*. The design of both broad and species-specific assays was based primarily on sequences of the *18S rRNA* and *GAPDH* genes which, as well as being conserved across species (Supplementary Figure 1), are highly expressed and contain regions that allowed for species discrimination. However, for *T. vivax*, the single-copy *IFX* gene was selected for its high specificity (**Romero-Ramirez et al., 2022**). The first simple version of our pan-trypanosomatid assay employed a single crRNA guide and was previously used to evaluated blood samples from gHAT seropositive individuals with a specificity of 93 % (95.4-98.6) (**N'Djetchi et al., 2024**). To increase the performances of this assay, we added a second crRNA guide into the reaction mix to target the same RPA amplicon during the SHERLOCK reaction, resulting in an increased test sensitivity (Figure 3 and Supplementary Figure 6). For *Trypanozoon* species, no sequences were identified to distinguish *T. b. brucei*, *T. evansi* (A and B) and *T. equiperdum* as these parasites cannot be easily differentiated using the *18S rRNA* gene. In the future, other genes such as *RoTat1.2 VSG* or *ESAG6/7* could be explored to discriminate *T. evansi* from other *Trypanozoon* by SHERLOCK (**Claes et al., 2004; Holland et al., 2001**). Overall, *in silico* and *in vitro* analyses enabled us to assess the analytical specificities of these

new SHERLOCK assays, confirming that the targets selected were sufficiently specific

to be used as diagnostic for the species and groups of species concerned.

Using parasite RNAs, the LoD of the SHERLOCK assays were estimated to range

between 10 to 10^3 parasites/mL for the pan-trypanosomatid assay, 10 parasites/mL for

the pan-*Trypanozoon* and *T. theileri* assays, from 10 to 10^3 parasites/mL for *T.*

congolense and 10^4-10^5 parasites/mL for *T. vivax*. Most PCR-derived molecular

techniques currently used for detecting animal trypanosomes have a detection limit of

1 to 10 parasites/mL using genus- or species-specific primers, 50 parasites/mL using

pan-species ITS1 primers (**Desquesnes et al., 2022a**) or 5 parasites/mL using 18S

rRNA-targeting primers (**Deborggraeve et al., 2006**). The ROC analyses of the *T.*

brucei, *T. congolense* and *T. vivax* SHERLOCK assays (Supplementary Figure 8C)

also shown sensitivity and specificity comparable to those of the recently developed

7SL sRNA diagnostic test for Animal trypanosomosis (**Contreras Garcia et al., 2022**).

Therefore, in total, the analytical sensitivities obtained by the pan-trypanosomatid, pan-

Trypanozoon, *T. theileri* and *T. congolense* SHERLOCK assays were comparable with

those of other existing molecular tools. Yet, a significant difference in favour of using

SHERLOCK4AAT is that it detects RNAs, probing for active infections. The *T. vivax*

SHERLOCK assay*,* targeting the single-copy *IFX* gene (**Autheman et al., 2021**), did

not meet the sensitivity requirements for the molecular diagnosis of animal

trypanosomes and would require the identification of another target. As no genetic

material from parasites was available for developing *T. simiae* and *T. suis* SHERLOCK

assays, LoDs were determined using synthetic RNA controls. Both assays were able

to detect 10 to 100 fM of the synthetic controls, far from the detection capacity of

SHERLOCK, which can detect synthetic targets at a concentration of 2 aM in optimal

conditions (**Gootenberg et al., 2017**).

In the current version of the SHERLOCK4AAT toolbox, each assay can be run in 2 h

30 min, in a two-step reaction, at 2.5 € (exclusive of initial sample processing) (**Sima**

et al., 2022). All tests are optimized for high throughput screening with a fluorescence

readout; hence, they can be performed in standard lab conditions by a technician

trained for basic molecular biology. SHERLOCK4AAT was not developed for direct

case diagnosis in the field, but rather for eco-epidemiological surveillance at large

scale. Considering the absence of trypanosome species-specific treatment for

domestic animals, the future development of a RDT for AAT would only include the

pan-trypanosomatid target or equivalent. This would however require further technical
optimizations (TNA extraction, sensitivity, LFA, …).

<u>SHERLOCK4AAT profiling in Guinea and Côte d'Ivoire.</u>

Trypanosome species identification can be used to finely map the risk of AAT parasite
transmission in these areas. After cattle, pigs are the main domestic hosts infected by
animal trypanosomes, but they can also be reservoirs of *T. b. gambiense. T. b.*
gambiense infections in pigs are frequently reported in Sub-Saharan Africa
(***Ebhodaghe et al., 2018; Okello et al., 2022a***) and these domestic animals likely
represent an underestimated reservoir of human-infective parasites *(**Büscher et al.,**
2018; Mehlitz and Molyneux, 2019). This is all the more important than pigs are often
raised in close proximity to humans, increasing the risk of animal to human
trypanosome transmission, reinforcing our efforts to integrate the detection of both
animal and human parasites in domestic pigs in a One Health perspective (***Rotureau***
et al., 2022).

In blood samples from Ivorian pigs, analyses with the SHERLOCK4AAT toolbox
revealed 37.9 % positive samples for *Trypanozoon* RNAs and 23.2 % positive samples
for *T. congolense* RNAs, with 15.2 % samples positive for both tests. In addition, no
sample was positive for *T. vivax, T. theileri* or *T. simiae*, but one sample was positive
in the *T. suis* assay and another positive for *T. b. gambiense*. These findings are
consistent with the most recent known distribution of animal trypanosomes in free-
ranging pigs raised in Côte d'Ivoire's historical gHAT foci. The use of conventional
molecular diagnostics on 97 free-range pigs resulted in an overall positivity rate of 57 %
for *Trypanozoon* DNA (PCR-TBR) and 24 % for *T. congolense forest* DNA (PCR-TCF),
with 14 % of pigs positive in both PCRs and none positive for *T. vivax* (PCR-TVW)
(***N'Djetchi et al., 2017; Traoré et al., 2021***). The epidemiological status of AAT in
Guinea is unclear, as no large-scale molecular studies have been published so far.
The blood samples from farmed pigs analysed here were collected in the prefecture of
N'zérékoré, a former endemic focus of gHAT. Results showed a positivity rate of
39.0 % for the detection of *Trypanozoon* RNAs, like that observed in Côte d'Ivoire, but
a lower rate of positivity for *T. congolense* RNAs (7.5 %), with 6.5 % of samples positive
for both tests. In addition, one sample tested positive for *T. b. gambiense,* but no *T.*
vivax, *T. simiae, T. theileri* nor *T. suis* was detected. The lack of any samples positive
for *T. vivax or T. theileri* in any study areas confirms that pigs are not frequent carriers

of these species (*Fetene et al., 2021*). However, the blood samples positive for *T. b. gambiense* in each study site provide further evidence of the ability of pigs to carry human trypanosomes and to act as potential reservoir. Similarly, the presence of *T. b. gambiense* genetic material in pig samples has already been suspected in Côte d'Ivoire (*Traoré et al., 2021*).

In both sites, a high prevalence of trypanosomatid parasites was observed using the multiplex pan-trypanosomatid and/or pan-*Trypanozoon* assays (42.0 % in Côte d'Ivoire and 48.5 % in Guinea). Having partially deconvolved the species present in these animals; there is still a small proportion (8.7 % in Côte d'Ivoire and 9.5 % in Guinea) of trypanosomatid parasites that remained uncharacterised by our tests on samples found only positive in the multiplex pan-trypanosomatid SHERLOCK assay. Firstly, these assays were designed by *in silico* determination of the most specific and conserved region to target the entire Trypanosomatidae family, and even Kinetoplastea class, or to discriminate between species, yet the limited number of sequences available for some species in the literature and in the online databases may under-represent the real genetic diversity of these parasites *in natura*. In line with this hypothesis, there may also exist uncharacterised trypanosome species circulating among these reservoirs, as described in small mammals in central and East Africa (*Votypka et al., 2022*), and that were only detected by the broad multiplex pan-trypanosomatid assay. Altogether, these results are consistent with those reported in the few available previous studies (*N'Djetchi et al., 2017; Traoré et al., 2021*) and confirm the effectiveness of the SHERLOCK4AAT toolbox in analysing the distribution of trypanosomes in hypo-endemic gHAT foci. During natural infections, trypanosome nucleic acids can be detected in the skin, adipose tissue and various organs even if the blood remains negative (*Amisigo et al., 2024*). Trypanosome DNA also persists in blood and skin, even after drug treatment (*Ekloh et al., 2023*). Hence, applying SHERLOCK detection on skin samples in the future would be useful for monitoring both blood and skin-dwelling parasites. The future development of multiplexed assays combined with adapted lateral flow tests would also be highly desirable for reaching field-applicable point-of-care molecular diagnostic versions of the tests useable in the field. The absence of point of care diagnostic test for AAT is currently a major hurdle for its surveillance and control (*Desquesnes et al., 2022b*).

T. b. gambiense is suspected to circulate among domestic animals raised in close
contact to humans, such as pigs. Here, *T. b. gambiense* has been detected in one
animal in each study site. Considering the resulting low prevalence of *T. b. gambiense*
infections in these two populations of domestic pigs (0.45 ± 0.9 % in Côte d'Ivoire to
0.50 ± 1.0 % in Guinea) and the short life expectancy of domestic pigs reared for
commercial purposes in these two rural areas (6 to 8 months), these animals do likely
not represent a significant risk to the elimination of gHAT. This is apparently confirmed
by the absence of new human cases as well as by the low abundance of tsetse flies in
the two sites (article in preparation). However, because of their proximity to humans
and their easy access for frequent sampling, pigs could be used as sentinels to monitor
T. b. gambiense circulation, possibly with the SHERLOCK4AAT toolbox.

Acknowledgements
The authors thank Annette MacLeod (University of Glasgow, Glasgow, UK) for
providing RNA from *T. b. rhodesiense*, Nick Van Reet and Philippe Büscher (Institute
of Tropical Medicine [ITM], Antwerp, Belgium) for providing RNA from *T. equiperdum*,
T. evansi types A and B, Gerald Spaeth and Artur Scherf (Institut Pasteur, Paris,
France) for providing RNA from *Leishmania major* and *Plasmodium falciparum* and
John Kelly (The London School of Hygiene & Tropical Medicine, London, UK) for
providing RNA from *T. cruzi*. We warmly thank all the colleagues at the University Jean
Lorougnon Guédé (Daloa, Côte d'Ivoire) and at the Institut Pierre Richet (Bouaké, Côte
d'Ivoire) as well as all the health workers in the districts of Sinfra and Bonon (Côte
d'Ivoire) and N'zérékoré (Guinée) for their strong support in the field.

Funding
This project was funded by the Agence Nationale pour la Recherche (ANR-21-CE17-
0022-01 SherPa), the Programme Investissement d'Avenir of the French Government
through the two Laboratoires d'Excellence, ANR-10-LABX-62-IBEID and ANR-11-
LABX-0024-PARAFRAP, and the Pasteur Network (Bourse Calmette-Yersin to AC).

Author contributions
RE, EPA, AC, LG and BR conceived and designed the experiments. RE, EPA, AC,
SB, ADH, MND, MT, YL, BD, ADC, MK, VJ and JMB performed the experiments. RE,
EPA, AC, PC, LG and BR analysed the data. MK, DK, MC, VJ, ST, JMB, BR and LG

contributed reagents, materials and analysis tools. RE, EPA, AC, LG and BR wrote the

manuscript. MND, MK, VJ, ST, and JMB reviewed the manuscript.

Figure legends

Figure 1. Pan and species-specific SHERLOCK assays for human and animal African

trypanosome infections. (A) Schematic of the target regions used in the

SHERLOCK4AAT toolbox. *T. congolense*, *T. theileri* (multiplex)*, T. simiae*, Pan-

Trypanozoon and Pan-trypanosomatid SHERLOCK assays were designed using the

18S rRNA gene as a target; *T. vivax* SHERLOCK assay using the *IFX* gene and *T.*

suis test using the *gGAPDH* gene. Gene specific accession number in brackets below

the assay name. Details of the regions amplified are provided for each SHERLOCK

assay (amplicon). RPA primers are represented by black arrows, the extension in the

forward primer represents the T7 polymerase promoter sequence. The regions

targeted by the crRNAs (CRISPR guides) are also indicated in the diagram. The

schematic representation is to scale. (B) Evaluation of the specificity of single pan-

trypanosomatid and pan-*Trypanozoon* SHERLOCK tests RNA from *T. b. brucei* (*Tbb*),

T. b. gambiense (*Tbg*), *T. b. rhodesiense (Tbr*), *T. evansi* A (*Tev*A) and B (*Tev*B), *T.*

equiperdum (*Teq*) *T. congolense* (*Tcg*), *T. vivax* (*Tvx*), *T. theileri* (*Tth*), *Leishmania*

major (*Lm*), *Plasmodium falciparum* (*Pfl*) and HEK cells. The boxplot means are

represented by single lines, and error bars correspond to minimum and maximum

values. N=3 for the RPA amplifications and three Cas-13a detections for each RPA

amplification, totalling 9 replicates for each assay. *T. theileri* was used in fewer

replicates as an input due to the limited genetic material available. (C) Evaluation of

the specificity of *T. congolense*, *T. vivax*, *T. theileri*, *T. simiae* and *T. suis*-specific

SHERLOCK tests on RNA from *Tbb*, *Tbg*, *Tcg*, *Tvx*, *Tth*. Synthetic controls for *T.*

simiae (*Tsm*) and *T. suis* (*Tsu*) were used to evaluate the species-specificity of the

assays. N=1 RPA amplification and three Cas-13a detections totalling 3 replicates for

each assay. Error bars represent standard deviations from the mean of the replicates.

All inputs were used at a concentration of 5 ng/µL.

Figure 2. Evaluation of the sensitivity of SHERLOCK assays targeting trypanosome

species causing AAT. (A) Limit of detection (LoD) assays for the single pan-

trypanosomatid, pan-*Trypanozoon*, *T. vivax, T. congolense and T. theileri* assays were

performed using RNAs from bloodstream parasites diluted to obtain equivalent

concentrations of parasites per mL, ranging from 10^7 to 0.1 parasites/mL. (B) LoD of

T. simiae and *T. suis* assays were performed on consecutive dilutions of the synthetic

controls, ranging from 10^{10} to 10^1 fM. N=24 for pan-trypanosomatid assay, N=12 for

pan-*Trypanozoon* and *T. theileri* assays, N=6 for *T. simiae* assay and N=3 for *T.*

congolense, T. vivax and *T. suis* assays. Error bars represent standard deviations from

the mean of the replicates; open circle represents each replicate.

Figure 3. Comparison of the single and multiplex 18S pan-trypanosomatid

SHERLOCK assays efficiency. (A) Schematic of the single (one crRNA) and multiplex

(two crRNAs) pan-trypanosomatid SHERLOCK assays designed using the *18S rRNA*

gene as a target. Gene specific accession number in brackets below the assay name.

Details of the amplified gene regions are provided (amplicon). RPA primers are

represented by black arrows; the forward primer extension represents the T7

polymerase promoter sequence. The regions targeted by the crRNAs are indicated in

the diagram. (B) Posterior sensitivity estimates of the single (green) and multiplex

(purple) pan-trypanosomatid SHERLOCK assays on the 224 pig blood samples from

Côte d'Ivoire presented in Figure 4. Estimates were obtained with the Monte Carlo

method (Gibbs sampling) proceeding in an iterative chain. (C) Results of pan-

Trypanozoon SHERLOCK assay plotted against results of the single (left panel) and

multiplex (right panel) 18S pan-trypanosomatid SHERLOCK assays on pig samples

from Côte d'Ivoire. The coloured dashed lines represent the detection thresholds set

for each of the tests: single pan-trypanosomatid (green), multiplex pan-trypanosomatid

(purple), and pan-*Trypanozoon* (orange). Numbers and proportions (%) of double

positive samples are indicated at the top right of each graph and only positive for pan-

trypanosomatid in the bottom right part of each graph.

Figure 4. Diversity of blood trypanosomatids among free-range pigs from Côte d'Ivoire

assessed by the SHERLOCK4AAT toolbox. (A) Map of the Côte d'Ivoire, the region

where the samples were collected is highlighted in purple. (B) Results of the

SHERLOCK4AAT screening of 224 pig samples collected in Côte d'Ivoire. The

coloured dashed lines represent the detection thresholds set for each assay. Each

colour represents a specific test: multiplex pan-trypanosomatid (green), pan-

Trypanozoon (orange), *T. b. gambiense* (yellow), *T. congolense* (light green), *T. vivax*

(purple), *T. theileri* (red), *T. simiae* (blue) and *T. suis* (black). (C) Table summarizing

the number of positive samples as well as the percentage (%) positive infections for

each test, calculated on the total number of samples analysed. The total number of

samples analysed were 219 for the multiplex pan-trypanosomatid assay, 173 for the

T. vivax-specific assay, and 224 for all other SHERLOCK assays. (D) Venn diagram

representing the crossover between the positive samples analysed with the three tests

(multiplex pan-trypanosomatid (green), pan-*Trypanozoon* (orange) and *T. congolense*

(purple)) giving the highest prevalence. The percentage of the total is shown in

brackets below the number of samples constituting each fraction.

Figure 5. Diversity of blood trypanosomatids among farm pigs from Guinea assessed

by the SHERLOCK4AAT toolbox. (A) Map of Guinea showing the N'zérékoré

prefecture where samples were collected in pink. (B) Results of the SHERLOCK4AAT

screening of 200 pig samples collected in Guinea. The coloured dashed lines represent

the detection thresholds set for each of the tests. Each colour represents a specific

test: multiplex pan-trypanosomatid (green), pan-*Trypanozoon* (orange), *T. b.*

gambiense (yellow), *T. congolense* (light green), *T. vivax* (purple), *T. theileri* (red), *T.*

simiae (blue) and *T. suis* (black). (C) Table summarizing the number of positive

samples as well as the percentage (%) positive infections for each test, calculated on

the total number of samples analysed (n=200). (D) Venn diagram representing the

crossover between the positive samples analysed with the three tests (multiplex pan-

trypanosomatid (green), pan-*Trypanozoon* (orange) and *T. congolense* (purple)) giving

the highest prevalence. The percentage of the total is shown in brackets below the

number of samples constituting each fraction.

Supplementary figure legends

Supplementary Figure 1. A: Representation of the proportions of identity between

18S rRNA (A) and *GAPDH* (B) genes from Microsporidia, Ciliophora, Haemosporida,

Diplomonadida, Symbiontida, Diplonemea, Euglenida and Kinetoplastea, focusing on

several species within last family.

Supplementary Figure 2. (A) Alignments for 18S Pan-trypanosomatid SHERLOCK

target region. The sequences used for the alignments are as follows: Microsporidia

(*Nosema adalia*, KC412706.1), Ciliophora (*Vorticella sp.*, DQ487201.2),

Haemosporida (*Plasmodium falciparum*, XR_002966679.1), Diplomonadida (*Giargia*

intestinalis, XR_005248679.1), Symbiontida (*Calkinsia aureus*, EU753419.1),

Diplonemea (*Diplonema ambulator*, AY425009.1), Euglenida (*Euglena sanguinea*,

JQ281806.1) and Kinetoplastea (*Trypanosoma* (*T*) *brucei brucei*, XR_002989632.1).

(B) Alignments for 18S Pan-*Trypanozoon* SHERLOCK target region. The sequences used for the alignments are as follows: Human (H, pdb|5VYC|i), *Plasmodium falciparum* (Pfl, XR_002966679.1), *Leishmania major* (Lm, GQ332361.1), *T. cruzi* (Tcz, AJ009147.1), *T. vivax* (Tvx, EU477537.1), *T. congolense* (Tcg, AJ009146.1), *T. theileri* (Tth, AJ009163.1), *T. simiae* (Tsm, AJ404608.1), *T. suis* (Tsu, MK962702.1) and *T. brucei brucei* (Tbb, XR_002989632.1). Non-conserved bases between sequences are highlighted in bold.

Supplementary Figure 3. Summary of BLAST analysis obtained against the 18S pan-trypanosomatid SHERLOCK amplicon and evaluation of the homology of the target region of each crRNA. The percentage identity of the target amplicon of pan-trypanosomatid RPA was evaluated using different degrees of exclusion, to assess its percentage of identity within the different groups and especially in the Phylum Euglenozoa, excluding the target group Kinetoplastea. The number of sequences evaluated is shown, as well as the percentage of sequences where the target regions of the pan-trypanosomatid guides (crRNAs) are conserved.

Supplementary Figure 4. Screening of different RPA pairs of primers and/or crRNAs guides for each SHERLOCK assay optimization. In each evaluation a positive control (RNA from a target species, coloured bar) and a negative control (RNA from a non-target species, grey bar) were used. RPA pairs of primers are named A, B or C, while crRNAs were name from 1 to 10. The time point used is indicated. [#] indicates the combination of RPA primers and crRNA selected for the rest of the analysis. All inputs were used at a concentration of 5 ng/µL.

Supplementary Figure 5. Alignments of the 18S *T. congolense* SHERLOCK target region. The sequences used for the alignments are from *T. simiae*, *T. godfreyi*, *T. sapaensis*, *T. vivax*, *T. theileri* and *T. thomasbancrofti*. Non-conserved bases between sequences are highlighted in grey. The percentage identity of each sequence was analysed against the RPA amplicon in the *T. congolense*-specific SHERLOCK assay.

Supplementary Figure 6. Comparison of the limit of detection of the single and multiplex 18S pan-trypanosomatid SHERLOCK assays. Single and multiplex 18S pan-trypanosomatid SHERLOCK assays on dilutions of *T. b. brucei* RNA and on RNA extracted from HEK human cells as negative control. N=9 for each dilution. Error bars represent standard deviations from the mean of the replicates.

Supplementary Figure 7. Background cross-reactivity of the 18S pan-trypanosomatid (A) and 18S pan-*Trypanozoon* (B) SHERLOCK assays using genetic material extracted from pig blood as input. Means are represented by single lines. Quantities of porcine and *T. b. brucei* genetic material extracts included in each trial are as follows: "-" none, "+" 1 ng/µL, "++" 2 ng/µL and "+++" 3 ng/µL.

Supplementary Figure 8. Threshold determination for all new SHERLOCK4AAT assays. (A) Compilation of positive and negative results obtained using multiplex pan-trypanosomatid and pan-*Trypanozoon* SHERLOCK tests and ROC curves obtained for each analysis. The red dotted line marks the selected threshold for each test. (B) Compilation of positive and negative results obtained using the species-specific SHERLOCK assays and ROC curve obtained for each analysis. The red dotted line marks the selected threshold for each test. (C) Summary of the analytical features of each SHERLOCK assay. Analytical sensitivity (95 % of CI), specificity (95 % of CI) and fold change cut-off.

Supplementary Figure 9. Posterior sensitivity estimates of three SHERLOCK assays (multiplex pan-trypanosomatid, pan-*Trypanozoon* and *T. congolense*-specific SHERLOCK assays, red curve) to the three parasitological tests (buffy coat examination, mAECT and /or isolation in mice, orange curve) performed in parallel on the 224 pig blood samples from Côte d'Ivoire. Estimates were obtained with the Monte Carlo method (Gibbs sampling) proceeding in an iterative chain.

Supplementary Table 1. Accession numbers of organisms used to evaluate the presence of single nucleotide polymorphism (SNPs) in the different target regions.

Supplementary Table 2. Primers and crRNA guides used for the SHERLOCK4AAT toolbox.

Supplementary Table 3. RPA primers used to produce the *in vitro* transcribed targets (positive control templates).

References

Álvarez-Rodríguez A, Jin B-K, Radwanska M, Magez S. 2022. Recent progress in diagnosis and treatment of Human African Trypanosomiasis has made the elimination of this disease a realistic target by 2030. *Frontiers in medicine* **9**:1037094. doi: 10.3389/fmed.2022.1037094.

Amisigo CM, Amegatcher G, Sunter JD, Gwira TM. 2024. Adipose and skin
distribution of African trypanosomes in natural animal infections. *Parasites & vectors*
17:215. doi: 10.1186/s13071-024-06277-7.

Autheman D, Crosnier C, Clare S, Goulding DA, Brandt C, Harcourt K, Tolley C,
Galaway F, Khushu M, Ong H, Romero-Ramirez A, Duffy CW, Jackson AP, Wright
GJ. 2021. An invariant Trypanosoma vivax vaccine antigen induces protective
immunity. *Nature* **595**:96–100. doi: 10.1038/s41586-021-03597-x.

Barrett MP, Priotto G, Franco JR, Lejon V, Lindner AK. 2024. Elimination of human
African trypanosomiasis: The long last mile. *PLoS neglected tropical diseases*
18:e0012091. doi: 10.1371/journal.pntd.0012091.

Boulangé A, Lejon V, Berthier D, Thévenon S, Gimonneau G, Desquesnes M, Abah
S, Agboho P, Chilongo K, Gebre T, Fall AG, Kaba D, Magez S, Masiga D, Matovu
E, Moukhtar A, Neves L, Olet PA, Pagabeleguem S, Shereni W, Sorli B, Taioe MO,
Tejedor Junco MT, Yagi R, Solano P, Cecchi G. 2022. The COMBAT project:
controlling and progressively minimizing the burden of vector-borne animal
trypanosomosis in Africa. *Open research Europe* **2**:67.
doi: 10.12688/openreseurope.14759.2.

Brotánková A, Fialová M, Čepička I, Brzoňová J, Svobodová M. 2022. Trypanosomes
of the Trypanosoma theileri Group: Phylogeny and New Potential Vectors.
Microorganisms **10**. doi: 10.3390/microorganisms10020294.

Brown LD, Cai TT, DasGupta A. 2001. Interval Estimation for a Binomial Proportion.
Statistical Science **16**. doi: 10.1214/ss/1009213286.

Budd LT. 2001. Trypanosomosis, Tsestse and Africa.
https://assets.publishing.service.gov.uk/media/57a08d23e5274a31e0001686/ZC01
35a.pdf.

Büscher P, Bart J-M, Boelaert M, Bucheton B, Cecchi G, Chitnis N, Courtin D,
Figueiredo LM, Franco J-R, Grébaut P, Hasker E, Ilboudo H, Jamonneau V, Koffi M,
Lejon V, MacLeod A, Masumu J, Matovu E, Mattioli R, Noyes H, Picado A, Rock KS,
Rotureau B, Simo G, Thévenon S, Trindade S, Truc P, van Reet N. 2018. Do Cryptic
Reservoirs Threaten Gambiense-Sleeping Sickness Elimination? *Trends in*
parasitology **34**:197–207. doi: 10.1016/j.pt.2017.11.008.

Claes F, Radwanska M, Urakawa T, Majiwa PA, Goddeeris B, Büscher P. 2004.
Variable Surface Glycoprotein RoTat 1.2 PCR as a specific diagnostic tool for the

detection of Trypanosoma evansi infections. *Kinetoplastid biology and disease* **3**:3.
doi: 10.1186/1475-9292-3-3.

Contreras Garcia M, Walshe E, Steketee PC, Paxton E, Lopez-Vidal J, Pearce MC,
Matthews KR, Ezzahra-Akki F, Evans A, Fairlie-Clark K, Matthews JB, Grey F,
Morrison LJ. 2022. Comparative Sensitivity and Specificity of the 7SL sRNA
Diagnostic Test for Animal Trypanosomiasis. *Frontiers in Veterinary Science* **9**.
doi: 10.3389/fvets.2022.868912.

Courtin F, Camara O, Camara M, Kagbadouno M, Bucheton B, Solano P, Jamonneau
V. 2019. Maladie du sommeil dans le foyer historique de Guinée forestière :
actualisation grâce à une méthode géographique. *Parasite (Paris, France)* **26**:61.
doi: 10.1051/parasite/2019061.

Deborggraeve S, Claes F, Laurent T, Mertens P, Leclipteux T, Dujardin JC, Herdewijn
P, Büscher P. 2006. Molecular dipstick test for diagnosis of sleeping sickness.
Journal of clinical microbiology **44**:2884–2889. doi: 10.1128/JCM.02594-05.

Desquesnes M, Gonzatti M, Sazmand A, Thévenon S, Bossard G, Boulangé A,
Gimonneau G, Truc P, Herder S, Ravel S, Sereno D, Jamonneau V, Jittapalapong
S, Jacquiet P, Solano P, Berthier D. 2022a. A review on the diagnosis of animal
trypanosomoses. *Parasites & vectors* **15**:64. doi: 10.1186/s13071-022-05190-1.

Desquesnes M, Sazmand A, Gonzatti M, Boulangé A, Bossard G, Thévenon S,
Gimonneau G, Truc P, Herder S, Ravel S, Sereno D, Waleckx E, Jamonneau V,
Jacquiet P, Jittapalapong S, Berthier D, Solano P, Hébert L. 2022b. Diagnosis of
animal trypanosomoses: proper use of current tools and future prospects. *Parasites*
& vectors **15**:235. doi: 10.1186/s13071-022-05352-1.

Ebhodaghe F, Ohiolei JA, Isaac C. 2018. A systematic review and meta-analysis of
small ruminant and porcine trypanosomiasis prevalence in sub-Saharan Africa (1986
to 2018). *Acta tropica* **188**:118–131. doi: 10.1016/j.actatropica.2018.08.034.

Ekloh W, Sunter JD, Gwira TM. 2023. African trypanosome infection patterns in cattle
in a farm setting in Southern Ghana. *Acta tropica* **237**:106721.
doi: 10.1016/j.actatropica.2022.106721.

Fetene E, Leta S, Regassa F, Büscher P. 2021. Global distribution, host range and
prevalence of Trypanosoma vivax: a systematic review and meta-analysis. *Parasites*
& vectors **14**:80. doi: 10.1186/s13071-021-04584-x.

Franco JR, Priotto G, Paone M, Cecchi G, Ebeja AK, Simarro PP, Sankara D, Metwally
SBA, Argaw DD. 2024. The elimination of human African trypanosomiasis:

Monitoring progress towards the 2021-2030 WHO road map targets. *PLoS neglected*
tropical diseases **18**:e0012111. doi: 10.1371/journal.pntd.0012111.

Giordani F, Morrison LJ, Rowan TG, Koning HP de, Barrett MP. 2016. The animal
trypanosomiases and their chemotherapy: a review. *Parasitology* **143**:1862–1889.
doi: 10.1017/S0031182016001268.

Gootenberg JS, Abudayyeh OO, Lee JW, Essletzbichler P, Dy AJ, Joung J, Verdine
V, Donghia N, Daringer NM, Freije CA, Myhrvold C, Bhattacharyya RP, Livny J,
Regev A, Koonin EV, Hung DT, Sabeti PC, Collins JJ, Zhang F. 2017. Nucleic acid
detection with CRISPR-Cas13a/C2c2. *Science (New York, N.Y.)* **356**:438–442.
doi: 10.1126/science.aam9321.

Holland WG, Claes F, My LN, Thanh NG, Tam PT, Verloo D, Büscher P, Goddeeris
B, Vercruysse J. 2001. A comparative evaluation of parasitological tests and a PCR
for Trypanosoma evansi diagnosis in experimentally infected water buffaloes.
Veterinary parasitology **97**:23–33. doi: 10.1016/S0304-4017(01)00381-8.

Joseph L, Gyorkos TW, Coupal L. 1995. Bayesian estimation of disease prevalence
and the parameters of diagnostic tests in the absence of a gold standard. *American*
journal of epidemiology **141**:263–272. doi: 10.1093/oxfordjournals.aje.a117428.

Karshima SN, Lawal IA, Bata SI, Barde IJ, Adamu PV, Salihu A, Dross PN, Obalisa
A. 2016. Animal reservoirs of Trypanosoma brucei gambiense around the old Gboko
sleeping sickness focus in Nigeria. *Journal of parasitology and vector biology*.

Kellner MJ, Koob JG, Gootenberg JS, Abudayyeh OO, Zhang F. 2019. SHERLOCK:
nucleic acid detection with CRISPR nucleases. *Nature protocols* **14**:2986–3012.
doi: 10.1038/s41596-019-0210-2.

Kristjanson PM, Swallow BM, Rowlands GJ, Kruska RL, Leeuw PN de. 1999.
Measuring the costs of African animal trypanosomosis, the potential benefits of
control and returns to research. *Agricultural Systems* **59**:79–98. doi: 10.1016/S0308-
521X(98)00086-9.

Magang EMK, Kamga RMN, Telleria J, Tichit M, Crouzols A, Kaboré J, Hardy D,
Bouaka CUT, Jamonneau V, Rotureau B, Kuete V, Bart J-M, Simo G. 2023.
Prevalence of blood and skin trypanosomes in domestic and wild fauna from two
sleeping sickness foci in Southern Cameroon. *PLoS neglected tropical diseases*
17:e0011528. doi: 10.1371/journal.pntd.0011528.

McGinnis S, Madden TL. 2004. BLAST: at the core of a powerful and diverse set of
sequence analysis tools. *Nucleic acids research* **32**:W20-5.
doi: 10.1093/nar/gkh435.

Mehlitz D, Molyneux DH. 2019. The elimination of Trypanosoma brucei gambiense?
Challenges of reservoir hosts and transmission cycles: Expect the unexpected.
Parasite epidemiology and control **6**:e00113. doi: 10.1016/j.parepi.2019.e00113.

Meyer A, Holt HR, Selby R, Guitian J. 2016. Past and Ongoing Tsetse and Animal
Trypanosomiasis Control Operations in Five African Countries: A Systematic
Review. *PLoS neglected tropical diseases* **10**:e0005247.
doi: 10.1371/journal.pntd.0005247.

Morrison LJ, Tait A, McCormack G, Sweeney L, Black A, Truc P, Likeufack ACL,
Turner CM, MacLeod A. 2008. Trypanosoma brucei gambiense Type 1 populations
from human patients are clonal and display geographical genetic differentiation.
Infection, genetics and evolution : journal of molecular epidemiology and
evolutionary genetics in infectious diseases **8**:847–854.
doi: 10.1016/j.meegid.2008.08.005.

N'Djetchi MK, Camara O, Koffi M, Camara M, Kaba D, Kaboré J, Tall A, Rotureau B,
Glover L, Traoré MB, Koné M, Coulibaly B, Adingra GP, Soumah A, Gassama M,
Camara AD, Compaoré CFA, Camara A, Boiro S, Anton EP, Bessell P, van Reet N,
Bucheton B, Jamonneau V, Bart J-M, Solano P, Biéler S, Lejon V. 2024. Specificity
of serological screening tests and reference laboratory tests to diagnose gambiense
human African trypanosomiasis: a prospective clinical performance study. *Infectious*
diseases of poverty **13**:53. doi: 10.1186/s40249-024-01220-5.

N'Djetchi MK, Ilboudo H, Koffi M, Kaboré J, Kaboré JW, Kaba D, Courtin F, Coulibaly
B, Fauret P, Kouakou L, Ravel S, Deborggraeve S, Solano P, Meeûs T de, Bucheton
B, Jamonneau V. 2017. The study of trypanosome species circulating in domestic
animals in two human African trypanosomiasis foci of Côte d'Ivoire identifies pigs
and cattle as potential reservoirs of Trypanosoma brucei gambiense. *PLoS*
neglected tropical diseases **11**:e0005993. doi: 10.1371/journal.pntd.0005993.

Okello I, Mafie E, Eastwood G, Nzalawahe J, Mboera LEG. 2022a. African Animal
Trypanosomiasis: A Systematic Review on Prevalence, Risk Factors and Drug
Resistance in Sub-Saharan Africa. *Journal of medical entomology* **59**:1099–1143.
doi: 10.1093/jme/tjac018.

Okello I, Mafie E, Eastwood G, Nzalawahe J, Mboera LEG, Onyoyo S. 2022b. Prevalence and Associated Risk Factors of African Animal Trypanosomiasis in Cattle in Lambwe, Kenya. *Journal of parasitology research* **2022**:5984376. doi: 10.1155/2022/5984376.

Piepenburg O, Williams CH, Stemple DL, Armes NA. 2006. DNA detection using recombination proteins. *PLoS biology* **4**:e204. doi: 10.1371/journal.pbio.0040204.

Radwanska M, Claes F, Magez S, Magnus E, Perez-Morga D, Pays E, Büscher P. 2002. Novel primer sequences for polymerase chain reaction-based detection of Trypanosoma brucei gambiense. *The American journal of tropical medicine and hygiene* **67**:289–295. doi: 10.4269/ajtmh.2002.67.289.

Richards S, Morrison LJ, Torr SJ, Barrett MP, Manangwa O, Mramba F, Auty H. 2021. Pharma to farmer: field challenges of optimizing trypanocide use in African animal trypanosomiasis. *Trends in parasitology* **37**:831–843. doi: 10.1016/j.pt.2021.04.007.

Rodrigues CMF, Garcia HA, Rodrigues AC, Pereira DL, Pereira CL, Viola LB, Neves L, Camargo EP, Gibson W, Teixeira MMG. 2020. Expanding our knowledge on African trypanosomes of the subgenus Pycnomonas: A novel Trypanosoma suis-like in tsetse flies, livestock and wild ruminants sympatric with Trypanosoma suis in Mozambique. *Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases* **78**:104143. doi: 10.1016/j.meegid.2019.104143.

Romero-Ramirez A, Casas-Sánchez A, Autheman D, Duffy CW, Brandt C, Clare S, Harcourt K, André MR, Almeida Castilho Neto KJG de, Teixeira MMG, Machado RZ, Coombes J, Flynn RJ, Wright GJ, Jackson AP. 2022. Vivaxin genes encode highly immunogenic, non-variant antigens on the Trypanosoma vivax cell-surface. *PLoS neglected tropical diseases* **16**:e0010791. doi: 10.1371/journal.pntd.0010791.

Rotureau B, Waleckx E, Jamonneau V, Solano P, Molia S, Debré P, Dellagi K, Morand S. 2022. Enhancing research integration to improve One Health actions: learning lessons from neglected tropical diseases experiences. *BMJ global health* **7**. doi: 10.1136/bmjgh-2022-008881.

Schisterman EF, Perkins NJ, Liu A, Bondell H. 2005. Optimal cut-point and its corresponding Youden Index to discriminate individuals using pooled blood samples. *Epidemiology (Cambridge, Mass.)* **16**:73–81. doi: 10.1097/01.ede.0000147512.81966.ba.

Serem EK, Mburu DM, Abdullahi OA, Bargul JL. 2024. A scoping review on tsetse fly
blood meal sources and its assay methods since 1956 to 2022. *Parasites & vectors*
17:52. doi: 10.1186/s13071-023-06114-3.

Sievers F, Higgins DG. 2018. Clustal Omega for making accurate alignments of many
protein sequences. *Protein science : a publication of the Protein Society* **27**:135–
145. doi: 10.1002/pro.3290.

Sima N, Dujeancourt-Henry A, Perlaza BL, Ungeheuer M-N, Rotureau B, Glover L.
2022. SHERLOCK4HAT: A CRISPR-based tool kit for diagnosis of Human African
Trypanosomiasis. *EBioMedicine* **85**:104308. doi: 10.1016/j.ebiom.2022.104308.

Simo G, Rayaisse JB. 2015. Challenges facing the elimination of sleeping sickness in
west and central Africa: sustainable control of animal trypanosomiasis as an
indispensable approach to achieve the goal. *Parasites & vectors* **8**:640.
doi: 10.1186/s13071-015-1254-y.

Traoré BM, Koffi M, N'Djetchi MK, Kaba D, Kaboré J, Ilboudo H, Ahouty BA, Koné M,
Coulibaly B, Konan T, Segard A, Kouakou L, Meeûs T de, Ravel S, Solano P, Bart
J-M, Jamonneau V. 2021. Free-ranging pigs identified as a multi-reservoir of
Trypanosoma brucei and Trypanosoma congolense in the Vavoua area, a historical
sleeping sickness focus of Côte d'Ivoire. *PLoS neglected tropical diseases*
15:e0010036. doi: 10.1371/journal.pntd.0010036.

Vourchakbé J, Tiofack ZAA, Kante TS, Mpoame M, Simo G. 2020. L'identification
moléculaire de Trypanosoma brucei gambiense chez les porcs, les chiens et les
petits ruminants naturellement infectés confirme les animaux domestiques comme
réservoirs potentiels de la maladie du sommeil au Tchad. *Parasite (Paris, France)*
27:63. doi: 10.1051/parasite/2020061.

Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012. Primer-
BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC*
bioinformatics **13**:134. doi: 10.1186/1471-2105-13-134.

1122

Figure 1

Figure 2

Figure 3

Figure 4

A

B

C

	No. positive samples	% infections (n=224)
Trypanosomatid	90	41.1 %
Trypanozoon	85	37.9 %
T. b. gambiense	1	0.45 %
T. congolense	52	23.2 %
T. vivax	0	0.0 %
T. theileri	0	0.0 %
T. simiae	0	0.0 %
T. suis	1	0.45 %

D

Figure 5

A

B

Nzérékoré

C

	No. positive samples	% infections (n=200)
Trypanosomatid	59	29.5 %
Trypanozoon	78	39.0 %
T. b. gambiense	1	0.50 %
T. congolense	15	7.5 %
T. vivax	0	0.0 %
T. theileri	0	0.0 %
T. simiae	0	0.0 %
T. suis	0	0.0 %

D

Supplementary Figure 1

Supplementary Figure 2

A

        <u>RPA Forward Primer</u>   target region crRNA 1   target region crRNA 3   <u>RPA Reverse Primer</u>

```
Microsporidia    CTTAATTTGACTCAACGCGAGGTAA[…]AATATTTTATTATTCAGCGAGAATGGAA---[…]ATAGTGGTGCATGGGCCGTTTTCAATGGA TGCT-GTAAAGT-TTTGATTAATTTC
Ciliophora       CTTAATTTGACTCAACACTGGGAAA[…]AGGGCAAGAA---GATTTTAGGATTGACAGA[…]GTGGTGGTGCATGGCCGTTCTTAGC-TGG TGGA-GTGAGATGTCTGGTTAATTCC
Haemosporida     CTTAATTTGACTCAACACGGGAAAA[…]AGTTTAAGAC---AAGAGTAGGATTGACAGA[…]ATGGTGATGCATGGCCGTTTTTAGT-TCG TGAATATGATTTGTCTGGTTAATTCC
Diplomonadida    CTCAATCTGACTCAACGCGCG-CAC[…]AGGCCCGGAC---GCGCGGAGGACCGACAGC[…]GCGGTGGTGCATGGCCGCTCCCAGC-CCG TGGC-GCGAGCCGTCTGCTCCATTGC
Symbiontida      CTTAATTTGACTCAACACGGGAAAT[…]AGATCAGGAC---GCCCGGTGGATTGACAGA[…]ATGGTGGTGCATGGCCGCTCCTGGT-TGG TGGA-GTGATTTGTCTGGTTGATTCC
Diplonemea       CTTAATTTGACTCAACACGGGGAAT[…]AGGTCAGGAC---AATGTAAGGATTGACAGA[…]GTGGTGGTGCATGGCCGCTCCTGGT-TGG TGGA-GTGATTTGTCTGGTTGATTCC
Euglenida        CTTAATTTGACTCAACGCGGGGAAT[…]AGGTCCGGAC---GCAGCTGGGATGGACAGA[…]ACGGTGGTGCATGGCCGCTCCTGAT-TGG TGGA-GTGATTTGTCTGGTTGATTCC
Kinetoplastea    TTTAATTTGACTCAACACGGGGAAC[…]AGATCCGGAC---AGGGTGAGGATTGACAGA[…]ATGGTGGTGCATGGCCGCTTTTGGT-CGG TGGA-GTGATTTGTTTGGTTGATTCC
```

B

        <u>Forward RPA Primer</u>   crRNA target region   <u>Reverse RPA Primer</u>

```
H, pdb|5VYC|i            -------------------------[…]------------------------[…]CCCGCCGGGCAGCTTC-------------
Pfl, XR_002966654.1      ------------------AAAATATAT[…]CATTTCTTAGGGAATGTTGATTTT----[…]TTCCTTCAGTACCTTA-------------
Lm, GQ332361.1           CCGCGCCCGCTTTTA---CCAACTTAC[…]TA----------TTCGGCCTTTAC----[…]TATCCTCAGCACGTTTCTG--TTTTTTC
Tcz, AJ009147.1          TCTCGCCAATGAATATATTAAAATTAC[…]TT----------TTTGGTCTTCGTCTTT[…]TATCCTCAGCACGTTATCTGA-CTTCTTC
Tvx, EU477537.1          GCGCTCCACCAACCA--------TCAC[…]TTTGG-------CGCGGCCCTCGC----[…]TGTCCTCAGCACGTCC------AACCGCC
Tcg, AJ009146.1          GTGCAGTCACAACAA---ACAACTTAC[…]TGCTGCCGTACGTTCGCCCCTTATTTT[…]TATCCTCAGCACGTTGTTTACATTTTTC
Tth, AJ009163.1          CCTCGCCAATGGATA---TCAATTTAC[…]TT----------TTCGGTCCTCGC----[…]TATCCTCAGCACGTTATCTGA-CTTCTTC
Tsm, AJ404608.1          CCACAGCCAACTACA---CAAACTTAC[…]TGCCGCGGGTCGCC--------------[…]TATCCTCAGCACGTTGTTGA--CTCATTC
Tsu, MK954469.1          CGGTCCCAGTACTTA---ACAATTTAC[…]TTATATTCCGCCTGGCGTGCTTCGTTGT[…]TATCCTCAGCAC
Tbb, XR_002989632.1      GCCCCAATTTATTTA---TCAATTTAC[…]TATCACCCCCGGTTCCCTCTTTTGAGGT[…]TATCCTCAGCACGTTTCTTA--CTTCTTC
```

Supplementary Figure 3

BLAST exclusion	Range of amplicon percentage identity (arithmetic mean)	Number of sequences included	Percentage of sequences with conserved crRNA		Sequences with partially conserved crRNA	
			crRNA 1	crRNA 3	crRNA 1	crRNA 3
Family **Trypanosomatidae**	82.27 - 100% (98.4%)	1644	59.4 %	97.1 %	667	48
Order **Trypanosomatida**	82.27 - 100% (98.4%)	1652	59.6 %	97.1 %	667	48
Subclass **Metakinetoplastia**	81.88 – 100% (97.9%)	1931	57.6 %	94.8 %	818	100
Class **Kinetoplastea**	81.88 – 100% (97.3%)	2051	55.1 %	90.3 %	920	199
Phylum **Euglenozoa**	69.72 – 100% (91.7%)	3330	34.0 %	55.8 %	2199	1477
Phylum **Euglenozoa** (excluded Kinetoplastea)	69.72 – 98.55% (82.9%)	1287	0.6 %	0.7 %	1279	1278

Supplementary Figure 4

Supplementary Figure 5

	Accession	Alignment against the crRNA	Identity (%)	Query coverage
T. simiae	KR024686.1	TCCCTGCTCCGGCGGCGGGGCGACTCGC	90.62	100%
	KP307021.1	TCCCTGCTCCGGCGGCGGGGCGACTCGC	90.62	100%
	AJ009162.1	TCCCTGCTCCGGCGGCGGGGCGACTCGC	90.62	100%
	AJ223569.1	TCCCTGCTCCGGCGGCGGGGCGACTCGC	90.62	100%
	U22320.1	TCCCTGCTCCGGCGGCGGGGCGACTCGC	90.62	100%
	KP307022.1	TCCCTGCTCCGGCGGCGGGGCAACTCGC	89.06	100%
	AJ404608.1	TCCCTACCTCGGCGGCGGGGCAACTCGC	86.72	100%
	U22318.1	TCCCTACCTCGGCGGCGGGGCAACTCGC	86.72	100%
T. godfreyi	AJ009155.1	TCCCTACCCCGGCGGCGGGGCGACTCGC	88.28	100%
T. sapaensis	AB242822.1	TCTCTGTTCCGGCTGCGGGGCAACTCGT	88.28	100%
Trypanosoma vivax	EU477537.1	TCCCTGCTCCGGCGGCGGGGCAACCCGC	87.50	100%
	KM391829.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	KM391828.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	KM391827.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	KM391826.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	KM391823.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	KM391822.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	HE573019.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.38	100%
	U22316.1	TCCCTGCTCCGGCGACGGGGCAACCCGC	84.13	98%
	KM391825.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391824.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391821.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391820.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391818.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391816.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	83.59	100%
	KM391819.1	TCCCTGCTCCGGCGACGGCTTCCACCGC	82.81	100%
T. theileri	KR024688.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	86.72	100%
	LC618031.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	86.72	100%
	AJ009163.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	86.72	100%
	AB007814.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	86.72	100%
	LC618030.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	79%
	JX178191.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.29	79%
	JX178190.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	84.31	79%
	MG255951.1	TCTCTGTTACGGCGGTGGGGCAACTCAC	85.42	75%
	MG255958.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	84.38	75%
Trypanosoma thomasbancrofti	KT728394.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728393.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728392.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728391.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728390.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728389.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728388.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728387.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728386.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728385.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728384.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728383.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728382.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728381.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728380.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728379.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728378.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728377.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728376.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728375.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728374.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728373.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	OM509729.1	TCTCTGTTCCGGCGGTGGGGCAACTCAC	85.94	100%
	KT728396.1	TCTCTGTTCCGGCGATGGGGCAACTCAC	84.38	100%
	KT728395.1	TCTCTGTTCCGGCGATGGGGCAACTCAC	84.38	100%
	crRNA 18STcg1	**TCCCTGCTACGGCCGCGGGGTGACTCGC**		

Supplementary Figure 6

Single pan-trypanosomatid

Double pan-trypanosomatid

Supplementary Figure 7

Supplementary Figure 8

A

B

C

SHERLOCK Assay	Sensitivity (95 % CI)	Specificity (95 % CI)	Fold Change cut-off
Pan-trypanosomatid	**92.14 %** (90.43 to 93.57 %)	**100.0 %** (99.02 to 100.0 %)	**>6.050**
Pan-*Trypanozoon*	**95.10 %** (91.21 to 97.32 %)	**100.0 %** (97.93 to 100.0 %)	**>11.70**
T. vivax	**99.11 %** (96.82 to 99.84 %)	**100.0 %** (97.29 to 100.0 %)	**>1.831**
T. congolense	**97.98 %** (92.93 to 99.64 %)	**100.0 %** (97.04 to 100.0 %)	**>1.923**
T. theileri	**100.0 %** (98.42 to 100.0 %)	**100.0 %** (97.99 to 100.0 %)	**>1.805**
T. simiae	**99.12 %** (95.20 to 99.96 %)	**100.0 %** (96.60 to 100.0 %)	**>1.546**
T. suis	**100.0 %** (88.65 to 100.0 %)	**100.0 %** (86.20 to 100.0 %)	**>2.609**

Supplementary Figure 9

Posterior sensitivity estimates