Autoregulation of *RPL7B* by inhibition of a structural splicing enhancer

David Granas, Ishan Gammadde Hewa, Michael A. White, Gary D. Stormo

ABSTRACT

Yeast ribosomal protein gene RPL7B is autoregulated by inhibition of splicing. The first intron has a "zipper stem" that brings the 5' splice site near the branch point and serves as an enhancer of splicing that is required for efficient splicing because it has non-consensus branch point sequence of UGCUAAC. The intron also contains an alternative, and mutually exclusive structure, that is conserved across many yeast species. That conserved structure is a binding site for the Rpl7 protein so that when the protein is in excess over what is required for ribosomes, the protein binds to the conserved structure which eliminates the enhancer structure and represses splicing and gene expression.

INTRODUCTION

Feedback regulation is common across biology. From product inhibition of enzymatic pathways to autoregulation of gene expression, it is an efficient means of maintaining homeostasis of biological systems (Gerhart, 2014; Savageau, 1974b). The first discovered example of transcriptional autoregulation was in Phage lambda (Ptashne et al., 1976; Ptashne et al., 1980; Ptashne et al., 1982). Two lambda transcription factors (TFs), Repressor and Cro, negatively regulate each other's expression, but in addition, Repressor initially activates its own expression by binding cooperatively to two adjacent binding sites, and then at higher concentration turns off its own expression by binding to a third site. This positive-negative autoregulation system keeps the concentration of Repressor in a narrow range, poised to become activated at the appropriate signal (Ptashne, 1986). Since then, many examples of autoregulation by TFs have been identified (Alon, 2007; Crews & Pearson, 2009; Leyva-Diaz & Hobert, 2019). Evolution of TF autoregulation is readily accomplished because non-coding DNA evolves rapidly and the occurrence of binding sites for the TF in the appropriate location can be selected if it offers an advantage, which autoregulation often does (Kalisky et al., 2007; Savageau, 1974a, 1975).

Post-transcriptional autoregulation has also been observed frequently. In eukaryotic cells, RNA-binding proteins (RBPs) that control gene expression and alternative splicing can evolve to autoregulation, similarly to TFs, by selecting for binding sites for the RBP in appropriate locations in the non-coding regions of the gene (Buratti & Baralle, 2011; Muller-McNicoll et al., 2019; Yeo et al., 2007). Many RBPs function as enhancers or silencers of splicing, often regulating the choice of alternative isoforms of genes, and in many cases regulate their own expression by affecting splicing choices (Singh & Valcarcel, 2005). In bacteria, autoregulation is commonly observed even for proteins that do not normally function as regulators of expression. Most bacterial ribosomal genes are post-transcriptionally regulated either through direct autoregulation, or by being part of an operon that is regulated by one of its constituents (Deiorio-Haggar et al., 2013; Draper, 1989; Nomura et al., 1984; Nomura et al., 1980). Some tRNA synthetase genes are regulated by a mimic of a tRNA structure that occurs near the ribosome binding site, so when the protein is in excess over the amount needed for its normal function, it represses its own expression (Romby & Springer, 2003; Ryckelynck et al., 2005; Torres-Larios et al., 2002). While ribosomal proteins and tRNA synthetases are both RNA binding proteins, there are other examples of autoregulation of proteins that do not bind RNA as their normal function. First discovered in bacteriophage T4, the ssDNA-binding protein (gene 32) and DNA polymerase (gene 43) are each autoregulated at the translation initiation step, even though neither binds RNA as its normal function

(Andrake et al., 1988; McPheeters et al., 1988; Russel et al., 1976; Shamoo et al., 1993; Tuerk et al., 1990; von Hippel et al., 1982). Many other examples have been found in bacteria of RNA structures being involved in gene regulation (Chelkowska-Pauszek et al., 2021; Schlax & Worhunsky, 2003). Because of the enormous potential for RNA to bind to diverse ligands, it can evolve to be an exquisite sensor of cellular states including the concentrations of any functional class of genes. Riboswitches are prime examples of RNAs that have evolved to control the fate of their RNA, either by attenuation, translation or degradation based on the concentration of small molecules in the cell, without any protein requirement (Haller et al., 2011; Mandal & Breaker, 2004; McCown et al., 2017; Miranda-Rios et al., 2001; Mironov et al., 2002; Winkler et al., 2002). The invention of methods such as SELEX further highlights the enormous capability of RNAs to bind to a vast repertoire of ligands, essentially any protein (Gold et al., 1995; Joyce, 1994; Tuerk & Gold, 1990), which offers the possibility of widespread roles in regulating gene expression (Cech & Steitz, 2014; Stormo, 2018; Stormo & Ji, 2001).

Previously, we utilized the yeast GFP-fusion collection (Huh et al., 2003) to search for yeast ribosomal proteins that showed feedback regulation (Roy et al., 2020). For each of those genes we synthesized a copy of the gene encoding the same protein but lacking any potential cis-regulatory elements ("cre-less" versions of the genes), whose expression could be induced from the *GAL1* promoter. Feedback regulation was detected by the ability of the cre-less gene to repress the GFP-tagged wild-type gene. Of the 95 genes tested, 30 showed at least a 3-fold reduction in GFP expression after induction of the cre-less gene. We delved into the mechanism of regulation of *RPL1B*, a gene that was repressed 10-fold upon induction. We found that it contains an RNA sequence and structure immediately 5' of the initiation codon that is conserved from bacteria to archaea to eukaryotes in both the binding sites of the protein on rRNA and in the mRNA of the gene (Anger et al., 2013; Bernier et al., 2014; Fu et al., 2013; Kohrer et al., 1998). We also showed that disrupting that conserved sequence or structure eliminated regulation. Those results show that *RPL1B* autoregulation in yeast is very similar to many examples of bacterial autoregulation, where binding of the protein inhibits translation initiation.

In addition to regulation by simply blocking translation initiation, yeast can autoregulate by inhibition of splicing. Several examples of autoregulation of ribosomal protein expression by inhibition of splicing are known: *RPL22B, RPS41B, RPL30 and RPS9A* and *B*, are all regulated in that manner (Fewell & Woolford, 1999; Gabunilas & Chanfreau, 2016; Macias et al., 2008; Plocik & Guthrie, 2012; Vilardell, Chartrand, et al., 2000; Vilardell, Yu, et al., 2000). The non-ribosomal protein Yra1, which is involved in nuclear export, is also autoregulated by inhibition of splicing (Preker & Guthrie, 2006). Most yeast genes do not contain introns, but ribosomal protein genes are the exception, with 76% of them containing introns. Only nine genes in yeast that two introns, three of which are ribosomal protein genes (Parenteau & Abou Elela, 2019). The length distribution of introns in yeast is bimodal, with ribosomal genes having mean intron lengths of about 400 bases, whereas non-ribosomal introns are generally much shorter, with an average length closer to 100 bases (Spingola et al., 1999). Each of the yeast introns has been deleted, and many of those strains have growth phenotypes under some conditions (Parenteau & Abou Elela, 2019; Parenteau et al., 2011; Parenteau et al., 2008; Parenteau et al., 2019; Petibon et al., 2016). The secondary structure of yeast pre-mRNAs has long been recognized as important in influencing splicing efficiency (Charpentier & Rosbash, 1996; Gahura et al., 2011; Goguel & Rosbash, 1993; Rogic et al., 2008; Warf & Berglund, 2010). A recent study of yeast intronic RNA structures, using a combination of *in vivo* structure probing experiments and computational prediction, showed that most introns are highly structured, and it is common for there to be "zipper stems" that effectively shorten the intron length and

bring the 5' splice site close to the branch point (BP) (Rangan et al., 2024). In *RPS17B* it was demonstrated that such an RNA structure served as an enhancer for splicing (Libri et al., 1995). Computational analyses have suggested that the intron length between the 5' SS and BP, taking into account secondary structures that effectively shorten that distance, is correlated with splicing efficiency with longer introns being less efficiently spliced (Rogic et al., 2008)

In this paper we focus on *RPL7B* (also known as uL30 (Ban et al., 2014)) because our previous work showed it is likely to be autoregulated (Roy et al., 2020) and because the *RPL7B* gene has several interesting features that we suspected could be involved in the regulation. *RPL7B* and *RPL7A* contain two introns (Wong et al., 2023), which is very rare in yeast, occurring in only one other ribosomal protein gene (Parenteau & Abou Elela, 2019; Spingola et al., 1999). Both paralogs contain a snoRNA in the second intron. The first intron of both paralogs contains a predicted RNA secondary structure that is conserved across many yeast species (Gao et al., 2021; Hooks et al., 2016) and the first intron also contains the unusual BP (BP) sequence UGCUAAC, which only occurs in one other yeast gene (Spingola et al., 1999). The deviation from the consensus BP sequence of UACUAAC has been shown in other work to cause inefficient splicing (Rain & Legrain, 1997; Schirman et al., 2021).

RESULTS

To test if the introns of *RPLA* and *RPL7B* are sufficient as the cis-regulatory elements (CREs) for regulation, we inserted them into a chromosomal copy of GFP under control of the constitutive *TEF2* promoter (Peng et al., 2015). We drove expression of the cre-less Rpl7a protein from a plasmid under the control of the *GAL1* promoter (Roy et al., 2020). We compared the GFP expression levels by flow cytometry with and without galactose induction of the Rpl7a protein. Figure 1a is a schematic of the experimental design where "Insert" refers to the intron variants that were tested for *cis*-regulatory activity. Figure 1b shows the fluorescence levels of cells without GFP, as background, and Figures 1 c-e for cells in which both introns or only intron 1 or intron 2 of *RPL7B* are inserted into the *GFP* gene, with and without induction. Regulation of about 7-fold (0.85 change in log(fluorescence)) is observed when *GFP* contains both introns and only intron 1. With only intron 2 there is only a slight reduction in expression after induction. Figure 1 f-g shows that there is no significant regulation when both introns, or only intron 1, of *RPL7A* are present. We therefore focus on testing variants of intron 1 of *RPL7B* to elucidate the mechanism of regulation. The paralogous proteins Rpl7a and Rpl7b differ at five amino acids and endow ribosomes with somewhat different properties (Ghulam et al., 2020; Kim et al., 2009; Komili et al., 2007; Malik Ghulam et al., 2022; Palumbo et al., 2017) so we tested the Rpl7b protein for regulatory function by expressing a cre-less version of the gene. We found it to be equivalent to Rpl7a protein in that it regulates only *RPL7B* and not *RPL7A* (Supplemental Fig. S1), so all further work was performed with the Rpl7a protein. (Expression values for all tested variants are shown in Supplemental Table S1.)

We sought to identify the regions of intron 1 that are necessary for regulation by testing whether deletions eliminated regulation. Figure 2a shows a schematic of *RPL7B* intron 1 with various deletions indicated. The entire intron is 409 bases long with the first U of the UGCUAAC BP at position 382 after the 5' splice site. There is a predicted secondary structure that is conserved between 36 species of fungi from bases 200 to 355 within the intron (Hooks et al., 2016), which we refer to as the Hooks structure (Figure 2b). U1-U3 are about 63-base segments upstream of the Hooks structure, but after the 5' splice site sequence, that were independently deleted. D1 is the segment between the Hooks structure and the

BP that was deleted. S1-S4 are various deletions of the Hooks structure: S1 deletes the entire structure and S2-S4 delete independent stems or loops, as shown in Figure 2b. We also replaced the wild-type BP sequence of UGCUAAC with the consensus UACUAAC in each of the deletion mutants to assess its role in regulation.

Figure 3 summarizes the primary results. The names of each mutant are listed across the top of the figure. The vertical axis is the log of the fluorescence measurement with 0 defined as the wild-type intron 1 in the uninduced (not repressed) state. Each the four vertices of each quadrilateral represent the induction and BP states. The upper left (star in the wild-type panel) indicates the GFP expression with the wild-type BP without induction of the cre-less Rpl7a protein. The lower left (triangle in the wild-type panel) is expression with the wild-type BP after induction. In every case induction of Rpl7a reduces GFP expression, but the amount varies widely. The upper right vertex of the quadrilateral (square in wild-type panel) is the fluorescence when the wild-type BP sequence is replaced by the consensus sequence, in the absence of repression. In every case the consensus BP leads to higher expression, but the amount of change varies widely. The lower right vertex (circle in wild-type panel) is the fluorescence after induction of introns containing the consensus BP sequence.

Deletion U1 greatly reduces expression in the uninduced state, even lower than the repressed wild-type intron. It is still regulated, to a reduced extent. Replacing the wild-type BP sequence with the consensus restores expression, even slightly above the wild-type, and repression is maintained at the reduced level. U2 and U3 are not shown. They both exhibited about 2-fold higher expression than wild-type, and regulation was nearly the same as wild-type. We suspect these mutants have increased expression simply due to the shorter intron, and they were not studied further. The fact that the sequence deleted in U1 is necessary for high expression only with the wild-type BP suggests it contains a splice enhancer element that compensates for the weak BP.

To test the importance of the Hooks structure for regulation, the entire structure was deleted in S1. It still contains the splice enhancer region so is expressed at slightly higher level than wild-type, but it has lost most of the regulation. With that deletion, expression is repressed about 2-fold upon induction of Rpl7, instead of the 7-fold for wild-type. The consensus BP increases expression even more than in wild-type, perhaps due to the shortening of the intron. The combination of deletions U1 and S1 is similar to S1 alone, but somewhat higher expression even in the absence of the enhancer region, possibly because it is much shorter than the other introns tested. Deletions S2 and S3 remove the first and second stem-loops of the Hooks structure, and have similar effects to S1, with increased expression but reduced regulation. Deletion S4, which removes the third stem of the Hooks structure, has a much more severe effect than any of the other S-deletions, even S1 that removes the entire Hooks structure. It is also nearly insensitive to repression and is rescued to a larger degree upon replacement of the BP with a consensus sequence.

Deletion D1 removes the entire sequence between the Hooks structure and the BP. It has essentially no expression, only 0.1 above the background level, and is insensitive to either repression or replacement of the BP (not shown). Although D1 retains the entire canonical BP sequence (UGCUAAC and UACUAAC) it must remove some critical element for splicing. We made smaller deletions, D2-D4 each with nine bases deleted, in conjunction with S1, to determine if they are required for the enhancer activity. S1-D4 is essentially the same as S1 alone (not shown), but S1-D2 and S1-D3 each reduce expression about 0.3 below S1 alone, and show a larger increase in expression with the consensus BP. A possible explanation

for these results is that the sequences under U1 and D2-D3 form a secondary structure that serves as a splicing enhancer which is required for high level expression with the wild-type BP.

Enhancer and regulatory structures

In a study of the *RPL7A* intron, Howe and Ares (Howe & Ares, 1997) proposed a secondary structure between the 5' end of the intron and the region just 5' of the BP that effectively shortens the distances between the 5' splice site and the BP. More recent work. (Rangan et al., 2024), using *in vivo* chemical probing and structure prediction, confirmed that structure and extended it to a more stable structure. The sequences covered by U1 and D2-D3 in *RPL7B* differ substantially from the sequences in *RPL7A*, but a similar possible structure can be formed in the *RPL7B* intron. Figure 4a outlines our model for a secondary structure that brings the 5' splice near the BP and can serve as a splice enhancer. Figure 4b shows that structure in detail on the left. The bases in yellow, in Figures 4a and b, are the same bases that can form alternative structures. The structure in the lower part of Figure 4a, and shown in detail on the right in Figure 4b, is part of the conserved Hooks structure. We postulate that the Rpl7 protein binds to the Hooks structure, shifting the equilibrium towards that structure and away from the enhancer structure, thereby reducing expression. The structure on the right is probably just a portion of the binding site for Rpl7 protein, as deletion of the other stems, S2 and S3, also greatly reduced the level of repression. The reason that deletion S4 reduces expression even more than S1, is that while they both remove the 3' portion of the enhancer structure, S1 also reduces the size of the intron by over 150 bases, whereas S4 only removes 40 bases, leaving the intron still quite long compared to most yeast introns. In both cases, and especially in S4, changing the BP to a consensus sequence eliminates the need for the enhancer structure, giving rise to higher expression than the wild-type intron.

To further test this model, instead of deleting the large segment U1, we changed just the first ten bases shown in the left of Figure 4b, from positions 49 to 58, to a non-complementary sequence. This mutant would reduce the stability of the enhancer structure and shift the equilibrium to the structure on the right. This intron mutation is labeled M1 in Figure 3, which shows that it behaves nearly identically to the large deletion U1. We then altered the sequence at the 3' end of yellow sequence, to compensate for the M1 mutation and reform the enhancer structure, while leaving the right structure unchanged. This version is labeled M1-C1 in Figure 3, which shows that it restores behavior of the intron to near wild-type activity. In fact, it expresses GFP at a somewhat higher level than the wild-type intron, and shows somewhat reduced repression, probably because the mutated and compensated structure is predicted to be somewhat more stable than the wild-type enhancer structure.

Direct protein binding to the intron

The model requires that the Rpl7 protein bind to the intron to inhibit splicing. To test this, we overexpressed a version of Rpl7a protein fused to a protein A ZZ domain tag (Rpl7a-zz) (Gelperin et al., 2005) in the strains containing either the wild-type intron or the Hooks deletion (S1) in *GFP*. Expression of Rpl7a-zz was induced with galactose and samples were taken at 0 and 15 minutes after induction. After UV crosslinking, Rpl7a was isolated by immunoprecipitation with IgG resin, followed by reverse transcription of the bound RNA. We then used gene-specific PCR primers to determine whether the *GFP* or native *RPL7A* or *B* introns were bound by Rpl7a-zz (Figure 5). No PCR product was present at time 0, as expected because the tagged protein has not yet been expressed. At 15 minutes, the wild-type intron of *GFP* is bound by the protein, but the S1 version of the intron, with the Hooks structure deleted, is not bound. The *RPL7A* intron is not bound in either strain, consistent with that gene not being regulated. The

RPL7B intron is bound in both wild-type and S1 strains, which is consistent with the model. The lanes labeled T are for total RNA extracted from the respective strains and show that all the intron sequences are present and amplified specifically by the designed primers.

DISCUSSION

In the last several decades our knowledge of different classes of RNAs and their functional roles has been greatly expanded (Cech & Steitz, 2014). Some of those roles include the regulation of gene expression, and even RNAs with well-defined functions, such as snoRNAs, have additional functional roles (Caputa & Schaffer, 2016; Cheng et al., 2024). The diverse functional roles derive from the diversity of specific interactions that RNA is capable of. It can bind to RNA and DNA substrates with high specificity due to its base-pairing potential. It can also bind to proteins in a sequence-specific manner, where in addition to sequence, structure can play a significant role in the specificity of binding to proteins and other ligands, such as small molecules (Chelkowska-Pauszek et al., 2021). An important feature of RNAs is their capability of folding into two, or more, alternative structures of nearly equal free energy, and for environmental factors to influence which structure predominates. This is especially clear in the case of riboswitches, where small molecules can shift the equilibrium between two alternative structures that have altered fates for the adjacent mRNA (Breaker, 2011; Haller et al., 2011; Mandal & Breaker, 2004; McCown et al., 2017). Riboswitches can even be sensitive to temperature to regulate gene expression (Abduljalil, 2018; Reining et al., 2013).

In the case of *RPL7B*, the predominance of the two alternative structures (Figure 4a) is determined by the concentration of Rpl7 protein. When its concentration is low, the enhancer structure predominates and leads to gene expression. When its concentration increases, the alternative structure becomes preferred and expression decreases. The need for the structural enhancer is due to the length of the intron as well as the suboptimal BP sequence. When the intron is shortened considerably, the enhancer is less important, as shown by the high expression of the U1-S1 intron variant. In every case, replacing the wild-type BP with a consensus BP increases expression and generally eliminates the need for the enhancer structure. Regulation is largely dependent on the Hooks structure, although not entirely. Even when the entire Hooks structure is deleted, there remains about a two-fold change in expression upon induction of the Rpl7 protein. The presence of the consensus BP greatly reduces the degree of repression, indicating that binding of the Rpl7 protein to the RNA does not act by blocking spliceosome access to the BP sequence. If regulation occurred by the bound protein blocking access to the BP, the wild-type and the consensus BP sequences would be equally repressed. Rather, the autoregulatory mechanism requires that the long intron has a suboptimal BP sequence so that it is inherently spliced inefficiently. The structural enhancer greatly increases its splicing efficiency so that high level expression is possible. But repression can occur when the protein is in excess, not by blocking splicing directly, but by inhibiting the formation of the structural enhancer. To our knowledge, this is a novel mechanism for autoregulation.

RPL7A is not regulated (Figure 1) nor bound by the Rpl7 protein (Figure 5) under the conditions of our experiments. The fact that it contains the conserved Hooks structure (Hooks et al., 2016) suggests that it may be bound by the protein under alternative conditions. The *RPL7A* intron 1 proposed enhancer, or zipper stem, structure is longer and more stable (Rangan et al., 2024) than our proposed structure for *RPL7B* (Figure 4), consistent with it not being bound under conditions where the *RPL7B* intron is bound.

The alternative conditions under which *RPL7A* intron 1 is bound may require additional factors that help destabilize the enhancer structure.

Rpl7a and Rpl7b proteins are not functionally equivalent. They localize differently within the cell (Kim et al., 2009; Komili et al., 2007). *RPL7A* is the more highly expressed gene of the pair, accounting for 75% to over 90% of the Rpl7 protein (Ghaemmaghami et al., 2003; Malik Ghulam et al., 2022). A strain deleted for *RPL7A* shows a significant growth defect, but deletion of *RPL7B* has no effect (Malik Ghulam et al., 2022; Palumbo et al., 2017). In one study, adding a second copy of *RPL7B* did not suppress the growth defect of the *RPL7A* deletion (Malik Ghulam et al., 2022), while another study reported that if the second copy of *RPL7B* is under control of the *RPL7A* promoter, that can suppress the growth defect (Palumbo et al., 2017). They also respond differently to drugs. Cells expressing only *RPL7A* are more sensitive than wild-type to staurosporine, and cells expressing only *RPL7B* are less sensitive. Deletion of *RPL7B* has no effect on sensitivity to hygromycin, but cells expressing only *RPL7B* are much more sensitive. Furthermore, the two paralogs preferentially translate different subsets of genes.

Our data show that the first intron of *RPL7B* reduces gene expression even in the absence of induction of the repressor protein. When the first intron of *RPL7B* is inserted into GFP it reduces expression about 7-fold compared to the first intron (or both introns) of *RPL7A*, even in the absence of Rpl7 induction (Figure 1, uninduced curves of panels c versus g). But that reduced expression can be overcome by deleting parts of the Hooks structure required for regulation g (Figure 3). Of course, Rpl7 protein, from both paralogs, exists in the cells from the normal expression of the *RPL7A* and *RPL7B* genes. That amount of protein is sufficient to reduce the expression of GFP and presumably also contributes to the reduced expression of the Rpl7b protein compared to the Rpl7a protein. These paralogs present an excellent example of ribosome specialization (Xue & Barna, 2012) that may explain the evolutionary advantage of maintaining both genes and their differential regulation. It will be interesting to identify conditions under which RPL7A is repressed and the balance of the two proteins is altered.

MATERIALS and METHODS

Yeast strains

Plasmids MJB1-RPL7A and MJB1-RPL7B were created to express the cre-less versions of these genes as proteins fused to mCherry at the C-terminus, and under control of the inducible GAL1 promoter. "Cre-less" refers to the fact that the 5' and 3' UTRs have been replaced, the introns have been removed, and the codons have been shuffled to give the same protein sequence from a different mRNA sequence. These cre-less sequences were synthesized by Synbio and cloned into the MJB1 vector (previously described in Roy et al., 2020) using NEB's HiFi DNA Assembly. These plasmids contain the URA3 gene for selection in ura- yeast strains.

The Rpl7a-zz protein was expressed from *the RPL7A* plasmid from the movable ORF collection (Gelperin et al., 2005). This plasmid expresses Rpl7a with 3 C-terminal tags: 6x-His, HA, and ZZ Protein A. It also contains URA3 for selection in ura- yeast strains.

Yeast strains were constructed to express the eGFP gene containing different intron variants. The wild-type introns were cloned into the plasmid BAC690-TEF (previously described in Roy et al., 2020) using NEB's HiFi DNA Assembly. Intron variants were made using NEB's Site-Directed Mutagenesis Kit. A PCR amplicon was then generated from the plasmid containing the TEF2 promoter, eGFP with intron(s), ADH1

terminator, kanamycin/G418 resistance marker, and homology arms for integration into the yeast genome at the location of the dubious ORF YBR032W.

Yeast transformation was done using the LiAc/SS carrier DNA/PEG method. Transformants were selected on YPD plates with 400 µg/mL G418 and verified by sequencing.

GFP assays

Liquid cultures were inoculated and grown overnight in 400 µL SC-URA with 2% raffinose in 96-deep-well plates at 30°C. Overnight cultures were diluted into both SC-URA with 2% raffinose (uninduced) and SD-URA with 2% raffinose and 0.2% galactose (induced). Cells were grown at 30°C for 10 hours. 200 µL cultures were transferred to 96-well plates and assayed on a CytoFLEX S (Beckman Coulter). The live cells were gated and 10,000 events were acquired.

Assaying Rpl7-zz protein binding to the intron sequences

The movable ORF RPL7A plasmid was transformed into the strains containing the wild-type and S1 versions of *RPL7B* intron upstream of GFP. Strains were inoculated separately into 3 mL of SC-URA medium supplemented with 2% glucose and incubated overnight at 30°C. A 100 µL aliquot from each overnight culture was transferred into 50 mL of pre-warmed SC-URA medium containing 2% raffinose and incubated at 30°C for 6 hours. Cultures were subsequently diluted and transferred into 200 mL of pre-warmed SC-URA medium supplemented with 2% raffinose to an initial OD_{600} of 0.02 and they were grown at 30°C until OD_{600} 0.7–0.8. From each culture, 100 mL was transferred into sterile 250 mL conical centrifuge bottles, placed on ice, and labelled as time 0 (uninduced). The remaining 100 mL of each culture was combined with 50 mL of pre-warmed SC-URA medium containing 6% galactose and incubated for 15 min to induce Rpl7a-zz expression. A volume of 150 mL aliquot of each galactose-induced culture was transferred to sterile 250 mL conical centrifuge bottles, placed on ice, and labelled as time 15 (induced). Samples were subjected to 1,200 mJ/cm² of 254 nm UV irradiation using a Stratalinker UV Crosslinker 2400 as described in (Garrido-Godino et al., 2021). Cells were pelleted by centrifugation at 1,500 g, 4°C, for 5 min, and resuspended in 200 µL of lysis buffer (20 mM HEPES, pH 7.4, 2 mM Mg(OAc)₂, 100 mM KOAc, 100 µg/mL cycloheximide, 0.5 mM DTT, and 1× protease inhibitor cocktail [Sigma]). An equal volume of glass beads (500 µm, BioSpec) was added, and cells were lysed by vortexing at 4°C for 15 min. Lysates were centrifuged at 10,000 rpm, 4°C, for 5 min, and the supernatants were collected in new 1.5 mL centrifuge tubes. A 200 µL aliquot of lysate from each sample was mixed with an equal volume of 2× ice-cold binding buffer (100 mM Tris-HCl, pH 7.5, 24 mM Mg(OAc)₂, 1 mM DTT, 1 mM PMSF, 1 U/mL RNase inhibitor [NEB]) and incubated with 700 µL of IgG Sepharose (Cytiva) in 1.5 mL microcentrifuge tubes. Binding was carried out at 4°C for 1 h with constant gentle rocking. Resins were washed three times with 800 µL of ice-cold 1× wash buffer (50 mM Tris-HCl, pH 7.5, 100 mM KCl, 12 mM Mg(OAc)₂, 1 mM DTT, 1 mM PMSF), followed by DNase I treatment in 500 µL of nuclease-free water containing 10 U of DNase I (NEB) in 1× DNase I reaction buffer at 37°C for 10 min. Resins were then washed three additional times with 800 µL of ice-cold 1× wash buffer, and the final pellet was resuspended in 500 µL TRIzol (ThermoFisher) for RNA extraction according to the manufacturer's protocol. Total RNA was extracted from a separate isolate of the T0 samples by the addition of 500 µl TRIzol. RNA was then purified using the Monarch Spin RNA Cleanup Kit (10 µg) (NEB) and eluted in 10 µL of nuclease-free water and quantified using a NanoPhotometer® NP50. For cDNA synthesis, SuperScript™ IV First-Strand Synthesis System (Invitrogen) was used with the following gene-specific primers: RIP-GFP_cDNA for GFP, RPL7A_cDNA for RPL7A, and RPL7B_pull_cDNA for RPL7B (Table 1). GFP

cDNA from total RNA (T), wild-type, and S1 was PCR-amplified using T7_hooks_structure_F and T7_hooks_structure_R primers. Similarly, RPL7A cDNA was amplified using RPL7A_Intron_F and RPL7A_Intron_R, while RPL7B cDNA was amplified using T7_hooks_structure_F and T7_hooks_structure_R primers using Q5 Hot Start High-Fidelity DNA Polymerase (NEB). The final PCR products were analyzed by electrophoresis on a 3% agarose gel.

Table 1

Primer Name	Sequence (5' → 3')	Target
RIP-GFP_cDNA	GTAAGTAGCATCACCTTCACCTTCACCGGAGACAG	GFP
RPL7A_cDNA	CAGCTCTTTCAGCAGCGACTTGTTCAGC	RPL7A
RPL7B_pull_cDNA	CTTGATTCTGACAACGAAGACCAAC	RPL7B
T7_hooks_structure_F	TAATACGACTCACTATAGGGTGAAGTAGTTTATCTTCTTCAAGTACC	GFP & RPL7B
T7_hooks_structure_R	CAGGAGTCAAGATTTTTCTATTCAAAAATG	GFP & RPL7B
RPL7A_Intron_F	TGGCCGCTGAGTATGTATACG	RPL7A
RPL7A_Intron_R	CCTTAGACTTCTTCAACTGAGATTCTGG	RPL7A

REFERENCES

Abduljalil, J. M. (2018). Bacterial riboswitches and RNA thermometers: Nature and contributions to pathogenesis. *Noncoding RNA Res*, *3*(2), 54-63. https://doi.org/10.1016/j.ncrna.2018.04.003

Alon, U. (2007). *An introduction to systems biology : design principles of biological circuits*. Chapman & Hall/CRC. Publisher description http://www.loc.gov/catdir/enhancements/fy0654/2005056902-d.html

Andrake, M., Guild, N., Hsu, T., Gold, L., Tuerk, C., & Karam, J. (1988). DNA polymerase of bacteriophage T4 is an autogenous translational repressor. *Proc Natl Acad Sci U S A*, *85*(21), 7942-7946. https://www.ncbi.nlm.nih.gov/pubmed/3054876

Anger, A. M., Armache, J. P., Berninghausen, O., Habeck, M., Subklewe, M., Wilson, D. N., & Beckmann, R. (2013). Structures of the human and Drosophila 80S ribosome. *Nature*, *497*(7447), 80-85. https://doi.org/10.1038/nature12104

Ban, N., Beckmann, R., Cate, J. H., Dinman, J. D., Dragon, F., Ellis, S. R., Lafontaine, D. L., Lindahl, L., Liljas, A., Lipton, J. M., McAlear, M. A., Moore, P. B., Noller, H. F., Ortega, J., Panse, V. G., Ramakrishnan, V., Spahn, C. M., Steitz, T. A., Tchorzewski, M., . . . Yusupov, M. (2014). A new system for naming ribosomal proteins. *Curr Opin Struct Biol*, *24*, 165-169. https://doi.org/10.1016/j.sbi.2014.01.002

Bernier, C. R., Petrov, A. S., Waterbury, C. C., Jett, J., Li, F., Freil, L. E., Xiong, X., Wang, L., Migliozzi, B. L., Hershkovits, E., Xue, Y., Hsiao, C., Bowman, J. C., Harvey, S. C., Grover, M. A., Wartell, Z. J., & Williams, L. D. (2014). RiboVision suite for visualization and analysis of ribosomes. *Faraday Discuss*, *169*, 195-207. https://doi.org/10.1039/c3fd00126a

Breaker, R. R. (2011). Prospects for riboswitch discovery and analysis. *Mol Cell*, *43*(6), 867-879. https://doi.org/10.1016/j.molcel.2011.08.024

Buratti, E., & Baralle, F. E. (2011). TDP-43: new aspects of autoregulation mechanisms in RNA binding proteins and their connection with human disease. *FEBS J*, *278*(19), 3530-3538. https://doi.org/10.1111/j.1742-4658.2011.08257.x

Caputa, G., & Schaffer, J. E. (2016). RNA Regulation of Lipotoxicity and Metabolic Stress. *Diabetes*, *65*(7), 1816-1823. https://doi.org/10.2337/db16-0147

Cech, T. R., & Steitz, J. A. (2014). The noncoding RNA revolution-trashing old rules to forge new ones. *Cell*, *157*(1), 77-94. https://doi.org/10.1016/j.cell.2014.03.008

Charpentier, B., & Rosbash, M. (1996). Intramolecular structure in yeast introns aids the early steps of in vitro spliceosome assembly. *RNA*, *2*(6), 509-522. https://www.ncbi.nlm.nih.gov/pubmed/8718681

Chelkowska-Pauszek, A., Kosinski, J. G., Marciniak, K., Wysocka, M., Bakowska-Zywicka, K., & Zywicki, M. (2021). The Role of RNA Secondary Structure in Regulation of Gene Expression in Bacteria. *Int J Mol Sci*, *22*(15). https://doi.org/10.3390/ijms22157845

Cheng, Y., Wang, S., Zhang, H., Lee, J. S., Ni, C., Guo, J., Chen, E., Wang, S., Acharya, A., Chang, T. C., Buszczak, M., Zhu, H., & Mendell, J. T. (2024). A non-canonical role for a small nucleolar RNA in ribosome biogenesis and senescence. *Cell*, *187*(17), 4770-4789 e4723. https://doi.org/10.1016/j.cell.2024.06.019

Crews, S. T., & Pearson, J. C. (2009). Transcriptional autoregulation in development. *Curr Biol*, *19*(6), R241-246. https://doi.org/10.1016/j.cub.2009.01.015

Deiorio-Haggar, K., Anthony, J., & Meyer, M. M. (2013). RNA structures regulating ribosomal protein biosynthesis in bacilli. *RNA Biol*, *10*(7), 1180-1184. https://doi.org/10.4161/rna.24151

Draper, D. E. (1989). How do proteins recognize specific RNA sites? New clues from autogenously regulated ribosomal proteins. *Trends Biochem Sci*, *14*(8), 335-338. https://www.ncbi.nlm.nih.gov/pubmed/2678632

Fewell, S. W., & Woolford, J. L., Jr. (1999). Ribosomal protein S14 of Saccharomyces cerevisiae regulates its expression by binding to RPS14B pre-mRNA and to 18S rRNA. *Mol Cell Biol*, *19*(1), 826-834. https://doi.org/10.1128/mcb.19.1.826

Fu, Y., Deiorio-Haggar, K., Anthony, J., & Meyer, M. M. (2013). Most RNAs regulating ribosomal protein biosynthesis in Escherichia coli are narrowly distributed to Gammaproteobacteria. *Nucleic Acids Res*, *41*(6), 3491-3503. https://doi.org/10.1093/nar/gkt055

Gabunilas, J., & Chanfreau, G. (2016). Splicing-Mediated Autoregulation Modulates Rpl22p Expression in Saccharomyces cerevisiae. *PLoS Genet*, *12*(4), e1005999. https://doi.org/10.1371/journal.pgen.1005999

Gahura, O., Hammann, C., Valentova, A., Puta, F., & Folk, P. (2011). Secondary structure is required for 3' splice site recognition in yeast. *Nucleic Acids Res*, *39*(22), 9759-9767. https://doi.org/10.1093/nar/gkr662

Gao, W., Jones, T. A., & Rivas, E. (2021). Discovery of 17 conserved structural RNAs in fungi. *Nucleic Acids Res*, *49*(11), 6128-6143. https://doi.org/10.1093/nar/gkab355

Garrido-Godino, A. I., Gupta, I., Gutierrez-Santiago, F., Martinez-Padilla, A. B., Alekseenko, A., Steinmetz, L. M., Perez-Ortin, J. E., Pelechano, V., & Navarro, F. (2021). Rpb4 and Puf3 imprint and post-transcriptionally control the stability of a common set of mRNAs in yeast. *RNA Biol*, *18*(8), 1206-1220. https://doi.org/10.1080/15476286.2020.1839229

Gelperin, D. M., White, M. A., Wilkinson, M. L., Kon, Y., Kung, L. A., Wise, K. J., Lopez-Hoyo, N., Jiang, L., Piccirillo, S., Yu, H., Gerstein, M., Dumont, M. E., Phizicky, E. M., Snyder, M., & Grayhack, E. J. (2005). Biochemical and genetic analysis of the yeast proteome with a movable ORF collection. *Genes Dev*, *19*(23), 2816-2826. https://doi.org/10.1101/gad.1362105

Gerhart, J. (2014). From feedback inhibition to allostery: the enduring example of aspartate transcarbamoylase. *FEBS J*, *281*(2), 612-620. https://doi.org/10.1111/febs.12483

Ghaemmaghami, S., Huh, W. K., Bower, K., Howson, R. W., Belle, A., Dephoure, N., O'Shea, E. K., & Weissman, J. S. (2003). Global analysis of protein expression in yeast. *Nature*, *425*(6959), 737-741. https://doi.org/10.1038/nature02046

Ghulam, M. M., Catala, M., & Abou Elela, S. (2020). Differential expression of duplicated ribosomal protein genes modifies ribosome composition in response to stress. *Nucleic Acids Res*, *48*(4), 1954-1968. https://doi.org/10.1093/nar/gkz1183

Goguel, V., & Rosbash, M. (1993). Splice site choice and splicing efficiency are positively influenced by pre-mRNA intramolecular base pairing in yeast. *Cell*, *72*(6), 893-901. https://doi.org/10.1016/0092-8674(93)90578-e

Gold, L., Polisky, B., Uhlenbeck, O., & Yarus, M. (1995). Diversity of oligonucleotide functions. *Annu Rev Biochem*, *64*, 763-797. https://doi.org/10.1146/annurev.bi.64.070195.003555

Haller, A., Souliere, M. F., & Micura, R. (2011). The dynamic nature of RNA as key to understanding riboswitch mechanisms. *Acc Chem Res*, *44*(12), 1339-1348. https://doi.org/10.1021/ar200035g

Hooks, K. B., Naseeb, S., Parker, S., Griffiths-Jones, S., & Delneri, D. (2016). Novel Intronic RNA Structures Contribute to Maintenance of Phenotype in Saccharomyces cerevisiae. *Genetics*, *203*(3), 1469-1481. https://doi.org/10.1534/genetics.115.185363

Howe, K. J., & Ares, M., Jr. (1997). Intron self-complementarity enforces exon inclusion in a yeast pre-mRNA. *Proc Natl Acad Sci U S A*, *94*(23), 12467-12472. https://doi.org/10.1073/pnas.94.23.12467

Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S., & O'Shea, E. K. (2003). Global analysis of protein localization in budding yeast. *Nature*, *425*(6959), 686-691. https://doi.org/10.1038/nature02026

Joyce, G. F. (1994). In vitro evolution of nucleic acids. *Curr Opin Struct Biol*, *4*, 331-336. https://doi.org/10.1016/s0959-440x(94)90100-7

Kalisky, T., Dekel, E., & Alon, U. (2007). Cost-benefit theory and optimal design of gene regulation functions. *Phys Biol*, *4*(4), 229-245. https://doi.org/10.1088/1478-3975/4/4/001

Kim, T. Y., Ha, C. W., & Huh, W. K. (2009). Differential subcellular localization of ribosomal protein L7 paralogs in Saccharomyces cerevisiae. *Mol Cells*, *27*(5), 539-546. https://doi.org/10.1007/s10059-009-0077-0

Kohrer, C., Mayer, C., Neumair, O., Grobner, P., & Piendl, W. (1998). Interaction of ribosomal L1 proteins from mesophilic and thermophilic Archaea and Bacteria with specific L1-binding sites on 23S rRNA and mRNA. *Eur J Biochem*, *256*(1), 97-105. https://doi.org/10.1046/j.1432-1327.1998.2560097.x

Komili, S., Farny, N. G., Roth, F. P., & Silver, P. A. (2007). Functional specificity among ribosomal proteins regulates gene expression. *Cell*, *131*(3), 557-571. https://doi.org/10.1016/j.cell.2007.08.037

Leyva-Diaz, E., & Hobert, O. (2019). Transcription factor autoregulation is required for acquisition and maintenance of neuronal identity. *Development*, *146*(13). https://doi.org/10.1242/dev.177378

Libri, D., Stutz, F., McCarthy, T., & Rosbash, M. (1995). RNA structural patterns and splicing: molecular basis for an RNA-based enhancer. *RNA*, *1*(4), 425-436. https://www.ncbi.nlm.nih.gov/pubmed/7493320

Macias, S., Bragulat, M., Tardiff, D. F., & Vilardell, J. (2008). L30 binds the nascent RPL30 transcript to repress U2 snRNP recruitment. *Mol Cell*, *30*(6), 732-742. https://doi.org/10.1016/j.molcel.2008.05.002

Malik Ghulam, M., Catala, M., Reulet, G., Scott, M. S., & Abou Elela, S. (2022). Duplicated ribosomal protein paralogs promote alternative translation and drug resistance. *Nat Commun*, *13*(1), 4938. https://doi.org/10.1038/s41467-022-32717-y

Mandal, M., & Breaker, R. R. (2004). Gene regulation by riboswitches. *Nat Rev Mol Cell Biol*, *5*(6), 451-463. https://doi.org/10.1038/nrm1403

McCown, P. J., Corbino, K. A., Stav, S., Sherlock, M. E., & Breaker, R. R. (2017). Riboswitch diversity and distribution. *RNA*, *23*(7), 995-1011. https://doi.org/10.1261/rna.061234.117

McPheeters, D. S., Stormo, G. D., & Gold, L. (1988). Autogenous regulatory site on the bacteriophage T4 gene 32 messenger RNA. *J Mol Biol*, *201*(3), 517-535. https://www.ncbi.nlm.nih.gov/pubmed/3262167

Miranda-Rios, J., Navarro, M., & Soberon, M. (2001). A conserved RNA structure (thi box) is involved in regulation of thiamin biosynthetic gene expression in bacteria. *Proc Natl Acad Sci U S A*, *98*(17), 9736-9741. https://doi.org/10.1073/pnas.161168098

Mironov, A. S., Gusarov, I., Rafikov, R., Lopez, L. E., Shatalin, K., Kreneva, R. A., Perumov, D. A., & Nudler, E. (2002). Sensing small molecules by nascent RNA: a mechanism to control transcription in bacteria. *Cell*, *111*(5), 747-756. https://www.ncbi.nlm.nih.gov/pubmed/12464185

Muller-McNicoll, M., Rossbach, O., Hui, J., & Medenbach, J. (2019). Auto-regulatory feedback by RNA-binding proteins. *J Mol Cell Biol*, *11*(10), 930-939. https://doi.org/10.1093/jmcb/mjz043

Nomura, M., Gourse, R., & Baughman, G. (1984). Regulation of the synthesis of ribosomes and ribosomal components. *Annu Rev Biochem*, *53*, 75-117. https://doi.org/10.1146/annurev.bi.53.070184.000451

Nomura, M., Yates, J. L., Dean, D., & Post, L. E. (1980). Feedback regulation of ribosomal protein gene expression in Escherichia coli: structural homology of ribosomal RNA and ribosomal protein MRNA. *Proc Natl Acad Sci U S A*, *77*(12), 7084-7088. https://www.ncbi.nlm.nih.gov/pubmed/7012833

Palumbo, R. J., Fuchs, G., Lutz, S., & Curcio, M. J. (2017). Paralog-Specific Functions of RPL7A and RPL7B Mediated by Ribosomal Protein or snoRNA Dosage in Saccharomyces cerevisiae. *G3 (Bethesda)*, *7*(2), 591-606. https://doi.org/10.1534/g3.116.035931

Parenteau, J., & Abou Elela, S. (2019). Introns: Good Day Junk Is Bad Day Treasure. *Trends Genet*, *35*(12), 923-934. https://doi.org/10.1016/j.tig.2019.09.010

Parenteau, J., Durand, M., Morin, G., Gagnon, J., Lucier, J. F., Wellinger, R. J., Chabot, B., & Elela, S. A. (2011). Introns within ribosomal protein genes regulate the production and function of yeast ribosomes. *Cell*, *147*(2), 320-331. https://doi.org/10.1016/j.cell.2011.08.044

Parenteau, J., Durand, M., Veronneau, S., Lacombe, A. A., Morin, G., Guerin, V., Cecez, B., Gervais-Bird, J., Koh, C. S., Brunelle, D., Wellinger, R. J., Chabot, B., & Abou Elela, S. (2008). Deletion of many yeast introns reveals a minority of genes that require splicing for function. *Mol Biol Cell*, *19*(5), 1932-1941. https://doi.org/10.1091/mbc.e07-12-1254

Parenteau, J., Maignon, L., Berthoumieux, M., Catala, M., Gagnon, V., & Abou Elela, S. (2019). Introns are mediators of cell response to starvation. *Nature*, *565*(7741), 612-617. https://doi.org/10.1038/s41586-018-0859-7

Peng, B., Williams, T. C., Henry, M., Nielsen, L. K., & Vickers, C. E. (2015). Controlling heterologous gene expression in yeast cell factories on different carbon substrates and across the diauxic shift: a comparison of yeast promoter activities. *Microb Cell Fact*, *14*, 91. https://doi.org/10.1186/s12934-015-0278-5

Petibon, C., Parenteau, J., Catala, M., & Elela, S. A. (2016). Introns regulate the production of ribosomal proteins by modulating splicing of duplicated ribosomal protein genes. *Nucleic Acids Res*, *44*(8), 3878-3891. https://doi.org/10.1093/nar/gkw140

Plocik, A. M., & Guthrie, C. (2012). Diverse forms of RPS9 splicing are part of an evolving autoregulatory circuit. *PLoS Genet*, *8*(3), e1002620. https://doi.org/10.1371/journal.pgen.1002620

Preker, P. J., & Guthrie, C. (2006). Autoregulation of the mRNA export factor Yra1p requires inefficient splicing of its pre-mRNA. *RNA*, *12*(6), 994-1006. https://doi.org/10.1261/rna.6706

Ptashne, M. (1986). *A genetic switch : gene control and phage [lambda]*. Cell Press ;

Blackwell Scientific Publications.

Ptashne, M., Backman, K., Humayun, M. Z., Jeffrey, A., Maurer, R., Meyer, B., & Sauer, R. T. (1976). Autoregulation and function of a repressor in bacteriophage lambda. *Science*, *194*(4261), 156-161. https://www.ncbi.nlm.nih.gov/pubmed/959843

Ptashne, M., Jeffrey, A., Johnson, A. D., Maurer, R., Meyer, B. J., Pabo, C. O., Roberts, T. M., & Sauer, R. T. (1980). How the lambda repressor and cro work. *Cell*, *19*(1), 1-11. https://www.ncbi.nlm.nih.gov/pubmed/6444544

Ptashne, M., Johnson, A. D., & Pabo, C. O. (1982). A genetic switch in a bacterial virus. *Sci Am*, *247*(5), 128-130, 132, 134-140. https://www.ncbi.nlm.nih.gov/pubmed/6216590

Rain, J. C., & Legrain, P. (1997). In vivo commitment to splicing in yeast involves the nucleotide upstream from the branch site conserved sequence and the Mud2 protein. *EMBO J*, *16*(7), 1759-1771. https://doi.org/10.1093/emboj/16.7.1759

Rangan, R., Huang, R., Hunter, O., Pham, P., Ares, M., & Das, R. (2024). RNA structure landscape of S. cerevisiae introns. *bioRxiv*, 2022.2007.2022.501175. https://doi.org/10.1101/2022.07.22.501175

Reining, A., Nozinovic, S., Schlepckow, K., Buhr, F., Furtig, B., & Schwalbe, H. (2013). Three-state mechanism couples ligand and temperature sensing in riboswitches. *Nature*, *499*(7458), 355-359. https://doi.org/10.1038/nature12378

Rogic, S., Montpetit, B., Hoos, H. H., Mackworth, A. K., Ouellette, B. F., & Hieter, P. (2008). Correlation between the secondary structure of pre-mRNA introns and the efficiency of splicing in Saccharomyces cerevisiae. *BMC Genomics*, *9*, 355. https://doi.org/10.1186/1471-2164-9-355

Romby, P., & Springer, M. (2003). Bacterial translational control at atomic resolution. *Trends Genet*, *19*(3), 155-161. https://doi.org/10.1016/S0168-9525(03)00020-9

Roy, B., Granas, D., Bragg, F., Jr., Cher, J. A. Y., White, M. A., & Stormo, G. D. (2020). Autoregulation of yeast ribosomal proteins discovered by efficient search for feedback regulation. *Commun Biol*, *3*(1), 761. https://doi.org/10.1038/s42003-020-01494-z

Russel, M., Gold, L., Morrissett, H., & O'Farrell, P. Z. (1976). Translational, autogenous regulation of gene 32 expression during bacteriophage T4 infection. *J Biol Chem*, *251*(22), 7263-7270. https://www.ncbi.nlm.nih.gov/pubmed/791948

Ryckelynck, M., Giege, R., & Frugier, M. (2005). tRNAs and tRNA mimics as cornerstones of aminoacyl-tRNA synthetase regulations. *Biochimie*, *87*(9-10), 835-845. https://doi.org/10.1016/j.biochi.2005.02.014

Savageau, M. A. (1974a). Comparison of classical and autogenous systems of regulation in inducible operons. *Nature*, *252*(5484), 546-549. https://doi.org/10.1038/252546a0

Savageau, M. A. (1974b). Optimal design of feedback control by inhibition. *J Mol Evol*, *4*(2), 139-156. https://www.ncbi.nlm.nih.gov/pubmed/4469274

Savageau, M. A. (1975). Significance of autogenously regulated and constitutive synthesis of regulatory proteins in repressible biosynthetic systems. *Nature*, *258*(5532), 208-214. https://www.ncbi.nlm.nih.gov/pubmed/1105191

Schirman, D., Yakhini, Z., Pilpel, Y., & Dahan, O. (2021). A broad analysis of splicing regulation in yeast using a large library of synthetic introns. *PLoS Genet*, *17*(9), e1009805. https://doi.org/10.1371/journal.pgen.1009805

Schlax, P. J., & Worhunsky, D. J. (2003). Translational repression mechanisms in prokaryotes. *Mol Microbiol*, *48*(5), 1157-1169. https://www.ncbi.nlm.nih.gov/pubmed/12787346

Shamoo, Y., Tam, A., Konigsberg, W. H., & Williams, K. R. (1993). Translational repression by the bacteriophage T4 gene 32 protein involves specific recognition of an RNA pseudoknot structure. *J Mol Biol*, *232*(1), 89-104. https://doi.org/10.1006/jmbi.1993.1372

Singh, R., & Valcarcel, J. (2005). Building specificity with nonspecific RNA-binding proteins. *Nat Struct Mol Biol*, *12*(8), 645-653. https://doi.org/10.1038/nsmb961

Spingola, M., Grate, L., Haussler, D., & Ares, M., Jr. (1999). Genome-wide bioinformatic and molecular analysis of introns in Saccharomyces cerevisiae. *RNA*, *5*(2), 221-234. https://doi.org/10.1017/s1355838299981682

Stormo, G. D. (2018). Is Autogenous Posttranscriptional Gene Regulation Common? In N. Rajewsky, S. Jurga, & J. Barciszewski (Eds.), *Systems Biology* (pp. 217-227). Springer International Publishing. https://doi.org/10.1007/978-3-319-92967-5_10

Stormo, G. D., & Ji, Y. (2001). Do mRNAs act as direct sensors of small molecules to control their expression? *Proc Natl Acad Sci U S A*, *98*(17), 9465-9467. https://doi.org/10.1073/pnas.181334498

Torres-Larios, A., Dock-Bregeon, A. C., Romby, P., Rees, B., Sankaranarayanan, R., Caillet, J., Springer, M., Ehresmann, C., Ehresmann, B., & Moras, D. (2002). Structural basis of translational control by Escherichia coli threonyl tRNA synthetase. *Nat Struct Biol*, *9*(5), 343-347. https://doi.org/10.1038/nsb789

Tuerk, C., Eddy, S., Parma, D., & Gold, L. (1990). Autogenous translational operator recognized by bacteriophage T4 DNA polymerase. *J Mol Biol*, *213*(4), 749-761. https://doi.org/10.1016/S0022-2836(05)80261-X

Tuerk, C., & Gold, L. (1990). Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science*, *249*(4968), 505-510. https://www.ncbi.nlm.nih.gov/pubmed/2200121

Vilardell, J., Chartrand, P., Singer, R. H., & Warner, J. R. (2000). The odyssey of a regulated transcript. *RNA*, *6*(12), 1773-1780. https://doi.org/10.1017/s135583820000145x

Vilardell, J., Yu, S. J., & Warner, J. R. (2000). Multiple functions of an evolutionarily conserved RNA binding domain. *Mol Cell*, *5*(4), 761-766. https://doi.org/10.1016/s1097-2765(00)80255-5

von Hippel, P. H., Kowalczykowski, S. C., Lonberg, N., Newport, J. W., Paul, L. S., Stormo, G. D., & Gold, L. (1982). Autoregulation of gene expression. Quantitative evaluation of the expression and function of the bacteriophage T4 gene 32 (single-stranded DNA binding) protein system. *J Mol Biol*, *162*(4), 795-818. https://www.ncbi.nlm.nih.gov/pubmed/6984860

Warf, M. B., & Berglund, J. A. (2010). Role of RNA structure in regulating pre-mRNA splicing. *Trends Biochem Sci*, *35*(3), 169-178. https://doi.org/10.1016/j.tibs.2009.10.004

Winkler, W. C., Cohen-Chalamish, S., & Breaker, R. R. (2002). An mRNA structure that controls gene expression by binding FMN. *Proc Natl Acad Sci U S A*, *99*(25), 15908-15913. https://doi.org/10.1073/pnas.212628899

Wong, E. D., Miyasato, S. R., Aleksander, S., Karra, K., Nash, R. S., Skrzypek, M. S., Weng, S., Engel, S. R., & Cherry, J. M. (2023). Saccharomyces genome database update: server architecture, pan-genome nomenclature, and external resources. *Genetics*, *224*(1). https://doi.org/10.1093/genetics/iyac191

Xue, S., & Barna, M. (2012). Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nat Rev Mol Cell Biol*, *13*(6), 355-369. https://doi.org/10.1038/nrm3359

Yeo, G. W., Van Nostrand, E. L., & Liang, T. Y. (2007). Discovery and analysis of evolutionarily conserved intronic splicing regulatory elements. *PLoS Genet*, *3*(5), e85. https://doi.org/10.1371/journal.pgen.0030085

Figure legends.

Figure 1. Experimental Design and Results

a. Schematic of experimental design. Rpl7 protein is induced by adding galactose to the medium. The *Rpl7a* gene is the *cre*-less version that has eliminated any potential *cis-regulatory elements* (Roy et al., 2020). "insert" refers to different potential regulatory elements inserted into the GFP gene. Proper splicing of those inserted elements leads to GFP protein production.

b.-f. Flow-cytometry histograms of fluorescence signals at 525 nm wavelength to detect GFP protein levels. Replicate experiments are overlaid and can be seen to be nearly identical.

b. yeast cells in the absence of GFP gene to show background fluorescence.

c. The insert contains both introns 1 and 2, and the short exon between them, of *Rpl7b*, under conditions of both Rpl7 induction and uninduced.

d. The insert contains only intron 1 of *Rpl7b*.

e. The insert contains only intron 2 of *Rpl7b*.

f. The insert contains introns 1 and 2 of *Rpl7a*.

g. The insert contains only intron 1 of *Rpl7a*.

Figure 2. Schematic of intron 1 with deletions.

a. The 5' splice site is at position 1 of the intron and the 3' splice site is at position 409. The BP (BP) sequence UGCUAAC is at positions 382-388. The positions of deletions u1-u3 and d1 are indicated, as is the position of the conserved Hooks structure.
b. A diagram of the conserved Hooks structure with the deletions s2-s4 indicated in colors.

Figure 3. Plots of fluorescence values for different intron inserts. The vertical lines show the change upon induction of the Rpl7 protein, and the horizontal lines in each quadrilateral show the changes upon substitution of the consensus BP (UACUAAC) for the wild-type BP (UGCUAAC).

Figure 4. The model for splicing regulation.

a. The intron can form a structure between sequences at near the 5' end (red) and sequences near the BP (yellow) that serves as a splicing enhancer. Rpl7 protein can bind to an alternative structure (Hooks structure) which include the orange and yellow sequences. This eliminates the enhancer structure and makes the BP more distant from the 5' end of the intron.
b. The figure on the left is the proposed structure of the *RPL7B* splicing enhancer, that pairs positions 49-84 with positions 331-356 of intron 1. The structure on the right is stem 3 of the Hooks structure (deleted in s4, shown in Figure 2b). The bases shown in yellow are the same bases shown in mutually exclusive, alternative structures.

Figure 5. Rpl7zz protein binding. Lane 1 is a size ladder with bands of 100 bp. GFP, *RPL7A* and *RPL7B* are the three genes assayed using gene-specific primers for PCR (Table 1). The two strains are GFP with the wild-type *RPL7B* intron 1 sequence (wt) or with the Hooks structure deletion (S1). T shows total RNA, demonstrating that the primers specifically amplify the designated gene. 0 is the RNA from the pull-down of the Rpl7-zz protein at time 0 after induction. 15 is the RNA from the pull-down of the Rpl7-zz protein at 15 minutes after induction.

Figure 1.

    a.   Schematic of assay

Figure 2

a

u1	u2	u3	Hooks structure	d1	

100   200   300   400

BP

(8-70)   (71-133)   (134-197)   (200-355)   (356-381) (382-388)

b

Figure 3

GFP fluorescence of intron variants with and without induction and with wild-type or consensus BP

	Wt BP, uninduced
	Wt BP, induced Rpl7
	Consensus BP, uninduced
	Consensus BP, induced Rpl7

Figure 4

Model of structure switching

a

b

Figure 5

Binding assay gel