MF-PENS: an efficient and versatile nuclear extraction method for plant single-cell sequencing

Lin Du[1, 2 #], Qing Liu[3, #], Hongling Zhou[4, #], Mengyue Zhao[5, #], Fuyan Liu[6], Cheng Tong[7], Mengjiao Zhang[7, 8], Hao Yu[9, *], Zhe Liang[7, *], Jingmin Kang[1, 5 *]

[1]University of Chinese Academy of Sciences, Beijing 100049, China.
[2]College of Life Sciences, Chinese Academy of Sciences, Beijing 100049, China.
[3]State Key Laboratory of North China Crop Improvement and Regulation, College of Life Sciences, Hebei Agricultural University, Baoding 071000, China.
[4]Shenzhen Branch, Guangdong Laboratory for Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of Agriculture and Rural Affairs, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, Guangdong 518116, China.
[5]Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China.
[6]OmicsGang Biotechnology Corporation, 18 Yuan Street, Beijing, 101318, China.
[7]Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, Beijing, 100081, China.
[8]College of Bioscience and Biotechnology, Yangzhou University, Yangzhou 225000，China.
[9]Department of Biological Sciences and Temasek Life Sciences Laboratory, National University of Singapore, Singapore.
[#]These authors contributed equally to this work.

*Correspondence: liangzhe@caas.cn (Zhe Liang), dbsyuhao@nus.edu.sg (Hao Yu), kinzime@hotmail.com (Jingmin Kang)

Abstract

The widespread adoption of single-cell transcriptomics in plants remains constrained by the challenges of isolating high-quality nuclei from tissues with rigid cell walls and complex metabolomes. Here, we present **m**ethanol **f**ixation and **P**ercoll-**e**nriched **n**uclei **s**orting (MF-PENS), a robust strategy designed to tackle these limitations. Distinct from conventional protocols, MF-PENS circumvents the need for fluorescence activated cell sorting (FACS) and effectively reduce RNA leakage, consistently yielding high-quality single-nucleus RNA sequencing (snRNA-seq) data. MF-PENS has been validated across various plant tissues, including fresh and frozen rice roots and leaves, as well as recalcitrant samples such as soybean root nodules, soybean seeds, and tomato fruits. Benchmarking results demonstrate that MF-PENS significantly enhances gene capture efficiency, cell clustering performance, and cell type annotation accuracy. With its broad applicability and superior performance, MF-PENS represents a robust and versatile tool for plant single-cell research, facilitating deeper insights into plant development and biology processes.

Key words: snRNA-seq; Nuclei isolation; MF-PENS; Plant single-cell; Methanol fixation

Introduction

Cells constitute the fundamental units of growth and development in multicellular organisms. The formation of complex tissues and organs is driven by the dynamic interactions among diverse cell types and their specialized functions.[1] Recent studies have highlighted that even morphologically similar cells within the same tissue can exhibit substantial heterogeneity, characterized by distinct gene expression signatures.[2,3] Single-cell RNA sequencing (scRNA-seq) technologies have revolutionized genomic and transcriptomic profiling by offering single-cell resolution. Unlike traditional bulk sequencing, scRNA-seq enables the detection of rare but biologically significant cell populations, the dissection of heterogeneity among phenotypically indistinguishable cells, and the exploration of stochastic variation within genetically identical populations.[4] Furthermore, these technologies facilitate precise cell type annotation, functional characterization, and the inference of cellular dynamics. Consequently, scRNA-seq has been widely adopted across diverse disciplines, including cancer biology, immunology, stem cell research, neuroscience, infectious diseases, and plant stress biology.[5-9]

In plant biology, pioneering scRNA-seq applications focused on establishing foundational resources, such as the construction of high-resolution cellular atlases. For instance, Liu *et al.* and Zhang *et al.* established comprehensive single-cell atlases of rice root tips.[10,11] More recently, the field has pivoted toward elucidating the molecular mechanisms governing plant responses to environmental stresses. Liu *et al.* utilized single-cell transcriptomics to dissect the response of *Arabidopsis thaliana* to osmotic stress,[9] while Li *et al.* characterized the cellular responses to salt stress in cotton. Additionally, Swift *et al.* leveraged these tools to unravel key mechanisms driving the evolution of efficient photosynthetic pathways in C4 plants.[12] These studies represent a paradigm shift in plant single-cell research—transitioning from cellular atlas construction to the investigation of biological functions and regulatory

mechanisms. However, the application of single-cell technologies in plants

continues to lag behind their widespread adoption in animal models.[13] A

primary impediment lies in the intrinsic structural and biochemical complexity

of plant cells[14,15]: they are encased in rigid cell walls and often contain high

concentrations of interfering complex secondary metabolites, such as

polysaccharides and polyphenols. These physicochemical features impose

substantial technical challenges for isolating high-quality protoplasts or nuclei

from plant tissues, which are essential for scRNA-seq or single-nucleus RNA

sequencing (snRNA-seq).[16-19]

Protoplast preparation typically entails prolonged enzymatic digestion to

degrade the cell wall matrix.[20-22] The efficiency of this process is highly

dependent on species and tissue characteristics, precluding the establishment

of universal, standardized protocols. Crucially, cells remain metabolically

active during enzymatic digestion; extended exposure to enzymatic stress can

induce transcriptional artifacts—such as the ectopic activation of

stress-response genes—preferential depletion of rare or enzyme-sensitive cell

populations. These confounding factors ultimately compromise fidelity and

completeness of the cellular landscape reconstructed by scRNA-seq.

Furthermore, protoplast isolation strictly mandates the use of fresh tissue,

imposing strict temporal constraints on sample processing. This requirement

limits experimental flexibility and prevents the retrospective analysis of

archived or frozen specimens.

To circumvent the limitations inherent to protoplast-based methods,

snRNA-seq has emerged as a promising alternative. Farmer *et al.*

demonstrated the reliability of snRNA-seq in plant research using *Arabidopsis*

root tissues, showing its capacity to resolve cell subtypes often obscured in

protoplast-derived datasets.[23] However, the isolation of high-quality plant

nuclei remains a significant technical bottleneck. Current approaches, such as

those applied to *Arabidopsis* and wheat roots,[23,24] typically involve crude

nuclear extraction followed by purification using fluorescence activated cell

sorting (FACS), which enables snRNA-seq. Despite their effectiveness, these methods depend on specialized instrumentation (flow cytometer) that is not universally accessible, thereby imposing a high barrier to entry for many laboratories[12,25,26]. Although some studies have attempted to perform snRNA-seq without FACS, they frequently struggle to maintain the integrity of nucleus.[27] Physical damage during extraction often results in RNA leakage, which manifests as low library complexity (few genes detected per nucleus) or the loss of nuclei during quality control due to insufficient transcript capture[28-31]. Notably, although a few studies have employed methanol fixation in snRNA-seq, their primary purpose was solely sample storage, with no indication that this treatment enhances data quality; furthermore, these methods still rely on FACS for nuclear purification. Moreover, methanol fixation protocols have not clearly specified critical parameters (such as fixation time and methanol concentration) or standardized operating procedures, limiting reproducibility and broad applicability.[32] Collectively, these technical hurdles continue to impede the widespread adoption of single-cell omics in plant research.

To address this gap, we report a robust and versatile nuclei isolation strategy named MF-PENS that integrates methanol fixation with density gradient centrifugation. Designed to circumvent existing technical barriers, this method provides a comprehensive solution featuring (1) full compatibility with frozen tissues, (2) a FACS–free purification workflow, and (3) a fixation-based preservation system that ensures optimal nuclear integrity and RNA stability. We demonstrate the universality of this approach across a diverse array of plant tissues—including roots, leaves, root nodules, fruits, seeds, and callus—consistently yielding high-quality input for snRNA-seq. By establishing a user-friendly, cost-effective, and highly reproducible standard, this method empowers the research community to fully leverage single-cell resolution for advancing plant developmental biology, functional genomics, and agricultural innovation.

Results

Establishment of the MF-PENS workflow for plant single-nucleus sequencing

To optimize a nuclei isolation protocol suitable for high-quality plant snRNA-seq, we systematically benchmarked four distinct fixation conditions combined with the Percoll-enhanced nuclei separation (PENS) purification: 0.1% formaldehyde fixation (0.1%FF-PENS), 1% formaldehyde fixation (1%FF-PENS), 100% methanol fixation (MF-PENS), and a no-fixation control (NT-PENS). In detail, plant tissues were sectioned into approximately 0.5 cm^2 fragments and subjected to their respective fixation treatment, with the exception of the NT-PENS method which was proceeded directly to the next step without fixation. Subsequently, nuclei were released and purified via Percoll density gradient centrifugation. The isolated nuclei then underwent microfluidic-based barcoding, library construction, and high-throughput sequencing **(Figure 1A)**. We evaluated the performance of these methods by accessing data quality metrics and downstream analytical utility, including clustering resolution and the identification of differentially expressed genes (DEGs). Furthermore, we validated the robustness and versatility of MF-PENS across a diverse array of sample types, ranging from standard vegetative tissues (roots and leaves) to recalcitrant reproductive and symbiotic tissues, including root nodules, fruits, seeds, and callus **(Figure 1A)**.

MF-PENS yields high-quality single-nucleus transcriptomes

To assess the impact of fixation methods on snRNA-seq data quality, we systematically benchmarked four approaches in rice (*Oryza sativa*) root tissues (**Figure 1B-E, Supplementary Table S1**). All experiments were performed in biological replicates to ensure reproducibility. During data preprocessing, stringent quality control (QC) criteria were applied, including the removal of doublets/multiplets, empty droplets, and low-quality nuclei exhibiting high mitochondrial content (> 5%) or signs of enzymatic degradation. Notably, MF-PENS retained the highest number of high-quality nuclei (9,170 and 8,468) following identical QC procedures**(Figure 1B)**. Sequencing saturation analysis also demonstrated that MF-PENS consistently outperformed other methods across all examined read depths. MF-PENS replicates exhibited superior gene detection curves, recovering significantly

more genes (P <0.05, Student's t-test) and producing higher unique molecular

identifier counts (UMIs) than other methods at equivalent depths. These

findings demonstrate that MF-PENS confers enhanced transcript capture

efficiency, establishing it as a robust approach for plant snRNA-seq **(Figure**

1C).

Quantitative assessment confirmed that MF-PENS generated significantly

higher post-QC UMIs and gene numbers compared to other methods (P

<0.05), with high reproducibility **(Figures 1D and S1A)**. To assess nuclear

purity, we quantified the proportions of chloroplast- and mitochondrial-derived

transcripts **(Figure 1E)**. All methods exhibited low levels of organelle-derived

transcripts, indicating minimal cytosolic contamination and high integrity of the

extracted nuclear RNA. Additionally, Pearson correlation analysis **(Figure S1B)**

demonstrated strong global similarity across the datasets. Interestingly, while

0.1%FF-PENS showed the highest correlation with NT-PENS, MF-PENS

exhibited a distinct transcriptomic profile. We hypothesize that this divergence

reflects the superior capacity of MF-PENS to preserve fragile, transcriptionally

active   nuclei,   thereby   capturing   biologically   relevant   patterns

underrepresented in other methods. Collectively, these results highlight the

superiority of MF-PENS performance in generating optimal snRNA-seq data

quality, characterized by higher nuclei yield, robust library complexity, and

enhanced transcriptomic fidelity.

MF-PENS enhance cell clustering performance and biological DEGs

identification

While UMIs and gene detection rates are fundamental quality indicators,

robust clustering and accurate marker identification are pivotal prerequisites

for resolving biological heterogeneity.[33-36] To evaluate clustering performance,

we integrated biological replicates for each method. Uniform manifold

approximation and projection dimensionality reduction (UMAP) visualization

revealed negligible batch effects, validating experimental reproducibility and

ensuring the rigor of downstream comparisons **(Figure 2A)**. At a fixed

resolution yielding nine clusters, MF-PENS and 1%FF-PENS produced

discrete, well-separated populations, whereas NT-PENS and 0.1%FF-PENS

exhibited   significant   aggregation   and   boundary   overlap,   indicating

compromised clustering capacity **(Figure 2B)**. This visual assessment was

corroborated quantitatively by the Modularity score[37-39], a graph-based metric

that quantifies the contrast between intra- and inter-cluster connectivity, with

higher values indicating better clustering result. MF-PENS achieved the

highest Modularity score surpassing all other methods **(Figure 2B)**, thereby

confirming its superior capacity to resolve distinct cell populations.

The advantages of MF-PENS in clustering resolution were further

substantiated by functional enrichment analysis. Compared to other methods,

MF-PENS captured a broader spectrum of biological pathways relevant to

specific root cell types. For instance, in cluster 7, MF-PENS uniquely enriched

for terms such as "response to external biotic stimulus" (GO:0043207),

"intraspecies interaction between organisms" (GO:0051703), "response to

bacteria" (GO:0009617)—functional categories consistent with the molecular

signature of root epidermis cells involved in environmental sensing and

defense.[40-42] Similarly, cluster 0 derived from MF-PENS exhibited significant

enrichment in meristem-associated pathways, including ribosome (ko03010),

spliceosome (ko03040), and ribosome biogenesis in eukaryotes (ko03008),

supporting the identification of this cluster as meristematic cells actively

engaged in protein synthesis and transcriptional regulation **(Figures S1C-F,**

S2A-D).[43,44] Collectively, these results demonstrate that the superior clustering

fidelity of MF-PENS directly translates into more accurate and biologically

meaningful cell type resolution.

We next performed differential expression analysis to evaluate marker

specificity across the four methods. As visualized in the bubble plot of the top

10 upregulated DEGs (Up-DEGs) **(Figure 2C)**, DEGs detected from MF-PENS

exhibited high specificity, with expression restricted to their corresponding

cluster. Conversely, markers from the other three methods displayed diffuse

expression patterns across multiple clusters. Statistically, MF-PENS

significantly outperformed other methods in detecting upregulated,

downregulated, and total differentially expressed genes **(Figure 2D)**. To

assess biological fidelity, we analyzed the top 100 up-DEGs; MF-PENS

identified the highest number of unique markers (70) compared to NT-PENS

(26), 0.1%FF-PENS (54), and 1%FF-PENS (38) **(Figure 2E, Table S3-4)**.

Crucially, cross-referencing these unique genes with the *PlantCellMarke*r

database[45] (https://www.tobaccodb.org/pcmdb/homePage) revealed that

MF-PENS achieved the highest validation rate (31.43%) (**Table S2**),

substantially surpassing other methods. These findings indicate that MF-PENS

yields more biologically meaningful, cluster-specific signatures, enhancing

downstream interpretability.

Collectively, these results highlight three pivotal advantages of MF-PENS in

plant single-cell transcriptomics: (1) enhanced performance in cell clustering,

(2) superior sensitivity in resolving cell-type-specific functional pathways, and

(3) improved fidelity in identifying biologically validated DEGs. Consequently,

MF-PENS not only ensures high data quality but also provides opportunity for

more effectively mining deep biological insights into plant cellular

heterogeneity.

MF-PENS enables robust nuclei isolation from both fresh and frozen

plant tissues

Compatibility with frozen samples is essential for experimental flexibility and

the utilization of biobanked specimens. To validate the performance of

MF-PENS on cryopreserved material, we selected rice roots and leaves as

representative belowground and aboveground tissues. To mimic real-world

transport and storage logistics, fresh samples were flash-frozen in liquid

nitrogen, stored on dry ice for 15 days, and subsequently transferred to -80°C

for another 15 days.

We rigorously benchmarked MF-PENS by processing matched fresh and

frozen samples (n=2 biological replicates each) from root **(Figures 3A–G)** and

leaf **(Figures 3H–P)** tissues, followed by snRNA-seq with stringent quality

control.

In root tissue, MF-PENS achieved robust gene detection efficiency across all

sequencing depths for both fresh and frozen replicates, confirming its

suitability for transcriptomic profiling archival materials **(Figure 3B)**.

Quantitative comparison of key quality metrics—including UMIs, detected

genes, and mitochondrial gene percentages — showed no significant

differences between fresh and frozen samples **(Figures 3C, D)**. These results

demonstrate that MF-PENS effectively preserves transcriptomic integrity

independent of storage conditions. Furthermore, UMAP confirmed the robust

data integration, revealing distinct cell clusters with negligible batch effects

attributable to storage conditions **(Figures 3E, S3A and S3B)**. Using reported

canonical marker genes, we annotated eight distinct cell types **(Figures 3F,**

S4A-B). Cellular composition remained highly conserved between conditions,

except for a specific reduction in root hair cells in frozen samples, likely due to

physical shearing induced by cryopreservation **(Figures 3G)**.

Parallel experiments in leaf tissues yielded results concordant with those

observed in roots. Fresh and frozen leaf samples exhibited high consistency

across all key metrics, including the proportions of chloroplast-derived

transcripts **(Figures 3I-M)**. Unsupervised clustering confirmed the effective

integration of fresh and frozen samples, resolving 12 clusters that were

successfully annotated to six major leaf cell types **(Figures 3N, O, S3C, S3D,**

S5A-B). Consistent with root tissues, cell type proportions in leaves remained

largely stable across storage conditions **(Figure 3P)**.

Collectively, by consistently yielding high-quality snRNA-seq data from both

fresh and frozen tissues (roots and leaves), MF-PENS demonstrates

remarkable adaptability across diverse tissue types and storage conditions.

This capacity to preserve transcriptomic integrity underscores the utility of

MF-PENS for flexible experimental designs, enabling the effective exploitation

of archived or transported plant resources.

MF-PENS demonstrates broad versatility across diverse and recalcitrant

plant tissues

Agronomically important crop tissues often harbor complex metabolic

profiles.[46,47] For instance, soybean seeds are replete with lipids[48], whereas

tomato fruits accumulate high concentrations of polysaccharides.[49,50] Isolating

high-quality nuclei from such recalcitrant tissues poses significant technical

hurdles that severely impede snRNA-seq applications.[19,51] To benchmark

MF-PENS across diverse biochemical contexts, we selected four
representative tissue types characterized by distinct physicochemical

properties **(Figure 4A)**: infected tissue (soybean root nodules)[52,53], oil-rich

tissue (soybean seeds)[54], polysaccharide-rich tissue (tomato fruit)[55], and

fragile tissue (maize callus).[56-58] Following stringent quality control, MF-PENS

consistently recovered high-quality nuclei from all biological replicates (n=2),
yielding between 12,725 and 17,565 nuclei per library **(Figure 4B)**.
328
Sequencing metrics further confirmed the robust performance of MF-PENS
across tissues of varying complexity. Specifically, soybean seeds yielded an
average of 3,815 genes and 7,446 UMIs per nucleus. Soybean root nodules

achieved an average of 2,136 genes and 3,128 UMIs per nucleus—a marked

improvement over previous datasets, which reported averages of only
~1,000–1,600 genes and ~1,600–2,200 UMIs across developmental

stages.[59,60] Similarly, tomato fruits exhibited robust capture (2,349 genes;

5,137 UMIs), while maize callus demonstrated the highest efficiency,
averaging 4,535 genes and 8,724 UMIs per nucleus. **(Figure 4C)**. Furthermore,
mitochondrial gene proportions remained consistently low across all libraries
(Figure 4D), indicating minimal cytoplasmic contamination and efficient
nucleus isolation. High concordance between biological replicates **(Figure 4E)**
further corroborated the robustness of MF-PENS for profiling biochemically
recalcitrant tissues.
343
To investigate tissue-specific gene expression patterns, we performed
unsupervised clustering on the integrated datasets. UMAP visualization
revealed distinct, well-separated cell populations **(Figures S3E-H)**,
demonstrating the preservation of cell-type identities. Differential expression
analysis identified robust cluster-specific gene expression signatures,
visualized via heatmaps of the top up-DEGs **(Figures 4F-I)**. These potential
markers exhibited exquisite cluster specificity with minimal background noise,
highlighting the resolution of MF-PENS in dissecting transcriptional
heterogeneity within complex tissues.
353

354 Collectively, these results demonstrate that MF-PENS effectively circumvents
355 the technical barriers associated with metabolically complex plant materials.
356 As a versatile and robust method, MF-PENS empowers researchers to
357 generate high-quality single-cell data from recalcitrant tissues, thereby
358 facilitating deeper insights into plant development and physiology.
359

MF-PENS enables single-nucleus level dissection of plant heat stress
responses
The scRNA-seq permits the high-resolution dissection of transcriptomic
landscapes, unveiling lineage-specific activation patterns of key
stress-response pathways.[61-63] Beyond enhancing nuclear yield, MF-PENS
confers a critical advantage via rapid metabolic inactivation, effectively "locks"
the intracellular transcriptional state at the precise moment of sampling[64]. This
instantaneous preservation mitigates transcriptional artifacts induced by
mechanical stress or environmental fluctuations during processing, ensuring
that sequencing data faithfully reflect the physiological status of the plant *in*
situ. This capability enable MF-PENS serves as a robust, high-fidelity tool for
elucidating the mechanisms of plant stress resilience. To evaluate this utility,
we profiled rice leaves under normal (Control) and heat stress (Heat)
conditions using both NT-PENS and MF-PENS, generating eight datasets
across two biological replicates.
375

At the sample level, integrated analysis demonstrated successful batch
correction and data integration for both methods **(Figures 5A, B)**. B
Differential expression analysis (Control vs. Heat) revealed that MF-PENS
captured a significantly broader repertoire of heat-responsive genes, with ~80%
of up-DEGs being exclusive to the MF-PENS datasets **(Figure 5C)**. Gene
ontology enrichment (GO) analysis further confirmed that these 1,247
MF-PENS-exclusive genes were significantly enriched in "response to heat"
(GO:0009408), whereas DEGs specific to NT-PENS lacked such functional
relevance **(Figures 5D, E and Table S4)**. These findings underscore the
superior sensitivity of MF-PENS in capturing the full spectrum of
stress-responsive transcriptional changes.
387

We next assessed resolution at the cell-type level via unsupervised clustering
and canonical marker-based annotation. NT-PENS datasets annotated only
five cell types and exhibited poor reproducibility between replicates, with
minimal shifts in cellular composition observed under heat stress **(Figures 5F**
S8A). Furthermore, cell populations in NT-PENS appeared mixed on the
UMAP, confounding downstream bioinformatic interrogation. In stark contrast,
MF-PENS successfully delineated nine distinct cell lineages with high
consistency across biological replicates**(Figures 5H, S8B)**.
396
At the level of transcriptional regulation, we computed DEGs for each
annotated cell type for both methods **(Figures S6C, D)**. Under identical
statistical thresholds, MF-PENS identified a substantially higher number of
DEGs per cell type. This enhanced sensitivity suggests that MF-PENS
captures cellular heterogeneity with greater fidelity, thereby accurately
reconstructing cell-type-specific activation patterns **(Figures 5G, I)**.
Comparative analysis of four shared cell types (mesophyll, phloem, epidermal,
and vascular) revealed that MF-PENS-specific DEGs were significantly
enriched in heat stress response-responsive pathways, while
NT-PENS-specific DEGs showed no such functional enrichment **(Figures 5J,**
S8E-F, S7A-J and Table S5, 6). Additionally, DEGs identified within the four
cell types uniquely resolved by MF-PENS (bulliform, fiber, guard, and initial
cells) also exhibited robust enrichment for heat-response functions **(Figure**
5K). These results further validate the superior capability of MF-PENS in
resolving cell type-specific transcriptional regulation patterns under stress.
412
413   In summary, our multi-level analysis demonstrates that MF-PENS provides a
414   comprehensive and high-fidelity representation of heat-induced transcriptomic
415   remodeling. MF-PENS can serve as a robust, high-resolution platform for
416   elucidating plant stress resilience mechanisms and environmental adaptation.
417
MF-PENS enables precise cell subtyping and nuanced transcriptional
profiles
Resolving cellular heterogeneity at the level of subpopulations is critical for
elucidating lineage trajectories, functional compartmentalization, and the
developmental roles of specific cell subsets within plant tissues.[19,65] For
instance, rice leaf epidermis comprises distinct upper and lower layers, each

characterized by divergent gene regulatory networks and developmental

functions.[66-68] However, due to their high transcriptional similarity, previous

scRNA-seq data struggled to reliably distinguish these subtypes.[69,70]

MF-PENS addresses this challenge by generating high-quality snRNA-seq

data capable of robustly separating transcriptionally similar cell populations.

First, following the strategy described by Xia Keke *et al.*[66] to distinguish upper

and lower epidermal cells, we generated high-resolution Stereo-seq data from

rice leaves. Based on cell wall morphology and spatial coordinates , we

annotated upper and lower epidermal cells *in situ* as ground truth **(Figure 6A)**.

We utilized these spatially resolved cells to derive specific up-DEGs for the

upper and lower epidermis, establishing them as Stereo-seq-derived markers.

Subsequently, we extracted all epidermal cells from NT-PENS (epidermal) and

MF-PENS (bulliform, guard, and epidermal) datasets under normal growth

conditions. Leveraging Stereo-seq-derived markers as highly variable gene

(HVGs) for cell clustering, we successfully resolved two epidermal

subpopulations for both datasets with highly consistent cell proportions across

biological replicates **(Figures 6A-B, E-F)**. GO analysis revealed that cluster 0

in both datasets was significantly enriched for biological processes associated

with upper epidermal cell function, including photosynthesis, light harvesting,

and photorespiration **(Figures S8C, D, G)**, validating cluster 0 as upper

epidermal cells. Using Stereo-seq-derived markers,[66] we established the

ground truth identities of the upper and lower epidermal cells for our

snRNA-seq dataset. Accordingly, we used these validated epidermal cell

identities as ground truth annotations for subsequent analysis **(Figures 6C, F)**.

Next, we performed unsupervised clustering using Seurat,[71] employing normal

HVGs rather than Stereo-seq-derived markers on both datasets. The resulting

clusters were then benchmarked against the Stereo-seq-defined ground truth

labels. MF-PENS achieved a classification accuracy of 84% for lower

epidermis and 78% for upper epidermis, significantly outperforming NT-PENS

(59% and 67%, respectively) **(Figure 6B)**. UMAP visualization confirmed that

MF-PENS clusters were discrete and aligned closely with ground truth labels,

whereas NT-PENS data exhibited considerable mixing **(Figures 6D, G)**. This
classification accuracy was highly reproducible across biological replicates,
supporting the robustness of MF-PENS in preserving fine-grained
transcriptional identities **(Figures 6E, H)**. Functional characterization of
MF-PENS-derived clusters confirmed that the upper epidermal population was
enriched for photosynthesis-related terms, while the lower epidermal
population exhibited enrichment for biosynthetic and metabolic processes,

consistent with their known functions[66] **(Figures 6I, J, and Table S7-9)**.

In summary, MF-PENS effectively surmounts the resolution limits of
conventional methods, enabling the high-precision identification of fine-grained
cell subtypes. This capability establishes MF-PENS as a powerful tool for
dissecting the functional heterogeneity and specialization of plant tissues.
470

Discussion
Despite the transformative impact of single-cell technologies in animal
research, their application in plant systems has been impeded by the structural
rigidity of cell walls and the prevalence of interfering secondary metabolites.
These physical and biochemical barriers often compromise nuclei integrity and
yield, limiting the utility of scRNA-seq. To circumvent these hurdles, we
developed MF-PENS, a novel nucleus isolation method that integrates
methanol fixation with Percoll density gradient centrifugation. Unlike traditional
protocols that rely on fresh tissue and FACS, our approach utilizes methanol
fixation to rigidify the nuclear envelope and arrest transcriptional activity
immediately upon sampling **(Figure 1A)**. MF-PENS demonstrates superior
performance in preserving nuclear morphology and enhancing transcript
capture efficiency compared to standard methods. Our data confirms that
MF-PENS not only improves sequencing metrics—such as nuclear yield and
integrity **(Figures 1B-E)**—but also significantly boosts the sensitivity of
differential gene expression analysis **(Figure 2)**.
487

A critical limitation of current plant snRNA-seq protocols is the reliance on
fresh tissue, which restricts experimental design and logistical flexibility. A
pivotal advantage of MF-PENS is its compatibility with cryopreserved samples.
We demonstrated that MF-PENS maintains high data fidelity in both fresh and
frozen tissues across distinct organs (leaves and roots) **(Figure 3)**. This

capability is transformative for plant science, as it enables the retrospective
analysis of biobanked specimens and facilitates sample collection from remote
field sites where immediate processing is impractical.
496
Furthermore, we validated the robustness of MF-PENS on biochemically
complex, "recalcitrant" tissues—including polysaccharide-rich tomato fruits,
lipid-laden soybean seeds, symbiont-containing root nodules, and
polyphenol-rich maize callus **(Figure 4A)**. By successfully generating
high-quality transcriptomic profiles from these diverse systems, MF-PENS
proves its broad applicability, overcoming the metabolite interference that
typically hampers plant nuclei isolation.
504
Beyond technical robustness, MF-PENS delivers the high-resolution fidelity
necessary to dissect fine-scale biological heterogeneity. In rice leaves, the
method achieved superior clustering consistency, resolving nine distinct cell
lineages **(Figure 5H)**. Crucially, MF-PENS demonstrates pivotal utility in
decoding plant stress responses: under heat stress, it exhibited exceptional
sensitivity, capturing a comprehensive repertoire of heat-responsive genes
and unveiling specific regulatory pathways that were obscured in lower-quality
datasets **(Figures 5J-K)**. This capability is essential for elucidating the
complex, cell-type-specific mechanisms underlying plant stress resilience.
514
Furthermore, we addressed the long-standing challenge of distinguishing
transcriptionally similar cell subtypes, specifically the adaxial (upper) and
abaxial (lower) epidermal layers. By leveraging Stereo-seq as referent ground
truth, we benchmarked the resolving power of MF-PENS against NT-PENS.
The results demonstrate that MF-PENS accurately discriminates between
these closely related epidermal lineages, capturing distinct gene signatures
and functional specializations **(Figure 6)**. This capacity to resolve subtle
transcriptional differences underscores the method's potential for investigating
complex developmental lineages.
524
While MF-PENS consistently yields robust snRNA-seq data across a variety of
tissues, several limitations remain. First, while we validated the method on four
major crop species, its applicability to a wider range of plant taxa, particularly
those with extreme secondary metabolite profiles, remains to be
comprehensively assessed. Second, this study did not include samples with

distinctive tissue characteristics such as bark, xylem, and strawberry fruit. Bark
and xylem are highly lignified tissues with extremely thickened cell walls rich in
lignin, cellulose, and other secondary metabolites, whereas tissues like
strawberry fruit contain substantially larger nuclei. These features may impede
efficient nuclear release, compromise nuclear integrity, or cause abnormal
sedimentation during Percoll density gradient centrifugation. For such
challenging tissues, we recommend targeted optimization strategies,

specifically extending the fixation duration to 45–60 minutes and increasing the

centrifugal force to 1000–1500 g during gradient centrifugation to ensure better

structural preservation and recovery. In addition, for plant organs where
obtaining sufficient sample material is challenging, we recommend
homogenization using steel beads directly within the centrifuge tube.
Compared to traditional mortar and pestle grinding, this method significantly
reduces sample loss; in such cases, a minimal sample amount (just enough to
cover the bottom of the tube, ~0.1 g) is sufficient to meet the requirements for
library construction. Third, it is important to note that the choice of fixative may
introduce selection bias. Specifically, methanol fixation may affect nuclear
recovery efficiency in certain cell types, such as highly vacuolated cells,
potentially leading to their underrepresentation in the final dataset. Fourth, this
study focused exclusively on transcriptomics. The compatibility of MF-PENS
with other single-nucleus multi-omics modalities, such as snATAC-seq and
snCUT&Tag, requires further experimental optimization and validation. Finally,
our study evaluated only methanol and formaldehyde as fixatives and has not
systematically compared or tested alternative fixatives—such as ethanol,
acetone, or mixed fixative solutions—for their effects on nuclear integrity, RNA
quality, and downstream applications, which represents an additional limitation
of the current approach.

Notably, in the context of rapidly advancing spatial omics and multi-omics
integration, MF-PENS demonstrates broad application potential. Of particular
importance is its strong synergy with high-resolution spatial transcriptomics
technologies such as Stereo-seq. While Stereo-seq provides precise spatial
localization, its per-spot (e.g., at bin20 resolution, corresponding to a 10 µm²
pixel) gene capture depth is generally lower than that of snRNA-seq. Our
analyses show that the high-quality snRNA-seq data generated by MF-PENS

exhibit strong concordance with Stereo-seq data in terms of cell type
annotation and gene expression profiles **(Figure 6)**, effectively compensating
for Stereo-seq's limited transcriptomic coverage. This complementarity
positions MF-PENS as an ideal upstream tool for integrated single-nucleus
and spatial transcriptomics workflows, laying a solid foundation for future
multidimensional characterization of cell types, functional states, and spatial
organization in plant tissues.
572

Conclusion

In summary, MF-PENS represents a low-barrier, high-performance solution for
plant single-nucleus transcriptomics. By effectively addressing the challenges
of nuclear isolation in metabolite-rich and frozen specimens, this method
significantly expands the accessibility and scope of plant single-cell research.
The enhanced sensitivity, reproducibility, and biological interpretability
provided by MF-PENS pave the way for a paradigm shift from bulk tissue
analyses to high-resolution, cell-type-specific investigations. We anticipate that
the widespread adoption of MF-PENS will facilitate a deeper mechanistic
understanding of plant development, stress resilience, and environmental
adaptation, ultimately contributing to advancements in basic plant biology and
sustainable crop improvement.
585

Materials and methods

Plant materials and growth conditions

The plant materials used in this study included: rice roots (*Oryza sativa* L. cv. Zhonghua 11, 3 days post-germination), rice leaves (*Oryza sativa* L. cv. Zhonghua 11, 14 days post-germination with an additional 3-hour treatment), soybean nodules (*Glycine max* L. cv. Williams 82, 24 days post-inoculation), soybean seeds *(Glycine max* L. cv. Zhonghuang 13, 1-day post-germination), tomato fruits (*Solanum lycopersicum* cv. Ailsa Craig, red fruits at 14 days post-anthesis), and maize callus tissues (*Zea mays* L. inbred line B73). For the salt-treated rice leaf samples, the culture medium was replaced with a solution containing 150 mM NaCl at 14 days post-germination, followed by an additional 3 hours of incubation before sample collection. The control group underwent the same procedure, except that a fresh culture medium without NaCl was used during the replacement step.

Plant tissue fixation

In this study, a standard initial sample size of approximately 0.2 g (fresh weight) was used for all experiments.

No-treatment group: Plant samples were rapidly cut into ~0.5 cm² fragments using a razor blade and subjected to no fixation. The tissue fragments were washed twice with NIBTA buffer (10 mM Tris-HCl (pH 8.0), 10 mM EDTA (pH 8.0), 80 mM KCl, 1 mM Spermine-4HCl, 1 mM Spermidine-3HCl, 2% PVP-40, 0.3% Triton X-100, 1 mM DTT, 1× ProtectRNA™ RNase Inhibitor, 1× Protease Inhibitor Cocktail) and immediately processed for subsequent nuclear isolation.

0.1% formaldehyde fixation group: Plant samples were rapidly cut into ~0.5 cm² fragments using a razor blade and fixed on ice for 30 minutes in ice-cold NIBTA buffer containing 0.1% formaldehyde. After fixation, the tissue was washed twice with ice-cold NIBTA buffer and immediately processed for nuclear isolation.

1% formaldehyde fixation group: Plant samples were rapidly cut into ~0.5 cm² fragments using a razor blade and fixed on ice for 30 minutes in ice-cold NIBTA buffer containing 1% formaldehyde. After fixation, the tissue was

washed twice with ice-cold NIBTA buffer and immediately processed for
nuclear isolation.

100% methanol fixation group: Fresh plant samples were rapidly cut into
~0.5 cm² fragments using a razor blade, or frozen plant samples were crushed
in liquid nitrogen with a pestle into fragments no larger than 0.5 cm² (avoiding
grinding into a powder), followed by fixation in ice-cold 100% methanol on ice
for 30 minutes. After fixation, the tissue was washed twice with ice-cold NIBTA
buffer and immediately processed for nuclear isolation.

Nucleus isolation
The tissue was transferred to a petri dish on ice and chopped with a Gillette
razor blade for approximately 2 minutes. The chopped tissue was
resuspended in NIBTA buffer and homogenized on ice for 5 minutes. The

homogenate was then filtered through a 40 µm strainer, and the filtrate was

collected into a new 50 mL centrifuge tube. The tube was centrifuged at 1,260

g for 10 minutes at 4℃. The supernatant was discarded, and the tissue pellet

was resuspended in 4 mL of NIBTA buffer. In a new 15 mL centrifuge tube, 70%
Percoll solution (3.5 mL Percoll and 1.5 mL NIBTA buffer) was added, and the
4 mL resuspended tissue homogenate was carefully overlaid onto the 70%

Percoll layer. The tube was centrifuged at 650 g for 30 minutes at 4℃, during

which most nuclei banded at the interface between the NIBTA buffer and the
Percoll layer. The nuclei band was gently collected into a new 15 mL centrifuge
tube, 10 mL of NIBTA buffer was added, and the tube was centrifuged again at
1,260 g for 5 minutes. The nuclei pellet was washed twice with NIBTA buffer,
then resuspended in PBS containing 0.04% BSA. After counting the nuclei, the
suspension was kept on ice for later use (it is recommended not to exceed 30
minutes).

Single-nucleus RNA library construction and sequencing
The single-nucleus RNA-seq libraries were prepared as previously described
with DNBelab C Series High-throughput Single-Cell RNA Library Preparation
Set V2.0 (MGI, 940-000508-00), and Sequencing was performed using
DNBSEQ T7.

Spatial transcriptomics library construction

Stereo-seq[72] was used to construct spatial transcriptomics libraries. The leaf was embedded in OCT and sectioned into 10-micron slices. These slices were then adhered to Stereo-seq chips. The chips containing tissue sections were incubated at 37 ° C for 3 min. Permeability, reverse transcription, tissue removal, cDNA release, and library preparation were next performed using the previous methods.[66] Finally, DIPSEQ T10 was used for sequencing.

Preprocessing of snRNA-seq data

The snRNA-seq data were processed and analyzed using DNBelab_C_Series_HT_single cell-analysis-software (version 2.0),[73] an open-source and flexible pipeline. Sequencing reads were aligned to respective reference genomes: rice reads to MSU v7.0,[74] soybean reads to GCF_000004515.6,[75] tomato reads to GCF_036512215.1,[76] and maize reads to GCF_902167145.1.[77] Stringent quality control measures were implemented across all samples. Potential doublets or multiplets were identified and filtered using the DoubletFinder package,[78] followed by the removal of unbound or apoptotic cells. We established strict inclusion criteria for cell quality: (1) cells expressing fewer than 200 or more than 6,000 genes were excluded; (2) cells with UMIs below 500 or exceeding 40,000 were removed; (3) cells demonstrating mitochondrial gene expression exceeding 5% of total expression were discarded. These rigorous quality control steps ensured the retention of high-quality cells for subsequent bioinformatics analysis.

Cell clustering and visualization

We processed the snRNA-seq data using the Seurat v4.3.2 package.[71] To normalize gene expression across cells, we applied the "NormalizeData" function, which utilizes a global-scaling normalization method. This approach involves dividing the gene expression values by the total expression per cell, scaling the results by a factor of 10,000 (default), and applying a natural log

transformation to stabilize the variance. To capture the most biologically
relevant transcriptional variations, we identified the top 2,000 highly variable
genes using the "FindVariableFeatures" function. To ensure robust integration
of datasets and minimize batch effects, we employed Canonical Correlation
Analysis (CCA) through the "FindIntegrationAnchors" and "IntegrateData"
functions, which align shared biological features across samples. For
dimensionality reduction, we performed principal component analysis (PCA)
using the "RunPCA" function (npcs = 40) to extract the most informative
components of variation. Subsequently, we applied graph-based clustering
with the Louvain algorithm via the "FindClusters" function (resolution = 0.3) to
group cells based on their transcriptional profiles. Finally, to visualize the
high-dimensional data in an interpretable two-dimensional space, we utilized
Uniform Manifold Approximation and Projection (UMAP) with the "RunUMAP"
function (dims = 1:40, min.dist = 0.3). This approach preserves the local and
global structure of the data, enabling clear visualization of distinct cell clusters
and their relationships.
703

Cell type annotation of rice leaf and root
In the rice leaf datasets, both NT-PENS and MF-PENS datasets were
clustered into ten clusters **(Figure S3B)**, which were subsequently assigned to
specific cell types. Epidermal cells were identified in clusters 3 and 4 based on
the marker genes *Os03g52390* and *Os02g39930*.[79] Cluster 8 was annotated
as fiber, supported by the specific expression of the fiber marker genes
Os11g10310, *Os11g10320*, *Os12g02320*, and *Os11g37900*.[70] Cluster 9 was
assigned as guard cell, as indicated by the high expression of the guard cell
marker *Os03g10090*.[70] Cluster 7 was annotated as phloem, characterized by
the expression of *Os09g28510*, *Os05g03934*, and *Os04g14220*.[70,79] Cluster 6
was identified as vascular, based on the expression of *Os10g14870*.[70] Finally,
clusters 0 and 1 were annotated as mesophyll, supported by the expression of
Os12g17600, *Os01g41710*, and *Os07g37240*.[80]
717

In the rice root datasets, NT-PENS and MF-PENS were also clustered into ten
clusters **(Figure S3D)**, and cell types were similarly annotated. Cluster 7 was
assigned as phloem, supported by the marker genes *Os09g28510*,
Os05g34980, *Os05g04000*, and *Os02g37070*.[45] Cluster 0 was annotated as
cortex, based on the expression of *Os11g02350*, *Os09g12970*, *Os04g59260*,
Os01g68580, *Os12g02300*, *Os04g46810*, *Os02g44310*, and *Os01g68589*.[45]
Clusters 6 and 8 were identified as endodermis, characterized by three marker
genes including *Os07g35480*, *Os06g13560*, and *Os05g12630*.[45] Cluster 9 was
annotated as xylem, supported by the expression of nine xylem-specific
markers such as *Os01g62490*, *Os01g54620*, and *Os01g06580*.[45] Cluster 3
was assigned as exodermis, based on the Plant Single Cell Transcriptome
Hub (PsctH: http://jinlab.hzau.edu.cn/PsctH/) markers *Os06g29994*,
Os04g41740, and *Os03g02460*.[81] Cluster 5 was annotated as epidermis,
supported by the expression of *Os10g31510*, *Os10g31660*, *Os03g06040*, and
Os06g20150.[79] Cluster 4 was identified as meristem, marked by *Os01g61920*
and *Os03g02780*.[79] Finally, cluster 2 was assigned as root hair, supported by
nine specific markers including *Os10g42750*, *Os07g44460*, and
Os04g57860.[79,81]

Identification and enrichment analyses of DEGs
To identify differentially expressed genes (DEGs) across different treatments,
we utilized the "FindMarkers" and "FindAllMarkers" functions in Seurat.
Significant DEGs for clusters and each cell type were identified using a
threshold of P_adjust < 0.05 and |$\log_2$ fold change| > = 0.25. In addition, Gene
Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG)
pathway analyses were performed on the DEGs to uncover their potential
functional enrichments and biological signal transductions. The former
includes biological processes, cellular components, and molecular functions,
while the latter includes metabolism, environmental information processing,
organismal systems, genetic information processing, and cellular processes.
Functional enrichment analyses were conducted using the clusterProfiler R

package,[82] with significantly enriched terms identified based on a false

discovery rate (FDR) threshold of < 0.05.

Analysis of rice leaf Stereo-Seq data

In the Stereo-Seq data of rice leaves, we employed Calcofluor White (CW)

staining images to visualize the cell walls and ssDNA staining images to reveal

nuclear distribution, thereby capturing the structural features of plant tissue

sections more comprehensively. Using CW-stained images aligned with the

sequencing data, we delineated the boundaries of individual cells and

assigned cell type labels, thus constructing ground-truth spatial transcriptomic

data. To accurately identify and trace the complete cell wall contours of each

epidermal cell, we combined manual cell outlining with the Cellpose[83]

automated cell segmentation software. This approach enabled precise

characterization of the boundaries of both upper and lower epidermal cells in

rice leaves **(Figure 6A)**. The spatial position of each cell on the sequencing

chip was represented by its centroid coordinates. These cell data served as

the ground truth for evaluating the ability of MF-PENS to discriminate between

different epidermal cell types.

Differential Expression Analysis and Benchmarking of Epidermal

Subpopulations

We first performed differential expression analysis on spatially resolved cells

identified via Stereo-seq. Using Scanpy, we conducted a Wilcoxon rank-sum

test to identify significantly up-DEGs specific to the upper and lower epidermal

layers, respectively (*P_adj* < 0.05, |log2FC| > 0.25). These Stereo-seq-derived

up-DEGs were subsequently employed to establish ground truth references for

the snRNA-seq datasets. We extracted all epidermal cells from the NT-PENS

dataset (annotated as epidermal cells) and the MF-PENS dataset (annotated

as bulliform, guard, and epidermal cells). Instead of unsupervised feature

selection, we manually set the identified Stereo-seq up-DEGs as the

"VariableFeatures" for these subsets. The data were then processed using the

standard Seurat pipeline: normalization and scaling via "ScaleData",

dimensionality reduction via "RunPCA" and "RunUMAP", neighbor

identification via "FindNeighbors", and clustering via "FindClusters". The

clustering resolution was adjusted to force the separation into two distinct

clusters, serving as the ground truth reference for upper and lower epidermal
identities **(Figures 6C and 6F)**. In parallel, we performed standard
unsupervised clustering on the same snRNA-seq subsets. For this analysis,
variable features were identified computationally using the
"FindVariableFeatures" function in Seurat (parameters: "selection.method" =
'dispersion', "nfeatures" = 800). The subsequent workflow (scaling, PCA,
UMAP, and clustering) remained identical, yielding clustering results derived
solely from the snRNA-seq data structure **(Figures 6D and 6G)**. By comparing
the unsupervised snRNA-seq clustering results against this
Stereo-seq-derived ground truth, we evaluated the effectiveness of MF-PENS
in resolving epidermal cell subtypes.
795

Quantitative and statistical analysis

797 All experiments were repeated at least two times, and the data are presented

798 as the mean ± standard deviation (SD). The significance of group differences

799 was evaluated with a two-tailed Student's t test or analysis of variance.

Data availability

802 All datasets generated in this study have been submitted to the China National
803 GeneBank (CNGB, accession number CNP0007006). All the other data and
804 materials are available from the corresponding authors upon request.
805

Code availability

807 We utilized the analysis of snRNA-seq data can be obtained from the
808 corresponding author upon request. We also thank the China National
809 GenBank (CNGB) for their generous support.
810

Acknowledgments
We thank X. Lin (Institute of Genetics and Developmental Biology, Chinese
Academy of Sciences) for providing the B73, Williams 82, and Zhonghuang 13
materials; F. Zhang (Institute of Genetics and Developmental Biology, Chinese
Academy of Sciences) for providing the Zhonghua 11 materials; and T. Lin
(College of Horticulture, China Agricultural University) for providing the Alisa

Craig materials. This work was supported by the Biological Breeding-National
Science and Technology Major Project [No. 2023ZD04076], the High-quality
Science and Technology Journal Construction Project of Guangdong Province
[No. 2025B1212100003] and [No. 2025B1212070003].

Author contributions

J.-M.K., H.Y. and Z.L. designed the research. J.-M.K. and L.D. wrote the
manuscript. J.-M.K., Q.L., and L.D. developed and tested the MF-PENS
technology. J.-M.K., Q.L., M.-Y.Z., C.T. and M.-J.Z. did experiments. L.D.,
J.-M.K. and H.-L.Z. performed data analysis. Q.L., H.-L.Z., F.-Y.L., H.Y. and
Z.L. revised the manuscript. L.D. and J.-M.K. prepared all the figures. All
authors discussed the results and commented on the manuscript.

Competing interests

831   The authors declare no competing interests.

Figures

835 Figure 1. Design principle and performance evaluation of MF-PENS on rice
836 roots.

(A) Workflow diagrams for NT-PENS, 0.1%FF-PENS, 1%FF-PENS, and MF-PENS,
illustrating distinct fixation and extraction strategies. The yellow shade indicates the
application of MF-PENS in various plants, demonstrating its wide adaptability and
robustness. **(B)** Cell counts of eight samples before (pink)/ after (blue) quality control,
with two biological replicates for each method. **(C)** Stepwise increases in the number
of detected genes and UMIs across down-sampled sequencing depths (sequencing
depth vs. gene detection efficiency). Points indicate mean values at the
corresponding depth, and colors represent snRNA-seq data generated by different
methods. **(D)** Violin plots showing the distributions of UMIs and detected genes
across four methods. *** indicates that MF-PENS yields significantly higher values (*P*

 < 0.05) compared with the other methods. **(E)** Mitochondrial and chloroplast

848 percentages (y-axis values × 10^{-4}%) for the eight biological samples corresponding

849 to the four methods.

Figure 2. MF-PENS-generated snRNA-seq data exhibit superior cell clustering performance.

(A) UMAP visualization of two biological replicates for each of the four methods (NT-PENS, 0.1%FF-PENS, 1%FF-PENS, and MF-PENS). **(B)** UMAP plots of Seurat

clustering results, showing nine clusters derived from snRNA-seq data generated by
the four methods, corresponding to **(A)**. Clustering performance was quantified using
a transformed Modularity score, defined as -(Modularity x 10^5). Lower scores
indicate superior clustering results. **(C)** Bubble plot of the top 10 upregulated
differentially expressed genes (up-DEGs) in each cluster based on the clustering
results across the four methods. **(D)** Bar plot of up-DEGs (*P_adj* < 0.05, $\log_2$FC >
0.25), down-DEGs (*P_adj* < 0.05, $\log_2$FC < -0.25), and total significant DEGs
(All-DEGs, *P_adj* < 0.05, |$\log_2$FC| > 0.25) identified by the four methods. **(E)** Venn
diagram showing the overlap of the top 100 up-DEGs among the four methods.
Columns in table display the count of method-specific up-DEGs, the subset of these
genes validated by the *PlantCellMarker* database, and the calculated validation rate
for each method.
868

Figure 3. MF-PENS is applicable for both fresh and cryopreserved plant samples.

(A) Workflow schematic of MF-PENS-based snRNA-seq for fresh and frozen rice root tissues. **(B)** Stepwise increases in mean detected genes and UMIs at different down-sampled sequencing depths for MF-PENS root data. **(C-D)** Violin plots showing UMIs, gene numbers, and mitochondrial percentages (y-axis × 10^{-4}) for four samples, including two replicates each of fresh and frozen methods. **(E-F)** UMAP visualizations of integrated clustering and corresponding cell type annotation for four MF-PENS samples. **(G)** Proportion of cell types in **(F)** across four samples.**(H)** Workflow

schematic of MF-PENS-based snRNA-seq for fresh and frozen rice leaf tissues. **(I)**
Stepwise increases in mean detected genes and UMIs at different down-sampled
sequencing depths for NT-PENS leaf data. **(J-M)** Violin plots showing UMIs, gene

numbers, mitochondrial percentages (y-axis × 10^{-4}%), and chloroplast percentages

(y-axis × 10^{-4}%) for four samples, including two replicates of fresh and frozen

conditions. **(M-O)** UMAP visualizations of integrated clustering and corresponding cell
type annotation for four NT-PENS samples. **(P)** Proportion of cell types in **(O)** across
four samples.
887

Figure 4. Universal applicability of MF-PENS across plant species and tissues.
(A) Application of MF-PENS-based snRNA-seq to diverse tissue types, including infected tissue (soybean root nodules), polysaccharide-rich tissue (tomato fruit), lipid-rich tissue (soybean seeds), and callus tissue (maize callus). **(B)** Cell number of eight samples after quality control. **(C)** Violin plots of gene numbers and UMIs for eight biological samples (y-axis × 10^3), including two replicates for each of the four tissue types. **(D)** Mitochondrial percentages (y-axis × 10^{-4}%) for the eight biological

896　samples. **(E)** Pearson correlation heatmap of the eight biological samples. **(F-G)**
897　Heatmaps of the top five differentially expressed genes in each cluster identified from
898　the four samples.
899

Figure 5. MF-PENS enables improved identification of cell types and detection

of heat stress–responsive genes.

(A-B) UMAP visualizations of integrated snRNA-seq data from NT-PENS and

MF-PENS under control and heat stress conditions, each with two biological

replicates. **(C)** Venn diagram showing the overlap of up-DEGs between control and

heat stress conditions in NT-PENS and MF-PENS datasets. **(D-E)** Top 20 enriched

biological processes based on the up-DEGs identified in **(C)**. **(F)** UMAP visualizations

of cell type annotations and percentage based on clustering results from **(A)**. **(G)** Bar

plot showing the number of up-/down-DEGs in each annotated cell type from **(F)**. **(H)**

UMAP visualizations of cell type annotations and percentage based on clustering

results from **(B)**. **(I)** Bar plot showing the number of up-/down-DEGs in each

annotated cell type from **(H)**. **(J)** Venn diagram of heat-responsive genes (control vs.

heat) shared by four commonly annotated cell types in both NT-PENS and MF-PENS

datasets. **(K)** Bubble plot of the top 10 enriched biological processes of

heat-responsive genes uniquely identified in four MF-PENS–specific cell types.

Figure 6. MF-PENS enables accurate identification of upper and lower epidermal cell subtypes in rice leaves.

(A) Ground truth spatial locations of upper and lower epidermal cells in rice leaf tissue based on Stereo-seq. **(B)** Accuracy of epidermal cell subtype identification using MF-PENS and NT-PENS datasets. *** indicates a significant difference ($P < 0.05$). **(C)** Ground truth annotation of upper and lower epidermal cells in the NT-PENS dataset using a previously reported strategy. **(D)** Cell type annotation after standard clustering analysis by Seruat of the NT-PENS dataset. **(E)** Proportions of upper and lower epidermal cells across two biological replicates based on **(D)**. **(F-H)** Application of the same strategy to MF-PENS data: ground truth upper and lower epidermal cells **(F)**,

928   clustering-based cell type annotation **(G)**, and cell type proportions across two
929   replicates **(H)**. **(I-J)** Bubble plots showing the top 10 enriched biological processes for
930   DEGs in the two epidermal subtypes identified in MF-PENS annotations **(G)**.
931

Reference

1. Birnbaum, K., Shasha, D.E., Wang, J.Y., Jung, J.W., Lambert, G.M., Galbraith, D.W., and Benfey, P.N. (2003). A gene expression map of the Arabidopsis root. Science *302*, 1956-1960. 10.1126/science.1090022.

2. Zhao, J., Zhang, S., Liu, Y., He, X., Qu, M., Xu, G., Wang, H., Huang, M., Pan, J., Liu, Z., et al. (2020). Single-cell RNA sequencing reveals the heterogeneity of liver-resident immune cells in human. Cell Discovery *6*, 22. 10.1038/s41421-020-0157-z.

3. Tsagiopoulou, M., Rashmi, S., Aguilar-Fernandez, S., Nieto, J., and Gut, I.G. (2024). Multi-organ single-cell transcriptomics of immune cells uncovered organ-specific gene expression and functions. Scientific Data *11*, 316. 10.1038/s41597-024-03152-z.

4. Jovic, D., Liang, X., Zeng, H., Lin, L., Xu, F., and Luo, Y. (2022). Single-cell RNA sequencing technologies and applications: A brief overview. Clinical and Translational Medicine *12*, e694. 10.1002/ctm2.694.

5. Bian, S., Hou, Y., Zhou, X., Li, X., Yong, J., Wang, Y., Wang, W., Yan, J., Hu, B., Guo, H., et al. (2018). Single-cell multiomics sequencing and analyses of human colorectal cancer. Science *362*, 1060-1063. 10.1126/science.aao3791.

6. Terekhova, M., Swain, A., Bohacova, P., Aladyeva, E., Arthur, L., Laha, A., Mogilenko, D.A., Burdess, S., Sukhov, V., Kleverov, D., et al. (2023). Single-cell atlas of healthy human blood unveils age-related loss of NKG2C(+)GZMB(-)CD8(+) memory T cells and accumulation of type 2 memory T cells. Immunity *56*, 2836-2854.e2839. 10.1016/j.immuni.2023.10.013.

7. Mazid, M.A., Ward, C., Luo, Z., Liu, C., Li, Y., Lai, Y., Wu, L., Li, J., Jia, W., Jiang, Y., et al. (2022). Rolling back human pluripotent stem cells to an eight-cell embryo-like stage. Nature *605*, 315-324. 10.1038/s41586-022-04625-0.

8. Ochocka, N., Segit, P., Walentynowicz, K.A., Wojnicki, K., Cyranowski, S., Swatler, J., Mieczkowski, J., and Kaminska, B. (2021). Single-cell RNA sequencing reveals functional heterogeneity of glioma-associated brain macrophages. Nature Communications *12*, 1151. 10.1038/s41467-021-21407-w.

9. Liu, Q., Ma, W., Chen, R., Li, S.T., Wang, Q., Wei, C., Hong, Y., Sun, H.X., Cheng, Q., Zhao, J., and Kang, J. (2024). Multiome in the Same Cell Reveals the Impact of Osmotic Stress on Arabidopsis Root Tip Development at Single-Cell Level. Advanced Science *11*, e2308384. 10.1002/advs.202308384.

10. Liu, Q., Liang, Z., Feng, D., Jiang, S., Wang, Y., Du, Z., Li, R., Hu, G., Zhang, P., Ma, Y., et al. (2021). Transcriptional landscape of rice roots at the single-cell resolution. Molecular Plant *14*, 384-394. 10.1016/j.molp.2020.12.014.

11. Zhang, T.Q., Chen, Y., Liu, Y., Lin, W.H., and Wang, J.W. (2021). Single-cell
transcriptome atlas and chromatin accessibility landscape reveal
differentiation trajectories in the rice root. Nature Communications *12*, 2053.
10.1038/s41467-021-22352-4.

12. Swift, J., Luginbuehl, L.H., Hua, L., Schreier, T.B., Donald, R.M., Stanley, S.,
Wang, N., Lee, T.A., Nery, J.R., Ecker, J.R., and Hibberd, J.M. (2024).
Exaptation of ancestral cell-identity networks enables C(4) photosynthesis.
Nature *636*, 143-150. 10.1038/s41586-024-08204-3.

13. Singh, S., Praveen, A., Dudha, N., Sharma, V.K., and Bhadrecha, P. (2024).
Single-cell transcriptomics: a new frontier in plant biotechnology research.
Plant Cell Rep *43*, 294. 10.1007/s00299-024-03383-9.

14. Chen, K., Chen, J., Pi, X., Huang, L.J., and Li, N. (2023). Isolation, Purification,
and Application of Protoplasts and Transient Expression Systems in Plants.
International Journal of Molecular Sciences *24*. 10.3390/ijms242316892.

15. Xu, X.F., Zhu, H.Y., Ren, Y.F., Feng, C., Ye, Z.H., Cai, H.M., Wan, X.C., and
Peng, C.Y. (2021). Efficient isolation and purification of tissue-specific
protoplasts from tea plants (Camellia sinensis (L.) O. Kuntze). Plant Methods
17, 84. 10.1186/s13007-021-00783-w.

16. Grones, C., Eekhout, T., Shi, D., Neumann, M., Berg, L.S., Ke, Y., Shahan, R.,
Cox, K.L., Jr., Gomez-Cano, F., Nelissen, H., et al. (2024). Best practices for
the execution, analysis, and data storage of plant single-cell/nucleus
transcriptomics. Plant Cell *36*, 812-828. 10.1093/plcell/koae003.

17. Liao, R.Y., and Wang, J.W. (2023). Analysis of meristems and plant
regeneration at single-cell resolution. Current Opinion in Plant Biology *74*,
102378. 10.1016/j.pbi.2023.102378.

18. Rodriguez-Villalon, A., and Brady, S.M. (2019). Single cell RNA sequencing
and its promise in reconstructing plant vascular cell lineages. Current Opinion
in Plant Biology *48*, 47-56. 10.1016/j.pbi.2019.04.002.

19. Shaw, R., Tian, X., and Xu, J. (2021). Single-Cell Transcriptome Analysis in
Plants: Advances and Challenges. Molecular Plant *14*, 115-126.
10.1016/j.molp.2020.10.012.

20. Yang, P., Sun, Y., Sun, X., Li, Y., and Wang, L. (2024). Optimization of
preparation and transformation of protoplasts from Populus simonii□x□P.
nigra leaves and subcellular localization of the major latex protein 328
(MLP328). Plant Methods *20*, 3. 10.1186/s13007-023-01128-5.

21. Zhang, T.Q., Chen, Y., and Wang, J.W. (2021). A single-cell analysis of the
Arabidopsis vegetative shoot apex. Developmental Cell *56*, 1056-1074.e1058.
10.1016/j.devcel.2021.02.021.

22. Li, C., Zhang, S., Yan, X., Cheng, P., and Yu, H. (2023). Single-nucleus
sequencing deciphers developmental trajectories in rice pistils. Developmental
Cell *58*, 694-708.e694. 10.1016/j.devcel.2023.03.004.

23. Farmer, A., Thibivilliers, S., Ryu, K.H., Schiefelbein, J., and Libault, M. (2021).

Single-nucleus RNA and ATAC sequencing reveals the impact of chromatin
accessibility on gene expression in Arabidopsis roots at the single-cell level.
Molecular Plant *14*, 372-383. 10.1016/j.molp.2021.01.001.

24.   Zhang, L., He, C., Lai, Y., Wang, Y., Kang, L., Liu, A., Lan, C., Su, H., Gao, Y.,
Li, Z., et al. (2023). Asymmetric gene expression and cell-type-specific
regulatory networks in the root of bread wheat revealed by single-cell
multiomics analysis. Genome Biology *24*, 65. 10.1186/s13059-023-02908-x.

25.   Marand, A.P., Chen, Z., Gallavotti, A., and Schmitz, R.J. (2021). A
cis-regulatory atlas in maize at single-cell resolution. Cell *184*,
3041-3055.e3021. 10.1016/j.cell.2021.04.014.

26.   Thibivilliers, S., Anderson, D., and Libault, M. (2020). Isolation of Plant Root
Nuclei for Single Cell RNA Sequencing. Current Opinion in Plant Biology *5*,
e20120. 10.1002/cppb.20120.

27.   Wang, K., Zhao, C., Xiang, S., Duan, K., Chen, X., Guo, X., and Sahu, S.K.
(2023). An optimized FACS-free single-nucleus RNA sequencing (snRNA-seq)
method for plant science research. Plant Science *326*, 111535.
10.1016/j.plantsci.2022.111535.

28.   Chung, J.Y., Braunschweig, T., Williams, R., Guerrero, N., Hoffmann, K.M.,
Kwon, M., Song, Y.K., Libutti, S.K., and Hewitt, S.M. (2008). Factors in tissue
handling and processing that impact RNA obtained from formalin-fixed,
paraffin-embedded tissue. Journal of Histochemistry & Cytochemistry *56*,
1033-1042. 10.1369/jhc.2008.951863.

29.   Srinivasan, M., Sedmak, D., and Jewell, S. (2002). Effect of fixatives and
tissue processing on the content and integrity of nucleic acids. The American
Journal of Pathology *161*, 1961-1971. 10.1016/s0002-9440(10)64472-0.

30.   Ichikawa, T., Wang, D., Miyazawa, K., Miyata, K., Oshima, M., and Fukuma, T.
(2022). Chemical fixation creates nanoscale clusters on the cell surface by
aggregating membrane proteins. Communications Biology *5*, 487.
10.1038/s42003-022-03437-2.

31.   SĘdek, Ł., Kulis, J., SŁota, Ł., Twardoch, M., Pierzyna-ŚwitaŁa, M., Perkowski,
B., and SzczepaŃski, T. (2020). The influence of fixation of biological samples
on cell count and marker expression stability in flow cytometric analyses.
Central European Journal of Immunology *45*, 206-213.
10.5114/ceji.2020.95858.

32.   Sunaga-Franze, D.Y., Muino, J.M., Braeuning, C., Xu, X., Zong, M.,
Smaczniak, C., Yan, W., Fischer, C., Vidal, R., Kliem, M., et al. (2021).
Single-nucleus RNA sequencing of plant tissues using a nanowell-based
system. Plant J *108*, 859-869. 10.1111/tpj.15458.

33.   Clarke, Z.A., Andrews, T.S., Atif, J., Pouyabahar, D., Innes, B.T., MacParland,
S.A., and Bader, G.D. (2021). Tutorial: guidelines for annotating single-cell
transcriptomic maps using automated and manual methods. Nature Protocols
16, 2749-2764. 10.1038/s41596-021-00534-0.

34. Luecken, M.D., and Theis, F.J. (2019). Current best practices in single-cell
RNA-seq analysis: a tutorial. Molecular Systems Biology *15*, e8746.
10.15252/msb.20188746.
35. Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Challenges in
unsupervised clustering of single-cell RNA-seq data. Nature Reviews
Genetics *20*, 273-282. 10.1038/s41576-018-0088-9.
36. Mereu, E., Lafzi, A., Moutinho, C., Ziegenhain, C., McCarthy, D.J.,
Álvarez-Varela, A., Batlle, E., Sagar, Grün, D., Lau, J.K., et al. (2020).
Benchmarking single-cell RNA-sequencing protocols for cell atlas projects.
Nature Biotechnology *38*, 747-755. 10.1038/s41587-020-0469-4.
37. Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Challenges in
unsupervised clustering of single-cell RNA-seq data. Nat Rev Genet *20*,
273-282. 10.1038/s41576-018-0088-9.
38. Alcalá-Corona, S.A., Sandoval-Motta, S., Espinal-Enríquez, J., and
Hernández-Lemus, E. (2021). Modularity in Biological Networks. Frontiers in
Genetics *12*, 701331. 10.3389/fgene.2021.701331.
39. Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden:
guaranteeing well-connected communities. Scientific Report *9*, 5233.
10.1038/s41598-019-41695-z.
40. Li, P., Liu, Q., Wei, Y., Xing, C., Xu, Z., Ding, F., Liu, Y., Lu, Q., Hu, N., Wang, T.,
et al. (2023). Transcriptional Landscape of Cotton Roots in Response to Salt
Stress at Single-cell Resolution. Plant Commun *5*, 100740.
10.1016/j.xplc.2023.100740.
41. Dolan, L., Janmaat, K., Willemsen, V., Linstead, P., Poethig, S., Roberts, K.,
and Scheres, B. (1993). Cellular organisation of the Arabidopsis thaliana root.
Development *119*, 71-84. 10.1242/dev.119.1.71.
42. Lundberg, D.S., Lebeis, S.L., Paredes, S.H., Yourstone, S., Gehring, J.,
Malfatti, S., Tremblay, J., Engelbrektson, A., Kunin, V., Del Rio, T.G., et al.
(2012). Defining the core Arabidopsis thaliana root microbiome. Nature *488*,
86-90. 10.1038/nature11237.
43. Xiao, G., and Zhang, Y. (2020). Adaptive Growth: Shaping Auxin-Mediated
Root System Architecture. Trends in Plant Science *25*, 121-123.
10.1016/j.tplants.2019.12.001.
44. Weis, F., Giudice, E., Churcher, M., Jin, L., Hilcenko, C., Wong, C.C., Traynor,
D., Kay, R.R., and Warren, A.J. (2015). Mechanism of eIF6 release from the
nascent 60S ribosomal subunit. Nature Structural & Molecular Biology *22*,
914-919. 10.1038/nsmb.3112.
45. Jin, J., Lu, P., Xu, Y., Tao, J., Li, Z., Wang, S., Yu, S., Wang, C., Xie, X., Gao, J.,
et al. (2022). PCMDB: a curated and comprehensive resource of plant cell
markers. Nucleic Acids Research *50*, D1448-d1455. 10.1093/nar/gkab949.
46. Dixon, R.A. (2001). Natural products and plant disease resistance. Nature *411*,
843-847. 10.1038/35081178.

47.   Dixon, R.A., Gang, D.R., Charlton, A.J., Fiehn, O., Kuiper, H.A., Reynolds, T.L.,
1101   Tjeerdema, R.S., Jeffery, E.H., German, J.B., Ridley, W.P., and Seiber, J.N.
1102   (2006). Applications of Metabolomics in Agriculture. Journal of Agricultural and
1103   Food Chemistry *54*, 8984-8994. 10.1021/jf061218t.

1104   48.   Duan, Z., Li, Q., Wang, H., He, X., and Zhang, M. (2023). Genetic regulatory
1105   networks of soybean seed size, oil and protein contents. Frontiers in Plant
1106   Science *14*, 1160418. 10.3389/fpls.2023.1160418.

1107   49.   Chylińska, M., Szymańska-Chargot, M., Deryło, K., Tchórzewska, D., and
1108   Zdunek, A. (2017). Changing of biochemical parameters and cell wall
1109   polysaccharides distribution during physiological development of tomato fruit.
1110   Plant Physiology and Biochemistry *119*, 328-337.
1111   https://doi.org/10.1016/j.plaphy.2017.09.010.

1112   50.   Quinet, M., Angosto, T., Yuste-Lisbona, F.J., Blanchard-Gros, R., Bigot, S.,
1113   Martinez, J.P., and Lutts, S. (2019). Tomato Fruit Development and
1114   Metabolism. Frontiers in Plant Science *10*, 1554. 10.3389/fpls.2019.01554.

1115   51.   Yang, M.C., Wu, Z.C., Huang, L.L., Abbas, F., and Wang, H.C. (2022).
1116   Systematic Methods for Isolating High Purity Nuclei from Ten Important Plants
1117   for Omics Interrogation. Cells *11*. 10.3390/cells11233919.

1118   52.   Cervantes-Pérez, S.A., Zogli, P., Amini, S., Thibivilliers, S., Tennant, S.,
1119   Hossain, M.S., Xu, H., Meyer, I., Nooka, A., Ma, P., et al. (2024). Single-cell
1120   transcriptome atlases of soybean root and mature nodule reveal new
1121   regulatory programs that control the nodulation process. Plant
1122   Communications *5*. 10.1016/j.xplc.2024.100984.

1123   53.   Sun, B., Wang, Y., Yang, Q., Gao, H., Niu, H., Li, Y., Ma, Q., Huan, Q., Qian,
1124   W., and Ren, B. (2023). A high-resolution transcriptomic atlas depicting
1125   nitrogen fixation and nodule development in soybean. Journal of Integrative
1126   Plant Biology *65*, 1536-1552. https://doi.org/10.1111/jipb.13495.

1127   54.   Rahman, M.M., Divi, U.K., Liu Qing, L.Q., Zhou XueRong, Z.X., Surinder
1128   Singh, S.S., and Aruna Kilaru, A.K. (2016). Oil-rich nonseed tissues for
1129   enhancing plant oil production. CABI Reviews, 1–11.
1130   10.1079/PAVSNNR201611021.

1131   55.   Philippe, G., Geneix, N., Petit, J., Guillon, F., Sandt, C., Rothan, C., Lahaye,
1132   M., Marion, D., and Bakan, B. (2020). Assembly of tomato fruit cuticles: a
1133   cross-talk between the cutin polyester and cell wall polysaccharides. New
1134   Phytologist *226*, 809-822. https://doi.org/10.1111/nph.16402.

1135   56.   Shillito, R.D., Carswell, G.K., Johnson, C.M., DiMaio, J.J., and Harms, C.T.
1136   (1989). Regeneration of Fertile Plants from Protoplasts of Elite Inbread Maize.
1137   Nature Biotechnology *7*, 581-587. 10.1038/nbt0689-581.

1138   57.   Welter, M.E., Clayton, D.S., Miller, M.A., and J, E.P. (1995). Morphotypes of
1139   friable embryogenic maize callus. Plant Cell Reports *14*, 725-729.
1140   10.1007/bf00232656.

58.   Souter, M., and Lindsey, K. (2000). Polarity and signalling in plant

embryogenesis. Journal of Experimental Botany *51*, 971-983.
10.1093/jexbot/51.347.971.

59. Liu, Z., Kong, X., Long, Y., Liu, S., Zhang, H., Jia, J., Cui, W., Zhang, Z., Song,
X., Qiu, L., et al. (2023). Integrated single-nucleus and spatial transcriptomics
captures transitional states in soybean nodule maturation. Nature Plants *9*,
515-524. 10.1038/s41477-023-01387-z.

60. Fan, J., Shen, Y., Chen, C., Chen, X., Yang, X., Liu, H., Chen, R., Liu, S.,
Zhang, B., Zhang, M., et al. (2025). A large-scale integrated transcriptomic
atlas for soybean organ development. Molecular Plant.
https://doi.org/10.1016/j.molp.2025.02.003.

61. Sun, X., Feng, D., Liu, M., Qin, R., Li, Y., Lu, Y., Zhang, X., Wang, Y., Shen, S.,
Ma, W., and Zhao, J. (2022). Single-cell transcriptome reveals dominant
subgenome expression and transcriptional response to heat stress in Chinese
cabbage. Genome Biology *23*, 262. 10.1186/s13059-022-02834-4.

62. Wang, T., Wang, F., Deng, S., Wang, K., Feng, D., Xu, F., Guo, W., Yu, J., Wu,
Y., Wuriyanghan, H., et al. (2025). Single-cell transcriptomes reveal
spatiotemporal heat stress response in maize roots. Nature Communications
16, 177. 10.1038/s41467-024-55485-3.

63. Li, Y., Ma, H., Wu, Y., Ma, Y., Yang, J., Li, Y., Yue, D., Zhang, R., Kong, J.,
Lindsey, K., et al. (2024). Single-Cell Transcriptome Atlas and Regulatory
Dynamics in Developing Cotton Anthers. Advanced Science *11*, e2304017.
10.1002/advs.202304017.

64. Alles, J., Karaiskos, N., Praktiknjo, S.D., Grosswendt, S., Wahle, P., Ruffault,
P.-L., Ayoub, S., Schreyer, L., Boltengagen, A., Birchmeier, C., et al. (2017).
Cell fixation and preservation for droplet-based single-cell transcriptomics.
BMC Biology *15*, 44. 10.1186/s12915-017-0383-5.

65. Du, L., Pan, W., Song, Y., Liu, F., Jia, Y., Wei, C., Li, X., Hou, Y., Nie, X., Sun,
H.-X., and Kang, J. Cell-Type-Specific and Variety-Specific Responses to Salt
Stress in Wheat Root Revealed by Single-Cell Transcriptomics. Plant
Biotechnology Journal *n/a*. https://doi.org/10.1111/pbi.70422.

66. Xia, K., Sun, H.X., Li, J., Li, J., Zhao, Y., Chen, L., Qin, C., Chen, R., Chen, Z.,
Liu, G., et al. (2022). The single-cell stereo-seq reveals region-specific cell
subtypes and transcriptome profiling in Arabidopsis leaves. Developmental
Cell *57*, 1299-1310.e1294. 10.1016/j.devcel.2022.04.011.

67. Narawatthana, S., Phansenee, Y., Thammasamisorn, B.O., and Vejchasarn, P.
(2023). Multi-model genome-wide association studies of leaf anatomical traits
and vein architecture in rice. Frontiers in Plant Science *14*, 1107718.
10.3389/fpls.2023.1107718.

68. Zuch, D.T., Doyle, S.M., Majda, M., Smith, R.S., Robert, S., and Torii, K.U.
(2022). Cell biology of the leaf epidermis: Fate specification, morphogenesis,
and coordination. Plant Cell *34*, 209-227. 10.1093/plcell/koab250.

69. Zhang, G.-H., Xu, Q., Zhu, X.-D., Qian, Q., and Xue, H.-W. (2009).

SHALLOT-LIKE1 Is a KANADI Transcription Factor That Modulates Rice Leaf
Rolling by Regulating Leaf Abaxial Cell Development The Plant Cell *21*,
719-735. 10.1105/tpc.108.061457.

70.   Wang, Y., Huan, Q., Li, K., and Qian, W. (2021). Single-cell transcriptome atlas
of the leaf and root of rice seedlings. Journal of Genetics and Genomics *48*,
881-898. https://doi.org/10.1016/j.jgg.2021.06.001.

71.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018).
Integrating single-cell transcriptomic data across different conditions,
technologies, and species. Nature Biotechnology *36*, 411-420.
10.1038/nbt.4096.

72.   Chen, A., Liao, S., Cheng, M., Ma, K., Wu, L., Lai, Y., Qiu, X., Yang, J., Xu, J.,
Hao, S., et al. (2022). Spatiotemporal transcriptomic atlas of mouse
organogenesis using DNA nanoball-patterned arrays. Cell *185*,
1777-1792.e1721. 10.1016/j.cell.2022.04.003.

73.   Huang, Y., Wu, Y., Han, S., Wang, Q., Cong, G., Liu, Z., Guan, S., Huang, X.,
Liu, Y., Yin, J., et al. (2024). Protocol for optimized nasal mucosa sample
processing to obtain high-quality scRNA-seq and scATAC-seq data. STAR
Protocols *5*, 103298. 10.1016/j.xpro.2024.103298.

74.   Kawahara, Y., de la Bastide, M., Hamilton, J.P., Kanamori, H., McCombie,
W.R., Ouyang, S., Schwartz, D.C., Tanaka, T., Wu, J., Zhou, S., et al. (2013).
Improvement of the Oryza sativa Nipponbare reference genome using next
generation sequence and optical map data. Rice (New York, N.Y.) *6*, 4.
10.1186/1939-8433-6-4.

75.   Schmutz, J., Cannon, S.B., Schlueter, J., Ma, J., Mitros, T., Nelson, W., Hyten,
D.L., Song, Q., Thelen, J.J., Cheng, J., et al. (2010). Genome sequence of the
palaeopolyploid soybean. Nature *463*, 178-183. 10.1038/nature08670.

76.   Sato, S., Tabata, S., Hirakawa, H., Asamizu, E., Shirasawa, K., Isobe, S.,
Kaneko, T., Nakamura, Y., Shibata, D., Aoki, K., et al. (2012). The tomato
genome sequence provides insights into fleshy fruit evolution. Nature *485*,
635-641. 10.1038/nature11119.

77.   Hayford, R.K., Haley, O.C., Cannon, E.K., Portwood, J.L., 2nd, Gardiner, J.M.,
Andorf, C.M., and Woodhouse, M.R. (2024). Functional annotation and
meta-analysis of maize transcriptomes reveal genes involved in biotic and
abiotic stress. BMC Genomics *25*, 533. 10.1186/s12864-024-10443-7.

78.   McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder:
Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial
Nearest Neighbors. Cell Systems *8*, 329-337.e324.
10.1016/j.cels.2019.03.003.

79.   Cao, S., He, Z., Chen, R., Luo, Y., Fu, L.Y., Zhou, X., He, C., Yan, W., Zhang,
C.Y., and Chen, D. (2023). scPlant: A versatile framework for single-cell
transcriptomic data analysis in plants. Plant Communications *4*, 100631.
10.1016/j.xplc.2023.100631.

80. Chen, Y.E., Ma, J., Wu, N., Su, Y.Q., Zhang, Z.W., Yuan, M., Zhang, H.Y.,
Zeng, X.Y., and Yuan, S. (2018). The roles of Arabidopsis proteins of Lhcb4,
Lhcb5 and Lhcb6 in oxidative stress under natural light conditions. Plant
Physiology and Biochemistry *130*, 267-276. 10.1016/j.plaphy.2018.07.014.
81. Xu, Z., Wang, Q., Zhu, X., Wang, G., Qin, Y., Ding, F., Tu, L., Daniell, H.,
Zhang, X., and Jin, S. (2022). Plant Single Cell Transcriptome Hub (PsctH): an
integrated online tool to explore the plant single-cell transcriptome landscape.
Plant Biotechnology Journal *20*, 10-12. 10.1111/pbi.13725.
82. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R
package for comparing biological themes among gene clusters. Omics *16*,
284-287. 10.1089/omi.2011.0118.
83. Pachitariu, M., and Stringer, C. (2022). Cellpose 2.0: how to train your own
model. Nature Methods *19*, 1634-1641. 10.1038/s41592-022-01663-4.
1239

 Supplementary figures

 Figure S1. MF-PENS is the optimal method for single-nucleus sequencing in
 root tissue samples.
 (A) Violin plots showing UMIs and gene numbers across four methods. **(B)** Pearson
 correlation heatmap of eight samples where color gradients from blue to red indicate
 correlation coefficients from 0.5 to 1.0. Hierarchical clustering was applied to the

1247   correlation matrix. **(C-F)** Bubble plots showing the top 10 enriched biological
1248   processes of up-DEGs in each of the nine cell clusters identified from rice root using
1249   four different methods. The significance scale (-log$_{10}$(P value) was capped at a
1250   maximum value of 30 (values >30 are displayed as 30). **Related to Figure 1 and**
1251   **Figure 2.**

1252

Figure S2. Comparative KEGG pathway enrichment analysis of rice root cell
clusters across four nuclear isolation methods.
Bubble plots display the top 10 significantly enriched KEGG pathways for the
up-DEGs in each of the nine cell clusters identified by NT-PENS, 0.1%FF-PENS,
1%FF-PENS, and MF-PENS, respectively. The size of the dots typically represents
the gene count or ratio, while the color gradient indicates the statistical significance
level. For visualization clarity, the significance scale (-log$_{10}$(P value) was capped at a
maximum value of 30 (values >30 are displayed as 30).

Related to Figure 1 and Figure 2.

1262

1263   **Figure S3. Cell clustering analysis of MF-PENS snRNA-seq data from diverse**
1264   **plant tissues.**

1265   **(A-B)** UMAP visualization of integrated fresh and frozen rice root samples using CCA
1266   and the resulting cell clustering. **(C-D)** UMAP visualization of integrated fresh and
1267   frozen rice leaf samples using CCA and the resulting cell clustering. **(E-H)** UMAP

 plots showing cell clustering of snRNA-seq data from four challenging tissue samples.
Related to Figure 3 and Figure 4.

Figure S4. Expression profiles of marker genes used for cell type annotation in rice root snRNA-seq data. (A) Dot plot illustrating the expression patterns of representative marker genes across eight identified cell types. **(B)** Feature plots displaying the distribution and expression levels of 29 selected marker genes. The color gradient indicates the relative expression level. **Related to Figure 3.**

1277

1278 **Figure S5. Expression profiles of marker genes used for cell type annotation in**
1279 **rice leaf snRNA-seq data. (A)** Dot plot illustrating the expression patterns of
1280 representative marker genes across six identified cell types. **(B)** Feature plots
1281 displaying the distribution and expression levels of 14 selected marker genes. The
1282 color gradient indicates the relative expression level. **Related to Figure 3.**

1283

1284

Figure S6. MF-PENS outperforms NT-PENS in capturing transcriptomic responses to heat stress in rice leaves.

(A–B) UMAP plots of cell clustering of epidermal cells in rice leaves profiled by NT-PENS **(A)** and MF-PENS **(B)**. **(C-D)** Volcano plots of differentially expressed

1289   genes identified within each cluster from NT-PENS **(C)** and MF-PENS **(D)**,

1290   corresponding to **(A)** and **(B)**, respectively. **(E-F)** Top 20 enriched Gene Ontology (GO)

1291   terms derived from heat-responsive genes identified in **Figure 5A-B**, including

1292   categories of Biological Process (BP), Cellular Component (CC), and Molecular

1293   Function (MF). **Related to Figure 5.**

1294

1295

1296 **Figure S7. Bubble plots of the top 20 enriched biological processes for heat**
1297 **stress-responsive genes in rice leaf snRNA-seq data.** The analysis was performed
1298 on five cell types shared between NT-PENS and MF-PENS datasets, comparing
1299 control versus heat stress conditions. **Related to Figure 5.**

1300

Figure S8. Identification of upper and lower epidermal cells in rice leaves using
Stereo-seq data as ground truth.
(A) UMAP plot of cell clustering from MF-PENS data, using marker genes of upper
and lower epidermal cells derived from Stereo-seq as feature genes. **(B)** Proportions
of the identified clusters across two biological replicates in MF-PENS data. **(C-D)** GO
enrichment analysis of cluster-specific genes for the two clusters identified in **(A). (E)**
UMAP plot of cell clustering from NT-PENS data using the same Stereo-seq-derived
marker genes as features. **(F)** Proportions of clusters across two biological replicates
in NT-PENS data. **(G)** GO enrichment analysis of cluster-specific genes from cluster 0
in **(E)**; no cluster-specific genes were identified for cluster 1. **Related to Figure 6.**

1311