Title: Variable performance of widely used bisulfite sequencing methods and read mapping

software for DNA methylation

Authors: Emily V. Kerns, Jesse N. Weber

Emily V. Kerns, University of Wisconsin-Madison, Department of Integrative Biology,

Kerns3@wisc.edu, https://orcid.org/0009-0002-0538-4632

Corresponding Author: Jesse N. Weber, University of Wisconsin-Madison, Department of

Integrative Biology, Jnweber2@wisc.edu, https://orcid.org/0000-0003-4839-6684

Funding Statement: This work was funded by a startup award to JNW from Department of

Integrative Biology at the University of Wisconsin-Madison, UW-Madison Integrative Biology

Graduate Summer Research Award (EVK), and National Institute of General Medical Sciences

5R35GM142891 (JNW).

Conflict of Interest Disclosure: We have no conflicts of interest.

Ethics Approval Statement: All animals were handled and euthanized in accordance with UW-

Madison IACUC protocol L006460-A01.

Abstract:

DNA methylation (DNAm) is the most commonly studied marker in ecological epigenetics, yet

the performance of popular library preparation strategies and bioinformatic tools is seldom

assessed and compared in genetically variable natural populations. We profiled DNAm using

reduced representation bisulfite sequencing (RRBS) and whole genome bisulfite sequencing

(WGBS), including technical and biological replicates from lab-reared and wild-caught

threespine stickleback (*Gasterosteus aculeatus*). We then compared how the most commonly

used read mapper and methylation caller (Bismark) performed relative to two alternative

pipelines (BWA mem or BWA meth read mappers analyzed with MethyDackel). BWA meth

provided 50% higher mapping efficiency than BWA mem and 45% higher efficiency than

Bismark. Despite differences in mapping efficiency, BWA meth and Bismark produced highly

similar methylation profiles, while BWA mem systematically discarded unmethylated cytosines.

Sequencing depth filters had large impacts on CpG sites recovered across multiple individuals,

with the largest impact on WGBS data. Notably, the prevalence of CpG sites with intermediate

methylation levels is greatly reduced in RRBS data compared to WGBS, which may have

important consequences for functional interpretations. We conclude by discussing how library

construction and bisulfite alignment wrappers can influence SNP filtering, genomic coverage,

and the abundance and reliability of data available for downstream analysis. Our analyses

suggest that researchers studying genetically variable populations may prioritize filtering SNPs

by constructing RRBS libraries with small insert sizes and paired end reads, which is counter to

conventional wisdom.

Key words: DNA methylation, RRBS, WGBS, Bismark, BWA meth, Ecological epigenetics

<u>Introduction:</u>

Epigenetics is an umbrella term for chemical and conformational modifications to DNA

and proteins, with DNA methylation (DNAm) being among the most well studied epigenetic

marks in ecologically focused studies (Banta & Richards, 2018; Hu & Barrett, 2017; Lea et al.,

2017; Rey et al., 2020). As molecular ecologists increasingly explore the role of methylation in

natural systems, there is a need for increased attention to how methodological choices impact

inferences of between and within individual variation. This is particularly important given that

most methylation profiling methods were originally developed and applied in inbred laboratory

model organisms.

There are multiple forms of DNAm, but the most abundant type in vertebrates is the

addition of a methyl group to cytosine at CpG dinucleotides, often found in CpG-dense regions

of the genome (CpG islands) (Jaenisch & Bird, 2003). DNAm has a broad range of genomic

effects, including silencing transposable elements (Marin et al., 2020), blocking transcription

factors and other proteins from binding to DNA, increasing the likelihood of spontaneous

deamination of methylated cytosines converting them to thymine (Ord et al., 2023), and slowing

down transcriptional machinery (Cholewa-Waclaw et al., 2019). While its role varies, increased

methylation is often associated with decreased gene expression (Zemach et al., 2010).

Multigenerational common garden experiments show that although the genomic loci that

undergo DNAm are largely heritable, they can also be modified by environmental stimuli (Bogan

et al., 2023; Heckwolf et al., 2020; Hu et al., 2021). Moreover, environmentally-induced shifts in

DNAm can be inherited across generations (Takahashi et al., 2023; Tobi et al., 2018). Recent

findings also suggest that methylated regions experience different selection pressures than

background genomic DNA (Ord et al., 2023). Despite accumulating evidence that DNAm may

play an important role in evolution, this topic is still an active area for debate (Adrian-

Kalchhauser et al., 2020; Charlesworth et al., 2017; Jaenisch & Bird, 2003; Otterdijk & Michels,

2016). As interest in the functional and evolutionary roles of DNAm continues to increase, more

attention needs to be directed toward developing a robust understanding of which loci are

methylated and at what rates.

DNAm is commonly measured using bisulfite sequencing methods. Treating DNA with

bisulfite converts unmethylated cytosines to uracil, which then becomes thymine during PCR

amplification. When compared to a reference genome, C/T mismatches are identified as

unmethylated cytosines, while any cytosines remaining are inferred to be methylated. Although

DNAm at any particular base pair is a binary trait (i.e., present/absent at a CpG site on each

read), the same location may not be methylated across cells or tissues. Some CpG sites will be

consistently methylated (or unmethylated) while others will have intermediate methylation (Hay

et al., 2023). When bisulfite sequencing is used for bulk sequencing of many cells within a

particular tissue, the three possible methylation states of a CpG site in a single cell

(present/absent/heterozygous) are summed and become a quantitative metric (i.e., proportion of

total sites that are methylated). The ability to accurately capture this quantitative variation is

highly contingent on the number of times that a given base pair is represented in the sequencing

data, often referred to as read depth.

The two most commonly used bisulfite sequencing methods are whole genome bisulfite

sequencing (WGBS) and reduced representation bisulfite sequencing (RRBS; Gu et al., 2011).

While WGBS offers the potential to profile all CpG sites, it also requires large amounts of

sequencing data in order to repeatedly sample individual DNAm sites. This requirement often

limits WGBS to experiments involving few biological replicates and relatively low read depth,

which reduces accuracy of methylation calls and statistical power to detect group-level

differences (Seiler Vellame et al., 2021). Additionally, because WGBS does not enrich for

methylated sites, much of the sequence data is uninformative for epigenetic studies. For example,

70-80% of mapped reads in human WGBS studies do not contain CpG dinucleotides (Elliott et

al., 2015). In contrast, RRBS uses methylation-insensitive restriction enzymes to target

sequencing toward CpG islands (e.g., the most commonly used enzyme is MspI, with a cut site

of CC/GG). CpG islands are regions of the genome with a high density of CpG sites and believed

to be functional hotspots for DNAm (Suzuki & Bird, 2008). Thus, RRBS not only enriches for

regions most likely to harbor functionally important methylation differences, but by also

profiling less than 10% of the genome for any individual it allows researchers to increase sample

sizes and read depth of regions covered (Bock et al., 2010; Laine et al., 2022). The large

differences in the types of data recovered from WGBS (wide breadth of the genome sequenced,

lower read depth, and smaller sample sizes) and RRBS (specifically targets CpG islands, higher

read depth, and larger sample sizes) make each approach better for particular types of research

questions. Because evolutionary and ecological studies often assess group-level variation, and

therefore require relatively large sample sizes, RRBS is becoming the preferred epigenetic

profiling method for these disciplines. However, few RRBS studies have directly compared its

results with WGBS, making it difficult to directly assess what biases or limitations are inherent

to each method.

Apart from the differences in library construction and experimental design, analysis

methods also differ for DNAm data. For example, bioinformatic tools apply different approaches

to account for unmethylated cytosines no longer matching the reference genome after bisulfite

treatment. Bismark (Krueger & Andrews, 2011), the most common tool for mapping bisulfite

converted sequences (Laine et al., 2022) (Table 1), handles mismatches by producing *in silico*

sense $(C \rightarrow T)$ and antisense $(G \rightarrow A)$ conversions at all CpG sites before aligning reads to a

reference genome that is itself converted using the same process. Reads are considered

ambiguous and discarded if they align with multiple versions of the *in silico* converted reference

genome. In addition to mapping reads, Bismark also streamlines data analysis by extracting

methylation information in a file format (i.e., bedGraph) that is compatible with downstream

statistical tools. However, this software relies on Bowtie2 (Langmead & Salzberg, 2012) for read

mapping, which often produces lower mapping efficiency (percent of reads mapped to the

reference) than other commonly used mapping algorithms such as BWA mem (Harrath et al.,

2019). Although tuning Bowtie2 mapping parameters can improve mapping rates, this process

can pose a substantial hurdle for novice users. Additionally, generating four intermediate *in*

silico conversions for both strands of the reference genome and sample reads leads to longer

computational run times and greater memory demands than alternative tools (Farrell et al., 2021;

Krueger & Andrews, 2011; Nunn et al., 2021).

BWA meth (Pedersen et al., 2014) is an alternative bisulfite sequence alignment script to

Bowtie2. This software is faster than Bismark because it only performs *in silico* conversion of

the reference genome prior to read mapping, not the sampled reads, and uses the BWA mem

algorithm (Nunn et al., 2021). However, a current limitation of BWA meth is that it stops after

mapping reads; the tool MethylDackel (https://github.com/dpryan79/MethylDackel) is

recommended for extracting methylation information. An advantage of MethylDackel is that it

uses overlaps between paired end sequencing data to discriminate between SNPs and

unmethylated cytosines. If a site is a bisulfite converted cytosine, the opposite strand should have

a G, otherwise it is likely a SNP. MethylDackel allows the user to specify the maximum

proportion of non-Gs permitted on the opposite strand before a site is considered a SNP and

excluded from the dataset. Importantly, this could impact estimates of DNAm variation if

polymorphic sites are frequently eliminated from analyses. To prevent erroneous filtering of sites

based on sequencing errors, the user can also specify the minimum depth the opposite strand

must have in order to be used for filtering. As polymorphism data are often unavailable for

natural populations, this feature has the potential to increase the reliability of DNAm research on

understudied systems.

Here, we provide a comparison of approaches for generating and analyzing DNAm data

that are particularly relevant to molecular ecologists. Specifically, we generated WGBS and

RRBS data from five populations of threespine stickleback (*Gasterosteus aculeatus*). Four

individuals were profiled using both WGBS and RRBS to assess how the methods diverge in: 1)

which and how many DNAm sites are recovered, 2) depth of coverage, and 3) distributions of

percentage DNAm per site. We then applied several bioinformatic pipelines to characterize the

performance of bisulfite alignment software and how they impact the mean and distribution of

percent DNAm for each individual. While studies have assessed performance of bioinformatic

pipelines for human and plant methylation data (Farrell et al., 2021; Nunn et al., 2021; Zhou et

al., 2024), to our knowledge this has not yet been done in ecologically-focused vertebrate

systems. We predicted that RRBS would provide fewer total DNAm sites, greater per-site read

depth, and that most DNAm would be found in CpG islands (Bock et al., 2010). We also

predicted that BWA mem-based read mappers would perform better than Bowtie-based mappers

in terms of total mapping efficiency, and that MethylDackel would efficiently remove

polymorphic sites. After explaining our findings, we discuss how to optimize sequencing

resources for ecological epigenetics, particularly for systems where DNA polymorphisms are

unknown. Finally, we consolidate information for each of our computational wrappers for

bisulfite alignment, providing open access code to encourage researchers to test various read

mapping methods, optimize their own bioinformatic pipelines, and increase the availability of

data for downstream analysis.

<u>Methods:</u>

Literature review of bisulfite library preparation and analysis methods

To assess how frequently Bismark, BWA meth, MethylDackel, and BWA mem are used

to analyze methylation data, we used "software name" AND "DNA methylation" to query Web

of Science and Google Scholar databases on August 21, 2024. We consolidated the number of

search results from each database in Table 1.

Sample Collection

We sampled wild and lab-reared threespine stickleback from five populations (Table 2).

RRBS was applied to all populations, while WGBS was used for common garden fish from two

Alaskan populations: Watson and Wik Lakes. Fish were euthanized with an overdose of buffered

MS-222 according to IACUC protocol L006460-A01. Field samples were dissected lakeside to

avoid introducing transportation stress that may influence DNAm profiles. We dissected and

stored liver tissue in RNALater (Sigma-Aldrich). We chose liver as our focal tissue because

previous work suggests liver-specific methylation is relatively conserved across vertebrates

(Klughammer et al., 2023). Liver also has relatively few cell types (Gumucio & Miller, 1981),

which should limit between sample variation due to cell type specific methylation. Two lab fish

from each of the Roberts, Gosling, and Sayward populations were exposed to the tapeworm

Schistocephalus solidus as part of a separate study, and one of the Sayward individuals was

successfully infected. None of these individuals displayed obvious differences in DNAm

patterns, and therefore were retained for further analysis (Supplemental Figure 1).

Sequencing

We extracted DNA from each liver sample using carboxyl coated magnetic beads

(BOMB.bio method: Oberacker et al., 2019). Bisulfite conversion, library preparation, and

sequencing were performed by Admera Health Biopharm Services. RRBS samples were bisulfite

treated and then converted to Illumina sequencing libraries and using Zymo-Seq RRBS Library

Kit. WGBS were similarly bisulfite converted using EZ DNA Methylation-Gold Kit (Zymo

Research, California, USA) and library preparation was done with xGen™ Methylation-

Sequencing DNA Library Preparation Kit (Integrated DNA Technologies, California, USA).

Sample quality was assessed using Quibit, agarose gel, and qPCR prior to sequencing. Libraries

were sequenced on Illumina Novaseq 6000 using paired end 150bp read chemistry. DNA

extractions and sequencing were performed in two batches. Each barcoded RRBS sample was

mixed into one of two sequencing pools (Table 2), which produced an average (± SEM) of 10.5

(± 1.3) million and 31.5 (± 15.4) million reads per sample following adapter trimming. All

WGBS samples were sequenced in one batch, resulting in 58.5 (± 1.0) million reads per sample

after adapter trimming.

Read Mapping and Methylation Calling

We trimmed adapters and barcodes using the -paired, -rrbs (for RRBS samples only, to

remove cytosines filled in at the end of reads as an artifact of library preparation), and -q 20

options in Trim Galore! v. 0.6.10 (https://github.com/FelixKrueger/TrimGalore). We then

aligned reads to the threespine stickleback reference genome (v.5; Nath et al., 2021) using

default parameters in three programs: BWA mem v. 0.7.17, BWA meth v. 0.2.7, and Bismark v.

0.24.2. Bismark resulted in a low mapping efficiency 28.10 (± 2.26%). We followed the

guidance in the documentation and aligned the data in single-end mode with the flag -

nondirectional, resulting in a mapping efficiency analogous to other studies. We therefore use the

data mapped in single end mode for the rest of the analysis. For BWA mem and BWA meth

alignments, samtools v. 1.9 (Danecek et al., 2021) was used to convert SAM outputs to BAM

format. Mapping efficiency is included in the Bismark output, and we used Bamtools v.2.5.2

(Barnett et al., 2011) to calculate it for BWA meth and BWA mem runs.

To extract methylation from BWA meth and BWA mem results we used MethylDackel v.

0.5.1 (https://github.com/dpryan79/MethylDackel). Specifically, we filtered SNPs, tabulated per

base methylation metrics, and created an output file compatible with methylKit in R (--

methylKit) (Akalin et al., 2012). As methylation call accuracy scales logistically with read depth

and becomes asymptotic at 10x (Seiler Vellame et al., 2021), we used --minDepth 10 to only

retain reads that were sequenced at least 10 times. It can be difficult to differentiate between

PCR replicates and true biological duplicates in RRBS data due to biased genomic fragmentation

(Laine et al., 2022). Therefore, for the RRBS samples only, we followed recommendations in the

MethylDackel documentation and applied the --keepDupes option. Three different SNP filtering

cut offs were tested (--maxVaraintFrac 0.25, 0.5, and 0.8) with --minOppositeDepth 5. To

prioritize capturing potential methylation variation, we used the most lenient SNP cut off (--

maxVariantFrac 0.8) for further analyses.

Statistical analyses

We applied a number of statistical tests in R v. 4.3.2 (R Core Team, 2023) to compare

read mapping approaches with both RRBS and WGBS data. We used the nonparametric

Kruskall-Wallis rank sum test to assess if mapping efficiency differed between Bismark, BWA

meth, and BWA mem. Post hoc pairwise comparisons were done using the Dunn test and p-

values were corrected for multiple comparisons using the Holm method in FSA v. 0.9.5 (Ogle et

al., 2023). Figures were generated using ggpubr v. 0.6.0 (Kassambara, 2023).

For RRBS samples, which were sequenced in two batches, we used Wilcox rank sum

exact tests to test whether batch affected mapping efficiency for each of the three read mapping

tools. Results were visualized using ggplot2 v. 3.5.1 (Wickham, 2016) and cowplot v. 1.1.3

(Wilke, 2024) (Supplemental Figure 2). The same procedure was used to analyze the number of

mapped reads (Supplemental Figure 3) and average read depth for each read mapping approach.

To determine if the resulting methylation profiles of individuals were in agreement

despite differences in read mapping and SNP calling, we compared both the mean and

distribution of percent methylation per individual obtained after using Bismark, BWA meth +

MethylDackel, and BWA mem + MethylDackel. We used methylKit v. 1.27.1 (Akalin et al.,

2012) to calculate the mean and visualize the distribution of percent methylation for each

sample. We then created a linear model in ggpmisc v. 0.6.1 (Aphalo, 2024) to compare the mean

percent methylation obtained using each pipeline. If, for every individual, the mean percent

methylation obtained from one pipeline reliably matches the mean percent methylation

calculated using another pipeline, then the slope and R^2 should be 1. If one method consistently

over or underestimates methylation compared to another, then the slope may be significantly

different from 1 while retaining a high R^2.

We next assessed differences between RRBS and WGBS. To determine if average read

depth per sample varied we used a Wilcox rank sum test. We then used ggvenn v.0.1.10 to

tabulate the proportion of CpG sites that were shared (1) between methods within an individual

and (2) between individuals using the same method. As we were interested in the breadth of

genomic coverage between library preparation methods, we did not filter SNPs prior to this

comparison and relaxed the minimum depth to test both 5x and 10x. Finally, the frequency

distribution of percent methylation per base was also used to visualize how the overall

methylation profile for an individual varied based on the sequencing method.

Results:

Read Mapping: BWA meth outperformed BWA mem and Bismark

BWA meth, BWA mem, and Bismark differed significantly in their mapping efficiency

using RRBS data (Kruskal-Wallis chi-squared = 75.95, df = 2, $p < 2.2 \times 10^{-16}$). With a mean ($\pm$

SD) mapping efficiency of 99.23% ($\pm$ 0.18) , BWA meth outperformed both Bismark (54.38 $\pm$

4.35%) and BWA mem (49.26 $\pm$ 4.46%) (Bismark - BWA meth: Z = -5.64, $p = 3.41 \times 10^{-8}$;

BWA mem - BWA meth: Z = -8.57, $p = 3.00 \times 10^{-17}$). The mapping rate for BWA mem was also

significantly lower than Bismark (Bismark - BWA mem: Z = 2.93, $p = 3.34 \times 10^{-3}$) (Figure 1).

Mapping efficiency for Bismark did not differ between sequencing batches (Wilcox rank sum

exact test, W = 142, p-value = 0.732), but there were small but significant batch effects for both

BWA meth (Wilcox rank sum exact test, W = 261, $p = 2.55 \times 10^{-8}$) and BWA mem (Wilcox rank

sum test, W = 253, p-value = 7.112×10^{-7}) (Supplemental Figure 2). Despite the batch effects,

BWA meth produced the highest mapping efficiency (>99%, 17.8 $\pm$ 13.5 million mapped reads

per RRBS sample), so its output was used for downstream analysis. For the WGBS samples,

BWA meth again performed the best with a mapping efficiency of 99.58 ($\pm$ 0.16%), resulting in

a mean of 58.6 ($\pm$ 1.1) million mapped reads. Although we had too few WGBS libraries to make

strong statistical inferences, the average WGBS mapping efficiency of BWA mem and Bismark

were again lower than BWA meth: 74.64% ($\pm$ 3.43) and 38.7% ($\pm$ 1.24), respectively.

BWA mem artificially inflates methylation

Despite significant differences in mapping efficiency and lack of SNP filtering with

Bismark, the mean methylation per putatively methylated site was highly similar between BWA

meth and Bismark (R^2 = 0.91, slope = 0.935; Figure 2a). BWA mem produced inflated

methylation rates compared to the other two methods (BWA meth - BWA mem: R^2 = 0.67,

Figure 2b; Bismark - BWA mem: R^2 = 0.64, Figure 2c). This bias was so extreme that the

median percent methylation for every sample mapped with BWA mem was 100%. We also

examined distribution of percent methylation per base to further characterize how read mapping

wrapper influenced which CpG sites were retained for analysis (Figures 3-5, Supplemental

Figures 4-8). BWA mem hardly recovered any CpG sites that were consistently unmethylated

(<10% of methylation) while these sites made up 51.6% ($\pm$ 14.8) and 43.7% ($\pm$ 13.9) of the data

in BWA meth and Bismark, respectively (Figure 4).

Differences in genomic breadth and depth of coverage between RRBS and WGBS

After applying BWA meth and MethylDackel the average median insert size of RRBS

reads was 38.06 ($\pm$ 2.44) compared to 242.25 ($\pm$ 22.47) for WGBS. Despite the small insert size,

RRBS resulted in a higher per site depth (mean of 16.39 $\pm$ 3.46) than WGBS (12.73 $\pm$ 0.06).

Batch 1 RRBS samples had a significantly lower read depth than batch 2 (14.73 $\pm$ 1.07 versus

19.44 $\pm$ 4.22; Wilcoxon rank sum exact test, W = 24.5, p-value = 0.0001151), but also a slightly

higher mapping efficiency (99.34 $\pm$ 0.00086% versus 99.03 $\pm$ 0.0013%; Wilcoxon rank sum

exact test, W = 261, p-value = 2.553e-08, Supplemental figures). This indicates that the

difference in depth of coverage between batches was driven by greater sequencing of batch 2, not

by better performance of read mapping software.

We next compared the effects of applying different library preparation methods to

technical replicates (i.e. RRBS vs WGBS; Table 3), as well as the

similarity of biological replicates from a single population (i.e.,

RRBS versus RRBS, WGBS versus WGBS; Table 4). Applying a minimum

depth filter of 10 reads per site on RRBS and WGBS libraries from the

same individual resulted in ≤1.0% shared CpG sites. Lowering the

minimum depth to 5 reads caused the proportion of shared CpG sites

to increase to 2.25% (± 0.42). Notably, at a minimum of 10x depth, >96%

of CpG sites were exclusive to WGBS, highlighting the large

difference in breadth of coverage between RRBS and WGBS (Table 3).

When assessing CpG sites shared among individuals using RRBS, the percent of overlap

depended on fish population. With a minimum of 10x depth, 13.1% and 27.2% of CpG sites

were shared between biological replicates from Watson and Wik Lakes, respectively. This

population effect was much reduced with WGBS; 18.8% of CpG sites were shared between

Watson Lake individuals and 19.7% between Wik individuals. Lowering the minimum depth to

5x increased the proportion of shared CpG sites using RRBS (28.4% for Watson, 40.1% for Wik)

and WGBS (57.6% for Watson, 59.0% for Wik) (Table 4).

Apart from comparing overlap in particular sites, the vast majority of the quantitative

CpG sites (i.e. the sites that are not always methylated or unmethylated), are not represented in

the RRBS data (Figures 3 - 5). At a minimum depth of 10x, the proportion of CpG sites with

methylation rates >10% and < 90% was twice as high in the WGBS samples ($55.63 \pm 2.49\%$)

than the RRBS samples ($28.56 \pm 8.37\%$). A higher proportion of intermediately methylated sites

were recovered in batch 2 of RRBS sequencing (Supplemental Figures), but none of these

samples were also analyzed via WGBS.

Discussion:

The increasing popularity of RRBS for methylation profiling has coincided with a rise in

the use of Bismark as the default method for read mapping and methylation calling, particularly

in ecologically focused experiments (Table 1). However, little effort has been dedicated to

understanding the strengths and limitations of these methodological choices, particularly in

genetically variable, wild organisms. By comparing RRBS and Bismark against less commonly

used library preparation and analysis alternatives, we identify several options for researchers to

maximize their sequencing resources.

Optimizing bisulfite alignment wrappers for ecological epigenetics

As predicted by previous studies (Farrell et al., 2021; Nunn et al., 2021; Zhou et al.,

2024), BWA meth outperformed both BWA mem and Bismark in its ability to map the majority

of sequenced reads to a reference genome (Figure 1). However, it is noteworthy that this

previous work focused on human and plant data. We found no previous studies assessing

bisulfite alignment wrappers in wild-caught/non-model systems. Despite differences in mapping

efficiency, BWA meth and Bismark provided consistent estimates of the mean percent of CpG

sites methylated per individual (Figure 2). In contrast, BWA mem substantially inflated estimates

of mean methylation rate (Figures 2 & 4). This difference reflects the optimization of particular

bioinformatic tools for bisulfite converted data. Because BWA mem does not account for C/T

mismatches induced during bisulfite conversion, reads enriched with converted thymines are

more likely to differ considerably from the reference sequence and be lost during mapping.

Importantly, BWA meth and Bismark only partially address this bias through bioinformatic

conversions that improve mapping. Bisulfite conversion damages DNA at unmethylated

cytosines during library preparation, which inherently inflates genome-wide methylation levels

(Olova et al., 2018). The magnitude of this bias is only recently becoming clear thanks to newer

library preparation approaches that forego harsh chemical reactions and instead rely on enzymes

to convert unmethylated cytosines (e.g., EM-Seq: Vaisvila et al., 2021). If precise estimates of

genomewide methylation are of key concern, then consideration should be given to using these

newer, non-bisulfite methods of library preparation. Importantly, EM-seq converts unmethylated

cytosines to thymine, making the sequencing data compatible with existing bioinformatic tools

for bisulfite converted data. Despite these advances, RRBS remains the most cost effective

method and is likely sufficient if the primary goal of a project is to measure or detect large

changes in methylation (i.e., those likely to have functional effects or are environmentally-

induced). Regardless of the library preparation method chosen, BWA meth or Bismark will

likely provide similar results.

For projects focusing on metrics related to per-site methylation, a key goal is to maximize

the number of usable sequencing reads. Bismark produced a mapping efficiency only slightly

higher than BWA mem, yet it remains the most commonly used read mapping method (Table 1).

Continued reliance on Bismark may partly be attributed to it being one of the first bioinformatic

tools designed to handle bisulfite converted data (Table 5). Because Bismark does *in silico*

conversions of all reads (sense and antisense, read and reference), its reliability is also agnostic

to library directionality. Although Bismark's ability to both map reads and call methylation

levels makes it particularly accessible for researchers new to epigenetics, the tool's strict default

parameters may cause difficulties when working with paired end reads, as we observed when we

initially had a 28% mapping efficiency. Although rarely discussed in the literature, low mapping

rates of paired end reads appears to be a relatively common issue with this tool (e.g.,: Nunn et

al., 2021; Zhou et al., 2024). Indeed, the developers provide thorough documentation to help

users resolve this mapping problem. The guidance highlights failure to trim adapters prior to read

mapping, potential contaminants, and high stringency under default parameters, among other

possible sources of error. Under default parameters, both paired end reads must map to one

unique version of the *in silico* converted reference genome, otherwise the paired end sequence

gets discarded. Following the advice in the documentation, and in-line with Nunn et al. (2021)

and Zhou et al. (2024), we relaxed this threshold by mapping our reads as non-directional. This

resulted in a mapping rate (54%) that is more in-line with previous DNAm studies in stickleback

(Heckwolf et al., 2020; Hu et al., 2021).

Despite requiring additional tools for methylation calling, BWA meth resulted in

significantly higher mapping rates than Bismark, leading to more data available for downstream

analysis. While a previous study found that BWA meth had higher error rates than alternative

tools (Farrell et al., 2021), this analysis was based on both directional and nondirectional

libraries. It is plausible that the increased error rate was driven by strand bias rather than being a

software limitation. Following BWA meth, we used MethylDackel to tabulate methylation and

filter SNPs. Conventional wisdom is that RRBS libraries should be single end, as paired end

reads may lead to bias by repeatedly sequencing the same sites (Krueger et al., 2012). However,

this bias can be reduced by normalizing the data based on median read depth, and if genetic

polymorphisms are unknown, the overlapping nature of paired end reads can be exploited by

software like MethylDackel to call SNPs. Additionally, using MethylDackel for SNP filtering

reduces errors associated with SNPs falsely being considered unmethylated cytosines. Similar to

most studies of ecological model organisms, we did not have genotype data for either the focal

individuals or populations, and were thus unable to verify the accuracy of SNP filtering using

MethylDackel.

Recently, there has been an influx of bisulfite alignment wrappers that combine the

strengths of BWA meth and Bismark. These software packages use BWA mem for read

mapping, are compatible with both directional and nondirectional libraries, and call methylation

(Table 5). Although our goal was to assess the most commonly used methods, future efforts

should aim to similarly compare the strengths and limitations of new software for ecologically-

focused experiments.

RRBS preferentially sequences stable methylation in comparison with WGBS

In addition to assessing how bisulfite alignment wrappers performed with RRBS data, we

examined how methylation profiles varied across technical and biological replicates using RRBS

and WGBS. Despite receiving relatively low amounts of RRBS data, which drove low read

depth, RRBS still reliably resulted in 29% higher sequencing depth than WGBS. However, few

CpG sites were commonly sequenced between technical and biological replicates using the two

sequencing methods (Tables 3 & 4). In line with previous research (Vaisvila et al., 2021), the

number of overlapping CpG sites between replicates greatly increased when the minimum depth

threshold was lower. When comparing the methylation patterns recovered from technical

replicates, the most striking difference between RRBS and WGBS was the proportion of

intermediately methylated sites recovered with both methods (Figures 3-5). Intermediate

methylation comprised 56% of the CpG sites sequenced with WGBS, while it was only 29% of

the CpG sites recovered with RRBS.

One potential explanation for the lack of congruence between sites sequenced with RRBS

and WGBS is that our sequencing parameters (i.e., 2x150bp reads) were too long for the small

inserts (~38bp) produced via RRBS. Therefore, using a longer read length resulted in extraneous

sequencing of adapters that were trimmed prior to mapping. However, the greater depth of batch

2 (19x compared to 14x for batch 1) suggests that investing in more sequencing can offset the

effects of adapter sequencing. Thus, we suggest selecting shorter read lengths that are still within

the constraints of mapping software (70bp) (Harrath et al., 2019). Accuracy of methylation calls

requires a minimum of 10x depth (Seiler Vellame et al., 2021). However, due to uneven

distribution of sequencing effort across the genome, as exhibited by the low proportion of

overlapping loci between our biological and technical replicates, we recommend using a

minimum depth threshold of 20x genomic coverage when selecting sequencing parameters.

Perhaps the most striking outcome from using technical replicates to contrast sequencing

methods was that the proportion of intermediately methylated sites sequenced with WGBS was

almost twice as high as intermediate methylation captured with RRBS (56% versus 29%,

respectively). Methylation can degrade if samples are collected multiple hours after mortality

(Rhein et al., 2015) or frozen for a long period of time (Lee et al., 2023), but the liver tissue we

sequenced was stored in RNALater immediately upon euthanasia and kept at -70°C for less than

one year. Therefore, degradation of intermediately methylated sites is unlikely the main driver of

this observation. Rather, there are technical and evolutionary explanations that likely interacted

to create this pattern. As previously mentioned, batch 2 RRBS samples recovered a much higher

proportion of intermediately methylated sites than batch 1 (Supplemental Figures), likely driven

by more sequencing. However, this higher rate ($37.83 \pm 5.78\%$) still appears to be substantially

lower than the average of WGBS. This pattern suggests that there may be fundamental,

biological differences in the methylation patterns surveyed by each method. Future studies could

use technical replicates to confirm this reduction of intermediately methylated sites in RRBS

data.

426   In considering how much sequencing to invest in RRBS, there may be evolutionary

arguments not to prioritize capturing intermediately methylated sites. A recent study in mouse

cell lines found that loci with intermediate methylation probabilistically, but not reliably,

transmitted methyl marks during mitosis. Furthermore, their methylation status had little effect

on gene expression (Hay et al., 2023). Another study in wild stickleback found that regions of

the genome containing stable methylation (<15% and >85% methylation) had stronger signatures

of selection than those containing intermediate methylation (Ord et al., 2023). These studies

suggest that because CpG islands are often enriched for methylation sites that have functional

importance (Jaenisch & Bird, 2003), they are more likely to be subject to selection, and thus may

have increased investment by regulatory machinery to maintain methylation state across all cells.

Thus, the lower proportion of intermediately methylated cytosines we sequenced with RRBS,

which targets CpG islands, may be picking up a biologically interesting pattern. Additionally,

given that molecular ecologists are likely interested in identifying functionally important

epigenetic effects, RRBS may provide a better option for detecting and measuring these changes.

As the field of ecological epigenetics continues to expand, we encourage researchers to continue

to interrogate which CpG sites are informative for evolutionary processes and consider how the

sequencing and bioinformatic methods they use may influence resolution of methylation profiles.

<u>Conclusions:</u>

We take a critical approach to analyzing RRBS and WGBS data using several

conventional methods. We highlight the strengths and weaknesses of both molecular and

bioinformatic methods, all of which should be considered during experimental design. If SNP

filtering in the absence of genotype data is a priority, researchers may consider choosing paired

end sequencing, contrary to traditional advice regarding RRBS library construction.

Additionally, the most commonly used tool for analyzing bisulfite sequencing data (Bismark)

performs significantly worse than alternative methods, such as BWA meth, in terms of

percentage of reads mapped. Researchers can increase the availability and reliability of data for

downstream analysis by considering alternative analysis pipelines (Table 5). Finally, we

highlight how different library preparation methods recover distinct signatures of per-site

methylation, and why these patterns warrant additional investigation for understanding how

epigenetics influence evolutionary processes.

Acknowledgements:

We thank Sean Schoville for computing resources, and Kristofer Sasser and Cole Wolf for

collecting fish from Vancouver Island. We recognize that the fish populations analyzed here

originate from lakes in the unceded territory of the Dena'ina and Coast Salish people and thank

them for their continued land stewardship. This work was funded by the Department of

Integrative Biology at UW-Madison and National Institute of General Medical Sciences

5R35GM142891-02.

References:

Adrian-Kalchhauser, I., Sultan, S. E., Shama, L. N. S., Spence-Jones, H., Tiso, S., Keller

Valsecchi, C. I., & Weissing, F. J. (2020). Understanding "Non-genetic" Inheritance:

Insights from Molecular-Evolutionary Crosstalk. *Trends in Ecology and Evolution*,

35(12), 1078–1089. https://doi.org/10.1016/j.tree.2020.08.011

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick,

A., & Mason, C. E. (2012). methylKit: A comprehensive R package for the analysis of

genome-wide DNA methylation profiles. *Genome Biology*, *13*(10), R87.

https://doi.org/10.1186/gb-2012-13-10-r87

Aphalo P (2024). ggpmisc: Miscellaneous Extensions to 'ggplot2'. R package version

0.6.1, https://CRAN.R-project.org/package=ggpmisc.

Banta, J.A., Richards, C.L. (2018). Quantitative epigenetics and evolution. *Heredity, 121,*

210-224. https://doi.org/10.1038/s41437-018-0114-x

Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P., & Marth, G. T. (2011).

BamTools: A C++ API and toolkit for analyzing and managing BAM files.

Bioinformatics, *27*(12), 1691–1692. https://doi.org/10.1093/bioinformatics/btr174

Bock, C., Tomazou, E. M., Brinkman, A. B., Müller, F., Simmer, F., Gu, H., Jäger, N.,

Gnirke, A., Stunnenberg, H. G., & Meissner, A. (2010). Quantitative comparison of

genome-wide DNA methylation mapping technologies. *Nature Biotechnology*, *28*(10),

1106–1114. https://doi.org/10.1038/nbt.1681

Bogan, S. N., Strader, M. E., & Hofmann, G. E. (2023). Associations between DNA

methylation and gene regulation depend on chromatin accessibility during

transgenerational plasticity. *BMC Biology*, *21*(1), 149. https://doi.org/10.1186/s12915-

023-01645-8

Charlesworth, D., Barton, N. H., & Charlesworth, B. (2017). The sources of adaptive

variation. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1855), 20162864.

https://doi.org/10.1098/rspb.2016.2864

Cholewa-Waclaw, J., Shah, R., Webb, S., Chhatbar, K., Ramsahoye, B., Pusch, O., Yu,

M., Greulich, P., Waclaw, B., & Bird, A. P. (2019). Quantitative modelling predicts the

impact of DNA methylation on RNA polymerase II traffic. *Proceedings of the National*

Academy of Sciences, *116*(30), 14995–15000. https://doi.org/10.1073/pnas.1903549116

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,

Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve

years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008.

https://doi.org/10.1093/gigascience/giab008

Elliott, G., Hong, C., Xing, X., Zhou, X., Li, D., Coarfa, C., Bell, R. J. A., Maire, C. L.,

Ligon, K. L., Sigaroudinia, M., Gascard, P., Tlsty, T. D., Harris, R. A., Schalkwyk, L. C.,

Bilenky, M., Mill, J., Farnham, P. J., Kellis, M., Marra, M. A., Milosavljevic, A., Hirst,

M., Stormo, G.D., Wang, T., & Costello, J. F. (2015). Intermediate DNA methylation is a

conserved signature of genome regulation. *Nature Communications*, *6*(1), 6363.

https://doi.org/10.1038/ncomms7363

Farrell, C., Thompson, M., Tosevska, A., Oyetunde, A., & Pellegrini, M. (2021).

BiSulfite Bolt: A bisulfite sequencing analysis platform. *GigaScience*, *10*(5).

https://doi.org/10.1093/gigascience/giab033

Gu, H., Smith, Z. D., Bock, C., Boyle, P., Gnirke, A., & Meissner, A. (2011). Preparation

of reduced representation bisulfite sequencing libraries for genome-scale DNA

methylation profiling. *Nature Protocols*, *6*(4), 468–481.

https://doi.org/10.1038/nprot.2010.190

Gumucio, J. J., & Miller, D. L. (1981). Functional implications of liver cell

heterogeneity. *Gastroenterology*, *80*(2), 393–403. https://doi.org/10.1016/0016-

5085(81)90732-0

Guo, W., Fiziev, P., Yan, W., Cokus, S., Sun, X., Zhang, M. Q., Chen, P.-Y., &

Pellegrini, M. (2013). BS-Seeker2: A versatile aligning pipeline for bisulfite sequencing

data. *BMC Genomics*, *14*(1), 774. https://doi.org/10.1186/1471-2164-14-774

Harrath, Y., Mahjoub, A., AbuBakr, F., & Azhar, M. (2019). Comparative Evaluation of

Short Read Alignment Tools for next Generation DNA Sequencing. *2019 International*

Conference on Innovation and Intelligence for Informatics, Computing, and Technologies

(3ICT), 1–6. https://doi.org/10.1109/3ICT.2019.8910272

Hay, A. D., Kessler, N. J., Gebert, D., Takahashi, N., Tavares, H., Teixeira, F. K., &

Ferguson-Smith, A. C. (2023). Epigenetic inheritance is unfaithful at intermediately

methylated CpG sites. *Nature Communications*, *14*(1), 5336.

https://doi.org/10.1038/s41467-023-40845-2

Heckwolf, M. J., Meyer, B. S., Häsler, R., Höppner, M. P., Eizaguirre, C., & Reusch, T.

B. H. (2020). Two different epigenetic information channels in wild three-spined

sticklebacks are involved in salinity adaptation. *Science Advances*, *6*(12), eaaz1138.

https://doi.org/10.1126/sciadv.aaz1138

Hu, J., & Barrett, R. D. H. (2017). Epigenetics in natural animal populations. *Journal of*

Evolutionary Biology, *30*(9), 1612–1632. https://doi.org/10.1111/jeb.13130

Hu, J., Wuitchik, S. J. S., Barry, T. N., Jamniczky, H. A., Rogers, S. M., & Barrett, R. D.

H. (2021). Heritability of DNA methylation in threespine stickleback (Gasterosteus

aculeatus). *Genetics*, *217*(1). https://doi.org/10.1093/GENETICS/IYAB001

Jaenisch, R., & Bird, A. (2003). Epigenetic regulation of gene expression: How the

genome integrates intrinsic and environmental signals. *Nature Genetics*, *33*(3S), 245–

254. https://doi.org/10.1038/ng1089

Kassambara A. (2023). ggpubr: 'ggplot2' Based Publication Ready Plots. R package

version 0.6.0, https://CRAN.R-project.org/package=ggpubr.

Klughammer, J., Romanovskaia, D., Nemc, A., Posautz, A., Seid, C. A., Schuster, L. C.,

Keinath, M. C., Lugo Ramos, J. S., Kosack, L., Evankow, A., Printz, D., Kirchberger, S.,

Ergüner, B., Datlinger, P., Fortelny, N., Schmidl, C., Farlik, M., Skjærven, K.,

Bergthaler, A., Liedvogel, M., Thaller, D., Burger, P.A., Hermann, M., Distel, M., Distel,

D.L., Kübber-Heiss, A., & Bock, C. (2023). Comparative analysis of genome-scale, base-

resolution DNA methylation profiles across 580 animal species. *Nature Communications*,

14(1), 232. https://doi.org/10.1038/s41467-022-34828-y

Krueger, F., & Andrews, S. R. (2011). Bismark: A flexible aligner and methylation caller

for Bisulfite-Seq applications. *Bioinformatics*, *27*(11), 1571–1572.

https://doi.org/10.1093/bioinformatics/btr167

Krueger, F., Kreck, B., Franke, A., & Andrews, S. R. (2012). DNA methylome analysis

using short bisulfite sequencing data. *Nature Methods*, *9*(2), 145–151.

https://doi.org/10.1038/nmeth.1828

Laine, V. N., Sepers, B., Lindner, M., Gawehns, F., Ruuskanen, S., & van Oers, K.

(2022). An ecologist's guide for studying DNA methylation variation in wild vertebrates.

Molecular Ecology Resources, 1755-0998.13624. https://doi.org/10.1111/1755-

0998.13624

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2.

Nature Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lea, A.J., Vilgalys, T.P., Durst, P.A.P., & Tung, J. (2017). Maximizing ecological and

evolutionary insight in bisulfite sequencing data sets. *Nature Ecology and Evolution, 1,*

1074–1083. https://doi.org/10.1038/s41559-017-0229-0

Lee, N. Y., Hum, M., Tan, G. P., Seah, A. C., Kin, P. T., Tan, N. C., Law, H.-Y., & Lee,

A. S. G. (2023). Degradation of methylation signals in cryopreserved DNA. *Clinical*

Epigenetics, *15*(1), 147. https://doi.org/10.1186/s13148-023-01565-y

Marin, P., Genitoni, J., Barloy, D., Maury, S., Gibert, P., Ghalambor, C. K., & Vieira, C.

(2020). Biological invasion: The influence of the hidden side of the (epi)genome.

Functional Ecology, *34*(2), 385–400. https://doi.org/10.1111/1365-2435.13317

Nath, S., Shaw, D. E., & White, M. A. (2021). Improved contiguity of the threespine

stickleback genome using long-read sequencing. *G3: Genes, Genomes, Genetics*, *11*(2).

https://doi.org/10.1093/g3journal/jkab007

Nunn, A., Otto, C., Stadler, P. F., & Langenberger, D. (2021). Comprehensive

benchmarking of software for mapping whole genome bisulfite data: From read

alignment to DNA methylation analysis. *Briefings in Bioinformatics*, *22*(5), bbab021.

https://doi.org/10.1093/bib/bbab021

Oberacker, P., Stepper, P., Bond, D. M., Höhn, S., Focken, J., Meyer, V., Schelle, L.,

Sugrue, V. J., Jeunen, G. J., Moser, T., Hore, S. R., von Meyenn, F., Hipp, K., Hore, T.

A., & Jurkowski, T. P. (2019). Bio-On-Magnetic-Beads (BOMB): Open platform for

high-throughput nucleic acid extraction and manipulation. *PLoS Biology*, *17*(1).

https://doi.org/10.1371/journal.pbio.3000107

Ogle DH, Doll JC, Wheeler AP, Dinno A (2023). *FSA: Simple Fisheries Stock*

Assessment Methods. R package version 0.9.5, https://CRAN.R-

project.org/package=FSA.

Olova, N., Krueger, F., Andrews, S., Oxley, D., Berrens, R. V., Branco, M. R., & Reik,

W. (2018). Comparison of whole-genome bisulfite sequencing library preparation

strategies identifies sources of biases affecting DNA methylation data. *Genome Biology*,

19(1). https://doi.org/10.1186/s13059-018-1408-2

Ord, J., Gossmann, T. I., & Adrian-Kalchhauser, I. (2023). High Nucleotide Diversity

Accompanies Differential DNA Methylation in Naturally Diverging Populations.

Molecular Biology and Evolution. https://doi.org/10.1093/molbev/msad068

Otterdijk, S. D., & Michels, K. B. (2016). Transgenerational epigenetic inheritance in

mammals: How good is the evidence? *The FASEB Journal*, *30*(7), 2457–2465.

https://doi.org/10.1096/fj.201500083

Pedersen, B. S., Eyring, K., De, S., Yang, I. V., & Schwartz, D. A. (2014). *Fast and*

accurate alignment of long bisulfite-seq reads (No. arXiv:1401.1129). arXiv.

http://arxiv.org/abs/1401.1129

R Core Team (2023). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rey, O., Eizaguirre, C., Angers, B., Baltazar-Soares, M., Sagonas, K., Prunier, J. G., &

Blanchet, S. (2020). Linking epigenetics and biological conservation: Towards a

conservation epigenetics perspective. *Functional Ecology*, *34*(2), 414–427.

https://doi.org/10.1111/1365-2435.13429

Rhein, M., Hagemeier, L., Klintschar, M., Muschler, M., Bleich, S., & Frieling, H.

(2015). DNA methylation results depend on DNA integrity—Role of post mortem

interval. *Frontiers in Genetics*, *6*. https://doi.org/10.3389/fgene.2015.00182

Seiler Vellame, D., Castanho, I., Dahir, A., Mill, J., & Hannon, E. (2021). Characterizing

the properties of bisulfite sequencing data: Maximizing power and sensitivity to identify

between-group differences in DNA methylation. *BMC Genomics*, *22*(1), 446.

https://doi.org/10.1186/s12864-021-07721-z

Suzuki, M. M., & Bird, A. (2008). DNA methylation landscapes: Provocative insights

from epigenomics. *Nature Reviews Genetics*, *9*(6), 465–476.

https://doi.org/10.1038/nrg2341

Takahashi, Y., Morales Valencia, M., Yu, Y., Ouchi, Y., Takahashi, K., Shokhirev, M.

N., Lande, K., Williams, A. E., Fresia, C., Kurita, M., Hishida, T., Shojima, K.,

Hatanaka, F., Nuñez-Delicado, E., Esteban, C. R., & Izpisua Belmonte, J. C. (2023).

Transgenerational inheritance of acquired epigenetic signatures at CpG islands in mice.

Cell, *186*(4), 715-731.e19. https://doi.org/10.1016/j.cell.2022.12.047

Tobi, E. W., Slieker, R. C., Luijk, R., Dekkers, K. F., Stein, A. D., Xu, K. M., Biobank-

based Integrative Omics Studies Consortium, Slagboom, P. E., van Zwet, E. W., Lumey,

L. H., & Heijmans, B. T. (2018). DNA methylation as a mediator of the association

between prenatal adversity and risk factors for metabolic disease in adulthood. *Science*

Advances, *4*(1), eaao4364. https://doi.org/10.1126/sciadv.aao4364

Vaisvila, R., Ponnaluri, V. K. C., Sun, Z., Langhorst, B. W., Saleh, L., Guan, S., Dai, N.,

Campbell, M. A., Sexton, B. S., Marks, K., Samaranayake, M., Samuelson, J. C., Church,

H. E., Tamanaha, E., Corrêa, I. R., Pradhan, S., Dimalanta, E. T., Evans, T. C., Williams,

L., & Davis, T. B. (2021). Enzymatic methyl sequencing detects DNA methylation at

single-base resolution from picograms of DNA. *Genome Research*, *31*(7), 1280–1289.

https://doi.org/10.1101/gr.266551.120

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York.

Wilke C (2024). cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R

package version 1.1.3, https://CRAN.R-project.org/package=cowplot.

Zemach, A., McDaniel, I. E., Silva, P., & Zilberman, D. (2010). Genome-Wide

Evolutionary Analysis of Eukaryotic DNA Methylation. *Science*, *328*(5980), 916–919.

https://doi.org/10.1126/science.1186366

Zhou, W., Johnson, B. K., Morrison, J., Beddows, I., Eapen, J., Katsman, E., Semwal, A.,

Habib, W. A., Heo, L., Laird, P. W., Berman, B. P., Triche, T. J., & Shen, H. (2024).

BISCUIT: An efficient, standards-compliant tool suite for simultaneous genetic and

epigenetic inference in bulk and single-cell studies. *Nucleic Acids Research*, *52*(6), e32–

e32. https://doi.org/10.1093/nar/gkae097

Author Contributions:

E.V.K. and J.N.W. conceived and designed the study and co-wrote the manuscript; E.V.K. wrote

the software wrappers, conducted analyses, and produced figures; J.N.W. supervised and funded

the work.

Tables and Figures:

Table 1: Search results for different bioinformatic tools associated with bisulfite converted DNA.
The search terms used were "software name" AND "DNA methylation," conducted on August
21, 2024.

Tool	Google Scholar Search Results	Web of Science Search Results
Bismark	6,030	30
BWA meth	313	7
BWA mem	3,570	0
MethylDackel	347	1

648 Table 2: Population, rearing environment, and sample sizes with associated batch of sequencing
649 for RRBS and WGBS. Abbreviations for populations and rearing environments are consistent
650 with figures in the results.

Population	Rearing Environment	Sample Size *(Sequencing Batch)*
Watson (WT, 60.54, -150.47)	Field (FD)	n=4 *(1)*
	Common Garden (CG)	n=4 *(1)*
		WGBS (n=2) *(1)*
Wik (WK, 60.72, -151.25)	Field	n=4 *(1)*
	Common Garden	n=4 *(1)*
		WGBS (n=2) *(1)*
Roberts (ROB, 50.22, -125.54)	Field	n=3 *(2)*
	Common Garden	n= 2 *(1)*, n = 2 *(2)*
Gosling (GOS, 50.05, -125.50)	Field	n=3 *(2)*
	Common Garden	n=2 *(1)*, n = 2 *(2)*
Sayward (SAY, 50.39, -125.95)	Common Garden	n=2 *(1)*, n = 2 *(2)*

655 Figure 1: Mapping efficiency of Bismark, BWA mem, and BWA meth on all **(A)** RRBS and **(B)**
656 WGBS samples.

659
660

Figure 2: Mean percent methylation per base for each individual sequenced with RRBS using **(A)** BWA meth and Bismark, **(B)** BWA meth and BWA mem, and **(C)** Bismark and BWA mem. A slope and R^2 that is close to one indicates agreement in mean percent methylation between methods.

Figure 3: Frequency histograms of percent methylation per CpG site for technical replicates from
Wik Lake **(A: RRBS, B: WGBS)** and Watson Lake **(C: RRBS, D: WGBS)**. The numbers on

675   top of each bin are the percentage of CpG sites that fall within that bin. Obtained from BWA
676   meth aligned reads with SNP filtering parameters in MethylDackel set to MaxVariantFrac 0.8.
677

681
682
683 Figure 4: Frequency histograms of percent methylation per CpG site for technical replicates from
684 Wik Lake **(A: RRBS, B: WGBS)** and Watson Lake **(C: RRBS, D: WGBS)**.. The numbers on
685 top of each bin are the percentage of CpG sites that fall within that bin. Obtained from BWA
686 mem aligned reads and SNP filtering parameters in MethylDackel set to MaxVariantFrac .8.

687
688
689

693
694
695 Figure 5: Histograms of percent methylation per CpG site for technical replicates from Wik Lake
696 **(A: RRBS, B: WGBS)** and Watson Lake **(C: RRBS, D: WGBS)**. The numbers on top of each

697 bin are the percentage of CpG sites that fall within that bin. Obtained from Bismark aligned
698 reads and no SNP filtering.
699
700 Table 3: CpG sites that were in common or unique to WGBS and RRBS for technical replicates
701 with a minimum of 10x depth and 5x depth.

Sample ID	Minimum Depth	Unique to WGBS (*Number of CpG sites*)	Unique to RRBS (*Number of CpG sites*)	Overlapping Sites (*Number of CpG sites*)
CG_WK_002	10x	96.6% *(5,417,478)*	2.4% *(134,797)*	1.0% *(55,001)*
	5x	96.0% *(18,501,995)*	1.1% *(219,188)*	2.8% *(546,037)*
CG_WK_003	10x	97.1% *(5,160,689)*	2.2% *(116,016)*	0.7% *(36,879)*
	5x	96.9% *(18,194,034)*	1.0% *(188,716)*	2.1% *(402,195)*
CG_WT_002	10x	98.8% *(4,966,649)*	0.9% *(46,124)*	0.3% *(14,544)*
	5x	97.1% *(17,915,424)*	1.1% *(194,016)*	1.8% *(337,685)*
CG_WT_004	10x	97.1% *(4,410,125)*	2.3% *(102,397*	0.7% *(31,069)*
	5x	96.5% *(17,312,003)*	1.2% *(209,426)*	2.3% *(418,643)*

703 Table 4: CpG sites that were in common between two individuals within the using RRBS and
704 WGBS with a minimum of 10x and 5x depth.

Sample Comparison	Sequencing Method	Minimum Depth	Unique to First Sample (*Number of CpG sites*)	Unique to Second Sample (*Number of CpG sites*)	Overlapping Sites (*Number of CpG sites*)
CG_WT_002 & CG_WT_004	RRBS	10x	22.2% *(38,133)*	64.6% *(110,931)*	13.1% *(22,535)*
		5x	30.4% *(274,939)*	41.1% *(371,307)*	28.4% *(256,762)*
	WGBS	10x	44.0% *(3,493,419)*	37.2% *(2,953,420)*	18.8% *(1,487,774)*
		5x	22.3% *(5,099,302)*	20.0% *(4,576,839)*	57.6% *(13,153,807)*
CG_WK_002 & CG_WK_003	RRBS	10x	43.2% *(116,469)*	29.5% *(79,556)*	27.2% *(73,329)*
		5x	39.0% *(377,396)*	21.0% *(203,082)*	40.1% *(387,829)*
	WGBS	10x	41.7% *(3,715,901)*	38.6% *(3,440,990)*	19.7% *(1,756,578)*

		5x	21.5% *(5,085,066)*	19.6% *(4,633,263)*	59.0% *(13,962,966)*

Table 5: A summary of the limitations and strengths of alignment wrappers available for bisulfite converted DNA. Qualitative descriptions of performance (Low, Moderate, High) are relative to the performance of other mapping tools for bisulfite converted libraries and collated from the results of multiple studies.

Tool	Algorithm	Mapping Efficiency	Library Type	SNP calling with bisulfite sequencing data	Sources
Bismark (Krueger & Andrews, 2011)	Bowtie2, HiSAT2	Low	Directional & Nondirectional	No	Present study, (Farrell et al., 2021; Guo et al., 2013; Nunn et al., 2021; Zhou et al., 2024)
BWA meth (Pedersen et al., 2014) + **MethylDackel** (https://github.com/dpryan79/MethylDackel)	BWA mem	High	Directional	Yes (can also use BISCUIT in place of MethylDackel)	Present study, (Farrell et al., 2021; Nunn et al., 2021; Zhou et al., 2024)
BISCUIT (Zhou et al., 2024)	BWA mem	High	Directional & Nondirectional	Yes	(Farrell et al., 2021; Zhou et al., 2024)
BiSulfite Bolt (Farrell et al., 2021)	BWA mem	High	Directional & Nondirectional	Yes	(Farrell et al., 2021; Zhou et al., 2024)

710