Variable performance of widely used bisulfite sequencing methods and read

mapping software for DNA methylation

Emily V. Kerns[1], Jesse N. Weber*[1]

Corresponding Author: Emily V. Kerns, Kerns3@wisc.edu, https://orcid.org/0009-0002-0538-

4632

Jesse N. Weber, Jnweber2@wisc.edu, https://orcid.org/0000-0003-4839-6684

[1]University of Wisconsin-Madison, Department of Integrative Biology, 430 Lincoln Drive,

Madison, WI 53706

<u>Funding Statement:</u> This work was funded by a startup award from the University of

Wisconsin-Madison Department of Integrative Biology (JNW), UW-Madison Department of

Integrative Biology Graduate Summer Research Award (EVK), and National Institute of General

Medical Sciences 5R35GM142891 (JNW).

<u>Conflict of Interest Disclosure:</u> We have no conflicts of interest.

<u>Ethics Approval Statement:</u> All animals were handled and euthanized in accordance with UW-

Madison IACUC protocol L006460-A01. Field sampling was done under the Alaska Department

of Fish & Game permit number SF2022-043 and BC Ministry of Forests permit number NA23-

787881.

<u>Abstract:</u>

DNA methylation (DNAm) is the most commonly studied marker in ecological

epigenetics, yet the performance of library preparation strategies and bioinformatic tools are

seldom assessed in genetically variable natural populations. We profiled DNAm in threespine

stickleback (*Gasterosteus aculeatus*) liver tissue, using reduced representation bisulfite

sequencing (RRBS) and whole genome bisulfite sequencing (WGBS) across technical and

biological replicates. We additionally collated publicly available RRBS and WGBS data from

taxonomically diverse organisms, and then compared how the most commonly used methylation

software (Bismark) performed relative to alternative pipelines (BWA meth, BiSulfite Bolt, and

Biscuit). Even after choosing parameters to maximize Bismark's mapping efficiency, it was still

outperformed by all other methods. Surprisingly, newer tools overrepresented DNAm compared

to older methods, highlighting the importance of testing methods on nonmodel organisms. There

were also distinct differences in DNAm profiles produced across library preparation methods,

with large impacts of population and read depth filters. Methylated sites unique to WGBS

predominantly mapped to introns and intergenic regions, while sites unique to RRBS primarily

overlapped with promoters and exons. Moreover, the prevalence of nucleotides with intermediate

methylation (within individuals) was greatly reduced in RRBS. Together, this suggests that

RRBS may be more useful for detecting functionally-relevant methylation differences. Based on

these results, we provide methodological recommendations for improving the reliability and

utility of DNAm profiles, particularly concerning the detection of functionally relevant DNAm

differences in genetically diverse natural populations.

Key words: DNA methylation, bisulfite sequencing, natural populations, bioinformatics,

Ecological epigenetics

Introduction:

Epigenetics is an umbrella term for chemical modifications to DNA and proteins, with

DNA methylation (DNAm) being among the most well studied epigenetic marks in ecologically

focused studies (Banta & Richards, 2018; Hu & Barrett, 2017; Lea et al., 2018; Rey et al., 2020).

As molecular ecologists increasingly explore the role of methylation in natural systems, there is a

need for increased attention to how methodological choices impact inferences of between and

within individual variation. This is particularly important given that most methylation profiling

methods were originally developed and applied in inbred laboratory model organisms.

There are multiple forms of DNAm, but the most abundant type in vertebrates is the

addition of a methyl group to cytosine at CpG dinucleotides (Jaenisch & Bird, 2003). DNAm has

a broad range of genomic effects, including silencing transposable elements (Marin et al., 2020),

blocking transcription factors and other proteins from binding to DNA, increasing the likelihood

of spontaneous deamination of methylated cytosines converting them to thymine (Ord et al.,

2023), and slowing down transcriptional machinery (Cholewa-Waclaw et al., 2019). While its

role varies, increased methylation is often associated with decreased gene expression (Zemach et

al., 2010). Multigenerational common garden experiments show that while DNAm profiles are

largely heritable they can also be modified by environmental stimuli (Bogan et al., 2023;

Brennan et al., 2025; Heckwolf et al., 2020; Hu et al., 2021). Moreover, environmentally-

induced shifts in DNAm can be inherited across generations (Takahashi et al., 2023; Tobi et al.,

2018). Recent findings also suggest that methylated regions experience different selection

pressures than background genomic DNA (Brennan et al., 2025; Ord et al., 2023). Despite

accumulating evidence that DNAm may play an important role in evolution, this topic is still an

active area for debate (Adrian-Kalchhauser et al., 2020; Charlesworth et al., 2017; Jaenisch &

Bird, 2003; Otterdijk & Michels, 2016). The increasing interest in understanding functional and

evolutionary roles of DNAm warrants greater focus on how methodological choices affect the

detection and quantification of DNAm.

DNAm is commonly measured using bisulfite sequencing methods (BS-seq). Treating

DNA with bisulfite converts unmethylated cytosines to uracil, which then becomes thymine

during PCR amplification. When compared to a reference genome, C/T mismatches are

identified as unmethylated cytosines, while any cytosines remaining are inferred to be

methylated. Although DNAm at any particular base pair is a binary trait (i.e., present/absent at a

CpG site on each read), the same location may not be methylated across cells or tissues. Some

CpG sites will be consistently methylated (or unmethylated) while others will have intermediate

methylation (Hay et al., 2023; Wang et al., 2020). When BS-seq is used for bulk sequencing of

many cells, the three possible methylation states of a CpG site in a single cell

(present/absent/heterozygous) are summed and become a quantitative metric (i.e., proportion of

total sites that are methylated). Accurately measuring this quantitative variation is contingent on

the number of times that a given base pair is represented in the sequencing data, often referred to

as read depth.

The two most commonly used BS-seq methods are whole genome (WGBS) and reduced

representation bisulfite sequencing (RRBS; Gu et al., 2011). While WGBS offers the potential to

profile all CpG sites, it requires large amounts of sequencing data in order to repeatedly sample

individual DNAm sites. This requirement often limits WGBS to experiments involving few

biological replicates and relatively low read depth, which reduces accuracy of methylation calls

and statistical power to detect group-level differences (Seiler Vellame et al., 2021). Additionally,

because WGBS does not enrich for CpG sites, much of the sequence data is uninformative for

epigenetic studies. For example, 70-80% of mapped reads in human WGBS studies do not

contain CpG dinucleotides (Elliott et al., 2015). In contrast, RRBS uses methylation-insensitive

restriction enzymes to target sequencing toward CpG islands (e.g., the most commonly used

enzyme is MspI, with a cut site of CC/GG). CpG islands are regions of the genome with a high

density of CpG sites and believed to be functional hotspots for DNAm (Suzuki & Bird, 2008).

Thus, RRBS not only enriches for regions most likely to harbor functionally important

methylation differences, but also by profiling <10% of the genome researchers can increase

sample sizes and read depth of regions covered (Bock et al., 2010; Laine et al., 2022). The large

differences in the types of data recovered from WGBS (wide breadth of the genome sequenced,

lower read depth, and smaller sample sizes) and RRBS (specifically targets CpG islands, higher

read depth, and larger sample sizes) make each approach better for particular types of research

questions. Because evolutionary and ecological studies often assess group-level variation, and

therefore require relatively large sample sizes, RRBS is becoming the preferred epigenetic

profiling method for these disciplines. However, few RRBS studies have directly compared its

results with WGBS, making it difficult to directly assess what biases or limitations are inherent

to each method.

Apart from the differences in library construction and experimental design, DNAm also

requires specialized analysis methods. For example, bioinformatic tools apply different

approaches to account for unmethylated cytosines no longer matching the reference genome after

bisulfite treatment. Bismark (Krueger & Andrews, 2011), the most common tool for mapping

bisulfite converted sequences (Laine et al., 2022) (Table 1), handles mismatches by producing *in*

silico sense $(C \rightarrow T)$ and antisense $(G \rightarrow A)$ conversions at all CpG sites before aligning reads

to a reference genome that is itself converted using the same process. Reads are considered

ambiguous and discarded if they align with multiple versions of the *in silico* converted reference

genome. However, this software relies on Bowtie2 (Langmead & Salzberg, 2012) for read

mapping, which often produces lower mapping efficiency (percentage of reads mapped to the

reference) than other commonly used mapping algorithms such as BWA mem (Harrath et al.,

2019). Notably, BWA mem uses local alignment, meaning that if a small portion of a sample

read does not align well to the reference genome it is clipped for a small penalty. Meanwhile,

Bowtie2 requires global alignment, requiring the entire read to successfully map without

additional trimming. Tuning Bismark mapping parameters can improve mapping efficiency but

may pose a substantial hurdle for novice users. Additionally, generating four intermediate *in*

silico conversions for both strands of the reference genome and sample reads leads to longer

computational run times and greater memory demands than alternative tools (Farrell et al., 2021;

Krueger & Andrews, 2011; Nunn et al., 2021).

Table 1: Search results for different bioinformatic tools associated with bisulfite converted
DNA. The search terms used were "software name" AND "DNA methylation," conducted on
September 8, 2025.

Tool	Google Scholar Search Results	Web of Science Search Results
Bismark	7,110	36
BWA meth	396	8
BiSulfite Bolt	74	1
Biscuit	372	5
MethylDackel	479	1

While Bismark remains the most popular BS-seq aligner, there are a number of

alternative software wrappers that use the BWA mem algorithm. BWA meth (Pedersen et al.,

2014), the oldest such wrapper, is faster than Bismark because it only performs *in silico*

conversion of the reference genome prior to read mapping, not the sampled reads (Nunn et al.,

2021; Sun et al., 2018). However, a current limitation of BWA meth is that it stops after mapping

reads, recommending users rely on MethylDackel (https://github.com/dpryan79/MethylDackel)

to tabulate methylation. BiSulfite Bolt (Farrell et al., 2021) and Biscuit (Zhou et al., 2024) are

more recent aligners that also perform methylation calling. Both BiSulfite Bolt and Biscuit create

a 3-base BWA-indexed reference genome that has been bisulfite converted *in silico* (Farell et al.,

2021; Zhou et al., 2024). However, Biscuit is unique in that it also creates a 4-base indexed

reference and uses asymmetrical alignment to increase mapping efficiency of unmethylated

cytosines. Biscuit will map T's (bisulfite-converted unmethylated cytosines on the forward

strand) and A's (bisulfite-converted unmethylated cytosines on the reverse strand) from the

sample read to C's and G's in the reference genome, but a G and C in the sample cannot be

aligned with an A and T in the reference. These methods have yet to be thoroughly compared

across a diverse set of wild organisms.

When working with BS-seq data from genetically variable organisms, C/T mismatches

between the sample read and reference can result from either an unmethylated cytosine or a

nucleotide polymorphism. While there are a number of available tools to filter SNPs from BS-

seq data (Lindner et al., 2022), here we focus on MethylDackel

(https://github.com/dpryan79/MethylDackel). MethylDackel allows the user to specify the

maximum proportion of non-Gs permitted on the opposite strand before a site is considered a

SNP and excluded from the dataset. Notably, depth influences the accuracy of SNP filtering with

this method, which may be constrained if higher depth was sacrificed for increased genomic

breadth (RRBS versus WGBS). Importantly, SNP filtering may impact estimates of DNAm

variation if polymorphic sites are frequently eliminated from analyses. This type of filtering may

be particularly important for quantifying DNAm in natural populations where individual

genotypes are unknown.

Here, we provide a comparison of approaches for generating and analyzing DNAm data

that are particularly relevant to molecular ecologists. Specifically, we generated WGBS and

RRBS data from five ecologically-divergent populations of threespine stickleback (*Gasterosteus*

aculeatus). Four individuals were profiled using both WGBS and RRBS to assess how the

methods diverge in: 1) which and how many DNAm sites are recovered, 2) depth of coverage,

and 3) distributions of percentage DNAm per site. We then compared the results of several

bioinformatic pipelines when applied our stickleback dataset and five other taxonomically-

diverse organisms (cichlid (*Astatotilapia calliptera*), purple sea urchin (*Strongylocentrotus*

purpuratus), great tit (*Parus major*), coral (*Acropora nana*), and stick insect (*Timema cristinae*)).

While studies have assessed performance of bioinformatic pipelines for human and plant

methylation data (Farrell et al., 2021; Nunn et al., 2021; Sun et al., 2018; Zhou et al., 2024), to

our knowledge this has not yet been done in a comparative context that considers natural animal

populations despite taxonomic variation in genome-wide methylation patterns (Klughammer et

al., 2023). We predicted that RRBS would provide fewer total DNAm sites, greater per-site read

depth, and that most DNAm would be found in CpG islands (Bock et al., 2010). We also

predicted that BWA mem-based read mappers would perform better than Bowtie-based mappers

in terms of total mapping efficiency. After explaining our findings, we discuss how to optimize

sequencing resources for ecological epigenetics, particularly for systems where DNA

polymorphisms are unknown. Finally, we consolidate information for each of our computational

wrappers for bisulfite alignment, providing open access code to encourage researchers to test

various read mapping methods, optimize their own bioinformatic pipelines, and increase the

availability of data for downstream analysis.

Methods:

Sample and Data Collection

We sampled liver tissue from wild and lab-reared threespine stickleback across five

populations (Supplemental Table 1; UW-Madison IACUC protocol L006460-A01; Alaska

Department of Fish & Game permit number SF2022-043, BC Ministry of Forests permit number

NA23-787881). RRBS was applied to all samples (n=34), while WGBS (n = 4) was used for

common garden fish from two Alaskan populations: Watson and Wik Lakes. We extracted DNA

from each sample using carboxyl coated magnetic beads (BOMB.bio method: Oberacker et al.,

2019). Bisulfite conversion, library preparation, and sequencing were performed by Admera

Health Biopharm Services. Some fish were exposed to the tapeworm *Schistocephalus solidus* as

part of a separate study. None of these individuals displayed obvious differences in DNAm

patterns, and therefore were retained for further analysis (Supplemental Figure 1). Notably, each

barcoded RRBS sample was mixed into one of two sequencing pools (Supplemental Table 1),

which produced an average (± SEM) of 10.5 (± 1.3) million and 31.5 (± 15.4) million reads per

sample following adapter trimming. Given that CpG islands comprise of ~8.5% of the threespine

stickleback genome (see *Genomic Annotations of RRBS-WGBS Technical Replicates* for CpG

island definition), this is predicted to result in a depth of 20.7x and 62.2x for batch 1 and 2,

respectively. Based on PC1 from a PCA (methylKit v.1.33.3, Akalin et al., 2012) separated

samples by sequencing batch, but batch effects dissipated along PC2 (Supplemental Figure 2).

All WGBS samples were sequenced in one batch, resulting in 58.5 (± 1.0) million reads per

sample after adapter trimming, which is predicted to result in an average depth of 9.8x.

In addition to generating our own data from stickleback, we used publicly available data

from five other animals to assess the performance of BS-seq bioinformatic pipelines

(Supplemental Table 2). Our four stickleback samples were the only data we could find in which

the same samples were used for both RRBS and WGBS. We gathered RRBS data from a cichlid

(*Astatotilapia calliptera*; n = 33; Vernaz et al., 2021), great tit (*Parus major*; n = 61; Viitaniemi

et al., 2019), and purple sea urchin (*Strongylocentrotus purpuratus*; n = 12; Bogan et al., 2023).

We used WGBS data from a cichlid (*Astatotilapia calliptera*; n = 6; Vernaz et al., 2021), coral

(*Acropora nana*; n = 30; Guerrero & Bay, 2024), and stick insect (*Timema cristinae*; n = 7; de

Carvalho et al., 2023). All of the publicly available RRBS datasets were single end reads, while

the WGBS data were paired end reads.

Read Mapping

We applied the same bioinformatic pipelines to all datasets. First, we used the -q 20

option in Trim Galore! v. 0.6.10 (https://github.com/FelixKrueger/TrimGalore) to trim adapters

and barcodes, also using the -paired and -rrbs options (where appropriate) to remove cytosines

filled in at the end of reads as an artifact of library preparation. We then aligned reads to the

reference genome of each organism (Supplemental Table 2) using default parameters in four

programs: BWA meth v. 0.2.7, Bismark v. 0.24.2, BiSulfite Bolt v. 1.4.8, and Biscuit v. 1.7.0.

Bismark resulted in a low mapping efficiency 28.10 (± 2.26%) for the RRBS stickleback

samples, so we followed the guidance in the documentation and aligned the data in single-end

mode with the flag -nondirectional, resulting in a mapping efficiency analogous to other studies.

As Bismark performs global alignment while all of the other tools perform local alignment, we

also used the -local option to have Bismark perform local alignment ("Bismark Local"), making

it more comparable to the other methods. For BWA meth, BiSulfite Bolt, and Biscuit alignments,

samtools v. 1.9 (Danecek et al., 2021) was used to convert SAM outputs to BAM format. We

used Bamtools v.2.5.2 (Barnett et al., 2011) to calculate mapping efficiency for each method

except Bismark, which includes mapping efficiency in the output. For all other methods, we

calculated mapping efficiency as the percentage of reads mapped to the genome - the percent of

mapped reads that failed QC.

Methylation and SNP Calling

We sought to understand how choice of read mapping method may influence downstream

methylation profiles. We therefore used MethylDackel v. 0.5.1

(https://github.com/dpryan79/MethylDackel) to call methylation and filter SNPs from the

threespine stickleback bam files produced with BWA meth, BiSulfite Bolt, and Biscuit, even

though BiSulfite Bolt and Biscuit have their own methylation calling software. Methylation call

accuracy scales logistically with read depth and becomes asymptotic at 10x (Seiler Vellame et

al., 2021), so we only retained sites that were sequenced at least 10 times. It can be difficult to

differentiate between PCR replicates and true biological duplicates in RRBS data due to biased

genomic fragmentation (Laine et al., 2022). Therefore, for the RRBS samples only, we followed

recommendations in the MethylDackel documentation and applied the --keepDupes option. To

prioritize capturing methylation variation, we used a lenient SNP filter (--maxVariantFrac 0.8, --

minDepth 5) in MethylDackel. The final methylation file was modified to be compatible as an

input to methylKit in R (Akalin et al., 2012). Bismark's bam files were incompatible with

MethylDackel, so methylation calls were performed using the standard Bismark algorithm.

Statistical analyses of BS-seq Alignment Wrappers

We applied a number of statistical tests in R v. 4.3.2 (R Core Team, 2023) to compare

read mapping approaches with both RRBS and WGBS data. We used the nonparametric

Friedman test of repeated measures to assess whether the mapping efficiency of individual

libraries differed between alignment software. Effect sizes were calculated using Kendall's W,

with scores of 0.1-0.3 indicating small effect, 0.3-0.5 moderate effect, and >0.5 is a large effect

Post hoc pairwise comparisons were done using a Wilcox signed-rank test, with p-values

corrected for multiple comparisons using the Bonferroni method in rstatix v. 0.7.2 and coin

v.1.4-3 (Hothorn et al., 2008; Kassambara, 2023).

To determine if the resulting methylation profiles of individuals were in agreement

despite differences in read mapping approach, we created a linear model in ggpmisc v. 0.6.1

(Aphalo, 2024) to compare the mean percent methylation obtained using each pipeline. A slope

near 1 and intercept near zero indicates high agreement between pipelines. We then visualized

the relationship between the residuals of each model and per-sample sequencing depth,

predicting that agreement between software would increase with depth.

Genomic Annotations of RRBS-WGBS Technical Replicates

We next assessed differences between RRBS and WGBS results in stickleback. We

tabulated the proportion of CpG sites that were shared either (1) between methods within an

individual (i.e. technical replicates; Supplemental Table 3) or (2) between individuals using the

same method (i.e. biological replicates; Supplemental Table 4). As we were interested in the

breadth of genomic coverage between library preparation methods, we did not filter SNPs prior

to this comparison and relaxed the minimum depth (tested at both 5x and 10x). The frequency

distribution of percent methylation per base was used to visualize how the overall methylation

profile for an individual varied based on the library preparation method.

To determine where RRBS and WGBS vary in their coverage of the genome, we

conducted genomic annotations on CpG sites that were sequenced with both methods, RRBS

only, and WGBS only with a minimum depth of 5x. TaJoCGI (Nell, 2020) identified CpG

islands in the v.5 threespine stickleback reference genome (Nath et al., 2021) based on the

algorithm from Takai & Jones (2002). We quantified the overlap between methylation sites and

CpG islands and shores (±2 kbp CpG islands) with genomation v. 1.34.0 (Akalin et al., 2015).

We next assessed differences in functional regions captured by the two BS-seq methods. We

specifically examined variation in 1) the proportion of CpG sites that overlapped with promoters,

gene bodies, and intergenic regions and 2) the proportion of all promoters and gene bodies in the

genome that were captured. For more information regarding library preparation, sequencing, and

analysis, refer to the supplemental methods.

Results:

Figure 1: Interspecific variation in mapping efficiency of five alignment tools using **(A)** RRBS
and **(B)** WGBS data. Layout refers to whether samples were sequenced using paired end or
single end reads. All WGBS data was sequenced with paired end read chemistry. Symbols
(Biorender) denote the read mapping software that resulted in the highest mapping efficiency for
each organism. Threespine stickleback are the dark blue fish, while cichlid are the dark gray fish.

Read Mapping: BWA mem-based algorithms outperform Bismark

The five sequence alignment tools produced significantly different mapping efficiency

(RRBS: Friedman test $p_{adj} = 1.30 \times 10^{-60}$, df = 4, Kendall's W = 0.514, Figure 1A; WGBS:

Friedman test $p_{adj} = 3.04 \times 10^{-39}$, df = 4, Kendall's W = 0.992, Figure 1B). The most consistent

pattern was that local mapping outperformed Bismark's default method of global alignment.

Simply running Bismark with the "-local" parameter greatly increased mapping efficiency

(RRBS Bismark - Bismark Local: $p_{adj} = 1.38 \times 10^{-23}$, effect size r = 0.868; WGBS Bismark -

Bismark Local: $p_{adj} = 2.47 \times 10^{-8}$, effect size r = 0.871). Despite this increase in performance,

BWA mem-based aligners tended to result in higher mapping efficiency than Bismark Local. For

example, nearly 100% of the Cichlid sample reads were mapped to the reference using Biscuit

and BWA meth (RRBS Biscuit: $99.5 \pm 0.103\%$ (SD), RRBS BWA meth: $98.7 \pm 0.564\%$),

compared to a mean mapping rate of $85.2 \pm 1.56\%$ RRBS reads with Bismark Local. Meanwhile,

BiSulfite Bolt was clearly the best method for Great Tits in terms of mapping efficiency ($93.0 \pm$

2.58%), with a mapping rate almost 12% higher than the second best aligner (Great Tit BiSulfite

Bolt - Bismark Local: $p_{adj} = 1.38 \times 10^{-23}$, effect size $r = 0.870$). Overall, Biscuit displayed the

highest mapping efficiency, closely followed by BiSulfite Bolt (RRBS Bisulfite Bolt - Biscuit:

$p_{adj} > 0.05$, effect size $r = 0.220$; WGBS: $p_{adj} = 1.85 \times 10^{-9}$, effect size $r = 0.804$) and BWA meth

(RRBS Biscuit - BWA meth: $p_{adj} = 1.24 \times 10^{-6}$, effect size $r = 0.444$; WGBS: $p_{adj} = 1.42 \times 10^{-13}$,

effect size $r = 0.871$).

Because a PCA revealed clear sequencing batch effects among the stickleback samples

(Supplemental Figure 2), we assessed whether variation in mapping efficiency across alignment

tools was connected to batch effects. Although Bismark's mapping efficiency did not differ

between sequencing batches (Wilcox rank sum exact test, $W = 142$, p-value $= 0.732$), there were

small but significant batch effects for every other BS-seq aligner (Supplemental Figure 3).

Figure 2: Mean percent methylation per base for individual stickleback sequenced with RRBS and analyzed using **(A)** BWA meth and Bismark, **(B)** Bismark global alignment and Bismark local alignment, **(C)** BWA meth and Biscuit, and **(D)** BiSulfite Bolt and Biscuit. A slope that is close to one and y-intercept near zero indicates agreement in mean percent methylation between read mapping methods.

Figure 3: Total sequences per sample compared to the residuals (Figure 2) from models comparing **(A)** BWA meth and Bismark, **(B)** Bismark global versus Bismark local alignment, **(C)** BWA meth versus Biscuit, and **(D)** BiSulfite Bolt versus Biscuit.

Read mapping software & sequencing effort influence reliability of downstream methylation

calls

We next tested if read mapping methods produced similar methylation profiles in

threespine stickleback sequenced with RRBS despite significant differences in mapping

efficiency. While Bismark resulted in the lowest mapping efficiency among all read aligners we

tested (54.4 ± 4.35%), the average percent methylation of each individual was remarkably

similar to that produced from BWA meth, which had an average mapping efficiency of $85.3 \pm$

3.23% (slope = 0.935, intercept = 9.24; Figure 2A). Next, because Bismark Local had a higher

mapping efficiency than Bismark ($65.2 \pm 5.55\%$ versus $54.4 \pm 4.35\%$, respectively; $p_{adj} = 5.63$ x

10^{-6}, effect size r = 0.873), we compared the average percent methylation of each sample

recovered from these two methods, resulting in highly similar methylation profiles (slope =

0.923, intercept = 4.52; Figure 2B). Surprisingly, there was large disagreement between the

results of older (Bismark and BWA meth) and newer methods (Biscuit and BiSulfite Bolt). Mean

percent methylation per fish was much higher in Biscuit compared to BWA meth (slope = 0.377,

intercept = 57.8, Figure 2C). While BiSulfite Bolt and Biscuit were more similar to each other

than they were to BWA meth or Bismark, the lower slope and higher intercept indicate that

average methylation was not as consistent between these two methods (slope = 0.793, intercept =

14.2, Figure 2D) as it was between BWA meth and Bismark.

We next leveraged the variation in sequencing effort among the stickleback sequencing

batches to assess whether the total number of reads per sample influenced the reliability of

methylation tabulation (Figure 3). With low sequence coverage there was wide variation between

methods. An exponential decay model was a better predictor of the data than a linear model for

all four methods comparisons. With the exception of the BWA meth-Biscuit residuals (Figure

3C), sequencing effort was a significant predictor of reduced residuals (BWA meth-Bismark: t =

-3.061, p = 0.00444, Figure 3A; Bismark-Bismark Local: t = -2.084, p = 0.0455, Figure 3B;

BiSulfite Bolt-Biscuit: t = -3.121, p = 0.00381, Figure 3D). Agreement in methylation calls

increased with sequence coverage until approximately 20 million reads per sample, but

increasing sequencing effort past this threshold provided little to no improvement (Figure 3).

We next examined distribution of percent methylation per base to further characterize

how read mapping software influenced which CpG sites were retained for analysis (Figure 4,

Supplemental Figures 4-6). Distributions obtained from older methods (Figure 4A-C) differed

greatly from newer methods (Figure 4D-E), consistent with the previously noted differences in

mean percent methylation obtained from each read mapping method (Figure 23). Notably, both

Biscuit and BiSulfite Bolt recovered almost no unmethylated cytosines (percent methylation <

10%), while BWA meth, Bismark, and Bismark Local were dominated by unmethylated

cytosines. Additionally, BWA meth, Bismark, and Bismark Local contained very few

intermediately methylated sites (10-90% methylation).

366
Figure 4: Frequency histograms of percent methylation per base for one RRBS sample from

Wik lake (sample ID: CG_WK_002), analyzed with either **(A)** BWA meth, **(B)** Bismark, **(C)**

Bismark Local, **(D)** Biscuit, or **(E)** BiSulfite Bolt.

Differences in genomic breadth and depth of coverage between RRBS and WGBS

After running BWA meth and MethylDackel, the average median insert size of RRBS

reads was 38.06 (± 2.44) compared to 242.25 (± 22.47) for WGBS. Despite the larger insert

sizes and greater sequencing effort per sample for WGBS (RRBS batch 1: 10.5 ± 1.3M reads;

RRBS batch 2: 31.5 ± 15.4M reads; WGBS: 58.5 ± 1.0M reads), RRBS still resulted in a higher

per site depth (16.39 ± 3.46) than WGBS (12.73 ± 0.06).

We next compared the effects of applying different library preparation methods to

technical replicates (i.e. RRBS vs WGBS; Supplemental Table 3). Applying a minimum depth

filter of 10 reads per site on RRBS and WGBS libraries from the same individual resulted in

≤1.0% shared CpG sites. Lowering the minimum depth to 5 reads increased the proportion of

shared sites to increase to 2.25% (± 0.42). Notably, at a minimum of 10x depth, >96% of CpG

sites were exclusive to WGBS, highlighting the large difference in breadth of coverage between

RRBS and WGBS (Supplemental Table 3).

When assessing CpG sites shared among biological replicates using RRBS, the percent of

overlap depended on fish population. With a minimum of 10x depth, 13.1% and 27.2% of CpG

sites were shared between individuals from either Watson or Wik Lakes, respectively. This

population effect was much reduced with WGBS; 18.8% of CpG sites were shared between

Watson Lake individuals and 19.7% between Wik individuals. Lowering the minimum depth to

5x increased the proportion of shared CpG sites using RRBS (28.4% for Watson, 40.1% for Wik)

and WGBS (57.6% for Watson, 59.0% for Wik) (Supplemental Table 4).

Notably, the vast majority of the intermediately methylated CpG sites (i.e. the sites that

are not always methylated or unmethylated), are not represented in the RRBS data (Figure 5). At

a minimum depth of 10x, the proportion of CpG sites with methylation rates >10% and < 90%

was nearly two-fold higher in WGBS (55.63 ± 2.49%, Figure 5A) than RRBS samples (28.56 ±

8.37%, Figure 5B). We recovered a higher proportion of intermediately methylated sites in batch

2 of RRBS sequencing (Supplemental Figure 7), but none of these samples were also analyzed

via WGBS.

Methodological differences in genomic breadth were also evident when assessing the

distribution of CpG sites across functional categories. We conducted annotations using a

minimum depth threshold of 5x to increase the number of shared CpG sites between library

preparation methods, enhancing our ability to meaningfully compare the methods. CpG sites

unique to WGBS recovered nearly all CpG islands, shores, and promoters, and 90% of exons and

introns present in the threespine stickleback reference genome (Figure 6C & F). Additionally, the

number of CpG sites available for functional annotations was 14.7x higher from sites unique to

WGBS than sites unique to RRBS (Figure 6B & E). Because of the large difference in

sequencing breadth, the intersection of both methods (i.e. CpG sites in common between WGBS

and RRBS) reflects the same patterns found from sites recovered in RRBS only. However, CpG

sites that were unique to RRBS still captured nearly 50% of CpG islands (Figure 6C) and 60% of

promoters in the stickleback genome. Relatively few gene bodies were captured with RRBS (i.e.,

<15% of exons and introns; Figure 6F).

While WGBS clearly provides a larger breadth of coverage, RRBS enriches for

functional genomic regions. When averaged across replicates from the same population (n=2),

almost half of CpG sites unique to RRBS were found within CpG islands and 37% were within

promoters. Meanwhile, a much smaller proportion of sites unique to WGBS overlapped with

CpG islands (27%) and promoters (15%) (Figure 6A & D). Additionally, 26% of WGBS-only

sites were in intergenic regions, compared to 17% of sites unique to RRBS. Overall, WGBS

predominantly profiles CpG sites in intergenic regions or introns, while the majority of CpG sites

sequenced by RRBS are in or near CpG islands, promoters, and exons.

Figure 5: The percent methylation per base of sample CG_WK_002 sequenced with **(A)** WGBS
and **(B)** RRBS after being read mapping with BWA meth.

Figure 6: Comparisons of CpG sites that were shared or unique across sequencing methods, with all metrics averaged across replicates from each population.**(A)** Proportion of shared and unique CpG sites that overlap with CpG islands and shores. **(B)** Total number of observed sites. **(C)** Proportion of sampled CpG islands and shores in the threespine stickleback reference genome. **(D)** Proportion of sites that overlap with promoters, exons, introns, and intergenic regions. **(E)** Total number of CpG sites used for functional annotations. **(F)** Proportion of sampled promoters, exons, and introns in the threespine stickleback reference genome.

Concordance between RRBS and WGBS methylation calls

To assess agreement in methylation calling between RRBS and WGBS, we merged datasets from technical replicates by genomic position and calculated differences in percent methylation at each base obtained with WGBS and RRBS. We found broad agreement on a nucleotide-level between library preparation methods (RRBS percent methylation ~ WGBS percent methylation: t = 2,467.4, p < 0.0001, R^2_{adj} = 0.735). Almost 60% of CpG sites had ≤10% difference in percent methylation, which is within the margin of error according to the methylKit documentation (Figure 7A). As read depth influences methylation call accuracy, we also assessed the difference in methylation as a function of depth at each CpG site for both WGBS

(Figure 7B) and RRBS (Figure 7C). We observed a significant but negligible relationship

between depth and disagreement in percent methylation ($R^2_{adj} < 0.01$).

Figure 7: Comparisons of per site methylation estimated from different library preparation
methods. **(A)** Difference in percent methylation per base obtained from RRBS and WGBS.
Negative values indicate CpG sites with higher methylation in RRBS than WGBS. The
relationship between the difference in percent methylation on a nucleotide level between library
preparation methods and the depth of **(B)** WGBS and **(C)** RRBS data.

Discussion:

The increasing popularity of RRBS for methylation profiling has coincided with a rise in

the use of Bismark as the default method for read mapping and methylation calling, particularly

in ecologically focused experiments (Table 1). However, little effort has been dedicated to

understanding the strengths and limitations of these methodological choices, particularly in

genetically variable, wild organisms. By comparing RRBS and Bismark against less commonly

used library preparation and analysis alternatives, we identify several options for researchers to

maximize their sequencing resources.

Optimizing bisulfite alignment software for ecological epigenetics

As predicted by previous studies (Farrell et al., 2021; Nunn et al., 2021; Zhou et al.,

2024), BWA mem-based read mappers outperformed and Bismark in their ability to map the

majority of sequenced reads to a reference genome (Figure 1). However, it is noteworthy that

this previous work focused on human and plant data. We could find no previous studies

assessing performance of bisulfite alignment wrappers across wild-caught/non-model systems.

Despite differences in mapping efficiency, BWA meth and Bismark provided consistent

estimates of the mean percent of CpG sites methylated per individual. However, there was a

clear disagreement between more recent read mapping software and older software, in which

Biscuit and BiSulfite Bolt estimated substantially higher methylation rates than BWA meth or

Bismark (Figure 2). When we explored this trend further, we found that it was driven by a

greater number of unmethylated cytosines captured by BWA meth and Bismark while Biscuit

and BiSulfite Bolt recovered more intermediately methylated cytosines (Figure 4). Disagreement

between read mappers could partially be alleviated through greater sequencing, but sequencing

effort alone was insufficient to correct for large differences in software output (Figure 3).

For projects focusing on metrics related to per-site methylation, a key goal is to maximize

the number of usable sequencing reads. Bismark resulted in the lowest mapping efficiency across

all organisms and both BS-seq library types analyzed here, yet it remains the most commonly

used read mapping method (Table 1). Continued reliance on Bismark may partly be attributed to

it being one of the first bioinformatic tools designed to handle bisulfite converted data. Because

Bismark does *in silico* conversions of all reads (sense and antisense, read and reference), its

reliability is also agnostic to library directionality. Although Bismark's ability to both map reads

and call methylation levels makes it particularly accessible for researchers new to epigenetics,

the tool's default parameters may cause difficulties when working with paired end reads, as we

observed when we initially had a 28% mapping efficiency for our stickleback RRBS data.

Although rarely discussed in the literature, low mapping efficiency of paired end reads appears

to be a relatively common issue with this tool (e.g., Nunn et al., 2021; Zhou et al., 2024). Indeed,

the developers provide thorough documentation to help users resolve this mapping problem. The

guidance highlights failure to trim adapters prior to read mapping, potential contaminants, and

high stringency under default parameters, among other possible sources of error. Under default

parameters, both paired end reads must map to one unique version of the *in silico* converted

reference genome, otherwise the paired end sequence gets discarded. Following the advice in the

documentation, and in-line with Nunn et al. (2021) and Zhou et al. (2024), we relaxed this

threshold by mapping our reads as non-directional. This resulted in a mapping rate (54%) that is

more in-line with previous DNAm studies in stickleback (Heckwolf et al., 2020; Hu et al., 2021).

An additional complication with Bismark is the decision of whether to use the default global

alignment method, in which sample reads must meet strict alignment requirements from

beginning to end, or allow soft clipping of read ends that do not align as well to the reference

genome as the rest of the insert (i.e. "local alignment"). The developers of Bismark argue that

higher mapping efficiency obtained from local alignment is due to erroneously mapped reads.

However, the other three BS-seq aligners tested here, including two that have been published

within the last five years, rely on local alignment. Indeed, Zhou et al. (2024) found that Bismark

not only had a lower mapping efficiency compared to other methods, but also resulted in the

lowest rate of reads optimally mapped to the genome. Additionally, Farrell et al. (2021) used

simulated data to determine Bismark had among the highest methylation call error rate,

particularly for genetically variable data. While we cannot confirm the accuracy of local

alignment because we used real BS-seq data, simply running Bismark as a local aligner resulted

in significantly higher mapping efficiency across all organisms and sequencing methods.

Despite requiring additional tools for methylation calling, BWA meth resulted in

significantly higher mapping efficiency than Bismark. While a previous study found that BWA

meth had higher error rates than alternative tools (Farrell et al., 2021), this analysis was based on

both directional and nondirectional libraries. It is plausible that the increased error rate was

driven by strand bias rather than being a software limitation. Following BWA meth, we used

MethylDackel to tabulate methylation and filter SNPs. Conventional wisdom is that RRBS

libraries should be sequenced using single end reads, as paired end reads may lead to bias by

repeatedly sequencing the same sites (Krueger et al., 2012). However, this bias can be reduced

by normalizing the data based on median read depth, and if genetic polymorphisms are unknown,

the overlapping nature of paired end reads can be exploited by software like MethylDackel to

call SNPs. Additionally, using MethylDackel for SNP filtering reduces errors associated with

SNPs falsely being considered unmethylated cytosines. In addition to facilitating SNP calling,

paired end sequencing ensures high mappability in regions with many genomic repeats, which

tend to be enriched in RRBS (Sun et al., 2018). Similar to most studies of ecological model

organisms, we did not have genotype data for either the focal individuals or populations, and

were thus unable to verify the accuracy of SNP filtering using MethylDackel. For a

comprehensive assessment of SNP filtering tools for BS-seq data, see Lindner et al. (2022),

although they do not include MethylDackel in their analysis.

We discovered that modern read mapping algorithms captured very few unmethylated

cytosines, while older wrappers captured few intermediately methylated sites. There are a

number of features inherent to RRBS that make it difficult to map to a reference genome.

Enzymatic shearing during library preparation targets CpG-dense regions of the genome while

bisulfite conversion increases the number of mismatches between the sample read and reference

genome. These processes cause RRBS reads to be repetitive with small insert sizes, consequently

making them particularly difficult to map accurately to a reference. It is possible that some

algorithms repeatedly align reads to the same regions of the genome, while other algorithms

spread the sample reads across the genome. Because we used real data, we are unable to

determine which read mappers resulted in the most accurate methylation profiles. However,

BWA meth and Bismark produced methylation profiles that are more analogous to the expected

bimodal distribution of percent methylation, with most sites being either unmethylated (≤10%

methylation) or fully methylated (≥90% methylated) (Akalin et al., 2012; Seiler Vellame et al.,

2021; Sun et al., 2018). As far as we are aware, this type of bias between these methods has not

been reported previously, highlighting how conventional methods optimized and tested only on

inbred model organisms and/or simulated data may not be suitable for genetically variable, wild

populations.

RRBS and WGBS vary in breadth of coverage and methylation calls

In addition to assessing how bisulfite alignment wrappers performed with RRBS data, we

examined how methylation profiles varied across technical and biological replicates using RRBS

and WGBS. Few CpG sites were commonly sequenced between technical (Supplemental Table

3) or biological replicates (Supplemental Table 4). When comparing the methylation patterns

recovered from technical replicates, the number of intermediately methylated sites (10-90%) was

much lower in RRBS compared to WGBS (Figure 5). However, RRBS enriches for functional

regions, while WGBS sequences across the entire genome (Figure 6). Although relatively few

sites were shared across datasets, those that were shared often had similar estimates of

methylation percentage (Figure 7).

One potential explanation for the lack of congruence between sites sequenced with RRBS

and WGBS is that our sequencing parameters (i.e., 2x150bp reads) were much longer than the

small inserts (~38bp) produced via RRBS. This resulted in only a fraction of the total expected

sequence data being useful for analyses. Importantly, increasing sequence effort led to greater

agreement in downstream methylation calling, regardless of read mapping methods. However,

this benefit plateaus at ~20 million reads per sample (Figure 3). A previous study used

simulations based on DNAm profiles from the cortex region of brains in an inbred mouse mouse

strain to determine that the relationship between depth of coverage and methylation call accuracy

increases rapidly until 10x depth of coverage, at which point the trend becomes asymptotic

(Seiler Vellame et al., 2021). Comparing our stickleback results with those of Seiler Vellame et

al. (2021) suggests that the optimal target coverage depth may vary among species, and also

likely depends on what level of precision is important to investigators.

Interestingly, the level of concordance between biological replicates depended on both

population and sequencing method. Individuals from Wik lake had over twice as many

overlapping CpG sites compared to samples from Watson lake when sequenced using RRBS

(Supplemental Table 4). Because this pattern was not present in WGBS, RRBS library

preparation is the likely culprit. Despite having a similar number of reads, Watson fish had fewer

CpG sites and a slightly higher read depth (14x vs 13x) than individuals from Wik. Taken

together, this suggests that fewer MspI cut sites (or a decrease in enzymatic cleavages) in Watson

fish led to less CpG sites following library preparation, resulting in the sites that were recovered

to be sequenced at a slightly higher depth. To test this prediction, we would need population-

specific genomes for *in silico* digestion and a larger sample size than n = 2/population. These

types of data limitations are common in studies of natural populations. Overall, these results

highlight the importance of assessing common methods across a diversity of taxa and

populations, as genetic variation can affect what data is available for downstream analysis.

When assessing methylation profiles of WGBS-RRBS technical replicates, the proportion

of intermediately methylated sites sequenced with WGBS was almost twice as high as

intermediate methylation captured with RRBS (56% versus 29%, respectively). Methylation can

degrade if tissue samples are collected multiple hours after mortality (Rhein et al., 2015) or

frozen for a long period of time (Lee et al., 2023), but the liver tissue we sequenced was stored in

RNALater immediately upon euthanasia and kept at -70□ for less than one year. Therefore,

degradation of intermediately methylated sites is unlikely the main driver of this observation.

Rather, there are technical and evolutionary explanations that likely interacted to create this

pattern. Batch 2 RRBS samples recovered a much higher proportion of intermediately

methylated sites than batch 1 (Supplemental Figure 7), likely driven by more sequencing.

However, this higher rate ($37.83 \pm 5.78\%$) is still substantially lower than the average of WGBS,

suggesting that there may be fundamental, biological differences in the methylation patterns

surveyed by each method. Indeed, while WGBS captured almost all of the CpG islands,

promoters, introns, and exons in the stickleback reference genome, over half of WGBS CpG sites

were found in introns or intergenic regions. Meanwhile, over half of the CpG sites unique to

RRBS were in promoters and exons. Given that molecular ecologists are likely interested in

identifying functionally important epigenetic effects, RRBS may provide a better option for

detecting and measuring these changes.

We found consensus when comparing methylation rates on a single base pair level

between technical replicates sequenced with RRBS and WGBS, confirming the validity of both

library preparation methods for assessing DNAm. This concordance would likely have been

higher if we used a more stringent SNP filter. Our SNP filtering cut off was rather lenient,

meaning that we likely called some sites as methylated cytosines when in reality they were due

to genetic variation. When RRBS and WGBS disagreed on methylation calls, it was equally

common for RRBS to call a site as being more or less methylated than WGBS, suggesting that

there is not a methodological bias. However, bisulfite conversion damages DNA at unmethylated

cytosines during library preparation, which inherently inflates genome-wide methylation levels

(Olova et al., 2018). The magnitude of this bias is only recently becoming clear thanks to newer

library preparation approaches that forego harsh chemical reactions and instead rely on enzymes

to convert unmethylated cytosines (e.g., EM-Seq: Vaisvila et al., 2021). If precise estimates of

genomewide methylation are of key concern, then consideration should be given to using these

newer, non-bisulfite methods of library preparation. Importantly, EM-seq converts unmethylated

cytosines to thymine, making the sequencing data compatible with existing bioinformatic tools

for bisulfite converted data. Despite these advances, RRBS remains the most cost effective

DNAm profiling method and is likely sufficient if the primary goal of a project is to measure or

detect relative changes in methylation (i.e., those likely to have functional effects or are

environmentally-induced). However, more studies should be done to assess agreement between

technical replicates sequenced with WGBS, RRBS, and EM-seq to validate these methods across

a variety of organisms.

Conclusions:

We take a critical approach to analyzing RRBS and WGBS data using several

conventional methods. We highlight the strengths and weaknesses of both molecular and

bioinformatic methods, all of which should be considered during experimental design. If SNP

filtering in the absence of genotype data is a priority, researchers may consider choosing paired

end sequencing, contrary to traditional advice regarding RRBS library construction. Given that

RRBS can result in exceedingly small insert sizes and there may be random variation in the

amount of sequencing effort for each sample, we suggest using relatively short read lengths and

deeply sequencing a small number of initial samples to determine the sequencing threshold

required to obtain accurate per site methylation metrics. In the case of threespine stickleback, 20

million reads/sample appears to be the optimal trade off between cost and accuracy.

Additionally, the most commonly used tool for analyzing BS-seq data (Bismark) performs

significantly worse than alternative methods in terms of percentage of reads mapped. However,

more work needs to be done to reconcile the difference between methylation profiles obtained

using older and newer read mapping software. Finally, we highlight how different library

preparation methods recover distinct methylation profiles, and why researchers may consider

alternative library preparation methods to BS-seq.

Acknowledgements:

We thank Sean Schoville for computing resources, and Kristofer Sasser and Cole Wolf for

collecting fish from Vancouver Island. We recognize that the fish populations analyzed here

originate from lakes in the unceded territory of the Dena'ina and Coast Salish people and thank

them for their continued land stewardship. This work was funded by the Department of

Integrative Biology at UW-Madison and National Institute of General Medical Sciences

5R35GM142891-02.

References:

Adrian-Kalchhauser, I., Sultan, S. E., Shama, L. N. S., Spence-Jones, H., Tiso, S., Keller
Valsecchi, C. I., & Weissing, F. J. (2020). Understanding "Non-genetic" Inheritance:
Insights from Molecular-Evolutionary Crosstalk. *Trends in Ecology and Evolution*,
35(12), 1078–1089. https://doi.org/10.1016/j.tree.2020.08.011
Akalin, A., Franke, V., Vlahoviček, K., Mason, C. E., & Schübeler, D. (2015). genomation: A
toolkit to summarize, annotate and visualize genomic intervals. *Bioinformatics*, *31*(7),
1127–1129. https://doi.org/10.1093/bioinformatics/btu775
Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., &
Mason, C. E. (2012). methylKit: A comprehensive R package for the analysis of genome-
wide DNA methylation profiles. *Genome Biology*, *13*(10), R87.
https://doi.org/10.1186/gb-2012-13-10-r87
Aphalo P (2024). ggpmisc: Miscellaneous Extensions to 'ggplot2'. R package version 0.6.0.9000,
https://github.com/aphalo/ggpmisc, https://docs.r4photobiology.info/ggpmisc/.
Banta, J.A., & Richards, C.L. (2018). Quantitative epigenetics and evolution. Heredity *121*, 210–
224. https://doi.org/10.1038/s41437-018-0114-x
Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P., & Marth, G. T. (2011).
BamTools: A C++ API and toolkit for analyzing and managing BAM files.
Bioinformatics, *27*(12), 1691–1692. https://doi.org/10.1093/bioinformatics/btr174
Bock, C., Tomazou, E. M., Brinkman, A. B., Müller, F., Simmer, F., Gu, H., Jäger, N., Gnirke,
A., Stunnenberg, H. G., & Meissner, A. (2010). Quantitative comparison of genome-wide
DNA methylation mapping technologies. *Nature Biotechnology*, *28*(10), 1106–1114.
https://doi.org/10.1038/nbt.1681
Bogan, S. N., Strader, M. E., & Hofmann, G. E. (2023). Associations between DNA methylation
and gene regulation depend on chromatin accessibility during transgenerational plasticity.
BMC Biology, *21*(1), 149. https://doi.org/10.1186/s12915-023-01645-8
Brennan, R. S., deMayo, J. A., Finiguerra, M., Baumann, H., Dam, H. G., & Pespeni, M. H.
(2025). Complementary genetic and epigenetic changes facilitate rapid adaptation to
multiple global change stressors. *Proceedings of the National Academy of Sciences*,
122(29), e2422782122. https://doi.org/doi/10.1073/pnas.2422782122

Charlesworth, D., Barton, N. H., & Charlesworth, B. (2017). The sources of adaptive variation.
Proceedings of the Royal Society B: Biological Sciences, *284*(1855), 20162864.
https://doi.org/10.1098/rspb.2016.2864
Cholewa-Waclaw, J., Shah, R., Webb, S., Chhatbar, K., Ramsahoye, B., Pusch, O., Yu, M.,
Greulich, P., Waclaw, B., & Bird, A. P. (2019). Quantitative modelling predicts the
impact of DNA methylation on RNA polymerase II traffic. *Proceedings of the National
Academy of Sciences*, *116*(30), 14995–15000. https://doi.org/10.1073/pnas.1903549116
Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,
Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools
and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008
de Carvalho, C. F., Slate, J., Villoutreix, R., Soria☐Carrasco, V., Riesch, R., Feder, J. L.,
Gompert, Z., & Nosil, P. (2023). DNA methylation differences between stick insect
ecotypes. *Molecular Ecology, 32*(24), 6809-6823. https://doi.org/10.1111/mec.17165
Elliott, G., Hong, C., Xing, X., Zhou, X., Li, D., Coarfa, C., Bell, R. J. A., Maire, C. L., Ligon,
K. L., Sigaroudinia, M., Gascard, P., Tlsty, T. D., Harris, R. A., Schalkwyk, L. C.,
Bilenky, M., Mill, J., Farnham, P. J., Kellis, M., Marra, M. A., Milosavljevic, A., Hirst,
M., Stormo, G. D., Wang, T., & Costello, J. F. (2015). Intermediate DNA methylation is
a conserved signature of genome regulation. *Nature Communications*, *6*(1), 6363.
https://doi.org/10.1038/ncomms7363
Farrell, C., Thompson, M., Tosevska, A., Oyetunde, A., & Pellegrini, M. (2021). BiSulfite Bolt:
A bisulfite sequencing analysis platform. *GigaScience*, *10*(5).
https://doi.org/10.1093/gigascience/giab033
Gu, H., Smith, Z. D., Bock, C., Boyle, P., Gnirke, A., & Meissner, A. (2011). Preparation of
reduced representation bisulfite sequencing libraries for genome-scale DNA methylation
profiling. *Nature Protocols*, *6*(4), 468–481. https://doi.org/10.1038/nprot.2010.190
Guerrero, L., & Bay, R. (2024). Patterns of methylation and transcriptional plasticity during
thermal acclimation in a reef☐building coral. *Evolutionary Applications, 17*(7), e13757.
https://doi.org/10.1111/eva.13757
Gumucio, J. J., & Miller, D. L. (1981). Functional implications of liver cell heterogeneity.
Gastroenterology, *80*(2), 393–403. https://doi.org/10.1016/0016-5085(81)90732-0
Harrath, Y., Mahjoub, A., AbuBakr, F., & Azhar, M. (2019). Comparative Evaluation of Short
Read Alignment Tools for next Generation DNA Sequencing. *2019 International
Conference on Innovation and Intelligence for Informatics, Computing, and Technologies
(3ICT)*, 1–6. https://doi.org/10.1109/3ICT.2019.8910272
Hay, A. D., Kessler, N. J., Gebert, D., Takahashi, N., Tavares, H., Teixeira, F. K., & Ferguson-
Smith, A. C. (2023). Epigenetic inheritance is unfaithful at intermediately methylated
CpG sites. *Nature Communications*, *14*(1), 5336. https://doi.org/10.1038/s41467-023-
40845-2
Heckwolf, M. J., Meyer, B. S., Häsler, R., Höppner, M. P., Eizaguirre, C., & Reusch, T. B. H.
(2020). Two different epigenetic information channels in wild three-spined sticklebacks

are involved in salinity adaptation. *Science Advances*, *6*(12), eaaz1138. https://doi.org/10.1126/sciadv.aaz1138

Hothorn T., Hornik K., van de Wiel M. A., & Zeileis, A. (2008). Implementing a class of permutation tests: The coin package. *Journal of Statistical Software, 28*(8), 1–23. https://doi.org/10.18637/jss.v028.i08.

Hu, J., & Barrett, R. D. H. (2017). Epigenetics in natural animal populations. *Journal of Evolutionary Biology*, *30*(9), 1612–1632. https://doi.org/10.1111/jeb.13130

Hu, J., Wuitchik, S. J. S., Barry, T. N., Jamniczky, H. A., Rogers, S. M., & Barrett, R. D. H. (2021). Heritability of DNA methylation in threespine stickleback (*Gasterosteus aculeatus*). *Genetics*, *217*(1). https://doi.org/10.1093/GENETICS/IYAB001

Jaenisch, R., & Bird, A. (2003). Epigenetic regulation of gene expression: How the genome integrates intrinsic and environmental signals. *Nature Genetics*, *33*(3S), 245–254. https://doi.org/10.1038/ng1089

Kassambara, A. ((2023). rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package version 0.7.2. https://rpkgs.datanovia.com/rstatix/.

Klughammer, J., Romanovskaia, D., Nemc, A., Posautz, A., Seid, C. A., Schuster, L. C., Keinath, M. C., Lugo Ramos, J. S., Kosack, L., Evankow, A., Printz, D., Kirchberger, S., Ergüner, B., Datlinger, P., Fortelny, N., Schmidl, C., Farlik, M., Skjærven, K., Bergthaler, A., Liedvogel, M., Thaller, D., Burger, P. A., Hermann, M., Distel, M., Distel, D. L., Kübber-Heiss, A., & Bock, C. (2023). Comparative analysis of genome-scale, base-resolution DNA methylation profiles across 580 animal species. *Nature Communications*, *14*(1), 232. https://doi.org/10.1038/s41467-022-34828-y

Krueger, F., & Andrews, S. R. (2011). Bismark: A flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics*, *27*(11), 1571–1572. https://doi.org/10.1093/bioinformatics/btr167

Krueger, F., Kreck, B., Franke, A., & Andrews, S. R. (2012). DNA methylome analysis using short bisulfite sequencing data. *Nature Methods*, *9*(2), 145–151. https://doi.org/10.1038/nmeth.1828

Laine, V. N., Sepers, B., Lindner, M., Gawehns, F., Ruuskanen, S., & van Oers, K. (2022). An ecologist's guide for studying DNA methylation variation in wild vertebrates. *Molecular Ecology Resources*, 1755-0998.13624. https://doi.org/10.1111/1755-0998.13624

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan, M. T., & Carey, V. J. (2013). Software for Computing and Annotating Genomic Ranges. *PLoS Computational Biology*, *9*(8), e1003118. https://doi.org/10.1371/journal.pcbi.1003118

Lea, A. J., Tung, J., Archie, E. A., & Alberts, S. C. (2018). Developmental plasticity: Bridging research in evolution and human health. *Evolution, Medicine, and Public Health, 2017*(1), 162-175. https://doi.org/10.1093/emph/eox019

Lee, N. Y., Hum, M., Tan, G. P., Seah, A. C., Kin, P. T., Tan, N. C., Law, H.-Y., & Lee, A. S. G. (2023). Degradation of methylation signals in cryopreserved DNA. *Clinical Epigenetics*, *15*(1), 147. https://doi.org/10.1186/s13148-023-01565-y

Lindner, M., Gawehns, F., Te Molder, S., Visser, M. E., Van Oers, K., & Laine, V. N. (2022). Performance of methods to detect genetic variants from bisulphite sequencing data in a non☐model species. *Molecular Ecology Resources*, *22*(2), 834–846. https://doi.org/10.1111/1755-0998.13493

Marin, P., Genitoni, J., Barloy, D., Maury, S., Gibert, P., Ghalambor, C. K., & Vieira, C. (2020). Biological invasion: The influence of the hidden side of the (epi)genome. *Functional Ecology*, *34*(2), 385–400. https://doi.org/10.1111/1365-2435.13317

Nath, S., Shaw, D. E., & White, M. A. (2021). Improved contiguity of the threespine stickleback genome using long-read sequencing. *G3 Genes|Genomes|Genetics, 11*(2), jkab007. https://doi.org/10.1093/g3journal/jkab007

Nell, L. (2020). lucasnell/TaJoCGI: Initial release (v1.0.0). *Zenodo*. https://doi.org/10.5281/zenodo.3731298

Nunn, A., Otto, C., Stadler, P. F., & Langenberger, D. (2021). Comprehensive benchmarking of software for mapping whole genome bisulfite data: From read alignment to DNA methylation analysis. *Briefings in Bioinformatics*, *22*(5), bbab021. https://doi.org/10.1093/bib/bbab021

Oberacker, P., Stepper, P., Bond, D. M., Höhn, S., Focken, J., Meyer, V., Schelle, L., Sugrue, V. J., Jeunen, G. J., Moser, T., Hore, S. R., von Meyenn, F., Hipp, K., Hore, T. A., & Jurkowski, T. P. (2019). Bio-On-Magnetic-Beads (BOMB): Open platform for high-throughput nucleic acid extraction and manipulation. *PLoS Biology*, *17*(1). https://doi.org/10.1371/journal.pbio.3000107

Olova, N., Krueger, F., Andrews, S., Oxley, D., Berrens, R. V., Branco, M. R., & Reik, W. (2018). Comparison of whole-genome bisulfite sequencing library preparation strategies identifies sources of biases affecting DNA methylation data. *Genome Biology*, *19*(1). https://doi.org/10.1186/s13059-018-1408-2

Ord, J., Gossmann, T. I., & Adrian-Kalchhauser, I. (2023). High Nucleotide Diversity Accompanies Differential DNA Methylation in Naturally Diverging Populations. *Molecular Biology and Evolution*. https://doi.org/10.1093/molbev/msad068

Otterdijk, S. D., & Michels, K. B. (2016). Transgenerational epigenetic inheritance in mammals: How good is the evidence? *The FASEB Journal*, *30*(7), 2457–2465. https://doi.org/10.1096/fj.201500083

Pedersen, B. S., Eyring, K., De, S., Yang, I. V., & Schwartz, D. A. (2014). *Fast and accurate alignment of long bisulfite-seq reads* (arXiv:1401.1129). arXiv. http://arxiv.org/abs/1401.1129

R Core Team (2023). R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria*. https://www.R-project.org/.

794   Rey, O., Eizaguirre, C., Angers, B., Baltazar☐Soares, M., Sagonas, K., Prunier, J. G., &
795   Blanchet,   S. (2020). Linking epigenetics and biological conservation: Towards a
796   *conservation   epigenetics* perspective. *Functional Ecology*, *34*(2), 414–427.
797   https://doi.org/10.1111/1365-2435.13429
798   Rhein, M., Hagemeier, L., Klintschar, M., Muschler, M., Bleich, S., & Frieling, H. (2015). DNA
799   methylation results depend on DNA integrity—Role of post mortem interval. *Frontiers in*
800   *Genetics*, *6*. https://doi.org/10.3389/fgene.2015.00182
801   Seiler Vellame, D., Castanho, I., Dahir, A., Mill, J., & Hannon, E. (2021). Characterizing the
802   properties of bisulfite sequencing data: Maximizing power and sensitivity to identify
803   between-group differences in DNA methylation. *BMC Genomics*, *22*(1), 446.
804   https://doi.org/10.1186/s12864-021-07721-z
805   Sun, X., Han, Y., Zhou, L., Chen, E., Lu, B., Liu, Y., Pan, X., Cowley, A. W., Liang, M., Wu,
806   Q., Lu, Y., & Liu, P. (2018). A comprehensive evaluation of alignment software for
807   reduced representation bisulfite sequencing data. *Bioinformatics*, *34*(16), 2715–2723.
808   https://doi.org/10.1093/bioinformatics/bty174
809   Suzuki, M. M., & Bird, A. (2008). DNA methylation landscapes: Provocative insights from
810   epigenomics. *Nature Reviews Genetics*, *9*(6), 465–476. https://doi.org/10.1038/nrg2341
811   Takahashi, Y., Morales Valencia, M., Yu, Y., Ouchi, Y., Takahashi, K., Shokhirev, M. N.,
812   Lande, K., Williams, A. E., Fresia, C., Kurita, M., Hishida, T., Shojima, K., Hatanaka, F.,
813   Nuñez-Delicado, E., Esteban, C. R., & Izpisua Belmonte, J. C. (2023). Transgenerational
814   inheritance of acquired epigenetic signatures at CpG islands in mice. *Cell*, *186*(4), 715-
815   731.e19. https://doi.org/10.1016/j.cell.2022.12.047
816   Takai, D., & Jones, P. A. (2002). Comprehensive analysis of CpG islands in human
817   chromosomes 21 and 22. *Proceedings of the National Academy of Sciences, 99*(6), 3740-
818   3745. https://doi.org/10.1073/pnas.052410099
819   Tobi, E. W., Slieker, R. C., Luijk, R., Dekkers, K. F., Stein, A. D., Xu, K. M., Biobank-based
820   Integrative Omics Studies Consortium, Slagboom, P. E., van Zwet, E. W., Lumey, L. H.,
821   & Heijmans, B. T. (2018). DNA methylation as a mediator of the association between
822   prenatal adversity and risk factors for metabolic disease in adulthood. *Science Advances*,
823   *4*(1), eaao4364. https://doi.org/10.1126/sciadv.aao4364
824   Vaisvila, R., Ponnaluri, V. K. C., Sun, Z., Langhorst, B. W., Saleh, L., Guan, S., Dai, N.,
825   Campbell, M. A., Sexton, B. S., Marks, K., Samaranayake, M., Samuelson, J. C., Church,
826   H. E., Tamanaha, E., Corrêa, I. R., Pradhan, S., Dimalanta, E. T., Evans, T. C., Williams,
827   L., & Davis, T. B. (2021). Enzymatic methyl sequencing detects DNA methylation at
828   single-base resolution from picograms of DNA. *Genome Research*, *31*(7), 1280–1289.
829   https://doi.org/10.1101/gr.266551.120
830   Vernaz, G., Malinsky, M., Svardal, H., Du, M., Tyers, A. M., Santos, M. E., Durbin, R., Genner,
831   M. J., Turner, G. F., & Miska, E. A. (2021). Mapping epigenetic divergence in the
832   massive radiation of Lake Malawi cichlid fishes. *Nature Communications, 12*(5870).
833   https://doi.org/10.1038/s41467-021-26166-2

Viitaniemi, H. M., Verhagen, I., Visser, M. E., Honkela, A., van Oers, K., & Husby, A. (2019).
Seasonal Variation in Genome-Wide DNA Methylation Patterns and the Onset of
Seasonal Timing of Reproduction in Great Tits. *Genome Biology and Evolution, 11*(3),
970-983. https://doi.org/10.1093/gbe/evz044

Wang, Q., Yu, G., Ming, X., Xia, W., Xu, X., Zhang, Y., Zhang, W., Li, Y., Huang, C., Xie, H.,
Zhu, B., & Xie, W. (2020). Imprecise DNMT1 activity coupled with neighbor-guided
correction enables robust yet flexible epigenetic inheritance. *Nature Genetics*, *52*(8),
828–839. https://doi.org/10.1038/s41588-020-0661-y

Zemach, A., McDaniel, I. E., Silva, P., & Zilberman, D. (2010). Genome-Wide Evolutionary
Analysis of Eukaryotic DNA Methylation. *Science*, *328*(5980), 916–919.
https://doi.org/10.1126/science.1186366

Zhou, W., Johnson, B. K., Morrison, J., Beddows, I., Eapen, J., Katsman, E., Semwal, A., Habib,
W. A., Heo, L., Laird, P. W., Berman, B. P., Triche, T. J., & Shen, H. (2024). BISCUIT:
An efficient, standards-compliant tool suite for simultaneous genetic and epigenetic
inference in bulk and single-cell studies. *Nucleic Acids Research*, *52*(6), e32–e32.
https://doi.org/10.1093/nar/gkae097

Data Accessibility and Benefit-Sharing:

Data Accessibility Statement: All scripts are available at

https://github.com/EmilyKerns/VariablePerformanceofBS-SeqandReadMappingSoftware.

Stickleback sequence data is available on Dryad at https://doi.org/10.5061/dryad.dncjsxm9h and

NCBI SRA (BioProject Accession Number PRJNA1439918).

Author Contributions:

E.V.K. and J.N.W. conceived and designed the study and co-wrote the manuscript; E.V.K. wrote

the scripts, conducted analyses, and produced figures; J.N.W. supervised and funded the work.