Genomic resources for curimba *Prochilodus lineatus* reveal a fractal pattern of hierarchical genetic population structure in mixed stocks across fish segregated by dams

Gabriel M. Yazbeck [1,2,3], **Gésica Aparecida Santana Nascimento** [1], **Leiliane Campos Carvalho** [2], **Raíssa Cristina Dias Graciano** [2], **Rosiane de Paula Santos** [3], **and Rafael Sachetto Oliveira** [4]

[1] Programa de Pós Graduação em Ecologia, Universidade Federal de São João Del Rei, São João Del Rei, MG, Brazil

[2] Programa de Pós Graduação em Biotecnologia, Universidade Federal de São João Del Rei, Divinópolis, MG, Brazil.

[3] Laboratório de Recursos Genéticos (LARGE), DEZOO, Universidade Federal de São João Del Rei, São João Del Rei, MG, Brazil.

[4] Departamento de Ciência da Computação, Universidade Federal de São João Del Rei, São João Del Rei, MG, Brazil.

Genomic resources, new microsatellite markers and a novel observation of a fractal pattern of the genetic population structure are presented for curimba *Prochilodus lineatus*, a suitable model organism for freshwater migratory species in South America. Our main goals were to investigate the presence of mixed fish stocks, the effects of isolation by dams and broodstocking over the genetic diversity for this species. We focused on the upper segment of the extremely fragmented Grande River, MG, Brazil, from the megadiverse continental La Plata Basin. Two main co-occurring genetically admixed stocks were identified by maximum likelihood estimation and iteratively unfolded with two further steps, through a hierarchical analysis of genetic structure. It consistently unveiled two newly defined internal groups, from the previously resolved clusters (2^3). We suggest this stratified pattern of mixed stocks from the main river channel may be attributed to the existence of different, well defined breeding grounds, nested within differential scales throughout this basin, arranged according to fractal geometry (1^{st} order river, 2^{nd} and 3^{rd} order tributaries, and so on). This would imply in homing behaviour for this species. Our work also provides massive molecular data for this fish, including its first draft genome assembly and a valuable panel for the rapid and low-cost development of new microsatellite DNA markers. These findings will likely benefit researchers and environmental managers in tackling challenging issues regarding the conservation of Neotropical migratory fishes, as they support the hypothesis of the isolating effects of dams and broodstocking over genetic variability. More importantly, the results presented here point to the potential role of tributaries of different orders in harbouring genetically diverse populations of migratory freshwater fishes.

Fractals | Microsatellites | Population genetics | Sequencing-by-synthesis | Upper Grande River MG

Correspondence: *dna@ufsj.edu.br*

Introduction

The curimba *Prochilodus lineatus* Valenciennes 1836 (Characiformes: Prochilodontidae), also known as sabálo, grumatá and other common names, is one of the most important freshwater migratory species for inland fisheries in South America. It corresponds to >50% of the fish biomass from the La Plata Basin (Bowen, 2022a,b), a megadiverse hydrological system (Cassemiro *et al.*, 2023), reaching Argentine, Bolivia, Brazil, Paraguay and Uruguay. This major Neotropical basin is heavily impacted by anthropogenic activity (Reis *et al.*, 2016). *Prochilodus lineatus* also naturally occurs in the Paraíba do Sul Basin (Brazil), and has been recorded as an invasive species in China (Endruweit, 2014) and Vietnam (Kalous *et al.*, 2012).

Prochilodus lineatus has a conspicuous specialized inverted funnel mouth, an adaptation from its detritivorous habits (Bowen, 2022b) and, due to its abundance and popularity, it is an emblematic example of the Neotropical potamodromous (*i.e.* freshwater migratory) life history. Its reproductive migrations are triggered by environmental cues and its fish runs are regionally denominated piracema, a term used to refer to both, the migratory process, and the fish species which partake in it. It typically occurs from around the early summer, until around the season's end (Carolsfeld *et al.*, 2003). Thus, this process is potentially sensible to ongoing climate change (Alix *et al.*, 2020).

Piracema fishes naturally spawn upon reaching upstream breeding grounds (Ribolli *et al.*, 2020). Larvae and juveniles then passively drift downstream and reach nursery and feeding floodplains, where they grow and mature (Petry *et al.*, 2003). Adults spawn several consecutive years during its live cycle (Carolsfeld *et al.*, 2003).

Species from the *Prochilodus* genus were among the first Neotropical migratory fishes to have its *ex situ* spawning mastered. This was motivated by the intensive exploitation of hydroelectricity generation and its impact over migratory species, aiming broodstock release operations, intended to counteract dwindling fisheries around regions affected by dams (Saraiva and Pompeu, 2016).

These stocking practices have been largely abandoned in some relevant places for hydroelectric production (*e.g.* the Brazilian state of Minas Gerais), during the last decade, in part because of general relaxation of environmental protection oriented regulations (Zellhuber, 2016), although a rigorous, holistic (*i.e.* ecological, genetic and anthropological) assessment of the activity's potential benefits or pitfalls is still lacking. Given its status as the main targeted fish for broodstocking, in the last 50 years or so, *P. lineatus* is a natural candidate as a model species for gauging the impacts of dams and the effectiveness of possible mitigation measures, regarding piracema fishes.

Molecular markers have been applied to access the influence of dams over Neotropical migratory fishes showing, for example, the existence of different stocks across fragmented basins (*e.g.* Oliveira-Farias *et al.*, 2022), or the lack of differentiation following recent isolation (*e.g.* García-Castro and Márquez, 2024). It was also used to show the absence of regional genetic structure (*e.g.* Sivasundar *et al.*, 2001), or to approach fish stocks delimitation (Collins *et al.*, 2013). DNA markers were also used to reveal life history traits of silurid catfish species, showing evidence for the occurrence of homing breeding behaviour (*i.e.* philopatric spawning), in the Amazon (Batista and Alves-Gomes, 2006; Carvajal-Vallejos *et al.*, 2014) and the La Plata Basin (Pereira *et al.*, 2009). Other questions worth a deeper assessment in South America include the efficiency of transposing mechanisms and practices for sustaining gene flow (*e.g.* Wilkes *et al.*, 2019); if broodstocking are inducing genetic effects (*e.g.* bottlenecks, domestication) from hatchery spawning (*e.g.* Howe *et al.*, 2024); if released fish lead to stocks introgression (*e.g.* Chang *et al.*, 2022), or jeopardize local adaptations (*e.g.* Shedd *et al.*, 2022), and so on.

Piracema fishes under the impacts of hydroelectric power production, thus, may benefit from objective genetic information raised through molecular markers such as microsatellites, which are amenable to simple genotyping routines for hatchery or wild fish monitoring operations (Wenne, 2023). The popularization of next-generation sequencing (NGS) techniques have made the characterization of new DNA markers increasingly accessible and straightforward (Yazbeck *et al.*, 2024).

Since the seminal work of Smouse *et al.* (1990), scientists and managers have applied the statistical inference of genetic structure (the form in which genetic polymorphisms are distributed), to characterize mixed stocks from fish samples, through DNA markers (Hallerman, 2003; Valenzuela-Quiñonez, 2016). Migratory runs potentially place individuals from different, well defined, random breeding populations (demes) together in the same local mixed fish stock, along commonly shared migratory routes (Collins *et al.*, 2013). Mixed fish stocks can follow a genetically mixed population model, characterized by each individual's genome originating from a single ancestral population source (genetic cluster), out of a K number of alternatives, where these demes do not share breeding grounds and events. Alternatively, mixed fish stocks might be portrayed by an admixture model. It considers different individual proportions of genomic heritage, out of several K ancestral genetic clusters, where previous generations have potentially engaged in common reproductive events (Pritchard *et al.*, 2000; Wang, 2022). Admixture, thus, provide a convenient framework for the problem of unravelling the genetic structure of piracema fishes, under the impact of dams and decades of stocking operations, largely conducted without planned considerations towards genetic issues.

Genetic investigations applying microsatellites and admixture models in *P. lineatus* were first presented by Rueda

et al. (2013), unveiling distinct seasonal segregation of stocks from the Lower Uruguay River. They have concluded this species should be deemed as exhibiting mixed stocks, for fisheries management purposes. Perini *et al.* (2021) described a genetic source-sink metapopulation dynamics involving the Pardo River and the Lower Grande River (Upper Paraná Basin). Both, Ferreira *et al.* (2017) and Rosa *et al.* (2022) also combined mitochondrial sequences along microsatellites. The former studied fish from the Middle and Upper Paraná River Basin and showed no evidence for homing in this species. The latter revealed temporal genetic structuring of a location of the Mogi-Guaçu River (Pardo River Basin). Ferreira *et al.* (2023) also showed temporal structuring and mixed stocks of *P. lineatus* from the Paranapanema River Basin, a major tributary of the Upper Paraná River.

In general, all these studies relate to lower regions of the same intensely dam fragmented system and discussed the importance of free-flowing tributaries for harbouring spawning groups and events in *P. lineatus*. All showed high levels of genetic variability and frequently reported on the absence of a single panmitic population associated with distinct localities. Instead, these studies point to local or temporal genetic substructuring and tend to resolve two or three genetic clusters, according to Bayesian admixture analysis. To date, except for observations provided by Yazbeck and Kalapothakis (2007), there are no genetic population structure analysis for *P. lineatus* from the Upper Grande River, the final and highest region from the Upper Paraná Basin and, therefore, the uppermost terminal region from the vast Upper La Plata River Basin's oriental flank.

The Grande River springs from an altitude of almost 2,000 m in Minas Gerais and runs through 1,360 km until its confluence with the Paranaíba River, where both coalesce onto the Paraná River (a major component of the La Plata Basin). It is an illustrative example of thoroughly impounded river, with 12 large dams, with the Camargos hydroelectric power plant being the last dam upbound, implemented around 1960. It is closely followed downstream by another dam, from the Itutinga power plant, operational since 1955, which delimits a singular, relatively small closed impoundment (without tributaries). There, an environmental hatchery used to actively operate for decades in broodstock production and release, mainly of *P. lineatus*, throughout the region, until 2018. It provides a singularly convenient stage for testing for the genetic effects of broodstoking. Next, further downstream, lies the relatively new Funil power plant dam (constructed circa the year 2000), the only of these dams accommodating a fish transposition mechanism, a lift (Suzuki *et al.*, 2011). These three dams are featured in Figure 1. The degree in which these large dams genetically isolate *P. lineatus* in the Upper Grande River is not currently known, nor are the genetic effects from broodstock release and fish transposition operations there.

Only a handful of genomic resources are currently available for prochilodontid fish species. The mitochondrial DNA sequence for *P. lineatus* was unveiled by Carmo *et al.* (2016), while other *Prochilodus* species' mitogenomes were presented by Chagas *et al.* (2016) and Santos *et al.* (2021). Regarding the nuclear genome, Stornioli *et al.* (2021) have disclosed NGS data for *P. lineatus* and Yepes-Blandón *et al.* (2022), in turn, have delivered the first draft of a full reference genome for *Prochilodus magdalenae*, a Colombian species, although not publicly releasing NGS data. Our core objectives in this work were *i)* testing the hypothesis of the occurrence of mixed stocks across *P. lineatus* populations segregated by the last three upbound dams from the Upper Grande River, MG, Brazil; *ii)* testing the hypothesis that dams are genetically isolating migratory fish in the region and; *iii)* testing the hypothesis that broodstocking practices have altered genetic diversity in populations of fish trapped between dams. We aimed at doing so through the development of a genome-wide set of microsatellite *loci*, for rapid marker development in *P. lineatus*, while making available new genomic resources for this fish, a model for Neotropical freshwater migratory species in general.

Figure 1. Geographical location of the studied population samples for curimba *Prochilodus lineatus* from the Upper Grande River Basin, MG, Brazil. Photography by André Seale.

Material and Methods

Ethics statement

All proceedings presented here were conducted under the guidance and licenses 27/2011 and 6991130223 of the Committee on Ethics in Animal Use (CEUA-UFSJ); along CGEN licences A9D0E5 and AC5A784; and SISBIO licenses 37222-2 and 85497-1. No experimental assays were conducted on live specimens.

Genomics, bioinformatics and microsatellite markers development

Genomic libraries of whole genome shotgun fragments (500 bp) were produced for NGS, out of a single adult female individual from the Itutinga Hatchery Station (21°19'20.7"S 44°36'56.9"W - Figure 1. The methodological steps and parameters for NGS and initial microsatellite characterization were the same as presented in Yazbeck *et al.* (2018), and carried out by an external service provider (SP), BGI (Tai Po, Hong Kong). The resulting 90 bases paired-end (PE) short-reads were organized as three sets of twin FASTQ files. We verified the quality of DNA short-reads using MultiQC (Ewels *et al.*, 2016), and computed all of its 21-mers (k-mer substrings) in Jellyfish (Marçais and Kingsford, 2011). These latter results were applied in the estimation of the expected genome size of *P. lineatus* through a frequency histogram, in R version 4.4.1 (R Core Team, 2024).

We used all the DNA short-reads, along a set of fasta format contigs containing microsatellites resolved by the SP upon the original undisclosed assembly, to execute a *de novo* genomic assembly, ourselves. For this step, we used MEGAHIT (Li *et al.*, 2015), using k-mer=47, to generate new contigs and, then, submitting these to the SOAPdenovo-fusion module from SOAPdenovo 2 (Luo *et al.*, 2012), to resolve scaffolds. This genome assembly was processed in the Sagarana high-performance computing cluster (CEPAD-ICB-UFMG, Belo Horizonte, Brazil). The National Center for Biotechnology Information's (Sayers *et al.*, 2022) contamination screening tool FCS (Astashyn *et al.*, 2024) was used by NCBI staff and ourselves for filtering out mitochondrial assembly, sequences from microorganisms and library adaptors found in the final scaffold level sequence. Assembly statistics were accessed through assemblyStats (http://gif.biotech.iastate.edu) and its visualization by assembly-stats 17.02 (Challis, 2017).

We mapped the microsatellites characterized in the steps above in the new genomic assembly with the strategy also delineated in Yazbeck *et al.* (2018). A sequence alignment map (SAM) was then produced with BBmap (Bushnell, 2014), by mapping short-reads unto the genomic assembly, and finally converted to a binary format (BAM) with SAMtools (Li *et al.*, 2009).

For this work we developed new species-specific microsatellite markers to be applied in the population genetics analysis. We used the results of the genomic microsatellite *loci* search by the SP for choosing candidate *loci* for empirical tests. The microsatellites were sorted by decreasing motif size and number of repeats, in order to maximize chances of finding polymorphic *loci* (Brandström and Ellegren, 2008). A total of 22 candidates primer pairs were picked (12 from the top sorted candidates and 10 at random) and then synthesized by an SP (Integration DNA Technologies, Coralville, USA). These were applied to wet bench assays in four individual DNA samples. Polymerase chain reaction (PCR) was carried in a final volume of 10 μl, with initial conditions of 1 μl of unquantified chelex extracted genomic DNA to seed the reaction; 10 pmol of each primer; $MgCl_2$ (2 mM); Tris-HCl (10 mM); KCl (50 mM); and 1 unity of Taq DNA polymerase enzyme (Phoneutria, Belo Horizonte, Brazil). Thermal PCR cycles were as following: initial genomic DNA denaturation at 94ºC for 4 min.; 30 iterated cycles of denaturation at 94ºC for 30 s, primers annealing temperature (specific for each primer pair) for 30 s, and amplicon elongation step at 72ºC for 30 s; followed by a final elongation step, at the same temperature, for 5 min.

The resulting PCR products were visualized by polyacrylamide gel electrophoresis at 10%, at 15 V/cm, loaded along with 25 bp DNA fragment size ladder, stained with ethidium bromide. *loci* yielding polymorphic patterns were then assayed in a 20 individuals sample, for optimization of PCR conditions. Gels were photographed under UVB light, and scored with GelAnalyzer. Scored DNA fragments were then converted to nominal allele classes, through a in house semi-automated (*i.e.* automated, under human supervision) binning approach, based on frequency histograms and formatted for downstream analysis.

Population genetics

A total of 100 *P. lineatus* individual tissue samples (ethanol preserved fin fragments) were used, from the DNA bank of the Laboratório de Recursos Genéticos (LARGE-UFSJ). These were collected in 2013 from three locations (Figure 1) from the Upper the Grande River, MG: downstream the Funil dam, upstream the Itutinga dam and upstream the Camargos dam (see Suzuki *et al.* 2011 for a more detailed characterization from these locations). An additional sample of 40 individuals was obtained from the Itutinga dam reservoir in 2023 (Table 1).

Following DNA marker development experiments, a total of 15 new empirically validated microsatellite *loci* were applied to 140 individuals of *P. lineatus*. Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) were tested through the methods implemented in GenePop 007 (Rousset, 2008), for each population sample, with default

Table 1. Evaluated *Prochilodus lineatus* individuals sampled, from the last three dams from the Upper Grande River, MG, Brazil.

Sample	ID	Location	Year	N
Camargos	1-20	21°20'27.1"S 44°36'34.3"W	2013	20
Itutinga 1	21-60	21°17'39.9"S 44°37'24.1"W	2013	40
Itutinga 2	61-100	21°17'30.4"S 44°37'11.4"W	2023	40
Funil	101-140	21°08'39.4"S 45°02'19.1"W	2013	40
Total				**140**

ID=Individual label; N=Sample size.

options for Markov chain Monte Carlo (MCMC) parameters. These were analysed under new significance threshold values for multiple tests, following sequential Bonferroni corrections. Microchecker (van Oosterhout *et al.*, 2004) was used for controlling scoring errors (stuttering, large allele dropout and null alleles). Genetic data handling, conversions and table visualizations were mainly conducted with the adegenet package for R (Jombart, 2008). Other genetic diversity and structure estimates for the population samples were obtained through GenAlex (Peakall and Smouse, 2012), including expected and observed heterozygosities (H_E and H_O, respectively); allelic richness, *i.e.* number of alleles (N_A); Jost's D; the fixation index $G"_{ST}$; and analysis of molecular variance (AMOVA), each with 999 random permutations for assessing probability values. For D and $G"_{ST}$, standard errors (SE) were calculated by jackknifing over *loci* and 95% confidence intervals (CI) were obtained by bootstraping over *loci* (Peakall and Smouse, 2012). Resampling jackknife estimates of total allelic richness, per population samples, were conducted with the aid of EstimateS (Colwell and Elsensohn, 2014) and its confidence intervals were calculated in R. Rarefaction analysis of allelic richness was conducted for controlling sample size bias, with the aid of R packages dplyr, tidyr and pegas (Paradis, 2010). R visualizations were plotted with the aid of ggplot2.

A discriminant analysis of principal components (DAPC - Jombart *et al.*, 2010) was conducted through adegenet, by *de novo* grouping, initially retaining 80% of total variation captured by principal components (PC) and with optimal k chosen according to the Bayesian information criterion (BIC). A cross-validation procedure, with 100 replicates, was executed and used to gauge the DAPC analysis reliability, with the best number of PC found. Finally, the DAPC was also conducted with population sample information, with the same optimal number of PC retained in cross-validation, in order to visually record the genetic diversity within and among groups.

Figure 2. Genome assembly statistical features for *Prochilodus lineatus*.
a) The radius inside the circular plot marks the length of the longest scaffold resolved and the grey segments represent the cumulative percentage of the assembly contained in scaffolds equal or longer than a particular length. **b**) Cumulative contig/scaffold length of the genomic assembly, as a function of the the number of sequences resolved. The point in the middle of the curve marks the L_{50} feature, the smallest number of sequences, which the sum surpass half the total genome assembly length (point at the end of the cuve). G=Gigabases and k=thousand of sequences.

Results

Genomics, bioinformatics and microsatellite markers development

Approximately 483.7 million short-reads (43.53 gigabases) were generated for *P. lineatus* and deposited at the the Sequence Read Archive (SRA) database, from NCBI (Bioproject accession code PRJNA1120875). Its CG content was 42% and per base quality value (Q) ranged from 31 to 39, with an average of 38 per read. Based on the k-mer counts, total genome size was estimated at 1.65 Gb.

The *de novo* genome assembly generated was 1.4 Gb (NCBI accession JBGFTF000000000), with 31X coverage. The total number of contigs/scaffolds was 738,832 sequences, with N_{50}=5,503 bp and L_{50}=52,608. The average length of contigs was 1,901 with the longest being 192,837 bp (Figure 2). The BAM file associated with it can also be retrieved from NCBI (PRJNA1120875). A total of 37,476 perfect di- to hexa-nucleotide microsatellite *loci* for *P. lineatus* were characterized *in silco* and compiled in Supporting Information (Supplementary Files 1 and 2). These were mapped upon the genome assembly, but around 12% of putative *loci* could not be traced back to it.

Of the 22 primer pairs tested, five (Prol14, Prol15, Prol18, Prol31 and Prol52) resulted in complex amplification patterns and two (Prol56 and Prol28) failed to amplify. Fifteen new microsatellite markers were successfully amplified and results for the pooled sample of 140 fish are presented in Table 2. The optimal PCR conditions were as initially tested, except for Prol02 and Prol19 (6.7 pmol of each primer), the latter using 35 cycles, and Prol54, which also used 35 cycles. The average number of alleles per locus was 22.7 (±1.3 standard error - SE).

The amplification/genotyping failure rate varied from 2.1% (Prol53) to 32.8% (Prol07), with average 18.6% (±7.9 SE), including instances where one allele was scored and the other deemed undetermined for the same diploid genotype. Six DNA samples (four from Camargos, one from Itutinga 1 and one from Funil) failed to amplify between five and ten *loci*. These were removed from data set and most downstream analyses were performed with and without it, in parallel. Marginal or no differences were observed when comparing results and, thus, these samples were retained.

Population genetics

Adjusted significance for 60 multiple HWE tests was set to p<0.008 following the sequential Bonferroni correction. By this criterion, only six *loci* were deemed in HWE. The only locus found to conform with HWE within every sample, Prol07, was not in equilibrium across all four population samples (p=0.002). All deviations were towards heterozygote deficits. Tests for LD within population samples, had an adjusted significance level α=0.0001. No LD was inferred, except for six instances: Prol06 and Prol54 for Camargos; Prol01 and Prol19, Prol05 and Prol57 for Itutinga 1; Prol01

Table 2. New validated microsatellite markers for *Prochilodus lineatus*, along observed amplified DNA fragment length intervals for each locus and its respective defined allelic classes.

Locus	Primer sequence	Motif	T_m (°C)	Size range (bp)	N_A	N
Prol01	F: CAAAAGAAAAACGAAAATAAAAGAAACA R: CCCTCTGGGATCAATATAGTATGTATGT	AGAT	54.0	109-261	24	117
Prol02	F: TTTAAACTCATTCAACCGGCCTAC R: GTTGGAGGTTACTTGATAGGGAGGTA	TATC	54.0	113-389	24	119
Prol03	F: TTCCCCATCTTTAGAAATGTCCTTC R: CTCCTGGGGATAAACCATCAATCT	AGAT	63.6	134-375	27	122
Prol04	F: TCTGAAACTATTTTTATAGGTAGGGGGA R: TCCTCTCACCTACAAACTTCCTCAA	TAA	56.4	113-264	25	125
Prol05	F: TCTATCCATCCAATCTATCCATCCA R: AAATTCCATAGTTTGGAGTGACGC	ATCT	58.8	89-276	31	114
Prol06	F: GGCCACTATAAAAGGTAGGCACAA R: CATACGCTGTCCTAAATGTTCAGC	AGAT	54.0	121-227	17	104
Prol07	F: TGGAGGCTGCAGGTATAAGTTTTT R: AAGGCTGGTTCCCTAACCTCC	TTA	56.4	102-205	20	94
Prol08	F: AAGGGCAAGATCAGACAAACAAG R: GAATACTACGCTCATTTCAGAAGAAAAA	TAT	63.6	146-288	16	122
Prol10	F: CACAGACAGACACATATGGGGA R: CAGGTATGTTCTATTGAAATGCTAACAC	AT	61.2	145-316	25	100
Prol11	F: AGAGCTTCAATAGGCTGAAGTCGT R: TGAATTTCCCTCTGGGATCAATAA	TAGA	56.4	113-222	23	110
Prol12	F: GTTGCACCTTTGTAGAGGGGTATG R: ACAGCCGTGGATATAGTCAGAAGC	TCA	56.4	117-280	23	107
Prol19	F: TAAAATGTGGTTTCCTGTGCAATG R: TTTGCATTTTATCCATCTTTTCCC	TAGA	60.0	87-200	18	106
Prol53	F: ATTGCCTGTGTCACTTAAGCATGT R: TGCTACTGGCTGCTAAATTTCCTT	AGAT	51.0	127-209	13	137
Prol54	F: CGAGACGTCTAGCTGACACCATAA R: TTGACTTCTTGCTGTCATGTCAAA	AGAT	49.0	96-309	28	110
Prol57	F: AATGAAACACAGTCAGGCTCCTTC R: TTCCTTCAGGCAATGCACTTTTAT	ATA	49.0	139-317	27	122

T_m=Primers annealing temperature; N=Sample size (successfully genotyped specimens); F=Forward primer; and R=Reverse primer; N_A=Number of alleles.

and Prol06, Prol04 and Prol07 for Itutinga 2; and finally, Prol53 and Prol57 for Funil. The following *loci* pairs exhibited LD for the pooled sample of fish: Prol01 and Prol06; Prol04 and Prol07; Prol01 and Prol19; Prol06 and Prol54; Prol05 and Prol57; and lastly, Prol53 and Prol57. There was no evidence for scoring errors due to stuttering or large allele dropout, but the presence of null alleles was suggested as possible, due to general excess of homozygotes for most allele size classes, for all *loci*, in all population samples, except for Prol07.

Allelic richness ranged from 10.8 (±0.8 SE) alleles for Camargos to 16.9 (±0.9 SE) alleles for and Itutinga 2 (Table 3). This trend was confirmed by the jackknife estimates of allelic richness (Supporting Information - Supplementary Figure 1), along the results for the rarefaction analysis (Supplementary Figure 2). Camargos also had the lowest expected heterozygosity, while displaying the highest inbreeding coefficient value, along the Itutinga 2 sample (Table 3). All population samples exhibited private alleles, ranging from 7 alleles for Camargos and 37 alleles for Itutinga 2 (average 19.7 ±12.7 SE).

Table 3. Genetic diversity results for each microsatellite maker across population samples in *Prochilodus lineatus*. Probability for Hardy-Weinberg test had significance set to α=0.008. All probability values (p) are significant, except where noted (ns), and values deemed zero are actually extremely low non-zero values p<0.00009.

Sample	*Locus*	N	N_A	H_O	H_E	F	HWE (p)
	Prol01	17	14	0.65	0.91	0.29	0.0023
	Prol02	15	14	0.60	0.89	0.33	0
	l**Prol03**	13	13	0.31	0.91	0.66	0
	Prol04	17	10	0.41	0.87	0.53	0
	Prol05	14	13	0.36	0.9	0.6	0
	Prol06	14	9	0.43	0.81	0.47	0
	Prol07	15	10	0.73	0.85	0.14	0.0191[ns]
Camargos	**Prol08**	13	4	0.08	0.65	0.88	0
	Prol10	16	10	0.69	0.85	0.19	0.0049
	Prol11	17	12	0.35	0.88	0.6	0
	Prol12	13	6	0	0.77	1	0
	Prol19	12	9	0.50	0.84	0.41	0.0004
	Prol53	18	9	0.33	0.77	0.57	0
	Prol54	17	15	0.41	0.91	0.55	0
	Prol57	17	11	0.41	0.87	0.53	0
	Over *loci*	15.2	10.6	0.42	0.85	0.52	0
	Prol01	33	16	0.27	0.91	0.7	0
	Prol02	33	18	0.55	0.92	0.41	0
	Prol03	31	16	0.52	0.91	0.43	0
	Prol04	36	13	0.31	0.9	0.66	0
	Prol05	26	22	0.62	0.94	0.34	0
	Prol06	34	11	0.21	0.87	0.76	0
	Prol07	20	12	0.95	0.88	-0.08	0.0157[ns]
Itutinga 1	**Prol08**	38	10	0.45	0.85	0.48	0
	Prol10	33	20	0.73	0.94	0.22	0
	Prol11	30	16	0.57	0.89	0.37	0
	Prol12	31	13	0.16	0.9	0.82	0
	Prol19	30	13	0.23	0.91	0.74	0
	Prol53	39	9	0.31	0.64	0.52	0
	Prol54	34	20	0.38	0.92	0.59	0
	Prol57	38	16	0.47	0.91	0.48	0
	Over *loci*	32.4	15	0.45	0.89	0.50	0
	Prol01	34	19	0.29	0.92	0.68	0
	Prol02	36	18	0.56	0.91	0.39	0
	Prol03	39	19	0.36	0.93	0.61	0
	Prol04	33	14	0.27	0.91	0.7	0
	Prol05	38	24	0.39	0.94	0.58	0
	Prol06	27	14	0.11	0.89	0.87	0
	Prol07	27	18	0.81	0.91	0.1	0.0498[ns]
Itutinga 2	**Prol08**	39	14	0.62	0.87	0.3	0
	Prol10	27	15	1	0.91	-0.1	0

		N	N_A	H_O	H_E	F	HWE
	Prol11	33	17	0.27	0.88	0.69	0
	Prol12	32	17	0.34	0.91	0.62	0
	Prol19	31	13	0.13	0.89	0.86	0
	Prol53	40	10	0.18	0.65	0.73	0
	Prol54	34	19	0.29	0.92	0.68	0
	Prol57	36	22	0.81	0.92	0.13	0.0210^{ns}
	Over *loci*	33.7	16.9	0.43	0.89	0.52	0
	Prol01	33	16	0.15	0.92	0.84	0
	Prol02	35	17	0.49	0.91	0.46	0
	Prol03	39	21	0.51	0.95	0.46	0
	Prol04	39	17	0.51	0.92	0.44	0
	Prol05	36	27	0.56	0.95	0.42	0
	Prol06	29	12	0.21	0.79	0.74	0
	Prol07	32	11	0.91	0.85	-0.07	0.4959^{ns}
Funil	**Prol08**	32	13	0.66	0.86	0.24	0
	Prol10	24	18	1	0.93	-0.08	0.7323^{ns}
	Prol11	30	18	0.4	0.92	0.57	0
	Prol12	31	19	0.29	0.9	0.68	0
	Prol19	33	15	0.15	0.91	0.83	0
	Prol53	40	11	0.23	0.68	0.67	0
	Prol54	25	14	0.72	0.91	0.2	0.0052
	Prol57	31	21	0.71	0.93	0.23	0
	Over *loci*	32.6	16.67	0.50	0.89	0.44	0
Overall		140	341	0.45	0.91	0.50	0

N=Sample size; N_A=Number of alleles; H_O=Observed heterozygosity; H_E=Expected heterozygosity; F=inbreeding coefficient; HWE=Exact test for Hardy-Weinberg equilibrium.

The AMOVA results showed that only 1% of total variation was explained by the four population samples, with the
remaining variation being explained by differences among individuals (52%) and within genomes (47%). All pairwise
fixation estimates for G''_{ST} and Jost's D were significantly different from 0, including between samples collected in the
same location, but at different times (Table 4). These Itutinga samples (2013 and 2023) had the lowest values, while
Camargos showed the highest overall pairwise variation and differentiation, for both respective indices. Considering
all 15 microsatellite *loci*, among the four population samples, the unbiased standardise fixation index G''_{ST} reached
0.16 (±0.03 SE; 0 to 0.26 CI, n=999; p=0.001) and the total genetic differentiation estimate of Jost's D, was 0.15
(±0.03; 0 to 0.24 CI, n=999; p=0.001).

Hierarchical analysis of the genetic population structure

Maximum likelihood estimation (MLE) of admixture (Figure 3) pointed to K=2 as the most likely number of clusters,
according to Wang's D_{LK2} metric. However, F_{STIS} suggested K=4 as the most likely number of groups (Supporting
Information - Supplementary Figure 3). The 1st order admixture analysis allowed the definition of two clusters (with
respective individual IDs - Figure 3a). Estimates of D_{LK2} for both these clusters yielded the value of K=2 as optimal
in the 2nd order analysis (Figure 3b), while F_{STIS} recorded values of K=5 and K=7 (Supplementary Figure 4). Finally,
these four newly defined individual rearrangements were studied in the 3rd order analysis (Figure 3c), and D_{LK2}
consistently pointed to K=2 for each and every subject cluster (Supplementary Figure 5; individual assignments for
all hierarchical orders can be found in Supplementary File 3). We addressed and discarded the possibility of panmixia
(K=1), at this and each hierarchical order level, by directly inspecting log likelihood values of the clustering analysis,
against the results for the assumed value for K (Supporting Information - Supplementary File 4 - Supplementary
Table 1).

Table 4. Summary of pairwise fixation index G''_{ST} (below diagonal) and Jost's genetic distance D (above diagonal) for microsatellites diversity among population samples of *Prochilodus lineatus* from the Upper Grande River, MG.

	Camargos	Itutinga 1	Itutinga 2	Funil
Camargos	-	0.156**	0.222**	0.203**
Itutinga 1	0.171**	-	0.066*	0.087**
Itutinga 2	0.241**	0.072*	-	0.167**
Funil	0.221**	0.095**	0.180*	-

Figure 3. Individual admixture coefficients (bars) for K=2, across inclusive levels of hierarchical genetic structure of *Prochilodus lineatus* populations from the Upper Grande River, MG, Brazil.

Different shades of grey represent alternative clusters of origin, with the proportional size of bars indicating the respective genomic contributions from ancestral demes. For each panel, the diagram above shows the sample in the original input sequence, while diagram below shows the sample organized as newly defined clusters. **a**) 1st order analysis, showing fish grouped according to population sample and original ID. **b**) 2nd order analysis from clusters defined in the 1st order. **c**) 3rd order analysis of 2nd order clusters. For 2nd and 3rd orders, the numbers below bars are arbitrary and not equivalent to original ID.

Figure 4. Density of individual clusters defined by the initial *de novo* discriminant analysis of principal components (DAPC) for *Prochilodus lineatus* from the Upper Grande River, MG.

These different MLE sample arrangements showed a consistent decrease in intrapopulation diversity parameters for the higher orders, such as expected heterozygosity and number of alleles (Supplementary File 4 - Supplementary Table 2), while interpopulation indicators ($G"_{ST}$ and Jost's D) steadily increased (Supplementary File 4 - Supplementary Tables 3-5). The HWE analysis of MLE defined clusters indeed showed a tendency towards Hardy-Weinberg expectations in higher order group arrangements. In average, 3rd order clusters had more *loci* in HWE, with one particular cluster showing eight *loci* in equilibrium (Supplementary File 5).

The DAPC performed without population sample information pointed for two main clusters, according to the BIC (Figure 4, Supplementary Figure 6). This initial DAPC also revealed that two alleles from Prol53 (03 and 04) had the most weight in defining variable contributions. The cross-validation assay pointed to 30 as the ideal number of PC to be retained (Supplementary Figure 7) and the ratification of the DAPC according to this parameter led to very convergent results (Supplementary Figure 8). The DAPC performed with this number of PC and information of population samples revealed an apparent level of genetic differentiation among locations (Figure 5).

The Bayesian approach to admixture also resulted in alternative values for best K estimates, with K=8 according to Pritchard's Pr(X|K) while, both, Evanno's ΔK and Wang's parsimony estimators resolved K=5 (Figure 6).

Figure 5. DAPC for *Prochilodus lineatus* from the Upper Grande River, MG, performed with population sample information, with three discriminant functions (scaled discriminant analysis eigenvalues showed on the bottom right).

Figure 6. Admixture plots from the STRUCTURE clustering results for a) K=5 and b) K=8 for *Prochilodus lineatus* from the Upper Grande River, MG. Each colour represents a different cluster. Note the apparent differentiation at Camargos and Itutinga 2.

Discussion

Genomics, bioinformatics and microsatellite markers development

Genome assemblies are becoming currently mandatory for many different biological disciplines (Jung *et al.*, 2020). Our work takes a new step towards this goal, by publicly presenting the first comprehensive genomic resources for this highly iconic species, *P. lineatus*. The volume of short-read DNA sequence data presented here surpass the only previously available FASTQ data for *P. lineatus* (Stornioli *et al.*, 2021) by a factor of 12. This number of sequences is, though, one order of magnitude below the short-reads data reported for the trans-Andean congeneric species, *P. magdalenae* (Yepes-Blandón *et al.*, 2022) which, however, is not currently publicly available. The CG content and quality rate of reads were the same as the ones previously presented for other piracema fish (*e.g.* Yazbeck *et al.*, 2018; Graciano *et al.*, 2022). It's also the most extensive dataset of short-reads publicly disclosed so far for any member of the Prochilodontidae family.

This work provides the first *de novo* genome assembly for *P. lineatus*. It's slightly longer then the described genome size reported in *P. magdalenae* (Yepes-Blandón *et al.*, 2022), despite its extensive fragmentation, and shorter than estimated by the k-mer method carried, which is not always precise (Pflug *et al.*, 2020). The limited length of the short-reads used here contributed to curtailing our genome assembly, primarily motivated by the exclusive search of microsatellite *loci*. The full sequence of this *P. lineatus* specimen's mitogenome was previously presented in Santos *et al.* (2021). This novel genome assembly might be explored for the initial annotation of longer characterized sequences (*e.g.* Graciano *et al.*, 2021).

The genome wide microsatellite panel produced here, with more than 37 thousand putative *loci*, can now be free and easily accessed for the rapid, low-cost development of potentially hundreds of new empirically validated markers. It may also be explored for technological development of practical tools for hatcheries and management of impacted areas, such as multiplexed sets of markers (*e.g.* Carvalho *et al.*, 2019). Some of these resources may also, arguably, be applied in different fishes, particularly in other prochilodontid and characid species (*e.g.* Barbosa *et al.*, 2006; Carmo *et al.*, 2015). The relatively high rate of success in empirical assays attests to the high quality of this panel. These 15 new DNA markers presented here increase by 84% the number of available microsatellites for *P. lineatus*. We observed a variable rate of amplification success among these new markers. Some *loci* presented a relatively high amplification/genotyping failure rate (*e.g.* Prol07). Some approaches (*e.g.* admixture assignments) are found to be reasonably insensible to missing data, specially when it is not due to a systematic cause (Falush *et al.*, 2007; Wang, 2022). Even null alleles might have only a marginal impact (Carlsson, 2008; Putman and Carbone, 2014; Budowle and Sajantila, 2024). For some other applications, the issue of missing data is more critical (*e.g.* estimating effective population size, Ne – Peel *et al.*, 2013). The microsatellite Prol53 had a particularly high rate of amplification and genotyping and high information content. Despite Microchecker suggesting the possible presence of null alleles for most *loci* in all populations, we consider this an artefact due to the inherent sample substructuring, which leads to critical confounding effects in this analysis (Waples, 2015).

Population genetics

A study by Wang *et al.* (2021) suggests that applying less than 11 or 12 microsatellite markers may lead to important bias in population genetics structure assessments. The results presented here are the most extensive efforts so far, concerning the number of analysed *loci*, in *P. lineatus*. It revealed high levels of microsatellite diversity in fish from the uppermost portion of the Grande River, MG, under the influence of large dams. This is a common result to other former studies regarding population genetics in *P. lineatus* with close genomic sampling efforts (*e.g.* Rueda *et al.*, 2013; Perini *et al.*, 2021).

The genetic diversity analysis of population samples showed a lower genetic richness for fish upstream the Camargos dam, the last artificial obstacle towards the upper end of the Grande River Basin. This is an area deemed as priority for conservation of migratory fishes in the state of Minas Gerais (Drummond, 2005) and has been segregated from the downstream system for the last 65 years, since the construction of the Camargos power plant dam. It does not bear a fish passage and larvae and juveniles are not likely to successfully drift downstream (Suzuki *et al.*, 2011). Fish transposition, using specimens caught downstream, and repopulation initiatives with hatchery broodstock (mainly *P. lineatus* - Saraiva and Pompeu, 2016) did occur in the region for the last decades, but was mostly executed without purposefully defined goals, other than fulfilling regulatory agencies' licensing demands or in the context of environmental education actions. This portion of the river is still connected to a last major free-flowing upstream Grande River tributary, the Aiuruoca River, MG. The Camargos dam is, therefore, isolating fish from the rest of the hydrographic system below and stocking operations might have been inefficient in providing appropriate gene flow connectivity for the system above (lack of migrants). Its local populations are most likely undergoing stronger random genetic drift, which leads to genetic impoverishment within demes and differentiation among them. This notion is corroborated by the results showing the largest amount of interpopulational variation and differentiation in Camargos, from all other samples. Thus, our results support the hypothesis that dams are importantly isolating

fish populations in the Upper Grande River. The DAPC performed with sample information also supports this notion, based on the visual inspection of microsatellite variation within and among population samples.

The little difference among population samples registered by the AMOVA is compatible with a single, almost panmitic population, or could be explained by more complex, cryptic patterns of genetic variability. Yet, it is deemed as less straightforward in the interpretation of hypervariable markers, such as microsatellites (Jost, 2008). The G''_{ST} fixation index, along the genetic differentiation estimator Jost's D, pointed to significant deviations from panmixia, with all comparisons significantly different from zero. This supports the hypothesis of isolation by dams, over fish unable to freely migrate.

The 2023 sample from the Itutinga location is the most genetically diverse, followed by fish caught downstream the Funil dam. The latter, as first raised in Yazbeck and Kalapothakis (2007), could be preventing migrating fish of potentially different stocks, from a relatively long downstream stretch of free flowing waters, from proceeding further upstream, despite its fish lift. This virtual *cul-de-sac* could be leading to a higher observed genetic diversity in the sample by accumulation, comparable to the trapped gene pool effect discussed by Perini *et al.* (2021). Upstream this dam lies the mouth from two major free-flowing tributaries of the Grande River, which could be natural candidates for harbouring different *P. lineatus* stocks (*e.g.* Suzuki *et al.*, 2011), the Mortes River, and the the Capivari River.

The *P. lineatus* population at Itutinga's reservoir, in turn, has undergone genetic variation and differentiation of around 7% in 10 years. Despite being closed and strongly isolated (without tributaries), it seems to have locally accumulated genetic variation (allelic richness), since 2013. A plausible explanation for this would be the broodstocking operations of *P. lineatus* from the hatchery situated on its banks. Repeated introduction (or escape) of broodstock fish, with varying sources for spawning matrices, along the marked tendency for the occurrence genetic bottlenecks, over particular reproductive efforts (single or few spawned couples), could be yielding alternative sets of new alleles being introduced into this limited stretch of the Grande River. While this location had the highest levels of diversity (N_A and H_E), it also had the highest inbreeding coefficient, reinforcing the notion of hatchery driven substructuring. For more than five years now, though, the hatchery station operates mainly for scientific research and has ceased its fish stocking in the Grande River, MG, and region. These results support the hypothesis that hatcheries can influence the genetic diversity in populations of fish trapped between dams.

Regarding our main hypothesis of the occurrence of mixed stocks in the Upper Grande River, MG, the extensive verified lack of adherence to Hardy-Weinberg proportions in all population samples, always towards heterozygotes deficit, along some verified LD, suggest strong genetic stratification in all sampled populations (*i.e.* more than one deme being sampled) and, thus, is in accordance to its prediction. The fact it is so pervasive across *loci* allows us to initially disregard other putative confounding factors, such as eventual null alleles. It can be biologically interpreted as an expression of the Wahlund effect, leading to a pronounced downward departure from the expected heterozygosity under HWE.

Hierarchical genetic structure analysis

The admixture model seems biologically meaningful and appropriate for studying potamodromous Neotropical fishes. These fish might experience multiple reproductive runs, through several years which could potentially result in variable breeding outcomes, regarding the moment and place of spawning. More importantly, potential genetically distinct individuals and stocks might end up co-occurring in the same locality and time during migratory runs, or trapped between adjacent dams. This would isolate them from their original breeding grounds, leading to subsequent exogamy among different demes.

The MLE approach for admixture, which estimates population genetics parameters that maximize a likelihood function (Wang, 2003, 2022), is a powerful and scalable alternative to the more hegemonic Bayesian inference. Bayesian analysis is based on the conditional probabilities of genotypes, given some population genetics parameter and which is, in turn, highly sensible to often subjective prior probabilities (van de Schoot *et al.*, 2021). Since admixture models are highly hyperdimensional (in our particular case, from 466 independent variables, for K=1, up to 6,384, for K=14), the MLE path has the advantage of high accuracy, coupled with lower computational demands (Wang, 2022), which allows for more extensive replication. Here, it showed a high degree of convergence, across replicates (many have zero variance), for different K.

We have adopted Wang's D_{LK2} estimator for optimal K, as it was shown to be generally more reliable, over varying investigated scenarios, in comparison with the F_{STIS} estimator (Wang, 2022). The 1st order MLE analysis of admixture and the DAPC both initially pointed for two groups, which did not correlate with sampling scheme. Both clusters contain members from all sampled populations. This strongly suggests the occurrence of mixed stocks of *P. lineatus*, from the main channel of the Upper Grande River. Caution was taken to dismiss panmixia as the best possibility, by directly examining the log likelihood values of alternative models, since D_{LK2} would not possibly capture K=1 (Wang, 2022), just like Evanno's ΔK for Bayesian estimation (Janes *et al.*, 2017), and given that the F_{STIS} estimator would always be zero (Wang, 2022). The results from the Bayesian analysis of admixture also reinforce the notion of several, different fish stocks in our total sample, not correlated with sampling locations or moments,

supporting the hypothesis of the occurrence of mixed stocks throughout the Upper Grande River Basin.

Different values of K can reveal multiple underlying aspects of population structure and distinct stages in the population's history (Gilbert, 2016; Janes *et al.*, 2017; Gehri *et al.*, 2021; Wang, 2022). The two major genetically differentiated groups defined in the 1st order MLE analysis could be the reflex of existence of two main ancient breeding grounds (*e.g.* the Mortes, Capivari or the Aiuruoca Rivers), defining forks along migratory routes, splitting groups in the Upper Grande River. The DAPC conducted without assigning samples to localities or groups also revealed particular alleles from one of the best performing markers as being important in explaining the variation between these two DAPC defined clusters, and the cross-validation attested for its robustness. However, this method might show irregular success across particular scenarios, such as with high migration rate (as would be expected in admixed populations) and low level of differentiation among demes (as might be expected of species with high mobility), often underestimating the presence of more internal groups (Miller *et al.*, 2020).

Together, these results confirm the probable existence of at least two mixed stocks along the studied area, not explained by the recent implementation of dams. Evidence for mixed stocks in *P. lineatus* were previously forwarded, for example, by Rueda *et al.* (2013), Avigliano *et al.* (2019) and indirectly by Perini *et al.* (2021). The Bayesian exploration of the admixture model suggested deeper levels of substructuring, pointing to as many as eight possible clusters in the whole sample of fish, a scenario compatible with the eight demes defined during the "peeling" of MLE layers.

Aiming to deepen the exploration of these seemingly co-occurring genetically mixed stocks of *P. lineatus* from the main channel of the Upper Grande River, the two clusters initially defined by MLE were iteratively unfolded, for two more analyses rounds. Consistent and surprisingly, every examined cluster revealed two newly characterized smaller subgroups, nested within clusters resolved in the immediate previous step (2^3). Judging by the sorted samples of newly defined clusters from Figure 3c, the F_{STIS} estimates from Supplementary Figure 5, or the HWE analysis from MLE clusters, there would be potentially more admixture degrees to be untangled, from further hierarchical analysis, although we would reach rather critical sample sizes. The MLE defined clusters showed the expected admixture model behaviour towards higher orders. The 3rd order had more genetic homogeneity within clusters and a maximization of genetic variation and distance among clusters. It also showed increasing numbers of *loci* in HWE, attesting to the meaningfulness of the hierarchical admixture analysis and the usefulness of these new microsatellite markers.

The observed stratified, self-similar, hidden pattern of the population genetic structure, across the larger sample of fish, may reveal the existence of potential different, well defined, ancestral breeding grounds, hierarchically nested within differential scale levels, throughout the basin, organized according to fractal geometry (e.g. 1st order river, 2nd and 3rd order tributaries, and so on). If this association is confirmed, we are led to conclude, it may very well constitute the first indirect genetic evidence for a strong degree of homing behaviour for a characiformes piracema fish species. Philopatric spawning, or homing behaviour, a thoroughly investigated issue in salmonid and other migratory species, is still poorly understood in the Neotropics. It has been documented for Neotropical catfishes (*e.g.* Batista and Alves-Gomes, 2006; Pereira *et al.*, 2009). But most importantly, it was suggested for the congeneric *P. argenteus*, based on radio-tagging results from the São Francisco River Basin (Godinho and Kynard, 2006) and discussed for *P. lineatus* itself by Careaga and Carvajal-Vallejos (2019), from morphometric data. Our work revives this area of inquiry, for migratory fish in South America.

Being particularly abundant and a conspicuously competent, far-ranging migratory species, we argue that in a pre-damming past, *P. lineatus* could probably reach the innermost free-flowing tributaries for spawning, upon the colossal La Plata Basin, which cuts through a diversified range of biomes and microhabitats. We hypothesize this would likely point towards philopatric behaviour as an evolutionary stable strategy, due to the possible occurrence of distinct local adaptations, as it has been shown in other fish species (*e.g.* Salles *et al.*, 2016; Mobley *et al.*, 2019; Vøllestad and Primmer, 2019).

In conservation biology, the understanding of degrees of admixture is important in management decisions (Wang, 2003, 2022). The confirmation of mixed stocks and the hierarchical genetically stratified nature of *P. lineatus* from the main channel of an important large and intensely dammed river will likely enlighten its future research and fisheries management. If further corroborated, these findings, particularly its ensuing propositions, would have direct implications to environmental and development decision-making. It highlights the possible central relevance of key smaller tributaries (actively sought for the implementation of new small hydroelectric power plants - Ferreira *et al.* 2022), for the conservation of genetic variation, in a socio-environmentally relevant species, as it highlights the importance of free-flowing tributaries of different orders throughout the basin.

On a broader level, we raise the question whether these apparent fractal layers from the genetic structure could imply in a more widespread phenomena for piracema, other migratory fishes, or even other non-aquatic migratory organisms, showing a temporary mixture, but segregated breeding grounds. We believe this is worthy of further investigation, since fractal structures are pervasive in nature, not only in biology (*e.g.* Bassingthwaighte, 1992; Leggett *et al.*, 2019; Meyer *et al.*, 2020; Sendker *et al.*, 2024), but it is also conspicuously present on major

geographical landscape attributes like mountain ranges, forest borders, rivers and coastlines (*e.g.* Xu *et al.*, 1993; Andronache *et al.*, 2019). The only direct discussion on the relation of fractals and genetic population structure were laid by Wingen *et al.* (2007), regarding computer simulations results on long-distance dispersal of clonal fungal spores, and by Lee (2020).

Our results will very likely benefit researchers and environmental managers in tackling challenging issues regarding the conservation of Neotropical migratory fishes, as they support the hypothesis of the isolating effects of dams and broodstocking over genetic variability. More importantly, the results presented here point to the potential role of tributaries of different orders in harbouring genetically diverse populations of migratory freshwater fishes.

Acknowledgements

We thank the fishermen couple Donizete Monteiro de Paula and Sônia Bárbara de Oliveira Paula, biologists and photographers couple Lucia and André Seale, biologists João de Magalhães Lopes, Miriam Aparecida de Castro, Raquel Coelho Loures and everyone at Peixe Vivo, José Mauro Ribeiro for bioinformatics assistance, Mônica Soares da Silva for assistance with marker development, Fausto Moreira da Silva Carmo for assistance with specimen collection and DNA extraction, Ian Cosenza Irsigler for map assistance, Pedro Manoel Galetti Jr for valuable comments and Evanguedes Kalapothakis. This work was financed by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq grant 303023/2023-6), Companhia Energética de Minas Gerais and Agência Nacional de Energia Elétrica (CEMIG-ANEEL grant GT345), by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (CAPES finance Code 001), and by Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG grant APQ-04569-10).

Contributions

G.M.Y. conceived, conducted and supervised the study, raised funding, performed bioinformatics and population genetics analyses, wrote and edited the manuscript; G.A.S.N. performed marker development, population genetics analysis and wrote the manuscript; L.C.C. performed marker development; R.C.D.G. performed bioinformatics analysis; R.P.S. performed bioinformatics analysis; R.S.O. edited the manuscript, supervised and performed bioinformatics pipelines. All contributors read and agreed upon the final manuscript.

Bibliography

Alix, M., Kjesbu, O. S., and Anderson, K. C. From gametogenesis to spawning: How climate-driven warming affects teleost reproductive biology. *Journal of Fish Biology*, 97(3):607–632, Sept. 2020. doi: 10.1111/jfb.14439.

Andronache, I., Marin, M., Fischer, R., Ahammer, H., Radulovic, M., Ciobotaru, A.-M., Jelinek, H. F., Di Ieva, A., Pintilii, R.-D., Drăghici, C.-C., Herman, G. V., Nicula, A.-S., Simion, A.-G., Loghin, I.-V., Diaconu, D.-C., and Peptenatu, D. Dynamics of Forest Fragmentation and Connectivity Using Particle and Fractal Analysis. *Scientific Reports*, 9(1):12228, Aug. 2019. doi: 10.1038/s41598-019-48277-z. Publisher: Nature Publishing Group.

Astashyn, A., Tvedte, E. S., Sweeney, D., Sapojnikov, V., Bouk, N., Joukov, V., Mozes, E., Strope, P. K., Sylla, P. M., Wagner, L., Bidwell, S. L., Brown, L. C., Clark, K., Davis, E. W., Smith-White, B., Hlavina, W., Pruitt, K. D., Schneider, V. A., and Murphy, T. D. Rapid and sensitive detection of genome contamination at scale with FCS-GX. *Genome Biology*, 25(1):60, Feb. 2024. doi: 10.1186/s13059-024-03198-7.

Avigliano, E., Pisonero, J., Bordel, N., Dománico, A., and Volpedo, A. V. Mixed-stock and discriminant models use for assessing recruitment sources of estuarine fish populations in La Plata Basin (South America). *Journal of the Marine Biological Association of the United Kingdom*, 99(6):1429–1433, Sept. 2019. doi: 10.1017/S002531541900016X.

Barbosa, A. C. D. R., Corrêa, T. C., Galzerani, F., Galetti Jr, P. M., and Hatanaka, T. Thirteen polymorphic microsatellite loci in the Neotropical fish Prochilodus argenteus (Characiformes, Prochilodontidae). *Molecular Ecology Notes*, 6(3):936–938, 2006. doi: 10.1111/j.1471-8286.2006.01406.x. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1471-8286.2006.01406.x.

Bassingthwaighte, J. B. FRACTAL VASCULAR GROWTH PATTERNS. *Acta stereologica*, 11(Suppl 1):305–319, 1992.

Batista, J. S. and Alves-Gomes, J. A. Phylogeography of Brachyplatystoma rousseauxii (Siluriformes - Pimelodidae) in the Amazon Basin offers preliminary evidence for the first case of "homing" for an Amazonian migratory catfish. *Genetics and molecular research*, 5(4):723–740, Dec. 2006.

Bowen, S. H. Digestion and assimilation of benthic biofilm by the Sábalo, *Prochilodus lineatus* . *Journal of Fish Biology*, 100(1):107–116, Jan. 2022a. doi: 10.1111/jfb.14924.

Bowen, S. H. The river flood pulse, benthic biofilm, and the nutrition of Prochilodus lineatus. *Environmental Biology of Fishes*, 105(2):213–230, Feb. 2022b. doi: 10.1007/s10641-022-01211-1.

Brandström, M. and Ellegren, H. Genome-wide analysis of microsatellite polymorphism in chicken circumventing the ascertainment bias. *Genome Research*, 18(6):881–887, June 2008. doi: 10.1101/gr.075242.107.

Budowle, B. and Sajantila, A. Short tandem repeats - how microsatellites became the currency of forensic genetics. *Nature Reviews. Genetics*, 25(7):450–451, July 2024. doi: 10.1038/s41576-024-00721-1.

Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. Technical Report LBNL-7065E, Lawrence Berkeley National Laboratory, Walnut Creek, Mar. 2014. URL https://escholarship.org/uc/item/1h3515gn.

Careaga, M. and Carvajal-Vallejos, F. Los sábalos (Characiformes: Prochilodontidae: Prochilodus) de Bolivia. pages 171–185. Sept. 2019.

Carlsson, J. Effects of microsatellite null alleles on assignment testing. *The Journal of Heredity*, 99(6):616–623, 2008. doi: 10.1093/jhered/esn048.

Carmo, F. M. d. S., Polo, M., Silva, M. A. d., and Yazbeck, G. d. M. Optimization of heterologous microsatellites in Piracanjuba. *Pesquisa Agropecuária Brasileira*, 50:1236–1239, Dec. 2015. doi: 10.1590/S0100-204X2015001200015. Publisher: Embrapa Secretaria de Pesquisa e Desenvolvimento; Pesquisa Agropecuária Brasileira.

Carmo, A. O., Brandão Dias, P. F. P., Martins, A. P. V., Bedore, A. G., and Kalapothakis, E. Complete mitochondrial genome sequence of Prochilodus lineatus (Characiformes, Prochilodontidae). *Mitochondrial DNA. Part A, DNA mapping, sequencing, and analysis*, 27(3):1946–1947, May 2016. doi: 10.3109/19401736.2014.971300.

Carolsfeld, J., Harvey, B., Carmen, R., and Anton, B. *Migratory Fishes of South America: Biology, Fisheries and Conservation Status*. IDRC, 2003. ISBN 978-0-9683958-2-0. Google-Books-ID: GCF3_kK6_8cC.

Carvajal-Vallejos, F. M., Duponchelle, F., Desmarais, E., Cerqueira, F., Querouil, S., Nuñez, J., García, C., and Renno, J.-F. Genetic structure in the Amazonian catfish Brachyplatystoma rousseauxii: influence of life history strategies. *Genetica*, 142 (4):323–336, Aug. 2014. doi: 10.1007/s10709-014-9777-2.

Carvalho, L., Ribeiro, J., and Yazbeck, G. Microsatellite multiplex systems for Brycon orbignyanus (Characiformes: Bryconidae). *Brazilian Journal of Biology*, 79(4):751–754, Nov. 2019. doi: 10.1590/1519-6984.192528.

Cassemiro, F. A. S., Albert, J. S., Antonelli, A., Menegotto, A., Wüest, R. O., Cerezer, F., Coelho, M. T. P., Reis, R. E., Tan, M., Tagliacollo, V., Bailly, D., da Silva, V. F. B., Frota, A., da Graça, W. J., Ré, R., Ramos, T., Oliveira, A. G., Dias, M. S., Colwell, R. K., Rangel, T. F., and Graham, C. H. Landscape dynamics and diversification of the megadiverse South American freshwater fish fauna. *Proceedings of the National Academy of Sciences*, 120(2):e2211974120, Jan. 2023. doi: 10.1073/pnas.2211974120. Publisher: Proceedings of the National Academy of Sciences.

Chagas, A. T. D. A., Carmo, A. O., Costa, M. A., Resende, L. C., Brandão Dias, P. F. P., Martins, A. P. V., and Kalapothakis, E. Description and comparison of two economically important fish species mitogenomes: *Prochilodus argenteus* and *Prochilodus costatus* (Characiformes, Prochilodontidae). *Mitochondrial DNA Part A*, 27(4):2852–2853, July 2016. doi: 10.3109/19401736.2015.1053125.

Challis, R. rjchallis/assembly-stats 17.02, Feb. 2017. URL https://zenodo.org/records/322347.

Chang, S. L., Ward, H. G. M., Elliott, L. D., and Russello, M. A. Genotyping-in-Thousands by sequencing panel development and
application for high-resolution monitoring of introgressive hybridization within sockeye salmon. *Scientific Reports*, 12(1):3441,
Mar. 2022. doi: 10.1038/s41598-022-07309-x.

Collins, S. M., Bickford, N., McIntyre, P. B., Coulon, A., Ulseth, A. J., Taphorn, D. C., and Flecker, A. S. Population Structure of a
Neotropical Migratory Fish: Contrasting Perspectives from Genetics and Otolith Microchemistry. *Transactions of the American
Fisheries Society*, 142(5):1192–1201, Sept. 2013. doi: 10.1080/00028487.2013.804005.

Colwell, R. K. and Elsensohn, J. E. EstimateS turns 20: statistical estimation of species richness and shared species
from samples, with non-parametric extrapolation. *Ecography*, 37(6):609–613, 2014. doi: 10.1111/ecog.00814. _eprint:
https://onlinelibrary.wiley.com/doi/pdf/10.1111/ecog.00814.

Drummond, G. M. *Biodiversidade em Minas Gerais: um atlas para sua conservação*. Fundação Biodiversitas, Belo Horizonte,
2005. ISBN 85-85401-21-4. Biodiversidade em Minas Gerais: um atlas para sua conservacão/ B615 / Gláucia Moreira
Drummond, ... [et al.]. 2. ed - Belo Horizonte: Fundação Biodiversitas, 2005. 222 p.: il color., fots., maps., grafs., tabs. Inclui
bibliografia e anexo. ISBN: 85-85401-21-4 1. Diversidade Biológica – Minas Gerais – Atlas. 2. Meio Ambiente – Minas Gerais
– Atlas. Drummond, Gláucia Moreira. CDU: 502.74.

Endruweit, M. Four new records of fish species (Cypriniformes: Nemacheilidae, Balitoridae; Characiformes: Prochilodontidae)
and corrections of two misidentified fish species (Tetraodontiformes: Tetraodontidae; Beloniformes: Belonidae) in Yunnan,
China. *Dong Wu Xue Yan Jiu = Zoological Research*, 35(1):51–58, Jan. 2014. doi: 10.11813/j.issn.0254-5853.2014.1.051.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. MultiQC: summarize analysis results for multiple tools and samples in a
single report. *Bioinformatics*, 32(19):3047–3048, Oct. 2016. doi: 10.1093/bioinformatics/btw354.

Falush, D., Stephens, M., and Pritchard, J. K. Inference of population structure using multilocus genotype data: dominant markers
and null alleles. *Molecular Ecology Notes*, 7(4):574–578, July 2007. doi: 10.1111/j.1471-8286.2007.01758.x.

Ferreira, D. G., Souza-Shibatta, L., Shibatta, O. A., Sofia, S. H., Carlsson, J., Dias, J. H. P., Makrakis, S., and Makrakis, M. C.
Genetic structure and diversity of migratory freshwater fish in a fragmented Neotropical river system. *Reviews in Fish Biology
and Fisheries*, 27(1):209–231, Mar. 2017. doi: 10.1007/s11160-016-9441-2.

Ferreira, M. E., Nogueira, S. H. d. M., Latrubesse, E. M., Macedo, M. N., Callisto, M., Bezerra Neto, J. F., and Fernandes, G. W.
Dams Pose a Critical Threat to Rivers in Brazil's Cerrado Hotspot. *Water*, 14(22):3762, Jan. 2022. doi: 10.3390/w14223762.
Number: 22 Publisher: Multidisciplinary Digital Publishing Institute.

Ferreira, D. G., Galindo, B. A., Limeira, D. M., Garla, A., Marques, A. J. C., Moreira, A. A., Rodrigues, M. V. H., Clemente,
R. C., Shimpo, J. Y., Martins, G. A., Almeida, F. S., Apolinário-Silva, C., Zanatta, A. S., Sofia, S. H., and Souza-Shibatta,
L. Different stocks of Prochilodus lineatus along the Cinzas River basin: contributions from a free-flowing tributary in the
Capivara Reservoir region, Paranapanema River basin (upper Paraná River). *Neotropical Ichthyology*, 21(4):e230080, 2023.
doi: 10.1590/1982-0224-2023-0080.

García-Castro, K. L. and Márquez, E. J. Temporal-scale assessment of population genetics of the freshwater fish Prochilodus
magdalenae in an area impacted by construction of a dam. *Hydrobiologia*, 851(6):1513–1531, Mar. 2024. doi: 10.1007/
s10750-023-05396-z.

Gehri, R. R., Gruenthal, K., and Larson, W. A. It's complicated: Heterogeneous patterns of genetic structure in five fish species
from a fragmented river suggest multiple processes can drive differentiation. *Evolutionary Applications*, 14(8):2079–2097, Aug.
2021. doi: 10.1111/eva.13268.

Gilbert, K. J. Identifying the number of population clusters with structure: problems and solutions.
Molecular Ecology Resources, 16(3):601–603, 2016. doi: 10.1111/1755-0998.12521. _eprint:
https://onlinelibrary.wiley.com/doi/pdf/10.1111/1755-0998.12521.

Godinho, A. L. and Kynard, B. Migration and Spawning of Radio-Tagged Zulega Prochilodus argenteus in a Dammed Brazilian
River. *Transactions of the American Fisheries Society*, 135(3):811–824, May 2006. doi: 10.1577/T04-176.1. Publisher: Taylor
& Francis _eprint: https://doi.org/10.1577/T04-176.1.

Graciano, R., Oliveira, R. S., Santos, I. M. d., and Yazbeck, G. d. M. Functional DNA annotation from a preliminary de novo
genome assembly of Brycon orbignyanus, an endangered Neotropical migratory fish. *Latin American Data in Science*, 1(2):
42–48, Nov. 2021. doi: 10.53805/lads.v1i2.12. Number: 2.

Graciano, R. C. D., Oliveira, R. S., Santos, I. M., and Yazbeck, G. M. Genomic Resources for Salminus brasiliensis. *Frontiers in
Genetics*, 13, Mar. 2022. doi: 10.3389/fgene.2022.855718. Publisher: Frontiers.

Hallerman, E. M. O. *Population genetics: principles and applications for fisheries scientists*. American Fisheries Society,
Bethesda, MD (USA), 2003. ISBN 1-888596-27-1. URL https://agris.fao.org/search/en/providers/122621/
records/647396cfe01106880097fed4.

Howe, N. S., Hale, M. C., Waters, C. D., Schaal, S. M., Shedd, K. R., and Larson, W. A. Genomic evidence for domestication
selection in three hatchery populations of Chinook salmon, *Oncorhynchus tshawytscha*. *Evolutionary Applications*, 17(2):
e13656, Feb. 2024. doi: 10.1111/eva.13656.

Janes, J. K., Miller, J. M., Dupuis, J. R., Malenfant, R. M., Gorrell, J. C., Cullingham, C. I., and Andrew, R. L. The K = 2 conundrum.
Molecular Ecology, 26(14):3594–3602, July 2017. doi: 10.1111/mec.14187.

Jombart, T., Devillard, S., and Balloux, F. Discriminant analysis of principal components: a new method for the analysis of
genetically structured populations. *BMC genetics*, 11:94, Oct. 2010. doi: 10.1186/1471-2156-11-94.

Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24(11):1403–1405, June
2008. doi: 10.1093/bioinformatics/btn129.

Jost, L. GST and its relatives do not measure differentiation. *Molecular Ecology*, 17(18):4015–4026, 2008. doi: 10.1111/j.
1365-294X.2008.03887.x. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-294X.2008.03887.x.

Jung, H., Ventura, T., Chung, J. S., Kim, W.-J., Nam, B.-H., Kong, H. J., Kim, Y.-O., Jeon, M.-S., and Eyun, S.-i. Twelve quick
steps for genome assembly and annotation in the classroom. *PLOS Computational Biology*, 16(11):e1008325, Nov. 2020. doi:
10.1371/journal.pcbi.1008325. Publisher: Public Library of Science.

Kalous, L., Bui, A. T., Petrtýl, M., Bohlen, J., and Chaloupková, P. The South American freshwater fish Prochilodus lineatus
(Actinopterygii: Characiformes: Prochilodontidae): new species in Vietnamese aquaculture: Prochilodus lineatus in Vietnam.
Aquaculture Research, 43(6):955–958, May 2012. doi: 10.1111/j.1365-2109.2011.02895.x.

Lee, C.-Y. The fractal dimension as a measure for characterizing genetic variation of the human genome. *Computational Biology*
and Chemistry, 87:107278, Aug. 2020. doi: 10.1016/j.compbiolchem.2020.107278.

Leggett, S. E., Neronha, Z. J., Bhaskar, D., Sim, J. Y., Perdikari, T. M., and Wong, I. Y. Motility-limited aggregation of mammary
epithelial cells into fractal-like clusters. *Proceedings of the National Academy of Sciences*, 116(35):17298–17306, Aug. 2019.
doi: 10.1073/pnas.1905958116. Publisher: Proceedings of the National Academy of Sciences.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. The Sequence
Alignment/Map format and SAMtools. *Bioinformatics*, 25(16):2078–2079, Aug. 2009. doi: 10.1093/bioinformatics/btp352.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. MEGAHIT: an ultra-fast single-node solution for large and complex
metagenomics assembly via succinct de Bruijn graph. *Bioinformatics (Oxford, England)*, 31(10):1674–1676, May 2015. doi:
10.1093/bioinformatics/btv033.

Luo, R., Liu, B., Xie, Y., Li, Z., Huang, W., Yuan, J., He, G., Chen, Y., Pan, Q., Liu, Y., Tang, J., Wu, G., Zhang, H., Shi, Y., Liu,
Y., Yu, C., Wang, B., Lu, Y., Han, C., Cheung, D. W., Yiu, S.-M., Peng, S., Xiaoqian, Z., Liu, G., Liao, X., Li, Y., Yang, H.,
Wang, J., Lam, T.-W., and Wang, J. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler.
GigaScience, 1:18, Dec. 2012. doi: 10.1186/2047-217X-1-18.

Marçais, G. and Kingsford, C. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics*,
27(6):764–770, Mar. 2011. doi: 10.1093/bioinformatics/btr011.

Meyer, H. V., Dawes, T. J. W., Serrani, M., Bai, W., Tokarczuk, P., Cai, J., de Marvao, A., Henry, A., Lumbers, R. T., Gierten, J.,
Thumberger, T., Wittbrodt, J., Ware, J. S., Rueckert, D., Matthews, P. M., Prasad, S. K., Costantino, M. L., Cook, S. A., Birney,
E., and O'Regan, D. P. Genetic and functional insights into the fractal structure of the heart. *Nature*, 584(7822):589–594, Aug.
2020. doi: 10.1038/s41586-020-2635-8. Publisher: Nature Publishing Group.

Miller, J. M., Cullingham, C. I., and Peery, R. M. The influence of a priori grouping on inference of genetic clusters: simulation
study and literature review of the DAPC method. *Heredity*, 125(5):269–280, Nov. 2020. doi: 10.1038/s41437-020-0348-2.
Publisher: Nature Publishing Group.

Mobley, K. B., Granroth-Wilding, H., Ellmen, M., Vähä, J.-P., Aykanat, T., Johnston, S. E., Orell, P., Erkinaro, J., and Primmer,
C. R. Home ground advantage: Local Atlantic salmon have higher reproductive fitness than dispersers in the wild. *Science*
Advances, 5(2):eaav1112, Feb. 2019. doi: 10.1126/sciadv.aav1112.

Oliveira-Farias, K. K. C., Farias, R. S., Neto, M. A. S., de Souza, M. E., Calado, L. L., Montes, M. A., Lima, A. P. S., Martinez, P.,
and Coimbra, M. R. M. Genetic structure and management of the Neotropical migratory fish Megaleporinus obtusidens on a
highly impacted river basin. *Hydrobiologia*, 849(7):1645–1658, Apr. 2022. doi: 10.1007/s10750-022-04808-w.

Paradis, E. pegas: an R package for population genetics with an integrated–modular approach. *Bioinformatics*, 26(3):419–420,
Feb. 2010. doi: 10.1093/bioinformatics/btp696.

Peakall, R. and Smouse, P. E. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an
update. *Bioinformatics*, 28(19):2537–2539, Oct. 2012. doi: 10.1093/bioinformatics/bts460.

Peel, D., Waples, R. S., Macbeth, G. M., Do, C., and Ovenden, J. R. Accounting for missing data in the estimation of contemporary
genetic effective population size (N(e)). *Molecular Ecology Resources*, 13(2):243–253, Mar. 2013. doi: 10.1111/1755-0998.
12049.

Pereira, L. H. G., Foresti, F., and Oliveira, C. Genetic structure of the migratory catfish Pseudoplatystoma corruscans
(Siluriformes: Pimelodidae) suggests homing behaviour. *Ecology of Freshwater Fish*, 18(2):215–225, 2009. doi: 10.1111/
j.1600-0633.2008.00338.x. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1600-0633.2008.00338.x.

Perini, V. R., Paschoalini, A. L., Bazzoli, N., Rizzo, E., and Carvalho, D. C. Metapopulation dynamics of the migratory fish
Prochilodus lineatus (Characiformes: Prochilodontidae) in a lotic remnant of the Grande River, Southeastern Brazil. *Neotropical*
Ichthyology, 19(4):e200046, 2021. doi: 10.1590/1982-0224-2020-0046.

Petry, A. C., Agostinho, A. A., and Gomes, L. C. Fish assemblages of tropical floodplain lagoons: exploring the role of connectivity
in a dry year. *Neotropical Ichthyology*, 1:111–119, Dec. 2003. doi: 10.1590/S1679-62252003000200005.

Pflug, J. M., Holmes, V. R., Burrus, C., Johnston, J. S., and Maddison, D. R. Measuring Genome Sizes Using Read-Depth, k-mers, and Flow Cytometry: Methodological Comparisons in Beetles (Coleoptera). *G3 (Bethesda, Md.)*, 10(9):3047–3060, Sept. 2020. doi: $10.1534/g3.120.401028$.

Pritchard, J. K., Stephens, M., and Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics*, 155 (2):945–959, June 2000. doi: $10.1093/genetics/155.2.945$.

Putman, A. I. and Carbone, I. Challenges in analysis and interpretation of microsatellite data for population genetic studies. *Ecology and Evolution*, 4(22):4399–4428, 2014. doi: $10.1002/ece3.1305$. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/ece3.1305.

R Core Team. R: A Language and Environment for Statistical Computing, 2024. URL https://www.R-project.org.

Reis, R. E., Albert, J. S., Di Dario, F., Mincarone, M. M., Petry, P., and Rocha, L. A. Fish biodiversity and conservation in South America. *Journal of Fish Biology*, 89(1):12–47, July 2016. doi: $10.1111/jfb.13016$.

Ribolli, J., Miño, C. I., Scaranto, B. M. S., Reynalte-Tataje, D. A., and Zaniboni Filho, E. Genetic evidence supports polygamous mating system in a wild population of *Prochilodus lineatus* (Characiformes: Prochilodontidae), a Neotropical shoal spawner fish. *Neotropical Ichthyology*, 18:e190123, June 2020. doi: $10.1590/1982-0224-2019-0123$.

Rosa, I. F. D., Oliveira, D. J. D., Cruz, V. P. D., Ashikaga, F. Y., Costa, G. O., Doretto, L. B., Senhorini, J. A., Rocha, R. D. C., Foresti, F. P., Oliveira, C., and Foresti, F. Temporal genetic structure of a stock of Prochilodus lineatus (Characiformes: Prochilodontidae) in the Mogi-Guaçu River ecosystem, São Paulo, Brazil. *Neotropical Ichthyology*, 20(2):e210156, 2022. doi: $10.1590/1982-0224-2021-0156$.

Rousset, F. genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources*, 8(1):103–106, 2008. doi: $10.1111/j.1471-8286.2007.01931.x$. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1471-8286.2007.01931.x.

Rueda, E. C., Carriquiriborde, P., Monzón, A. M., Somoza, G. M., and Ortí, G. Seasonal variation in genetic population structure of sábalo (Prochilodus lineatus) in the Lower Uruguay River. *Genetica*, 141(7-9):401–407, Sept. 2013. doi: $10.1007/s10709-013-9739-0$.

Salles, O. C., Pujol, B., Maynard, J. A., Almany, G. R., Berumen, M. L., Jones, G. P., Saenz-Agudelo, P., Srinivasan, M., Thorrold, S. R., and Planes, S. First genealogy for a wild marine fish population reveals multigenerational philopatry. *Proceedings of the National Academy of Sciences*, 113(46):13245–13250, Nov. 2016. doi: $10.1073/pnas.1611797113$. Publisher: Proceedings of the National Academy of Sciences.

Santos, R. P., Melo, B. F., Yazbeck, G. M., Oliveira, R. S., Hilário, H. O., Prosdocimi, F., and Carvalho, D. C. Diversification of *Prochilodus* in the eastern Brazilian Shield: Evidence from complete mitochondrial genomes (Teleostei, Prochilodontidae). *Journal of Zoological Systematics and Evolutionary Research*, 59(5):1053–1063, July 2021. doi: $10.1111/jzs.12475$.

Saraiva, S. O. and Pompeu, P. S. Fish hatchering and its effects on the morphology of Prochilodus lineatus (Actinopterygii: Prochilodontidae). *Brazilian Journal of Biology*, 76(1):209–217, Jan. 2016. doi: $10.1590/1519-6984.18514$.

Sayers, E. W., O'Sullivan, C., and Karsch-Mizrachi, I. Using GenBank and SRA. In Edwards, D., editor, *Plant Bioinformatics: Methods and Protocols*, pages 1–25. Springer US, New York, NY, 2022. ISBN 978-1-0716-2067-0. doi: $10.1007/978-1-0716-2067-0\_1$. URL https://doi.org/10.1007/978-1-0716-2067-0_1.

Sendker, F. L., Lo, Y. K., Heimerl, T., Bohn, S., Persson, L. J., Mais, C.-N., Sadowska, W., Paczia, N., Nußbaum, E., del Carmen Sánchez Olmos, M., Forchhammer, K., Schindler, D., Erb, T. J., Benesch, J. L. P., Marklund, E. G., Bange, G., Schuller, J. M., and Hochberg, G. K. A. Emergence of fractal geometries in the evolution of a metabolic enzyme. *Nature*, 628(8009):894–900, Apr. 2024. doi: $10.1038/s41586-024-07287-2$. Publisher: Nature Publishing Group.

Shedd, K. R., Lescak, E. A., Habicht, C., Knudsen, E. E., Dann, T. H., Hoyt, H. A., Prince, D. J., and Templin, W. D. Reduced relative fitness in hatchery-origin Pink Salmon in two streams in Prince William Sound, Alaska. *Evolutionary Applications*, 15 (3):429–446, Mar. 2022. doi: $10.1111/eva.13356$.

Sivasundar, A., Bermingham, E., and Ortí, G. Population structure and biogeography of migratory freshwater fishes (Prochilodus: Characiformes) in major South American rivers. *Molecular Ecology*, 10(2):407–417, Feb. 2001. doi: $10.1046/j.1365-294x.2001.01194.x$.

Smouse, P. E., Waples, R. S., and Tworek, J. A. A Genetic Mixture Analysis for use with Incomplete Source Population Data. *Canadian Journal of Fisheries and Aquatic Sciences*, 47(3):620–634, Mar. 1990. doi: $10.1139/f90-070$. Publisher: NRC Research Press.

Stornioli, J. H. F., Goes, C. A. G., Calegari, R. M., dos Santos, R. Z., Giglio, L. M., Foresti, F., Oliveira, C., Penitente, M., Porto-Foresti, F., and Utsunomia, R. The B Chromosomes of Prochilodus lineatus (Teleostei, Characiformes) Are Highly Enriched in Satellite DNAs. *Cells*, 10(6):1527, June 2021. doi: $10.3390/cells10061527$.

Suzuki, F. M., Pires, L. V., and Pompeu, P. S. Passage of fish larvae and eggs through the Funil, Itutinga and Camargos Reservoirs on the upper Rio Grande (Minas Gerais, Brazil). *Neotropical Ichthyology*, 9:617–622, 2011. doi: $10.1590/S1679-62252011000300014$.

Valenzuela-Quiñonez, F. How fisheries management can benefit from genomics? *Briefings in Functional Genomics*, 15(5): 352–357, Sept. 2016. doi: $10.1093/bfgp/elw006$.

van de Schoot, R., Depaoli, S., King, R., Kramer, B., Märtens, K., Tadesse, M. G., Vannucci, M., Gelman, A., Veen, D., Willemsen, J., and Yau, C. Bayesian statistics and modelling. *Nature Reviews Methods Primers*, 1(1):1–26, Jan. 2021. doi: 10.1038/s43586-020-00001-2. Publisher: Nature Publishing Group.

van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M., and Shipley, P. micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4(3):535–538, 2004. doi: 10.1111/j.1471-8286.2004.00684.x. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1471-8286.2004.00684.x.

Vøllestad, L. A. and Primmer, C. R. Understanding local adaptation in a freshwater salmonid fish: evolution of a research programme. *ICES Journal of Marine Science*, 76(6):1404–1414, Dec. 2019. doi: 10.1093/icesjms/fsz037.

Wang, H., Yang, B., Wang, H., and Xiao, H. Impact of different numbers of microsatellite markers on population genetic results using SLAF-seq data for Rhododendron species. *Scientific Reports*, 11(1):8597, Apr. 2021. doi: 10.1038/s41598-021-87945-x. Publisher: Nature Publishing Group.

Wang, J. Maximum-Likelihood Estimation of Admixture Proportions From Genetic Data. *Genetics*, 164(2):747–765, June 2003. doi: 10.1093/genetics/164.2.747.

Wang, J. Fast and accurate population admixture inference from genotype data from a few microsatellites to millions of SNPs. *Heredity*, 129(2):79–92, Aug. 2022. doi: 10.1038/s41437-022-00535-z.

Waples, R. S. Testing for Hardy–Weinberg Proportions: Have We Lost the Plot? *Journal of Heredity*, 106(1):1–19, Jan. 2015. doi: 10.1093/jhered/esu062.

Wenne, R. Microsatellites as Molecular Markers with Applications in Exploitation and Conservation of Aquatic Animal Populations. *Genes*, 14(4):808, Mar. 2023. doi: 10.3390/genes14040808.

Wilkes, M. A., Webb, J. A., Pompeu, P. S., Silva, L. G. M., Vowles, A. S., Baker, C. F., Franklin, P., Link, O., Habit, E., and Kemp, P. S. Not just a migration problem: Metapopulations, habitat shifts, and gene flow are also important for fishway science and management. *River Research and Applications*, 35(10):1688–1696, Dec. 2019. doi: 10.1002/rra.3320.

Wingen, L. U., Brown, J. K. M., and Shaw, M. W. The population genetic structure of clonal organisms generated by exponentially bounded and fat-tailed dispersal. *Genetics*, 177(1):435–448, Sept. 2007. doi: 10.1534/genetics.107.077206.

Xu, T., Moore, I. D., and Gallant, J. C. Fractals, fractal dimensions and landscapes — a review. *Geomorphology*, 8(4):245–262, Dec. 1993. doi: 10.1016/0169-555X(93)90022-T.

Yazbeck, G. M. and Kalapothakis, E. Isolation and characterization of microsatellite DNA in the piracema fish Prochilodus lineatus (Characiformes). *Genetics and molecular research: GMR*, 6(4):1026–1034, Oct. 2007.

Yazbeck, G. M., Oliveira, R. S., Ribeiro, J. M., Graciano, R. D., Santos, R. P., Carmo, F. M. S., and Lavenier, D. A broad genomic panel of microsatellite loci from Brycon orbignyanus (Characiformes: Bryconidae) an endangered migratory Neotropical fish. *Scientific Reports*, 8(1):8511, May 2018. doi: 10.1038/s41598-018-26623-x. Publisher: Nature Publishing Group.

Yazbeck, G. M., Graciano, R. C. D., Santos, R. P., and Oliveira, R. S. Let them eat cake: when the small aims at being LARGE or the empowering effects of bioinformatics in NGS wonderland. *Journal of Information and Data Management*, 15(1):45–50, Feb. 2024. doi: 10.5753/jidm.2024.2626. Number: 1.

Yepes-Blandón, J. A., Bian, C., Benítez-Galeano, M. J., Aristizabal-Regino, J. L., Estrada-Posada, A. L., Mir, D., Vásquez-Machado, G., Atencio-García, V. J., Shi, Q., and Rodríguez-Osorio, N. Draft genome assembly for the colombian freshwater bocachico fish, Prochilodus magdalenae. *Frontiers in Genetics*, 13:989788, 2022. doi: 10.3389/fgene.2022.989788.

Zellhuber, A. Environmental Policy in Brazil. Tensions Between Conservation and the Ideology of Growth. In de la Fontaine, D. and Stehnken, T., editors, *The Political System of Brazil*, pages 329–350. Springer, Berlin, Heidelberg, 2016. ISBN 978-3-642-40023-0. doi: 10.1007/978-3-642-40023-0_18. URL https://doi.org/10.1007/978-3-642-40023-0_18.

Supporting Information

Supplementary File 1: Microsatellite DNA sequences and primers presented in the plain text format (txt) for *Prochilodus lineatus*. The order of columns is "Locus"; "Original fasta label"; "Motif"; "Repeats"; "5'-flank"; "3'-flank"; "F-primer"; "F-Tm"; "R-Primer"; "R-Tm"; "PCR product"; "Product length"; "Location"; "Scaffold or contig label in the newly presented assembly, when mapped". This file is available at fisgshare under doi: 10.6084/m9.figshare.26848654. https://doi.org/10.6084/m9.figshare.26848654.

Supplementary File 2: List of DNA sequences presented in the fasta format for contigs from the original assembly from where microsatellite *loci* for *Prochilodus lineatus* were originally characterized. This file is available at fisgshare under doi: 10.6084/m9.figshare.26848015 https://doi.org/10.6084/m9.figshare.26848015.

Supplementary File 3: *Prochilodus lineatus* population samples, fish IDs and its rearrangements according to 1st, 2nd, and 3rd order MLE defined clusters. This file is available at fisgshare under doi: 10.6084/m9.figshare.26871031. https://doi.org/10.6084/m9.figshare.26871031.

Supplementary File 4: Supplementary Tables, including log likelihood values of the clustering analysis for panmixia and genetic diversity estimates for *Prochilodus lineatus* MLE defined clusters from the 1st, 2nd, and 3rd orders of hierarchical analysis. This file is available at fisgshare under doi: 10.6084/m9.figshare.27020857. https://doi.org/10.6084/m9.figshare.27020857.

Supplementary File 5: Hardy-Weinberg exact tests for MLE defined clusters of *Prochilodus lineatus*, from the 1st, 2nd, and 3rd orders of hierarchical analysis. This file is available at fisgshare under doi: 10.6084/m9.figshare.27020869. https://doi.org/10.6084/m9.figshare.27020869.

Supplementary Figures 1 – 12: Supplementary Figures 1 through 12 are presented in a single portable document format (pdf) file, with respective captions. This file is available at figshare under doi: 10.6084/m9.figshare.26889478 https://doi.org/10.6084/m9.figshare.26889478.