Short-term exposure to Cook, Vitrolife, and KSOM media shapes early calcium oscillations in ICSI-fertilized mouse oocytes and impacts adult phenotype

Bernadette Banrezes[1,2], Thierry Sainte Beuve[1,2], Anne Frambourg[1,2] and Alice Jouneau[1,2], [1] Université Paris-Saclay, UVSQ, INRAE, BREED, 78350, Jouy-en-Josas, France, [2] Ecole Nationale Vétérinaire d'Alfort, BREED, 94700, Maisons-Alfort, France, Corresponding Author:(bernadette.banrezes@inrae.fr)

Abstract

The influence of culture media used during in vitro fertilization (IVF) on the offspring phenotype remains controversial and not subject to consensus. However, the specific effects of short exposure time and duration immediately after fertilization remain underexplored. In this study, we investigated Ca^{2+} oscillatory responses in mouse oocytes fertilized by intracytoplasmic sperm injection (ICSI) as an indicator of early metabolic activity. We evaluated the long-term effects of a short but critical 4-hour exposure to three different culture media: Cook and Vitrolife - commonly used in human IVF - and KSOM, a standard mouse embryo culture medium. Our results indicate that culture media significantly modulate Ca^{2+} oscillations during oocyte activation. ICSI-fertilized oocytes cultured in Cook and Vitrolife exhibited fewer oscillations, lower frequency, and reduced inter-individual variability compared to KSOM, which produced a more diverse oscillation pattern. Notably, these early differences correlated with long-term developmental outcomes: females derived from Cook and Vitrolife embryo cultures exhibited were heavier throughout their growth period and had larger adult organ sizes compared to those derived from KSOM. These findings highlight the critical influence of this short post-fertilization window in shaping Ca^{2+} dynamics, which serve as biomarkers of metabolic variation and ultimately influence adult phenotype. Further research is needed to explore additional IVF media and refine culture conditions to better understand the metabolic pathways linking fertilization environment to offspring development. Optimization of embryo culture protocols in assisted reproductive technologies may improve developmental outcomes in IVF.

Keywords: IVF, ICSI, calcium, oscillation, culture medium, embryo, development

1. Introduction

The number of assisted reproductive technology (ART) treatment cycles continues to increase, reflecting growing reliance on in vitro fertilization (IVF) and related techniques [1], [2]. Among IVF techniques, intracytoplasmic sperm injection (ICSI) has become the dominant method, accounting for 72.7% of procedures in Europe [1] and 75.7% in the United States [3]. Given the widespread use of ICSI, it is essential to understand how the

different culture media used immediately after fertilization can influence the development of ICSI-fertilized oocytes.

ICSI involves the direct injection of a single sperm into an oocyte, triggering a cascade of Ca^{2+} oscillations that activate mitochondrial metabolism and stimulate embryo development [4], [5]. As mitochondrial function is critical during early embryogenesis, the surrounding culture environment may play a key role in modulating these metabolic processes. The composition of the embryo culture medium, as well as the duration of exposure, may influence implantation potential and long-term developmental outcomes [6], [7].

Despite the recognized importance of embryo culture conditions, their precise role in shaping long-term development remains debated. Several human studies suggest that differences in the composition of embryonic culture media are associated with variations in birth weight, two-year weight, and up to nine years of age in children conceived by ART [8], [9], [10], [11]. However, other studies report no significant association [12], [13], [14]. These conflicting findings suggest that additional factors—such as differences in medium formulation, fertilization technique, and patient variability—may contribute to outcome variability, underscoring the need for standardized ART laboratory protocols.

In animal models, the composition of culture media has been shown to influence metabolic activity, modulate developmental trajectories, and even shape adult body weight [15], [16], [17], [18], [19], [20]. However, one of the main challenges in evaluating the functional impact of commercial embryo culture media is the lack of transparency regarding their exact chemical composition. Relevant studies have identified significant differences in the concentrations of more than 39 chemical components across sixteen commercial human embryo culture media [21], [22]. A more recent analysis of fifty-six human embryo culture media [23] has further raised concerns about how this variability may influence embryonic responses and long-term developmental potential. [8]

Before being marketed, embryo culture media undergo safety testing, such as the Mouse Embryo Assay (MEA), which ensures that at least 80% of mouse embryos fertilized in vivo reach the blastocyst stage [24]. However, the MEA has notable limitations: it does not assess ICSI-fertilized oocytes or examine the long-term effects of culture media on embryonic development. Given the increasing reliance on ICSI, it is crucial to evaluate how post-fertilization culture conditions influence both early embryonic response and long-term phenotypic outcomes. A key determinant during this critical post-fertilization period is the regulation of mitochondrial activation and genome reprogramming by Ca^{2+} oscillations [4], [25]. These oscillations, which persist for approximately four hours before ceasing at pronucleus (PN) formation, serve as a real-time indicator of how embryos respond to their culture media formulation. These findings emphasize the need to integrate alternative assessment methods into the MEA, incorporating both early metabolic markers and long-term phenotypic evaluations to improve ART outcome prediction.

To address these knowledge gaps, we investigated how different ART culture media affect the initial metabolic responses of mouse oocytes after ICSI and their long-term developmental effects. Specifically, we compared two widely used ART media - Cook and Vitrolife - one of which has been associated in humans with differences in birth weight,

weight at two years old, and even up to nine years of age in children conceived by ART [8], [9], [10], [11]. As a reference, we used KSOM, a well-established mouse embryo culture medium [26].

We first analyzed Ca^{2+} oscillation patterns in ICSI-fertilized oocytes cultured for four hours in these three culture media. We then evaluated whether these early metabolic responses correlated with in vivo developmental potential following embryo transfer. Key parameters included Ca^{2+} oscillation dynamics, embryonic development rates, postnatal growth and organ and tissue weights.

By integrating these analyses, we aimed to determine whether early Ca^{2+} responses in ICSI-fertilized oocytes could serve as predictive markers of long-term developmental potential. This study provides new insights into how human embryo culture media composition influences a critical four-hour post-fertilization window and its lasting effects on offspring phenotype.

2. Materials and methods

2.1. Ethics statement

Animal experiments were conducted in strict adherence to the Code for Methods and Welfare Considerations in Behavioral Research (Directive 2010/63/EU). The study received approval from the French Ministry of Higher Education, Research, and Innovation, as well as the local Ethics Committee for Animal Experiments at the author's institution, under permit numbers #12-165 and #16-22 INRAE.

2.2. Gametes production and collection

Mature oocytes were recovered from superovulated B6CBAF1/JRj female mice as described previously in Ozil *et al.* [17]. Metaphase II oocytes were collected in a Petri dish containing hyaluronidase (90 i.u. ml^{-1}) in Hepes-buffered KSOM medium [26] to remove cumulus cells. After washing and selection, the oocytes were placed in a drop of KSOM medium under mineral oil and incubated in a 5% CO_2 atmosphere at 37°C. Spermatozoa from B6CBAF1/JRj males were extracted from the caudal epididymis on each day of the experiment and suspended in KSOM medium [17].

2.3. Culture media formulation

The *in vitro* culture was carried out either in three different media. Cook and Vitrolife media were obtained from their respective suppliers, Cook Medical (K-SICM) and Vitrolife (G1™ Version 5 PLUS). The compositions of these two commercial media were analysed for concentrations by Morbeck *et al.* [21] and presented in Tables 1&2. The KSOM was made in the laboratory [26]. All media contained proteins, human serum albumin (HSA) already mixed in the Cook and Vitrolife, and bovine serum albumin (BSA) added to the KSOM (see Table 1). The KSOM contains no amino acids (see Table 2). All inorganic and organic components were sourced from Sigma Chemical Company.

#Table 1: Comparison of electrolytes, carbohydrates and proteins in embryo culture media.

Compounds	Cook	Vitrolife	KSOM
Calcium (Ca) (mM)	1.1	1.1	1.7
Phosphate (mM)	0.2	0.3	0.35
Potassium (mM)	4.8	5.8	2.85
Chloride (mM)	122	124	99.2
Sodium (mM)	140	152	130
Magnesium (Mg) (mM)	1.5	1.7	0.2
Mg/Ca ratio	1.4	1.5	0.1
Glucose (mM)	0.3	0.5	0.2
Citrate (mM)	0	0.08	0
Lactate (L) (mM)	1.8	10.8	10
Pyruvate (P) (mM)	0.36	0.3	0.2
L/P ratio	5	36	50
Proteins (mg/ml)	5	5	1

Source: Morbeck et al. (2014) for Cook and Vitrolife; Lawitts and Biggers (1993) for KSOM.

#Table 2: Comparison of amino acids in embryo culture media.

Compounds	Cook	Vitrolife	KSOM
Essential (µM)			
Arg	25	0	0
Cys	2	0	0
His	8	0	0
Ile	17	0	0
Leu	18	0	0
Lys	18	0	0
Met	4	0	0
Phe	8	0	0
Thr	18	0	0
Trp	2	0	0
Tyr	12	0	0
Val	17	0	0
Nonessential (µM)			
Ala	135	148	0
Asn	88	126	0
Asp	81	0	0
Glu	90	0	0
Gln	30	0	0
Gly	6647	135	0
Pro	85	112	0
Ser	92	127	0
Tau	6489	131	0

Source: Morbeck et al. (2014) for Cook and Vitrolife; Lawitts and Biggers (1993) for KSOM.

2.4.   Intra-cytoplasmic sperm injection

Oocytes were fertilized via ICSI at room temperature using a NIKON TMD microscope, following the protocol described by Ozil et al. [17]. After sperm head injection, the plasma membrane of the oocyte was mechanically compressed against the zona pellucida to ensure sealing and prevent ion influx during membrane repair. ICSI-fertilized oocytes were immediately placed into 40-µL drops of either Cook, Vitrolife or KSOM at 37°C and cultured for 4 hours. After this short different incubation medium, eggs with two pronuclei (2PNs) were further cultured in KSOM until they were transferred into pseudo-pregnant recipients (Figure 1).

#**Figure 1** Schematic experimental design. Oocytes were fertilized by ICSI with Ca^{2+} dye for Ca^{2+} recording or without dye for in vitro culture. After ICSI with dye, Ca^{2+} responses were recorded in a microfluidic device until Ca^{2+} oscillations ceased. Eggs were examined for the presence of 2PNs. Ca^{2+} recordings were then analyzed. After ICSI without dye, ICSI-fertilized oocytes are immediately placed in one of the three culture media and cultured for 4 hours at 37°C in the incubator. Eggs with 2PNs were transferred into female recipients at the 1-cell stage as a priority or at 2-cell stage, depending on the availability of recipients. The survival and growth rate of the animals was recorded until 8 weeks of age.

2.5. Ca^{2+} signal recording

For Ca^{2+} signal recordings, sperm injections were performed in ICSI medium containing 500 µM Fura-2 dextran. After injection, two ICSI-fertilized oocytes were placed in a microfluidic chamber mounted on a Nikon TE2000 inverted microscope equipped with a Fluor 40× oil immersion objective and an EMCCD Photometrics camera. The ICSI-fertilized oocytes were first washed with a rapid flow of KSOM medium at approximately 10 µL s^{-1} to remove residual ICSI medium, followed by stabilization at a slower medium flow rate of 1.4 µL s^{-1} to establish baseline fluorescence measurements. The KSOM medium was then replaced with one of the three experimental culture media. Intracellular Ca^{2+} signals were monitored using fluorescence excitation provided by a monochromator at alternating wavelengths of 340 nm and 380 nm, with emission recorded at 510 nm. The intracellular Ca^{2+} concentration was expressed as the fluorescence ratio (F340/F380) and recorded at a sampling frequency of 0.5 Hz, beginning 1 min after the ICSI-fertilized oocytes were placed in the microfluidic chamber and continuing until the Ca^{2+} oscillations ceased. Stable experimental conditions were maintained during recording by a constant medium flow, a

temperature of 37°C, and regulated gas concentrations. Upon completion of Ca^{2+} recordings, eggs were observed to confirm PN formation.

2.6. Embryo transfer

Female mice (RjOrl:SWISS) were used as recipients. These females were mated with vasectomized males, and on the morning of the experimental day, they were examined for the presence of a vaginal plug. Eggs were then transferred in groups of 8 into the left oviduct of the recipient females, either at the one-cell stage or the two-cell stage depending on the availability of recipients.

2.7. Postnatal phenotype

The postnatal phenotype of the offspring was assessed through weight monitoring, with the animals being weighed weekly starting from the first week after birth. At the 4th week, the pups were weaned, and males and females were separated, with groups of up to five animals housed in individual cages. At 8 weeks of age, the animals were euthanized by cervical dislocation. The liver, kidneys, heart, lungs, brain, and fat were then dissected and weighed.

2.8. Statistical analysis

Statistical analyses were performed using R software (version 4.3.1; https://www.R-project.org). Figures and graphs were generated using the ggplot2 package (version 3.5.1; https://ggplot2.tidyverse.org). Group differences were evaluated with the Kruskal-Wallis test, a non-parametric method appropriate for comparing multiple independent groups. Pairwise comparisons between groups were conducted using the Dunn test, with Bonferroni correction applied to account for multiple comparisons. Additionally, for Table 3, the chi-square (χ^2) test was used to analyze contingency tables and compare proportions as specified in the legend. Most of the results derived from KSOM media have been previously published by Ozil et al. [17].

3. Results

3.1. Impact of culture media composition on Ca²⁺ responses in mouse ICSI-fertilized oocytes

The composition of three media used for in vitro embryo culture was compared (Table 1). The media differ primarily in the ratio of the two main energy sources for early embryos, lactate and pyruvate, as well as in their amino acid content. Specifically, Cook contains both essential and non-essential amino acids, while Vitrolife contains only essential amino acids, and KSOM lacks amino acids entirely. In addition, potassium and magnesium concentrations are higher in Cook and Vitrolife than in KSOM.

To assess the impact of these different media on Ca^{2+} responses in oocytes fertilized by ICSI, we evaluated several key parameters. These parameters included the Ca^{2+} oscillation

patterns observed in vitro, the total number of calcium peaks over four hours, the duration of each Ca²⁺ oscillation event, and its frequency over one hour.

Typical individual Ca²⁺ responses following oocyte fertilization in Cook, Vitrolife or KSOM media are shown in Figure 2. Time zero corresponds to the start of calcium recording in the microfluidic chamber, initiated 2-3 minutes after ICSI. In each response, the first oscillation was characterized by a series of super-oscillations and lasted longer than subsequent ones, as described by Ozil *et al.* [17]. This initial oscillation was followed by repeated oscillations that could be recorded for several hours. In particular, the number of Ca²⁺ oscillations (Ca²⁺ peaks) was lower in the Cook and Vitrolife media compared to the KSOM medium (Figure 2).

#Figure 2 Representative individual Ca²⁺ responses in ICSI-fertilized oocytes cultured in three media for 4 hours. Typical Ca²⁺ responses are shown in Cook (yellow with 6 Ca²⁺oscillations), Vitrolife (blue with 5 Ca²⁺oscillations), and KSOM (green with 16

Ca^{2+}oscillations) media. Time zero corresponds to the start of Ca^{2+} recording in the microfluidic chamber, initiated within 2-3 minutes following ICSI.

Compilation of multiple recordings revealed that the mean number of Ca^{2+} oscillations in ICSI-fertilized oocytes was three times lower in Cook or Vitrolife than in KSOM, 5.6 ± 1.6 and 5.1 ± 1.6 vs. 15.6 ± 5.9, respectively (P < 0.05; Figure 3A). There was no difference between Cook and Vitrolife. The dispersion of the number of oscillations is lower in Cook or Vitrolife compared to the KSOM medium (Figure 3A). Furthermore, the mean number of Ca^{2+} oscillations within the first hour was twice lower in Cook or Vitrolife than in KSOM medium, 3.1 ± 0.8 vs. 6.4 ± 2.4, respectively (P < 0.05; Figure 3B). This significant difference was even more pronounced over the following hours: threefold in the second and third hours, and sevenfold in the fourth hour (P < 0.05; Figure 3C through 3E).

#Figure 3 Number of Ca²⁺ oscillations in ICSI-fertilized oocytes cultured in three media. Figure 3A shows the mean number of Ca²⁺ responses for ICSI-fertilized oocytes cultured for 4 hours in Cook, Vitrolife or KSOM media. Each black dot represents a complete Ca²⁺ record, from the initial Ca²⁺ peak to the cessation of oscillations. Figures 3B through 3E show the mean number of Ca²⁺ responses from hour 1 through hour 4 for ICSI-fertilized oocytes cultured in Cook, Vitrolife, or KSOM media. The mean number of responses for each medium is indicated by a colored square, and the standard deviation is indicated by the error bars in red. P-values obtained by Dunn's post-hoc test (with Bonferroni correction) are shown.

In addition to the lower number of oscillations in Cook and Vitrolife compared to the other medium, the total duration of the ICSI-fertilized oocyte response was also reduced, as was the frequency of oscillations (Figure 4). The mean durations in Cook and Vitrolife were 2 hours, 26 minutes, and 59 seconds and 2 hours, 5 minutes, and 5 seconds, respectively, compared to 3 hours, 17 minutes, and 46 seconds for KSOM. The mean frequencies of Ca^{2+} responses in Cook and Vitrolife were 4.7 ± 1.2 and 2.5 ± 1.0, respectively, vs. 2.7 ± 1.0 in KSOM. However, there was no difference between Cook and Vitrolife in either duration or frequency.

#Figure 4 Duration and frequency of Ca^{2+} oscillations in ICSI-fertilized oocytes cultured in three media. 4A shows the duration and 4B the frequency of Ca^{2+} responses for ICSI-fertilized oocytes cultured in Cook, Vitrolife, or KSOM media. Each black dot represents a complete Ca^{2+} record, from the initial Ca^{2+} peak to the cessation of oscillations. The mean duration of responses for each medium is indicated by a colored square, with the standard deviation represented by the error bars in red. P-values obtained by Dunn's post-hoc test (with Bonferroni correction) are shown.

3.2. Development to term and postnatal growth

After ICSI, embryos were cultured in the three different media for four hours until 2PNs were formed. They were then transferred to recipient females (Table 3). The rate of term development was 31 to 40%, with no statistical difference between the three media. A few pups died during the first week. Thereafter, the growth of male and female pups was followed through week 8 by measuring their weight once a week.

The rates of pregnancy, survival to term, and pup survival were not statistically different among the three groups according to the chi-squared test. In addition, mean litter size at birth was not significantly different between groups (Table 3).

Media	No. eggs transferred	No. pregnant/No. recipient	No. pups from eggs transferred in pregnant	No. live pups	Litter size	Ratio F/M
Cook	184	14/23 (61%)	35/112(31%)	32/35(91%)	2.6+1.8	1.00 (16/16)
KSOM	200	16/25 (64%)	42/128(33%)	40/42(95%)	3.3+2.6	0.86 (18/21)
Vitrolife	152	10/19 (53%)	32/80(40%)	27/32(84%)	2.7+1.7	0.93 (13/14)

#Table 3: Development to term of embryos after ICSI eggs cultured in three different media.

3.3. Postnatal growth

3.3.1. *Growth differences in pups from ICSI eggs cultured in KSOM compared to Cook and Vitrolife media*

Pups were weighed one week after birth and then for eight weeks as the mice grew to adulthood. No differences were observed between the mean growth curves of 'Cook' and 'Vitrolife' females and males. The mean growth curve of 'Cook' females was significantly higher than that of 'KSOM' females at every time point ($P < 0.001$; Figure 5A), although there were two individual growth curves that were very low in the Cook medium (Supplementary Figure 1A). The mean growth curve of 'Vitrolife' females was significantly higher than that of 'KSOM' females from week 2 to week 8 ($P < 0.05$; Figure 5A). The same significant difference is observed for males, but with a less pronounced effect, with 'KSOM' males having a lower weight than those in the other two media at most time points, except for week 1 in Cook and weeks 3-5 in Vitrolife ($P < 0.05$; Figure 5B). There was an individual growth curve for the male that was very low in the Cook medium (Supplementary Figure 1B).

#**Figure 5** Postnatal growth of pups up to 8 weeks of age. 5A and 5B show the mean growth curve with SD of females (A) and males (B). Cook's growth profile is shown in the yellow line, Vitrolife in the blue line, and KSOM in the green line. Pairwise comparison of animal weight in different media from 1 to 8 weeks of age. Statistical analyses were performed separately for females and males using Dunn's post-hoc test, with Bonferroni correction applied to all pairwise comparisons for each week. The p-values resulting from these comparisons are indicated by different labels. Comparison KSOM - Vitrolife: ° $p \leq 0.05$, °° $p \leq 0.01$, °°° $p \leq 0.001$. Comparison Cook - Vitrolife : ¤ $p \leq 0.05$, ¤¤ $p \leq 0.01$, ¤¤¤ $p \leq 0.001$. Comparison Cook - KSOM : * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$.

3.4.   Organ and tissue weights of mice at the 8th week of age

3.4.1. *Differences in specific organ-to-body weight ratios in female offspring*

At 8 weeks of age, Cook and Vitrolife female mice had significantly higher organ weights, including brain, heart, kidney, liver, and lung, compared to KSOM female mice, as shown in Supplementary Figure 2. They also had greater fat deposition around the kidney and gonads. After normalization for body weight, Cook and Vitrolife females had a higher kidney/body weight ratio than KSOM females. Cook females had a higher white fat to body weight ratio and a lower brain to body weight ratio than KSOM females. Vitrolife females had a higher liver-to-body weight ratio than KSOM females, as shown in Figure 6.

#Figure 6 Mean organ/body weight ratios of female offspring at 8 weeks of age. Box plots illustrate the distribution of organ and tissue weight ratios for each group. The red square in each box represents the mean ratio, while jittered round colored dots indicate individual data points. Statistical significance is indicated by horizontal brackets and p-values, with significant differences (p < 0.05) indicated and non significant comparisons marked "ns".

3.4.2. *No differences in specific organ-to-Body weight ratios among male offspring*

At 8 weeks of age, Cook and Vitrolife male mice had significantly higher absolute organ weights for brain and heart compared to KSOM male mice, as shown in Supplementary Figure 3. However, only Cook male mice had a significantly higher absolute kidney weight compared to KSOM. After normalization for body weight, no significant differences in organ-to-body weight ratios were observed between the male groups, as shown in Figure 7.

#**Figure 7** Mean organ-to-body weight ratios of male offspring at 8 weeks of age.

4. Discussion

Sperm entry during fertilization triggers a rapid increase in intracellular free Ca^{2+} concentration in the oocyte. This Ca^{2+} spike initiates a cascade of physiological and biochemical changes that activate the developmental program of the early embryo [5]. Our previous studies demonstrated that the composition of two mouse embryo culture media,

M16 and KSOM—both containing the same chemical components in different proportions—modulates the pattern of calcium oscillations, leading to long-term consequences on development [17].

Despite the increasing number of children born worldwide through ICSI, the impact of ART culture media composition on the metabolic activity of ICSI-fertilized oocytes and its long-term effects after embryo transfer remain largely unexplored in mice. Investigating these interactions at the earliest developmental stage immediately after fertilization and over a brief 4-hour period, corresponding to Ca^{2+} oscillations and PN formation could provide new insights into how culture media influence embryonic metabolism. These findings could ultimately contribute to optimizing assisted reproductive technology protocols.

Determining whether culture conditions at the time of fertilization influence phenotypic characteristics, such as body weight, is crucial. If proven, this would support the hypothesis that the embryonic environment from fertilization onward induces metabolic changes with long-term consequences, aligning with the Developmental Origins of Health and Disease (DOHaD) theory [27].

In this study, we compared three culture media, including two commonly used in ART, and identified distinct Ca^{2+} response patterns immediately after fertilization. These differences correlated with variations in postnatal growth and organ weight. Specifically, our findings reveal that the two ART media induce a different short-term response and long-term developmental outcomes compared to KSOM.

Ca^{2+} response

Our results confirm and extend previous findings in mice [17], showing that sperm-induced Ca^{2+} oscillations are more efficiently generated in the presence of a low Mg^{2+}/high Ca^{2+} culture medium, such as KSOM. Collectively, these data indicate that the external $[Mg^{2+}]/[Ca^{2+}]$ ratio plays a key role in shaping the pattern of Ca^{2+} oscillations by regulating Ca^{2+} influx. Modifying this ratio using different culture media—KSOM, Cook, and Vitrolife—during egg activation significantly impacts the Ca^{2+} response. KSOM, containing 0.2 mM Mg^{2+} and 1.7 mM Ca^{2+}, promotes a distinct oscillatory pattern compared to Cook and Vitrolife, which contain 1.5 mM and 1.7 mM Mg^{2+} with 1.1 mM Ca^{2+}, respectively [21]. These results suggest that Mg^{2+} levels antagonize Ca^{2+} influx, thereby modulating oscillation frequency and amplitude.

During ICSI, sperm-specific phospholipase C zeta (PLCζ) hydrolyzes phosphatidylinositol 4,5-bisphosphate, generating inositol trisphosphate (IP_3) and diacylglycerol [28]. IP_3 then binds to its receptors on the endoplasmic reticulum, triggering the release of stored Ca^{2+} into the cytosol, initiating a series of oscillations lasting several hours [29]. However, sustaining these oscillations requires Ca^{2+} influx across the oocyte plasma membrane [30], primarily mediated by CaV3.2 and TRPM7 channels [31]. Notably, TRPM7 also plays a role in Mg^{2+} homeostasis, potentially competing with Ca^{2+} influx [32].

Our study confirms that higher Mg^{2+} concentrations reduce both the frequency and variability of Ca^{2+} oscillations, whereas lower Mg^{2+} levels permit a broader oscillatory

range with greater interindividual variability. This pattern is evident when comparing Cook/Vitrolife with KSOM.

Another key difference between KSOM and the ART media (Cook/Vitrolife) is the presence of amino acids and higher protein content in Cook/Vitrolife. However, experiments using Global medium (which contains amino acids and higher protein levels but a low $[Mg^{2+}]/[Ca^{2+}]$ ratio: 0.2/1.9 mM) confirmed that Ca^{2+} oscillation differences persisted, suggesting that the Mg^{2+}/Ca^{2+} balance remains the primary determinant of oscillatory dynamics (unpublished data).

Furthermore, variations in lactate and pyruvate concentrations, as well as the lactate/pyruvate ratio, differentiate KSOM from Cook/Vitrolife. However, despite Global medium having a lower lactate/pyruvate ratio [21], it still induces a higher Ca^{2+} response than Vitrolife, reinforcing the conclusion that ionic composition —particularly the Mg^{2+}/Ca^{2+} ratio— dominates the regulation of Ca^{2+} oscillations at this stage.

Thus, modulating extracellular Mg^{2+} concentration directly influences the dynamics of Ca^{2+} oscillations post-fertilization. Careful control of this balance in embryo culture media is therefore crucial for optimizing ART protocols [17].

Growth Rates and Organ/Tissue Weights

Since Ca^{2+} oscillations play a key role in initiating embryonic development, alterations in their pattern could have long-term consequences. To explore this, we analyzed postnatal growth and organ weights in mice cultured in different media.

Our results reveal significant effects of the culture media on growth in both males and females, aligning with previous mouse studies and human IVF reports. Even a brief exposure to specific media between fertilization and PN formation can influence postnatal development, from birth to adulthood.

Notably, females from the Cook and Vitrolife groups exhibited increased organ and tissue weights, while in males, differences were restricted to three organs: the brain, heart, and kidney. This suggests a sex-specific impact of culture media on organ development. After normalization by body weight, females from Cook and Vitrolife showed an increased kidney and fat ratio, while the brain ratio decreased.

Strikingly, we observed significant differences between Cook, KSOM, and Vitrolife after only 4 hours of culture. These differences persisted into adulthood, affecting both body weight and organ/body weight ratios in female mice. This underscores the potential for even a short exposure to specific culture conditions to induce long-lasting developmental effects.

Similar trends have been reported in humans: embryos cultured in Vitrolife medium for 2–3 days had significantly higher birth weights than those cultured in Cook medium [8]. Follow-up studies confirmed that these differences persisted into early childhood, with children from the Vitrolife group exhibiting higher weight, height, BMI, waist circumference, and truncal adiposity at age nine [10], [11].

These findings reinforce the profound and lasting impact of embryo culture conditions on postnatal growth and metabolic development. Understanding these mechanisms is essential to refine ART protocols and improve long-term health outcomes for individuals conceived through assisted reproduction.

5. Conclusion

This study shows that a 4-hour post-fertilization exposure of ICSI-fertilized oocytes to different culture media influences both early Ca^{2+} signaling and long-term development in mice. Cook and Vitrolife media, with high $[Mg^{2+}]/[Ca^{2+}]$ ratios, generate distinct Ca^{2+} oscillation patterns compared to KSOM, emphasizing the role of ionic balance in oocyte activation. These early differences result in significant phenotypic variations in adulthood, highlighting the critical influence of the short window between ICSI and PN formation. However, species-specific differences in embryonic development and epigenetic regulation limit direct extrapolation to humans. Future research should further investigate the interactions between ionic balance, metabolism, and epigenetic modifications to optimize ART culture conditions and improve long-term outcomes.

Author contributions

B.B. designed research; B.B. and T.SB. performed research; B.B., T.SB and A.F analyzed data; and B.B. and A.J. wrote the paper.

Funding

This research was supported in part by INRAE department PHASE and by agence de la biomedicine (AMP, diagnostic prénatal et diagnostic génétique 2014 'Measurement of the metabolic impact of the culture media used for in vitro fertilization').

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements The authors would like to thank Sophie Calderari for critical reading of the manuscript.They thank the unit of Infectiology of Fishes and Rodents Facility (IERP-UE907, Jouy-en-Josas Research Center: https://doi.org/10.15454/1.5572427140471238E12; member of the National Infrastructure EMERG'IN), for excellent animal care and help with the animal experiments.

Appendix A. Supplementary data The following are the supplementary data to this article:

#**Supplementary Figure 1** Postnatal growth of pups up to 8th week of age. 1A and 1B show individual growth curve of females (A) and males (B). Cook's growth profile is shown in the yellow line, Vitrolife's in the blue line and KSOM's in the green line.

#**Supplementary Figure 2** Supplementary Figure 2: Mean organ and tissue weights of female offspring at 8 weeks of age. Box plots illustrate the distribution of organ and tissue weights for each group. The red square in each box represents the mean organ weight, while jittered round colored dots indicate individual data points. Statistical significance is indicated by horizontal brackets and p-values, with significant differences (p < 0.05) marked and nonsignificant comparisons marked "ns". The figure shows that Cook and Vitrolife females had significantly higher organ weights (brain, heart, kidney, liver, and lung) and greater fat deposition (white fat around the kidney and gonads) compared to KSOM females.

#**Supplementary Figure 3** Supplementary Figure 3. Mean organ and tissue weights of male offspring at 8 weeks of age. The figure shows that while Cook and Vitrolife males had significantly higher brain and heart weights than KSOM males, only Cook males had a significantly higher kidney weight. Additionally, both Cook and Vitrolife males exhibited greater fat deposition (white fat around the kidney and gonads).

References

[1]   J. Smeenk *et al.*, "ART in europe, 2019: Results generated from european registries by ESHRE," *Human Reproduction (Oxford, England)*, vol. 38, no. 12, pp. 2321–2338, Oct. 2023, doi: 10.1093/humrep/dead197.

[2]   S. Sunderam, Y. Zhang, A. Jewett, and D. M. Kissin, "State-Specific Assisted Reproductive Technology Surveillance, United States: 2019 Data Brief," Oct. 2021.

[3]   S. Sunderam *et al.*, "Assisted Reproductive Technology Surveillance United States, 2018," Feb. 2022.

[4]   R. Dumollard, P. Marangos, G. Fitzharris, K. Swann, M. Duchen, and J. Carroll, "Sperm-triggered [Ca2+] oscillations and Ca2+homeostasis in the mouse egg have an absolute requirement for mitochondrial ATP production," *Development*, vol. 131, no. 13, pp. 3057–3067, Jul. 2004, doi: 10.1242/dev.01181.

[5]   Z. Machaty, "The signal that stimulates mammalian embryo development," *Frontiers in Cell and Developmental Biology*, vol. 12, Sep. 2024, doi: 10.3389/fcell.2024.1474009.

[6]   H. Zandstra, A. P. A. Van Montfoort, and J. C. M. Dumoulin, "Does the type of culture medium used influence birthweight of children born after IVF?" *Human Reproduction*, vol. 30, no. 3, pp. 530–542, Mar. 2015, doi: 10.1093/humrep/deu346.

[7]   R. Sciorio and P. Rinaudo, "Culture conditions in the IVF laboratory: state of the ART and possible new directions," *Journal of Assisted Reproduction and Genetics*, vol. 40, no. 11, pp. 2591–2607, Nov. 2023, doi: 10.1007/s10815-023-02934-5.

[8]   J. C. Dumoulin *et al.*, "Effect of in vitro culture of human embryos on birthweight of newborns," *Human Reproduction*, vol. 25, no. 3, pp. 605–612, Mar. 2010, doi: 10.1093/humrep/dep456.

[9]   E. C. Nelissen *et al.*, "Further evidence that culture media affect perinatal outcome: findings after transfer of fresh and cryopreserved embryos," *Human Reproduction*, vol. 27, no. 7, pp. 1966–1976, Jul. 2012, doi: 10.1093/humrep/des145.

[10]   S. H. M. Kleijkers *et al.*, "IVF culture medium affects post-natal weight in humans during the first 2 years of life," *Human Reproduction*, vol. 29, no. 4, pp. 661–669, Apr. 2014, doi: 10.1093/humrep/deu025.

[11]   H. Zandstra *et al.*, "Association of culture medium with growth, weight and cardiovascular development of IVF children at the age of 9 years," *Human Reproduction*, vol. 33, no. 9, pp. 1645–1656, Sep. 2018, doi: 10.1093/humrep/dey246.

[12]   B. Carrasco, M. Boada, I. Rodríguez, B. Coroleu, P. N. Barri, and A. Veiga, "Does culture medium influence offspring birth weight?" *Fertility and Sterility*, vol. 100, no. 5, pp. 1283–1288, Nov. 2013, doi: 10.1016/j.fertnstert.2013.07.006.

[13]   D. Wunder *et al.*, "Effect of embryo culture media on birthweight and length in singleton term infants after IVF-ICSI," *Swiss Medical Weekly*, Oct. 2014, doi: 10.4414/smw.2014.14038.

[14]   C. G. Petersen *et al.*, "Randomized comparison of two commercial culture media (Cook and Vitrolife) for embryo culture after IMSI," *JBRA Assisted Reproduction*, 2019, doi: 10.5935/1518-0557.20180058.

[15]   D. J. Ecker *et al.*, "Long-term effects of culture of preimplantation mouse embryos on behavior," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 6, pp. 1595–1600, Feb. 2004, doi: 10.1073/pnas.0306846101.

[16]   B. Banrezes, T. Sainte-Beuve, E. Canon, R. M. Schultz, J. Cancela, and J.-P. Ozil, "Adult body weight is programmed by a redox-regulated and energy-dependent process during the pronuclear stage in mouse," *PloS One*, vol. 6, no. 12, p. e29388, 2011, doi: 10.1371/journal.pone.0029388.

[17]   J.-P. Ozil, T. Sainte-Beuve, and B. Banrezes, "[Mg2+]o/[Ca2+]o determines Ca2+ response at fertilization: tuning of adult phenotype?" *Reproduction*, vol. 154, no. 5, pp. 675–693, Nov. 2017, doi: 10.1530/REP-16-0057.

[18]   T. P. Fleming *et al.*, "Origins of lifetime health around the time of conception: causes and consequences," *The Lancet*, vol. 391, no. 10132, pp. 1842–1852, May 2018, doi: 10.1016/S0140-6736(18)30312-X.

[19]   A. F. Ermisch *et al.*, "A novel culture medium with reduced nutrient concentrations supports the development and viability of mouse embryos," *Scientific Reports*, vol. 10, no. 1, p. 9263, Jun. 2020, doi: 10.1038/s41598-020-66019-4.

[20]   M. A. Velazquez, A. Idriss, P. Chavatte-Palmer, and T. P. Fleming, "The mammalian preimplantation embryo: Its role in the environmental programming of postnatal health and performance," *Animal Reproduction Science*, vol. 256, p. 107321, Sep. 2023, doi: 10.1016/j.anireprosci.2023.107321.

[21]   D. E. Morbeck, R. L. Krisher, J. R. Herrick, N. A. Baumann, D. Matern, and T. Moyer, "Composition of commercial media used for human embryo culture," *Fertility and Sterility*, vol. 102, no. 3, pp. 759–766.e9, Sep. 2014, doi: 10.1016/j.fertnstert.2014.05.043.

[22]   D. E. Morbeck, N. A. Baumann, and D. Oglesbee, "Composition of single-step media used for human embryo culture," *Fertility and Sterility*, vol. 107, no. 4, pp. 1055–1060.e1, Apr. 2017, doi: 10.1016/j.fertnstert.2017.01.007.

[23]   M. S. Zagers *et al.*, "The composition of commercially available human embryo culture media," *Human Reproduction*, vol. 40, no. 1, pp. 30–40, Jan. 2025, doi: 10.1093/humrep/deae248.

[24]   L. Delaroche *et al.*, "Embryotoxicity testing of IVF disposables: how do manufacturers test?" *Human Reproduction*, vol. 35, no. 2, pp. 283–292, Feb. 2020, doi: 10.1093/humrep/dez277.

[25]   T. Ducibella, R. Schultz, and J. Ozil, "Role of calcium signals in early development," *Seminars in Cell & Developmental Biology*, vol. 17, no. 2, pp. 324–332, Apr. 2006, doi: 10.1016/j.semcdb.2006.02.010.

[26]   J. A. Lawitts and J. D. Biggers, "[9] Culture of preimplantation embryos," vol. 225, Elsevier, 1993, pp. 153–164. doi: 10.1016/0076-6879(93)25012-Q.

[27]   D. J. Barker, "The fetal and infant origins of adult disease." *BMJ*, vol. 301, no. 6761, pp. 1111–1111, Nov. 1990, doi: 10.1136/bmj.301.6761.1111.

[28]   C. M. Saunders *et al.*, "PLCζ: a sperm-specific trigger of Ca2+ oscillations in eggs and embryo development," *Development*, vol. 129, no. 15, pp. 3533–3544, Aug. 2002, doi: 10.1242/dev.129.15.3533.

[29]   K. S. R. Cuthbertson and P. H. Cobbold, "Phorbol ester and sperm activate mouse oocytes by inducing sustained oscillations in cell Ca2+," *Nature*, vol. 316, no. 6028, pp. 541–542, Aug. 1985, doi: 10.1038/316541a0.

[30]   K. Swann, "Sperm-Induced Ca2+ Release in Mammalian Eggs: The Roles of PLCζ, InsP3, and ATP," 2023.

[31]   M. L. Bernhardt *et al.*, "TRPM7 and Ca_V 3.2 channels mediate Ca^{2+} influx required for egg activation at fertilization," *Proceedings of the National Academy of Sciences*, vol. 115, no. 44, Oct. 2018, doi: 10.1073/pnas.1810422115.

[32]   I. Carvacho, G. Ardestani, H. C. Lee, K. McGarvey, R. A. Fissore, and K. Lykke-Hartmann, "TRPM7-like channels are functionally expressed in oocytes and modulate post-fertilization embryo development in mouse," *Scientific Reports*, vol. 6, no. 1, p. 34236, Sep. 2016, doi: 10.1038/srep34236.