1

citrOgen: a synthesis-free polysaccharide and protein antigen-presentation to antibody-induction platform

Joshua L C Wong[#,*], Julia Sanchez-Garrido[#], Jaie Rattle, Jonathan Bradshaw, Vishwas Mishra and Gad Frankel[*]

Department of Life Sciences, Imperial College London, SW7 2AZ, UK

[#]These authors contributed equally to this work

[*]For correspondence:

Electronic mail: joshua.wong@imperial.ac.uk and g.frankel@imperial.ac.uk

Keywords

Antibodies, *Citrobacter rodentium, Klebsiella pneumoniae,* type III secretion system (T3SS), EspO

Abstract

Existing technologies employed to generate antibodies against bacterial polysaccharides and proteins rely on the availability of purified or synthetic antigens. Here we present a genetics-based platform that utilises *Citrobacter rodentium* (CR), an enteric mouse pathogen, to both produce and present complex heterologous polysaccharide and protein antigens during natural infection. As a proof of concept, we use the lipopolysaccharide (O1), capsular polysaccharide (K2) and type 3 fimbrial (T3F) antigens expressed by the WHO critical priority pathogen *Klebsiella pneumoniae* (KP). Following one infection cycle (28 days) CR induces specific IgG antibodies against KPO1, KPK2 and KPT3F. We demonstrate that the antibodies are functional in downstream applications including protection against pathogenic KP challenge, KP capsular serotyping and KP biofilm inhibition. Whilst KP antigens were used as prototypical examples, this modular platform is now readily adaptable to generate antibodies against diverse polysaccharide and complex protein antigens, with basic science, public health and therapeutic applications.

Introduction

The specificity of antibody/antigen (Ab/Ag) binding has been widely leveraged in science, biotechnology, and medicine. Both polyclonal (pAb) and monoclonal (mAb) antibodies are routinely employed across these disciplines and have shaped the face of the modern practice. There is, therefore, significant value in expanding the technologies available to generate new Abs that bind specific Ags of interest.

Abs are employed widely in the field of infectious disease with both diagnostic and, more recently, therapeutic applications. One key use in diagnostics is bacterial serotyping, which has played important roles in pathogen epidemiology. Therapeutic Abs are licensed for both the prevention and treatment of viral and bacterial infections in humans. Palivizumab, a humanised monoclonal IgG Ab that binds Respiratory Syncytial Virus, was the first mAb licensed for the prophylaxis of infectious disease in 1998[1]. Bezlotoxumab, a mAb against the enteric pathogen *C. difficile*, the sole mAb licensed in bacterial disease, was licensed in 2016 for the secondary prevention of *C. difficile* colitis [2]. More recently, during the SARS-CoV-2 pandemic, alongside vaccination, both pAb administration (convalescent sera)[3] and mAbs against spike protein (e.g. bamlanivimab, etesevimab, casirivimab and imdevimab)[4] were used in the prevention and treatment of COVID-19 disease. We currently have limited therapeutic options to combat the antimicrobial resistance (AMR) pandemic and anti-bacterial Abs (α-bAbs) are a promising modality to tackle this threat.

In target-specific approaches α-bAbs binding predetermined Ags are generated and selected for downstream use. Target choice can be informed *a priori* by pathogen epidemiology and genomics, selecting bacterial Ags that are both widespread and conserved across a bacterial species. Ags representing virulence determinants can also be selectively targeted as Ab-binding has the potential to functionally disrupt pathogenic mechanisms[5]. The immunisation of experimental animals or human subjects is a common method to generate target-specific Abs. This results in sera containing pAb secreted from reactive B-cells. These cells can be immortalised or Ab-encoding regions subcloned to

produce mAbs *in vitro*. Irrespective of host species, the generation of such responses is fundamentally

reliant on a sufficient source of conformationally correct immunogen. Generating these immunogens

represents a key bottleneck in target-specific Ab generation[6].

Polysaccharide Ags are diverse and typically composed of variable sugar subunits, each covalently

connected in the correct order by specific glycosidic links. These linkages can be linear or include

branching arrangements that determine the final polysaccharide structure. In isolation,

polysaccharides typically require carrier protein glycoconjugation to generate formulations that are

sufficiently immunogenic[6]. Production of sufficient quantities of protein Ags, which are typically affinity

purified following *de novo* synthesis in cells, is largely restricted to protein monomers and not

applicable to multiprotein complexes. The correct formation of the latter not only requires the

presence of chaperones and signal sequences, directing subunit assembly in the correct subcellular

space, but also necessitates conformationally correct monomers to interact with each other in the

correct stochiometric ratio. Bacterial genomes have compartmentalised the genes required to

synthesise both polysaccharide and multimeric protein Ags into operons, providing cloneable linear

DNA fragments that we exploit in the development of the technology we describe here.

Citrobacter rodentium (CR) causes colitis in mice with an infection strategy mirroring the human

pathogen Enteropathogenic *E. coli* (EPEC). CR attaches intimately to intestinal epithelial cells (IECs)

forming attaching and effacing lesions and actin-rich pedestal-like structures, a prerequisite for

productive infection and colitis[7]. A major virulence factor enabling CR infection is the type 3 secretion

system (T3SS)[8] which translocates 31 effector proteins into IECs, facilitating CR adherence and

proliferation [9]. The resolution of CR infection coincides with the onset of IgG secretion[10,11]. However,

until this adaptive immune response clears the infection, CR maintains high mucosal-associated

burdens, inducing significant local inflammation with associated immune cell recruitment[9]. We

hypothesised that this natural host-pathogen context represents an evolutionarily pre-optimised

environment to present alternative (heterologous) Ags to the mouse host, thereby generating mature

target-specific Ab responses. This hypothesis forms the basis of our platform which we term citrOgen.

citrOgen is a flexible and modular *in vivo* Ag presentation platform co-opting CR to present

heterologous Ags. It represents a single unifying genetics-based system that can be used to generate

functional target-specific α-bAb responses to divergent Ags in mice. We selected three structurally

diverse and validated Ags from the WHO critical priority pathogen *Klebsiella pneumoniae* (KP) to test

the platform [12]. The choice includes two surface polysaccharides, lipopolysaccharide (LPS – KP O1) and

capsular polysaccharide (CPS – KP K2) and a protein complex, type 3 fimbriae (KP T3F). Substituting

the endogenous CR O152 LPS O-Ag with KP O1 resulted in lower gut colonisation, which we overcome

by overexpressing the CR T3SS effector EspO. Functionally, EspO-augmented α-LPS responses potently

protected mice from sepsis-induced organ failure following KP challenge. We then successfully

generated α-CPS Abs, which we employed to serotype a diverse KP collection of 100 clinical isolates.

Finally, we demonstrate the platforms' ability to raise Abs to a large cell surface filamentous structure,

KP T3F, the regulated expression of which we designed to coincide with CR attachment to host

epithelia. These α-KP T3F Abs effectively blocked T3F-mediated KP biofilm formation. Together these

results demonstrate the ability of the platform to rapidly generate functional target-specific antibodies

against a broad range of major bacterial Ags.

Results

Heterologous KP O1 O-antigen expression causes an in vivo colonisation defect

Our goal was to express diverse heterologous Ags in CR to generate target-specific Abs in mice. As model Ags we chose O1, K2 and T3F, the natural surface Ag configuration encoded by our prototype KP strain ICC8001 (herein KP$_{WT}$). The *rfb* locus typically encodes all the genes required for O-Ag synthesis and export in Gram-negative bacteria[13]. An exception is the KP O1 Ag, where *rfb*-derived O2 Ag is modified by the glycosyltransferase activity of two enzymes, WbbY and WbbZ, which are encoded at a distinct chromosomal position[14]. Moreover, KP employs an ABC-transporter instead of the more common *wzy*-dependent O-Ag export process used by CR$_{WT}$[15].

We started by identifying the *rfb* locus in CR$_{WT}$, which is encoded on the reverse strand and flanked by *galF* (5') and *gnd* (3') genes (Fig 1A). We used *in silico* serotyping to confirm that CR$_{WT}$ encodes an *E. coli*-like O152 Ag[16]. The endogenous O152 genes were replaced with those comprising the KP O2 *rfb* locus (cloned en bloc) by homologous recombination and the *wbbYZ* gene cluster inserted into the CR genome by Tn7 transposon mutagenesis (Fig 1A)[17]. This strain, CR$_{KPO1}$, contains all the genetic determinants required to synthesise and export KP O1 Ag, albeit using a non-native ABC-transporter export pathway. We did not modify the CR gene products that synthesise lipid A or core oligosaccharide (Fig 1B) and, therefore, CR$_{KPO1}$ is predicted to produce a hybrid LPS molecule; the endogenous CR Raetz pathway and CR *rfa* locus producing lipid A and core regions, respectively, with KP O1 Ag attached to the native CR lipooligosaccharide scaffold (Fig 1B). We purified LPS from KP$_{WT}$, CR$_{WT}$ and CR$_{KPO1}$ to confirm O-antigen replacement by Western blot. As anticipated α-O152 pAb sera recognised CR$_{WT}$ but was not reactive against KP$_{WT}$ and CR$_{KPO1}$ (Fig 1C). We then used an α-KP O1 mAb (C13)[18], which bound to both KP$_{WT}$ and CR$_{KPO1}$, confirming heterologous O-antigen expression in CR. The homologous recombination vector employed to substitute the KP O2 locus was designed to include the endogenous 5' CR promoter, ensuring that future O-antigen locus substitutions can be achieved by using the same vector, simply modified with O-antigen genes of choice.

We tested if replacing CR O152 with KP O1 affected infectivity *in vitro* and confirmed that CR_{KPO1}, like

$CR_{WT,}$ induced actin-rich pedestal formation following the infection of mouse fibroblasts cells.

Furthermore, detectable KP O1 Ag expression was restricted to CR_{KPO1} (Supplementary Fig 1A). We

therefore proceeded to infect CD-1 mice with 10^9 colony forming units (CFUs) of CR_{WT} or CR_{KPO1} by oral

gavage (Fig 1D). At 8 days post infection (dpi), when stool CR shedding peaks, faecal CR_{KPO1} CFU counts

were significantly lower than CR_{WT} (Fig 1D). Staining of colonic sections revealed that whilst CR_{KPO1}

maintained heterologous KP O1 expression *in vivo,* there were substantially fewer adhered bacteria

compared to CR_{WT} (Fig. 1E, Supplementary Figure 1B). These data indicated an *in vivo* colonisation

defect imposed by heterologous Ag expression that needed to be addressed to develop an optimal

platform.

EspO overexpression restores CR_{KPO1} colonisation

A mutant in the *Shigella flexneri* T3SS effector *ospE*, a homologue of the CR effector *espO*, results in a

colonisation defect in a guinea pig gut infection model, as OspE suppresses epithelial detachment[19].

We therefore hypothesised that over expression of EspO would promote greater bacterial attachment

to IECs during infection. Indeed, while deletion of *espO* in CR_{WT} showed a tendency towards decreased

proportion of CR-associated IECs, overexpression of EspO increased the proportion from 14.6% to

22.9% (Supplementary Fig 2A-C). We then overexpressed EspO in CR_{KPO1} (CR_{KPO1}+p*espO*) and infected

CD-1 mice; infections with CR_{WT} and CR_{KPO1} were used as controls. Enumeration of faecal CFUs now

revealed no difference between the faecal CFUs between CR_{WT} and CR_{KPO1}+p*espO* at 8 dpi (Fig 2A).

Staining of colonic sections demonstrated higher levels of CR_{KPO1}+p*espO* adhesion to the epithelia

compared to CR_{KPO1}, which maintained heterologous KP O1 expression *in vivo* (Fig 2B). Based on these

results, EspO overexpression (+p*espO*) was included in all subsequent experiments to facilitate CR

colonisation in the context of heterologous Ag expression.

152 **CR_{KPO1}+p*espO* infection elicits KP O1 specific IgG antibodies**

153 We inoculated mice with CR_{KPO1}+p*espO* and CR_{WT} by oral gavage. After CR clearance (c. 28 dpi) we

154 collected blood and assessed α-KP_{WT} IgG antibody titres by enzyme-linked immunosorbent assay

155 (ELISA), using a capture plate coated with heat-killed KP_{WT}. While control CR_{WT} mice sera had negligible

156 α-KP_{WT} IgG binding, this increased significantly in sera from CR_{KPO1}+p*espO*-infected mice, indicating

157 that we have successfully raised α-KP_{WT} Abs (Fig. 2C). To confirm the specificity of the Ab response we

158 used an isogenic rough (absent O-antigen) KP mutant, $KP_{Δrfb}$, and assessed binding to live bacteria by

159 flow cytometry (Fig. 2D). Consistent with the ELISA data, sera from CR_{WT}-infected mice bound to

160 neither KP_{WT} nor $KP_{Δrfb}$. However, sera from CR_{KPO1}+p*espO*-infected mice bound specifically to KP_{WT} and

161 not $KP_{Δrfb}$, confirming the presence of α-KP O1 Abs (Fig 2D, Supplementary Fig 2D). We then assessed

162 the pattern of binding to KP_{WT} by immunofluorescence (IF) microscopy (Fig 2E). This confirmed the

163 same pattern of binding of CR_{KPO1}+p*espO* sera and the reference α-KP O1 Ab C13. No detectable signal

164 was observed in sera from CR_{WT}-infected mice.

165 **Anti-KPO1 IgG Abs are functionally protective against KP pulmonary challenge**

166 We next tested the functionality of the CR_{KPO1}+p*espO*-induced α-KP O1 pAbs by lung challenge with

167 KP_{WT}. Mice that had been infected and cleared CR_{KPO1}+p*espO* were intubated and administered 500

168 CFU of KP_{WT} directly into the lungs. In preliminary experiments with CD-1 mice this KP_{WT} inoculum was

169 found to result in a progressive increase in KP CFUs in the lungs and blood resulting in a severe disease

170 phenotype at 72 h post infection (hpi) (Supplementary Fig 3 A-C). Mice that had been infected by and

171 subsequently cleared CR_{WT} were also challenged as controls. We evaluated infection outcomes by

172 assessing KP_{WT} replication and protection against host organ failure employing markers of

173 haematological, lung and renal injury in line with international consensus[20,21]. At 72 hpi KP_{WT} burdens

174 in the lungs of CR_{WT}-preinfected mice reached a median of c. 10^9 CFUs (Fig 3A). In contrast,

175 CR_{KPO1}+p*espO*-preinfected mice had significantly reduced lung CFUs with KP_{WT} only detectable in the

176 lungs of 30% (5/14) of mice. KP bacteraemia was absent in all CR_{KPO1}+p*espO*-preinfected mice in

177 comparison to 69% (9/13) of CR_{WT}-preinfected control mice (Fig 3B).

We then assessed if the reduction of KP_{WT} burden translated into a positive effect on host physiology.

Only 3 mice (20%, 3/15) in the CR_{KPO1}+p*espO*-preinfected group lost more than 10% of their starting

weight when challenged with KP_{WT}, whereas all mice in the CR_{WT}-preinfected group lost weight after

KP challenge with 86% (12/14) losing more than 10% of their starting weight (Fig 3C). Total white cell

counts, lymphocyte counts and platelets in the blood demonstrated significantly positive

improvements in CR_{KPO1}+p*espO*-preinfected mice compared to CR_{WT}-preinfected animals (Fig 3D-F).

Specifically, the extent of sepsis-induced leucopaenia, lymphopaenia and thrombocytopaenia were

abrogated by the CR_{KPO1}+p*espO* preinfection, while the number of neutrophils did not change between

the mouse groups (Fig 3G). Moreover, the positive acute phase reactants (APR) Serum amyloid A and

C-reactive protein were significantly lower in CR_{KPO1}+p*espO*-preinfected mice compared CR_{WT}-

preinfected animals (Fig 3H&I). This beneficial modulation in APRs translated into higher levels of the

negatively regulated APR, albumin, in CR_{KPO1}+p*espO* preinfected mice (Fig 3J).

We then used three markers to assess lung injury: IL1β and myeloperoxidase (MPO) in lung

homogenate and surfactant protein D (SPD) levels in the serum[20]. This revealed that CR_{KPO1}+p*espO*

preinfection resulted in significant reductions in IL1β and SPD levels compared to CR_{WT} preinfection,

indicating reduced lung inflammation and improved maintenance of the alveolar capillary barrier

respectively (Fig 3K&L). Preinfection with CR_{KPO1}+p*espO* also resulted in significantly reduced

neutrophilic alveolitis, evidenced by reduced lung MPO levels compared to CR_{WT}-preinfected mice (Fig

3M). Finally, we assessed acute kidney injury by measuring serum creatinine and cystatin C levels.

While high levels of creatinine and cystatin C were found in KP_{WT} challenged CR_{WT}-preinfected mice,

improvements in both excretory renal function markers were seen in CR_{KPO1}+p*espO*-preinfected mice

(Fig 3N&O). These data indicate that α-KP O1 Abs raised by citrOgen were protective against KP

challenge, restricting both bacterial replication and dissemination, and preventing KP sepsis-induced

organ failure.

citroGen presentation of heterologous KP capsular polysaccharide antigen

We next explore expanding the citrOgen platform to generate α-CPS Abs. KP_{WT} produces K2 CPS, encoded by a c. 22Kb K2 *cps* locus, lying immediately upstream of the *rfb* locus in the KP_{WT} genome. As upstream of the CR O152 *rfb* locus lie the genes involved in synthesis and export of the extracellular polysaccharide colanic acid (Fig. 4A), we replaced the CR_{WT} genes between *ROD21991* and *galF* (c. 24kb) with the KP KL2 operon from *wzi* to *manB* by homologous recombination, generating strain CR_{KPK2}+p*espO* (Fig. 4A). We chose this insertion site as we anticipate that it will tolerate other *cps* locus insertions in future applications of this technology. This established vector can be used as the template together with *cps* locus of choice (e.g. KP *cps* loci vary from 10-30kb and include internal promoters[22]).

We infected mouse fibroblasts to evaluate if the expression K2 capsule impacted on CR T3SS functionality. CR_{KPK2} formed actin-rich pedestals and heterologous K2 expression was confirmed by staining with reference K2 antiserum (Supplementary Fig 4A). We then infected CD-1 mice by oral gavage with CR_{KPK2}+p*espO,* or CR_{WT} as a control, which were shed at comparable levels (Fig 4B). Following CR_{KPK2}+p*espO* and CR_{WT} clearance, we collected blood samples and α-KP IgG levels were assessed by ELISA, using KP_{WT} and an acapsular *wcaJ* mutant, $KP_{\Delta wcaJ}$. No binding to either KP_{WT} or $KP_{\Delta wcaJ}$ was observed with sera from CR_{WT}-infected mice. In contrast, sera from mice infected with CR_{KPK2}+p*espO* bound to KP_{WT} but not to $KP_{\Delta wcaJ}$ (Fig 4C). This indicated that KP K2 was successfully presented by the citrOgen platform *in vivo* and that α-KP IgG was specifically raised against the heterologous KP K2 antigen.

KP strains demonstrate broad diversity in CPS Ags with over 147 described to date in circulating strains[22]. However, despite their key role in virulence and frequent use in genome-based epidemiology, no reference antisera are currently being produced for *in vitro* applications. We used sera derived from CR_{KPK2}+p*espO* infection, together with a reference K2 antiserum as a control, to stain KP_{WT}. No detectable binding was seen in sera from CR_{WT}-infected mice. In contrast, binding was seen when KP_{WT} was stained with the CR_{KPK2}+p*espO* sera, which phenocopied the pattern of binding seen with the

228   reference K2 antiserum (Fig. 4D). Together with the ELISA results, this provides strong evidence for

229   specific heterologous α-KP K2 Abs induced by CR$_{KPK2}$+p*espO* infection.

KP serotyping using the citrOgen KP K2 sera

We next tested if the citrOgen KP K2 sera could be used for serotyping. To this end, we used a publicly

available collection of 100 KP isolates, representing 54 distinct CPS types determined by genomic

analysis[23]. We incubated the strains with citrOgen KP K2 sera and analysed binding by flow cytometry

(Fig 4E, Supplementary Fig. 5B). We included KP$_{WT}$ and KP$_{\Delta wcaJ}$ as controls and measured the median

fluorescence intensity (MFI) across the collection. The citrOgen KP K2 sera specifically bound KL2

isolates and KP$_{WT}$ with minimal cross reactivity and absent binding to KP$_{\Delta wcaJ}$. To confirm the KL

assignments, we selected 9 isolates for IF with the reference K2 antisera (Fig 4F). We observed a

pattern of binding consistent with capsule in the KL2 strain but not the surrounding non-KL2 strains.

We concluded that the citrOgen KP K2 sera can be used for serotyping and validated the collection's

KL2 assignments.

citroGen presentation of heterologous protein complex antigens

We next assessed the utility of citrOgen in generating Abs against heterologous proteins, using the

well-conserved KP T3F as a model protein complex[24]. T3F are a KP virulence factor that mediate binding

to abiotic surfaces (such as plastic and glass) and are involved in the first steps of biofilm formation.

The major T3F subunit is MrkA which together with MrkD (adhesin tip) and MrkF (unknown function)

are externalised via a β-barrel transmembrane usher protein, MrkC, and assemble into T3F on the KP

cell surface (reviewed in Murphy et al. 2012[25]). In the periplasm a chaperone, MrkB, binds MrkA to

prevent ectopic pilin polymerisation. Therefore, MrkABCDF, are all required for T3F biogenesis. T3F

expression is tightly regulated in KP by an adjacent operon (*mrkHIJ*) via the second messenger cyclic-

di GMP[26]. To coordinate expression of the T3F with CR attachment to IECs *in vivo,* we constructed a

citrOgen strain in which *mrkABCDF* (excluding *mrkHIJ*) was driven by the promoter of the T3SS effector

map (CR$_{T3F}$)[27]. The recombinant operon was inserted into the CR *glmS* site (Fig 5A). We perceived two

significant advantages of this strategy, firstly, maximising T3F expression during intimate attachment

to host cells and secondly to limit the metabolic burden of T3F expression in CR. To confirm accurate

gene regulation, we quantified *mrkA* transcript levels in T3SS-suppressing (LB media, static growth)

and T3SS-inducing (DMEM, static growth) conditions (Supplementary Fig 5A)[28]. qRT-PCR analysis

demonstrated low *mrkA* transcripts from CR_{T3F}+*pespO* grown in LB which increased significantly in

DMEM (Fig 5B), a change also observed for the endogenous *map* gene used as a positive control

(Supplementary Fig. 5A). No *mrkA* transcripts were detected in the CR_{WT} negative control. We then

confirmed that CR_{T3F} formed actin-rich pedestals *in vitro* and expressed KP T3F by staining with an α-

MrkA pAb (Supplementary Figure 5B). Finally, we tested if T3F conferred CR_{T3F} the ability to adhere to

glass under T3SS inducing conditions. This revealed a stark increase in the number of adhered CR_{T3F}

observed on glass coverslips compared to CR_{WT} when both were grown statically in DMEM, indicating

functional heterologous T3F expression (Fig 5C). Whilst this vector was designed to insert and provide

stringent control of T3F expression it can be readily be used for the expression of other operons

encoding mono/multimeric protein Ags to facilitate future iterations of this technology.

We next used oral gavage to infect CD-1 mice with CR_{T3F}+p*espO*, which colonised similarly to CR_{WT} at 8

dpi (Fig 5D). We used the sera collected from mice who had cleared the infection for ELISA, with plates

coated with heat killed KP_{WT} and $KP_{\Delta mrkA-F}$, a deletion mutant in the T3F structural genes. Sera collected

from the control CR_{WT}-infected mice were not reactive against KP_{WT} or the isogenic mutant (Fig 5E).

Conversely, sera from CR_{T3F}+*pespO* -infected mice bound $KP_{WT,}$ with absent binding to $KP_{\Delta mrkA-F.}$ This

data confirmed the α-KP IgG antibodies were directed towards T3F. Using pooled sera from the

CR_{T3F}+*pespO* infected mice to stain KP_{WT} revealed a binding pattern that matched that of the reference

α-MrkA pAb (Fig 5F). As T3F mediate attachment to abiotic surfaces and biofilm formation, we then

tested if the sera from could prevent KP_{WT} attachment. We first pre-adsorbed the sera from

CR_{T3F}+p*espO*- and CR_{WT}-infected mice against PFA-fixed CR_{WT} to remove α-CR Abs (10x 24h adsorption

cycles against fresh PFA-fixed CR_{WT}) (Supplementary Fig. 5C). This resulted in sera that negligibly bound

CR_{WT} (Supplementary Fig. 5D&E). We preincubated these two preadsorbed sera, which had equivalent

concentrations of total IgG (Supplementary Fig 5F), with KP_{WT} expressing sfGFP and measured

attachment after static growth at 37°C. Sera from CR_{T3F}+p*espO*-infected mice resulted in 3-fold

reduction in KP attachment compared to sera from CR_{WT}-infected mice (Fig. 5G &H). These data

demonstrate the generation of α-KP T3F Abs that functionally inhibit surface attachment, the first step

in KP biofilm formation.

Discussion

LPS, CPS and proteins are common bacterial targets where generating specific Abs is desirable. Whilst techniques already exist to generate such target-specific Abs, individually tailored and time-consuming approaches are required to prepare the Ags. Here, instead, we co-opt the mouse pathogen CR for this purpose and combine the steps of Ag preparation and immunisation into a single platform that both expresses heterologous Ags and presents them continuously, in their native conformation, to the host over the entire infection cycle (Figure 6). This was achieved by recombination-based genome editing in which whole operons were cloned from KP and inserted or substituted into the CR; thus, all genetically encoded components are inherently included, ensuring that Ag assembly takes place in a fashion mirroring the pathogen of choice. It should be noted that, whilst we achieve this for three diverse Ags, they all come from KP, another Gram-negative organism, which will have favoured our approach. Nevertheless, we were able to achieve specific and functional Ab responses for LPS, CPS and a large protein complex. If we consider the latest iteration of the WHO Bacterial Priority Pathogens List[12] then three out of four critical pathogens are Gram-negatives, indicating the clear and present danger posed by this group of bacterial threats. The data we present already supports using citrOgen for these organisms. We anticipate that further modifications will be required to synthesise and present Gram-positive bacterial Ags. As well as the ability to present complex polysaccharide structures and multicomponent protein Ags in their native location and configuration, one key advantage of citrOgen is time (Fig 6). In this work we have already validated substitution/insertion sites and generated the appropriate genome editing vectors. Changing the inserts encoded by these vectors (to alternative LPS, CPS and protein Ags) is routine and together with subsequent CR genome editing can be achieved over a period of days. Whilst we cloned sequences directly from KP DNA we propose that this technology could also be applied to generate diagnostic or therapeutic Abs in an outbreak scenario. The now routine use of *de novo* gene synthesis would enable us to engineer strains based on early sequencing data alone. This would allow us to use this platform to generate pAb sera in as little as 28 days- a complete CR infection cycle in mice. In more controlled settings we envisage generating

citrOgen strains encoding multiple Ags simultaneously. Ethically, this reduces the number of

experimental animals for use in research and development and would permit multiple target-specific

Ab responses to be obtained following a single infection.

We overcame two major obstacles to develop a viable platform for broader applicability. The first was

the potential fitness cost imposed by heterologous Ag expression. Indeed, we observed lower gut

colonisation in CR_{KPO1} than CR_{WT}, a defect that only becomes evident under the pressure of the host-

pathogen-microbiome environment. We overcame this obstacle by overexpressing the T3SS effector

EspO, which promotes IEC attachment and mucosal colonisation. The second was the metabolic cost

to CR of dysregulated heterologous protein expression which we overcame by placing expression

under the control of the endogenous CR promoter for the *map* gene. This restricts heterologous

protein expression to T3SS-inducing conditions and promotes presentation at host mucosal surfaces.

In the case of KP O1 Ag we combined citrOgen with subsequent KP lung challenge, which allowed us

to quantitively evaluate protection from organ failure conferred by the α-bAb response to this Ag (Fig

6). Given the current lack of correlates of protection for bacterial infections we propose that citrOgen

strains may represent an efficient tool to evaluate future proposed Ags. Strains engineered to express

heterologous Ags (singly or in combination) can be used to evaluate Ab responses following a CR

infection cycle. In the same cohort of mice, subsequent pathogenic challenge can then evaluate the

protection conferred by the response. Expanding the translational applicability of current priority

bacterial infection models would significantly enhance the data we could achieve in this approach.

CR infections are host restricted to mice, conferring platform safety and preventing infection risk to

humans. This is especially important as future citrOgen strains could include additional virulence

factors as Ags for Ab generation. We currently use the citrOgen platform in conventional mice

generating murine immunoglobulins. Advances in molecular genetics have enabled the generation of

'humanised' Abs initially raised in non-human species and could be applied to the current iteration of

the platform[29]. This improves IgG tolerability and durability when administered to human patients

largely by reducing the immunogenicity provoked by non-human Ig regions. Furthermore, extensive

work in the field of mouse genetics has enabled the generation of fully human Abs in transgenic

animals[30]. Combining these transgenic mice with citrOgen strain infection would result in human Ig

induction in one CR infection cycle (Fig 6). Existing downstream workflows, such as hybridoma

generation or IgG loci subcloning are already well defined and employed to produce mAbs from B-cells

derived from these mice. Whilst our antibiotic treatment options are rapidly diminishing the

application of this technology may alleviate future infectious diseases deaths by generating new Ab-

based diagnostic tools and therapeutics.

Acknowledgements

We would like to thank Professor Richard Strugnell and Dr Jonathan Wilksch at the University of Melbourne for providing the reference K2 antisera and C13 monoclonal antibody for use in this work. We would like to thank our colleagues Dr Abigail Clements, Dr Andrew Conway Morris, Professor Victor de Lorenzo, Professor Gordon Dougan and Professor Jose Penadés for their critical assessment and feedback in writing this manuscript. We would like to thank our colleagues Dr Mei Chong and Mr Edmond Yau in the Imperial College London Enterprise team for their support in the patent application covering the technology described in this manuscript.

Conflicts of interest

None declared.

Inclusion and diversity

We support inclusive, diverse, and equitable conduct of research.

Patent

The technology underlying citrOgen has a patent pending.

Funding

This project has been supported by grants from The Royal Society (IC160080) and Medical Research Council (MR/R02671).

Figure Legends

Figure 1. Heterologous KP O1 O-antigen expression causes an *in vivo* colonisation defect.

A. The *rfb locus* in CR_{WT} encodes a O152-like *E. coli* O-antigen locus located between *galF* and *gnd*. In

CR_{KPO1} the CR_{WT} *rfb* locus was substituted with the genes encoding KP O2a O-antigen and *wbbYZ*

(converted KP O2a to KP O1v1) inserted at the 3' *glmS* site.

B. CR_{KPO1} expresses a hybrid lipopolysaccharide molecule with native CR lipid A and core

lipooligosaccharide linked to KP O1 Ag. The western blot is representative of 2 biological repeats.

C. Western blot confirmation of heterologous KP O1 Ag in CR_{KPO1} performed on purified LPS

preparations. α-O152 Ab are immunoreactive with CR_{WT} but not KP_{WT} or CR_{KPO1} whereas α-KP O1 mAb

(C13) is immunoreactive with both KP_{WT} and CR_{KPO1}.

D. At 8 dpi following oral gavage, faecal CFUs are significantly lower in CR_{KPO1} compared with CR_{WT}.

n=10 mice per group, 2 biological repeats. Central tendency: median. Mann-Whitney test, ** exact

p=0.0095. GoF, gram of faeces.

E. Colonic histology was taken 8 dpi and stained for CR, KP O1 Ag; DAPI staining used as a structural

marker. CR_{WT} confluently adhering to the luminal surface of the colon with no detectable KP O1

staining. Sparse staining for CR_{KPO1} is observed with detectable KP O1 staining however, where present,

KP O1 Ag expression detectable. Scale bar = 200 μm. Images are representative of 3 biological repeats.

Figure 2. CR_{KPO1} overexpressing EspO maintains infectivity and induces specific anti-KP O1 Ag IgG responses.

A. At 8 dpi *espO* overexpression in CR_{KPO1}+p*espO* reverses the colonisation defect observed in CR_{KPO1} restoring colonisation to that of the parental CR_{WT} strain. n=15 mice per group, 3 biological repeats. Central tendency: median. Ordinary one-way ANOVA with Dunnett's multiple comparison. *adjusted p=0.0146 ns=non-significant. GoF, gram of faeces.

B. Colonic histology was taken 8 dpi and stained for CR and KP O1 Ag; DAPI staining used as a structural marker. CR_{KPO1}+p*espO* confluently adhering to the luminal surface with KP O1 Ag expression detectable and co-localising to CR staining. Sparse staining is observed for CR_{KPO1} with KP O1 Ag expression. Scale bar = 200 μm. Images are representative of 3 biological repeats.

C. At 30 dpi blood was taken from the tail vein of mice that cleared CR_{WT} and CR_{KPO1}+p*espO*. ELISA, using heat-killed KP_{WT} (expressing KP O1 Ag), revealed α-KP_{WT} IgG in mice following infection with CR_{KPO1} +p*espO* but not CR_{WT} controls. Central tendency: mean. Mann-Whitney test, **** exact p<0.0001.

D. Sera from mice was tested for binding to live KP_{WT} and $KP_{\Delta rfb}$. Binding was assessed by flow cytometry and expressed as the percentage of the population positive for mouse IgG (mIgG+ve). Sera from CR_{WT}-infected mice demonstrated absent binding to either KP_{WT} or $KP_{\Delta rfb}$, whereas sera from CR_{KPO1}+p*espO* -infected selectively bound KP_{WT} expressing KP O1 Ag. Central tendency: mean. Unpaired T-test. ns=non-significant. **** p<0.0001.

E. Sera from CR_{WT} and CR_{KPO1}+p*espO* -infected mice was used to stain KP_{WT} and image by immunofluorescence microscopy. An α-KP O1 Ag mAb (C13) was used as a positive control for KP O1 Ag staining. CR_{WT} derived sera results in no detectable staining of KP_{WT} whereas sera from CR_{KPO1} +p*espO* -infected mice and the control C13 mAb phenocopy each other with membrane-associated staining of KP_{WT}.

Figure 3. Antibodies to KP O1 O-antigen protect against KP pulmonary challenge.

Mice that had cleared pre-infection with CR$_{WT}$ and CR$_{KPO1}$+p$espO$ were challenged by intratracheal infection with 500 CFU of KP$_{WT}$. At 72 hpi we analysed KP replication (A&B) and host-derived severity indicators (C-O). n=15 mice per group (starting) over 3 biological repeats.

A&B. KP$_{WT}$ replication and bacteraemia are significantly reduced in the lungs (A) and blood (B) of CR$_{KPO1}$+p$espO$-preinfected mice compared to CR$_{WT}$ controls. . Central tendency: median. Mann-Whitney tests. ****exact p<0.0001 ***exact p=0.0002.

C. KP$_{WT}$ induces significant weight loss in CR$_{WT}$-preinfected mice which is reversed in CR$_{KPO1}$+p$espO$ preinfection. Central tendency: mean, error: standard deviation (SD). Mann-Whitney test. ****exact p<0.0001

D-F. Clinical haematological markers of disease severity are significantly improved in CR$_{KPO1}$+p$espO$ -preinfected mice compared to CR$_{WT}$ controls. Total white cell counts (D), lymphocyte counts (E) and platelets counts (F) are significantly lower in CR$_{WT}$-preinfected controls indicating increased diseases severity. The innate responders, neutrophils (G) are unchanged between the CR$_{WT}$ and CR$_{KPO1}$+p$espO$ preinfected groups. Central tendency: mean, error: SD. Unpaired T-tests **p<0.0065, n=non-significant.

H-J. The acute phase reactants (Serum amyloid A (H), C-reactive protein (I) and Albumin (J)) demonstrate a reduced inflammatory state in CR$_{KPO1}$+p$espO$ -preinfected mice compared to CR$_{WT}$ controls. Central tendency: mean, error: SD. H&J). Mann-Whitney test, I) Unpaired T-test ****p<0.0001, p**=0.0002.

K-M. Acute lung injury markers (lung homogenate IL1β (K), serum surfactant B (L) and lung homogenate myeloperoxidase (M)) are improved by pre-infection with CR$_{KPO1}$+p$espO$ compared to CR$_{WT}$ preinfection. Significant reductions in lung iIL1β, improvement in the alveolar capillary barrier (surfactant protein D) and neutrophilic alveolitis (myeloperoxidase) indicate α-KP O1 Abs protect against lung injury. Central tendency: mean, error: SD. K&M) Mann-Whitney test, L) Unpaired T-test ****p<0.0001, ***p=0.0001, *p=0.0109.

431 **N&O.** Acute kidney injury and excretory renal failure are reduced in mice preinfected with

432 CR_{KPO1}+p*espO* compared to CR_{WT}. Serum creatinine (N) and serum cystatin C (O) are reduced in KP_{WT}

433 infected mice who have cleared preinfection with CR_{KPO1}+p*espO* compared to CR_{WT}. Central tendency:

434 mean, error: SD. Unpaired T-tests ****$p<0.0001$, ***$p=0.0006$.

Figure 4. citrOgen derived anti-KP K2 antibodies can be used for serotyping.

A. The *cps* in KP_{WT} is c.22kb long with 15 open reading frames. This was inserted into CR_{WT} by substitution of the colanic acid biosynthesis locus generating the strain CR_{KPK2}.

B. At 8 dpi faecal CR_{KPK2}+p*espO* levels are similar to CR_{WT} . Central tendency: median. n=15 mice per group, 3 biological repeats. Unpaired T-test, ns=non-significant. GoF, gram of faeces.

C. At 30 dpi blood was taken from mice that cleared infection with CR_{WT} or CR_{KPK2}+p*espO*-. An ELISA was conducted against heat-killed KP_{WT} (expressing KP K2) and an isogenic acapsular mutant ($KP_{\Delta wcaJ}$). CR_{WT} infected mice had no detectable IgG responses to KP_{WT} or $KP_{\Delta wcaJ}$. CR_{KPK2} infected mice had IgG responses specifically against $KP_{WT.}$ n=15 mice per group, 3 biological repeats. Central tendency: mean. Mann Whitney test **** p<0.0001 ns=non-significant.

D. Sera from CR_{WT} and CR_{KPK2}+p*espO*-infected mice were used to stain KP_{WT} and image by immunofluorescence microscopy. An α-KP K2 pAb was used as a positive control. CR_{WT} derived sera results in no detectable staining of KP_{WT} whereas sera from CR_{KPK2}+p*espO*-infected mice and the α-KP K2 pAb phenocopy each other with punctate membrane-associated staining of KP_{WT}.

E. Sera from CR_{KPK2}+p*espO*-infected mice were used to serotype a collection of 100 diverse KP strains with serotypes assigned by genomic analysis. Each box represents one strain with the cps (KL) type described. The boxes are coloured according to the strength of binding by CR_{KPK2}+p*espO*-infected mice sera, which was assessed by flow cytometry. The median fluoresence intensity (MFI) key is shown in the bottom right together with staining of KP_{WT} and $KP_{\Delta wcaJ}$ controls. A box, centred on a KL2 isolate is shown.

F. The strains contained within the highlighted box in (E) were stained with the reference α-KP K2 pAb and imaged by immunofluorescence microscopy. The KL2 strain in the centre demonstrate punctate membrane staining in keeping with K2 capsule. The other strains (non-KL2) demonstrate no staining. Bacterial cell membranes were counterstained with FM-4-64 for imaging.

Figure 5. citrOgen derived anti-KP T3F antibodies block biofilm formation.

A. To generate a strain which expresses KP T3F the structural genes *mrkABCDF* were inserted at the 3'

glmS site in CR under the control of the T3SS effector protein Map promoter. This generated the strain

CR_{3F}.

B. *mrkA* transcripts were assessed by qRT-PCR from bacterial cultures of CR_{T3F} under T3SS suppressing

(LB) and inducing (DMEM) conditions. CR_{WT} was used as a negative control. T3SS inducing conditions

specifically induce *mrkA* transcription in CR_{T3F}. 4 biological replicates, central tendency mean, error

bars SEM. Paired T-test between LB and DMEM-grown CR_{T3F}; * $p<0.5$.

C. mCherry expressing CR_{T3F} and CR_{WT} were grown in T3SS inducing conditions on glass coverslips

bottomed slides. Following 6 h attachment to glass was assessed and the adhered bacteria were

stained for MrkA. A dense lawn of CR_{T3F} is observed binding to the glass slide with MrkA detected on

the cell surface indicating functional surface KP T3F expression. The parental CR_{WT} strain adheres to

glass slides only sparsely.

D. At 8 dpi faecal CR_{T3F}+p*espO* levels are similar to CR_{WT}. Central tendency: median. n=15 mice per

group, 3 biological repeats. Mann-Whitney test, ns=non-significant. GoF, gram of faeces.

E. At 30 dpi blood was taken from mice that cleared infection with CR_{WT} or CR_{T3F}+p*espO*. An ELISA was

conducted against heat-killed KP_{WT} (expressing KP T3F) and an isogenic mutant lacking T3F expression

($KP_{\Delta A-F}$). CR_{WT} infected mice had no detectable IgG responses to KP_{WT} or $KP_{\Delta mrkA-F}$. CR_{T3F}+p*espO* infected

mice had specific IgG responses against $KP_{WT.}$ n=15 mice per group, 3 biological repeats. Central

tendency: mean. Mann Whitney test, **** $p<0.0001$, ns=non-significant.

F. Sera from CR_{WT} and CR_{T3F}+p*espO*-infected mice were used to stain KP_{WT} and image by

immunofluorescence microscopy. An α-MrkA pAb was used as a positive control. CR_{WT} derived sera

results in no detectable staining of KP_{WT} whereas sera from CR_{T3F}+p*espO*-infected mice and the α-MrkA

pAb phenocopy each other with spiculated membrane-associated staining of KP_{WT}.

G. CR preadsorbed sera from CR_{WT} and CR_{T3F}+p*espO*-infected mice were preincubated with KP_{WT} or

$KP_{\Delta mrkA-F}$ expressing sfGFP. A no-antibody was used as a control. Following 6 h of growth in M9 minimal

media sfGFP signal was quantified on a plate reader and presented as the KP_{WT} binding normalised to

$KP_{\Delta mrkA-F}$ binding. The preadsorbed sera collected from CR_{T3F}+p$espO$-infected mice specifically inhibited

binding compared to $CR_{WT.}$ 4 biological replicates, central tendency: mean, error bars: SEM. Repeated-

measures One-way ANOVA corrected with Dunnett's multiple comparisons test, * $p<0.05$.

H. The biofilm formation from (G) was assessed by imaging using sfGFP fluorescence. A large reduction

in adhered KP_{WT} is observed when CR_{T3F} p$espO$-infected cross adsorbed sera was incubated with KP_{WT}

compared to CR_{WT} cross adsorbed sera.

Figure 6. The citrOgen platform as a versatile, quick and cost-efficient method of Ab production

A. citrOgen strain generation. Selected Ag sequences are incorporated into the appropriate gene

editing vectors designed for optimal chromosomal expression of heterologous O-Ag, capsule or

complex protein Ags; p*espO* is then transformed to finalise the new citrOgen strain.

B. In vivo Ab production. The CR infection cycle and immune responses are co-opted by the citrOgen

platform: (i) EspO-mediated boosting of CR attachment to intestinal epithelial cells, (ii) continuous

exposure of immune cells to the heterologous Ags, (iii) B cell affinity maturation results in Abs,

primarily IgG, which mediate CR clearance by 28 dpi and can be obtained as pAb sera from blood.

C. Downstream applications. Following citrOgen infection/immunisation, the functionality of Abs can

be assessed *in vitro* or by *in vivo* challenge studies, the pAb sera can be used for serotyping and the B

cells can be used for mAb production, either by using hybridoma technology to immortalise single-cell

cloned B-cells or by expressing heterologous Ag-specific Ab sequences in cell lines to obtain

recombinant mAb.

Ab: Antibodies; mAb: monoclonal antibodies; pAb: polyclonal antibodies; Ag: Antigen; CR: *Citrobacter*

rodentium.

Materials and Methods

Bacterial culture and antibiotics

Bacteria were routinely cultured in LB Broth (Miller) (84649.05000, VWR, UK). Saturated (overnight,

18 h) cultures took place by inoculating broth and incubation at 37°C and shaking at 200rpm.

Antibiotics were added to LB broth or LB broth solidified with agar (1.10283.0500, VWR, UK) at the

following concentrations:

Gentamicin 10mcg/ml (cloning) (G1272, Merck, UK)

Streptomycin 50mcg/ml (cloning) (S6501, Merck, UK)

Kanamycin 50mcg/ml (cloning) Merck, UK)

Nalidixic acid 50mcg/ml (selecting CR and CFU enumeration from *in vivo* sample) (N4382, Merck, UK)

Rifampicin 50mcg/ml (selecting KP and CFU enumeration from *in vivo* sample) (R3501, Merck, UK)

Cloning

High-fidelity polymerase chain reaction was conducted using Phanta Flash (2x Phanta Flash Master Mix

(Dye Plus), Vazyme, China). Screening and colony polymerase chain reaction was conducted using

Rapid Taq (2x Rapid Taq Master Mix, Vazyme, China).

PCR purification (Monarch PCR & DNA Cleanup Kit (5ug), NEB) and plasmid extraction were conducted

with kit-based methods.

All mutagenesis vectors were generated by PCR products ligation via Gibson Assembly (NEB, UK)

according to the manufacturer's guidance.

Genome editing

The whole genome references sequences used for CR strain ICC168 (Nalidixic acid-sensitive version of

ICC169) is FN543502. The whole genome reference sequence used for ICC8001 is the closely related

ATCC43186-deriviitve KPPR1 is CP009208.1. A list of strains generated in this work can be found in

Supplementary Table 1.

Insertions at the CR 3' *glmS* site were conducted using Tn7 mutagenesis with methods published in

Choi et al. Nat Methods 2005 Jun;2(6):443-8[17]. Seamless, markerless genome editing at other sites was

conducted by homologous recombination based on the pSEVA612S system (https://seva-
plasmids.com/) in both CR and KP as previously published in detail[9,31]. To improve the efficiency of
pSEVA612S for Gibson Assembly we further modified this plasmid by reversing the sequence of the 3'
ISceI endonuclease digestion site generating pSEVA612S(R). This modification improves efficiency by
reducing homology-based empty vector recombination.

All mutagenesis vectors are summarised in Supplementary Table 2. These will be submitted to a
publicly accessible NCBI depository upon publication. In the case where more than one vector was
used these are listed against the respective strain. To generate CR_{KPK2} we initially cloned a construct
(pSEVA612S(R)_CRColanic::K2cpsΔwbbJ_ORF) without the genes wbbJ and an annotated open reading
frame of unknown function according to Feldman et al Proc Natl Acad Sci U S A 2019 Sep
10;116(37):18655-18663[32]. We subsequently inserted these genes into our final CR_{KPK2} strain
(pSEVA612S(R)_ΔwbbJ::Ins_wbbJ_ORF) as new data become available according to Walker et al 2020
mBio 2020 Sep 22;11(5):e01750-20[33] and included the native KP promoter
(pSEVA612S(R)_K2:col_Pvariant_Porf2).

Transformation of pACYC184_espO was conducted using room temperature competent cells according
to our previously published protocol[31].

Primers sequences available from corresponding authors upon request.

Sequencing

Sanger sequencing of PCR products and plasmids was conducted by Eurofins Genomics.

In silico serotyping

In silico serotyping of CR was conducted using SerotypeFinder 2.0
(https://cge.food.dtu.dk/services/SerotypeFinder/) on genome FN543502.

LPS extraction and Western Blotting

LPS was extracted from 3OD units of saturated overnight cultures using a modified phenol method kit
(iNtRON Biotechnology, Korea) according to the manufacturer's guidance with no modification.

Samples were mixed with 5x Laemmli buffer and 5% 2-mercaptoethanol and separated by SDS–

polyacrylamide gel electrophoresis (PAGE) using a 12% acrylamide gel and tris-glycine buffer system

and transferred to polyvinylidene difluoride membranes (#1620177, Bio-Rad Laboratories) using a

TransBlot semi-dry electrophoretic transfer machine (Bio-Rad). Membranes were blocked for 2 h at

room temperature (RT) in 10% fat-free milk PBS–0.1% Tween 20 (PBS-T) and incubated overnight at

4°C with 1:500 anti-O152 or 1:2000 anti-KP O1 (Supplementary Table 3) in PBS-T with 5% BSA (A3059-

50G, Sigma). Secondary antibodies (Supplementary Table 3) were added diluted 1:10000 in 5% fat-free

milk in PBST and incubated for 1 h at RT. Immunoblots were developed with Clarity Western enhanced

chemiluminescence (ECL, Bio-Rad Laboratories) and images were acquired on a ChemiDoc XRS+ and

visualised using Image Lab software (Bio-Rad Laboratories).

In vitro CR_{KPO1}, $CR_{KP\ K2}$ and CR_{T3F} infections of fibroblasts

Murine Swiss 3T3 fibroblasts (ATCC® CCL-92™) were maintained in DMEM containing 4500   mg l^{-1}

glucose (D6546, Sigma) and supplemented with 10% fetal calf serum (F9665, Sigma) and 2 mM

glutaMAX (#35050061, ThermoFisher Scientific) at 37°C in 5% (v/v) CO_2 and seeded on glass coverslips

in 24-well plates (density $2x10^5$cells/well) for immunofluorescence. Indicated strains of CR expressing

mCherry from a pUltra plasmid[34] were grown to saturation in LB supplemented with gentamicin at

37°C with 200 rpm agitation for 8 h before subculturing 1:500 for 16 h in LB or DMEM with 1000 mg

l^{-1} glucose (# D5546, Sigma) at 37°C, 5% (v/v) CO_2, static. Cells were infected with the static culture at

a multiplicity of infection (MOI) of 100, confirmed by retrospective plating. The plate was centrifuged

at 1000 rpm for 5 min at RT to synchronize infection and incubated for 4 h at 37°C in a 5% CO_2

atmosphere. The infection was stopped with three washes with sterile phosphate-buffered saline (PBS)

and cells were fixed in PBS with 4% (w/v) paraformaldehyde for 15 mins at RT and washed three times

in PBS. Cells were then permeabilized in 0.1% (v/v) of Triton X100 in PBS for 10 min, incubated in 50

mM NH_4Cl in PBS as a quenching agent for 10 min, washed in PBS, and blocked in 3% BSA in PBS for 1

h at RT before staining with one of the following primary antibodies in 1% BSA-PBS for 1 h at RT: anti-

KPO1 C13 (1:200), anti-mrkA (1:100) or anti-KP K2 (1:100) (Supplementary Table 3). Coverslips were

washed in PBS and incubated with Alexafluor488-conjugated secondary antibodies (1:200),

Tetramethylrhodamine (TRITC)–conjugated phalloidin (1:500) and 4',6-diamidino-2-phenylindole

(DAPI) (1:1000) to label the heterologously-expressed antigen, F-actin and DNA, respectively

(Supplementary Table 3) in 1% BSA-PBS for 1 h. Coverslips were washed in PBS and once in water

before being mounted using Gold Pro-Long Anti-fade medium (#P36930, ThermoFisher Scientific) and

left to cure overnight at RT in the dark. Immunofluorescence images were acquired using a Zeiss

AxioImager Z1 microscope and processed using ZEN3.1 software (ZEN lite, Zeiss).

Reverse transcription quantitative PCR (RT-qPCR)

The indicated strains of CR were grown to saturation overnight in LB supplemented with the

corresponding antibiotics. 0.5 ml of the overnight culture were washed once in PBS, incubated in

RNAprotect Bacteria reagent (#76506, Qiagen) for 10 mins at RT and centrifuged at 3000 xg for 5 min.

The bacterial pellet was digested in 100 µl TE buffer ((30 mM Tris·Cl, 1 mM EDTA, pH 8.0) with 15 mg/ml

Lysozyme (L6876, Sigma-Aldrich) and 20 µl Proteinase K (#19131, Qiagen) for 10 min according to

manufacturer's guidelines. RNA was isolated using the RNeasy minikit (Qiagen) following the

manufacturer's instructions, and RNA concentration and purity was determined using a Nanodrop

2000 spectrophotometer (ThermoFisher Scientific). 1 µg of RNA was treated with RNase-free DNase

(#M6101) for 1 h at 37 °C and cDNA was then synthesized using a Moloney murine leukemia virus (M-

reverse transcriptase (M-MLV, #M1705) with random primers (#C1181) following the manufacturer´s

instructions (all from Promega). To confirm the absence of undigested DNA, a reaction mixture without

the M-MLV reverse transcriptase was also included (NRT). qPCR was performed using the Power Up

SYBR Green master mix (#A25778, ThermoFisher Scientific) and the relevant primers, whose primer

efficiency had been evaluated previously and determined to be 95-105% (Supplementary table 4). The

assay was run on a Quantstudio 1 System (Applied Biosystems, ThermoFisher Scientific) and results

were analysed using the QuantStudio Design & Analysis Software v1.5.2 (Applied Biosystems,

ThermoFisher Scientific). The relative expression level of each mRNA was evaluated using the 2-

ΔCtmethod, where Ct is the threshold cycle and 16S rRNA is used as the control.

Whole bacterial cell ELISA

To prepare heat-killed KP the indicated strains were grown overnight in LB at 37°C at 200 rpm. The culture was then normalised to an optical density of 1 at 600 nm in sterile PBS, heat inactivated in the presence of EDTA-free protease inhibitor cocktail (#4693159001, Sigma) at 60°C for 1 h and stored at 4°C until use (short-term). ELISA Flat-Bottom Maxisorp Immuno 96- Well Plates (#10547781, Fisher Scientific) were incubated with 100 µL/well heat-killed CR for our serum samples or with 100 µL/well goat anti-mouse IgG antibody for the standards (2 µg/µL) (Supplementary Table 3) overnight at 4°C. Plates were washed three times with PBS-T, blocked for 1 h at RT with 5% BSA-PBS-T, and incubated overnight at 4°C with serum samples or serially diluted mouse IgG (Supplementary Table 3) as standard. Plates were washed as described above and incubated with horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG antibody (Supplementary Table 3) for 1 h at room temperature. Plates were washed and developed by the addition of 100 µL 3,3',5,5'-tetramethylbenzidine (TMB; Thermo Fisher Scientific) for 5-15 mins before stopping the reaction with 2 N H_3PO_4. The absorbances at 450 and 540 nm were measured in a plate reader (FLUOstar Omega; BMG Labtech), and anti-KP IgG levels were quantified in samples by comparison to a mouse IgG standard curve.

Determination of O-antigen-binding specificity: KP_{WT} expressing mCherry and $KP_{\Delta rfb}$ expressing sfGFP were grown as saturated overnight cultures in LB supplemented with gentamicin. On the following day, both strains were mixed 1:1 (v/v) after confirming the had grown to the same optical density (600nm) and transferred to a V-bottom 96-well plate (Greiner Bio-one) for staining. Single strain controls were also included to ensure correct gating during flow cytometry analysis. Samples were first blocked in 5% BSA/PBS for 1 h at RT shaking, centrifuged at 4000 rpm, 4 °C, 10 mins and then incubated overnight with sera from CR_{WT} or CR_{KPO1}+pespO-infected mice diluted 1in100 (v/v) in 1% BSA/PBS. Bacterial samples were centrifuged, washed twice in cold PBS and resuspended in 1% BSA/PBS containing AlexaFluor647-conjugated anti-mouse IgG (1:200, Supplementary Table 3) for 1 h at RT shaking. Cells were washed twice in cold PBS, fixed in 1% PFA/PBS and kept in the dark until flow cytometry analysis.

Samples were acquired (minimum of 100000 bacterial cells per sample) on a Cytek Amnis CellStream

and data was analysed using FlowJo v10.10.0 (gating strategy shown in Supplementary Fig 3A).

K2 serotyping: The KP MRSN Diversity Panel (NR-55604) was obtained through BEI Resources, NIAID,

NIH[23]. The 100 strains from the MRSN collection, KP_{WT} and $KP_{\Delta wcaJ}$ (positive and negative controls

respectively) were grown overnight in 1mL of LB in deep well 96-well plates (VWR). 100ul of each

overnight was transferred to a V-bottom plate and used for staining. Plates were centrifuged at 4000

rpm for 10 mins at 4°C, resuspended in 5% BSA/PBS and blocked for 1 h at RT shaking using a

microplate mixer. Bacterial samples were then centrifuged and stained overnight at 4°C shaking using

sera from CR_{KPK2}+pespO-infected mice diluted 1in100 (v/v) in 1% BSA/PBS. On the following day,

bacterial samples were washed two times in cold PBS and stained for 1 h at RT shaking with AlexaFluor-

488-conjugated anti-mouse IgG (1:300; Supplementary Table 3). Cells were washed twice in cold PBS,

fixed in 1% PFA/PBS and kept in the dark until flow cytometry analysis. Samples were acquired on a

Cytek Amnis CellStream (minimum of 50000 bacterial cells per sample) and data was analysed using

FlowJo v10.10.0.

Bacterial cell immunofluorescence microscopy (adherent protocol and suspension protocol)

Adherent protocol (all ICC8001 staining)

mCherry expressing ICC8001 was grown as a saturated overnight culture which was sub-cultured

(1;1000 v/v) into M9 media (M9 salts M6030, Sigma, UK) supplemented with 0.4% glucose w/v (G8270,

Sigma, UK). 400ul was added into each well for imaging in a coverslip slide (μ-Slide 8 Well high Glass

Bottom, ibidi). Slides were grown statically at 37°C for 4 h and cells were washed in phosphate-buffered

saline before staining. Primary staining (sera, antibodies) took place overnight at 4°C in 0.22um filtered

3% bovine serum albumin (A3059, Sigma, UK)/ PBS on a rocking platform. Primary antibody was

removed and slides were washed 3x in PBS followed by incubation with fluorescently labelled

secondary antibodies (Supplementary Table 3) in 3%BSA/PBS at room temperature for 1 h on a rocking

platform. Slides were washed 3 times before imaging.

Suspension protocol (MRSN collection, Figure 4D)

MRSN collection strains were grown as saturated overnight cultures. 100ul of overnight was used for

staining. Following centrifugation (5000g, 5 minutes) cells were washed 3 times in PBS. Primary

staining (K2 antisera) took place overnight at 4°C in 3%BSA/PBS in 400ul volume with agitation. Cells

were washed by sequential spins (5000g, 5 minutes) and resuspension in PBS (3 washes). Secondary

antibody (Supplementary Table 3) was added for 45mins at room temperature in a 100ul volume with

agitation. Cells were washed by sequential spins (5000g, 5 minutes) and resuspension in PBS (3

washes). Cell membranes were then labelled with FM® 4-64 Dye (T13320, ThermoFisher Scinentific,

UK) at a final concentration of 5ug/ml in 100ul of PBS for 10 mins at room temperature in the dark.

Cells were washed once and resuspended in 100ul of fresh PBS before imaging. Cell suspension was

imaged on an 1% agarose (in dH$_2$0) pad according to the following protocol (https://biotium.com/tech-

tips/tech-tip-imaging-bacteria-using-agarose-pads/).

CR$_{T3F}$+p*espO* glass binding assay

mCherry expressing CR$_{T3F}$+p*espO* and CR$_{WT}$ were grown as saturated overnight cultures and sub-

cultured (1:100) into DMEM low glucose (D5546, Merck, UK) into glass coverslip slides (μ-Slide 8 Well

high Glass Bottom, ibidi). Slides were grown statically at 37°C for 6 h and cells were washed in

phosphate-buffered saline before fixing in 4% paraformaldehyde/PBS (R1026, Agar Scientific, UK) for

20 mins at room temperature. Cells were stained for MrkA according to the adherent protocol outlined

above.

Pre-adsorption of CR sera

Serum samples from mice infected with CR$_{T3F}$+pespO with anti-KP antibodies above 200ng/mL were

pooled; sera from CR$_{WT}$-infected mice were also pooled separately as a control. To inactivate

complement, pooled serum samples were heated at 55°C for 1 h and then were pre-adsorbed onto

CR$_{WT}$ expressing sfGFP to remove CR-specific antibodies, thus enriching for anti-KP antibodies. This was

done by serially incubating the pooled sera samples, both from CR$_{WT}$- and CR$_{T3F}$+pespO-infected mice,

with ~5x10^8 CR$_{WT}$ for 24 h at 4°C on a rotating wheel for 10 rounds of pre-adsorption. After each 24h-

incubation, the serum samples with CRWT were centrifuged at 3000 xg for 5 mins to pellet the

bacteria, which were kept at 4°C until flow cytometry analysis; the supernatant containing the

antibodies was subsequently incubated with a fresh batch of ~5x10^8 CR$_{WT}$ expressing sfGFP. At the end

of the 8 rounds, sfGFP-expressing bacterial samples from each round of pre-adsorption with sera from

CR$_{WT}$- and CR$_{T3F}$+*pespO*-infected mice were stained with AF555-conjugated anti-mouse IgG (1:400;

Supplementary Table 3) in 2% BSA-PBS for 30 mins and analysed using a Cytek Amnis CellStream.

Obtained data was subsequently analysed using FlowJo v10.10.0.

Dot blot

Bacterial cultures were grown overnight in LB (CR) or M9 supplemented with glucose (KP), normalised

to an OD600 of 1 and centrifuged to pellet the cells. The pellet was resuspended in 1 ml PBS and equal

volumes of bacteria were then spotted onto a nitrocellulose membrane (#88018, ThermoFisher

Scientific) and allowed to air dry. The membrane was blocked at RT for 2 h in 5% milk-PBS-T and

incubated overnight at 4°C with the indicated sera diluted in 5% BSA-PBS-T to a concentration of

2µg/ml of total IgG. The primary was washed three times in PBS-T and a secondary HRP-conjugated

anti-mouse antibody (Supplementary Table 3) was added in 5% milk-PBS-T, incubated for one hour at

RT, and washed three times. Detection was carried out using ECL substrate (Bio-Rad) and images were

acquired on a ChemiDoc XRS+ and visualised using Image Lab software (Bio-Rad Laboratories).

Fluorescent biofilm formation assay

KP$_{WT}$ and KP$_{\Delta mrkA-F}$ expressing sfGFP were grown overnight in M9 media and diluted into fresh M9 1:500

(v/v) and preincubated for 30 mins with no sera or with ~50 µg/mL IgG (1:100 dilution) of the pre-

adsorbed sera from either CR$_{WT}$- and CR$_{T3F}$+p*espO*-preinfected mice at RT on a rotating wheel. 100 µl

of each sample in triplicate were transferred into wells in a black/clear-bottom 96-well tissue-culture

treated plate (#353219, SLS) and incubated statically at 37°C for 6 h. Samples were then washed 5

times with sterile PBS and bottom optic fluorescence readings were taken at 485/520 (Ex/Em) using a

FLUOstar Omega plate reader. Fluorescence values were normalised to those obtained with the

KP$_{\Delta mrkA-F}$ strain, which does not attach, to remove any non-specific fluorescence and reduce batch-to-

batch variation.

Animal work

All animal work was conducted at Imperial College London (Association for Assessment and

Accreditation of Laboratory Animal Care accredited unit) under the auspices of the Animals (Scientific

Procedures) Act (UK) 1986 (PP7392693). The animal work was approved locally by the institutional

ethics committee.

Female CD-1 and C57BL/6 mice were purchased from Charles River, UK. Mice were purchased at 29-

31 g weight (5-7 weeks) (CD-1 mice) and 18-20 g weight (6-8 weeks) (C57BL/6 mice).

Husbandry

Animal technicians within the facility received mice and blindly separated them into groups of 5 into

high-efficiency particulate air (HEPA)–filtered cages with sterile bedding. Bedding was changed weekly,

and enrichment supplied. Mice were housed under a 12/12-h light/dark cycle with access to food and

water *ad libitum*. Researchers (JSG and JLCW) conducting the work in this study were not involved in

selection of animals into groups as this was undertaken by technical staff prior to starting experiments.

Animals were identified by ear notching (conducted under restraint with no anaesthesia).

CR infection by oral gavage

For infections, the indicated CR strains were grown overnight in 15 ml of LB (50 ml capped Falcon

tubes) at 37°C and 200 rpm. The next day, the tubes were centrifuged, and the pellet was resuspended

in 1.5 ml of sterile PBS. Mice were infected by oral gavage with 200 μl of this solution (1x10^9 CFU

confirmed by viable counts as previously described[10]). CR faecal shedding was determined by collecting

stool samples at the specified days post-infection (DPI), homogenising the samples in PBS and plating

serial dilutions onto LB agar containing nalidixic acid. Plates were incubated at 37°C overnight and

colonies were counted on the following day.

Mice were culled at 6 (C57BL/6) or 8 dpi (CD-1) and colons were extracted for downstream processing.

When characterising antibody responses, mice were followed until clearance of CR infection and then

culled to obtain sera or challenged with KP infection.

Colonic histology processing and immunostaining

At 8 dpi, the colon was extracted, and the distal 0.5 cm was fixed in 4% PFA for 2 hours and then kept

in 70% ethanol until processing. Fixed tissues were processed, paraffin embedded, and sectioned at 5

µm. For immunofluorescence, sections were dewaxed by submersion in HistoClear II solution

(#101412-884, VWR) twice for 10 min, 100% ethanol twice for 10 min, 95% ethanol twice for 10 min,

80% ethanol once for 3 min, and PBS-0.1% Tween 20-0.1% saponin (PBS-TS) twice for 3 min.

Subsequently, sections were heated for 30 min in demasking solution (0.3% trisodium citrate-0.05%

Tween 20 in distilled H2O). Once cooled, slides were first blocked in PBS-TS supplemented with 10%

normal donkey serum (NDS, D9663, Sigma) for a minimum of 1 h in a humid chamber, before being

incubated with anti-CR antibody (1:50) and anti-KP O1 (1:100) diluted in PBS-TS with 10% NDS

overnight at 4°C. Slides were rinsed twice for 10 min each time in PBS-TS, followed by incubation with

Alexafluor488-conjugated secondary antibodies (1:100) and DAPI (1:1000) (Supplementary Table 3).

Washing steps were repeated before slides were mounted with ProLong Gold antifade mountant.

Images were acquired using a Zeiss AxioVision Z1 microscope using a Hamamatsu microscope camera

and processed using ZEN3.1 software (ZEN lite, Carl Zeiss MicroImaging GmbH, Germany).

cIEC isolation and flow cytometry analysis

Colonic intestinal epithelial cells (cIECs) from mock-infected and CR-infected C57BL/6 mice were

isolated from 2-cm distal colonic tissue samples as previously described[35]. Briefly, the 2-cm colonic

tissue sample was opened longitudinally and washed in 1× Hanks' balanced salt solution (HBSS)

without Mg and Ca (# 14185045, ThermoFisher Scientific) and incubated at 37°C, 200 rpm for 45 min

in IEC dissociation buffer (1× HBSS, 10 mM HEPES, 1 mM EDTA, and 5 µL/mL 2-β-mercaptoethanol).

The remaining tissue was removed, and detached cells were collected by centrifugation (3000 rpm for

10 min), followed by two DPBS (#D8537, Sigma) washes at 4°C. Cells were then incubated 5 mins at RT

with 50 µg/mL DNase I (#11284932001, Sigma) to digest extracellular DNA and passed through a 70-

µm cell strainer on ice to ensure a single-cell suspension, pooling samples from the same group and

rinsing with 20 mL of cold DPBS. All steps from here onwards were performed on ice/at 4°C to maintain

cell viability. cIECs were counted using trypan blue to exclude dead cells and ~1x10^7 live cells were

centrifuged, resuspended in fluorescence-activated cell sorter (FACS) buffer (5% FBS, 2mM EDTA in

DPBS) supplemented with Fc block (#130-092-575, Miltenyi Biotec) and incubated for 10 min. Samples

were then incubated with FACS buffer containing anti-CR (1:1000) for 20 mins, washed twice in and

then incubated with EpCAM-APC (1:100) and phycoerythrin (PE) anti-rabbit IgG (Supplementary Table

3) for 40 min, washed, and fixed in 1% paraformaldehyde in PBS prior to analysis on a BD LSRFortessa

cell analyser (BD Biosciences). Data were analysed using FlowJo v10.10.0 and the gating strategy is

shown in Supplementary Fig. 2A.

Tail vein bleeds and serum separation

Small volume bleeds (<100ul) were conducted on conscious, restrained mice using a Tail Vein Injection

platform (Hugeiron, MouseSerum, China). Blood was taken from the lateral tail vein by puncture with

a 25G needle and collected via pipette into a paediatric serum collection tube (Microvette® 100 Serum

Cat, Starstedt). Following 1 hour of clotting at RT, sera was separated by centrifugation at 20 000g for

3 minute. Separated sera was stored at -80°C until use.

KP pulmonary infections

Anaesthesia (ketamine 80mg/kg and medetomidine 0.8mg/kg) was delivered by intraperitoneal

injection. Body temperature was maintained with a heat mat throughout until recovery or during the

intubation procedure itself.

Intubation was achieved with a 20G cannula (BD Insyte, BD) using the commercially supplied

intubation kit (Endotracheal Mouse Intubation Kit, ETI-MSE, Kent Scientific). 50ul of inoculum (diluted

in phosphate buffered saline) was delivered into the hub of the 20G cannula and drawn into the lungs

by spontaneous ventilation.

Recovery took place at 32°C in a forced air heating cabinet. At 15 minutes post anaesthesia induction

medetomidine was reversed by the administration of 0.8mg/kg atipamezole, delivered by

subcutaneous injection into the neck scruff.

The total volume of administered drugs was designed to deliver 20mg/kg of total fluids to animals, a

human paediatric resuscitation dose.

Mice were returned to their home individual ventilated cages after they were spontaneously moving

and checked at regular intervals post infection.

Cardiac puncture

Mice were weighed at end-point (after clearance for CR infected animals and 72 hpi for KP-challenged

animals).

Anaesthesia (ketamine 100mg/kg and medetomidine 1mg/kg) was delivered by intraperitoneal

injection. Mice were checked and following the absence of pedal reflexes placed in dorsal recumbency.

Paws were taped to a board and scissors used to remove the skin over the xiphisternum. Blood was

collected from the right ventricle (closed approach) with a 25G needle inserted at 30° to the skin

surface until blood was freely aspirated into a 1ml syringe.

When blood was obtained for determination of IgG titres (i.e mice infected with indicated CR strains

expressing heterologous antigens), the 25G needle was removed (to prevent haemolysis) and the

blood was transferred to a serum collection tube containing a clot activator (BD Microtainer SST with

separating gel, BD). Blood in the serum tube was allowed to clot for 1 hour at RT before centrifugation

(20 000 xg) for 3 minutes. The sera was divided into aliquots and stored at -80°C until use.

When blood was obtained from KP challenged mice, 20ul of whole blood was taken for CFU

enumeration by plating 10-fold serial dilutions (in PBS supplemented with 1mmol EDTA to prevent

clotting) on LB Agar plates containing rifampicin 50ug/ml. The 25G needle was again removed and

approximately 50-100 μl placed immediately into a tube coated with EDTA for anticoagulation

(Microvette® 100 EDTAK3E, 100ul, Sarstedt) for haematological analysis. The remaining blood

(approximately 800 μl) was allowed to clot for 1 hour at room temperature in a serum collection tube

as described above, before centrifugation (20 000g) for 3 minutes. The sera was divided into aliquots

and stored at -80°C until use.

Lung homogenate processing

Post-mortem lungs were removed and placed in 3ml of phosphate buffered saline in a gentleMACS C-

tube (Miltenyi Biotec). These were homogenised on a gentleMACS Octo Dissociator (Miltenyi Biotec)

using the program m_lung_2. The program was run twice in succession to achieve uniform

homogenisation.

30ul of lung homogenate was taken for CFU enumeration by plating 10-fold serial dilutions on LB Agar

plates containing rifampicin 50ug/ml.

1ml of lung homogenate was transferred into a 1.5ml microcentrifuge tube and spun at 10 000 xg for

5 minute at 4°C to pellet out particulate matter. The supernatant was stored -80°C for future use.

HM5 analysis of EDTA anticoagulated whole blood

Anticoagulated blood was run on the HM5 Vetscan analyser according to the manufacturer's

instruction. Outcome variables of infection were assessed using the kits summarised in Supp Table 5

according to the manufacturer's guidelines with the following specifications.

Capture ELISAs: ELISAs were performed using ELISA Clear Flat-Bottom Maxisorp Immuno 96- Well

Plates (#10547781, Fisher Scientific). Samples were added to coated and blocked plates and incubated

overnight at 4 degrees for all ELISAs. The range of sample dilutions used for each ELISA is indicated in

Supplementary Table 5; serial dilutions in the corresponding assay diluent were performed when

samples needed to be diluted further than 1in50. After addition of the TMB substrate solution (#00-

4201-56, ThermoFisher Scientific) to the wells at the end of the assay, these were incubated at RT until

a clear gradation in colour could be seen from the highest to the lowest standard, at which point the

reaction was stopped by adding 1M phosphoric acid (Sigma). All absorbance readings were obtained

using a using a FLUOstar Omega plate reader and readings at 540 nm were subtracted from absorbance

readings at 450nm to correct for optical imperfections of the plate. All ELISAs included blanks and

standard curves with at least 7 points within range. Sample values were obtained by interpolation into

the standard curve; any samples with values out of the range of the standard were assayed again at a

dilution(s) that ensured the value was within range.

Albumin: All reactions were done in a total volume of 200 µl in an untreated 96 well plate (Greiner

Bio-one) to allow vigorous shaking after reagent addition. Samples and standard were diluted 1:5 in

PBS (0.01 M, pH 7.4) and 5 µl were added per well; 5 µl of double-distilled water were added to the

blank. Subsequently 195 µl of stock working solution (prepared according to the manufacturer's

instructions) were added into each well, mixed thoroughly using a plate mixer and then left to stand

at RT for 10 mins before measuring the absorbance of each well at 628 nm using a FLUOstar Omega

plate reader.

Creatinine: Serum samples did not require deproteinization. The assay was performed using the

colorimetric reaction mix, and readings were acquired by measuring absorbance at 570 nm using a

FLUOstar Omega plate reader. High levels of haemolysis lead to high background due to the colour;

background control samples (ie sample with no creatininase) were essential in those cases. 40 µl of

serum sample and 10 µl of assay buffer were used per sample well; standards were prepared according

to the manufacturer's instructions. After interpolating into the standard curve and correcting for

sample dilution, the values are given as nmol/ 50 µl sample; these were transformed to mg/dL, where

the molecular weight of creatinine is 113.12 mg/mmol.

Data analysis and visualisation

Graphs and statistical analysis was carried out in GraphPad Prism version 10.4.1 for Windows and

version 10.4.0 for Mac (GraphPad Software, Boston, Massachusetts USA, www.graphpad.com). Data

were analysed for normal distribution based on D'Agostino-Pearson or Shapiro-Wilk normality tests

before performing any statistical analyses. Normally distributed data was analysed with the parametric

tests detailed in the figure legends; indicated non-parametric tests were applied to data that failed to

meet the normality criteria. When more than three comparisons were made, p values were adjusted

for multiple comparisons as indicated in the figure legends (Dunnett's post hoc test). For ELISAs, two

or three technical replicates were used to estimate experimental mean. Statistical significance was

only determined in datasets where each group had three or more biologically independent repeats. A

p value <0.05 is considered statistically significant.

Schematics in Supplementary Fig. 5C and Fig. 6 were made using some images from BioRender

(created with BioRender.com). Images were edited and compiled into figures in Affinity Designer V2.

(Serif Europe Ltd).

References

1.   El-Atawi, K. *et al.* Efficacy and Safety of Palivizumab as a Prophylaxis for Respiratory Syncytial Virus (RSV) Disease: An Updated Systemic Review and Meta-Analysis. *Cureus* **15**, e51375 (2023).

2.   Thandavaram, A. *et al.* The Efficacy of Bezlotoxumab in the Prevention of Recurrent Clostridium difficile: A Systematic Review. *Cureus* **14**, e27979 (2022).

3.   Self, W. H. *et al.* Neutralizing COVID-19 Convalescent Plasma in Adults Hospitalized With COVID-19: A Blinded, Randomized, Placebo-Controlled Trial. *Chest* **162**, 982–994 (2022).

4.   Focosi, D., Franchini, M., Casadevall, A. & Maggi, F. An update on the anti-spike monoclonal antibody pipeline for SARS-CoV-2. *Clin Microbiol Infect* **30**, 999–1006 (2024).

5.   Tkaczyk, C. *et al.* Identification of anti-alpha toxin monoclonal antibodies that reduce the severity of Staphylococcus aureus dermonecrosis and exhibit a correlation between affinity and potency. *Clin Vaccine Immunol* **19**, 377–85 (2012).

6.   Zhu, H., Rollier, C. S. & Pollard, A. J. Recent advances in lipopolysaccharide-based glycoconjugate vaccines. *Expert Rev Vaccines* **20**, 1515–1538 (2021).

7.   Mundy, R., MacDonald, T. T., Dougan, G., Frankel, G. & Wiles, S. Citrobacter rodentium of mice and man. *Cell Microbiol* **7**, 1697–706 (2005).

8.   Sanchez-Garrido, J., Ruano-Gallego, D., Choudhary, J. S. & Frankel, G. The type III secretion system effector network hypothesis. *Trends Microbiol* **30**, 524–533 (2022).

9.   Ruano-Gallego, D. *et al.* Type III secretion system effectors form robust and flexible intracellular virulence networks. *Science* **371**, (2021).

10.   Crepin, V. F., Collins, J. W., Habibzay, M. & Frankel, G. Citrobacter rodentium mouse model of bacterial infection. *Nat Protoc* **11**, 1851–76 (2016).

11.   Dennis, A. *et al.* The p50 subunit of NF-kappaB is critical for in vivo clearance of the noninvasive enteric pathogen Citrobacter rodentium. *Infect Immun* **76**, 4978–88 (2008).

12.   World Health Organization. *WHO Bacterial Priority Pathogens List, 2024: Bacterial Pathogens of Public Health Importance to Guide Research, Development and Strategies to Prevent and Control Antimicrobial Resistance.* (2024).

13.   Whitfield, C. & Trent, M. S. Biosynthesis and export of bacterial lipopolysaccharides. *Annu Rev Biochem* **83**, 99–128 (2014).

14.   Vinogradov, E. *et al.* Structures of lipopolysaccharides from Klebsiella pneumoniae. Eluicidation of the structure of the linkage region between core and polysaccharide O chain and identification of the residues at the non-reducing termini of the O chains. *J Biol Chem* **277**, 25070–81 (2002).

15.   Bronner, D., Clarke, B. R. & Whitfield, C. Identification of an ATP-binding cassette transport system required for translocation of lipopolysaccharide O-antigen side-chains across the cytoplasmic membrane of Klebsiella pneumoniae serotype O1. *Mol Microbiol* **14**, 505–19 (1994).

16.   Joensen, K. G., Tetzschner, A. M. M., Iguchi, A., Aarestrup, F. M. & Scheutz, F. Rapid and Easy In Silico Serotyping of Escherichia coli Isolates by Use of Whole-Genome Sequencing Data. *J Clin Microbiol* **53**, 2410–26 (2015).

17.   Choi, K.-H. *et al.* A Tn7-based broad-range bacterial cloning and expression system. *Nat Methods* **2**, 443–8 (2005).

18.   Clements, A. *et al.* Targeting subcapsular antigens for prevention of Klebsiella pneumoniae infections. *Vaccine* **26**, 5649–53 (2008).

19.   Kim, M. *et al.* Bacteria hijack integrin-linked kinase to stabilize focal adhesions and block cell detachment. *Nature* **459**, 578–82 (2009).

20.   Matute-Bello, G. *et al.* An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol Biol* **44**, 725–38 (2011).

21. Osuchowski, M. F. *et al.* Minimum quality threshold in pre-clinical sepsis studies (MQTiPSS):
an international expert consensus initiative for improvement of animal modeling in sepsis.
Intensive Care Med Exp **6**, 26 (2018).

22. Lam, M. M. C., Wick, R. R., Judd, L. M., Holt, K. E. & Wyres, K. L. Kaptive 2.0: updated capsule
and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex. *Microb*
Genom **8**, (2022).

23. Martin, M. J. *et al.* A panel of diverse Klebsiella pneumoniae clinical isolates for research and
development. *Microb Genom* **9**, (2023).

24. Old, D. C. & Adegbola, R. A. A new mannose-resistant haemagglutinin in Klebsiella. *J Appl*
Bacteriol **55**, 165–72 (1983).

25. Murphy, C. N. & Clegg, S. Klebsiella pneumoniae and type 3 fimbriae: nosocomial infection,
regulation and biofilm formation. *Future Microbiol* **7**, 991–1002 (2012).

26. Yang, J. *et al.* Transcriptional activation of the mrkA promoter of the Klebsiella pneumoniae
type 3 fimbrial operon by the c-di-GMP-dependent MrkH protein. *PLoS One* **8**, e79038 (2013).

27. Ma, C. *et al.* Citrobacter rodentium infection causes both mitochondrial dysfunction and
intestinal epithelial barrier disruption in vivo: role of mitochondrial associated protein (Map).
Cell Microbiol **8**, 1669–86 (2006).

28. Leverton, L. Q. & Kaper, J. B. Temporal expression of enteropathogenic Escherichia coli
virulence genes in an in vitro model of infection. *Infect Immun* **73**, 1034–43 (2005).

29. Riechmann, L., Clark, M., Waldmann, H. & Winter, G. Reshaping human antibodies for
therapy. *Nature* **332**, 323–7 (1988).

30. Lee, E.-C. *et al.* Complete humanization of the mouse immunoglobulin loci enables efficient
therapeutic antibody discovery. *Nat Biotechnol* **32**, 356–63 (2014).

31. Wong, J. L. C. *et al.* OmpK36-mediated Carbapenem resistance attenuates ST258 Klebsiella
pneumoniae in vivo. *Nat Commun* **10**, 3957 (2019).

32. Feldman, M. F. *et al.* A promising bioconjugate vaccine against hypervirulent Klebsiella
pneumoniae. *Proc Natl Acad Sci U S A* **116**, 18655–18663 (2019).

33. Walker, K. A., Treat, L. P., Sepúlveda, V. E. & Miller, V. L. The Small Protein RmpD Drives
Hypermucoviscosity in Klebsiella pneumoniae. *mBio* **11**, (2020).

34. Mavridou, D. A. I., Gonzalez, D., Clements, A. & Foster, K. R. The pUltra plasmid series: A
robust and flexible tool for fluorescent labeling of Enterobacteria. *Plasmid* **87–88**, 65–71
(2016).

35. Hopkins, E. G. D., Roumeliotis, T. I., Mullineaux-Sanders, C., Choudhary, J. S. & Frankel, G.
Intestinal Epithelial Cells and the Microbiome Undergo Swift Reprogramming at the Inception
of Colonic Citrobacter rodentium Infection. *mBio* **10**, (2019).

FIGURE 1.

A.

CR_WT: galF, rmlB, rmlD, rmlA, rmlC, wfgB, wzx, wfgC, wzy, wfgD, wfgE, gnd
CR O152 O-antigen biosynthesis gene cluster (*rfb* locus)

CR_KPO1: galF, wzm, wzt, wbbM, glf, wbbN, wbbO, kfoC, gnd
KP O2a O-antigen biosynthesis gene cluster (*rfb* locus)

glmS, wbbY, wbbZ
Glycosyltransferases converting O2a to O1v1 (O1)

CR origin
KP origin
1000bp

B.

KP O1 Ag — KP WbbY & WbbZ

CR O152 — CR O152 *rfb* locus

KP O2a Ag — KP O2a *rfb* locus

Core — *rfa* locus

Lipid A — Raetz pathway genes

CR_WT LPS

CR_KPO1 LPS

CR origin
KP origin

C.

KP_WT, CR_WT, CR_KPO1   KP_WT, CR_WT, CR_KPO1

α-CR O152   α-KP O1

D.

Day 8 stool shedding

CFU/GoF

**

CR_WT   CR_KPO1

E.

MERGE   CR   KP O1 Ag   DAPI

CR_WT

CR_KPO1

FIGURE 2.

A.

CR stool shedding

B.

| MERGE | CR | KP O1 Ag | DAPI |

CR_{KPO1}

$CR_{KPO1 +pespO}$

C.

α-KP antibody titres
(Day 30 post CR infection)

D.

α-KP antibodies
bind KP O1 Ag

○ $KP_{\Delta rfb}$
● KP_{WT}

E.

KP_{WT} anti-CR_{WT} IgG

KP_{WT} anti-KP O1 Ag

KP_{WT} anti-CR_{KPO1} IgG

zoom of (i)

zoom of (ii)

zoom of (iii)

FIGURE 3.

FIGURE 4.

A.

CR colanic acid biosynthesis locus

KP K2 capsule biosynthesis locus

CR origin
KP origin

1000bp

B.

Day 8 stool shedding

ns

C.

α-KP antibody titres
(Day 30 post CR infection)

ns

****

CR_WT
CR_KPK2
p*espO*

D.

E.

MFI signal key

F.

FIGURE 5.

A.

CR$_{T3F}$

P$_{map}$

glmS | mrkA | mrkB | mrkC | mrkD | mrkF

KP type 3 fimbriae (T3F) structural gene operon

CR origin
KP origin

1000bp

B.

qRT-PCR *mrkA* expression

2$^{-\Delta Ct}$

*

CR$_{T3F}$ p*espO* LB
CR$_{T3F}$ p*espO* DMEM
CR$_{WT}$ DMEM

C.

CR$_{WT}$ CR$_{T3F}$ MrkA

i   ii

zoom of (i)   zoom of (ii)

D.

Day 8 stool shedding

CFU/GoF

ns

CR$_{WT}$ CR$_{T3F}$ p*espO*

E.

α-KP antibody titres
(Day 30 post CR infection)

IgG (ng/ml)

ns   ****

CR$_{WT}$
CR$_{T3F}$ p*espO*

KP ΔmrkA-F   KP$_{WT}$

F.

KP$_{WT}$ anti-CR$_{WT}$ IgG   KP$_{WT}$ anti-KP MrkA ref   KP$_{WT}$ anti-CR$_{T3F}$ IgG

i   ii   iii

zoom of (i)   zoom of (ii)   zoom of (iii)

G.

Functional inhibition of biofilm
by CR$_{T3F}$ sera

AU (normalised to KP$_{\Delta mrkA-F}$)

*

KP$_{WT}$ no Ab
KP$_{WT}$ + CR$_{WT}$ sera
KP$_{WT}$ + CR$_{T3F}$ sera

H.

KP$_{WT}$ + CR$_{WT}$ sera   KP$_{WT}$ + CR$_{T3F}$ sera

20 μm   20 μm

FIGURE 6.

A. citrOgen strain generation ~7 days

choice of heterolgous Ag
gene editing vectors

galF | LPS O-Ag | gnd

ROD21991 | Capsule-Ag | galF

Pmap
glmS | Complex protein Ag

Single Ag or multiple Ags selected

✓ Validated CR insertion/substitution sites
✓ *in vitro* evaluation tools for QC

B. *In vivo* Ab production ~28 days

EspO boost
of mucosal
adherence

Continuous
heterologous Ag
presentation over
the infection cycle

Ab-mediated
CR clearance

CR stool shedding (10^0 to 10^{10}) vs dpi (0 to 30)

Blood collection

Antibody testing
~3 days

✓ Heterologous antigen Abs

C.

pAb applications
(sera)

Functional assays
e.g. serum killing
biofilm inhibition

Serotyping

mAb development
(B-cells)

Recombinant
mAb

Hybridoma
mAb

Evaluation of protection
(challenge studies)

citrOgen Platform

Downstream applications

FIGURE 1.

A.

CR$_{WT}$

galF — rmlB — rmlD — rmlA — rmlC — wfgB — wzx — wfgC — wzy — wfgD — wfgE — gnd

CR O152 O-antigen biosynthesis gene cluster (*rfb* locus)

CR$_{KPO1}$

galF — wzm — wzt — wbbM — glf — wbbN — wbbO — kfoC — gnd

KP O2a O-antigen biosynthesis gene cluster (*rfb* locus)

glmS — wbbY — wbbZ

Glycosyltransferases converting O2a to O1v1 (O1)

CR origin
KP origin

1000bp

B.

CR O152 — CR O152 *rfb* locus
Core — *rfa* locus
Lipid A — Raetz pathway genes

CR$_{WT}$ LPS

KP O1 Ag — KP WbbY & WbbZ
KP O2a Ag — KP O2a *rfb* locus
rfa locus
Raetz pathway genes

CR$_{KPO1}$ LPS

CR origin
KP origin

C.

α-CR O152   α-KP O1

D.

Day 8 stool shedding

CFU/GoF

**

CR$_{WT}$ CR$_{KPO1}$

E.

MERGE | CR | KP O1 Ag | DAPI

CR$_{WT}$

CR$_{KPO1}$

FIGURE 2.

A.

CR stool shedding

B.

C.

**α-KP antibody titres
(Day 30 post CR infection)**

D.

**α-KP antibodies
bind KP O1 Ag**

E.

FIGURE 3.

FIGURE 4.

A.

CR$_{WT}$

CR colanic acid biosynthesis locus

CR$_{KPK2}$

KP K2 capsule biosynthesis locus

CR origin
KP origin

1000bp

B. Day 8 stool shedding

ns

CFU/GoF

CR$_{WT}$ CR$_{KPK2}$ p$espO$

C. α-KP antibody titres
(Day 30 post CR infection)

ns   ****

IgG (ng/ml)

CR$_{WT}$
CR$_{KPK2}$ p$espO$

KP Δ$wcaJ$ KP$_{WT}$

D.

KP$_{WT}$ anti-CR$_{WT}$ IgG
KP$_{WT}$ anti-KPK2 ref
KP$_{WT}$ anti-CR$_{KPK2}$ IgG

zoom of (i)   zoom of (ii)   zoom of (iii)

E.

MFI signal key

62.5
31.3

KP$_{WT}$
KP$_{ΔwcaJ}$

F.

KP
anti-KPK2

KL30   KL119   KL110
KL64   KL2   KL52
KL107   KL15   KL113

FIGURE 5.

A.

CR$_{T3F}$

P$_{map}$

glmS · mrkA · mrkB · mrkC · mrkD · mrkF

KP type 3 fimbriae (T3F) structural gene operon

CR origin
KP origin

1000bp

B.

qRT-PCR *mrkA* expression

$2^{-\Delta Ct}$

CR$_{T3F}$ p*espO* LB
CR$_{T3F}$ p*espO* DMEM
CR$_{WT}$ DMEM

C.

CR$_{WT}$ · CR$_{T3F}$ MrkA

i · ii

zoom of (i) · zoom of (ii)

D.

Day 8 stool shedding

CFU/GoF

ns

CR$_{WT}$ · CR$_{T3F}$ p*espO*

E.

α-KP antibody titres
(Day 30 post CR infection)

IgG (ng/ml)

ns · ****

CR$_{WT}$
CR$_{T3F}$ p*espO*

KP ΔmrkA-F · KP$_{WT}$

F.

KP$_{WT}$ anti-CR$_{WT}$ IgG · KP$_{WT}$ anti-KP MrkA ref · KP$_{WT}$ anti-CR$_{T3F}$ IgG

i · ii · iii

zoom of (i) · zoom of (ii) · zoom of (iii)

G.

Functional inhibition of biofilm
by CR$_{T3F}$ sera

AU (normalised to KP$_{\Delta mrkA-F}$)

*

KP$_{WT}$ no Ab
KP$_{WT}$ + CR$_{WT}$ sera
KP$_{WT}$ + CR$_{T3F}$ sera

H.

KP$_{WT}$ + CR$_{WT}$ sera · KP$_{WT}$ + CR$_{T3F}$ sera

FIGURE 6.

citrOgen Platform

A. citrOgen strain generation ~7 days

choice of heterolgous Ag
gene editing vectors

galF → LPS O-Ag — gnd

ROD21991 → Capsule-Ag — galF

P_map
glmS → Complex protein Ag

Single Ag or multiple Ags selected

✓ Validated CR insertion/substitution sites
✓ *in vitro* evaluation tools for QC

B. *In vivo* Ab production ~28 days

CR stool shedding vs dpi

- EspO boost of mucosal adherence
- Continuous heterologous Ag presentation over the infection cycle
- Ab-mediated CR clearance

Blood collection

Antibody testing ~3 days

✓ Heterologous antigen Abs

Downstream applications

C.

pAb applications (sera)

Functional assays
e.g. serum killing
biofilm inhibition

Serotyping

mAb development (B-cells)

Recombinant mAb

Hybridoma mAb

Evaluation of protection (challenge studies)