citrOgen: a synthesis-free polysaccharide and protein antigen-presentation to antibody-induction platform

Short title

A *Citrobacter rodentium*-based antigen presentation platform

Joshua L C Wong[#,*], Julia Sanchez-Garrido[#], Jaie Rattle, Jonathan Bradshaw, Vishwas Mishra and Gad Frankel[*]

Department of Life Sciences, Imperial College London, SW7 2AZ, UK

[#]These authors contributed equally to this work

[*]For correspondence:

Electronic mail: joshua.wong@imperial.ac.uk and g.frankel@imperial.ac.uk

Keywords

Antibodies, *Citrobacter rodentium, Klebsiella pneumoniae,* type III secretion system (T3SS), EspO

1   **Abstract**
2
Existing technologies employed to generate antibodies against bacterial polysaccharides and proteins

rely on the availability of purified or synthetic antigens. Here we present a genetics-based platform

that utilises *Citrobacter rodentium* (CR), an enteric mouse pathogen, to both produce and present

complex heterologous polysaccharides and protein antigen complexes during natural infection. As

proof of concept, we use lipopolysaccharides (O), capsular polysaccharides (K) and type 3 fimbrial

(T3F) antigens expressed by the WHO critical priority pathogens *Klebsiella pneumoniae* (KP) and

Escherichia coli (EC). Following one infection cycle (28 days) CR induces specific IgG antibodies

against KPO1, ECO25b, KPK2 and KPT3F. We demonstrate that the antibodies are functional in

downstream applications including protection against pathogenic KP challenge, KP capsular

serotyping and KP biofilm inhibition. Whilst KP and EC antigens were used as prototypical examples,

this modular platform is now readily adaptable to generate antibodies against diverse polysaccharide

and protein antigens, with basic science, public health and therapeutic applications.

Introduction

The specificity of antibody/antigen (Ab/Ag) binding has been widely leveraged in science, biotechnology, and medicine. Both polyclonal (pAb) and monoclonal (mAb) antibodies are routinely employed across these disciplines and have shaped the face of the modern practice. There is, therefore, significant value in expanding the technologies available to generate new Abs that bind specific Ags of interest.

Abs are employed widely in the field of infectious disease with both diagnostic and, more recently, therapeutic applications. One key use in diagnostics is bacterial serotyping, which has played important roles in pathogen epidemiology. Therapeutic Abs are licensed for both the prevention and treatment of viral and bacterial infections in humans. Palivizumab, a humanised monoclonal IgG Ab that binds Respiratory Syncytial Virus, was the first mAb licensed for the prophylaxis of infectious disease in 1998[1]. Bezlotoxumab, a mAb against the enteric pathogen *C. difficile*, the sole mAb licensed in bacterial disease, was licensed in 2016 for the secondary prevention of *C. difficile* colitis [2]. More recently, during the SARS-CoV-2 pandemic both pAb administration (convalescent sera)[3] and mAbs against spike protein (e.g. bamlanivimab, etesevimab, casirivimab and imdevimab)[4] were used in the prevention and treatment of COVID-19 disease.

We currently have limited therapeutic options to combat the antimicrobial resistance (AMR) pandemic with both anti-bacterial Abs (⬚-bAbs) and vaccines representing promising modalities to tackle this contemporary threat. Vaccines rely on inducing an enduring memory response through the immunisation of naïve populations with target bacterial Ags. This allows adaptive host immunity to briskly reactivate upon subsequent pathogen exposure, clearing infection and reducing disease severity. In the case of extracellular bacteria, the effector arm of this response is centrally mediated by Abs. In contrast, ⬚-bAb therapeutics are produced *in vitro* and administered to patients as prophylaxis or in treatment settings. Uniquely, Fc-enhanced effector functions can be molecularly introduced to ⬚-bAbs that prolong intravascular half-life or promote hexamerisation and

complement-mediated cytotoxicity[5,6]. There are merits and disadvantages to both approaches, for

example, whilst ▯-bAb administration ensures that the effector arm of the host response is delivered

directly to the patient, it fails to induce immunological memory. However, memory responses are

often blunted in the high-risk populations that would benefit most from vaccination, such as the

elderly[7]. As such, both approaches are being actively pursued and could be used in combination or to

target different populations.

In target-specific ▯-bAb campaigns Abs that bind predetermined Ags are generated and selected for

downstream applications. Target choice can be informed *a priori* by pathogen epidemiology and

genomics, selecting bacterial Ags that are both widespread and conserved across a bacterial species.

Ags representing virulence determinants can also be selectively targeted as Ab-binding has the

potential to functionally disrupt pathogenic mechanisms[8]. The immunisation of experimental animals

or human subjects is a common method to generate target-specific Abs. This results in sera

containing pAb secreted from reactive B-cells. These cells can be immortalised or Ab-encoding

regions subcloned to produce mAbs *in vitro*. Irrespective of host species or Ag, the generation of

such responses is fundamentally reliant on a sufficient source of conformationally correct

immunogen. Generating these immunogens represents a key bottleneck in both target-specific Ab

generation[9] and vaccine development.

Polysaccharide Ags are diverse and typically composed of variable sugar subunits, each covalently

connected in the correct order by specific glycosidic links. These linkages can be linear or include

branching arrangements that determine the final polysaccharide structure. In isolation,

polysaccharides typically require carrier protein glycoconjugation to generate formulations that are

sufficiently immunogenic[9]. Production of sufficient quantities of protein Ags, which are typically

affinity purified following *de novo* synthesis in cells, is largely restricted to protein monomers and not

applicable to multiprotein complexes. The correct formation of the latter not only requires the

presence of chaperones and signal sequences, directing subunit assembly in the correct subcellular

space, but also necessitates conformationally correct monomers to interact with each other in the

correct stochiometric ratio. Bacterial genomes have compartmentalised the genes required to

synthesise both polysaccharide and multimeric protein Ags into operons, providing cloneable linear

DNA fragments that we exploit in the development of the technology we describe here.

Citrobacter rodentium (CR) causes colitis in mice with an infection strategy mirroring the human

pathogen Enteropathogenic *Escherichia coli* (EPEC). CR attaches intimately to intestinal epithelial

cells (IECs) forming attaching and effacing lesions and actin-rich pedestal-like structures, a

prerequisite for productive infection and colitis[10]. A major virulence factor enabling CR infection is

the type 3 secretion system (T3SS)[11] which translocates 31 effector proteins into IECs, facilitating CR

adherence and proliferation[12]. The resolution of CR infection coincides with the onset of IgG

secretion[13,14]. However, until this adaptive immune response clears the infection, CR maintains high

mucosal-associated burdens, inducing significant local inflammation with associated immune cell

recruitment[12]. We hypothesised that this natural host-pathogen context represents an evolutionarily

pre-optimised environment to present alternative (heterologous) Ags to the mouse host, thereby

generating mature target-specific Ab responses. This hypothesis forms the basis of our platform

which we term citrOgen.

Heterologous Ag delivery employing alternative bacterial chassis has been evaluated in previous

studies and largely focussed on live attenuated vaccine candidates. Common candidates include

Salmonella typhimurium and *Shigella spp*, where heterologous Ag expression aims to broaden the

vaccines range to now include pathogens beyond the attenuated candidate itself[15–17]. This field has

been hampered by the low stability of heterologous Ags *in vivo* and failure to induce protective

responses. Indeed, when the licensed live-attenuated *Salmonella typhi* vaccine strain Ty21a

expressing *Shigella sonnei* O2a Ag was administered to human subjects, only 19-27% (n=121

participants) had detectable *S. sonnei* IgG responses[18]. A subgroup of participants and controls then

underwent *S. sonnei* challenge and clinical shigellosis developed at the same rate in both groups,

providing no evidence of protection or amelioration of disease.

citrOgen is a flexible and modular *in vivo* Ag presentation platform co-opting CR to present

heterologous Ags. It represents a single unifying genetics-based system that can be used to generate

functional target-specific ⍺-bAb responses to divergent major Ag subtypes in mice. We selected four

structurally diverse and validated Ags from the WHO critical priority pathogens *Klebsiella*

pneumoniae (KP) and *E. coli* (EC) to test the platform [19]. The choice includes two surface

polysaccharide families, lipopolysaccharide (LPS – KP O1 & EC O25b) and capsular polysaccharide

(CPS – KP K2) and a protein complex, type 3 fimbriae (KP T3F). Substituting the endogenous CR O152

LPS O-Ag with KP O1 resulted in lower gut colonisation, which we overcome by overexpressing the

CR T3SS effector EspO. Functionally, EspO-augmented ⍺-KP O1 Ab responses potently protected mice

from sepsis-induced organ failure following KP challenge. Furthermore, we were able to rapidly

modify CR to express EC O25b eliciting specific ⍺-EC O25b IgG responses after a single infection cycle

(28 days) in mice. We then successfully generated ⍺-CPS Abs, which we employed to serotype a

diverse KP collection of 100 clinical isolates. Finally, we demonstrate citrOgen's ability to raise Abs to

a large cell surface filamentous structure, KP T3F, the regulated expression of which we designed to

coincide with CR attachment to host epithelia. These ⍺-KP T3F Abs effectively blocked T3F-mediated

KP biofilm formation. Together these results demonstrate the ability to rapidly generate functional

target-specific Abs against a broad range of major bacterial Ags.

Results

Heterologous KP O1 O-antigen expression causes an in vivo colonisation defect

Our goal was to express diverse heterologous Ags in CR to generate target-specific Abs in mice. The

rfb locus typically encodes all the genes required for O-Ag synthesis and export in Gram-negative

bacteria[20]. An exception is the KP O1 Ag, where *rfb*-derived O2 Ag is modified by the

glycosyltransferase activity of two enzymes, WbbY and WbbZ, which are encoded at a distinct

chromosomal position[21]. Moreover, KP employs an ABC-transporter instead of the more common

wzy-dependent O-Ag export process used by CR_{WT}[22].

We started by identifying the *rfb* locus in CR_{WT}, which is encoded on the reverse strand and flanked

by *galF* (5') and *gnd* (3') genes (Fig 1A). We used *in silico* serotyping to confirm that CR_{WT} encodes an

E. coli-like O152 Ag[23]. The endogenous O152 genes were replaced with those comprising the KP O2a

rfb locus (cloned en bloc) by homologous recombination and the *wbbYZ* gene cluster inserted into

the CR genome by Tn7 transposon mutagenesis (Fig 1A)[24]. Tn7 inserts at a neutral insertion site,

downstream of the *glmS* site *(attTn7* site) and we have already used this approach to successfully

express multiple different cargos in vivo[25]. This strain, CR_{KPO1}, contains all the genetic determinants

required to synthesise and export KP O1 Ag, albeit using a non-native ABC-transporter export

pathway. We did not modify the CR gene products that synthesise lipid A or core oligosaccharide (Fig

1B) and, therefore, CR_{KPO1} is predicted to produce a hybrid LPS molecule; the endogenous CR Raetz

pathway and CR *rfa* locus producing lipid A and core regions, respectively, with KP O1 Ag attached to

the native CR lipooligosaccharide scaffold (Fig 1B). We purified LPS from KP_{WT}, CR_{WT} and CR_{KPO1} and

confirmed the presence of a substituted O-antigen CR_{KPO1} by silver stain, with a banding pattern

distinct from CR_{WT} and similar to KP_{WT} (Supplementary Fig 1A). The specificity of O-antigen

substitution was confirmed Western blot. As anticipated ⊡-O152 pAb sera recognised CR_{WT} but was

not reactive against KP_{WT} and CR_{KPO1} (Fig 1C). We then used an ⊡-KP O1 mAb (C13)[26], which bound to

both KP_{WT} and CR_{KPO1}, confirming heterologous O-antigen expression in CR. The homologous

recombination vector employed to substitute the KP O2 locus was designed to include the

endogenous 5' CR promoter, ensuring that future O-antigen locus substitutions can be achieved by

using the same vector, simply modified with O-antigen genes of choice.

We tested if replacing CR O152 with KP O1 affected infectivity *in vitro* and confirmed that CR_{KPO1}, like

$CR_{WT,}$ induced actin-rich pedestal formation following the infection of mouse fibroblasts cells.

Furthermore, detectable KP O1 Ag expression was restricted to CR_{KPO1} (Supplementary Fig 1B). We

therefore proceeded to infect CD-1 mice with 10^9 colony forming units (CFUs) of CR_{WT} or CR_{KPO1} by

oral gavage (Fig 1D). At 8 days post infection (dpi), when stool CR shedding peaks, faecal CR_{KPO1} CFU

counts were significantly lower than CR_{WT} (Fig 1D). Staining of colonic sections revealed that whilst

CR_{KPO1} maintained heterologous KP O1 expression *in vivo,* there were substantially fewer adhered

bacteria compared to CR_{WT} (Fig. 1E, Supplementary Figure 1C). These data indicated an *in vivo*

colonisation defect imposed by heterologous Ag expression that needed to be addressed to develop

an optimal platform.

EspO overexpression restores CR_{KPO1} colonisation

A mutant in the *Shigella flexneri* T3SS effector *ospE*, a homologue of the CR effector *espO*, results in a

colonisation defect in a guinea pig gut infection model, as OspE suppresses epithelial detachment[27].

We therefore hypothesised that over expression of EspO would promote greater bacterial

attachment to IECs during infection. Indeed, while deletion of *espO* in CR_{WT} showed a tendency

towards decreased proportion of CR-associated IECs, overexpression of EspO increased the

proportion from 14.6% to 22.9% (Supplementary Fig 2A-C). We then overexpressed EspO in CR_{KPO1}

(CR_{KPO1}+p*espO*) and infected CD-1 mice; infections with CR_{WT} and CR_{KPO1} were used as controls.

Enumeration of faecal CFUs now revealed no difference between the faecal CFUs between CR_{WT} and

CR_{KPO1}+p*espO* at 8 dpi (Fig 2A). Staining of colonic sections demonstrated higher levels of

CR_{KPO1}+p*espO* adhesion to the epithelia compared to CR_{KPO1}, which maintained heterologous KP O1

expression *in vivo* (Fig 2B).

160 **CR_{KPO1}+p*espO* infection elicits KP O1 specific IgG antibodies**

161 We inoculated mice with CR_{KPO1}+p*espO* and CR_{WT} by oral gavage. After CR clearance (c. 28 dpi) we

162 collected blood and assessed α-KP_{WT} IgG antibody titres by enzyme-linked immunosorbent assay

163 (ELISA), using a capture plate coated with heat-killed KP_{WT}. While control CR_{WT} mice sera had

164 negligible α-KP_{WT} IgG binding, this increased significantly in sera from CR_{KPO1}+p*espO*-infected mice,

165 indicating that we had successfully raised α-KP_{WT} Abs (Fig. 2C). To confirm the specificity of the Ab

166 response we used an isogenic rough (absent O-antigen) KP mutant, $KP_{\Delta rfb}$, and assessed binding to

167 live bacteria by flow cytometry (Fig. 2D). Consistent with the ELISA data, sera from CR_{WT}-infected

168 mice bound to neither KP_{WT} nor $KP_{\Delta rfb}$. However, sera from CR_{KPO1}+p*espO*-infected mice bound

169 specifically to KP_{WT} and not $KP_{\Delta rfb}$, confirming the presence of α-KP O1 Abs (Fig 2D, Supplementary

170 Fig 2D). We then assessed the pattern of binding to KP_{WT} by immunofluorescence (IF) microscopy (Fig

171 2E). This confirmed the same pattern of binding of CR_{KPO1}+p*espO* sera and the reference α-KP O1 Ab

172 C13. No detectable signal was observed in sera from CR_{WT}-infected mice.

173 **Anti-KPO1 IgG Abs are functionally protective against KP pulmonary challenge**

174 We next tested the functionality of the CR_{KPO1}+p*espO*-induced α-KP O1 pAbs by lung challenge with

175 KP_{WT}. Mice that had been infected and cleared CR_{KPO1}+p*espO* were intubated and administered 500

176 CFU of KP_{WT} directly into the lungs. In preliminary experiments with CD-1 mice this KP_{WT} inoculum

177 was found to result in a progressive increase in KP CFUs in the lungs and blood resulting in a severe

178 disease phenotype at 72 h post infection (hpi) (Supplementary Fig 3 A-C). Mice that had been

179 infected by and subsequently cleared CR_{WT} were also challenged as controls. We evaluated infection

180 outcomes by assessing KP_{WT} replication and protection against host organ failure employing markers

181 of haematological, lung and renal injury in line with international consensus[28,29]. At 72 hpi KP_{WT}

182 burdens in the lungs of CR_{WT}-preinfected mice reached a median of c. 10^9 CFUs (Fig 3A). In contrast,

183 CR_{KPO1}+p*espO*-preinfected mice had significantly reduced lung CFUs with KP_{WT} only detectable in the

184 lungs of 30% (5/14) of mice. KP bacteraemia was absent in all CR_{KPO1}+p*espO*-preinfected mice in

185 comparison to 69% (9/13) of CR_{WT}-preinfected control mice (Fig 3B).

We then assessed if the reduction of KP_{WT} burden translated into a positive effect on host physiology.

Only 3 mice (20%, 3/15) in the CR_{KPO1}+p*espO*-preinfected group lost more than 10% of their starting

weight when challenged with KP_{WT}, whereas all mice in the CR_{WT}-preinfected group lost weight after

KP challenge with 86% (12/14) losing more than 10% of their starting weight (Fig 3C). Total white cell

counts, lymphocyte counts and platelets in the blood demonstrated significantly positive

improvements in CR_{KPO1}+p*espO*-preinfected mice compared to CR_{WT}-preinfected animals (Fig 3D-F).

Specifically, the extent of sepsis-induced leucopaenia, lymphopaenia and thrombocytopaenia were

abrogated by the CR_{KPO1}+p*espO* preinfection, while the number of neutrophils did not change

between the mouse groups (Fig 3G). Moreover, the positive acute phase reactants (APR) Serum

amyloid A and C-reactive protein were significantly lower in CR_{KPO1}+p*espO*-preinfected mice

compared CR_{WT}-preinfected animals (Fig 3H&I). This beneficial modulation in APRs translated into

higher levels of the negatively regulated APR, albumin, in CR_{KPO1}+p*espO* preinfected mice (Fig 3J).

We then used three markers to assess lung injury: IL1β and myeloperoxidase (MPO) in lung

homogenate and surfactant protein D (SPD) levels in the serum[28]. This revealed that CR_{KPO1}+p*espO*

preinfection resulted in significant reductions in IL1β and SPD levels compared to CR_{WT} preinfection,

indicating reduced lung inflammation and improved maintenance of the alveolar capillary barrier

respectively (Fig 3K&L). Preinfection with CR_{KPO1}+p*espO* also resulted in significantly reduced

neutrophilic alveolitis, evidenced by reduced lung MPO levels compared to CR_{WT}-preinfected mice

(Fig 3M). Finally, we assessed acute kidney injury by measuring serum creatinine and cystatin C

levels. While high levels of creatinine and cystatin C were found in KP_{WT} challenged CR_{WT}-preinfected

mice, improvements in both excretory renal function markers were seen in CR_{KPO1}+p*espO*-preinfected

mice (Fig 3N&O). These data indicate that ⍺-KP O1 Abs raised by citrOgen were protective against KP

challenge, restricting both bacterial replication and dissemination, and preventing KP sepsis-induced

organ failure.

citroGen can rapidly be adapted to present alternative O-antigens

Having overcome the fitness cost imposed by heterologous O Ag expression in CR and demonstrated

functional responses we wanted to further test the platform's modular nature and efficiency. To that

end, we selected *E. coli* O25b Ag (EC O25b), frequently encoded by ESBL-producing ST131 *E. coli*

strains (EC958 representing the prototype[30]) and used our vector system to generate CR_{ECO25b} (Figure

4A). O25b Ag was selected for two reasons, firstly, it is expressed an alternative WHO critical priority

pathogen and secondly, it is exported by the most common wzy-dependent export pathway. The

additional gene, *wzz*, which lies outside the *rfb* locus, is a chain length determinant in wzy-

dependent systems; we did not replace this and opted to keep the native CR_{WT}-encoded variant. We

confirmed specific O25b expression by agglutination, using reference ⬚-O25b pAb sera, which

agglutinated EC958 and CR_{ECO25b} but not $CR_{WT.}$ (Fig 4B). Silver staining of LPS preparations confirmed

the presence of O Ag in CR_{ECO25b}, which now runs at a higher molecular weight, as dictated by the

novel wzz(CR)/wzy(EC958) interaction (Supplementary Fig 4). Mice were orally infected with

CR_{ECO25b}+p*espO* and at 8 dpi no colonisation defect was observed compared to CR_{WT} (Fig4C). Blood

was collected from mice after CR clearance and sera used in an ELISA against EC958 and an isogenic

rough mutant (EC958$_{\Delta rfb}$). Absent Ab responses were detected from CR_{WT}-infected mice against either

EC958 or EC958$_{\Delta rfb}$. However, sera from mice infected with CR_{ECO25b}+p*espO* specifically reacted to

EC958 with absent binding to EC958$_{\Delta rfb}$ (Fig 4D). This specificity confirmed that citrOgen can be used

against additional O Ags, irrespective of export pathway. Moreover, it confirms that Ag selection to

antibody production is achievable in c. 35 days.

citroGen presentation of heterologous KP capsular polysaccharide antigen

We next explore expanding the citrOgen platform to generate ⬚-CPS Abs. KP_{WT} produces K2 CPS,

encoded by a c. 22Kb K2 *cps* locus, lying immediately upstream of the *rfb* locus in the KP_{WT} genome.

As upstream of the CR O152 *rfb* locus lie the genes involved in synthesis and export of the

extracellular polysaccharide colanic acid (Fig. 5A), we replaced the CR_{WT} genes between *ROD21991*

and *galF* (c. 24kb) with the KP KL2 operon from *wzi* to *manB* by homologous recombination,

generating strain CR_{KPK2}+p*espO* (Fig. 5A). We chose this insertion site as we anticipate that it will

tolerate other *cps* locus insertions in future applications of this technology. This established vector

can be used as the template together with *cps* locus of choice (e.g. KP *cps* loci vary from 10-30kb and

include internal promoters[31]).

We infected mouse fibroblasts to evaluate if the expression K2 capsule impacted on CR T3SS

functionality. CR_{KPK2} formed actin-rich pedestals and heterologous K2 expression was confirmed by

staining with reference K2 antiserum (Supplementary Fig 5A). This suggests that the K2 capsule does

not obstruct the T3SS needle, likely due to the fact that it is extended by a long EspA filament[32]. We

confirmed that EspO overexpression was required for optimal colonisation in the context of

heterologous CPS expression, as mice orally infected with CR_{KPK2} colonised at a lower level compared

to CR_{WT} at 8dpi (Supplementary Fig. 5B). We then proceeded to infect CD-1 mice with CR_{KPK2}+p*espO,*

or CR_{WT} as a control, which were shed at comparable levels (Fig 5B). Following CR_{KPK2}+p*espO* and CR_{WT}

clearance, we collected blood samples and ⍺-KP IgG levels were assessed by ELISA, using KP_{WT} and an

acapsular *wcaJ* mutant, $KP_{\Delta wcaJ}$. No binding to either KP_{WT} or $KP_{\Delta wcaJ}$ was observed with sera from

CR_{WT}-infected mice. In contrast, sera from mice infected with CR_{KPK2}+p*espO* bound to KP_{WT} but not to

$KP_{\Delta wcaJ}$ (Fig 5C). This indicated that KP K2 was successfully presented by the citrOgen platform *in vivo*

and that ⍺-KP IgG was specifically raised against the heterologous KP K2 antigen.

KP strains demonstrate broad diversity in CPS Ags with over 147 described to date in circulating

strains[31]. However, despite their key role in virulence and frequent use in genome-based

epidemiology, no reference antisera are currently being produced for *in vitro* applications. We used

sera derived from CR_{KPK2}+p*espO* infection, together with a reference K2 antiserum as a control, to

stain KP_{WT}. No detectable binding was seen in sera from CR_{WT}-infected mice. In contrast, binding was

seen when KP_{WT} was stained with the CR_{KPK2}+p*espO* sera, which phenocopied the pattern of binding

seen with the reference K2 antiserum (Fig. 5D). Together with the ELISA results, this provides strong

evidence for specific heterologous ⍺-KP K2 Abs induced by CR_{KPK2}+p*espO* infection.

KP serotyping using the citrOgen KP K2 sera

We next tested if the citrOgen KP K2 sera could be used for serotyping. To this end, we used a

publicly available collection of 100 KP isolates, representing 54 distinct CPS types determined by

genomic analysis[33]. We incubated the strains with citrOgen KP K2 sera and analysed binding by flow

cytometry (Fig 5E, Supplementary Fig. 5C). We included KP_{WT} and $KP_{\Delta wcaJ}$ as controls and measured

the median fluorescence intensity (MFI) across the collection. The citrOgen KP K2 sera specifically

bound KL2 isolates and KP_{WT} with minimal cross reactivity and absent binding to $KP_{\Delta wcaJ}$. This binding

was comparable to that observed when staining the KL2 strains with our reference K2 antiserum (Fig

5E, Supplementary Fig. 5D). To confirm the KL assignments, we selected 9 isolates for IF with the

reference K2 antisera (Fig 5F). We observed a pattern of binding consistent with capsule in the KL2

strain but not the surrounding non-KL2 strains. We concluded that the citrOgen KP K2 sera can be

used for serotyping and validated the collection's KL2 assignments.

citroGen presentation of heterologous protein complex antigens

We next assessed the utility of citrOgen in generating Abs against heterologous proteins, using the

well-conserved KP T3F as a model protein complex[34]. T3F are a KP virulence factor that mediate

binding to abiotic surfaces (such as plastic and glass) and are involved in the first steps of biofilm

formation. The major T3F subunit is MrkA which together with MrkD (adhesin tip) and MrkF

(unknown function) are externalised via a β-barrel transmembrane usher protein, MrkC, and

assemble into T3F on the KP cell surface (reviewed in Murphy et al. 2012[35]). In the periplasm a

chaperone, MrkB, binds MrkA to prevent ectopic pilin polymerisation. Therefore, MrkABCDF, are all

required for T3F biogenesis. T3F expression is tightly regulated in KP by an adjacent operon (*mrkHIJ*)

via the second messenger cyclic-di GMP[36]. To coordinate expression of the T3F with CR attachment

to IECs *in vivo,* we constructed a citrOgen strain in which *mrkABCDF* (excluding *mrkHIJ*) was driven by

the promoter of the T3SS effector *map* (CR_{T3F})[37]. The recombinant operon was inserted into the CR

glmS site (Fig 6A). We perceived two significant advantages of this strategy, firstly, maximising T3F

expression during intimate attachment to host cells and secondly to limit the metabolic burden of

T3F expression in CR. To confirm accurate gene regulation, we quantified *mrkA* transcript levels in

T3SS-suppressing (LB media, static growth) and T3SS-inducing (DMEM, static growth) conditions

(Supplementary Fig 6A)[38]. qRT-PCR analysis demonstrated low *mrkA* transcripts from $CR_{T3F}+pespO$

grown in LB which increased significantly in DMEM (Fig 6B), a change also observed for the

endogenous *map* gene used as a positive control (Supplementary Fig. 6A). No *mrkA* transcripts were

detected in the CR_{WT} negative control. We then confirmed that CR_{T3F} formed actin-rich pedestals *in*

vitro and expressed KP T3F by staining with an ⬚-MrkA pAb (Supplementary Figure 6B). Finally, we

tested if T3F conferred CR_{T3F} the ability to adhere to glass under T3SS inducing conditions. This

revealed a stark increase in the number of adhered CR_{T3F} observed on glass coverslips compared to

CR_{WT} when both were grown statically in DMEM, indicating functional heterologous T3F expression

(Fig 6C). Whilst this vector was designed to insert and provide stringent control of T3F expression it

can be readily be used for the expression of other operons encoding mono/multimeric protein Ags to

facilitate future iterations of this technology.

We started by confirming if EspO overexpression was required in the context of complex protein Ag

expression and infected mice with CR_{T3F} and CR_{WT}. A similar colonisation defect was evident at 8dpi

that we had encountered when KP O1 or KP K2 were expressed in CR_{WT} (Supplementary Fig. 6C). We

next used oral gavage to infect CD-1 mice with $CR_{T3F}+pespO$, which colonised similarly to CR_{WT} at 8

dpi (Fig 6D). We used the sera collected from mice who had cleared the infection for ELISA, with

plates coated with heat killed KP_{WT} and $KP_{\Delta mrkA-F}$, a deletion mutant in the T3F structural genes. Sera

collected from the control CR_{WT}-infected mice were not reactive against KP_{WT} or the isogenic mutant

(Fig 6E). Conversely, sera from $CR_{T3F}+pespO$ -infected mice bound KP_{WT}, with absent binding to

$KP_{\Delta mrkA-F}$. This data confirmed the ⬚-KP IgG antibodies were directed towards T3F. Using pooled sera

from the $CR_{T3F}+pespO$ infected mice to stain KP_{WT} revealed a binding pattern that matched that of

the reference ⬚-MrkA pAb (Fig 6F). As T3F mediate attachment to abiotic surfaces and biofilm

formation, we then tested if the sera from could prevent KP_{WT} attachment. We first pre-adsorbed the

sera from $CR_{T3F}+pespO$- and CR_{WT}-infected mice against PFA-fixed CR_{WT} to remove ⬚-CR Abs (10x 24h

adsorption cycles against fresh PFA-fixed CR_{WT}) (Supplementary Fig. 6D). This resulted in sera that

negligibly bound CR_{WT} (Supplementary Fig. 6E&F). We preincubated these two preadsorbed sera,

which had equivalent concentrations of total IgG (Supplementary Fig 6G), with KP_{WT} expressing sfGFP

and measured attachment after static growth at 37°C. Sera from CR_{T3F}+p*espO*-infected mice resulted

in 3-fold reduction in KP attachment compared to sera from CR_{WT}-infected mice (Fig. 6G&H). These

data demonstrate the generation of ⍰-KP T3F Abs that functionally inhibit surface attachment, the

first step in KP biofilm formation.

Discussion

LPS, CPS and proteins are common bacterial targets where generating specific Abs is desirable.

Whilst techniques already exist to generate such target-specific Abs, individually tailored and time-

consuming approaches are required to prepare the Ags. Here, instead, we co-opt the mouse

pathogen CR for this purpose and combine the steps of Ag preparation and immunisation into a

single platform that both expresses heterologous Ags and presents them continuously, in their native

conformation, to the host over the entire infection cycle (Figure 7). This was achieved by

recombination-based genome editing in which whole operons were cloned from KP and inserted or

substituted into the CR; thus, all genetically encoded components are inherently included, ensuring

that Ag assembly takes place in a fashion mirroring the pathogen of choice. It should be noted that,

whilst we achieve this for four diverse Ags, they all come from Gram-negative *Enterobacterales*,

which will have favoured our approach. Nevertheless, we were able to achieve specific and

functional Ab responses for LPS, CPS and a large protein complex. If we consider the latest iteration

of the WHO Bacterial Priority Pathogens List[19] then three out of four critical pathogens are Gram-

negatives, indicating the clear and present danger posed by this group of bacterial threats. The data

we present already supports using citrOgen for bacterial Ag targets in these key organisms. We

anticipate that further modifications will be required to synthesise and present Gram-positive

bacterial Ags.   As well as the ability to present complex polysaccharide structures and

multicomponent protein Ags in their native location and configuration, one key advantage of

citrOgen is time (Fig 7). In this work we have already validated substitution/insertion sites and

generated the appropriate genome editing vectors. Changing the inserts encoded by these vectors

(to alternative LPS, CPS and protein Ags) is routine and together with subsequent CR genome editing

can be achieved over a period of days. Whilst we cloned sequences directly from EC and KP DNA we

propose that this technology could also be applied to generate diagnostic or therapeutic Abs in an

outbreak scenario. The now routine use of *de novo* gene synthesis would enable us to engineer

strains based on early sequencing data alone. This would allow us to use this platform to generate

pAb sera in as little as 28 days- a complete CR infection cycle in mice. In more controlled settings we

envisage generating citrOgen strains encoding multiple Ags simultaneously. Ethically, this reduces the

number of experimental animals for use in research and development and would permit multiple

target-specific Ab responses to be obtained following a single infection.

We overcame two major obstacles to develop a viable platform for broader applicability. The first

was the potential fitness cost imposed by heterologous Ag expression. Indeed, we observed lower

gut colonisation in $CR_{KPO1,}$ CR_{KPK2} and CR_{T3F} compared to CR_{WT}, a defect that only becomes evident

under the pressure of the host-pathogen-microbiome environment. We overcame this obstacle by

overexpressing the T3SS effector EspO, which promotes IEC attachment and mucosal colonisation.

The second was the metabolic cost to CR of dysregulated heterologous protein expression which we

overcame by placing expression under the control of the endogenous CR promoter for the *map* gene.

This restricts heterologous protein expression to T3SS-inducing conditions and promotes

presentation at host mucosal surfaces, a strategy that can be used for the expression of any

heterologous protein, where expression needs to be tightly controlled and/or where regulatory

genes are absent in CR. Further modifications for expression can now easily be implemented given

the rich genetic toolset provided here or that already available for *E. coli,* a close relative of CR, which

shares similar genetic tractability.

In the case of KP O1 Ag we combined citrOgen with subsequent KP lung challenge, which allowed us

to quantitively evaluate protection from organ failure conferred by the ⍰-bAb response to this Ag (Fig

3). Given the current lack of correlates of protection for bacterial infections we propose that citrOgen

strains may represent an efficient tool to evaluate future proposed Ags.   Strains engineered to

express heterologous Ags (singly or in combination) can be used to evaluate Ab responses following a

CR infection cycle. In the same cohort of mice, subsequent pathogenic challenge can then evaluate

the protection conferred by the response. Expanding the translational applicability of current priority

bacterial infection models would significantly enhance the data we could achieve in this approach.

CR infections are host restricted to mice, conferring platform safety and preventing infection risk to

humans. This is especially important as future citrOgen strains could include additional virulence

factors as Ags for Ab generation. We currently use the citrOgen platform in conventional mice

generating murine immunoglobulins. Advances in molecular genetics have enabled the generation of

'humanised' Abs initially raised in non-human species and could be applied to the current iteration of

the platform[39]. This improves IgG tolerability and durability when administered to human patients

largely by reducing the immunogenicity provoked by non-human Ig regions. Furthermore, extensive

work in the field of mouse genetics has enabled the generation of fully human Abs in transgenic

animals[40]. Combining these transgenic mice with citrOgen strain infection would result in human Ig

induction in one CR infection cycle (Fig 7). Existing downstream workflows, such as hybridoma

generation or IgG loci subcloning are already well defined and employed to produce mAbs from B-

cells derived from these mice. Whilst our antibiotic treatment options are rapidly diminishing the

application of this technology may alleviate future infectious diseases deaths by generating new Ab-

based diagnostic tools and therapeutics.

 Figure Legends

Figure 1. Heterologous KP O1 O-antigen expression causes and *in vivo* colonisation defect.

A. The *rfb locus* in CR_{WT} encodes a O152-like *E. coli* O-antigen locus located between *galF* and *gnd*. In

CR_{KPO1} the CR_{WT} *rfb* locus was substituted with the genes encoding KP O2a O-antigen and *wbbYZ*

(converted KP O2a to KP O1v1) inserted at the 3' *glmS* site.

B. CR_{KPO1} expresses a hybrid lipopolysaccharide molecule with native CR lipid A and core

lipooligosaccharide linked to KP O1 Ag. The western blot is representative of 2 biological repeats.

C. Western blot confirmation of heterologous KP O1 Ag in CR_{KPO1} performed on purified LPS

preparations. α-O152 Ab are immunoreactive with CR_{WT} but not KP_{WT} or CR_{KPO1} whereas α-KP O1 mAb

(C13) is immunoreactive with both KP_{WT} and CR_{KPO1}.

D. At 8 dpi following oral gavage, faecal CFUs are significantly lower in CR_{KPO1} compared with CR_{WT}.

n=10 mice per group, 2 biological repeats. Central tendency: median. Mann-Whitney test. **,

p<0.01. GoF, gram of faeces.

E. Colonic histology was taken 8 dpi and stained for CR, KP O1 Ag and DAPI staining used as a

structural marker. CR_{WT} is observed confluently adhering to the luminal surface of the colon with no

KP O1 Ag expression detectable. Sparse staining for CR_{KPO1} is observed however, where present, KP

O1 Ag expression detectable. Scale bar = 200 µm. Images are representative of 3 biological repeats.

Figure 2. CR_{KPO1} overexpressing EspO maintains infectivity and induces specific anti-KP O1 Ag IgG

responses.

A. At 8 dpi *espO* overexpression in CR_{KPO1}+p*espO* reverses the colonisation defect observed in CR_{KPO1}

restoring colonisation to that of the parental CR_{WT} strain. n=15 mice per group, 3 biological repeats.

Central tendency: median. Ordinary one-way ANOVA with Dunnett's multiple comparison.

B. Colonic histology was taken 8 dpi and stained for CR, KP O1 Ag and DAPI staining used as a

structural marker. CR_{KPO1}+p*espO* now confluently adhering to the luminal surface with KP O1 Ag

expression detectable and co-localising to CR staining. Sparse staining is observed for CR_{KPO1} with KP

O1 Ag expression. Scale bar = 200 µm. Images are representative of 3 biological repeats.

C. At 30 dpi blood was taken from the tail vein of mice that cleared CR_{WT} and CR_{KPO1}+p*espO*. ELISA,

using heat-killed KP_{WT} (expressing KP O1 Ag) revealed α-KP_{WT} IgG in mice following infection with

CR_{KPO1}+p*espO* but not CR_{WT} controls. Central tendency: mean. Mann-Whitney test.

D. Sera from mice was tested for binding to live KP_{WT} and $KP_{\Delta rfb}$. Binding was assessed by flow

cytometry and expressed as the percentage of the population positive for mouse IgG (mIgG+ve). Sera

from CR_{WT}-infected mice demonstrated absent binding to either KP_{WT} or $KP_{\Delta rfb}$, whereas sera from

CR_{KPO1}+p*espO* -infected selectively bound KP_{WT} expressing KP O1 Ag. Central tendency: mean.

Unpaired T-test.

E. Sera from CR_{WT} and CR_{KPO1}+p*espO* -infected mice was used to stain KP_{WT} and image by

immunofluorescence microscopy. An α-KP O1 Ag mAb (C13) was used as a positive control for KP O1

Ag staining. CR_{WT} derived sera results in no detectable staining of KP_{WT} whereas sera from CR_{KPO1}

+p*espO* -infected mice and the control C13 mAb phenocopy each other with membrane-associated

staining of KP_{WT}.

436
437   ns=non-significant; *, p<0.5; ****, p<0.0001.
438
439   **Figure 3. Antibodies to KP O1 O-antigen protect against KP pulmonary challenge**.
440
441   Mice that had cleared pre-infection with CR_{WT} and CR_{KPO1}+p*espO* were challenged by intratracheal
442   infection with 500 CFU of KP_{WT}. At 72 hpi we analysed KP replication (A&B) and host-derived severity
443   indicators (C-O). n=15 mice per group (starting) over 3 biological repeats.
444
445   A&B. KP_{WT} replication and bacteraemia are significantly reduced in the lungs (A) and blood (B) of
446   CR_{KPO1}+p*espO*-preinfected mice compared to CR_{WT} controls. Central tendency: median. Mann-
447   Whitney test.
448
449   C. KP_{WT} induces significant weight loss in CR_{WT}-preinfected mice which is reversed in CR_{KPO1}+p*espO*
450   preinfection. Central tendency: mean, error: standard deviation (SD). Mann-Whitney test.
451
452   D-F. Clinical haematological markers of disease severity are significantly improved in CR_{KPO1}+p*espO* -
453   preinfected mice compared to CR_{WT} controls. Total white cell counts (D), lymphocyte counts (E) and
454   platelets counts (F) are significantly lower in CR_{WT}-preinfected controls indicating increased diseases
455   severity. The innate responders, neutrophils (G) are unchanged between the CR_{WT} and CR_{KPO1}+p*espO*
456   preinfected groups. Central tendency: mean, error: SD. Unpaired T-test.
457
458   H-J. The acute phase reactants (Serum amyloid A (H), C-reactive protein (I) and Albumin (J))
459   demonstrate a reduced inflammatory state in CR_{KPO1}+p*espO* -preinfected mice compared to CR_{WT}
460   controls. Central tendency: mean, error: SD. H&J). Mann-Whitney test, I) Unpaired T-test.
461
462   K-M. Acute lung injury markers (lung homogenate IL1β (K), serum surfactant B (L) and lung
463   homogenate myeloperoxidase (M)) are improved by pre-infection with CR_{KPO1}+p*espO* compared to
464   CR_{WT} preinfection. Significant reductions in lung iIL1β, improvement in the alveolar capillary barrier
465   (surfactant protein D) and neutrophilic alveolitis (myeloperoxidase) indicate α-KP O1 Abs protect
466   against lung injury. Central tendency: mean, error: SD. K&M) Mann-Whitney test, L) Unpaired t test.
467
468   N&O. Acute kidney injury and excretory renal failure are reduced in mice preinfected with
469   CR_{KPO1}+p*espO* compared to CR_{WT}. Serum creatinine (N) and serum cystatin C (O) are reduced in KP_{WT}
470   infected mice who have cleared preinfection with CR_{KPO1}+p*espO* compared to CR_{WT}. Central tendency:
471   mean, error: SD. Unpaired t test.
472
473   ns=non-significant; *, p<0.5 **; p<0.01, ***, p<0.001; ****, p<0.0001.
474
475   **Figure 4. Rapid citrOgen modification to present *E. coli* O25b O-antigen**
476
477   A. The CR *rfb locus* editing vector can be rapidly modified to insert the O25b *rfb* locus from EC958
478   (ST131 ESBL producing strain) into the CR genome, generating CR_{ECO25b}.
479
480   B. Agglutination of EC958 and CR_{ECO25b} with polyclonal sera to O25b O-antigen. CR_{WT} forms a tight
481   pellet indicating no agglutination of this strain. In control wells (PBS) all strains form tight pellets in
482   the absence of agglutination.
483
C. At 8 dpi following oral gavage, faecal CFUs are not significantly lower in CR_{ECO25b}+p*espO* compared
with CR_{WT}. n=10 mice per group, 2 biological repeats. Central tendency: median. Unpaired T-test.
GoF, gram of faeces.

D. At 30 dpi blood was taken from mice that cleared infection with CR_{WT} or CR_{ECO25b}+p$espO$. An ELISA

was conducted against heat-killed $EC958_{WT}$ (expressing O25b) and an isogenic rough mutant

($EC958_{\Delta rfb}$). CR_{WT} infected mice had no detectable IgG responses to $EC958_{WT}$ or $EC958_{\Delta rfb}$.

CR_{ECO25b}+p$espO$ infected mice had IgG responses specifically against $EC958_{WT.}$ n=10 mice per group, 3

biological repeats. Central tendency: mean. Unpaired t tests.

ns=non-significant; ****, p<0.0001.

Figure 5. citrOgen derived anti-KP K2 antibodies can be used for serotyping.

A. The *cps* in KP_{WT} is c.22kb long with 15 open reading frames. This was inserted into CR_{WT} by

substitution of the colanic acid biosynthesis locus generating the strain CR_{KPK2}.

B. At 8 dpi faecal CR_{KPK2}+p$espO$ levels are similar to CR_{WT}. Central tendency: median. n=15 mice per

group, 3 biological repeats. Unpaired T-test, ns=non-significant. GoF, gram of faeces.

C. At 30 dpi blood was taken from mice that cleared infection with CR_{WT} or CR_{KPK2}+p$espO$. An ELISA

was conducted against heat-killed KP_{WT} (expressing KP K2) and an isogenic acapsular mutant ($KP_{\Delta wcaJ}$).

CR_{WT} infected mice had no detectable IgG responses to KP_{WT} or $KP_{\Delta wcaJ}$. CR_{KPK2}+p$espO$ infected mice

had IgG responses specifically against $KP_{WT.}$ n=15 mice per group, 3 biological repeats. Central

tendency: mean. Mann Whitney test.

D. Sera from CR_{WT} and CR_{KPK2}+p$espO$-infected mice were used to stain KP_{WT} and image by

immunofluorescence microscopy. An α-KP K2 pAb was used as a positive control. CR_{WT} derived sera

results in no detectable staining of KP_{WT} whereas sera from CR_{KPK2}+p$espO$-infected mice and the α-KP

K2 pAb phenocopy each other with punctate membrane-associated staining of KP_{WT}.

E. Sera from CR_{KPK2}+p$espO$-infected mice were used to serotype a collection of 100 diverse KP strains

with serotypes assigned by genomic analysis. Each box represents one strain with the cps (KL) type

described. The boxes are coloured according to the strength of binding by CR_{KPK2}+p$espO$-infected

mice sera, which was assessed by flow cytometry. The median fluoresence intensity (MFI) key is

shown in the bottom right together with staining of KP_{WT} and $KP_{\Delta wcaJ}$ controls. A box, centred on a KL2

isolate is shown.

F. The strains contained within the highlighted box in (E) were stained with the reference α-KP K2

pAb and imaged by immunofluorescence microscopy. The KL2 strain in the centre demonstrate

punctate membrane staining in keeping with K2 capsule. The other strains (non-KL2) demonstrate no

staining. Bacterial cell membranes were counterstained with FM-4-64 for imaging.

ns=non-significant; ****, p<0.0001.

Figure 6. citrOgen derived anti-KP T3F antibodies block biofilm formation.

A. To generate a strain which expresses KP T3F the structural genes *mrkABCDF* were inserted at the

3' *glmS* site in CR under the control of the T3SS effector protein Map promoter. This generated the

strain CR_{3F}.

B. *mrkA* transcripts were assessed by qRT-PCR from bacterial cultures of CR_{T3F} under T3SS

suppressing (LB) and inducing (DMEM) conditions. CR_{WT} is also shown as a negative control. T3SS

inducing conditions specifically induce *mrkA* transcription in CR_{T3F}. 4 biological replicates, central
tendency mean, error bars SEM. Paired T-test between LB and DMEM-grown CR_{T3F}.

C. mCherry expressing CR_{T3F} and CR_{WT} were grown in T3SS inducing conditions on glass coverslips
bottomed slides. Following 6 h attachment to glass was assessed and the adhered bacteria were
stained for MrkA. A dense lawn of CR_{T3F} is observed binding to the glass slide with MrkA detected on
the cell surface indicating functional surface KP T3F expression. The parental CR_{WT} strain adheres to
glass slides only sparsely.

D. At 8 dpi faecal CR_{T3F}+p*espO* levels are similar to CR_{WT}. Central tendency: median. n=15 mice per
group, 3 biological repeats. Mann-Whitney test, ns=non-significant. GoF, gram of faeces.

E. At 30 dpi blood was taken from mice that cleared infection with CR_{WT} or CR_{T3F}+p*espO*. An ELISA
was conducted against heat-killed KP_{WT} (expressing KP T3F) and an isogenic mutant lacking T3F
expression ($KP_{\Delta A\text{-}F}$). CR_{WT} infected mice had no detectable IgG responses to KP_{WT} or $KP_{\Delta mrkA\text{-}F}$. CR_{T3F}
+p*espO* infected mice had specific IgG responses against $KP_{WT.}$ n=15 mice per group, 3 biological
repeats. Central tendency: mean. Mann Whitney test.

F. Sera from CR_{WT} and CR_{T3F}+p*espO*-infected mice were used to stain KP_{WT} and image by
immunofluorescence microscopy. An α-MrkA pAb was used as a positive control. CR_{WT} derived sera
results in no detectable staining of KP_{WT} whereas sera from CR_{T3F}+p*espO*-infected mice and the α-
MrkA pAb phenocopy each other with spiculated membrane-associated staining of KP_{WT}.

G. CR preadsorbed sera from CR_{WT} and CR_{T3F}+p*espO*-infected mice were preincubated with KP_{WT} or
$KP_{\Delta mrkA\text{-}F}$ expressing sfGFP. A no-antibody control was used as an additional control. Following 6 h of
growth in M9 minimal media sfGFP signal was assessed on a plate reader and presented as the KP_{WT}
binding normalised to $KP_{\Delta mrkA\text{-}F}$ binding. The preadsorbed sera collected from CR_{T3F}+p*espO*-infected
mice specifically inhibited binding compared to $CR_{WT.}$ 4 biological replicates, central tendency: mean,
error bars: SEM. Repeated-measures One-way ANOVA corrected with Dunnett's multiple
comparisons test.

H. The biofilm formation from (G) was assessed by imaging using sfGFP fluorescence. A large
reduction in adhered KP_{WT} is observed when CR_{T3F} p*espO*-infected cross adsorbed sera was incubated
with KP_{WT} compared to CR_{WT} cross adsorbed sera.

ns=non-significant; *, $p < 0.5$; ****, $p < 0.0001$.

Figure 7. The citrOgen platform as a versatile, quick and cost-efficient method of Ab production
and multiple downstream applications

A. citrOgen strain generation. Selected Ag sequences are incorporated into the appropriate gene
editing vectors designed for optimal chromosomal expression of heterologous O-Ag, capsule
or complex protein Ags, the latter under the map promoter to ensure timely expression
during mucosal infection. p*espO* is then transformed to finalise the new citrOgen strain. QC
checks include assessment of heterologous expression and pedestal formation *in vitro* (as shown in
Supp Fig 1A, 4A and 5B) before proceeding to *in vivo* Ab production.

B. In vivo Ab production. The CR infection cycle and immune responses are co-opted by the citrOgen
platform: (i) EspO-mediated boosting of CR attachment to intestinal epithelial cells, (ii) continuous
exposure of immune cells to the heterologous Ags, (iii) B cell affinity maturation results in Abs,
primarily IgG, which mediate CR clearance by 28 dpi and can be obtained as pAb sera from blood.

C. Downstream applications.

Following citrOgen infection/immunisation, the functionality of Abs can be assessed *in vitro* or by *in*
vivo challenge studies, the pAb sera can be used for serotyping and the B cells can be used for mAb
production, either by using hybridoma technology to immortalise single-cell cloned B-cells or by
expressing heterologous Ag-specific Ab sequences in cell lines to obtain recombinant mAb.

Ab: Antibodies; mAb: monoclonal antibodies; pAb: polyclonal antibodies; Ag: Antigen; CR:
Citrobacter rodentium.

Materials and Methods

Bacterial culture and antibiotics

Bacteria were routinely cultured in LB Broth (Miller) (84649.05000, VWR, UK). Saturated (overnight, 18 hour) cultures took place by inoculating broth and incubation at 37°C and shaking at 200rpm.

Antibiotics were added to LB broth or LB broth solidified with agar (1.10283.0500, VWR, UK) at the following concentrations:

Gentamicin 10mcg/ml (cloning) (G1272, Merck, UK)
Streptomycin 50mcg/ml (cloning) (S6501, Merck, UK)
Kanamycin 50mcg/ml (cloning) Merck, UK)
Nalidixic acid 50mcg/ml (selecting CR and CFU enumeration from *in vivo* sample) (N4382, Merck, UK)
Rifampicin 50mcg/ml (selecting KP and CFU enumeration from *in vivo* sample) (R3501, Merck, UK)

Cloning

High-fidelity polymerase chain reaction was conducted using Phanta Flash (2x Phanta Flash Master Mix (Dye Plus), Vazyme, China). Screening and colony polymerase chain reaction was conducted using Rapid Taq (2x Rapid Taq Master Mix, Vazyme, China).

PCR purification (Monarch PCR & DNA Cleanup Kit (5ug), NEB) and plasmid extraction were conducted with kit-based methods.

All mutagenesis vectors were generated by PCR products ligation via Gibson Assembly (NEB, UK) according to the manufacturer's guidance.

Genome editing

The whole genome references sequences used for CR strain ICC168 (Nalidixic acid-sensitive version of ICC169) is FN543502. The whole genome reference sequence used for ICC8001 is the closely related ATCC43186-derivitive KPPR1 is CP009208.1. A list of strains generated in this work can be found in Supplementary Table 1.

Two methods of genome editing were conducted in this work. Insertions at the CR 3' *glmS* site were conducted using Tn7 mutagenesis with methods published in Choi et al. Nat Methods 2005 Jun;2(6):443-8[24]. Seamless, markerless genome editing at other sites was conducted by homologous recombination based on the pSEVA612S system (https://seva-plasmids.com/) in both CR and KP as previously published in detail[12,25]. To improve the efficiency of pSEVA612S for Gibson Assembly we further modified this plasmid by reversing the sequence of the 3' ISceI endonuclease digestion site generating pSEVA612S(R). This modification improves efficiency by reducing homology-based empty vector recombination.

All mutagenesis vectors are summarised in Supplementary Table 2 and are publicly available at https://doi.org/10.6084/m9.figshare.c.7869659.v1. In the case where more than one vector was used these are listed against the respective strain. To generate CR_{KPK2} we initially cloned a construct (pSEVA612S(R)_CRColanic::K2cpsΔwbbJ_ORF) without the genes wbbJ and an annotated open reading frame of unknown function[41]. We subsequently inserted these genes into our final CR_{KPK2} strain (pSEVA612S(R)_ΔwbbJ::Ins_wbbJ_ORF) as new data become available[42] and included the native KP promoter (pSEVA612S(R)_K2:col_Pvariant_Porf2).

644
645 Transformation of pACYC184_espO was conducted using room temperature competent cells
646 according to our previously published protocol[25].

647
648 All primers are appended to GenBank files and publicly available at
649 https://doi.org/10.6084/m9.figshare.c.7869659.v1.

650
651 **Sequencing**

652
653 Sanger sequencing of PCR products and plasmids was conducted by Eurofins Genomics.

654
655 **In silico serotyping**

656
657 In silico serotyping of CR was conducted using SerotypeFinder 2.0
658 (https://cge.food.dtu.dk/services/SerotypeFinder/) on genome FN543502.

659
660 **Crude LPS preparation and Silver stain**

661
1 ml of overnight bacterial cultures normalised to OD_{600} of 1.2 were pelleted by centrifugation at
10,000 g and resuspended in 200 µl Laemmli buffer, followed by boiling at 100°C for 5 minutes.
Proteinase K was added to a final concentration of 1 mg/ml, and samples were incubated at 60°C for
2 hours. Subsequently, 5% β-mercaptoethanol was added, and samples were incubated for an
additional 5 minutes. Crude LPS samples were separated by SDS- polyacrylamide gel electrophoresis
(PAGE) using a 12% gel, and the resulting gels were fixed in 5% acetic acid/40% isopropanol
overnight. Silver staining was then performed using the SilverQuest™ Silver Staining Kit (LC6070,
Thermo Fisher Scientific) according to the manufacturer's protocol. All steps were carried out at
room temperature with gentle agitation. Images were acquired on a ChemiDoc XRS+ and visualised
using Image Lab software (Bio-Rad Laboratories).

672
673 **LPS extraction and Western Blotting**

674
675 LPS was extracted from 3OD units of saturated overnight cultures using a modified phenol method
676 kit (iNtRON Biotechnology, Korea) according to the manufacturer's guidance with no modification.

677
Samples were mixed with 5x Laemmli buffer and 5% 2-mercaptoethanol and separated by SDS–PAGE
using a 12% acrylamide gel and tris-glycine buffer system. and transferred to polyvinylidene
difluoride membranes (#1620177, Bio-Rad Laboratories) using a TransBlot semi-dry electrophoretic
transfer machine (Bio-Rad). Membranes were blocked for 2 h at room temperature (RT) in 10% fat-
free milk PBS–0.1% Tween 20 (PBS-T) and incubated overnight at 4°C with 1:500 anti-O152 or 1:2000
anti-KP O1 (Supplementary Table 3) in PBS-T with 5% BSA (A3059-50G, Sigma). Secondary antibodies
(Supplementary Table 3) were added diluted 1:10000 in 5% fat-free milk in PBST and incubated for 1
h at RT. Immunoblots were developed with Clarity Western enhanced chemiluminescence (ECL, Bio-
Rad Laboratories) and images were acquired on a ChemiDoc XRS+ and visualised using Image Lab
software (Bio-Rad Laboratories).

688
689 **In vitro $CR_{KPO1,}$ $CR_{KP K2}$ and CR_{T3F} infections of fibroblasts**

690
Murine Swiss 3T3 fibroblasts (ATCC® CCL-92™) were maintained in DMEM containing 4500 mg l^{-1}
glucose (D6546, Sigma) and supplemented with 10% fetal calf serum (F9665, Sigma) and 2 mM
glutaMAX (#35050061, ThermoFisher Scientific) at 37°C in 5% (v/v) CO_2 and seeded on glass
coverslips in 24-well plates (density $2x10^5$ cells/well) for immunofluorescence. Indicated strains of CR

expressing mCherry from a pUltra plasmid[43] were grown to saturation in LB supplemented with
gentamicin at 37°C with 200 rpm agitation for 8 hours before subculturing 1:500 for 16 hours in LB or
DMEM with 1000 mg l^{-1} glucose (# D5546, Sigma) at 37°C, 5% (v/v) CO_2, static. Cells were infected
with the static culture at a multiplicity of infection (MOI) of 100, confirmed by retrospective plating.
The plate was centrifuged at 1000 rpm for 5 min at RT to synchronize infection and incubated for 4
hours at 37°C in a 5% CO_2 atmosphere. The infection was stopped with three washes with sterile
phosphate-buffered saline (PBS) and cells were fixed in PBS with 4% (w/v) paraformaldehyde for 15
mins at RT and washed three times in PBS. Cells were then permeabilized in 0.1% (v/v) of Triton X100
in PBS for 10 min, incubated in 50 mM NH_4Cl in PBS as a quenching agent for 10 min, washed in PBS,
and blocked in 3% BSA in PBS for 1 h at RT before staining with one of the following primary
antibodies in 1% BSA-PBS for 1 h at RT: anti-KPO1 C13 (1:200), anti-mrkA (1:100) or anti-KP K2
(1:100) (**Supplementary Table 3**). Coverslips were washed in PBS and incubated with Alexafluor488-
conjugated secondary antibodies (1:200), Tetramethylrhodamine (TRITC)–conjugated phalloidin
(1:500) and 4ᐧ,6-diamidino-2-phenylindole (DAPI) (1:1000) to label the heterologously-expressed
antigen, F-actin and DNA, respectively (**Supplementary Table 3**) in 1% BSA-PBS for 1 h. Coverslips
were washed in PBS and once in water before being mounted using Gold Pro-Long Anti-fade medium
(#P36930, **ThermoFisher Scientific**) and left to cure overnight at RT in the dark. Immunofluorescence
images were acquired using a Zeiss AxioImager Z1 microscope and processed using ZEN3.1 software
(ZEN lite, Zeiss).

714
715 **Reverse transcription quantitative PCR (RT-qPCR)**
716
The indicated strains of CR were grown to saturation overnight in LB supplemented with the
corresponding antibiotics. 0.5 ml of the overnight culture were washed once in PBS, incubated in
RNAprotect Bacteria reagent (#76506, Qiagen) for 10 mins at RT and centrifuged at 3000 xg for 5
min. The bacterial pellet was digested in 100 µl TE buffer ((30 mM Tris·Cl, 1 mM EDTA, pH 8.0) with
15ᐧmg/ml Lysozyme (L6876, Sigma-Aldrich) and 20 µl Proteinase K (#19131, Qiagen) for 10ᐧmin
according to manufacturer's guidelines. RNA was isolated using the RNeasy minikit (Qiagen)
following the manufacturer's instructions, and RNA concentration and purity was determined using a
Nanodrop 2000 spectrophotometer (ThermoFisher Scientific). 1 µg of RNA was treated with RNase-
free DNase (#M6101) for 1 h at 37 °C and cDNA was then synthesized using a Moloney murine
leukemia virus (M- reverse transcriptase (M-MLV, #M1705) with random primers (#C1181) following
the manufacturer's instructions (all from Promega). To confirm the absence of undigested DNA, a
reaction mixture without the M-MLV reverse transcriptase was also included (NRT). qPCR was
performed using the Power Up SYBR Green master mix (#A25778, ThermoFisher Scientific) and the
relevant primers, whose primer efficiency had been evaluated previously and determined to be 95-
105% (**Supplementary Table 4**). The assay was run on a Quantstudio 1 System (Applied Biosystems,
ThermoFisher Scientific) and results were analysed using the QuantStudio Design & Analysis
Software v1.5.2 (Applied Biosystems, ThermoFisher Scientific). The relative expression level of each
mRNA was evaluated using the 2-ΔCtmethod, where Ct is the threshold cycle and 16S rRNA is used
as the control.

736
737 **O25 antisera agglutination**
738
739 Heat-killed agglutination was performed on the indicated strains after overnight growth in LB. The
740 anti-O25 sera (85022, SSI Diagnostica) was used according to the manufacturer's protocol. Images
741 were acquired on a ChemiDoc XRS+ and visualised using Image Lab software (Bio-Rad Laboratories).
742
743
744 **Whole bacterial cell ELISA**

745
To prepare heat-killed KP the indicated strains were grown overnight in LB at 37°C at 200 rpm. The
culture was then normalised to an optical density of 1 at 600 nm in sterile PBS, heat inactivated in
the presence of EDTA-free protease inhibitor cocktail (#4693159001, Sigma) at 60°C for 1 h and
stored at 4°C until use (short-term). ELISA Flat-Bottom Maxisorp Immuno 96- Well Plates
(#10547781, Fisher Scientific) were incubated with 100 µL/well heat-killed CR for our serum samples
or with 100 µL/well goat anti-mouse IgG antibody for the standards (2 µg/µL) (Supplementary Table
3) overnight at 4°C. Plates were washed three times with PBS-T, blocked for 1 h at RT with 5% BSA-
PBS-T, and incubated overnight at 4°C with serum samples or serially diluted mouse IgG
(Supplementary Table 3) as standard. Plates were washed as described above and incubated with
horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG antibody (Supplementary Table 3) for
1 h at room temperature. Plates were washed and developed by the addition of 100 µL 3,3′,5,5′-
tetramethylbenzidine (TMB; Thermo Fisher Scientific) for 5-15 mins before stopping the reaction
with 2 N H_3PO_4. The absorbances at 450 and 540 nm were measured in a plate reader (FLUOstar
Omega; BMG Labtech), and anti-KP IgG levels were quantified in samples by comparison to a mouse
IgG standard curve.
761
Determination of O-antigen-binding specificity: KP_{WT} expressing mCherry and $KP_{\Delta rfb}$ expressing sfGFP
were grown as saturated overnight cultures in LB supplemented with gentamicin. On the following
day, both strains were mixed 1:1 (v/v) after confirming the had grown to the same optical density
(600nm) and transferred to a V-bottom 96-well plate (Greiner Bio-one) for staining. Single strain
controls were also included to ensure correct gating during flow cytometry analysis. Samples were
first blocked in 5% BSA/PBS for 1 h at RT shaking, centrifuged at 4000 rpm, 4 °C, 10 mins and then
incubated overnight with sera from CR_{WT} or CR_{KPO1}+pespO-infected mice diluted 1in100 (v/v) in 1%
BSA/PBS. Bacterial samples were centrifuged, washed twice in cold PBS and resuspended in 1%
BSA/PBS containing AlexaFluor647-conjugated anti-mouse IgG (1:200, Supplementary Table 3) for 1 h
at RT shaking. Cells were washed twice in cold PBS, fixed in 1% PFA/PBS and kept in the dark until
flow cytometry analysis. Samples were acquired (minimum of 100000 bacterial cells per sample) on a
Cytek Amnis CellStream and data was analysed using FlowJo v10.10.0 (gating strategy shown in
Supplementary Fig 3A).
775
K2 serotyping: The KP MRSN Diversity Panel (NR-55604) was obtained through BEI Resources, NIAID,
NIH[33]. The 100 strains from the MRSN collection, KP_{WT} and $KP_{\Delta wcaJ}$ (positive and negative controls
respectively) were grown overnight in 1mL of LB in deep well 96-well plates (VWR). 100ul of each
overnight was transferred to a V-bottom plate and used for staining. Plates were centrifuged at 4000
rpm for 10 mins at 4°C, resuspended in 5% BSA/PBS and blocked for 1 h at RT shaking using a
microplate mixer. Bacterial samples were then centrifuged and stained overnight at 4°C shaking using
sera from CR_{KPK2}+pespO-infected mice diluted 1in100 (v/v) in 1% BSA/PBS. On the following day,
bacterial samples were washed two times in cold PBS and stained for 1 h at RT shaking with
AlexaFluor-488-conjugated anti-mouse IgG (1:300; Supplementary Table 3). Cells were washed twice
in cold PBS, fixed in 1% PFA/PBS and kept in the dark until flow cytometry analysis. Samples were
acquired on a Cytek Amnis CellStream (minimum of 50000 bacterial cells per sample) and data was
analysed using FlowJo v10.10.0.
788
789 **Bacterial cell immunofluorescence microscopy (adherent protocol and suspension protocol)**
790
791 We used two protocols for immunofluorescent staining of KP.
792
Adherent protocol (all ICC8001 staining)
mCherry expressing ICC8001 was grown as a saturated overnight culture which was sub-cultured
(1;1000 v/v) into M9 media (M9 salts M6030, Sigma, UK) supplemented with 0.4% glucose w/v

(G8270, Sigma, UK). 400ul was added into each well for imaging in a coverslip slide (μ-Slide 8 Well
high Glass Bottom, ibidi). Slides were grown statically at 37°C for 4 hours and cells were washed in
phosphate-buffered saline before staining. Primary staining (sera, antibodies) took place overnight at
4°C in 0.22um filtered 3% bovine serum albumin (A3059, Sigma, UK)/ PBS on a rocking platform.
Primary antibody was removed and slides were washed 3x in PBS followed by incubation with
fluorescently labelled secondary antibodies (Supplementary Table 3) in 3%BSA/PBS at room
temperature for 1 hour on a rocking platform. Slides were washed 3 times before imaging.
803
804   **Suspension protocol (MRSN collection, Figure 4D)**
805
MRSN collection strains were grown as saturated overnight cultures. 100ul of overnight was used for
staining. Following centrifugation (5000g, 5 minutes) cells were washed 3 times in PBS. Primary
staining (K2 antisera) took place overnight at 4°C in 3%BSA/PBS in 400ul volume with agitation. Cells
were washed by sequential spins (5000g, 5 minutes) and resuspension in PBS (3 washes). Secondary
antibody (Supplementary Table 3) was added for 45mins at room temperature in a 100ul volume
with agitation. Cells were washed by sequential spins (5000g, 5 minutes) and resuspension in PBS (3
washes). Cell membranes were then labelled with FM® 4-64 Dye (T13320, ThermoFisher Scinentific,
UK) at a final concentration of 5ug/ml in 100ul of PBS for 10 mins at room temperature in the dark.
Cells were washed once and resuspended in 100ul of fresh PBS before imaging. Cell suspension was
imaged on an 1% agarose (in dH_2O) pad according to the following protocol
(https://biotium.com/tech-tips/tech-tip-imaging-bacteria-using-agarose-pads/).
817
818   **CR_{T3F}+p*espO* glass binding assay**
819
mCherry expressing CR_{T3F}+p*espO* and CR_{WT} were grown as saturated overnight cultures and sub-
cultured (1:100) into DMEM low glucose (D5546, Merck, UK) into glass coverslip slides (μ-Slide 8 Well
high Glass Bottom, ibidi). Slides were grown statically at 37°C for 6 hours and cells were washed in
phosphate-buffered saline before fixing in 4% paraformaldehyde/PBS (R1026, Agar Scientific, UK) for
20 mins at room temperature. Cells were stained for MrkA according to the adherent protocol
outlined above.
826
827   **Pre-adsorption of CR sera**
828
Serum samples from mice infected with CR_{T3F}+pespO with anti-KP antibodies above 200ng/mL were
pooled; sera from CR_{WT}-infected mice were also pooled separately as a control. To inactivate
complement, pooled serum samples were heated at 55°C for 1 h and then were pre-adsorbed onto
CR_{WT} expressing sfGFP to remove CR-specific antibodies, thus enriching for anti-KP antibodies. This
was done by serially incubating the pooled sera samples, both from CR_{WT}- and CR_{T3F}+pespO-infected
mice, with ~5x10^8 CR_{WT} for 24 h at 4°C on a rotating wheel for 10 rounds of pre-adsorption. After
each 24h-incubation, the serum samples with CRWT were centrifuged at 3000 xg for 5 mins to pellet
the bacteria, which were kept at 4°C until flow cytometry analysis; the supernatant containing the
antibodies was subsequently incubated with a fresh batch of ~5x10^8 CR_{WT} expressing sfGFP. At the
end of the 8 rounds, sfGFP-expressing bacterial samples from each round of pre-adsorption with sera
from CR_{WT}- and CR_{T3F}+*pespO*-infected mice were stained with AF555-conjugated anti-mouse IgG
(1:400; Supplementary Table 3) in 2% BSA-PBS for 30 mins and analysed using a Cytek Amnis
CellStream. Obtained data was subsequently analysed using FlowJo v10.10.0.
842
Dot blot

Bacterial cultures were grown overnight in LB (CR) or M9 supplemented with glucose (KP),
normalised to an OD600 of 1 and centrifuged to pellet the cells. The pellet was resuspended in 1 ml
PBS and equal volumes of bacteria were then spotted onto a nitrocellulose membrane (#88018,
ThermoFisher Scientific) and allowed to air dry. The membrane was blocked at RT for 2 h in 5% milk-
PBS-T and incubated overnight at 4°C with the indicated sera diluted in 5% BSA-PBS-T to a
concentration of 2μg/ml of total IgG. The primary was washed three times in PBS-T and a secondary
HRP-conjugated anti-mouse antibody (Supplementary Table 3) was added in 5% milk-PBS-T,
incubated for one hour at RT, and washed three times. Detection was carried out using ECL substrate
(Bio-Rad) and images were acquired on a ChemiDoc XRS+ and visualised using Image Lab software
(Bio-Rad Laboratories).
854
855 **Fluorescent biofilm formation assay**
856
KP_{WT} and $KP_{\Delta mrkA-F}$ expressing sfGFP were grown overnight in M9 media and diluted into fresh M9
1:500 (v/v) and preincubated for 30 mins with no sera or with ~50 μg/mL IgG (1:100 dilution) of the
pre-adsorbed sera from either CR_{WT}- and CR_{T3F}+p*espO*-preinfected mice at RT on a rotating wheel.
100 μl of each sample in triplicate were transferred into wells in a black/clear-bottom 96-well tissue-
culture treated plate (#353219, SLS) and incubated statically at 37°C for 6 h. Samples were then
washed 5 times with sterile PBS and bottom optic fluorescence readings were taken at 485/520
(Ex/Em) using a FLUOstar Omega plate reader. Fluorescence values were normalised to those
obtained with the $KP_{\Delta mrkA-F}$ strain, which does not attach, to remove any non-specific fluorescence
and reduce batch-to-batch variation.

 Animal work
867
868 All animal work took place at Imperial College London (Association for Assessment and Accreditation
869 of Laboratory Animal Care accredited unit) under the auspices of the Animals (Scientific Procedures)
870 Act (UK) 1986 (PP7392693). Work was approved locally by the institutional ethics committee.
871
872 **Animals**
873
874 Female CD-1 and C57BL/6 mice were purchased from Charles River, UK. Mice were purchased at 29-
875 31 g weight (5-7 weeks) (CD-1 mice) and 18-20 g weight (6-8 weeks) (C57BL/6 mice). For each
876 experiment, mice were randomly assigned to experimental groups. Investigators were not blind to
877 the allocation. C57BL/6 were used for the data shown in Supplementary Figure 2C, all experiments
878 directly related to the citrOgen platform were performed in CD-1.
879
880 **Husbandry**
881
Animal technicians within the facility received mice and blindly separated them into groups of 5 into
high-efficiency particulate air (HEPA)–filtered cages with sterile bedding. Bedding was changed
weekly, and enrichment supplied. Mice were housed under a 12/12-hour light/dark cycle with
access to food and water *ad libitum*. Researchers (JSG and JLCW) conducting the work in this study
were not involved in selection of animals into groups as this was undertaken by technical staff prior
to starting experiments. Animals were identified by ear notching (conducted under restraint with no
anaesthesia).
889
890 **CR infection by oral gavage**
891
For infections, the indicated CR strains were grown overnight in 15 ml of LB (50 ml capped Falcon
tubes) at 37°C and 200 rpm. The next day, the tubes were centrifuged, and the pellet was
resuspended in 1.5 ml of sterile PBS. Mice were infected by oral gavage with 200 µl of this solution
($1x10^9$ CFU confirmed by viable counts as previously described[13]). CR faecal shedding was
determined by collecting stool samples at the specified days post-infection (DPI), homogenising the
samples in PBS and plating serial dilutions onto LB agar containing nalidixic acid. Plates were
incubated at 37°C overnight and colonies were counted on the following day.
899
900 Mice were culled at 6 (C57BL/6) or 8 dpi (CD-1) and colons were extracted for downstream
901 processing. When characterising antibody responses, mice were followed until clearance of CR
902 infection and then culled to obtain sera or challenged with KP infection.
903
904 **Colonic histology processing and immunostaining**
905
At 8 DPI, the colon was extracted, and the distal 0.5 cm was fixed in 4% PFA for 2 hours and then kept
in 70% ethanol until processing. Fixed tissues were processed, paraffin embedded and sectioned at 5
µm. For immunofluorescence, sections were dewaxed by submersion in HistoClear II solution
(#101412-884, VWR) twice for 10 min, 100% ethanol twice for 10 min, 95% ethanol twice for 10 min,
80% ethanol once for 3 min, and PBS-0.1% Tween 20-0.1% saponin (PBS-TS) twice for 3 min.
Subsequently, sections were heated for 30 min in demasking solution (0.3% trisodium citrate-0.05%
Tween 20 in distilled H2O). Once cooled, slides were first blocked in PBS-TS supplemented with 10%
normal donkey serum (NDS, D9663, Sigma) for a minimum of 1 h in a humid chamber, before being
incubated with anti-CR antibody (1:50) and anti-KP O1 (1:100) diluted in PBS-TS with 10% NDS
overnight at 4°C. Slides were rinsed twice for 10 min each time in PBS-TS, followed by incubation
with Alexafluor488-conjugated secondary antibodies (1:100) and DAPI (1:1000) (Supplementary

Table 3). Washing steps were repeated before slides were mounted with ProLong Gold antifade
mountant. Images were acquired using a Zeiss AxioVision Z1 microscope using a Hamamatsu
microscope camera and processed using ZEN3.1 software (ZEN lite, Carl Zeiss MicroImaging GmbH,
Germany).
921

922 **cIEC isolation and flow cytometry analysis**
923

Colonic intestinal epithelial cells (cIECs) from mock-infected and CR-infected C57BL/6 mice were
isolated from 2-cm distal colonic tissue samples as previously described[44]. Briefly, the 2-cm colonic
tissue sample was opened longitudinally and washed in 1× Hanks' balanced salt solution (HBSS)
without Mg and Ca (# 14185045, ThermoFisher Scientific) and incubated at 37°C, 200 rpm for
45 min in IEC dissociation buffer (1× HBSS, 10 mM HEPES, 1 mM EDTA, and 5 μL/mL 2-β-
mercaptoethanol). The remaining tissue was removed, and detached cells were collected by
centrifugation (3000 rpm for 10 min), followed by two DPBS (#D8537, Sigma) washes at 4°C. Cells
were then incubated 5 mins at RT with 50 μg/mL DNase I (#11284932001, Sigma) to digest
extracellular DNA and passed through a 70-μm cell strainer on ice to ensure a single-cell suspension,
pooling samples from the same group and rinsing with 20 mL of cold DPBS. All steps from here
onwards were performed on ice/at 4°C to maintain cell viability. cIECs were counted using trypan
blue to exclude dead cells and ~$1x10^7$ live cells were centrifuged, resuspended in fluorescence-
activated cell sorter (FACS) buffer (5% FBS, 2mM EDTA in DPBS) supplemented with Fc block (#130-
092-575, Miltenyi Biotec) and incubated for 10 min. Samples were then incubated with FACS buffer
containing anti-CR (1:1000) for 20 mins, washed twice in and then incubated with EpCAM-APC
(1:100) and phycoerythrin (PE) anti-rabbit IgG (Supplementary Table 3) for 40 min, washed, and
fixed in 1% paraformaldehyde in PBS prior to analysis on a BD LSRFortessa cell analyser (BD
Biosciences). Data were analysed using FlowJo v10.10.0 and the gating strategy is shown in
Supplementary Fig. 2A.
943

944 **Tail vein bleeds and serum separation**
945

946 Small volume bleeds (<100ul) were conducted on conscious, restrained mice using a Tail Vein
947 Injection platform (Hugeiron, MouseSerum, China). Blood was taken from the lateral tail vein by
948 puncture with a 25G needle and collected via pipette into a paediatric serum collection tube
949 (Microvette® 100 Serum Cat, Starstedt). Following 1 hour of clotting at RT, sera was separated by
950 centrifugation at 20 000g for 3 minute. Separated sera was stored at -80°C until use.
951

952 **KP pulmonary infections**
953

954 Anaesthesia (ketamine 80mg/kg and medetomidine 0.8mg/kg) was delivered by intraperitoneal
955 injection. Body temperature was maintained with a heat mat throughout until recovery or during the
956 intubation procedure itself.
957

958 Intubation was achieved with a 20G cannula (BD Insyte, BD) using the commercially supplied
959 intubation kit (Endotracheal Mouse Intubation Kit, ETI-MSE, Kent Scientific). 50ul of inoculum
960 (diluted in phosphate buffered saline) was delivered into the hub of the 20G cannula and drawn into
961 the lungs by spontaneous ventilation.
962

963 Recovery took place at 32°C in a forced air heating cabinet. At 15 minutes post anaesthesia induction
964 medetomidine was reversed by the administration of 0.8mg/kg atipamezole, delivered by
965 subcutaneous injection into the neck scruff.
966

The total volume of administered drugs was designed to deliver 20mg/kg of total fluids to animals, a human paediatric resuscitation dose.

Mice were returned to their home individual ventilated cages after they were spontaneously moving and checked at regular intervals post infection.

Cardiac puncture

Mice were weighed at end-point (after clearance for CR infected animals and 72 hpi for KP-challenged animals).

Anaesthesia (ketamine 100mg/kg and medetomidine 1mg/kg) was delivered by intraperitoneal injection. Mice were checked and following the absence of pedal reflexes placed in dorsal recumbency. Paws were taped to a board and scissors used to remove the skin over the xiphisternum. Blood was collected from the right ventricle (closed approach) with a 25G needle inserted at 30° to the skin surface until blood was freely aspirated into a 1ml syringe.

When blood was obtained for determination of IgG titres (i.e mice infected with indicated CR strains expressing heterologous antigens), the 25G needle was removed (to prevent haemolysis) and the blood was transferred to a serum collection tube containing a clot activator (BD Microtainer SST with separating gel, BD). Blood in the serum tube was allowed to clot for 1 hour at RT before centrifugation (20 000 xg) for 3 minutes. The sera was divided into aliquots and stored at -80°C until use.

When blood was obtained from KP challenged mice, 20ul of whole blood was taken for CFU enumeration by plating 10-fold serial dilutions (in PBS supplemented with 1mmol EDTA to prevent clotting) on LB Agar plates containing rifampicin 50ug/ml. The 25G needle was again removed and approximately 50-100 μl placed immediately into a tube coated with EDTA for anticoagulation (Microvette® 100 EDTAK3E, 100ul, Sarstedt) for haematological analysis. The remaining blood (approximately 800 μl) was allowed to clot for 1 hour at room temperature in a serum collection tube as described above, before centrifugation (20 000g) for 3 minutes. The sera was divided into aliquots and stored at -80°C until use.

Lung homogenate processing

Post-mortem lungs were removed and placed in 3ml of phosphate buffered saline in a gentleMACS C-tube (Miltenyi Biotec). These were homogenised on a gentleMACS Octo Dissociator (Miltenyi Biotec) using the program m_lung_2. The program was run twice in succession to achieve uniform homogenisation.

30ul of lung homogenate was taken for CFU enumeration by plating 10-fold serial dilutions on LB Agar plates containing rifampicin 50ug/ml.

1ml of lung homogenate was transferred into a 1.5ml microcentrifuge tube and spun at 10 000 xg for 5 minute at 4°C to pellet out particulate matter. The supernatant was stored -80°C for future use.

HM5 analysis of EDTA anticoagulated whole blood

Anticoagulated blood was run on the HM5 Vetscan analyser according to the manufacturer's instruction.

1017   Outcome variables of infection were assessed using the kits summarised in Supp Table 5 according to
1018   the manufacturer's guidelines with the following specifications.
1019
Capture ELISAs: ELISAs were performed using ELISA Clear Flat-Bottom Maxisorp Immuno 96- Well
Plates (#10547781, Fisher Scientific). Samples were added to coated and blocked plates and
incubated overnight at 4 degrees for all ELISAs. The range of sample dilutions used for each ELISA is
indicated in Supplementary Table 5; serial dilutions in the corresponding assay diluent were
performed when samples needed to be diluted further than 1in50. After addition of the TMB
substrate solution (#00-4201-56, ThermoFisher Scientific) to the wells at the end of the assay, these
were incubated at RT until a clear gradation in colour could be seen from the highest to the lowest
standard, at which point the reaction was stopped by adding 1M phosphoric acid (Sigma). All
absorbance readings were obtained using a using a FLUOstar Omega plate reader and readings at
540 nm were subtracted from absorbance readings at 450nm to correct for optical imperfections of
the plate. All ELISAs included blanks and standard curves with at least 7 points within range. Sample
values were obtained by interpolation into the standard curve; any samples with values out of the
range of the standard were assayed again at a dilution(s) that ensured the value was within range.
1033
Albumin: All reactions were done in a total volume of 200 µl in an untreated 96 well plate (Greiner
Bio-one) to allow vigorous shaking after reagent addition. Samples and standard were diluted 1:5 in
PBS (0.01 M, pH 7.4) and 5 µl were added per well; 5 µl of double-distilled water were added to the
blank. Subsequently 195 µl of stock working solution (prepared according to the manufacturer's
instructions) were added into each well, mixed thoroughly using a plate mixer and then left to stand
at RT for 10 mins before measuring the absorbance of each well at 628 nm using a FLUOstar Omega
plate reader.
1041
Creatinine: Serum samples did not require deproteinization. The assay was performed using the
colorimetric reaction mix, and readings were acquired by measuring absorbance at 570 nm using a
FLUOstar Omega plate reader. High levels of haemolysis lead to high background due to the colour;
background control samples (ie sample with no creatininase) were essential in those cases. 40 µl of
serum sample and 10 µl of assay buffer were used per sample well; standards were prepared
according to the manufacturer's instructions. After interpolating into the standard curve and
correcting for sample dilution, the values are given as nmol/ 50 µl sample; these were transformed
to mg/dL, where the molecular weight of creatinine is 113.12 mg/mmol.
1050
1051   **Data analysis and visualisation**
1052
Graphs and statistical analysis was carried out in GraphPad Prism version 10.4.1 for Windows and
version 10.4.0 for Mac (GraphPad Software, Boston, Massachusetts USA, www.graphpad.com). Data
were analysed for normal distribution based on D'Agostino-Pearson or Shapiro-Wilk normality tests
before performing any statistical analyses. Normally distributed data was analysed with the
parametric tests detailed in the figure legends; indicated non-parametric tests were applied to data
that failed to meet the normality criteria. When more than three comparisons were made, p values
were adjusted for multiple comparisons as indicated in the figure legends (Dunnett's post hoc test).
For ELISAs, two or three technical replicates were used to estimate experimental mean. Statistical
significance was only determined in datasets where each group had three or more biologically
independent repeats. A p value <0.05 is considered statistically significant. Each parameter was
measured only once in each distinct sample, but more than one parameter could be measured in the
same sample (data in Fig.3). value <0.05 is considered statistically significant.

Schematics in Supplementary Fig. 5C and Fig. 6 were made using some images from BioRender

(created with BioRender.com). Images were edited and compiled into figures in Affinity Designer V2.

(Serif Europe Ltd).

Data availability

All plasmid vector sequences are publicly available at
https://doi.org/10.6084/m9.figshare.c.7869659.v1. The data that support this study are available in
the supplementary Figures and Tables and from the corresponding authors upon request.

Acknowledgements

We would like to thank Professor Richard Strugnell and Dr Jonathan Wilksch at the University of
Melbourne for providing the reference K2 antisera and C13 monoclonal antibody for use in this work.
We would like to thank Professor Matthew Upton and Dr Katie Muddiman at the University of
Plymouth for the EC958 isolate.

We would like to thank our colleagues Dr Abigail Clements, Dr Andrew Conway Morris, Professor
Victor de Lorenzo, Professor Gordon Dougan and Professor Jose Penadés for their critical assessment
and feedback in writing this manuscript.

We would like to thank our colleagues Dr Mei Chong and Mr Edmond Yau in the Imperial College
London Enterprise team for their support in the patent application covering the technology
described in this manuscript.

Conflicts of interest

The authors declare no competing interests.

Patent

The technology underlying citrOgen has a patent pending.

Funding

This project has been supported by grants from The Royal Society (IC160080) and Medical Research
Council (MR/R02671)

References

1091

1.   El-Atawi, K. *et al.* Efficacy and Safety of Palivizumab as a Prophylaxis for Respiratory
Syncytial Virus (RSV) Disease: An Updated Systemic Review and Meta-Analysis. *Cureus*
15, e51375 (2023).
2.   Thandavaram, A. *et al.* The Efficacy of Bezlotoxumab in the Prevention of Recurrent
Clostridium difficile: A Systematic Review. *Cureus* **14**, e27979 (2022).
3.   Self, W. H. *et al.* Neutralizing COVID-19 Convalescent Plasma in Adults Hospitalized
With COVID-19: A Blinded, Randomized, Placebo-Controlled Trial. *Chest* **162**, 982–994
(2022).
4.   Focosi, D., Franchini, M., Casadevall, A. & Maggi, F. An update on the anti-spike
monoclonal antibody pipeline for SARS-CoV-2. *Clin Microbiol Infect* **30**, 999–1006
(2024).
5.   van der Horst, H. J. & Mutis, T. Enhancing <scp>Fc</scp> -mediated effector functions
of monoclonal antibodies: The example of <scp>HexaBodies</scp>. *Immunol Rev* **328**,
456–465 (2024).
6.   Stone, C. A., Spiller, B. W. & Smith, S. A. Engineering therapeutic monoclonal
antibodies. *Journal of Allergy and Clinical Immunology* **153**, 539–548 (2024).
7.   Henry, C. *et al.* Influenza Virus Vaccination Elicits Poorly Adapted B Cell Responses in
Elderly Individuals. *Cell Host Microbe* **25**, 357-366.e6 (2019).
8.   Tkaczyk, C. *et al.* Identification of anti-alpha toxin monoclonal antibodies that reduce
the severity of Staphylococcus aureus dermonecrosis and exhibit a correlation
between affinity and potency. *Clin Vaccine Immunol* **19**, 377–85 (2012).
9.   Zhu, H., Rollier, C. S. & Pollard, A. J. Recent advances in lipopolysaccharide-based
glycoconjugate vaccines. *Expert Rev Vaccines* **20**, 1515–1538 (2021).
10.   Mundy, R., MacDonald, T. T., Dougan, G., Frankel, G. & Wiles, S. Citrobacter rodentium
of mice and man. *Cell Microbiol* **7**, 1697–706 (2005).
11.   Sanchez-Garrido, J., Ruano-Gallego, D., Choudhary, J. S. & Frankel, G. The type III
secretion system effector network hypothesis. *Trends Microbiol* **30**, 524–533 (2022).
12.   Ruano-Gallego, D. *et al.* Type III secretion system effectors form robust and flexible
intracellular virulence networks. *Science* **371**, (2021).
13.   Crepin, V. F., Collins, J. W., Habibzay, M. & Frankel, G. Citrobacter rodentium mouse
model of bacterial infection. *Nat Protoc* **11**, 1851–76 (2016).
14.   Dennis, A. *et al.* The p50 subunit of NF-kappaB is critical for in vivo clearance of the
noninvasive enteric pathogen Citrobacter rodentium. *Infect Immun* **76**, 4978–88
(2008).
15.   DiGiandomenico, A., Rao, J. & Goldberg, J. B. Oral vaccination of BALB/c mice with
Salmonella enterica serovar Typhimurium expressing Pseudomonas aeruginosa O
antigen promotes increased survival in an acute fatal pneumonia model. *Infect Immun*
72, 7012–21 (2004).
16.   Bridge, D. R., Whitmire, J. M., Makobongo, M. O. & Merrell, D. S. Heterologous
Pseudomonas aeruginosa O-antigen delivery using a Salmonella enterica serovar
Typhimurium wecA mutant strain. *Int J Med Microbiol* **306**, 529–540 (2016).

17. Zhao, X. *et al.* Recombinant attenuated Salmonella Typhimurium with heterologous
expression of the Salmonella Choleraesuis O-polysaccharide: high immunogenicity
and protection. *Sci Rep* **7**, 7127 (2017).
18. Herrington, D. A. *et al.* Studies in volunteers to evaluate candidate Shigella vaccines:
further experience with a bivalent Salmonella typhi-Shigella sonnei vaccine and
protection conferred by previous Shigella sonnei disease. *Vaccine* **8**, 353–7 (1990).
19. World Health Organization. *WHO Bacterial Priority Pathogens List, 2024: Bacterial*
Pathogens of Public Health Importance to Guide Research, Development and
Strategies to Prevent and Control Antimicrobial Resistance. (2024).
20. Whitfield, C. & Trent, M. S. Biosynthesis and export of bacterial lipopolysaccharides.
Annu Rev Biochem **83**, 99–128 (2014).
21. Vinogradov, E. *et al.* Structures of lipopolysaccharides from Klebsiella pneumoniae.
Eluicidation of the structure of the linkage region between core and polysaccharide O
chain and identification of the residues at the non-reducing termini of the O chains. *J*
Biol Chem **277**, 25070–81 (2002).
22. Bronner, D., Clarke, B. R. & Whitfield, C. Identification of an ATP-binding cassette
transport system required for translocation of lipopolysaccharide O-antigen side-
chains across the cytoplasmic membrane of Klebsiella pneumoniae serotype O1. *Mol*
Microbiol **14**, 505–19 (1994).
23. Joensen, K. G., Tetzschner, A. M. M., Iguchi, A., Aarestrup, F. M. & Scheutz, F. Rapid
and Easy In Silico Serotyping of Escherichia coli Isolates by Use of Whole-Genome
Sequencing Data. *J Clin Microbiol* **53**, 2410–26 (2015).
24. Choi, K.-H. *et al.* A Tn7-based broad-range bacterial cloning and expression system.
Nat Methods **2**, 443–8 (2005).
25. Wong, J. L. C. *et al.* OmpK36-mediated Carbapenem resistance attenuates ST258
Klebsiella pneumoniae in vivo. *Nat Commun* **10**, 3957 (2019).
26. Clements, A. *et al.* Targeting subcapsular antigens for prevention of Klebsiella
pneumoniae infections. *Vaccine* **26**, 5649–53 (2008).
27. Kim, M. *et al.* Bacteria hijack integrin-linked kinase to stabilize focal adhesions and
block cell detachment. *Nature* **459**, 578–82 (2009).
28. Matute-Bello, G. *et al.* An official American Thoracic Society workshop report:
features and measurements of experimental acute lung injury in animals. *Am J Respir*
Cell Mol Biol **44**, 725–38 (2011).
29. Osuchowski, M. F. *et al.* Minimum quality threshold in pre-clinical sepsis studies
(MQTiPSS): an international expert consensus initiative for improvement of animal
modeling in sepsis. *Intensive Care Med Exp* **6**, 26 (2018).
30. Schembri, M. A. *et al.* Molecular Characterization of the Multidrug Resistant
Escherichia coli ST131 Clone. *Pathogens* **4**, 422–30 (2015).
31. Lam, M. M. C., Wick, R. R., Judd, L. M., Holt, K. E. & Wyres, K. L. Kaptive 2.0: updated
capsule and lipopolysaccharide locus typing for the Klebsiella pneumoniae species
complex. *Microb Genom* **8**, (2022).
32. Knutton, S. A novel EspA-associated surface organelle of enteropathogenic Escherichia
coli involved in protein translocation into epithelial cells. *EMBO J* **17**, 2166–2176
(1998).
33. Martin, M. J. *et al.* A panel of diverse Klebsiella pneumoniae clinical isolates for
research and development. *Microb Genom* **9**, (2023).

34. Old, D. C. & Adegbola, R. A. A new mannose-resistant haemagglutinin in Klebsiella. *J*
Appl Bacteriol **55**, 165–72 (1983).
35. Murphy, C. N. & Clegg, S. Klebsiella pneumoniae and type 3 fimbriae: nosocomial
infection, regulation and biofilm formation. *Future Microbiol* **7**, 991–1002 (2012).
36. Yang, J. *et al.* Transcriptional activation of the mrkA promoter of the Klebsiella
pneumoniae type 3 fimbrial operon by the c-di-GMP-dependent MrkH protein. *PLoS*
One **8**, e79038 (2013).
37. Ma, C. *et al.* Citrobacter rodentium infection causes both mitochondrial dysfunction
and intestinal epithelial barrier disruption in vivo: role of mitochondrial associated
protein (Map). *Cell Microbiol* **8**, 1669–86 (2006).
38. Leverton, L. Q. & Kaper, J. B. Temporal expression of enteropathogenic Escherichia coli
virulence genes in an in vitro model of infection. *Infect Immun* **73**, 1034–43 (2005).
39. Riechmann, L., Clark, M., Waldmann, H. & Winter, G. Reshaping human antibodies for
therapy. *Nature* **332**, 323–7 (1988).
40. Lee, E.-C. *et al.* Complete humanization of the mouse immunoglobulin loci enables
efficient therapeutic antibody discovery. *Nat Biotechnol* **32**, 356–63 (2014).
41. Feldman, M. F. *et al.* A promising bioconjugate vaccine against hypervirulent Klebsiella
pneumoniae. *Proc Natl Acad Sci U S A* **116**, 18655–18663 (2019).
42. Walker, K. A., Treat, L. P., Sepúlveda, V. E. & Miller, V. L. The Small Protein RmpD Drives
Hypermucoviscosity in Klebsiella pneumoniae. *mBio* **11**, (2020).
43. Mavridou, D. A. I., Gonzalez, D., Clements, A. & Foster, K. R. The pUltra plasmid series:
A robust and flexible tool for fluorescent labeling of Enterobacteria. *Plasmid* **87–88**,
65–71 (2016).
44. Hopkins, E. G. D., Roumeliotis, T. I., Mullineaux-Sanders, C., Choudhary, J. S. & Frankel,
G. Intestinal Epithelial Cells and the Microbiome Undergo Swift Reprogramming at the
Inception of Colonic Citrobacter rodentium Infection. *mBio* **10**, (2019).
1205

FIGURE 1.

A.

CR_WT

galF — rmlB — rmlD — rmlA — rmlC — wfgB — wzx — wfgC — wzy — wfgD — wfgE — gnd

CR O152 O-antigen biosynthesis gene cluster (*rfb* locus)

CR_KPO1

galF — wzm — wzt — wbbM — glf — wbbN — wbbO — kfoC — gnd

KP O2a O-antigen biosynthesis gene cluster (*rfb* locus)

glmS — wbbY — wbbZ

Glycosyltransferases converting O2a to O1v1 (O1)

CR origin
KP origin

1000bp

B.

KP O1 Ag — KP WbbY & WbbZ

CR O152 — CR O152 *rfb* locus

KP O2a Ag — KP O2a *rfb* locus

Core — *rfa* locus

Lipid A — Raetz pathway genes

CR_WT LPS

CR_KPO1 LPS

CR origin
KP origin

C.

250
190
134 — 100

α-CR O152   α-KP O1

D.

Day 8 stool shedding

CFU/GoF

**

CR_WT   CR_KPO1

E.

MERGE   CR   KP O1 Ag   DAPI

CR_WT

CR_KPO1

FIGURE 2.

A.

CR stool shedding

B.

C.

α-KP antibody titres
(Day 30 post CR infection)

D.

α-KP antibodies
bind KP O1 Ag

E.

FIGURE 3.

FIGURE 4.

A.

B.

	EC958	CR_WT	CR_ECO25b
α-O25b pAb			
Control (PBS)			

C. Day 8 stool shedding

D. α-EC958 antibody titres
(Day 30 post CR infection)

FIGURE 5.

A.

CR_WT

ROD21991 · wza · wzb · wzc · wcaA · wcaB · wcaC · wcaD · wcaE · wcaF · gmd · fcl · nudD · wcaI · manC · manB · wcaJ · wzxC · wcaK · wcaL · wcaM · galF

CR colanic acid biosynthesis locus

CR_KPK2

ROD21991 · wzi · wza · wzb · wzc · wcuF · wcuD · wclX · wzy · wzx · hypothetical · wbbJ · wcaJ · gnd · manC · manB · galF

KP K2 capsule biosynthesis locus

CR origin
KP origin
1000bp

B.

Day 8 stool shedding

CFU/GoF

ns

CR_WT   CR_KPK2 +p*espO*

C.

α-KP antibody titres (Day 30 post CR infection)

IgG (ng/ml)

ns   ****

○ CR_WT
● CR_KPK2 +p*espO*

KP_ΔwcaJ   KP_WT

D.

KP_WT anti-CR_WT IgG (i)
KP_WT anti-KPK2 ref (ii)
KP_WT anti-CR_KPK2 IgG (iii)

KP_WT anti-CR_WT IgG — zoom of (i)
KP_WT anti-KPK2 ref — zoom of (ii)
KP_WT anti-CR_KPK2 IgG — zoom of (iii)

E.

KL25	KL31	KL38	KL38	KL148	KL30	KL62	KL17	KL3	KL25
KL23	KL10	KL3	KL17	KL142	KL60	KL62	KL113	KL42	KL63
KL2	KL14	KL28	KL24	KL28	KL105	KL151	KL102	KL30	KL52
KL19	KL30	KL119	KL110	KL3	KL154	KL146	KL27	KL15	KL125
KL51	KL64	KL2	KL52	KL57	KL5	KL3	KL25	KL51	KL57
KL21	KL107	KL15	KL113	KL58	KL51	KL10	KL2	KL30	KL1
KL43	KL30	KL64	KL61	KL27	KL9	KL46	KL36	KL116	KL62
KL2	KL35	KL20	KL28	KL13	KL61	KL2	KL51	KL112	KL14
KL102	KL62	KL39	KL30	KL46	KL15	KL14	KL65	KL54	KL107
KL122	KL46	KL15	KL2	KL2	KL128	KL62	KL107	KL64	KL107

MFI signal key

2000	
1000	
500	
250	
125	
KP_WT	62.5
KP_ΔwcaJ	31.3

F.

KP anti-KPK2

KL30   KL119   KL110
KL64   KL2   KL52
KL107   KL15   KL113

FIGURE 6.

A.

CR_{T3F}

P_{map}

glmS | mrkA | mrkB | mrkC | mrkD | mrkF

KP type 3 fimbriae (T3F) structural gene operon

CR origin (CR origin)
KP origin (KP origin)

1000bp

B.

qRT-PCR *mrkA* expression

$2^{-\Delta Ct}$

*

CR_{T3F} +p*espO* LB
CR_{T3F} +p*espO* DMEM
CR_{WT} DMEM

C.

CR_{WT} CR_{T3F} MrkA

i   ii

zoom of (i)   zoom of (ii)

D.

Day 8 stool shedding

CFU/GoF

ns

CR_{WT} CR_{T3F} +p*espO*

E.

α-KP antibody titres
(Day 30 post CR infection)

IgG (ng/ml)

ns   ****

CR_{WT}
CR_{T3F} +p*espO*

$KP_{\Delta mrkA-F}$ KP_{WT}

F.

KP_{WT} anti-CR_{WT} IgG
KP_{WT} anti-KP MrkA ref
KP_{WT} anti-CR_{T3F} IgG

i   ii   iii

KP_{WT} anti-CR_{WT} IgG
KP_{WT} anti-KP MrkA ref
KP_{WT} anti-CR_{T3F} IgG

zoom of (i)   zoom of (ii)   zoom of (iii)

G.

Functional inhibition of biofilm
by CR_{T3F} sera

AU (normalised to $KP_{\Delta mrkA-F}$)

*

KP_{WT} no Ab
KP_{WT} + CR_{WT} sera
KP_{WT} + CR_{T3F} sera

H.

KP_{WT} + CR_{WT} sera KP_{WT} + CR_{T3F} sera

FIGURE 7.

A. citrOgen strain generation ~7 days

choice of heterolgous Ag gene editing vectors

galF — LPS O-Ag — gnd

ROD21991 — Capsule-Ag — galF

P_{map}
glmS — Complex protein Ag

Single Ag or multiple Ags selected

✓ Validated CR insertion/substitution sites
✓ *in vitro* evaluation tools for QC

B. *In vivo* Ab production ~28 days

CR stool shedding

EspO boost of mucosal adherence

Continuous heterologous Ag presentation over the infection cycle

Ab-mediated CR clearance

dpi

Blood collection

Antibody testing ~3 days

✓ Heterologous antigen Abs

C.

pAb applications (sera)

Functional assays e.g. serum killing biofilm inhibition

Serotyping

mAb development (B-cells)

Recombinant mAb

Hybridoma mAb

Evaluation of protection (challenge studies)

citrOgen Platform

Downstream applications