**Dissecting the effects of single amino acid substitutions in SARS-CoV2 Mpro**

Shwetha Sreenivasan, Joseph D. Fontes, and Liskin Swint-Kruse*

Department of Biochemistry and Molecular Biology, University of Kansas Medical Center, Kansas City, Kansas, 66160, United States

* To whom correspondence should be addressed. Tel: 1 913 588 0399; Fax: +1 913 588 9896; Email: lswint-kruse@kumc.edu

Running title: Dissecting effects of single substitution in Mpro

Number of pages:

- Manuscript: 28 pages
- Supplementary material: 19 pages
- Tables: 1 page
- Figures: 5 figures

Description of supplementary material including filenames:

- Mpro_6positions_fnVSabundance_supplementary_14Mar25.docx (19 pages)

**ABSTRACT**

Successfully predicting the effects of amino acid substitutions on protein function and stability remains challenging. Recent efforts to improve computational models have included training and validation on high-throughput experimental datasets, such as those generated by deep mutational scanning (DMS) approaches. However, DMS signals typically conflate a substitution's effects on protein function with those on *in vivo* protein abundance; this limits the resolution of mechanistic insights that can be gleaned from DMS data. Distinguishing functional changes from abundance-related effects is particularly important for substitutions that exhibit intermediate outcomes (*e.g.,* partial loss-of-function), which are difficult to predict. Here, we explored changes in *in vivo* abundance for substitutions at representative positions in the SARS-CoV-2 Main Protease (Mpro).  For this study, we used previously-published DMS results to identify "rheostat" positions, which are defined by having substitutions that sample a broad range of intermediate outcomes. We generated 10 substitutions at each of six positions and separately measured effects on function and abundance. Results revealed an ~45-fold range of change for abundance, demonstrating that it can make significant contributions to DMS outcomes. Moreover, the six tested positions showed diverse substitution sensitivies for function and abundance. Some positions influenced only one parameter; others exhibited rheostatic effects on both parameters, which to our knowledge, provides the first example of such behaviour. Since effects on function and abundance may arise through different biophysical bases, these results underscore the need for datasets that independently measure these parameters in order to build predictors with enhanced mechanistic insights.

**ABSTRACT for broader audience**

Changing one amino acid in a protein can affect its function, its abundance, or both. Understanding these separate effects will help scientists predict how protein changes alter biology, which is important for understanding pathogen evolution, improving personalized medicine, and bioengineering. This work reports a study to experimentally separate these effects for the Main protease of SARS-CoV-2 and suggests strategies for building better prediction models for emerging variants of this key viral protein.

**INTRODUCTION**

Accurately predicting the outcomes of single amino acid substitutions is essential for advancing protein evolution, personalized medicine, and bioengineering.[1; 2] However, despite extensive efforts over three decades, advances in computational predictions have generally been incremental.[3-6] This indicates that the fundamental biochemical and biophysical mechanisms underlying the protein sequence-structure-function relationship – which form the bases of algorithmic assumptions – are not yet fully understood.

Understanding these mechanisms is particularly challenging for substitution outcomes that are not binary. Instead of leading to complete loss or retention of wild-type (WT) function, many substitutions result in intermediate or enhanced outcomes. Such outcomes are prevalent at "rheostat" positions, where the set of possible substitutions results in a graded functional change (*e.g.,* **Supplemental figure 1** and **Supplemental table 1**).[7-9] Indeed, predictions for substitution outcomes at rheostat positions are worse than predictions for other types of positions.[9]

To address these challenges, the field has recognized that training and validating computational models requires datasets that are not biased by selections for specific substitutions or positions, to capture the full spectrum of possible outcomes. However, experimentally testing all 19 substitutions at every position in a protein using conventional biochemical assays is challenging. To circumvent this, high-throughput techniques like deep mutational scanning (DMS)[10-12] have emerged as tools that enable the systematic assessment of substitution effects on proteins.

In DMS, the first step is to create a large library of protein substitution variants via site saturating mutagenesis[13] across the full-length protein or a region/domain of interest.[14;

[15] In a commonly-used DMS approach, the substitution libraries are then introduced into a biological system such as yeast,[16] bacteria,[17] or mammalian cells[18]. The cells are then placed under selective conditions, such as drug exposure, and screened for a desired cellular phenotype or reporter outcome.[19; 20] Finally, high-throughput sequencing techniques[21; 22] are used to identify substitutions that impacted each cell's fitness (and thereby infer the effects on the protein) (*e.g.,* [23]) or *in vivo* activity of the reporter (*e.g.,* [24; 25]). Regardless of experimental design, measured outputs are often referred to as "functional scores".[26]

Several factors complicate interpreting the scores from high-throughput DMS assays.[8; 27] First, different choices of selection pressures can change the fitness/growth measured for substitution variants[28], and none of these pressures may actually represent physiological conditions.[27] Second, many studies assume that fitness or growth levels directly correlate with protein function.[29] For example, variants with worse fitness are interpreted as having worse function, whereas those with enhanced function (beyond that of the wild-type variant) exhibit improved function.[10] However, a fitness or growth advantage might not be directly proportional to a protein's functional changes.[26; 30] Finally, in many DMS studies, the measured outcomes reflect the net effect of each substitution. That is, changes in the observed "functional scores" could arise from affecting any of several protein parameters[31], including solubility,[32] stability,[33; 34] binding affinity,[35-37] allosteric regulation,[38-40] ligand specificity[41; 42], folding rate[43], translation rate,[44] and/or enzyme activity[38; 45; 46]. Thus, the outcomes monitored by DMS cannot be assigned to individual aspects of a protein's function.

The use of "combined" signals presents a new challenge for training and validating computational predictors: Individual protein parameters can be altered through distinct

biophysical processes, and each parameter may necessitate specialized predictive models that must then be integrated to generate an overall prediction model. Thus, constructing such models requires experimental datasets that, at the very least, capture changes in both function *and in vivo* abundance. To date, only a limited number of DMS studies report both parameters.[18; 47-49] Instead, many studies explicitly assume that single amino acid substitutions do not alter *in vivo* abundance, equating overall growth or fitness changes solely to functional changes.

The use of "combined effects" for building computational predictors may be especially problematic for substitutions with intermediate outcomes. To explore this, we independently measured the effects of single amino acid substitutions on function and *in vivo* abundance in the SARS-CoV-2 main protease (Mpro). Our studies included several positions that showed a high frequency of intermediate outcomes in a published DMS study.[50] Strikingly, despite sampling only 60 of the 5814 possible Mpro substitutions, this set captured a ~45-fold range of change in *in vivo* abundance. Moreover, among the six positions tested, we observed diverse combinations of substitution sensitivities for both function and abundance, highlighting the complexity of substitution effects that can arise from single substitutions. To our knowledge, this work provides the first examples of rheostat positions that control both protein function and *in vivo* abundance. As such, these results highlight another complex outcome – the simultaneous tuning of multiple protein parameters – that can arise from substitutions at rheostat positions.

**RESULTS**

**In prior DMS studies, Mpro positions exhibited diverse substitution sensitivities**

Mpro, also known as the 3C-like protease or Nsp5,[51] is a highly conserved cysteine protease essential for the replication and transcription of coronaviruses.[52] Structurally, Mpro exists as a homodimer, and dimerization is essential for its proteolytic activity.[52] Given Mpro's role in SARS-CoV-2 replication and the number of changes observed over the course of the pandemic (with single amino acid substitutions observed for at least 287 of 306 positions)[53; 54], it is useful to understand how single amino acid substitutions alter its function and *in vivo* abundance.

In a previously published DMS study of Mpro, Flynn *et al.*,[50] created 5782 of the 5814 possible single amino acid substitutions.  Their goal was to identify positions where most substitutions abolished its activity, with the rationale that they might be good drug targets. In this context, abundance changes were perhaps unimportant and did not affect the interpretations made by the authors. However, by their nature, DMS outcomes cannot yield information on the molecular mechanisms by which substitutions alter Mpro activity.

In the work of Flynn *et al.*, each substitution's phenotype was assessed with three assays; results were assessed using deep sequencing to measure the enrichment or depletion of each substitution in the library, which was then interpreted as the substitutions' effects on Mpro activity.  Results were generally in good agreement with each other; approximately 60% of substitutions exhibited Mpro activities equivalent to wild-type (WT), and ~25% of substitutions showed complete inactivation **(Supplementary figure 2)**. No substitutions with enhanced activity (greater than WT) were detected. The remaining substitutions sampled the full range of possible intermediate values (between the WT and

inactive Mpro; **Figure 1** and **Supplementary figure 2).** The greatest number of positions with intermediate activities were detected using the growth assay, so we used this data set as a reference for our current work.

Here, we further tested whether single Mpro substitutions can separately affect function (proteolytic activity) and *in vivo* abundance. To identify candidate positions for our work, we used the DMS growth assay to classify each Mpro position based on its overall substitution sensitivity (**Figure 1, Supplementary table 2**). Examples for the five classes of substitution sensitivities are shown in **Supplementary figure 3**. This classification scheme is allows us to identify the set of rheostat positions that is not readily apparent from the average values of a position substitution outcomes.[8]

Of the 306 Mpro positions, 125 positions were rheostat positions (**Figure 1c-d** and **Supplementary figure 3b**). Of these rheostat positions, we further selected four positions (48, 52, 77, and 234), for which previously-published computations of dynamic flexibilities predicted different biophysical characteristics[55] **(Supplementary figure 4)**. Finally, we chose two DMS-neutral positions (110 and 50), also with different flexibilities, to serve as controls for positions with fewer iintermediate outcomes. At these six positions, we made ten random amino acid substitutions, which we previously showed was sufficient to assign a position's substitution sensitivity class.[7]

**Mammalian cell-based assays of MPro function**

To dissect the effects of the 60 single amino acid substitutions on Mpro function and *in vivo* abundance, we designed HEK293 cell-based assays **(Supplementary figure 5)**. A

plasmid expressing a cyclic-luciferase reporter enzyme, engineered to contain an Mpro cleavage site, was stably integrated into HEK293 cells to enable constitutive reporter expression. The cyclic-luciferase enzyme expressed from this gene was inactive unless cleaved by Mpro. Cells were then transiently transfected with Mpro substitution variants and the luciferase assay was performed with cell lysate (**Supplementary figure 6**, first column). Since the values measured in the lysate could be sensitive to changes in either Mpro's proteolytic activity or its cellular abundance, these luciferase assay outputs are equivalent to DMS growth outcomes. In our assays, the "combined effects" for the set of 60 substitutions sampled >90% of the accessible range of luciferase outputs **(Figure 2a),** as defined by the inactive C145A substitution and the maximum observed outcome (V77S, **Figure 3a**).

When the data from the previous[50] and current studies were compared, the values for the 60 substitution variants showed a Spearman correlation of r = 0.51 **(Supplementary figure 7)**. One key difference is that our assays identified substitutions with activity levels higher than WT, which were not detected in the DMS study**.** Excluding these enhanced outcomes substantially improved the comparison (Spearman r = 0.64). Such enhanced outcomes have previously been observed at rheostat positions in other proteins.[1; 7; 56] In addition, we compared each *position*'s overall substitution sensitivity between the two studies. Our substitution sensitivities matched those observed in the DMS growth assay for both neutral positions (50 and 110) and all four rheostat positions (48, 52, 77 and 234) (**Table 1; Supplementary figure 6**, first column).

**The relationship between effects on abundance and function is complex**

To determine the contributions of altered *in vivo* protein abundance to changes in luciferase outputs, we independently measured Mpro abundance using an indirect sandwich enzyme-linked immunosorbent assay (ELISA) **(Supplementary figure 5)**. We note that the ELISA cannot differentiate between functional homodimers and non-functional monomers. Furthermore, changes in *in vivo* protein abundance can arise from effects on translation rates, folding efficiency, stability, solubility, and degradation pathways. We then used the abundance values to normalize the luciferase output values and thereby isolate substitution effects on Mpro's protease activity. Results from the abundance normalized-function might reflect changes in enzyme catalytic rates, substrate binding affinity, ligand specificity, and/or dimerization. These three outcomes are shown in **Figure 2** for the set of 60 substitutions, and example plots for position 77 are shown in **Figure 3**. Data for all other positions are shown in **Supplementary figure 6**.

Even though the un-normalized luciferase output suggested that several substitutions had very low biological activity **(Figure 2a),** no substitutions showed extremely low *in vivo* abundance **(Figure 2b)** or was catalytically inactive **(Figure 2c)**. The set of 60 substitutions showed an ~45-fold range of changes in abundance **(Figure 2b)** and includes substitutions that are both lower and higher than the WT level. The abundance normalized-function sampled only 60% of the accessible functional range and also included substitutions with enhanced function **(Figure 2c).**

Notably, for any of the six positions tested, the rank-order of substitution outcomes for the un-normalized luciferase output does *not* match the order for abundance or normalized-function **(Figure 3; Supplementary figure 6)**. For example, at position 77,

substitutions D, G, and M showed a significant decrease in abundance, whereas the normalized-functions of these substitutions were within the range of WT values **(Figure 3)**. Conversely, at position 234, substitutions V and K exhibited significantly lower functional outcomes compared to the WT, although their abundances were near WT levels **(Supplementary figure 6p-r)**. Indeed, when changes in abundance were compared to changes in the un-normalized luciferase outputs for the full dataset, no correlation was observed **(Figure 4a)**. Thus, for Mpro, changes in protein abundance can significantly modulate the observed biological outcomes, and in some cases, may even overshadow changes in catalytic activity of the substitution itself.

Interestingly, a comparison of the changes in abundance and normalized-function revealed a moderate correlation for the set of 60 substitutions (Spearman r = -0.68, **Figure 4b**). When each of the six positions was examined individually, a strong relationship was observed for positions 48, 50 and 52 (Spearman r = -0.9; **Supplementary figure 8**). However, this trend was not observed for Mpro position 77, which had a Spearman correlation coefficient of -0.5 **(Supplementary figure 8, Figure 5**, magenta circles). At this position, individual substitutions showed a variety of outcomes: V77H increased abundance and decreased function, V77M decreased abundance and increased function, V77D decreased both parameters, and other substitutions affected only one of the two parameters.

**Substitution sensitivities of individual Mpro positions reveal distinct combinations of neutral and rheostat behaviors for function and abundance**

Finally, we determined the overall substitution sensitivity classes (**Supplementary table 1**) for normalized-function and abundance for the six Mpro positions **(Table 1; Figure**

**5**). Despite examining just six positions, the data reveals a wide range of behaviors. Positions 48 and 77 showed strong rheostat effects for both parameters, whereas position 52 displayed moderate rheostat behavior for both, and position 110 displayed neutral behavior for both. Position 234 was neutral for abundance but rheostat for function, and position 50 was a moderate rheostat for abundance and neutral for function.

The different sensitivities of these six positions are not readily explained by their locations on the Mpro structure **(Figure 5)**. Positions with different types of outcomes are adjacent to each other, and none are near the active site or the dimerization interface. We also considered solvent exposure, secondary structure, and proximity to the catalytic dyad. However, none of these clearly accounted for the observed differences in substitution sensitivities **(Supplementary table 2).**  This matches our studies of rheostat positions in other proteins, which also are distributed widely across protein structures and lack obvious relationships to structural features.[1; 7; 40; 57]

**DISCUSSION**

In this set of 60 Mpro substitutions, some altered function, some altered abundance, and some altered both parameters – to different magnitudes, and in same or different directions. The fact that multiple combinations of substitution sensitivities were observed for just six positions suggests that such variability is likely widespread across Mpro positions.

Our result is in striking contrast to the assumption underlying many DMS studies, that single amino acid substitutions do not greatly alter abundance,[11; 58; 59] but it is in agreement with other studies that demonstrate single substitutions alter abundance in

other proteins (e.g., [18; 45; 47; 60; 61]). For some applications, the distinction between changes in function and abundance might not matter. As noted above, Flynn *et al.*, strove to identify Mpro positions that did not tolerate substitutions. In another example, the DMS study of p53 was used to identify missense mutations that could lead to cancerous phenotypes.[62] In both of these studies, the "combined effects" from DMS are useful to understand biological outcomes and the mechanism by which the protein was inactivated was irrelevant.

However, to understand the general principles by which proteins *work*, which is critical for developing accurate prediction algorithms, the DMS assay design is not sufficient. Indeed, although the substitution classes for each Mpro position (rheostat or neutral, **Table 1**) for combined effects from DMS/luciferase outputs appear to agree with those for abundance-normalized function, the rank order of substitution outcomes differ. This highlights the fact that different *substitutions* can make diverse contributions to individual protein parameters.

Another complex relationship illustrated by the current results is that protein *positions* can play distinct roles in regulating function and abundance. The examples here for Mpro positions 48 and 77 are, to our knowledge, the first documentation some protein positions can have rheostatic effects on both parameters. This parallels the observations in other proteins, which have rheostat positions that alter multiple *functional* parameters. For example, rheostat positions in human liver pyruvate kinase affected up to five different functional parameters (binding affinity to its substrate, two different allosteric effectors, and two allosteric coupling constants).[63] In a second example, a subset of rheostat positions in *Escherichia coli* LacI affected both transcription repression and response to allosteric ligand.[64]

Studies of single substitutions in other proteins also show complex relationships between function and abundance, with the two parameters altered to different extents and in different directions.[65; 66] In the current study, the sets of substitutions for three positions show a clear trade-off, with increased function offset by decreased abundance. It is interesting to think about the possible mechanisms underlying this trade-off. Abundance changes might derive from changes in protein stability, which has been observed in many studies[47; 67; 68] and is often expected in protein engineering[69-72]. Alternatively, the function/abundance trade-off could arise from a cellular response, with the HEK293 cells limiting the overall level of Mpro activity.

However, we reason that a cell-based response should equally affect all substitutions at all positions. In contrast, the stability/function trade-off is not universal.[34; 45; 73; 74] Thus, the *lack* of correlation for Mpro position 77 supports the hypothesis that positions 48, 50, and 52 experience a stability/function trade-off. These results lead us to further hypothesize that some positions may act as rheostats for both function and stability. Prior to this work, stability rheostat positions were known for (i) *Zymomonas mobilis* pyruvate kinase, but they had few detectable effects on function[75; 76] and (ii) GB1[1; 77], for which effects on function are not known.

In conclusion, this study illustrates that single substitutions can separately affect different protein parameters. Each parameter can have different biophysical bases, which likely requires a unique model to predict. Ultimately, integrating parameter-specific contributions is critical to build biophysical models that accurately capture actual protein behavior.

**MATERIALS AND METHODS**

**Identification of rheostat positions in the Mpro DMS dataset**

Raw data from the previously published deep mutational scanning (DMS) dataset for the main protease (Mpro) of SARS-CoV2[50] (from supplementary file "elife-77433-fig2-data1-v2") was analysed using the RheoScale calculator[78]. The rheostat, toggle, and neutral scores thus obtained were used to assign the position types as discussed in **Supplementary table 1**. Such class assignment for all Mpro positions from the DMS growth assay is available in **Supplementary figure 1c**. For these analyses, the range was defined by substitution outcomes of P9I and S1T, and the bin number was set to 10.

**DNA constructs**

The plasmid pCS-NSP5, which encodes wild-type Mpro (Uniparc ID: UPI00137481FB) was used to generate Mpro variants. For each targeted amino acid position, 10 randomly-chosen, single amino acid substitutions were produced by Twist Biosciences (San Francisco, CA). Plasmids were transformed into DH5-alpha competent cells (Catalog # C2987H, New England Biolabs Ipswich, MA), and purified using QIAprep Spin Mini Kit (Catalog #27104, Hilden, Germany).

The cyclic-luciferase reporter plasmid pCycLucNSP5, which was used to assay Mpro activity, was created by modifying the plasmid cycLuc_TEVS (a gift from Roman Jerala, Addgene plasmid #119207).[79] This included replacing the cleavage site for the tobacco etch virus protease with the cleavage site for SARS-CoV2 Mpro (SAVLQ↓SGFRK)[80] and removing the PEST sequence on the C-terminus of the cLUC. This Mpro expression plasmid is currently being deposited with Addgene.

**Cell culture**

HEK293 cells (American Type Culture Collection, Manassas, VA; catalog number CRL-1573) were cultured in Dulbecco's Modified Eagle Medium with 4.5 g/L glucose, supplemented with 10% fetal bovine serum, 100 µg/mL penicillin, and 100 U/mL streptomycin. Stable cell lines expressing the reporter were maintained in 300 µg/ul G418 (CAS # 108321-42-2; Goldbio, St. Louis, MO). Cells were maintained at 37 °C under a humid atmosphere containing 5% $CO_2$. Growth was monitored daily, and cells were split at 80-90% confluency. The growth media was changed as required. Cells were used for a maximum of 20 passages. Subcultures for transfection were prepared in 96-well plates with $2.5 \times 10^4$ cells/well and 200 µL complete medium.

**Stable cell line generation**

The reporter plasmid pCycLucNSP5 was transfected into HEK293 cells using Turbofect transfection reagent (Thermo Fisher Scientific, Waltham, MA). Selection for stable expression of the plasmid was performed with 600 µg/ml G418. Multiple clones (HEK293-CLucNSP5) were isolated and utilized in experiments.

**Transfection of Mpro substitutions to assay protease activity**

HEK293-CLucNSP5 cells were transfected in 96-well plates utilizing Turbofect transfection reagent (Thermo Fisher Scientific, Waltham, MA). Each transfection contained 15 ng of pCS-NSP5 expressing WT or variant Mpro, 10 ng pTK-Renilla (when encoded *Renilla* luciferase that was used as a transfection efficiency control), and 175 ng pCMV-LacZ (as stuffer plasmid). Culture media was changed four hours post transfection. Cells were incubated overnight before performing the luciferase assay or ELISA.

**Luciferase assay**

Luciferase was measured using the Bright-Glo™ Luciferase Assay System kit (Catalog # E2620) per manufacturer's instructions except as noted below, and results were assayed using a Tecan plate reader (Mannedorf, Zurich, Switzerland). A modification of the reagent volumes for the luciferase assay was employed, using 50 µl of LAR II reagent per well for firefly luciferase activity. After measuring firefly luciferase signal, 50 µl of 1X Stop & Glo reagent was added per well to quench the firefly luciferase signal and measure *Renilla* luciferase activity. Raw data for all Mpro substitutions tested in this study have been deposited in

https://github.com/ShwethaSreenivasan/Mpro_abundance_vs_function/blob/main/Mpro_variant_data_22Dec25.xlsx .

**ELISA to measure Mpro abundance changes**

HEK293-CLucNSP5 cells transfected with Mpro were lysed with 25 µl of 1 X ELISA Cell Extraction Buffer (Catalog #69905, Cell Signaling Technology, Danvers, MA) per well in a 96-well plate. Indirect sandwich ELISA was performed using a kit from Elabscience® (Catalog # E-ELIR-K001, Houston, TX) in clear 96-well ELISA plates (Catalog # 15031, Thermo Fisher Scientific, Waltham, MA). The plates were coated with SARS-CoV-2 3C-Like Protease capture antibody at 1 µg/mL in coating buffer and incubated at 4 °C overnight. Wells were washed and blocked, and cell lysate (diluted 1:3 in the blocking buffer) was added. For detection, 2 ng/mL Strep II tag antibody (Catalog # A01737-100, GenScript, Piscataway, NJ) was added to wells and incubated at room temperature for two hours with agitation. Following washing, goat anti-mouse IgG (H + L)-HRP Conjugate (Catalog #1706516, Bio-Rad, Hercules, CA) was

added and incubated for an hour at room temperature, followed by addition of 100 μL of

TMB substrate to each well for detection. Following incubation for 12-15 minutes, stop

solution was added and absorbance at 450 nm measured within 30 minutes using a Tecan

Plate Reader (Mannedorf, Zurich, Switzerland). Raw data have been deposited in

https://github.com/ShwethaSreenivasan/Mpro_abundance_vs_function/blob/main/Mpro_

variant_data_22Dec25.xlsx.

**Computing abundance normalized-function**

Three experimental measurements were made for each Mpro WT and variants

(i) Mpro's *in vivo* activity was measured using the firefly luciferase activity. (ii) Transfection

differences were controlled by measuring the *Renilla* luciferase signal intensity from the

control plasmid (pTK-Renilla). (iii) Mpro abundance was determined from ELISA. For each

variant, 4-6 technical replicates for two biological replicates were assayed. These three

measurements were then used to generate three different outcomes for each Mpro variant.

*Mpro outcome 1: Combined effects*.  For each Mpro variant, the "combined effect"

was obtained by normalizing the firefly luciferase activity to the *Renilla* luciferase activity.

This was then normalized to each plate's WT average with appropriate error propagation

(Equation 1 below), to control day-to-day technical variation.

*Mpro outcome 2: Abundance changes*.  The abundance changes for each Mpro

variant were obtained by normalizing results from the indirect ELISA assay to each plate's

WT average, with appropriate error propagation (Equation 1 below), to control for day-to-

day technical variation.

*Mpro outcome 3: Abundance normalized-function*.  Abundance normalized-function for each variant was obtained by calculating the ratio of the normalized combined effects to normalized abundance changes, with appropriate error propagation (Equation 1).

For all error propagation during normalizations, we used the following formula:

$$\frac{\sigma_R}{R} = \sqrt{\left(\frac{\sigma_A}{A}\right)^2 + \left(\frac{\sigma_B}{B}\right)^2 + 2\sigma_R^2\left(\frac{\sigma_{AB}}{AB}\right)} \qquad \text{(Equation 1)}$$

where, $R$ is the ratio of outcomes $A$ and $B$, and $\sigma_R$, $\sigma_A$ and $\sigma_B$ are their respective standard deviations; $\sigma_{AB}$ is determined by ratio of the standard error of means of $A$ and $B$ as reported by GraphPad Prism version 9.4.1.

The independent measurements for these three outcomes are reported in

https://github.com/ShwethaSreenivasan/Mpro_abundance_vs_function/blob/main/Mpro_variant_data_22Dec25.xlsx. All data plots were made using GraphPad Prism version 9.4.1.

**Statistical analyses of experimental data**

To identify which variants differed from WT, statistical analyses were carried out using GraphPad Prism version 9.4.1.  The data acquired followed a log-normal distribution, as indicated by D'Agostino-Pearson tests. Thus, all values were transformed to log10 prior to statistical tests.  One-way ANOVA was performed with Dunnett's correction for multiple comparison of each substitution at a position to the value collected for WT of that position's data set; statistically significant variants were identified based on Dunnett's correction (****p <0.0001, ***p<0.001, **p<0.01, *p<0.1); complete reports of these statistical analyses are included in **Supplementary table 3**.

**Assigning overall substitution outcomes to each tested Mpro position**

The measurements for combined effects, abundance-normalized function, and abundance changes were assessed using the RheoScale calculator to obtain the rheostat, toggle, and neutral scores for each of these parameters at the seven positions tested in this study **(Table 1)**. For these analyses, the histogram ranges were determined by the minimum and maximum values for each of the three outcomes measured in this study, and the bin number was set to 10. The scores used to assign the position types are described in **Supplementary table 1**.

**DATA AVAILABILITY.** The number of replicates, averages and standard deviations for the values shown in **Figure 3** and **Supplementary figure 6** are available on https://github.com/ShwethaSreenivasan/Mpro_abundance_vs_function/blob/main/Mpro_variant_data_22Dec25.xlsx

**SUPPLEMENTARY DATA.** Supplementary Data are available at Protein Science. Additional citations associated with these data include[69-72; 81; 82].

**FUNDING.** This work was supported by funding from the National Institute of Health GM147635 to LSK and JDF.

**CONFLICT OF INTEREST**. The authors have no conflict of interest to declare.

**ACKNOWLEDGEMENTS.** We thank Halya Fedosyuk (KUMC) for assistance with initial experiments and Dr. Aron Fenton, Dr. Pierce O'Neil, Anastasiia Sivchenko and Carter Grey (KUMC) for discussion on the manuscript. We thank Jordan Baker (KUMC, Department of Biostatistics and Data Science) for providing suggestions for data analyses. Finally, we thank Dr. Paul Campitelli and Dr. Banu Ozkan (Arizona State University) for sharing and discussing their published DFI scores computed for Mpro.

**REFERENCES**

1. Swint-Kruse L, Fenton AW (2024) Rheostats, toggles, and neutrals, Oh my! A new framework for understanding how amino acid changes modulate protein function. J Biol Chem 300:105736.
2. Swint-Kruse L (2016) Using Evolution to Guide Protein Engineering: The Devil IS in the Details. Biophysical Journal 111:10-18.
3. Tsishyn M, Hermans P, Pucci F, Rooman M (2025) Residue conservation and solvent accessibility are (almost) all you need for predicting mutational effects in proteins. bioRxiv:2025.2002.2003.636212.
4. Munro D, Singh M (2020) DeMaSk: a deep mutational scanning substitution matrix and its use for variant impact prediction. Bioinformatics 36:5322-5329.
5. Nagar N, Tubiana J, Loewenthal G, Wolfson HJ, Ben Tal N, Pupko T (2023) EvoRator2: Predicting Site-specific Amino Acid Substitutions Based on Protein Structural Information Using Deep Learning. Journal of Molecular Biology 435:168155.
6. Cheng J, Novati G, Pan J, Bycroft C, Žemgulytė A, Applebaum T, Pritzel A, Wong LH, Zielinski M, Sargeant T, Schneider RG, Senior AW, Jumper J, Hassabis D, Kohli P, Avsec Ž (2023) Accurate proteome-wide missense variant effect prediction with AlphaMissense. Science 381:eadg7492.
7. Meinhardt S, Manley MW, Parente DJ, Swint-Kruse L (2013) Rheostats and Toggle Switches for Modulating Protein Function. PLoS ONE 8:e83502.
8. Fenton AW, Page BM, Spellman-Kruse A, Hagenbuch B, Swint-Kruse L (2020) Rheostat positions: A new classification of protein positions relevant to pharmacogenomics. Medicinal Chemistry Research 29:1133-1146.
9. Miller M, Bromberg Y, Swint-Kruse L (2017) Computational predictors fail to identify amino acid substitution effects at rheostat positions. Scientific Reports 7:41329.
10. Fowler DM, Fields S (2014) Deep mutational scanning: a new style of protein science. Nature Methods 11:801-807.
11. Fowler DM, Stephany JJ, Fields S (2014) Measuring the activity of protein variants on a large scale using deep mutational scanning. Nature Protocols 9:2267-2284.
12. Esposito D, Weile J, Shendure J, Starita LM, Papenfuss AT, Roth FP, Fowler DM, Rubin AF (2019) MaveDB: an open-source platform to distribute and interpret data from multiplexed assays of variant effect. Genome Biology 20.
13. Siloto RMP, Weselake RJ (2012) Site saturation mutagenesis: Methods and applications in protein engineering. Biocatalysis and Agricultural Biotechnology 1:181-189.
14. Araya CL, Fowler DM (2011) Deep mutational scanning: assessing protein function on a massive scale. Trends Biotechnol 29:435-442.
15. Kinney JB, McCandlish DM (2019) Massively Parallel Assays and Quantitative Sequence-Function Relationships. Annu Rev Genomics Hum Genet 20:99-127.
16. Sharma P, Procko E, Kranz DM (2022) Engineering Proteins by Combining Deep Mutational Scanning and Yeast Display. Methods Mol Biol 2491:117-142.
17. Dewachter L, Brooks AN, Noon K, Cialek C, Clark-Elsayed A, Schalck T, Krishnamurthy N, Versées W, Vranken W, Michiels J (2023) Deep mutational scanning of essential bacterial proteins can guide antibiotic development. Nature Communications 14.
18. Maes S, Deploey N, Peelman F, Eyckerman S (2023) Deep mutational scanning of proteins in mammalian cells. Cell Rep Methods 3:100641.
19. Wei H, Li X (2023) Deep mutational scanning: A versatile tool in systematically mapping genotypes to phenotypes. Frontiers in Genetics 14.
20. Shin H, Cho BK (2015) Rational Protein Engineering Guided by Deep Mutational Scanning. Int J Mol Sci 16:23094-23110.
21. Reuter JA, Spacek DV, Snyder MP (2015) High-throughput sequencing technologies. Mol Cell 58:586-597.

22. Alser M, Lindegger J, Firtina C, Almadhoun N, Mao H, Singh G, Gomez-Luna J, Mutlu O (2022) From molecules to genomic variations: Accelerating genome analysis via intelligent algorithms and architectures. Comput Struct Biotechnol J 20:4579-4599.

23. Chen L, Zhang Z, Li Z, Li R, Huo R, Chen L, Wang D, Luo X, Chen K, Liao C, Zheng M (2023) Learning protein fitness landscapes with deep mutational scanning data from multiple sources. Cell Systems 14:706-721.e705.

24. Coyote-Maestas W, Nedrud D, He Y, Schmidt D (2022) Determinants of trafficking, conduction, and disease within a K+ channel revealed through multiparametric deep mutational scanning. eLife 11:e76903.

25. Park Y, Metzger BPH, Thornton JW (2022) Epistatic drift causes gradual decay of predictability in protein evolution. Science 376:823-830.

26. Gerasimavicius L, Livesey BJ, Marsh JA (2023) Correspondence between functional scores from deep mutational scans and predicted effects on protein stability. Protein Sci 32:e4688.

27. Mathis AD, Otto RM, Reynolds KA (2020) A simplified strategy for titrating gene expression reveals new relationships between genotype, environment, and bacterial growth. Nucleic Acids Research 49:e6-e6.

28. Mavor D, Barlow K, Thompson S, Barad BA, Bonny AR, Cario CL, Gaskins G, Liu Z, Deming L, Axen SD, Caceres E, Chen W, Cuesta A, Gate RE, Green EM, Hulce KR, Ji W, Kenner LR, Mensa B et al. (2016) Determination of ubiquitin fitness landscapes under different chemical stresses in a classroom setting. Elife 5.

29. Melamed D, Young DL, Gamble CE, Miller CR, Fields S (2013) Deep mutational scanning of an RRM domain of the Saccharomyces cerevisiae poly(A)-binding protein. Rna 19:1537-1551.

30. Choudhury A, Gachet B, Dixit Z, Faure R, Gill RT, Tenaillon O (2023) Deep mutational scanning reveals the molecular determinants of RNA polymerase-mediated adaptation and tradeoffs. Nature Communications 14:6319.

31. Chi PB, Liberles DA (2016) Selection on protein structure, interaction, and sequence. Protein Science 25:1168-1178.

32. Vecchi G, Sormanni P, Mannini B, Vandelli A, Tartaglia GG, Dobson CM, Hartl FU, Vendruscolo M (2020) Proteome-wide observation of the phenomenon of life on the edge of solubility. Proc Natl Acad Sci U S A 117:1015-1020.

33. Rocklin GJ, Chidyausiku TM, Goreshnik I, Ford A, Houliston S, Lemak A, Carter L, Ravichandran R, Mulligan VK, Chevalier A, Arrowsmith CH, Baker D (2017) Global analysis of protein folding using massively parallel design, synthesis, and testing. Science 357:168-175.

34. O'Neil PT, Swint-Kruse L, Fenton AW (2024) Rheostatic contributions to protein stability can obscure a position's functional role. Protein Science 33:e5075.

35. Ricatti J, Acquasaliente L, Ribaudo G, De Filippis V, Bellini M, Llovera RE, Barollo S, Pezzani R, Zagotto G, Persaud KC, Mucignat-Caretta C (2019) Effects of point mutations in the binding pocket of the mouse major urinary protein MUP20 on ligand affinity and specificity. Scientific Reports 9:300.

36. Zhan H, Taraban M, Trewhella J, Swint-Kruse L (2008) Subdividing repressor function: DNA binding affinity, selectivity, and allostery can be altered by amino acid substitution of nonconserved residues in a LacI/GalR homologue. Biochemistry 47:8058-8069.

37. Swint-Kruse L, Martin TA, Page BM, Wu T, Gerhart PM, Dougherty LL, Tang Q, Parente DJ, Mosier BR, Bantis LE, Fenton AW (2021) Rheostat functional outcomes occur when substitutions are introduced at nonconserved positions that diverge with speciation. Protein Sci 30:1833-1853.

38. Tang Q, Fenton AW (2017) Whole-protein alanine-scanning mutagenesis of allostery: A large percentage of a protein can contribute to mechanism. Hum Mutat 38:1132-1143.

39. Wu T, Swint-Kruse L, Fenton AW (2019) Functional tunability from a distance: Rheostat positions influence allosteric coupling between two distant binding sites. Scientific reports 9:16957.

40. Swint-Kruse L, Dougherty LL, Page B, Wu T, O'Neil PT, Prasannan CB, Timmons C, Tang Q, Parente DJ, Sreenivasan S, Holyoak T, Fenton AW (2023) PYK-SubstitutionOME: an integrated database containing allosteric coupling, ligand affinity and mutational, structural, pathological, bioinformatic and computational information about pyruvate kinase isozymes. Database (Oxford) 2023.

41. Winkler K, Kramer A, KüTtner G, Seifert M, Scholz C, Wessner H, Schneider-Mergener J, HöHne W (2000) Changing the Antigen Binding Specificity by Single Point Mutations of an Anti-p24 (HIV-1) Antibody. The Journal of Immunology 165:4505-4514.

42. Ruggiero MJ, Malhotra S, Fenton AW, Swint-Kruse L, Karanicolas J, Hagenbuch B (2021) A clinically-relevant polymorphism in the Na(+)/taurocholate cotransporting polypeptide (NTCP) occurs at a rheostat position. J Biol Chem 296:100047.

43. Vila JA (2023) Protein folding rate evolution upon mutations. Biophys Rev 15:661-669.

44. Berg MD, Zhu Y, Ruiz BY, Loll-Krippleber R, Isaacson J, San Luis BJ, Genereaux J, Boone C, Villén J, Brown GW, Brandl CJ (2021) The amino acid substitution affects cellular response to mistranslation. G3 (Bethesda) 11.

45. Markin CJ, Mokhtari DA, Sunden F, Appel MJ, Akiva E, Longwell SA, Sabatti C, Herschlag D, Fordyce PM (2021) Revealing enzyme functional architecture via high-throughput microfluidic enzyme kinetics. Science 373.

46. Otto RM, Turska-Nowak A, Brown PM, Reynolds KA (2024) A continuous epistasis model for predicting growth rate given combinatorial variation in gene expression and environment. Cell Systems 15:134-148.e137.

47. Cagiada M, Johansson KE, Valanciute A, Nielsen SV, Hartmann-Petersen R, Yang JJ, Fowler DM, Stein A, Lindorff-Larsen K (2021) Understanding the Origins of Loss of Protein Function by Analyzing the Effects of Thousands of Variants on Activity and Abundance. Molecular Biology and Evolution 38:3235-3246.

48. Ruggiero MJ, Malhotra S, Fenton AW, Swint-Kruse L, Karanicolas J, Hagenbuch B (2021) A clinically relevant polymorphism in the Na(+)/taurocholate cotransporting polypeptide (NTCP) occurs at a rheostat position. J Biol Chem 296:100047.

49. Wu X, Go M, Nguyen JV, Kuchel NW, Lu BGC, Zeglinski K, Lowes KN, Calleja DJ, Mitchell JP, Lessene G, Komander D, Call ME, Call MJ (2024) Mutational profiling of SARS-CoV-2 papain-like protease reveals requirements for function, structure, and drug escape. Nature Communications 15:6219.

50. Flynn JM, Samant N, Schneider-Nachum G, Barkan DT, Yilmaz NK, Schiffer CA, Moquin SA, Dovala D, Bolon DNA (2022) Comprehensive fitness landscape of SARS-CoV-2 Mpro reveals insights into viral resistance mechanisms. eLife 11:e77433.

51. Bai C, Zhong Q, Gao GF (2022) Overview of SARS-CoV-2 genome-encoded proteins. Sci China Life Sci 65:280-294.

52. Jin Z, Du X, Xu Y, Deng Y, Liu M, Zhao Y, Zhang B, Li X, Zhang L, Peng C, Duan Y, Yu J, Wang L, Yang K, Liu F, Jiang R, Yang X, You T, Liu X et al. (2020) Structure of Mpro from SARS-CoV-2 and discovery of its inhibitors. Nature 582:289-293.

53. Hu Y, Lewandowski EM, Tan H, Zhang X, Morgan RT, Zhang X, Jacobs LMC, Butler SG, Gongora MV, Choy J, Deng X, Chen Y, Wang J (2023) Naturally Occurring Mutations of SARS-CoV-2 Main Protease Confer Drug Resistance to Nirmatrelvir. ACS Central Science 9:1658-1669.

54. Miao M, Clercq E, Li G (2021) Genetic Diversity of SARS-CoV-2 over a One-Year Period of the COVID-19 Pandemic: A Global Perspective. Biomedicines 9.

55. Campitelli P, Lu J, Ozkan SB (2022) Dynamic allostery highlights the evolutionary differences between the CoV-1 and CoV-2 main proteases. Biophys J 121:1483-1492.

56. Ruggiero MJ, Malhotra S, Fenton AW, Swint-Kruse L, Karanicolas J, Hagenbuch B (2021) A clinically relevant polymorphism in the Na+/taurocholate cotransporting polypeptide (NTCP) occurs at a rheostat position. Journal of Biological Chemistry 296:100047.

57. Sreenivasan S, Heffren P, Suh KS, Rodnin MV, Kosa E, Fenton AW, Ladokhin AS, Smith PE, Fontes JD, Swint-Kruse L (2024) The intrinsically disordered transcriptional activation domain of CIITA is functionally tuneable by single substitutions: An exception or a new paradigm? Protein Sci 33:e4863.

58. Gray VE, Hause RJ, Fowler DM (2017) Analysis of Large-Scale Mutagenesis Data To Assess the Impact of Single Amino Acid Substitutions. Genetics 207:53-61.

59. Boyle GE, Sitko KA, Galloway JG, Haddox HK, Bianchi AH, Dixon A, Wheelock MK, Vandi AJ, Wang ZR, Thomson RES, Garge RK, Rettie AE, Rubin AF, Geck RC, Gillam EMJ, DeWitt WS, Matsen FAt, Fowler DM (2024) Deep mutational scanning of CYP2C19 in human cells reveals a substrate specificity-abundance tradeoff. Genetics 228.

60. Matreyek KA, Starita LM, Stephany JJ, Martin B, Chiasson MA, Gray VE, Kircher M, Khechaduri A, Dines JN, Hause RJ, Bhatia S, Evans WE, Relling MV, Yang W, Shendure J, Fowler DM (2018) Multiplex assessment of protein variant abundance by massively parallel sequencing. Nature Genetics 50:874-882.

61. Schulze TK, Lindorff-Larsen K. Effects of residue substitutions on the cellular abundance of proteins. (2024). eLife Sciences Publications, Ltd.

62. Giacomelli AO, Yang X, Lintner RE, McFarland JM, Duby M, Kim J, Howard TP, Takeda DY, Ly SH, Kim E, Gannon HS, Hurhula B, Sharpe T, Goodale A, Fritchman B, Steelman S, Vazquez F, Tsherniak A, Aguirre AJ et al. (2018) Mutational processes shape the landscape of TP53 mutations in human cancer. Nat Genet 50:1381-1387.

63. Wu T, Swint-Kruse L, Fenton AW (2019) Functional tunability from a distance: Rheostat positions influence allosteric coupling between two distant binding sites. Sci Rep 9:16957.

64. Campitelli P, Ross D, Swint-Kruse L, Ozkan SB (2024) Dynamics-based protein network features accurately discriminate neutral and rheostat positions. Biophys J 123:3612-3626.

65. Teilum K, Olsen JG, Kragelund BB (2011) Protein stability, flexibility and function. Biochim Biophys Acta 1814:969-976.

66. Klesmith JR, Bacik J-P, Wrenbeck EE, Michalczyk R, Whitehead TA (2017) Trade-offs between enzyme fitness and solubility illuminated by deep mutational scanning. Proceedings of the National Academy of Sciences 114:2265-2270.

67. Unger T, Nau MM, Segal S, Minna JD (1992) p53: a transdominant regulator of transcription whose function is ablated by mutations occurring in human cancer. Embo j 11:1383-1390.

68. Kooistra AJ, Mordalski S, Pándy-Szekeres G, Esguerra M, Mamyrbekov A, Munk C, Keserű GM, Gloriam David E (2021) GPCRdb in 2021: integrating GPCR sequence, structure and function. Nucleic Acids Research 49:D335-D343.

69. Xie WJ, Warshel A (2022) Natural Evolution Provides Strong Hints about Laboratory Evolution of Designer Enzymes. Proc Natl Acad Sci U S A 119:e2207904119.

70. Xie WJ, Asadi M, Warshel A (2022) Enhancing computational enzyme design by a maximum entropy strategy. Proceedings of the National Academy of Sciences 119:e2122355119.

71. Khersonsky O, Röthlisberger D, Dym O, Albeck S, Jackson CJ, Baker D, Tawfik DS (2010) Evolutionary optimization of computationally designed enzymes: Kemp eliminases of the KE07 series. J Mol Biol 396:1025-1042.

72. Hou Q, Rooman M, Pucci F (2023) Enzyme Stability-Activity Trade-Off: New Insights from Protein Stability Weaknesses and Evolutionary Conservation. Journal of Chemical Theory and Computation 19:3664-3671.

73. Mu GC, Nie Y, Mu XQ, Xu Y, Xiao R (2015) Single Amino Acid Substitution in the Pullulanase of Klebsiella variicola for Enhancing Thermostability and Catalytic Efficiency. Applied Biochemistry and Biotechnology 176:1736-1745.

74. Upadhyay V, Lucas A, Panja S, Miyauchi R, Mallela KMG (2021) Receptor binding, immune escape, and protein stability direct the natural selection of SARS-CoV-2 variants. J Biol Chem 297:101208.

75. O'Neil PT, Swint-Kruse L, Fenton AW (2024) Rheostatic contributions to protein stability can obscure a position's functional role. Protein Sci 33:e5075.

76. Page BM, Martin TA, Wright CL, Fenton LA, Villar MT, Tang Q, Artigues A, Lamb A, Fenton AW, Swint-Kruse L (2022) Odd one out? Functional tuning of Zymomonas mobilis pyruvate kinase is narrower than its allosteric, human counterpart. Protein Sci 31:e4336.

77. Nisthal A, Wang CY, Ary ML, Mayo SL (2019) Protein stability engineering insights revealed by domain-wide comprehensive mutagenesis. Proceedings of the National Academy of Sciences 116:16367-16377.

78. Hodges AM, Fenton AW, Dougherty LL, Overholt AC, Swint-Kruse L (2018) RheoScale: A tool to aggregate and quantify experimentally determined substitution outcomes for multiple variants at individual protein positions. Hum Mutat 39:1814-1826.

79. Fink T, Lonzarić J, Praznik A, Plaper T, Merljak E, Leben K, Jerala N, Lebar T, Strmšek Ž, Lapenta F, Benčina M, Jerala R (2019) Design of fast proteolysis-based signaling and logic circuits in mammalian cells. Nat Chem Biol 15:115-122.

80. Yaghi RM, Andrews CL, Wylie DC, Iverson BL (2024) High-Resolution Substrate Specificity Profiling of SARS-CoV-2 Mpro; Comparison to SARS-CoV Mpro. ACS Chemical Biology 19:1474-1483.

81. Fenton KD, Meneely KM, Wu T, Martin TA, Swint-Kruse L, Fenton AW, Lamb AL (2022) Substitutions at a rheostat position in human aldolase A cause a shift in the conformational population. Protein Sci 31:357-370.

82. Høie MH, Kiehl EN, Petersen B, Nielsen M, Winther O, Nielsen H, Hallgren J, Marcatili P (2022) NetSurfP-3.0: accurate and fast prediction of protein structural features by protein language models and deep learning. Nucleic Acids Research 50:W510-W515.

83. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE (2004) UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem 25:1605-1612.

84. Douangamath A, Fearon D, Gehrtz P, Krojer T, Lukacik P, Owen CD, Resnick E, Strain-Damerell C, Aimon A, Ábrányi-Balogh P, Brandão-Neto J, Carbery A, Davison G, Dias A, Downes TD, Dunnett L, Fairhead M, Firth JD, Jones SP et al. (2020) Crystallographic and electrophilic fragment screening of the SARS-CoV-2 main protease. Nat Commun 11:5047.

**FIGURES**

**Figure 1. Mpro positions can be classified into five types of substitution sensitivities. (a)** Outcomes previously measured in the DMS growth assay[50] fully sample the measurable range of the reported "combined effects". On this histogram, the bin corresponding to the value for inactive Mpro (C145A) is marked with a magenta circle and that for wild-type (WT) Mpro is marked with a green circle. The error range from replicate WT samples is indicated by the green horizontal bar. Numbers on the X-axis correspond to the upper bin value. **(b)** The percent of each substitution class identified from Mpro DMS growth assay: WT-like Mpro substitutions have values that overlap the WT error range (green line in (a)), inactive Mpro substitutions occupy the bin marked by the magenta circle in (a), and substitutions with intermediate activity occupy the other bins. Similar results were obtained for the other two assays described by *Flynn et al.,*[50] (**Supplementary Figure 1**). **(c)** Each position's substitution sensitivity was analysed with the RheoScale calculator[78] and assigned to one of the five classes defined in **Supplement table 1**. The five classes are coloured on the Mpro structure using UCSF Chimera[83] according the legend shown in (d) (PDB: 5R7Y[84]). Note that rheostat positions (black) are widely distributed across the protein. **(d)** The percent of Mpro positions in each substitution class; assignments for individual positions in this study are listed in **Supplementary table 1c**.

**Figure 2. Outcomes measured for the 60 single amino acid Mpro substitutions assessed in this study**. **(a)** Luciferase assay output, which measured the combined effects on function and *in vivo* abundance; **(b)** abundance changes were measured by indirect sandwich ELISA; **(c)** function changes were computed by normalizing the measured combined effects with abundance changes. The lower bounds for these three ranges are delineated by inactive Mpro (C145A; a and c) or no protein (b) and are shown with magenta circles. The interval encompassing replicate assays for WT-like variants is marked by the green circle. Note that all three assays showed substitutions with gain-of-function outcomes.

**Figure 3. Outcomes measured for the set of substitutions at Mpro position 77. (a)** Combined effects of substitutions were assessed using a luciferase reporter assay. **(b)** Abundance changes were independently measured using indirect sandwich ELISA. **(c)** Measurements for combined effects **(a)** and abundance changes **(b)** were used to compute the abundance-normalized function for each substitution. Note that the X-axes for the three plots are ordered the same. For all three data sets, substitution outcomes were normalized to the WT value (dashed line at y=1). The green regions delineate the standard deviation of all replicate WT values measured during this study. For each substitution, six technical replicates for each of two biological replicates were measured (open circles); bars show the average for each variant and error bars are standard deviations. Error propagation for (c) is described in Methods. Variants with outcomes statistically different from WT were identified using one-way ANOVAs with Dunnett's correction (****p <0.0001, ***p<0.001, **p<0.01, *p<0.1); significance asterisks are shown on the X-axis labels. Data for the other five Mpro positions are shown in **Supplementary figure 5** and average values for all variants are available in **Supplementary data**.

**Figure 4. Changes in abundance can make significant contributions to changes in the luciferase assay output.** Each data point corresponds to the average values shown in **Supplementary figure 5** and listed in **Supplementary data**; error bars correspond to the standard deviations for each measurement. **(a)** "Combined effects" of the luciferase assay output versus *in vivo* abundance. The Spearman correlation coefficient is 0.04. **(b)** Normalized function versus *in vivo* abundance. For the whole data set, Spearman r is -0.68, which indicates that enhanced proteolytic activity is often accompanied by diminished Mpro abundance. However, individual plots (**Supplementary figure 8**) show distinct behaviours for individual positions: Positions 48, 50 and 52 show strong relationships, whereas positions 110 and 234 show moderate relationships. Position 77, shown here with large magenta circles, has the weakest correlation (Spearman r = -0.5).

**Figure 5. Distinct combinations of substitution sensitivities were observed at the six tested Mpro positions.** The six Mpro positions tested in this work are indicated with coloured spheres on one monomer of PDB 5R7Y[84]; colours correspond to the different combinations of substitution sensitivities measured for abundance and normalized function **(Table 1)**. The histidine and cysteine side chains forming the catalytic dyad (at positions 41 and 145) are in red and the active site is indicated by the red star. Orange position 110: neutral for both abundance and normalized-function. Green positions 48 and 77: rheostatic outcomes for both abundance and normalized function. Blue position 52: moderate rheostat for both abundance and normalized-function. Magenta position 50: moderate rheostat for abundance and rheostat for normalized-function. Cyan position 234: neutral for abundance and moderate rheostat for normalized-function. Figure rendered using UCSF Chimera.[83]