SpikeMAP: An unsupervised pipeline for the identification of cortical excitatory and
inhibitory neurons in high-density multielectrode arrays with ground-truth validation

Giraud, E. [2], Lynn, M. [1,3], Vincent-Lamarre, P. [1, 2], Béïque, J-C. [1, 3], Thivierge, J-P. [1, 2]
[1] uOttawa Brain and Mind Research Institute Center for Neural Dynamics
[2] School of Psychology, University of Ottawa
[3] Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa

Abstract Large-scale extracellular recording techniques represent a major advance in interrogating the
structure and dynamics of neuronal circuits. However, methods that can resolve cell-type identity in a
principled way, while simultaneously scaling to thousands of neurons, are currently lacking. Here, we
introduce spikeMAP, a pipeline for the analysis of large-scale recordings of *in vitro* cortical activity that
not only allows for the detection of spikes produced by single neurons (spike sorting), but also allows for
the reliable distinction between genetically determined cell types by utilizing viral and optogenetic
strategies as ground-truth validation. This approach tightly integrates the data analysis pipeline to an
optogenetic, viral, and pharmacological protocol allowing for the dynamical probing of distinct cell-types
while simultaneously recording from large populations. The novelty of spikeMAP is to combine a stream
of well-established analysis techniques in an end-to-end fashion, creating a unified framework as
follows. First, individual spike waveforms are fitted by spline interpolation to estimate their half-
amplitude and peak-to-peak durations. These values are then entered in a principal component analysis
with k-means clustering to identify uncorrelated signals from single channels on the array. Optimal
separability of clusters is assessed by linear discriminant analysis. Finally, each channel's source location
is identified using spatiotemporal characteristics of spike waveforms across the array. We show that
spikeMAP can resolve cell type identity in high-density arrays by analyzing activity monitored from
mouse prefrontal cortex *in vitro* slices with an array of 4,096 closely-spaced channels. Using an
optotagging functional strategy, we show an effective distinction of regular-spiking excitatory neurons
from fast-spiking inhibitory interneurons using measures of action potential waveform, Fano factor, and
spatially-dependent cross-correlations. In sum, the approach introduces a toolbox, validated by an
experimental pipeline, that allows for a comprehensive characterization of neuronal activity obtained
from different cell-types in high-density multielectrode recordings. This provides a scalable approach to
investigate the interplay between distinct cell types in microcircuits of the brain.

Summary We introduce spikeMAP, an automated cell-type specific spike sorting pipeline tailored to
high-density, large-scale multielectrode array recordings. We integrate this analysis pipeline to an
experimental protocol allowing for the dynamical probing of distinct cell-types while simultaneously
recording from large populations of neurons. Using high-density multielectrode arrays containing 4,096
closely spaced electrodes, we monitor the spiking activity of PFC neurons in acute slice preparations.
Our spike sorting technique combines spline interpolation and principal component analysis to
distinguish regular-spiking excitatory neurons from fast-spiking inhibitory interneurons. We provide an
experimental validation of our sorting algorithm using a targeted combination of pharmacological, viral
and optogenetic strategies.

Introduction

The advent of high-density, large-scale multielectrode array recordings has advanced our understanding of how population of neurons represent and process information, offering crucial insights into complex network dynamics (Riquelme et al., 2023; Hemberger et al., 2019; Shein-Idelson et al., 2017, Thivierge et al., 2022). Furthermore, the use of silicon recording probes such as Neuropixels probes *in vivo* has helped uncovered foundational principles of neural circuits in awake behaving mice (Lak et al., 2020; Peters et al., 2021). However, it remains a challenge to distinguish neuronal cell types based on their physiological properties alone, despite a growing appreciation for the importance of heterogeneous cell types in circuit-level computations (Sylwestrak et al., 2022; Liu et al., 2021; Bugeon et al., 2022). The scalability of large-scale electrophysiological recording techniques depends not only on the detection and isolation of spikes produced by single neurons (spike sorting), but also on the reliable distinction between those triggered by different cell-types.

The recording of extracellular potentials is a popular method to investigate neural activity, offering a high spatiotemporal resolution whilst allowing for the monitoring of action potentials from thousands of cells simultaneously (Magland et al., 2020; Buzsáki, 2004). Cell-type sorting based on extracellular action potential waveforms has been extensively explored. It has been proposed that excitatory (E) and inhibitory (I) neurons can be distinguished based on action potential waveforms (Hafizi et al., 2022; Ren et al., 2020; Bathó et al., 2004). These methods typically extract features of action potentials such as peak-to-peak time difference, and full-width-half amplitude maximum, and match these features to reported values for each cell types to distinguish them. The rationale for using these features as a proxy for cell-type identity stems from intracellular recording studies, which report that inhibitory cells, such as parvalbumin-expressing GABAergic interneurons, have short-duration action potentials (Nowak et al., 2003; Peyrache and Destexhe, 2019; Kawaguchi, 1993; Connors and Gutnick, 1990), whereas excitatory pyramidal neurons have longer-duration action potentials (McCormick et al., 1985). Narrow and broad action potentials have been distinguished from one another in several neocortical areas, including prefrontal cortex (Mitchell et al., 2007; Wilson et al., 1994; Constantinidis and Goldman-Rakic, 2002; Gonzalez-Burgos et al., 2005; Hasenstaub et al., 2005, Bartho et al., 2004, Tamura et al., 2004). These differences have been hypothesized to result from the fact that repolarization of the membrane potential following a spike is slower among broad-spiking pyramidal neurons (McCormick et al., 1985; Nowak et al., 2003; Hasenstaub et al., 2005). In the extracellular signal, which resembles the negative derivative of the intracellular membrane potential, this gives both a broader and shallower peak following the initial trough (Henze et al., 2000).

Modern extracellular recording techniques allow researchers to monitor the simultaneous activity of large, intact networks of neurons. However, in dense preparations, the activity of a single cell may be captured by several electrodes, and each electrode may report the activity of several nearby cells. Thus, to segregate the activity of single cells, effective spike-sorting techniques must be employed. Spike sorting typically includes filtering the raw extracellular activity, detecting action potentials using threshold crossings, extracting spike features, and clustering to distinguish amongst neurons (Rey et al., 2015).

Several methods have been developed , each with specialized features, including Kilosort4 (Pachitariu et al., 2023), SpyKING CIRCUS (Yger et al., 2018), MountainSort (Chung et al., 2017), Herding Spikes (Hilgen et al., 2017), YASS (Lee et al., 2017), and JRCLUST (Jun et al., 2017). A detailed comparison of these methods is presented in Table 1. A common lacking feature in these methods is the ability to identify

cell-types in a biologically validated fashion. Despite evidence showing differences in action potential
kinetics for distinct cell-types as well as the use of optogenetics (Hilgen et al., 2017), there exists no
large-scale validation efforts, to our knowledge, showing that extracellular waveforms can be used to
reliably distinguish cell-types. Methods leveraging genetic differences between cell types offer statistical
measures predictive of cell types, such as Fano Factor (Becchetti et al., 2012). These methods have
provided researchers with the ability to infer, in a post-hoc fashion, the most probable cell types that
were sampled. Yet, a fully integrated suite offering an open-source data analysis pipeline paired with a
validated experimental protocol allowing for high-density cell type-specific activation during recordings
is needed. Such method will enable the investigation of dense and intricate circuits in a flexible, scalable,
and generalizable manner.

Here, we introduce spikeMAP, an open-source, unsupervised and scalable spike sorting analysis pipeline
which performs cell-type classification as validated on high density MEAs. Crucially, we tightly integrate
our analysis with an experimental protocol utilizing a stabilized step function opsin to dynamically probe
distinct populations and validate sorting while avoiding the light-induced artefacts common on CMOS-
based chip designs (Fiscella et al., 2012). Our analysis pipeline, taking advantage of dense sampling,
estimates cell-body location for each detected spike, yielding well separated clusters associated with
single neurons. Spike waveform features were extracted utilizing signal interpolation to estimate half-
amplitude and peak-to-peak durations. These values were then entered in a principal component
analysis with k-means clustering to identify uncorrelated signals from single channels on the array.
Optimal separability of clusters was assessed by linear discriminant analysis. Each channel's source
location was identified using spatiotemporal characteristics of spike waveforms across the array. We
further classified cells as putative excitatory or inhibitory based on their waveform properties. As a final
validation, we introduced a method taking advantage of long timescale optogenetics (step function
opsins) to perform cell-type-specific stimulation in acute prefrontal cortex slices from mice, providing a
biological verification to our statistical methods. Together, spikeMAP provides a comprehensive toolkit
for experimenters to flexibly interrogate cell type-specific computations in a variety of neural circuits
throughout the brain.

Results

Spike sorting

Our spike sorting pipeline (Figure 1) performs three main tasks: (i) the detection of voltage deflections
associated with putative somatic spikes, (ii) the identification of individual putative neurons using spike
clustering, and (iii) the spatial localization of single cells in high-density recordings. We demonstrate the
use of this pipeline through targeted high-density recording from medial prefrontal cortex of mouse
(Figure 2A; see Materials and Methods). First, putative spike times were detected as threshold crossings
(Figure 2Bi; Lewicki, 1998), identifying negative signal peaks from the mean of the filtered signal (see
Materials and Methods). Optimal threshold choice is key in discriminating spikes from noise, as the
number of spikes detected per electrode is highly dependent on the threshold used (Figure 2C). Here,
the optimal threshold for detection was selected by fitting the voltage histogram of each channel with a
Gaussian distribution to separate spiking activity from normally distributed noise (Figure 2D). At a
threshold of -3σ, the signal could be largely accounted for by Gaussian noise, while a separation
between signal and noise began around a threshold of -4σ. To avoid multiple detections of the same
event, an absolute refractory period was imposed to discard successive events within <2 ms. Although
filtering is crucial in event detection, it can introduce signal distortion (Yael and Bar-Gad, 2017). To

resolve this issue, the unfiltered voltage around the time of each spike was extracted (Figure 2Bii),
yielding a series of high-frequency (18 kHz) waveforms for each channel.

In high density preparations, a single electrode can report the activity of multiple cells. To address this
issue, we employed a principal component analysis on the identified spike waveforms from individual
channels (Rossant et al., 2016), which revealed distinct clusters putatively corresponding to individual
cells (Figure 2E), indicating that more than one neuron likely contributes to the voltage recorded on
some channels. The SpikeMAP suite also offers a routine to select a radius around individual channels in
order to enter groups of adjacent channels in PCA. Next, we employed a k-means cluster analysis to
extract clusters amongst the principal components. To determine how well the data was separated by
this approach, we applied a linear discriminant analysis (LDA) to the labeled principal components (Hill
et al., 2011), allowing our method to remain completely unsupervised. In SpikeMAP, the optimal
number of k-means clusters can be chosen by a Calinski-Harabasz criterion (Calinski and Harabasz, 1974)
or pre-selected by the user.

To ensure the activity of each single cell was only reported once, we set a radius of 250 µm around the
electrode where a given cell was detected. This radius corresponds to the approximate spread of
extracellular electrical field around the soma of individual neurons (Jia et al., 2019). Within this radius,
the channel with the largest negative deflection in mean waveform was identified. Previous work
observed that the largest negative amplitude of extracellular signal originates from the axon initial
segment (AIS) (Bakkum et al., 2019; Obien et al., 2015). Utilizing spatial localization of the AIS, the
channel showing the greatest negative deflection was likely the closest to the putative soma of a given
cell (Jia et al., 2019). Activity at this channel was consistent with the somatic activity, showing a decay in
peak voltage as we moved away from the putative soma (Figure 2H; Ruz et Schultz, 2014; Pettersen and
Einevoll, 2008; Linden et al., 2011; Mechler et al., 2011) (Supplemental Figure 1). Visualization of
adjacent electrodes revealed a complex structure of voltage activity, possibly due to activity propagation
within the compartments of single neuron (Figure 2F; Shein-Idelson et al., 2017; Obien et al., 2015). As
an additional verification step, SpikeMAP allows the computation of spike-count correlations between
putative neurons located within a user-defined radius. Signals that exceed a defined threshold of
correlation can be rejected as they likely reflect the same underlying cell. To further verify that single
cells located near the same electrode could be distinguished, we estimated the precise soma locations
between adjacent recording electrodes using a center-of-mass analysis (see Materials and Methods).
This revealed subtle changes in position, reflected by distinct weights on each nearby electrodes (Figure
2I. Despite these subtle changes, the soma location could still be assigned to same electrode since
changes in localization across spikes were far less than the inter-electrode distance.

In sum, SpikeMAP provides an end-to-end pipeline to perform spike-sorting on high-density
multielectrode arrays. Some elements of this pipeline are similar to related approaches (Table 1),
including the use of voltage filtering, PCA, and k-means clustering. Other elements are novel, including
the use of spline interpolation, LDA, and the ability to identify putative excitatory and inhibitory cells. At
different steps in the process, conditions for rejecting spikes can be tailored by applying: (1) a stringent
threshold to filtered voltages; (2) a minimal cut-off on the signal-to-noise ratio of voltages (see
Supplemental Figure 2); (3) an LDA for cluster separability; (4) a minimal spike rate to putative neurons;
(5) a Hartigan statistical dip test to detect spike bursting; (6) a decrease in voltage away from putative

somas; and (7) a maximum spike-count correlation for nearby channels. Together, these criteria allow
SpikeMAP users the ability to precisely control parameters relevant to automated spike sorting.

Cell type classification based on action potential waveform kinetics

Despite technological advances, the ability to identify distinct cell types based solely on extracellular
voltage has remained challenging. Current data processing methods often rely on waveform kinetics
(Bathó et al., 2004) to return a classification of contributing excitatory and inhibitory units. As
technological progress continuously allows for denser arrays, waveform spike analyses are increasingly
sensitive to noise generated by the signal of proximal units. Here, we first implemented a classification
method using current knowledge about cell-type specific waveform kinetics and spline interpolation to
distinguish putative excitatory and inhibitory neurons in dense networks. We then validated this
classification using a combination of viral strategies, optogenetics and pharmacology while
simultaneously recording from stimulated mouse prefrontal cortex networks.

To provide a more precise description of the shape of each spike, we first fitted each waveform using
piecewise cubic spline interpolation, a method allowing for the statistical inference of points within the
boundaries of a set of known points. While several interpolation bases are possible, cubic splines allow
for smooth interpolation between known data points and a high accuracy in terms of curve fitting
(Ramsay & Silverman, 1997). Using query points at 90 kHz, we computed the mean fitted waveform
(Blanche, 2006) for each putative cell (see Materials and Methods). While we found that a resolution of
90 kHZ provided a reasonable estimate of spike waveforms, this value can be adjusted as a parameter in
SpikeMAP.

Next, we identified putative excitatory and inhibitory neurons using two time constants obtained from
mean spike waveforms at somatic locations: (1) the full width (FW) of spikes at half-maximum
amplitude; and (2) the peak-to-peak (PP) duration (Figure 3A). (Sirota et al., 2008; Sakata & Harris, 2009,
Barthó et al., 2004; Insel & Barnes, 2015). Although some work has cast doubt on the accuracy of spike-
width based classification (Moore and Wehr, 2013), many studies have shown good classification
accuracy (Nowak et al., 2003; Kawaguchi and Kubota, 1997; Barthó et al., 2004; Madisen et al., 2012;
Stark et al., 2013; Cardin et al., 2009; Cohen and Mizrahi, 2015). K-means clustering was performed on
values of FW and PP obtained across all simultaneously recorded somatic electrodes, where the optimal
number of clusters was chosen with a Calinski-Harabasz criterion (Calinski and Harabasz, 1974;
Vendramin et al, 2009). For validation purposes, strict criteria were applied at each step of the pipeline,
resulting in the low the number of neurons reported here. Clusters corresponding to putative excitatory
(E) and inhibitory (I) neurons were identified based on FW and PP values (Figure 3A). I cells were
identified as a single cluster with rapid time constants for both FW and PP (Barthó et al., 2004). We
extracted the putative soma locations of excitatory and inhibitory neurons on the MEA (Figure 3B), and
visualized firing activity in a raster plot (Figure 3C) (see Supplemental Figure 3 for firing rate
distributions). It is possible that the spatial location of putative I cells reflects the site of injection of the
opsin in medial prefrontal cortex. Other properties of putative I cells were consistent with past work,
including a higher Fano factor (Figure 3D; Becchetti et al., 2012) and a rapid spatial decay of pairwise
correlation (Figure 3E; Peyrache et al., 2012). Further, SpikeMAP contains a routine to perform a
Hartigan statistical dip test on the inter-spike distribution of individual cells to detect putative bursting
neurons. These results extend previous findings to large-scale recordings, yet a ground-truth validation

of cell type is still required to assess the extent to which this approach offers a reasonable proxy of
biological identity, as examined next.

A step-function opsin validation approach for cell classification

While multielectrode arrays have allowed the simultaneous recording from thousands of electrodes, the
reliance on light-sensitive complementary metal-oxide-semiconductor (CMOS) circuits has limited the
ability of researchers to use them in conjunction with optogenetics. As the ability to rapidly perturb
distinct populations in large network provides valuable insight as to how local circuits represent
information (Lemon et al., 2021; Buzsáki, 2010), it is of broad interest to validate methodologies that
allow for optogenetic manipulation of defined cell populations in multielectrode arrays.

We validated an experimental protocol using a stabilized step-function opsin (Bi-stable neural state
switches; Berndt et al., 2009), which is activated in a step-like fashion for prolonged periods (minutes)
by brief photostimulation with blue light and inactivated by green-yellow light. This viral and
optogenetic strategy, used in conjuncture with CMOS based multielectrode arrays, allows for the
monitoring of induced spiking beyond the initial light-generated artefacts (Supplemental Figure 4). We
demonstrated this by inducing spiking selectively in targeted PV inhibitory neurons, as more than 90% of
them have been reported to be fast-spiking (Stringer et al., 2016), to verify whether cells previously
identified as putative inhibitory interneurons indeed correspond to light-activated cells.

As an obligate validation step, we injected a viral vector expressing the step function opsin, pAAV-Ef1a-
DIO hChR2 (C128S/D156A)-EYFP, in medial prefrontal cortex of mice, and after waiting 2-3 weeks for
transfection, performed whole-cell recordings from pyramidal neurons located near identified PV
interneurons using *in vitro* slice preparations (Figure 4A; see Materials and Methods). Photostimulation
with blue light was sufficient to activate PV interneurons, as monitored by the robust, rapid and
consistent appearance of light-induced inhibitory synaptic postsynaptic currents (IPSCs) recorded from
pyramidal neurons (note the absence of direct photocurrent from these cells; Figure 4B). The
subsequent delivery of short pulses of yellow light reduced the frequency of IPSCs onto recorded
pyramidal neurons, indicating that the step-function opsin could be reliably activated and inactivated in
a step-like manner by phasic delivery of light (Figure 4B). Upon PV photoactivation, recorded pyramidal
neurons reliably received high-frequency barrages of 10 to 30 pA IPSCs, followed by a gradual reduction
in amplitude and elevated rates of inhibitory input that persisted until deactivation with green-yellow
light, reflecting some desensitization of the step-function opsin (Figure 4B).

To characterize the refractory period of the step-function opsin, we delivered repeated
photostimulation at various inter-trial intervals and measured steady-state activation (Figure 4C to E).
Longer inter-trial intervals (≥120 seconds) (Figure 4E) allowed expressing cells to recover and prevented
the reduction in steady-state activation observed with 20-second and 60-second inter-trial intervals
(Figure 4C and D). Specifically, the mean IPSCs/s for 20-second, 60-second, and 120-second intervals
were 12.64 ± 0.27, 17.32 ± 0.15, and 26.29 ± 0.31, respectively. Statistical comparisons, performed using
the Wilcoxon rank sum test for non-parametric data on the PSTH, revealed significant differences
between the 20-second and 120-second conditions ($P = 1.29 \times 10^{-6}$, n = 4500 and 3000 trials,
respectively) as well as between the 60-second and 120-second conditions ($P = 0.032$, n = 3000 trials per
condition). The number of trials conducted for each condition was 5 for 20 seconds, 6 for 60 seconds,
and 6 for 120 seconds.

Furthermore, the reduction in firing rate upon opsin deactivation was more prominent for longer
interval trials (Figure 4E). These results provided key insights to inform multielectrode array recording
protocols, allowing us to selectively activate and inactivate PV interneurons in a temporally precise
fashion using tools compatible with multielectrode array technology.

Optogenetic validation of E/I classification on multielectrode arrays

To validate our prior waveform-based cell-type classification in mouse prefrontal cortex slices, we
selectively activated transfected PV interneurons with the step-function opsin construct while
performing multielectrode recordings (see Supplemental Table 1). To prevent the propagation of
optogenetic activation throughout the network, we first recorded in the presence of 10 μM
cyanquixaline (CNQX) and 100 μM picrotoxin (PTX) to block AMPA receptors and GABA$_A$ receptors,
respectively (leading to a reduction in FR of 0.0937Hz, $p < 0.001$ from baseline). We photostimulated
with brief 100 millisecond pulses of blue light and examined post-stimulation responses of all identified
neurons recorded ($n = 8189$, 3 slices, from 3 mice) (Figure 5A). In the presence of synaptic blockers,
optogenetic activation of targeted PV interneurons led to a significant increase in mean firing rate for
the population of cells that had been classified as inhibitory based on their waveform properties (I
neurons; Figure 5B, $n = 430$, $p < 0.003$). Neurons classified as excitatory neurons based on their
waveform kinetics, on the other hand, showed only a subtle decrease in mean firing rate (Figure 5B, $n = $
7759, $p < 0.001$). A clustering analysis on the spike waveform properties of putative excitatory and
inhibitory neurons provides further evidence of classification (Supplemental Material Figure 5). The
overall firing rates in slice recordings were relatively low and thus, were not permissive to the use of
cross-correlations in validating putative excitatory and inhibitory neurons. While our pharmacological
manipulations attempted to largely eliminate network effects, the slight reduction in firing rate of
putative excitatory neurons may reflect the involvement of GABA$_B$ receptor-mediated inhibition (Kohl &
Paulsen, 2010).

Next, we examined in further detail the response profiles of individual neurons to photostimulation in
order to validate our algorithm at the single cell level. In the absence of synaptic blockers, individual
neurons displayed a diverse range of responses to light stimulation, including neurons showing no
significant change in firing rate (ΔFR ~ 0), neurons showing an increase in firing rate (ΔFR > 0), and
neurons showing a decrease in firing rate (ΔFR < 0) (Figure 5C). We applied a strict criterion to separate
neurons showing a statistically robust increase in firing rate (see Materials and Methods). We found that
56% of neurons that had been classified as I displayed a light-induced increase in firing rate. Conversely,
we found that only 8% of neurons which had been classified as E showed an increase in firing rate after
photostimulation. The percentage of I cells responsive to light may be influenced by our viral strategy,
which targets only a subset of PV-expressing interneurons largely located in the infralimbic and
prelimbic cortical subregions. Additionally, non-PV inhibitory interneurons, and some excitatory neurons
could have been initially classified as inhibitory based on their waveform properties. These experiments,
however, unambiguously identified a subset of genetically-identified PV neurons: those that showed a
transient photostimulation-induced increase in firing rate. Through a comparison with FWHM and PTP
features, these results demonstrate that a step-function opsin approach is a viable and scalable method
for validating spike waveform classification on multielectrode arrays.

Together, this provides evidence for cell classification of excitatory and inhibitory neurons based on
their extracted waveform properties using spikeMAP, showing that cells identified as inhibitory
displayed optogenetic-specific change in responses.

Discussion

Extracellular electrophysiological recordings have allowed to monitor the spiking of thousands of neurons simultaneously. To meaningfully interpret these dynamics, spike sorting is a crucial step in the analysis of those recordings. The analysis of network dynamics can be severely impacted by improper interpretation of the underlying neural activity (Rey et al., 2015; Carlson and Carin, 2019), highlighting the need for optimal methods. With technological progress constantly allowing for denser arrays, assigning spikes to single cells and determining cell-type identity requires an automated solution given that the size of the data prevents manual curation. Most spike sorting methods are at least somewhat supervised (Marre et al., 2012; Prentice et al., 2011), which is not feasible on large-scale arrays. Here, we described an open-source, unsupervised, and scalable suite of computational tools that perform spike sorting and cell type classification on large-scale, high-density recordings. We integrated our analysis pipeline with an experimental protocol, allowing for the validation of our statistical method and for the optogenetic manipulation of distinct cell-types in dense networks.

Previous work has detailed cell-type classification based on waveform analysis, but experimental validation has been lacking. To address this and provide biological evidence that this classification indeed correlates with cell-type, we selectively activated PV interneurons using a combination of viral strategies and optogenetics while simultaneously recordings from the whole network. Analysis of optogenetically stimulated trials revealed that cells that had been classified as inhibitory were more likely to display a change in firing rate during optogenetic stimulation.

We found that clustering in two dimensions, representing the full width of spikes at half-maximum amplitude and the peak-to-peak, achieved a high level of success in classifying cell-types. By sampling electrodes from PV-positive transfected cells, we distinguished their action potentials from other cell-types. Our work provides evidence that extracellular recordings can offer insights on the distinct neuronal populations that make up complex networks. Our pipeline encourages the integration of viral strategies, optogenetics and electrophysiological recordings to manipulate complex networks to better understand the contribution of each cell type.

While SpikeMAP is the only known method to enable the identification of putative excitatory and inhibitory neurons on high-density multielectrode arrays (Table 1), several aspects of SpikeMAP included in the spike sorting pipeline (Figure 1) overlap with existing methods, as these constitute required steps prior to performing E/I identification. To enable users the ability to integrate SpikeMAP with existing toolboxes, we provide a modularized suite of protocols so that E/I identification can be performed separately from preliminary spike sorting steps. In this way, a user could carry out spike sorting through Kilosort or another package before importing their data to SpikeMAP for E/I identification.

Cortical inhibitory neurons are heterogeneous, and the fact that fast-spiking PV interneurons represent the largest population exhibiting landmark features of narrow-spike made them an attractive candidate to validate our classification. However, future work could include different inhibitory interneurons such as somatostatin (SOM) and vasoactive intestinal polypeptide (VIP) neurons to improve the classification of inhibitory cell types. Another avenue could involve applying SpikeMAP on artificially generated spike data (Buccino & Einevoll 2021; Laquitaine et al., 2024). Nevertheless, it is possible that distinguishing all inhibitory cell types based on extracellular waveforms alone may not be achievable. Despite this limitation, incremental advancements such as those presented here are crucial for refining our understanding and improving analytical approaches. Given that spike sorting features in vitro can differ

from properties exhibited in vivo (Henze et al., 2000; Nowak et al., 2003; Gonzalez-Burgos et al., 2005),
future work should address if these classification/waveform properties can be generalized to behaving
organisms, for example with tetrodes, optrodes or implanted eletrode arrays.

In sum, our work described a fully integrated method with an open-source, unsupervised and scalable
spike sorting analysis pipeline performing cell-type classification, integrated with an experimental
protocol utilizing a stabilized step function opsin to validate sorting on large-scale MEAs. These tools
collectively provide a framework for investigating large-scale neural dynamics of genetically defined
neural populations.

Materials and Methods; Supplemental

Animals. PV-CRE mice were kept on a 12:12-h light/dark cycle, with access to food and water ad libitum.
All experiments and procedures were performed in accordance with approved procedures and
guidelines set forth by the University of Ottawa Animal Care and Veterinary Services (protocol # 3471).

Stereotaxic Injections. For stereotaxic injections, P21 mice were injected with 0.05 mg/kg
buprenorphine and anesthetized by inhalation of isoflurane. Injections were performed using a 10-µL
Hamilton syringe with a 33-gauge needle, and 1 µL was injected per coordinate. Stereotaxic coordinates
are as follows, from the bregma landmark: mPFC [anterior–posterior (AP), +2.4 —> +2.6; medial–lateral
(ML), ± 0.42; dorsal–ventral (DV), -1.5 —> -1.8]. The DV coordinate was calculated from the surface of
the pia. For ChR2 expression, 400 nL of pAAV-Ef1a-DIO hChR2 (C128S/D156A)-EYFP was injected, per
site, (bilaterally) and animals were left to recover for 2–3 weeks before ex vivo electrophysiological
recordings.

Slices. mPFC slices were prepared from 35 to 42 day old PV-CRE mice, 2–3 weeks poststereotaxic
injections. For slice preparation, mice were anesthetized by inhalation of isoflurane (Baxter Corporation)
and killed by decapitation. The brain was removed, and coronal brain slices were sectioned while
immersed in an ice-cold choline chloride-based cutting solution of the following composition: 119 mM
choline-Cl, 2.5 mM KCl, 1 mM CaCl2, 4.3 mM MgSO4-7H2O, 1 mM NaH2PO4, 1.3 mM sodium L-
ascorbate, 26.2 mM NaHCO3, and 11 mM glucose, and equilibrated with 95% O2 and 5% CO2. Slices
were recovered in a chamber containing standard artificial cerebrospinal fluid solution (aCSF) of the
following composition: 119 mM NaCl, 2.5 mM CaCl2, 1.3 mM MgSO4- 7H2O, 1 mM NaH2PO4, 26.2 mM
NaHCO3, and 11 mM glucose, at a temperature of 37 °C, continuously bubbled with a mixture of 95% O2
and 5% CO2. Slices were recovered for 1 h in the recovery chamber and equilibrated to a temperature of
~25 °C before the recordings were performed. The tissue was continuously perfused with fresh aCSF.
The recordings shown in this paper were 24 minutes long and were as follow: baseline (2 mins), opto-on
(2 mins), opto-off (2 mins), CNQX wash-on (5 mins), CNQX + opto-on (2 mins), CNQX + opto-off (2 mins),
PTX wash-on (5 mins), CNQX and PTX + opto-on (2 mins), CNQX and PTX + opto-off (2 mins). MEA-chip
baseline (no slice) were recorded (30 secs) as control prior to adding each slice. Slices were imaged on
the multi-electrode array and preserved for immunohistochemistry verification of virus expression site.

Whole-Cell Electrophysiology. Neurons were visualized for whole-cell recordings using an upright
microscope: BX61WU upright microscope (60X, 1.0 NA objective; Olympus, Melville, NY). PV neurons
were visually identified by EYFP expression. Whole-cell recordings were performed using borosilicate
glass patch electrodes (3–6 MΩ; World Precision In- struments) pulled on a Narishige PC-10 pipette
puller. Electrical signals were recorded using an Axon Multiclamp 700B amplifier, filtered at 2 kHz, and

digitized at 10 kHz with an Axon Digidata 1440A digitizer. All experiments were performed at room
temperature in Ringer's solution containing (in mM) 119 NaCl, 2.5 CaCl2, 1.3 MgSO4-7H20, 1 NaH2PO4,
and 26.2 NaHCO3, and 11 glucose saturated with 95% O2 and 5% CO2 (pH 7.3, 295–310 mOsm/L).
Electrodes were filled with an intracellular solution of the following compositions (in mM): 115 K-
gluconate, 20 KCl, 10 HEPES, 4 ATP-Mg, 0.5 GTP, 10 Na-phosphocreatine (pH = 7.2-7.3; 270-290
mOsm/L).

Multielectrode array recordings. Extracellular voltage was recorded using High-Density Multi Electrode
Arrays (HD-MEAs) from 3 Brain, Switzerland (www.3brain.com). A detailed description of the technology
behind this system can be found in Imfeld et al.,(2008) and in Berdondini et al.,(2009). High density
recordings from the mouse's prefrontal cortex were performed in vitro using the BioCAM X platform
with HD-MEA Arena chips (3Brain GmbH, Switzerland). This system records using 4,096 square
microelectrodes (pitch = 42 μm) positioned in a 64 x 64 layout (2.67mm x 2.67mm area) with a sampling
rate fs of 18 kHz when recording from the entire 64 x 64 array. Activity was recorded at 12 bits
resolution per channel. The data was visualized and acquired with the BrainWaveX software provided by
3Brain.

Spike sorting

A code library for SpikeMAP in Python and Matlab is available as open source software from:
https://figshare.com/articles/software/_b_SpikeMAP_An_unsupervised_spike_sorting_pipeline_for_cor
tical_excitatory_and_inhibitory_neurons_in_high-density_multielectrode_arrays_b_/28458749

as well as from GitHub: https:// github.com/evgiraud

Filtering. The extracellular voltages recorded by multielectrode arrays typically reflect the dynamics of
multiple cells in proximity to a given electrode (Rey et al., 2015), raising the problem of how to isolate
individual units. The unfiltered voltage at individual channels was characterized by sharp deflections
(Figure 2Bi) that are typical of in vitro extracellular activity in spontaneously active cortical circuits
(Hemberger et al., 2019). As a necessary first step to isolate individual units, we filtered the extracellular
voltage using a second order Butterworth bandpass filter (300-5,000 Hz) (Figure 2Bii) to remove slowly
changing field potentials and high frequency noise (Bullman et al., 2019). Recording artefacts (voltage
deflection > +- 500 uV) were removed and set to the mean voltage.

Spike detection. Putative spike times were detected using a voltage-threshold method (Lewicki, 1998),
identifying negative signal peaks below a threshold of x standard deviation (SD) from the mean of the
filtered signal. Here, x is calculated for each recording, by plotting the filtered voltage histogram for
each channel and fitting it with a Gaussian curve to determine the SD value at which non-normal,
putative spike-related activity is found. An absolute refractory period was applied and successive events
within <2 ms were discarded. Next, the unfiltered voltage around the time of each spike (from -5 to +5
ms) was extracted (Figure 2Bii), yielding a series of high-frequency (18 kHz) waveforms for each channel.
For the data shown here, all 3 recordings used had $x = 4.5$ SD

Isolation of single units. To isolate single-cell activity, we employed a principal component analysis on
individual channels, which revealed distinct clusters putatively corresponding to individual cells (Figure
2E). Next, clusters amongst the principal components of rrSVD were extracted using a k-means cluster
analysis (k=2) (Xu and Wunsch, 2005). A linear discriminant analysis (LDA) was then applied to the

labeled principal components (Hill et al., 2011) to determine how well the data were separated by this
approach. LDA provides an estimate of linear classification error between the two clusters assuming
equal class covariance. Ten-fold cross-validation was employed, where chance corresponds to an error
estimate of 0.5 (Figure 2G). Clusters with error lower than chance were merged by applying the same
label to all waveforms. Each cluster was considered as a putative neuron.

Center-of-mass triangulation. In order to precisely estimate soma locations between each recording
electrode, we used a center-of-mass analysis (Muthmann et al., 2015; Hilgen et al., 2017). For each
putative soma identified previously, a 3-by-3 matrix with elements a_{ij} was constructed by extracting
the waveforms on adjacent electrodes. Location (2,2) of this matrix was the amplitude of the negative
peak in mean waveform of the somatic electrode. Surrounding locations were populated with the
amplitude of the negative peaks of all electrodes immediately adjacent to the somatic electrode. The
row of the center-of-mass is given by

$$r = \frac{\sum_i \sum_j (i \cdot a_{ij})}{\sum_{ij} a_{ij}}$$

and the column is

$$c = \frac{\sum_i \sum_j (j \cdot a_{ij})}{\sum_{ij} a_{ij}} .$$

In addition to center-of-mass triangulation, SpikeMAP includes protocols to perform monopolar
triangulation and grid-based convolution, offering additional options to estimate putative soma
locations based on waveform amplitudes.

Spline Interpolation. The mean waveform for each single cell were resampled at a higher rate using a
piecewise cubic spline interpolation (Matlab R2015b), constructing new points within the boundaries of
a set of known points using query points at 90 kHz.

E/I Classification. We used a combination of cell-type specific viral strategies and optogenetics to
address cell-type distinction. We injected PV-CRE mice (B6.Cg-Pvalb<tm1.1(cre)Aibs>/J) with a stabilized
step-function opsin (pAAV-Ef1a-DIO hChR2 (C128S/D156A)-EYFP) bilaterally in the medial prefrontal
cortex (mPFC) (Berndt et al., 2009). This allows for selective, prolonged activation of parvalbumin (PV)
interneurons with a 100 ms pulse of 465 nm blue light and deactivation with a 10 s pulse of 545-580 nm
green-yellow light. Next, we computed a z-score for each neuron corresponding to photostimulation-
evoked change in firing rate, expressed in units of SD from the baseline mean (Figure 5D). We isolated
cells showing a robust increase in firing rate (z score >3). A total of (n = 8189, 3 slices, from 3 mice) were
classified in this fashion. Overall, 7759 (94.75 %) were reliably classified as excitatory and 430 (5.25 %)
were reliably identified as inhibitory. This number is close to the expected percentage of PV
interneurons in cortex (4-6%) (Markram et al. 2004).

LED Photostimulation. LED photostimulation was delivered using a PlexBright LED module and
controller (465 nm; Plexon). Light was delivered through a 200-µm patch fiber cable with a bare fiber tip
with ~1 cm of glass exposed. Photodeactivation was delivered using a yellow activation wideband 545-
580 nm filter (BP545-580; Olympus).

Immunohistochemistry. Brain slices used for immunohistochemistry were prepared from 35 to 42 day old mice with 1× PBS used as the solvent for all solutions. Mice were anesthetized by inhalation of isoflurane. They were then intracardially perfused with with 5mL of PBS pH=7.4 followed by 10mL of 4% paraformaldehyde/ 0.1M PBS, pH 7.4 for 10 min. Postfixation in 4% PFA was carried out for 2 h before sequential cryoprotection steps in 15% and 30% sucrose over the course of 2 d. Brains were frozen in −30 °C isopentane, mounted in cryomatrix, and cut to 40-μm slices on a cryostat (CM 3050S; Leica). Free-floating sections were blocked in 0.5% PBS-Triton-X-100 (TX-100) for 45 min, and incubated in primary anti-body against GFP (GFP-1020; 1:1000, AvesLabs) in blocking solution at 20 °C. On the following morning, slices were washed three times, for 5 min each time, with 0.5% PBS-Triton-X-100 before incubation in the secondary antibody goat anti-chicken–conjugated AlexaFluor 488 (Jackson Labs, 703-546-155) for 3 h at room temperature. Slices were washed with 0.1% Tx-100 three times, for 5 min each time, before mounting onto slides with Vectashield (Vector Labs). Additionally, all slices used for recordings were saved, kept immerged in Formalin 10% in PBS pH=7.4, for immunohistochemistry to verify viral expression. The protocol is the same as above.

Acknowledgements

This work was supported by a discovery grant from the Natural Sciences and Engineering Council of Canada (NSERC grants 210977; 210989; 2020-06901), the Canadian Institute for Health Research (175325) by the Fonds de recherche du Québec in Nature and technologie (doctoral scholarship), by the Natural Sciences and Engineering Council of Canada (NSERC CGS M scholarship) and the University of Ottawa Brain and Mind Institute.

References

1. Bakkum, D. J., Obien, M. E. J., Radivojevic, M., Jäckel, D., Frey, U., Takahashi, H. & Hierlemann, A. (2019). The Axon Initial Segment is the Dominant Contributor to the Neuron's Extracellular Electrical Potential Landscape. Advanced Biosystems, 3(2), 1800308. https://doi.org/10.1002/adbi.201800308
2. Barthó, P., Hirase, H., Monconduit, L., Zugaro, M., Harris, K. D. & Buzsáki, G. (2004). Characterization of Neocortical Principal Cells and Interneurons by Network Interactions and Extracellular Features. Journal of Neurophysiology, 92(1), 600–608. https://doi.org/10.1152/jn.01170.2003
3. Becchetti, A., Gullo, F., Bruno, G., Dossi, E., Lecchi, M. & Wanke, E. (2012). Exact distinction of excitatory and inhibitory neurons in neural networks: a study with GFP-GAD67 neurons optically and electrophysiologically recognized on multielectrode arrays. Frontiers in Neural Circuits, 6, 63. https://doi.org/10.3389/fncir.2012.00063
4. Berdondini, L., Imfeld, K., Maccione, A., Tedesco, M., Neukom, S., Koudelka-Hep, M. & Martinoia, S. (2009). Active pixel sensor array for high spatio-temporal resolution electrophysiological recordings from single cell to large scale neuronal networks. Lab on a Chip, 9(18), 2644–2651. https://doi.org/10.1039/b907394a
5. Berndt, A., Yizhar, O., Gunaydin, L. A., Hegemann, P. & Deisseroth, K. (2009). Bi-stable neural state switches. Nature Neuroscience, 12(2), 229–234. https://doi.org/10.1038/nn.2247
6. Blanche, T. J., & Swindale, N. V. (2006). Nyquist interpolation improves neuron yield in multiunit recordings. *Journal of neuroscience methods*, *155*(1), 81-91. https://doi.org/10.1016/j.jneumeth.2005.12.031

7. Buccino, A.P., Einevoll, G.T. (2021). MEArec: A fast and customizable testbench simulator for ground-truth extracellular spiking activity. *Neuroinformatics* 19, 185-204. https://doi.org/10.1007/s12021-020-09467-7

8. Bugeon, S., Duffield, J., Dipoppa, M., Ritoux, A., Prankerd, I., Nicoloutsopoulos, D., Orme, D., Shinn, M., Peng, H., Forrest, H., Viduolyte, A., Reddy, C. B., Isogai, Y., Carandini, M., Harris, K. D. A transcriptomic axis predicts state modulation of cortical interneurons. (2022). *Nature* 607, 330–338 https://doi.org/10.1038/s41586-022-04915-7

9. Bullmann, T., Radivojevic, M., Huber, S. T., Deligkaris, K., Hierlemann, A. & Frey, U. (2019). Large-Scale Mapping of Axonal Arbors Using High-Density Microelectrode Arrays. Frontiers in Cellular Neuroscience, 13, 404. https://doi.org/10.3389/fncel.2019.00404

10. Buzsáki, G. (2004). Large-scale recording of neuronal ensembles. Nature Neuroscience, 7(5), 446–451. https://doi.org/10.1038/nn1233

11. Buzsáki G. (2010). Neural Syntax: Cell Assemblies, Synapsembles, and Readers. *Neuron* 68:362–385. doi:10.1016/j.neuron.2010.09.023

12. Caliński, T., & Harabasz, J. (1974). A dendrite method for cluster analysis. Communications in Statistics-theory and Methods, 3(1), 1-27.

13. Cardin, J. A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H. & Moore, C. I. (2009). Driving fast-spiking cells induces gamma rhythm and controls sensory responses. Nature, 459(7247), 663–667. https://doi.org/10.1038/nature08002

14. Carlson, D. & Carin, L. (2019). Continuing progress of spike sorting in the era of big data. *Current Opinion in Neurobiology*, 55, 90–96. https://doi.org/10.1016/j.conb.2019.02.007

15. Chung, J. E., Magland, J. F., Barnett, A. H., Tolosa, V. M., Tooker, A. C., Lee, K. Y., Shah, K. G., Felix, S. H., Frank, L. M. & Greengard, L. F. (2017). A Fully Automated Approach to Spike Sorting. *Neuron*, 95(6), 1381-1394.e6. https://doi.org/10.1016/j.neuron.2017.08.030

16. Cohen, L. & Mizrahi, A. (2015). Plasticity during Motherhood: Changes in Excitatory and Inhibitory Layer 2/3 Neurons in Auditory Cortex. The Journal of Neuroscience, 35(4), 1806–1815. https://doi.org/10.1523/jneurosci.1786-14.2015

17. Connors, B. W. & Gutnick, M. J. (1990). Intrinsic firing patterns of diverse neocortical neurons. Trends in Neurosciences, 13(3), 99–104. https://doi.org/10.1016/0166-2236(90)90185-d

18. Constantinidis, C. & Goldman-Rakic, P. S. (2002). Correlated Discharges Among Putative Pyramidal Neurons and Interneurons in the Primate Prefrontal Cortex. Journal of Neurophysiology, 88(6), 3487–3497. https://doi.org/10.1152/jn.00188.2002

19. Fiscella M, Farrow K, Jones IL, Jäckel D, Müller J, Frey U, Bakkum DJ, Hantz P, Roska B, Hierlemann A. 2012. Recording from defined populations of retinal ganglion cells using a high-density CMOS-integrated microelectrode array with real-time switchable electrode selection. *J Neurosci Meth* **211**:103–113. doi:10.1016/j.jneumeth.2012.08.017

20. González-Burgos, G., Krimer, L. S., Povysheva, N. V., Barrionuevo, G. & Lewis, D. A. (2005). Functional Properties of Fast Spiking Interneurons and Their Synaptic Connections With Pyramidal Cells in Primate Dorsolateral Prefrontal Cortex. Journal of Neurophysiology, 93(2), 942–953. https://doi.org/10.1152/jn.00787.2004

21. Hafizi, H., Nigam, S., Barnathan, J., Ren, N., Stevenson, I. H., Masmanidis, S. C., Newman, E. L., Sporns, O. & Beggs, J. M. (2022). Inhibition-Dominated Rich-Club Shapes Dynamics in Cortical Microcircuits. *bioRxiv*, 2021.05.07.443074. https://doi.org/10.1101/2021.05.07.443074

22. Hasenstaub, A., Shu, Y., Haider, B., Kraushaar, U., Duque, A. & McCormick, D. A. (2005). Inhibitory Postsynaptic Potentials Carry Synchronized Frequency Information in Active Cortical Networks. Neuron, 47(3), 423–435. https://doi.org/10.1016/j.neuron.2005.06.016

23. Hemberger, M., Shein-Idelson, M., Pammer, L. & Laurent, G. (2019). Reliable Sequential
Activation of Neural Assemblies by Single Pyramidal Cells in a Three-Layered Cortex. Neuron,
104(2), 353-369.e5. https://doi.org/10.1016/j.neuron.2019.07.017
24. Henze, D. A., Borhegyi, Z., Csicsvari, J., Mamiya, A., Harris, K. D. & Buzsáki, G. (2000). Intracellular
Features Predicted by Extracellular Recordings in the Hippocampus In Vivo. Journal of
Neurophysiology, 84(1), 390–400. https://doi.org/10.1152/jn.2000.84.1.390
25. Hilgen, G., Sorbaro, M., Pirmoradian, S., Muthmann, J.-O., Kepiro, I. E., Ullo, S., Ramirez, C. J.,
Encinas, A. P., Maccione, A., Berdondini, L., Murino, V., Sona, D., Zanacchi, F. C., Sernagor, E. &
Hennig, M. H. (2017). Unsupervised Spike Sorting for Large-Scale, High-Density Multielectrode
Arrays. Cell Reports, 18(10), 2521–2532. https://doi.org/10.1016/j.celrep.2017.02.038
26. Hill, D. N., Mehta, S. B. & Kleinfeld, D. (2011). Quality Metrics to Accompany Spike Sorting of
Extracellular Signals. The Journal of Neuroscience, 31(24), 8699–8705.
https://doi.org/10.1523/jneurosci.0971-11.2011
27. Imfeld, K., Neukom, S., Maccione, A., Bornat, Y., Martinoia, S., Farine, P.-A., Koudelka-Hep, M. &
Berdondini, L. (2008). Large-Scale, High-Resolution Data Acquisition System for Extracellular
Recording of Electrophysiological Activity. IEEE Transactions on Biomedical Engineering, 55(8),
2064–2073. https://doi.org/10.1109/tbme.2008.919139
28. Insel, N. & Barnes, C. A. (2015). Differential Activation of Fast-Spiking and Regular-Firing Neuron
Populations During Movement and Reward in the Dorsal Medial Frontal Cortex. Cerebral Cortex,
25(9), 2631–2647. https://doi.org/10.1093/cercor/bhu062
29. Jia, X., Siegle, J. H., Bennett, C., Gale, S. D., Denman, D. J., Koch, C. & Olsen, S. R. (2019). High-
density extracellular probes reveal dendritic backpropagation and facilitate neuron
classification. Journal of Neurophysiology, 121(5), 1831–1847.
https://doi.org/10.1152/jn.00680.2018
30. Jun, J. J., Mitelut, C., Lai, C., Gratiy, S. L., Anastassiou, C. A. & Harris, T. D. (2017). Real-time spike
sorting platform for high-density extracellular probes with ground-truth validation and drift
correction. bioRxiv, 101030. https://doi.org/10.1101/101030
31. Kawaguchi, Y. & Kubota, Y. (1993). Correlation of physiological subgroupings of nonpyramidal
cells with parvalbumin- and calbindinD28k-immunoreactive neurons in layer V of rat frontal
cortex. Journal of Neurophysiology, 70(1), 387–396. https://doi.org/10.1152/jn.1993.70.1.387
32. Kohl, M. M., & Paulsen, O. (2010). The roles of GABAB receptors in cortical network
activity. Advances in pharmacology, 58, 205-229.
33. Lak, A., Okun, M., Moss, M. M., Gurnani, H., Farrell, K., Wells, M. J., Reddy, C. B., Kepecs, A.,
Harris, K. D. & Carandini, M. (2020). Dopaminergic and prefrontal basis of learning from sensory
confidence and reward value. Neuron, 105(4), 700-711.
34. Laquitaine, S., Imbeni, M., Tharayil, J., Isbister, J.B., Reimann, M.W. (2024). Spike sorting biases
and information loss in a detailed cortical model. BioRxiv.
https://doi.org/10.1101/2024.12.04.626805
35. Lee, J., Carlson, D., Shokri, H., Yao, W., Goetz, G., Hagen, E., Batty, E., Chichilnisky, E., Einevoll, G.
& Paninski, L. (2017). YASS: Yet Another Spike Sorter. bioRxiv, 151928.
https://doi.org/10.1101/151928
36. Lemon, R. N., Baker, S. N. & Kraskov, A. (2021). Classification of Cortical Neurons by Spike Shape
and the Identification of Pyramidal Neurons. Cerebral Cortex, 31(11), bhab147-.
https://doi.org/10.1093/cercor/bhab147
37. Lewicki, M. (1998). A review of methods for spike sorting: the detection and classification of
neural action potentials. Network: Computation in Neural Systems, 9(4), R53–R78.
https://doi.org/10.1088/0954-898x/9/4/001

38. Lindén, H., Tetzlaff, T., Potjans, T. C., Pettersen, K. H., Grün, S., Diesmann, M. & Einevoll, G. T.
(2011). Modeling the Spatial Reach of the LFP. Neuron, 72(5), 859–872.
https://doi.org/10.1016/j.neuron.2011.11.006
39. Liu, Y. H, Smith, S., Mihalas, S. & Sumbul, U. (2021). Cell-type–specific neuromodulation guides
synaptic credit assignment in a spiking neural network. PNAS, 118, e2111821118.
40. Madisen, L., Mao, T., Koch, H., Zhuo, J., Berenyi, A., Fujisawa, S., Hsu, Y.-W. A., Garcia, A. J., Gu,
X., Zanella, S., Kidney, J., Gu, H., Mao, Y., Hooks, B. M., Boyden, E. S., Buzsáki, G., Ramirez, J. M.,
Jones, A. R., Svoboda, K., … Zeng, H. (2012). A toolbox of Cre-dependent optogenetic transgenic
mice for light-induced activation and silencing. Nature Neuroscience, 15(5), 793–802.
https://doi.org/10.1038/nn.3078
41. Magland, J., Jun, J. J., Lovero, E., Morley, A. J., Hurwitz, C. L., Buccino, A. P., Garcia, S. & Barnett,
A. H. (2020). SpikeForest, reproducible web-facing ground-truth validation of automated neural
spike sorters. ELife, 9, e55167. https://doi.org/10.7554/elife.55167
42. Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., Wu, C. (2004).
Interneurons of the neocortical inhibitory system. Nat Rev Neurosci, 5, 793-807. https://doi.org/
10.1038/nrn1519
43. Marre, O., Amodei, D., Deshmukh, N., Sadeghi, K., Soo, F., Holy, T. E. & Berry, M. J. (2012).
Mapping a Complete Neural Population in the Retina. The Journal of Neuroscience, 32(43),
14859–14873. https://doi.org/10.1523/jneurosci.0723-12.2012
44. McCormick, D. A., Connors, B. W., Lighthall, J. W. & Prince, D. A. (1985). Comparative
electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. Journal of
Neurophysiology, 54(4), 782–806. https://doi.org/10.1152/jn.1985.54.4.782
45. Mechler, F., Victor, J. D., Ohiorhenuan, I., Schmid, A. M. & Hu, Q. (2011). Three-dimensional
localization of neurons in cortical tetrode recordings. Journal of Neurophysiology, 106(2), 828–
848. https://doi.org/10.1152/jn.00515.2010
46. Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. (2007). Differential Attention-Dependent
Response Modulation across Cell Classes in Macaque Visual Area V4. Neuron, 55(1), 131–141.
https://doi.org/10.1016/j.neuron.2007.06.018
47. Moore, A. K. & Wehr, M. (2013). Parvalbumin-Expressing Inhibitory Interneurons in Auditory
Cortex Are Well-Tuned for Frequency. The Journal of Neuroscience, 33(34), 13713–13723.
https://doi.org/10.1523/jneurosci.0663-13.2013
48. Muthmann, J.-O., Amin, H., Sernagor, E., Maccione, A., Panas, D., Berdondini, L., Bhalla, U. S. &
Hennig, M. H. (2015). Spike Detection for Large Neural Populations Using High Density
Multielectrode Arrays. Frontiers in Neuroinformatics, 9, 28.
https://doi.org/10.3389/fninf.2015.00028
49. Nowak, L. G., Azouz, R., Sanchez-Vives, M. V., Gray, C. M. & McCormick, D. A. (2003).
Electrophysiological Classes of Cat Primary Visual Cortical Neurons In Vivo as Revealed by
Quantitative Analyses. Journal of Neurophysiology, 89(3), 1541–1566.
https://doi.org/10.1152/jn.00580.2002
50. Obien, M. E. J., Deligkaris, K., Bullmann, T., Bakkum, D. J. & Frey, U. (2015). Revealing neuronal
function through microelectrode array recordings. Frontiers in Neuroscience, 8, 423.
https://doi.org/10.3389/fnins.2014.00423
51. Pachitariu, M., Sridhar, S. & Stringer, C. (2023). Solving the spike sorting problem with
Kilosort. bioRxiv, 2023.01.07.523036. https://doi.org/10.1101/2023.01.07.523036
52. Peters, A. J., Fabre, J. M., Steinmetz, N. A., Harris, K. D., & Carandini, M. (2021). Striatal activity
topographically reflects cortical activity. Nature, 591(7850), 420-425.

53. Pettersen, K. H. & Einevoll, G. T. (2008). Amplitude Variability and Extracellular Low-Pass Filtering of Neuronal Spikes. Biophysical Journal, 94(3), 784–802. https://doi.org/10.1529/biophysj.107.111179

54. Peyrache, A., Dehghani, N., Eskandar, E. N., Madsen, J. R., Anderson, W. S., Donoghue, J. A., Hochberg, L. R., Halgren, E., Cash, S. S. & Destexhe, A. (2012). Spatiotemporal dynamics of neocortical excitation and inhibition during human sleep. Proceedings of the National Academy of Sciences, 109(5), 1731–1736. https://doi.org/10.1073/pnas.1109895109

55. Peyrache, A. & Destexhe, A. (2019). Electrophysiological monitoring of inhibition in mammalian species, from rodents to humans. *Neurobiology of Disease*, *130*, 104500. https://doi.org/10.1016/j.nbd.2019.104500

56. Prentice, J. S., Homann, J., Simmons, K. D., Tkačik, G., Balasubramanian, V. & Nelson, P. C. (2011). Fast, Scalable, Bayesian Spike Identification for Multi-Electrode Arrays. PLoS ONE, 6(7), e19884. https://doi.org/10.1371/journal.pone.0019884

57. Ramsay, J. O. & Silverman, B. W. (1997). Functional Data Analysis, *Springer*, New York.

58. Ren, N., Ito, S., Hafizi, H., Beggs, J. M. & Stevenson, I. H. (2020). Model-based detection of putative synaptic connections from spike recordings with latency and type constraints. *Journal of Neurophysiology*, *124*(6), 1588–1604. https://doi.org/10.1152/jn.00066.2020

59. Rey, H. G., Pedreira, C. & Quiroga, R. Q. (2015). Past, present and future of spike sorting techniques. Brain Research Bulletin, 119(Pt B), 106–117. https://doi.org/10.1016/j.brainresbull.2015.04.007

60. Riquelme, J. L., Hemberger, M., Laurent, G. & Gjorgjieva, J. (2023) Single spikes drive sequential propagation and routing of activity in a cortical network *eLife* 12:e79928. https://doi.org/10.7554/eLife.79928.

61. Rossant, C., Kadir, S. N., Goodman, D. F. M., Schulman, J., Hunter, M. L. D., Saleem, A. B., Grosmark, A., Belluscio, M., Denfield, G. H., Ecker, A. S., Tolias, A. S., Solomon, S., Buzsaki, G., Carandini, M., Harris K. D. 2016. Spike sorting for large, dense electrode arrays. *Nat Neurosci* **19**:634–641. doi:10.1038/nn.4268

62. Ruz, I. D. & Schultz, S. R. (2014). Localising and classifying neurons from high density MEA recordings. *Journal of Neuroscience Methods*, *233*, 115–128. https://doi.org/10.1016/j.jneumeth.2014.05.037

63. Sakata, S. & Harris, K. D. (2009). Laminar Structure of Spontaneous and Sensory-Evoked Population Activity in Auditory Cortex. Neuron, 64(3), 404–418. https://doi.org/10.1016/j.neuron.2009.09.020

64. Shein-Idelson, M., Pammer, L., Hemberger, M. & Laurent, G. (2017). Large-scale mapping of cortical synaptic projections with extracellular electrode arrays. *Nature Methods*, *14*(9), 882–890. https://doi.org/10.1038/nmeth.4393

65. Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M. & Buzsáki, G. (2008). Entrainment of Neocortical Neurons and Gamma Oscillations by the Hippocampal Theta Rhythm. Neuron, 60(4), 683–697. https://doi.org/10.1016/j.neuron.2008.09.014

66. Stark, E., Eichler, R., Roux, L., Fujisawa, S., Rotstein, H. G. & Buzsáki, G. (2013). Inhibition-Induced Theta Resonance in Cortical Circuits. Neuron, 80(5), 1263–1276. https://doi.org/10.1016/j.neuron.2013.09.033

67. Stringer, C., Pachitariu, M., Steinmetz, N. A., Okun, M., Bartho, P., Harris, K. D., Sahani, M., Lesica, N. A. (2016). Inhibitory control of correlated intrinsic variability in cortical networks. *Elife*, *5*, e19695. https://doi.org/10.7554/eLife.19695

68. Sylwestrak, E. L., Jo, Y. J., Vesuna, S., Wang, X., Holcomb, B., Tien, R. H., Kim, D. K., Fenno, L.,
Ramakrishnan, C., Allen, W. E., Chen, R., Shenoy, K. V., Sussillo, D. & Deisseroth, K. (2022). Cell-
type-specific population dynamics of diverse reward computations. *Cell* 185, 3568–3587.
69. Tamura, H., Kaneko, H., Kawasaki, K. & Fujita, I. (2004). Presumed Inhibitory Neurons in the
Macaque Inferior Temporal Cortex: Visual Response Properties and Functional Interactions With
Adjacent Neurons. Journal of Neurophysiology, 91(6), 2782–2796.
https://doi.org/10.1152/jn.01267.2003
70. Thivierge, J. P., Giraud, E., Lynn, M., & Théberge Charbonneau, A. (2022). Key role of neuronal
diversity in structured reservoir computing. *Chaos: An Interdisciplinary Journal of Nonlinear*
Science, *32*(11), 113130.
71. Vendramin, L., Campello, R. J., & Hruschka, E. R. (2009, April). On the comparison of relative
clustering validity criteria. In Proceedings of the 2009 siam international conference on data
mining (pp. 733-744). Society for Industrial and Applied Mathematics.
72. Wilson, F. A., O'Scalaidhe, S. P. & Goldman-Rakic, P. S. (1994). Functional synergism between
putative gamma-aminobutyrate-containing neurons and pyramidal neurons in prefrontal cortex.
Proceedings of the National Academy of Sciences, 91(9), 4009–4013.
https://doi.org/10.1073/pnas.91.9.4009
73. Xu, R. & D. Wunsch. (2005). Survey of Clustering Algorithms. IEEE Transactions on Neural
Networks, 16(3), 645–678. https://doi.org/10.1109/tnn.2005.845141
74. Yael, D. & Bar-Gad, I. (2017). Filter based phase distortions in extracellular spikes. PLOS ONE,
12(3), e0174790. https://doi.org/10.1371/journal.pone.0174790
75. Yger, P., Spampinato, G. L., Esposito, E., Lefebvre, B., Deny, S., Gardella, C., Stimberg, M., Jetter,
F., Zeck, G., Picaud, S., Duebel, J. & Marre, O. (2018). A spike sorting toolbox for up to thousands
of electrodes validated with ground truth recordings in vitro and in vivo. *eLife*, *7*, e34518.
https://doi.org/10.7554/elife.34518

Tables

	lang.	# elect.	data preproc.	event/ channel loc.	waveform extrac.	e/i sorting	valida tion	experim. prot.	unsu per.	source code available
spikeMAP	Python and Matlab	4096	+	+	+	+	+	+	+	+
Kilosort4 (Pachitariu et al., 2023) github.com/cortex-lab/KiloSort	Python		+	+	-	-	-	-	-	+
SpyKING CIRCUS (Yger et al., 2018) spyking-circus.rtfd.org	Python	4225	-	+	+	-	(+)	-	+	+
MountainSort (Chung et al., 2017) github.com/flatironinstitute/mountainsort	Matlab and C++	16	+	+	-	-	-	-	+	+
Herding Spikes (Hilgen et al., 2017) github.com/mhhennig/HS2	Python	4096	+	+	+	-	(+)	+	+	+
YASS (Lee et al., 2017) yass.readthedocs.io	Python	512	-	+	-	-	-	-	+	+
JRCLUST (Jun et al., 2017) jrclust.org	Matlab and CUDA	120 (1000)	+	+	-	-	(+)	-	-	+

Table 1.

lang.: Programming language, **# elect.**: Maximum number of electrodes tested on, **data preproc.**: Data
preprocessing (formatting, filtering, artefact removal) included, **event/channel loc.**: Localization of
single events, **waveform extrac.**: Spike features (such as peak to peak distance and full-width half max)
are extracted from waveforms, **e/i sorting**: excitatory and inhibitory cell sorting, **validation**: validation of
cell-type sorting *(+) = ground truth validation via intracellular recordings, but not e/i validation*, **experim.**
prot.: integrates an experimental protocol to allow for network manipulation, **unsuper.**: unsupervised,
requiring no manual user-intervention, **source code available**: entire code is open-source and available.

Figures

Figure 1. Workflow diagram of the proposed method. (**A**) Filtering of the data is achieved through
bandpass filtering (300 to 5000 Hz). Spike detection is done on filtered voltages, however raw voltages
around each spike to avoid signal distortion. (**B**) Spike sorting on each electrode. (**Bi**) Events are detected
using a threshold method (6 SD from the mean of the filtered signal). (**Bii**) PCA and k-means on
individual waveforms (k = 2) (**C**) Precise soma location on the array. (**D**) Excitatory and inhibitory sorting.
(**Di**) Spike time constants kinetics are extract at the full width at half maximum and peak to peak
distances. (**Dii**) Interpolated voltage for increased resolution (**Diii**) k-means to distinguish cell-type based
on their waveform kinetics.

Figure 2. Isolation of single-cell action potential for each electrode. (**A**) Multielectrode array recording
set up (**B**) Extracellular voltage on a single electrode (**Bi**) Spike detection on filtered voltage (**Bii**) Raw
voltage extraction around detected spike (**C**) Firing rate per electrode depends on threshold selection.
Inset shows firing rate values at the selected threshold (4 SD) **D**) Threshold selection to part cellular
activity from normally distributed noise (**E**) Spike sorting on a single electrode (**F**) Current sink signal
from single neuron spans multiple electrodes (4 different neurons are shown in different colors) (**G**)
Histogram distribution of ten-fold cross-validation with chance performance indicated in red for all
electrodes in a single recording. (**H**) Voltage amplitude as a function of distance from highest electrode
peak, inset shows the directions measured from the center electrode (**I**) Center of mass location of all
spikes for a single soma, mean is indicated in red.

Figure 3. Identification of putative excitatory and inhibitory units. (**A**) Distinct waveform kinetics for E
and I cells (**B**) Respective cell location (**C**) Raster plot of E and I activity. (**D**) Fano factor for putative E and
I cells. The shaded area represents the standard error of the mean (SEM).The x-axis shows temporal bins
of increasing durations employed to compute the fano factor (**E**) Mean cross-correlation as a function of
spatial distance between pairs of electrodes. Non-overlapping bins of 100 ms were employed to

743   compute correlations. The resulting values were averaged by spatial distance between all pairs of

744   neurons.

Figure 4. Validation of step-function opsin by whole-cell recordings.(**A**) Visually identified PV neuron (**B**) Nearby pyramidal neuron receiving inhibitory currents following PV opto activation (**C**) IPSCs onto pyramidal cell with 20 s rest between trial (**D**) IPSCs onto pyramidal cell with 60 s rest between trial (**E**) IPSCs onto pyramidal cell with 120 s rest between trial

Figure 5. Optogenetic targeting of inhibitory cells for spike sorting (**A**) Average firing rate for putative excitatory and inhibitory neurons. The shaded area represents the SEM. (**B**) Quantification of change in firing rate following optogenetic stimulation. Average firing rates are taken over four recordings obtained from 3 mice. (**C**) Delta change in firing rate (on – drugs) (**D**) Opto-evoked change in firing rate

Supplemental Figures

Supplemental Figure 1: (A) Voltage amplitude as a function of distance from highest electrode peak.
Putative somas with an increase in voltage away from electrode peak were excluded from this analysis.
(B-C) Distribution of slopes and intercepts across all putative somas obtained from one recording
(*N*=1,950).

Supplemental Figure 2: (A) Distribution of signal-to-noise ratio (SNR) over all spike-sorted clusters of a
single recording (*N*=8,059). **(B)** Examples of mean voltages (black lines) and individual voltages (grey
lines) around individual spikes.

Supplemental Figure 3: Distribution of mean firing rates across a population of putative excitatory (**A**)
(*N*=570) and inhibitory (**B**) (*N*=54) neurons. (**C**) Mean firing rate across neurons. Vertical bars: SEM.

Supplemental Figure 4: Light artefact in a recording from mouse prefrontal cortex showing that the
application of a brief (1 ms) pulse of light leads to saturation lasting about 5 ms post-stimulation.

Supplemental Figure 5: Waveform kinetics for E and I cells obtained during the baseline window of the optogenetic protocol. Insets show cartoons of individual spikes with broader (E) and narrower (I) waveforms.

File name	Animal #	Protocol #	# of E cells identified	# of I cells identified	total recording time (min)
R20211221_Slice1_Vars3.mat	1	1	3687	47	8
R20211213_Slice2_Vars3.mat	2	1	2943	350	8
R20211209_Slice3_Vars3.mat	3	1	1994	712	8
R20211219_Slice3_Vars3.mat	4	2	1138	32	8

Supplemental Table 1: Summary of SpikeMAP open-source data. Datasets were obtained from four mice. The optogenetic stimulation protocol (see Materials and Methods) was run once per dataset. Data is available from: dx.doi.org/10.6084/m9.figshare.29416472

A

EYFP

Alexa 594

B

3 IPSCs/s

20 pA

20 s

10 pA
500 ms

C

20 s rest

Trial #

2 IPSCs/s
10 s

D

60 s rest

Trial #

5 IPSCs/s
20 s

E

120 s rest

Trial #

3 IPSCs/s
20 s

A

norm. count

0.05

0   10   20

SNR

B

SNR=9.79   *SNR*=0.46   *SNR*=11.36

25 µV

1 ms