Structural variants contribute to the gut microbiome, blood metabolic traits, and their causal relationships

Qun Zhou[1,†], Keqing Wang[2,†], Junhong Chen[1,†], Xinjiang Tan[2], Xun Xu[2], Xin Jin[2], Shaoqiao Zhang[1], Liang Xiao[2,4], Tao Zhang[1,5*], Xiaomin Liu[1,5*]

1. BGI Research, Wuhan 430074, China;
2. BGI Research, Shenzhen 518083, China;
3. James D. Watson Institute of Genome Sciences, Hangzhou 310058, China;
4. Shenzhen Engineering Laboratory of Detection and Intervention of Human Intestinal Microbiome, BGI-Shenzhen, Shenzhen 518083, China;
5. Shenzhen Key Laboratory of Human Commensal Microorganisms and Health Research, BGI Research, Shenzhen 518083, China

[†] These authors contributed equally.
* To whom correspondence should be addressed: X.L., liuxiaomin@genomics.cn   and T.Z., tao.zhang@genomics.cn

Highlights

• Provides a catalog of 138,859 high-confidence SVs and 49.57% of which were novel

• Identify 30 genome-wide significant "SVs—microbial taxa/pathways" and 48 "SVs—blood metabolites" associations, respectively

• An Asian-specific 6.5k deletion at *TENT2* gene associated with the species *Coprobacillus sp. 29_1*

• A deletion spanning *HBA1/HBA2/HBM/HBQ118* associated with 9 blood metabolic traits

• The majority of SVs demonstrated a more pronounced impact than neighboring SNPs in the associations with microbiome/metabolites

• SVs help to reveal the host-microbe causal relationships such as *Clostridium ramosum's* causal effects on serum uric acid and hyperuricemia

Abstract

Research on genetic variants affecting the gut microbiome and blood metabolites has mainly

focused on single nucleotide variants (SNPs), with structural variants (SVs) remaining largely

unexamined. In this study, we utilized an advanced structural variant detection pipeline on

whole-genome sequencing data from 2,002 individuals with an average sequencing depth of

42 folds, identifying 138,859 high-confidence SVs, 49.57% of which were previously

unreported. We performed genome-wide association studies on 22,519 common SVs with 616

gut microbial features and 121 blood metabolic traits, uncovering 30 significant links between

SVs and microbial taxa/pathways and 48 connections between SVs and blood metabolites.

Notable results include 19 associations reaching study-wide significance, such as an

Asian-specific 6.5k deletion at the *TENT2* gene, which is strongly associated with the

abundance of *Coprobacillus sp. 29_1* (β=0.158, p=2.5$\times$10^{-9}), and a deletion covering

HBA1/HBA2/HBM/HBQ118 linked to nine blood metabolic traits, most significantly with

mean corpuscular volume (MCV; β=-0.570, p=1.22$\times$10^{-91}). Most SVs showed a greater

impact than nearby SNPs in their associations with the microbiome/metabolites. Analysis of

rare SVs also identified multiple genes associated with the microbiome/metabolites.

Additionally, Mendelian randomization using SVs demonstrated causal relationships,

including effects from the microbiome to metabolites, such as *Clostridium ramosum*'s impact

on serum uric acid and hyperuricemia, and from metabolites to the microbiome, like the

influence of mean corpuscular hemoglobin concentration on MF0064: triacylglycerol

degradation. This study provides a detailed SV catalog for the Chinese population and

highlights the significant role of SVs in the gut microbiome, blood metabolites, and their

causal relationships.

Keywords: structural variants, genome-wide association analysis, gut microbiome, blood

metabolites, host-microbe causal relationships

Introduction

The human gut microbiome plays a pivotal role in numerous physiological processes and is closely linked to human health and the development of diseases such as obesity[1,2], diabetes[3,4], inflammatory bowel disease[5,6], rheumatoid arthritis[7,8], cardiovascular diseases[9], and mental disorders[10,11]. Apart from being influenced by factors such as environment, diet, lifestyle, and diseases, the gut microbiome is also controlled to some extent by host genetic factors[12,13]. In recent years, many microbial genome-wide association studies (mbGWAS) have identified various host genetic variants that influence the gut microbiome[14-18], with only the *LCT* and *ABO* loci being frequently discussed and well replicated.

Despite the progress made in understanding the relationship between host genetics and the gut microbiome, most mbGWAS studies have primarily focused on single nucleotide polymorphisms (SNPs) using genotyping arrays, except for the human microbiome project (HMP) and our prior research[19,20]. This approach has overlooked the potential impact of more extensive and complex genomic variations, such as structural variations (SVs). SVs, which constitute a significant form of genomic variation, including alterations exceeding 50 base pairs (bp) in length such as deletions (DEL), insertions (INS), inversions (INV), duplications (DUP), and copy number variants (CNV). These SVs are prevalent in the human genome and have been associated with various disease phenotypes[21-24]. A recent study identified a correlation between CNV of the human salivary amylase gene (*AMY1*) and the relative abundance of the genus *Ruminococcus* in the gut microbiome[25]. However, this study solely relied on PCR technology to explore the impact of a single human *AMY1* gene copy number variation on the microbiome. Given that few studies had both whole genome and whole metagenome data in the same cohort, thus to date, the impact of host SVs on the gut microbiome remains largely unexplored.

In addition to the gut microbiome, the genome-wide contributions of host SVs to blood metabolic traits have been underexplored. Very limited evidence has been reported. For example, Chen et al. identified a deletion of the *ALB* promoter and a CNV at *PDPR* that were significantly associated with blood metabolic traits in the Finish population[26]. The NHLBI TOPMed program has reported 21 independent SVs or 41 SV-trait associations across a total of 24 quantitative hematologic traits[27]. While these studies have identified associations between SVs and metabolic traits in Finnish and European populations, there is a notable gap in research focusing on Asian populations. Recent efforts to address this gap include a study using long-read sequencing in a Chinese population, which identified 22 SVs on 14 chromosomes associated with 13 phenotypes[28]. However, the association analysis was conducted on a maximum of 327 individuals. This underscores the need for a comprehensive investigation of SV contributions to blood metabolic traits in larger Asian populations. Moreover, comprehensively and systematically excavating the impact of host SVs on both the microbiome and metabolites using high-depth whole-genome data from a well-designed

102 cohort will enhance our understanding of host-microbiome interaction mechanisms.

104 To address the research gap and investigate the role of SVs in the gut microbiome and blood

105 metabolic traits, we constructed and characterized a high-quality SV profile of the Chinese

106 population based on 2,002 individuals from the 4D-SZ cohort[14,19,20,29-31], a multi-omics cohort

107 comprising high-depth whole genome and metagenomics sequencing, blood metabolites, and

108 detailed questionnaires. Subsequently, we conducted genome-wide association analyses to

109 identify common SVs and genes harboring rare SVs that significantly contributed to gut

110 microbial taxa/pathways and blood metabolic traits, respectively. Moreover, we examined the

111 intricate relationship between gut microbiome and blood metabolites through their associated

112 SVs as mediators (see **Graphic abstract**). The primary objective of this study was to

113 characterize the landscape of SVs in the Chinese population and identify novel SVs

114 associated with the microbiome and metabolites, while offering valuable insights into the

115 mechanisms governing host-microbiome interactions.

117 **Results**

118 **Characterization of SVs identified in the Chinese population**

119 We utilized high-depth whole-genome sequencing (WGS) data to identify SVs in 2,002

120 Chinese individuals from the 4D-SZ cohort. Among the sequenced samples, 53% were

121 females, with an average age of 30 years ranging from 21 to 67 years (**Supplementary Fig.**

122 **1**). The mean sequencing depth per individual was 42×, amounting to an average of 122.46

123 Gb of sequenced data (**Supplementary Fig. 2a**). The base mapping rate varied from 96.74%

124 to 100%, with an average of up to 99.89% (**Supplementary Fig. 2b**). An advanced Genome

125 Analysis Toolkit's Structural Variant (GATK-SV) pipeline[32,33] was employed to detect SVs

126 (**Supplementary Fig. 3**; Methods). Following rigorous quality control measures for both

127 samples and variants, a total of 1,839 samples were retained. We detected a total of 138,859

128 high-confidence SVs with quality scores surpassing 250 or marked as "PASS," while

129 excluding "BND" variant types (**Supplementary Table 1**). These SVs comprised 73,288

130 insertions (INSs), 38,782 deletions (DELs), 25,030 duplications (DUPs), 1,153 complex SVs

131 (CPXs), 378 inversions (INVs), and 228 copy number variations (CNVs) (**Fig. 1a**). The most

132 prevalent SV type was INSs, accounting for 52.78% of all SVs, followed by DELs,

133 representing 27.93% (**Fig. 1a**).

135 Despite the overall higher count of INSs than DELs in the population, each individual

136 exhibited a greater number of DELs than INSs, consistent with an earlier study[34]. On average,

137 each genome harbored 11,493 SVs events (ranging from 7,851 to 21,324), exceeding the SV

138 count documented in the 1000 genome phase3 (9,679 SVs; average depth of 30×) and the

139 Genome Aggregation Database (gnomAD[32]; 7,439 SVs; average depth of 32×), yet lower than

140 a long-read sequencing study[28] (18,489 SVs; average depth of 17×). DELs and INSs

141 continued to be the most prevalent SV types, constituting 41.5% and 35.5% of all identified

SVs, respectively. The average numbers of DELs, INSs, DUPs, CNVs, CPXs, and INVs per
sample were 4,769, 4,075, 2,177, 237, 174, and 62, respectively (**Fig. 1b**).

To evaluate the identified SVs, we applied the same sequencing and analysis methods to the
cell line obtained from reference sample NA12878 (**Methods**). Through the application of the
GATK-SV pipeline, 10,624 SVs were detected. A comparison of the detected SVs with the
gold standard truth set of NA12878[35] revealed a precision of 79.5% for deletions, 64.2% for
duplications, 69.4% for insertions, and 75% for inversions. Additionally, the impact of
sequencing depth on SV detection was assessed, as shown in the **Supplementary Fig. 4**.
Overall, a higher sequencing depth was correlated with an increased number of SVs, with a
gradual rise in SV counts from 20× to 50×**.** However, this trend declined beyond 50×,
potentially owing to the limited number of samples used.

Across all population-detected SVs, 71.55% were rare, with minor allele frequencies (MAF)
below 1%. Most SVs exhibited lengths distributed within the range of 50~100bp (61.02%)
and 100-500bp (15.77%; **Fig. 1c**). The expected peaks corresponding to the insertion of Alu,
SVA, and LINE1 mobile elements were identified at approximately 300 bp, 2 kb, and 6 kb,
respectively (**Fig. 1d**). These SV patterns were consistent with those reported in previous
studies.

Regarding the functional prediction of SVs, a significant proportion of SVs were located
in the intergenic (57.32%) or intronic regions (36.62%; **Fig. 1e**). Only 6.06% of SVs
breakpoints intersected with functional elements, including exons (n = 3381), predicted
loss-of-function (pLoF; n = 1895), copy gain of entire genes (CG; n=1170), intragenic exonic
duplication (IED; n=1308), and promoters (n = 655). On average, each genome harbored a
mean of 173.96 SVs that altered the functions of 213 genes (121.37 pLoF affected 122.7
genes, 26.09 CG affected 65.4 genes, and 25.69 IED affected 25.1 genes; **Fig. 1f**), indicating
that approximately 1.1% of human protein-coding genes per individual could experience
functional abnormalities due to SVs.

To ascertain whether the SVs identified in this study had been previously documented, we
compared them with five publicly available SV databases: gnomAD[32], Audano et al. 2019[36],
the Human Genome Structural Variation Consortium(HGSVC2)[37], Wu et al.[28], and dbVar[38]
(accessed on September 30, 2023). Initially, we ensured that the SV coordinates in these
databases were converted to GRCh38 to align them with our datasets. An SV in our study was
considered previously reported if the overlap with SVs from published datasets was $\geq$ 50%.
Finally, our dataset exhibited the greatest number of overlapping SVs (48,027, overlapping
rate 34.58%; **Fig. 1g**) with the dataset constructed by Wu et al., which was based on long-read
WGS results from 405 Chinese individuals. The shared ancestry of all Chinese individuals
likely contributed to this substantial overlap. Furthermore, a significant number of
overlapping SVs were also detected in the gnomAD (43,755; 31.51%) and dbVar (40,206;

28.95%) datasets, two commonly used reference datasets known for their diverse ethnic
population representation. In contrast, the overlaps with the HGSVC2 (25,643; 18.47%) and
Audano et al. datasets (19,231; 13.85%) were relatively low. Overall, 49.57%
(68,830/138,859) of the SVs discovered in our study were novel and had not been previously
documented (**Fig. 1g**). These novel SVs were characterized by a higher fraction of insertions
(p < 2.2×10^{-16}, fisher's exact test), particularly in intergenic and intronic regions (both p <
2.2×10^{-16}, fisher's exact test; **Fig. 1h**). Additionally, they showed a lower frequency of overlap
with functional elements, affecting an average of 20.61 SV-altered genes per genome,
including 6.95 pLoF, 2.24 CG, and 11.41 IED (**Fig. 1i**).
192
Associations of SVs count and common SVs with the gut microbiome
After quality control during SV calling, 1,406 of the 1,839 samples had available fecal
shotgun metagenomic sequencing data. We first assessed the impact of an individual's SVs
count on gut microbial diversity through SV burden analysis using a generalized linear
regression model adjusted for age, gender, and the first 10 principal components. A weak but
statistically significant positive correlation was observed between the SV burden and
α-diversity indices (Shannon index: $\beta = 0.08$, $p = 0.011$; Simpson index: $\beta = 0.06$, $p = 0.012$;
Supplementary Fig. 5). We then calculated the associations of 22,519 common SVs (MAF ≥
1%) with α-diversity indices and identified no significant associations after multiple test
correction ($p < 2.22 \times 10^{-6} = 0.05/22,519$), consistent with previous conclusions in the SNP
GWAS for α-diversity. The most significant SV was an insertion in the genetic region of
chr19:3502006-3502028 associated with the Shanno index ($p= 7.55 \times 10^{-5}$) and Simpson
index ($p=4.02 \times 10^{-6}$; **Supplementary Fig. 6**). This insertion is near the Fizzy-related protein
homolog 1 (*FZR1*) gene, which mainly regulates the cell cycle in association with the
anaphase-promoting complex/cyclosome (APC/C), and plays an important role in
neurodevelopment[39] and tumorigenesis[40].
209
Next, a microbial genome-wide association study (mbGWAS) was conducted on 22,519
common SVs and 616 microbial features comprising 517 gut microbial taxa and 99 gut
metabolic modules (GMMs) present in over 10% of the individuals (**Supplementary Table 2**).
A total of 30 SVs (DEL = 16, INS = 7, DUP = 6, INV=1) were identified to be associated
with gut microbial features with genome-wide significance ($p < 2.22 \times 10^{-6}$; **Fig. 2a;
Supplementary Table 3**). The genomic inflation factor (λ) ranged from 0.84 to 1.33, with a
median value of 0.98 (**Fig. 2b**), indicating no obvious evidence of inflation, except for some
specific mbGWAS tests. Only one association achieved the study-wide significance level ($p <
3.60 \times 10^{-9}$, adjusting for 22,519 SVs and 616 microbial features).
219
The most significant association was observed for a 6.5kb deletion (chr5:79657532-79664046)
within the intron region of the gene *TENT2*, which was positively correlated with the relative
abundance of the species *Coprobacillus sp. 29_1* ($\beta = 0.158$, $p = 2.50 \times 10^{-9}$; **Fig. 2c**).

Carriers of this SV exhibited significantly higher mean relative abundances of *Coprobacillus*
sp. 29_1 (homozygous DEL: 0.15%; heterozygous DEL: 0.024%) than the non-carriers
(0.015%; **Fig. 2d**). This GWAS test exhibited no inflation, with a λ value of 0.87. The
identified SV had an MAF of 0.014 in this cohort and was specific to East Asian (MAF=0.028
for ASN; MAF=0 for other populations) according to the 1000 Genomes database[41]. Further
analysis revealed that this SV was correlated with concentrations of glycine ($p = 0.015$),
ethanol ($p = 0.022$), monocytes ($p = 0.027$), and albumin ($p = 0.028$) by investigating its
associations with blood parameters in this cohort (**Supplementary Fig. 7**). Interestingly,
Coprobacillus sp. 29_1 associated with SV also showed a correlation with monocyte
concentration, suggesting potential interactions among the three. By calculating the pairwise
linkage disequilibrium (LD) between this SV and its neighboring SNPs, we found that this SV
was in a strong LD with the SNP chr5:79674158_C/T ($r^2 = 0.973$; **Fig. 2c**), which also resided
in the intronic region of *TENT2* and showed a significant association with *Coprobacillus sp.*
29_1 ($\beta = 0.143$, $p = 2.18 \times 10^{-7}$). Consistent with SV, this SNP had a MAF of 0.013 in this
cohort and 0.027 in the ASN population, but was absent in other populations from the 1000
Genomes database. This SNP was identified as a susceptibility mutation for schizophrenia[42],
and *Coprobacillus sp. 29_1* has been consistently reported to be enriched in schizophrenia
patients[11]. Due to the strong LD between the SV and the SNP, the association with
Coprobacillus sp. 29_1 became insignificant after conditioning on either of them. Both the
SV and SNP were located in the intronic region of gene *TENT2*, which is well known to
regulate miRNA stability and abundance via diverse mechanisms[43]. The existing literature
indicated the significant role of host miRNAs in modulating the gut microbiota[44]. We
hypothesized that SV or SNP may affect the gene expression of *TENT2*, potentially regulating
miRNA to influence the gut microbial abundance.
247
In addition to the study-wide significant signal, 22 SVs were associated with microbial taxa at
genome-wide significance (**Supplementary Table 3**). Among them, three SVs lied in the
genes that were extremely intolerant to pLoF variants with pLI ≥ 0.99, as annotated by the
AnnotSV tool. Specifically, a 61 bp duplication within the *CDH8* gene was negatively
associated with the abundance of *Streptococcus sanguinis* ($\beta = -0.137$, $p = 3.37 \times 10^{-7}$), a
predominant bacterium in the human oral cavity known to be linked to the development of
infective endocarditis[45]. Similarly, a 108 bp duplication in the *ANKRD11* gene was negatively
associated with the abundance of *Turicibacter* ($\beta = -0.133$, $p = 4.91 \times 10^{-7}$). The SV was
overlapped with pathogenic loss of *ANKRD11*, which caused KBG syndrome, a disease
characterized by intellectual disability, skeletal malformations, and macrodontia[46].
Turicibacter has been reported to be a highly heritable taxon in both human and mouse
genetic studies[12,47,48], supporting the association between SV and *Turicibacter* abundance
found in this study. Conversely, a 751 bp deletion in the *GRID2* gene was positively
associated with the presence of *Eubacterium cellulosolvens* ($\beta = 0.755$, $p = 6.85 \times 10^{-7}$), an
anaerobic cellulolytic bacterium in the gut. The *GRID2* gene encodes the glutamate receptor
channel delta-2 subunit, and its deletion leads to a recessive syndrome of cerebellar ataxia and

tonic upgaze in humans[49].

In addition to the three SVs within disease-associated genes, several SVs were found to
exhibit significant effects on potential probiotics that could contribute to host health. For
example, a duplication event in the *LOC107985978* gene on chromosome 7 (SV length: 68 bp)
was positively associated with the abundance of *Roseburia intestinalis* (β = 0.139, p = 2.20 $\times$
10^{-7}), a promising probiotic known for its anti-inflammatory properties by producing
butyrate[50,51]. An insertion on chromosome 15 involving an ALU element (SV length: 262 bp)
demonstrated a significantly negative correlation with the emerging probiotic
Parabacteroides distasonis[52] (β = -0.137, p = 2.51 $\times$ 10^{-7}).

In addition to microbial taxa, seven associations were identified for the GMMs, also known as
microbial pathways. For instance, the glyoxylate bypass pathway (MF0063) was negatively
associated with an insertion on chromosome 2 (SV length: 111 bp, β = -0.147, p = 2.92 $\times$ 10^{-8}).
The pyruvate dehydrogenase complex (MF0072) exhibited a negative association with a
deletion on chromosome 13 (SV length: 234 bp, β = -0.132, p = 3.26 $\times$ 10^{-7}). The pectin
degradation II pathway (MF0004) exhibited a negative association with a deletion on
chromosome 9 (SV length: 82 bp, β = -0.131, p = 5.19 $\times$ 10^{-7}). Notably, all the SVs associated
with the microbiome were located in the intronic and intergenic noncoding regions, except for
an inversion of 4.78Mb spanning 14 genes, consistent with previous findings in the
SNP-based mbGWAS[53].

Conditional analyses and functional annotations of microbiome-associated SVs
To further investigate the independence of the identified microbiome-associated SVs from
nearby SNPs in influencing changes in microbial features, pairwise LD between SVs and
their neighboring SNPs within the ±500kbp flanking region of SVs were calculated. We found
that eight of the 30 microbiome-associated SVs exhibited at least moderate LD ($r^2 \geq 0.5$) with
one SNP or indel associated with the same microbial feature (**Fig. 3a**; **Supplementary Table
3-4**). Subsequently, conditional analyses were performed for each SV, adjusting for the
tagging SNP that showed the strongest LD with SV. These analyses confirmed that eight
SV-microbiome associations were no longer significant after adjusting for SNP/indel at the
same locus (p > 0.01; **Fig. 3b**), whereas the remaining 22 SVs maintained their significance.
Notably, 11 of these 22 SVs retained genome-wide significance following conditional
analyses (p < 2.2$\times$$10^{-6}$). These independent SV signals included two SVs that were potentially
linked to the probiotics—*Roseburia intestinalis* and *Parabacteroides distasonis*. These
findings indicate that the majority of SVs (73%), rather than nearby SNPs, were the primary
causal agents driving changes in microbial abundance.

AnnotSV[54-56] was employed to annotate the 30 microbiome-associated SVs, resulting in the
identification of 24 genes that harbored these SVs. Subsequently, gene enrichment analysis

was conducted for these 24 genes. Gene Ontology (GO) enrichment analysis revealed 10
significantly enriched GO terms (p.adjust < 0.05), including one biological process (BP), six
cellular components (CC), and one molecular function (MF) (**Supplementary Fig. 8a**;
Supplementary Table 5). The enriched gene functions were predominantly related to the
structure and function of neuronal synapses, encompassing the synaptic membrane, synaptic
cleft, sodium-independent organic anion transport, sodium-independent organic anion
transmembrane transporter activity, postsynaptic specialization membrane, postsynaptic
density membrane, glutamatergic synapse, and excitatory synapse. Additionally, Kyoto
Encyclopedia of Genes and Genomes (KEGG) enrichment analysis indicated significant
enrichment of these genes in four pathways: nicotinate and nicotinamide metabolism,
long-term depression, the peroxisome, and the phosphatidylinositol signaling system
(**Supplementary Fig. 8b**; **Supplementary Table 6**). These findings suggested a pivotal role
of microbiome-associated SVs in neurotransmission, ion transport, and cell signaling
pathways.
318

Associations of rare SVs and the gut microbiome
To further explore the relationship between rare SVs (MAF < 0.01) and the gut microbiome,
we performed a gene-based sequence kernel association test (SKAT) analysis of overall

diversity and microbial abundances. With a p-value threshold of 8×10^{-6} by adjusting for the

detected 5861 genes, we did not observe a significant association between overall diversity
and rare SV burden. However, 56 gene loci were associated with microbial features

(**Supplementary Table 7**). Two associations reached the study-wide significance ($p<1.38\times$

10^{-9} by adjusting for 5861 genes and 616 microbial features). The strongest signal was
observed for the Janus kinase and microtubule interacting protein 2 (*JAKMIP2*) gene
(harboring four rare SVs) associated with the abundance of MF0103: mucin degradation
($p=8.28\times10^{-10}$). The *JAKMIP2* gene has been reported as a potential risk factor for Graves'
disease in the Chinese Han population[57], and its nearby long noncoding RNA, *JAKMIP2-AS1*,
promoted the growth of colorectal cancer and indicated poor prognosis[58]. Accordingly, mucin
degradation has been linked to intestinal inflammation and tumor incidence[59]. The second
strongest signal lied in the gene *SLC9A3* (encoding NHE3: sodium-hydrogen
antiporter/exchanger 3; harboring seven rare SVs) associated with the abundance of MF0021:
xylose degradation ($p=1.28\times10^{-9}$). This gene had altered expression in the colonic mucosa of
adult IBD patients[60], showed strong epigenetic interaction with microbiome[61],
and *SLC9A3* knockout mice developed colitis[62].
338

In addition to these pathways, some genes were associated with specific microbial taxa. For
example, several genes, including *INTS9*, *HLA-DQA1*, and *MTMR3*, were associated with the
species *C. intestinalis* from the genus *Collinsella*, which was causally linked to serum
cholesterol level[63] and liver-related disease[64]. The potassium voltage-gated channel subfamily
B member 1 (*KCNB1*) SV burden was correlated with the abundance of *Streptococcus*

salivarius (p=3.88×10^{-7}), *S. vestibularis* (*p*=6.06×10^{-7}) and *S. sp.C150* (p=4.23×10^{-6})
(**Supplementary Table 7**). A mouse study demonstrated that *S. salivarius* modulates sucrose
metabolism via exopolysaccharide (EPS)-dependent short-chain fatty acid production[65], while
KCNB1 variants impaired pancreatic β-cell function and increased the risk of type 2 diabetes
(T2D)[66]. This dual regulation—microbial EPS-SCFA axis and host ion channel
activity—suggests a synergistic mechanism whereby *KCNB1* SVs exacerbate T2D risk by
disrupting microbial carbohydrate processing and pancreatic glucose sensing. These "host
genes-gut microbiome" links need to be further investigated in the future.

Associations of SVs with blood metabolites

Next, we conducted a GWAS for 22,519 common SVs and 121 blood metabolic traits. A total
of 48 associations involving 33 SVs (DEL = 11, INS = 14, DUP = 8) reached genome-wide
significance ($p < 2.2×10^{-6} = 0.05/22519$; **Fig. 4a**; **Supplementary Table 8**). Among these, 17
associations reached study-wide significance ($p < 1.835×10^{-8}$, adjusting for 22,519 SVs and
121 blood metabolite traits). The genomic inflation factor λ was calculated, with a median
value of 1 (ranging from 0.80 to 1.33; **Supplementary Fig. 9**).

The strongest association signal was observed for a deletion on chromosome 16
(chr16:165396-184700; SV length: 19,304 bp) spanning the *HBA1*, *HBA2*, *HBM*, and *HBQ1*
genes, known as the widely reported Southeast Asian deletion (--^{SEA}), which is predominantly
distributed and associated with anemia in southern China and Southeast Asia[67,68]. In the
Chinese population in this study, the MAF of the SEA deletion was 0.01. Consistently, the
deletion was mainly detected in East Asian (ASN, MAF=0.019) and South Asian (SAN,
MAF=0.013) populations in the 1000 Genomes dataset. The deletion showed remarkably
significant associations with nine blood metabolic traits, particularly with mean corpuscular
volume (MCV, β = -0.570, $p = 1.22×10^{-91}$), mean corpuscular hemoglobin (MCH, β = -0.522,
$p = 4.19×10^{-76}$), mean corpuscular hemoglobin concentration (MCHC, β = -0.319, $p =$
$4.31×10^{-29}$), and red blood cell distribution (RBC, β = -0.198, $p = 2.43×10^{-11}$; **Fig. 4**;
Supplementary Table 8). Beyond its established correlations with blood cell parameters, the
deletion was also associated with blood microelement levels. Carriers of the deletion
exhibited elevated zinc and copper concentrations, whereas magnesium, iron, and manganese
levels were decreased compared to non-carriers (**Fig. 4b**). In addition to the
16:165396-184700 deletion, a 114 bp duplication on chromosome 9
(chr9:97255555-97255669) within the *SUGT1P4-STRA6LP-CCDC180* gene was also
associated with MCV (β= -0.174, $p = 7.89×10^{-9}$) and MCH (β= -0.174, $p = 7.89×10^{-9}$) levels.
This DUP had an MAF of 0.0125 in this study, which has not been previously reported.

Among the total 48 genome-wide associations, cystine, previously reported as the most
heritable metabolite in Chinese[69], exhibited the highest number of associated SVs, with 16
SVs linked to it (**Fig. 5a**). Three SVs reached study-wide significance, including a 9.2kb
deletion on chromosome 5 (5:46048650-46057870; β= -0.163, $p = 6.47×10^{-10}$) and two

insertions on chromosome 4 (β= -0.161, p =2.83×10^{-9}) and 12 (β= -0.161, p = 2.83×10^{-9}),

respectively (**Supplementary Table 8**).

The results of GWAS and LD analysis revealed that three out of the 33 SVs appeared to have

weaker associations with their respective metabolites compared to neighboring SNPs and

could be tagged by neighboring SNPs with r^2 >0.5 (**Supplementary Fig. 10**). Conditional

analysis further confirmed that the associations of the three SVs with metabolites were

attenuated and even not significant after adjusting for neighboring SNPs. For example, the

HBA1/HBA2/HBM/HBQ1 deletion was in moderate LD with the lead SNP chr16:199621

across the nine metabolic traits (r^2=0.76). After conditioning on the SNP, the association of

the deletion with these traits was notably weakened, although it remained statistically

significant for seven of them (p<0.01; **Supplementary Table 8**), highlighting the significance

of this SV in influencing host metabolic processes that are not completely accounted for by

nearby SNPs. Furthermore, a 64bp insertion (chr19:15866773-15866783) associated with

vitamin K concentration was in strong LD with its tagging SNP (r^2=0.94), while a ~6kb

insertion (chr12:20861159-20861190) associated with the direct bilirubin level was in

moderate LD with its tagging SNP (r^2=0.76). The associations between the two insertions and

metabolites were not significant after adjusting for neighboring SNPs. Apart from these three

SVs, the remaining 30 SVs continued to exhibit significant links with their respective

metabolites after conditional analysis, underscoring the potential driven and regulatory roles

of these SVs in blood metabolic traits (**Supplementary Table 8-9**).

Gene-based analysis of the rare SVs identified 22 unique genes associated with blood

metabolic metabolites and further confirmed the most correlations with cystine

(**Supplementary Table 10**). SV burdens involving six genes, including *CLN3*, *ELF2*, *WDFY4*,

RBPMS, *IL6ST*, and *SORCS3*, were linked to cystine levels. The most significant and the only

one passing multiple test correction (p< 7.6×10^{-8} by adjusting for 5861 genes and 121

metabolites) was for *the CLN3* gene, which encodes a protein that is involved in lysosomal

function and is prenylated most likely at cysteine 435[70]. The other candidate associations were

listed in the **Supplementary Table 10**.

GO enrichment analysis of the gene sets harboring significant SVs identified 26 BP, 16 CC,

and 30 MF terms (**Fig. 5b; Supplementary Table 11**). The enriched gene sets predominantly

featured biological process pathways encompassing gas transport, detoxification and redox,

protein and receptor transport, metabolism, and degradation, indicating that SVs associated

with blood metabolites play a pivotal role in these biological processes. Additionally, KEGG

enrichment analysis revealed a notable enrichment of the gene sets in pathways associated

with infectious diseases (e.g., African trypanosomiasis, malaria), bile secretion, and

glycosylation-related metabolic pathways (**Fig. 5c; Supplementary Table 12**), implying that

SVs located near these genes played crucial roles in these metabolic pathways.

Replication of previously reported metabolites-associated SVs

In the present study, we investigated previously reported metabolites-associated SVs to determine their replication. Out of a total of 58 autosomal SVs identified in previous studies[26-28], 33 were detected in our samples, with 16 showing associations with metabolites ($p < 0.05$; **Supplementary Table 13-15**). Notably, following a stringent Bonferroni correction threshold of 0.00152 (0.05/33), three SVs remained statistically significant. The most robust signal was observed for *HBA1/HBA2/HBM/HBQ1* (16:165396-184701) SV, which was associated with MCV/MCH, as confirmed by our study, the NHLBI TOPMed program, and the study by Wu et al. Additionally, the SV within the growth hormone receptor (GHR) gene, previously linked to the small round epithelial cell (SRC) count in the study by Wu et al., was found to be associated with multiple metabolites, including cystine, aspartic acid, and glutamic acid in our study. Moreover, the duplication involving the T cell receptor alpha variable (TRAV) genes, known to be associated with C-reactive protein (CRP) levels in the study by Chen et al., was also correlated with valine levels in our study. These replicated associations across different studies underscore the importance of some specific SVs in human metabolism, emphasizing the need for further validation in larger populations to confirm additional potential associations and to advance our understanding in this domain.

Host SVs associated with both microbiome and metabolites help understand their interactions

Furthermore, we investigated the potential role of host SVs to intricate interactions between the gut microbiome and blood metabolites. Initially, we observed that host SVs showed pleiotropic associations with both microbial features and plasma metabolites, demonstrating significant p-values reaching genome-wide significance ($p < 2.22 \times 10^{-6}$) for one phenotype and $p < 0.05$ for the other. Specifically, out of the 30 SVs significantly linked to the gut microbiome, 29 were found to be correlated with at least one blood metabolite (**Fig. 6a**; **Supplementary Table 16**). Similarly, among the 33 SVs significantly associated with blood metabolites, 31 were also correlated with at least one gut microbial feature (**Fig. 6b**; **Supplementary Table 17**).

Next, we directly calculated the phenotypic correlations between gut microbial features and blood metabolic traits. This approach resulted in the identification of 21 "microbial feature - metabolic trait" pairs meeting stringent significance criteria (associated SV with $p < 2.22 \times 10^{-6}$ for microbiome, $p < 0.05$ for the metabolic trait, and $p < 0.05$ for correlation between them; **Supplementary Table 16**). For example, the deletion on chromosome 8 (SV length: 141 bp) significantly associated with *Clostridium ramosum* ($\beta = 0.131$, $p = 7.47 \times 10^{-7}$) was correlated with the level of serum uric acid ($\beta = 0.073$, $p = 1.52 \times 10^{-3}$). The SV was also linked to an increased risk of hyperuricemia ($\beta = 1.608$, $p = 4.01 \times 10^{-4}$; **Supplementary Table 18**).

Moreover, significant positive correlations were observed between *Clostridium ramosum* and
serum uric acid ($\beta = 0.060$, $p = 3.61 \times 10^{-3}$; **Fig. 6c**) as well as hyperuricemia ($\beta = 0.124$,
$p=0.02$; **Fig. 6d**). Another instance highlighted an insertion on chromosome 5 (SV length: 96
bp) significantly linked to both *Citrobacter* ($\beta = -0.129$, $p = 1.33 \times 10^{-6}$) and 1-methylhistidine
($\beta=-0.076$, $p = 4.22 \times 10^{-3}$), unveiling intricate relationships within *Citrobacter* and
1-methylhistidine ($\beta = 0.048$, $p = 0.023$).

Likewise, we also identified 325 "metabolic trait—microbial feature" pairs that met the
significance criteria (associated SV with $p<2.22 \times 10^{-6}$ for metabolic trait, $p<0.05$ for the
microbial feature, and $p<0.05$ for correlation between them; **Supplementary Table 17**). For
example, the *HBA1/HBA2/HBM/HBQ1* deletion (chr16:165396-184700_DEL), significantly
associated with nine blood metabolites such as MCHC and zinc, was also linked to the
relative abundance of the microbial pathway MF0064: the triacylglycerol degradation ($\beta = -0.082$, $p = 6.47 \times 10^{-10}$). A significant association was observed between MCHC and MF0064:
the triacylglycerol degradation ($\beta = 0.075$, $p = 2.85 \times 10^{-3}$; **Fig. 6e**). A significant correlation
between zinc and MF0064: the triacylglycerol degradation was also observed ($\beta = -0.039$, $p = 4.95 \times 10^{-2}$; **Fig. 6f**). These results indicated a potential regulatory network involving microbial
and metabolic components, underscoring the importance of considering SVs in the context of
microbial/metabolic associations.

To infer causal relationships, we utilized SVs as instrumental variables (IVs) in a Mendelian
randomization (MR) analysis. This approach revealed 24 significant causal relationships
involving 21 microbial traits and 14 metabolic traits (**Supplementary Table 19**).
Observational correlation and MR analysis consistently confirmed the causal effect of
Clostridium ramosum on increased serum uric acid level and increased risk of hyperuricemia
(**Fig. 6c-d**), as well as that of *Citrobacter* on increased 1-methylhistidine level. Moreover,
blood MCHC, zinc, and manganese showed positive causal effects on the microbial pathway
MF0064: the triacylglycerol degradation (**Fig. 6e-f**). Additionally, tyrosine was found to be
causally correlated with abundances of diverse gut microbiota, including *Salmonella enterica*,
Bacteroides intestinalis, MF0052: arginine degradation II, Gammaproteobacteria, *Providencia*
rettgeri, MF0093: propionate production I, and Enterobacteriaceae. These results highlight
how host SVs could genetically determine the levels of blood metabolites and the gut
microbiome, even when the effect of one phenotype was mediated by the abundance of the
other phenotype, facilitating a deeper understanding of the interactive mechanisms shaping
these biological phenotypes.

Discussion

This study represents a comprehensive investigation into the role of SVs in the gut
microbiome, blood metabolites, and their interactions. Through high-depth whole-genome
sequencing of 2,002 individuals, we identified 138,859 high-confidence SVs, nearly half of

which were novel. Association analysis revealed 30 significant links between SVs and
microbial taxa/pathways, as well as 48 significant associations between SVs and blood
metabolites. Among these, 18 reached the study-wide significance. These findings highlight
the intricate interaction between host genetic variation and the gut microbiome and
metabolites, emphasizing the critical role of SVs in this process. This comprehensive SV
dataset serves as a valuable genomic resource for the Chinese population and can function as
a reference panel for SVs imputation in GWAS, offering broad utility for future phenotype
association studies, including those focusing on diseases.

This study revealed a strong association between specific SVs, particularly a deletion in the
TENT2 gene, and the relative abundance of *Coprobacillus sp. 29_1*. The *TENT2* gene is
required for the mono-adenylation of certain miRNAs, such as the liver-specific miRNA-122,
which is involved in the regulation of fatty acid homeostasis[71,72]. *Coprobacillus* has been
reported to be less abundant and could be a protective species against psoriasis and psoriatic
arthritis disease[73-75]. Previous studies have shown that individuals with high *AMY1* copy
numbers (*AMY1-CN*) may promote the enrichment of *Ruminococcaceae* microbes (including
Ruminococcus and *Oscillospira*) by consuming a diet rich in resistant starch, facilitating the
efficient fermentation of resistant starch and positively impacting host health[25]. Our analysis
identified nine SVs in the *AMY (AMY1, AMY2A, and AMY2B)* gene regions, with one
spanning *AMY2A* (DUP, chr1:103611001-103620001), which was significantly associated
with *Ruminococcaceae* (β = -0.054, p = 0.043). Additionally, a deletion near the *AMY* gene
region (DEL, chr1:102630524-102630828) was identified as significantly negatively
correlated with *Ruminococcus champanellensis* (β = -0.111, p = 1.74×10^{-5}) and *unclassified*
Ruminococcaceae bacterium_D16 (β = -0.097, p = 2.04×10^{-4}). Moreover, gene-based
analysis further confirmed that *AMY2A/AMY2B* harboring four rare SVs was associated with
the microbial pathway of MF0008:maltose degradation. These results consistently supported
the correlation between the SV of *AMY* gene region and microbial taxa and pathways,
highlighting the role of microbiota in the human genetic adaptation[76].

Among the "SV-blood metabolite" associations, the most significant association was observed
for the 19kb deletion in the *HBA1/HBA2/HBM/HBQ1* gene region. The deletion as the
well-known SEA variant exhibited a "thalassemia-like" pattern of red blood cell phenotypic
associations, characterized by lower MCH, MCV and RBC. These findings were consistent
with previous reports that this deletion is a key SV leading to alpha-thalassemia and is linked
to blood cell traits[27,28,77]. In addition to confirming associations with previously reported
phenotypes (MCV, MCH, MCHC, and RBC), our findings revealed novel correlations with
trace elements, such as copper, zinc, and iron, underscoring the potential utility of this region
in disease screening and diagnosis. Beyond the *HBA1/HBA2/HBM/HBQ1* gene region, other
associations were partially validated. One study reported that recurrent exon deletions in the
haptoglobin (*HP*) gene contribute to lower cholesterol levels[78]. We identified a deletion in the
HP gene (DEL, chr16:72049453-72076286) significantly associated with total cholesterol

(TC, $\beta = 0.061$, $p = 0.04595$). We also found an insertion in the *TOX2* gene (INS,
chr20:44061952-44061987) significantly associated with TC ($\beta = 0.130$, $p = 2.04 \times 10^{-5}$),
LDLC ($\beta = 0.118$, $p = 8.44 \times 10^{-5}$), and HDLC ($\beta = 0.056$, $p = 0.044$). The *TOX2* gene
encodes a transcription factor involved in immune regulation[79]. Cholesterol and low-density
lipoproteins are strongly linked to systemic inflammation[80,81]. This suggests that this SV may
influence systemic inflammation and immune function by regulating blood lipid levels, which
requires further analysis to clarify the mechanism of this association.

A noteworthy outcome of our study is that the majority of SVs appeared to exert a dominant
influence on microbial abundances and metabolic phenotypes, whereas the effects of nearby
SNPs appeared to be relatively limited. Traditional genome-wide association studies (GWAS)
have primarily focused on the role of SNPs; however, many diseases such as rare and
neurodevelopmental disorders[82-84] have been shown to be caused by SVs. Our study
emphasizes the importance of SVs in the gut microbiome and associated metabolite changes,
suggesting that future research should not concentrate exclusively on SNP-centric GWAS.
Furthermore, most SV loci were found to affect both blood metabolites and the gut microbiota,
revealing the important role of SVs in the interplay between microbiota and blood metabolites.
Previous studies have established significant observational correlations between the
microbiome and host metabolism[85-87]. Our study highlights the crucial role of host genetics at
the SV level within this context.

Despite providing valuable insights into SV-microbiota-metabolite interactions, our study had
several limitations. Although our sample size was relatively large, particularly for microbiome
SV GWAS, further studies are needed in an independent cohort or broader populations to
ensure the generalizability of our findings. Moreover, the functional mechanisms of SVs
remain incompletely understood. To gain deeper insights into how SVs influence the
microbiome and regulate host metabolism, additional research using mouse models or
functional experiments is required. As the number of reliable catalogs of SV loci continues to
grow, we will gain deeper insights into the impact of host SVs on the gut microbiome and
blood metabolites, and a more nuanced understanding of the relationship between the gut
microbiota, blood metabolites, and the host's physiological state. As more robust genetic
variants have been identified, future studies should further explore the causal relationships
between microbiota and metabolites, thereby elucidating their potential roles in host health
and disease.

In summary, this study established a reference SV dataset for the Chinese population and
presented compelling evidence regarding the associations of SV with the gut microbiome and
blood metabolites, laying the groundwork for future investigations and enriching our
knowledge of human health and diseases.

Methods

Study cohort and samples

Samples used in this study were obtained from the 4D-SZ omics-cohort, a cohort that has been described elsewhere[14,19,20,29-31]. In the year 2017, a total of 2002 individuals provided blood samples, and 1539 of these individuals also provided matched fecal samples as part of the annual health check program. The blood samples were subjected to high-depth whole genome sequencing and blood metabolic traits testing, whereas the fecal samples were subjected to shotgun metagenomic sequencing. The collection protocols for blood and stool samples, as well as the sequencing procedures, were consistent with those used in earlier studies. Buffy coats were isolated from blood samples, and DNA was extracted using the HiPure Blood DNA Mini Kit (Magen, catalog no. D3111) according to the manufacturer's protocol. Fecal samples were collected using the MGIEasy kit, and stool DNA was extracted using the MetaHIT protocol. Subsequently, DNA concentrations were quantified using Qubit (Invitrogen), and 200 ng of DNA from each sample was used for library preparation. Paired-end 100bp and single-end 100bp sequencing were performed for blood and stool samples, respectively, on the BGISEQ-500 platform[88].

Ethical approval for the study was obtained from the Institutional Review Boards (IRB) of BGI-Shenzhen, and all participants provided written informed consent.

High-depth whole genome sequencing and SNP/indel detection

A total of 2,002 individuals in this study were sequenced to a mean sequencing depth of 42× for the whole genome. The reads were aligned to the human reference genome GRCh38/hg38 using BWA (v.0.7.15)[89] with the default parameters. The adapter sequences and bases with quality value below five were removed. The alignments were indexed in the BAM format using Samtools (v.0.1.18)[90] and PCR duplicates were marked for downstream filtering using Picardtools (v.1.62). The Genome Analysis Toolkit's (GATK v.4.3[91]) BaseRecalibrator created recalibration tables to screen known SNPs and insertions or deletions (INDELs) in the BAM files from dbSNP (v.150). GATKlite (v.2.2.15) was used for the subsequent base quality recalibration and removal of read pairs with improperly aligned segments as determined by Stampy. GATK's HaplotypeCaller was used for the variant discovery. GVCFs containing SNVs and INDELs from GATK HaplotypeCaller were combined (CombineGVCFs), genotyped (GenotypeGVCFs), variant score recalibrated (VariantRecalibrator) and filtered (ApplyRecalibration). During the GATK VariantRecalibrator process, we took our variants as inputs and used four standard SNP sets to train the model: (1) HapMap3.3 SNPs; (2) dbSNP build 150 SNPs; (3) 1000 Genomes Project SNPs from Omni 2.5 chip and (4) 1000 G phase1 high confidence SNPs. The sensitivity thresholds of 99.9% to SNPs and 99% to INDELs were applied for variant selection after optimizing for Transition to Transversion (TiTv) ratios using the GATK ApplyRecalibration command.

A stringent inclusion threshold for SNP/INDEL variants was established: (i) mean depth >8×;

(ii) Hardy-Weinberg equilibrium (HWE) $P > 10^{-5}$; and (iii) genotype calling rate > 98%.
Furthermore, the sample criteria were as follows: (i) mean sequencing depth > 20×; (ii)
variant calling rate > 98%; (iii) no population stratification by performing principal
components analysis (PCA) analysis implemented in PLINK (v1.9) [92] and (iv) excluding
related individuals by calculating pairwise identity by descent (IBD, Pi-hat threshold of
0.1875) in PLINK. Only ten samples were removed after quality control filtering. After
variant and sample quality control, 1,992 individuals with 8.5 million common (MAF ≥ 1%)
SNP/indel variants were left for association analyses.

SV detection
Due to the complexity of SV detection from short-read sequencing data, the use of multiple
SV calling tools is a common practice to promote accuracy and sensitivity. For SV detection,
a   pipeline   was   designed   according   to   the   SV-Adjudicator[33]
(https://github.com/talkowski-lab/SV-Adjudicator)   and   GATK-SV   cohort   mode
(https://github.com/broadinstitute/gatk-sv). In the pipeline, the BAM file is obtained from the
preceding alignment and filtering process as its primary input. Subsequently, Manta (v1.6.0)[93],
Wham (v1.7.0)[94], Svaba (v1.1.3)[95], and Delly (v0.8.7)[96] were run for SV detection, xTea[97] was
run for mobile element insertion discovery, and cn.MOPS (v1.40.0)[98] and GATK gCNV[99]
were run for CNVs calling (**Supplementary Fig. 3**). The algorithms were selected according
to Kosugi et al[100]. Finally, the SV calls from the five algorithms and the CNV calls of each
sample were standardized and integrated using the GATK-SV pipeline. All computations were
completed locally.

Sample batching and quality control for SV
Following the batching strategy delineated by Collins et al[32], all samples that passed the
quality control (QC) were subdivided into five batches, with ~380 samples in each batch
according to the dosage score (WGD). The distribution of female and male samples in each
batch was maintained at a ratio of approximately 1:1, ensuring balance across all batches.
For filtering after GATK-SV integration, outlier samples exhibiting either fewer or
higher-than-average SV calls per contig for each caller, or samples manifesting autosomal
aneuploidies based on the per-batch binned coverage of each chromosome, were excluded.
Additionally, individuals with missing data or those who failed the preceding SNP/indel
filtering were omitted. Finally, 1,839 samples were remained in the final SV callset.

SV filtering
To curate a high-quality SV dataset for subsequent genetic analyses, further filtration of the
SVs was conducted post-GATK-SV pipeline execution and sample quality assessment. SVs
with quality scores (GQ) > 250 or marked as "PASS" were retained, while variants of type
"BND" were excluded. SVs with a genotyping rate >95% and Hardy-Weinberg equilibrium p

value $> 1 \times 10^{-6}$ were kept. For GWAS analysis, only SVs with MAF $>1\%$ were retained,

resulting in a final set of 22,519 SVs left for association analysis.

SV Annotation

The SVs that passed the quality control were annotated using GATK SVAnnotate (GATK

v4.6.0) for functional prediction. The protein-coding GTF file was downloaded from

GENCODE (hg38.v37). Only the protein-coding genes were retained for annotation. The

BED files of the non-coding regions were downloaded from GATK-SV. AnnotSV (v3.3.8)

was applied for additional SV annotation, utilizing the GRCh38 database build to incorporate

information such as the pLI score of disrupted genes.

Validation of SV using NA12878

To validate the SV detection process, the NA12878 human genome reference standard

sequenced on the BGISEQ-500 platform

(https://db.cngb.org/search/experiment/CNX0094812/) was added as a demo into the cohort

for GATK-SV integration. To evaluate the SV detection, the final SV set of the demo was

compared with the reference SV sets of NA12878 constructed with the datasets from the

Database of Genomic Variants (http://dgv.tcag.ca/dgv/app/home) and the long read-derived

SV data[35]. The evaluation was performed using EvalSVcallers

(https://github.com/stat-lab/EvalSVcallers).

Comparisons with published SV datasets

We examined whether the SVs detected in this study have been reported in five publicly

available SV databases, including gnomAD, Audano et al., 2019, HGSVC, Wu et al., and

dbVar (spanning from 2018 to September 30, 2023). The gnomAD project is a high-coverage

(average depth of 32X) short-read sequencing (SRS) dataset comprising 14,891 global

samples. The dbVar database catalogs SVs from multiple platforms. We first ensured that the

coordinates of the SVs in these public databases were in GRCh38. Our strategy involved the

utilization of the R package (v2.3.0) and a custom program to compare our identified SVs

with those in public databases, considering a reciprocal overlap rate of 50% or higher.

Shotgun metagenomic sequencing and profiling

For fecal samples obtained from the SZ-4D cohort, DNA libraries were constructed and

sequenced using the BGISEQ-500-DNA sequencer for subsequent metagenomic analysis.

Raw data were subjected to quality control and adapter trimming using fastp (v0.23.2)

software. To filter out human reads, sequencing reads were aligned to the hg38 reference

genome using SOAP2 (v2.22[101]) with an identity threshold of ≥ 0.9. The gene profiles were

generated by aligning high-quality sequencing reads to the integrated gene catalog (IGC) by

using SOAP2 (identity ≥ 0.95) as previously described[102]. The relative abundance profiles of

phylum, order, family, class, genera, and species were determined from the gene abundances.
To eliminate the influence of sequencing depth in the comparative analyses, we downsized the
unique IGC mapped reads to 20 million for each sample. The relative abundance profiles of
genes, phyla, orders, families, classes, genera and species were determined accordingly using
the downsized mapped reads per sample.
Gut metabolic modules (GMMs) characterize the bacterial and archaeal metabolism specific
to the human gut, emphasizing anaerobic fermentation processes[103]. The current GMM set
was developed through a comprehensive review of the literature and metabolic databases,
including MetaCyc[104] and KEGG, followed by expert curation to define modules and
alternative pathways. Ultimately, we identified 616 common microbial taxa and GMMs
present in at least 10% of the samples.
714
Microbiome association analysis with common SVs
We examined the relationship between common host SVs (MAF $\geq$ 1%) and gut taxa/GMMs
by utilizing linear or logistic models based on the abundance of each taxon/GMM. The
abundance of taxon/GMM with an occurrence rate exceeding 95% in the cohort was
log-transformed, and outliers (deviating over four standard deviations from the mean) were
excluded to ensure a reliable quantitative trait analysis. Alternatively, the taxon/GMM was
categorized into presence/absence pattern to mitigate zero inflation issues, treating the
abundance as a dichotomous trait. Subsequently, for SNP/indel and SV identified in this
cohort, we conducted a standard single variant-based mbGWAS analysis using PLINK (v1.9)
[92] with a linear model for quantitative traits or a logistic model for dichotomous traits.
Considering the influence of diet and lifestyle on microbial characteristics, we incorporated
age, gender, BMI, defecation frequency, stool form, 12 diet/lifestyle factors, and the top four
principal components (PCs) as covariates for mbGWAS analysis.
728
Microbiome association analysis with rare SVs
The SV burden test was conducted via generalized linear regression (glm) model with adding
age, gender, BMI, defecation frequency, stool form, 12 diet/lifestyle factors, along with the
top four PCs as covariates. The gene-based sequence kernel association test (SKAT) analysis
was performed in rare SVs (MAF < 0.01) using the SKAT (v2.2.4) package in R with method
= 'skato'. Age, gender, BMI, defecation frequency, stool form, 12 diet/lifestyle factors, and
the top four PCs were added into the model as covariates. The rare SVs were filtered using
plink v1.9 with options '--hwe 1e-4 --max-maf 0.01 --geno 0.02 --mind 0.02'. The gene-based
SV annotation was performed as described in SV Annotation section. An SV was annotated
within the gene if it overlapped with the gene region or was predicted to affect the promoter
of the gene.
740
Metabolic trait profiling
In this study, metabolic traits, including anthropometric characteristics and blood metabolites,

were assessed in all participants during physical examinations. Anthropometric data,
including height, weight, waist circumference, and hip circumference, were recorded by
nurses, while age and gender were self-reported. Clinical evaluations, such as blood tests and
urinalysis, were conducted by a certified physical examination organization. Hormones were
measured in a similar manner, with 250 µl of plasma and atmospheric pressure chemical
ionization (APCI). Trace elements were analyzed using a 7700x ICP-MS (Agilent
Technologies) equipped with octupole reaction system (ORS) technology to reduce spectral
interferences, using 200 µl of whole blood. Water-soluble vitamins were assessed by UPLC
coupled to a Xevo TQ-S Triple Quad mass spectrometer (Waters) with ESI in positive ion
mode, utilizing 200 µl of plasma. Fat-soluble vitamins were measured with UPLC linked to
an AB Sciex Qtrap 4500 mass spectrometer using APCI and 250 µl of plasma. Blood amino
acids were quantified using ultra-high pressure liquid chromatography (UHPLC) coupled
with an AB Sciex Qtrap 5500 mass spectrometer, utilizing electrospray ionization (ESI) in
positive ion mode with 40 µl of plasma. Hyperuricemia is defined as an elevated serum uric
acid level, with greater than 6 mg/dL in women and 7 mg/dL in men.

Metabolites association analysis
The log10-transformed median-normalized values of 121 anthropometric and metabolic traits
were used as quantitative traits. Samples with missing values or values greater than 4 standard
deviations from the mean were excluded from the association analysis. Each of the common
SNP/indel or common SV (MAF ≥ 1%) was independently tested using a linear model for
quantitative traits in PLINK v1.9. The gene-based sequence kernel association test (SKAT)
analysis was performed for rare SVs (MAF < 0.01) using the SKAT (v2.2.4) package in R, as
done in the microbiome association analysis. Covariates included age, gender, and the top
four PCs.

Gene enrichment analysis

The genes harboring significantly associated SVs were annotated using AnnotSV[54-56]. Gene
Ontology (GO) and Kyoto Encyclopaedia of Genes and Genomes (KEGG) gene functional
enrichment analysis of genes harboring significantly associated SVs were executed by using
the R packages including clusterProfiler, enrichplot, ggplot2 and org.Hs.eg.db. A standardized
metric (*p*-value < 0.05, Q-value < 0.2) was used to prioritize the top functional items and
pathways. Biological process (BP), cellular component (CC), and molecular function (MF)
were the three categories included in GO analysis.

Observational correlation analyses
All microbial features and metabolic traits were transformed using log-transformed
median-normalized function to reduce the skewness of the distributions. The observational
correlations between these variables were evaluated using multivariable linear regression

analysis while adjusting for age and gender. The beta coefficient and p-value were obtained.

One-sample MR analysis.

To explore the potential causal relationships between microbial traits and metabolic
characteristics within the same cohort, we conducted a one-sample bidirectional MR analysis
in this cohort with both metabolite and microbiome data. We selected SV instruments based
on a stringent genome-wide significance threshold ($p < 2.22 \times 10^{\square\square}$) and performed
clumping with an LD r² < 0.1 to ensure independence among genetic variants for MR analysis.
An unweighted polygenic risk score (PRS) was calculated for each participant based on
independent genetic variants derived from GWAS data. For each SV, we coded the risk alleles
as 0, 1, or 2, depending on the number of trait-specific risk alleles carried by the individual.
Instrumental variable (IV) analyses were conducted using two-stage least squares (TSLS)
regression. In the first stage, we assessed the association between the PRS and the
observational phenotype using linear regression, from which we derived fitted values
representing the predicted exposure. During this process, the *F*-statistics and the variance
explained by IVs were calculated. We required the *F*-statistics over 10 to ensure a strong IV
for analysis. In the second stage, we regressed the outcome trait on the genetically predicted
exposure from the first stage. Both stages were adjusted for potential confounders, including
age, gender, and BMI. TSLS was performed using the 'ivreg' function from the AER package
in R (v.1.2-9; https://www.rdocumentation.org/packages/AER/versions/1.2-9/topics/ivreg).

Data Availability

All summary statistics that support the findings of this study including the population SV
dataset were available in the **Supplementary Tables**. The gut metagenomic sequencing data
after removing host reads have been publicly available in the GSA from
https://ngdc.cncb.ac.cn/gsa-human/browse/HRA000907. All the metabolites and
individual-level host genetics data have been uploaded to the GSA database
(https://ngdc.cncb.ac.cn/bioproject/browse/PRJCA005334). According to the Human Genetic
Resources Administration of China regulation and the institutional review board of
BGI-Shenzhen related to protecting individual privacy, these data are controlled-access and
are available via an application on request. The human reference hg38 datasets are publicly
available from http://hgdownload.soe.ucsc.edu/goldenPath/hg38/bigZips/. The datasets of
common SVs are available at the following link: GnomAD SV:
https://datasetgnomad.blob.core.windows.net/dataset/release/4.1/genome_sv/gnomad.v4.1.sv.
sites.vcf.gz, dbvar SV: https://gnomad.broadinstitute.org/, HGSVC SV:
http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/HGSVC2/release/v1.0/integrated_
callset/freeze3.sv.alt.vcf.gz, Wu et al SV:
https://ftp.ncbi.nlm.nih.gov/pub/dbVar/data/Homo_sapiens/by_study/csv/all_variants_for_nst

d212.csv.gz, Audano et al. 2019：

http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/hgsv_sv_discovery/working/20181

025_EEE_SV-Pop_1.

Author Contributions

X.L and T.Z. conceived and organized this study. X.X., H.Y., Y.Z., W.L, X.J., and L.X. led the

organization of the cohort, the sample collection, and the questionnaire collection. M.H. and

H.L led the DNA extraction and sequencing. X.L and X.T. processed the whole genome data.

L.Z., Y.J., and M.L. processed the metagenome data. X.L., X.T., and L.Z. performed the

metagenome-genome-wide association analyses and Mendelian randomization analyses. X.L.

wrote the raw manuscript. All authors contributed to data and texts in this manuscript.

Competing interests

The authors declare no competing interest.

Acknowledgments

We sincerely thank the support by the National Natural Science Foundation of China (No.

32200548) and the China National GeneBank. We thank all the volunteers for their time and

for self-collecting the oral samples using our kit. The schematic workflow diagram of the

study was drawn by Figdraw (https://www.figdraw.com).

Reference

1　Liu, R. *et al.* Gut microbiome and serum metabolome alterations in obesity and after

weight-loss intervention. *Nat Med* **23**, 859-868 (2017).

https://doi.org:10.1038/nm.4358

2　Wu, C. *et al.* Obesity-enriched gut microbe degrades myo-inositol and

promotes lipid absorption. *Cell Host & Microbe* **32**, 1301-1314.e1309 (2024).

https://doi.org:10.1016/j.chom.2024.06.012

3　Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2

diabetes. *Nature* **490**, 55-60 (2012). https://doi.org:10.1038/nature11450

4   Mei, Z. *et al.* Strain-specific gut microbial signatures in type 2 diabetes identified in a

cross-cohort analysis of 8,117 metagenomes. *Nature Medicine* **30**, 2265-2276 (2024).

https://doi.org:10.1038/s41591-024-03067-7

5   Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel

diseases. *Nature* **569**, 655-662 (2019). https://doi.org:10.1038/s41586-019-1237-9

6   Zheng, J. *et al.* Noninvasive, microbiome-based diagnosis of inflammatory bowel

disease. *Nature Medicine* (2024). https://doi.org:10.1038/s41591-024-03280-4

7   Zhang, X. *et al.* The oral and gut microbiomes are perturbed in rheumatoid arthritis

and partly normalized after treatment. *Nature Medicine* **21**, 895-905 (2015).

https://doi.org:10.1038/nm.3914

8   Alpizar-Rodriguez, D. *et al.* Prevotella copri in individuals at risk for rheumatoid

arthritis. *Ann Rheum Dis* **78**, 590-593 (2019).

https://doi.org:10.1136/annrheumdis-2018-214514

9   Jie, Z. *et al.* The gut microbiome in atherosclerotic cardiovascular disease. *Nat*

Commun **8**, 845 (2017). https://doi.org:10.1038/s41467-017-00900-1

10   Zhu, F. *et al.* Transplantation of microbiota from drug-free patients with schizophrenia

causes schizophrenia-like abnormal behaviors and dysregulated kynurenine

metabolism in mice. *Mol Psychiatry* (2019).

https://doi.org:10.1038/s41380-019-0475-4

11   Zhu, F. *et al.* Metagenome-wide association of gut microbiome features for

schizophrenia. *Nature Communications* **11**, 1612 (2020).

https://doi.org:10.1038/s41467-020-15457-9

12   Goodrich, J. K. *et al.* Genetic Determinants of the Gut Microbiome in UK Twins. *Cell*

Host Microbe **19**, 731-743 (2016). https://doi.org:10.1016/j.chom.2016.04.017

13   Goodrich, J. K., Davenport, E. R., Waters, J. L., Clark, A. G. & Ley, R. E.

Cross-species comparisons of host genetic associations with the microbiome. *Science*

352, 532-535 (2016). https://doi.org:10.1126/science.aad9379

14   Liu, X. *et al.* Mendelian randomization analyses support causal relationships between

blood metabolites and the gut microbiome. *Nat Genet* **54**, 52-61 (2022).

https://doi.org:10.1038/s41588-021-00968-y

15   Rühlemann, M. C. *et al.* Genome-wide association study in 8,956 German individuals

identifies influence of ABO histo-blood groups on gut microbiome. *Nature genetics* **53**,

147-155 (2021). https://doi.org:10.1038/s41588-020-00747-1

16   Kurilshikov, A. *et al.* Large-scale association analyses identify host factors influencing

human gut microbiome composition. *Nature genetics* **53**, 156-165 (2021).

https://doi.org:10.1038/s41588-020-00763-1

17   Lopera-Maya, E. A. *et al.* Effect of host genetics on the gut microbiome in 7,738

participants of the Dutch Microbiome Project. *Nature Genetics* **54**, 143-151 (2022).

https://doi.org:10.1038/s41588-021-00992-y

18   Qin, Y. *et al.* Combined effects of host genetics and diet on human gut microbiota and

incident disease in a single population cohort. *Nature genetics* **54**, 134-142 (2022).

https://doi.org:10.1038/s41588-021-00991-z

19   Liu, X. *et al.* A genome-wide association study for gut metagenome in Chinese adults

illuminates complex diseases. *Cell Discovery* **7**, 9 (2021).

https://doi.org:10.1038/s41421-020-00239-w

20   Liu, X. *et al.* Metagenome-genome-wide association studies reveal human genetic

impact on the oral microbiome. *Cell Discovery* **7**, 117 (2021).

https://doi.org:10.1038/s41421-021-00356-0

21   Stankiewicz, P. & Lupski, J. R. Structural Variation in the Human Genome and its Role

in Disease. *Annual Review of Medicine* **61**, 437-455 (2010).

https://doi.org:https://doi.org/10.1146/annurev-med-100708-204735

22   D'Haene, E. & Vergult, S. Interpreting the impact of noncoding structural variation in

neurodevelopmental disorders. *Genet Med* **23**, 34-46 (2021).

https://doi.org:10.1038/s41436-020-00974-1

23   Fu, J. M. *et al.* Rare coding variation provides insight into the genetic architecture and

phenotypic context of autism. *Nature genetics* **54**, 1320-1331 (2022).

https://doi.org:10.1038/s41588-022-01104-0

24   Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature*

578, 112-121 (2020). https://doi.org:10.1038/s41586-019-1913-9

25   Poole, A. C. *et al.* Human Salivary Amylase Gene Copy Number Impacts Oral and Gut

Microbiomes. *Cell Host Microbe* **25**, 553-564 e557 (2019).

https://doi.org:10.1016/j.chom.2019.03.001

26   Chen, L. *et al.* Association of structural variation with cardiometabolic traits in Finns.

The American Journal of Human Genetics **108**, 583-596 (2021).

https://doi.org:https://doi.org/10.1016/j.ajhg.2021.03.008

27   Wheeler, M. M. *et al.* Whole genome sequencing identifies structural variants

contributing to hematologic traits in the NHLBI TOPMed program. *Nature*

Communications **13**, 7592 (2022). https://doi.org:10.1038/s41467-022-35354-7

28 Wu, Z. *et al.* Structural variants in the Chinese population and their impact on

phenotypes, diseases and population adaptation. *Nature Communications* **12**, 6501

(2021). https://doi.org:10.1038/s41467-021-26856-x

29 Jie, Z. *et al.* A transomic cohort as a reference point for promoting a healthy human

gut microbiome. *Medicine in Microecology* **8**, 100039 (2021).

https://doi.org:https://doi.org/10.1016/j.medmic.2021.100039

30 Liu, X. *et al.* A genome-wide association study reveals the relationship between

human genetic variation and the nasal microbiome. *Communications Biology* **7**, 139

(2024). https://doi.org:10.1038/s42003-024-05822-5

31 Liu, X. *et al.* Sex differences in the oral microbiome, host traits, and their causal

relationships. *iScience* **26**, 105839 (2023). https://doi.org:10.1016/j.isci.2022.105839

32 Collins, R. L. *et al.* A structural variation reference for medical and population genetics.

Nature **581**, 444-451 (2020). https://doi.org:10.1038/s41586-020-2287-8

33 Werling, D. M. *et al.* An analytical framework for whole-genome sequence association

studies and its implications for autism spectrum disorder. *Nature genetics* **50**, 727-736

(2018). https://doi.org:10.1038/s41588-018-0107-y

34 Abel, H. J. *et al.* Mapping and characterization of structural variation in 17,795 human

genomes. *Nature* **583**, 83-89 (2020). https://doi.org:10.1038/s41586-020-2371-0

35 Kosugi, S. *et al.* Comprehensive evaluation of structural variation detection algorithms

for whole genome sequencing. *Genome Biology* **20**, 117 (2019).

941   https://doi.org:10.1186/s13059-019-1720-5

942 36 Audano, P. A. *et al.* Characterizing the Major Structural Variant Alleles of the Human

943   Genome. *Cell* **176**, 663-675.e619 (2019). https://doi.org:10.1016/j.cell.2018.12.019

944 37 Ebert, P. *et al.* Haplotype-resolved diverse human genomes and integrated analysis of

945   structural variation. *Science* **372**, eabf7117 (2021).

946   https://doi.org:doi:10.1126/science.abf7117

947 38 Lappalainen, I. *et al.* dbVar and DGVa: public archives for genomic structural variation.

948   *Nucleic acids research* **41**, D936-D941 (2012). https://doi.org:10.1093/nar/gks1213

949 39 Manivannan, S. N. *et al.* De novo FZR1 loss-of-function variants cause developmental

950   and epileptic encephalopathies. *Brain* **145**, 1684-1697 (2022).

951   https://doi.org:10.1093/brain/awab409

952 40 Li, H. U. I., Zhou, C., Kuang, M. E. I., Liu, Y. U. N. & Chen, J. The role of FZR1 in

953   tumorigenesis: Focus on cell-cycle control. *Biocell* **47**, 2177-2186 (2023).

954   https://doi.org:https://doi.org/10.32604/biocell.2023.029373

955 41 Byrska-Bishop, M. *et al.* High-coverage whole-genome sequencing of the expanded

956   1000 Genomes Project cohort including 602 trios. *Cell* **185**, 3426-3440.e3419 (2022).

957   https://doi.org:10.1016/j.cell.2022.08.004

958 42 Lam, M. *et al.* Comparative genetic architectures of schizophrenia in East Asian and

959   European populations. *Nature* *genetics* **51**, 1670-1678 (2019).

960   https://doi.org:10.1038/s41588-019-0512-x

961 43 Yang, A. *et al.* TENT2, TUT4, and TUT7 selectively regulate miRNA sequence and

962   abundance. *Nature* *Communications* **13**, 5260 (2022).

https://doi.org:10.1038/s41467-022-32969-8

44   Liu, S. *et al.* The Host Shapes the Gut Microbiota via Fecal MicroRNA. *Cell Host*

Microbe **19**, 32-43 (2016). https://doi.org:10.1016/j.chom.2015.12.005

45   Martini, A. M., Moricz, B. S., Woods, L. J. & Jones, B. D. Type IV Pili of Streptococcus

sanguinis Contribute to Pathogenesis in Experimental Infective Endocarditis.

Microbiology Spectrum **9**, e01752-01721 (2021).

https://doi.org:doi:10.1128/Spectrum.01752-21

46   Sirmaci, A. *et al.* Mutations in ANKRD11 cause KBG syndrome, characterized by

intellectual disability, skeletal malformations, and macrodontia. *Am J Hum Genet* **89**,

289-294 (2011). https://doi.org:10.1016/j.ajhg.2011.06.007

47   Benson, A. K. *et al.* Individuality in gut microbiota composition is a complex polygenic

trait shaped by multiple environmental and host genetic factors. *Proc Natl Acad Sci U*

S A **107**, 18933-18938 (2010). https://doi.org:10.1073/pnas.1007028107

48   Org, E. *et al.* Genetic and environmental control of host-gut microbiota interactions.

Genome research **25**, 1558-1569 (2015). https://doi.org:10.1101/gr.194118.115

49   Hills, L. B. *et al.* Deletions in GRID2 lead to a recessive syndrome of cerebellar ataxia

and tonic upgaze in humans. *Neurology* **81**, 1378-1386 (2013).

https://doi.org:10.1212/WNL.0b013e3182a841a3

50   La Rosa, S. L. *et al.* The human gut Firmicute Roseburia intestinalis is a primary

degrader of dietary β-mannans. *Nature Communications* **10**, 905 (2019).

https://doi.org:10.1038/s41467-019-08812-y

51   Louis, P. & Flint, H. J. Formation of propionate and butyrate by the human colonic

microbiota. *Environ. Microbiol.* **19**, 29-41 (2017).

https://doi.org:https://doi.org/10.1111/1462-2920.13589

52   Zhao, Q. *et al.* Parabacteroides distasonis ameliorates hepatic fibrosis potentially via

modulating intestinal bile acid metabolism and hepatocyte pyroptosis in male mice.

Nature Communications **14**, 1829 (2023).

https://doi.org:10.1038/s41467-023-37459-z

53   Markowitz, R. H. G. *et al.* Microbiome-associated human genetic variants impact

phenome-wide disease risk. *Proc Natl Acad Sci U S A* **119**, e2200551119 (2022).

https://doi.org:10.1073/pnas.2200551119

54   Geoffroy, V. *et al.* The AnnotSV webserver in 2023: updated visualization and ranking.

Nucleic acids research **51**, W39-w45 (2023). https://doi.org:10.1093/nar/gkad426

55   Geoffroy, V. *et al.* AnnotSV and knotAnnotSV: a web server for human structural

variations annotations, ranking and analysis. *Nucleic acids research* **49**, W21-w28

(2021). https://doi.org:10.1093/nar/gkab402

56   Geoffroy, V. *et al.* AnnotSV: an integrated tool for structural variations annotation.

Bioinformatics **34**, 3572-3574 (2018). https://doi.org:10.1093/bioinformatics/bty304

57   Li, J., Teng, W., Yu, Y., Hou, X. & Shan, Z. Linkage Analysis of the Chromosome

5q31-33 Region Identifies JAKMIP2 as a Risk Factor for Graves' Disease in the

Chinese Han Population. *Med Sci Monit* **25**, 1439-1451 (2019).

https://doi.org:10.12659/msm.911489

58   Ma, T. *et al.* Long Noncoding RNA JAKMIP2-AS1 Promotes the Growth of Colorectal

Cancer and Indicates Poor Prognosis. *Onco Targets Ther* **14**, 763-772 (2021).

1007    https://doi.org:10.2147/ott.S289617

1008 59  Baxter, N. T., Zackular, J. P., Chen, G. Y. & Schloss, P. D. Structure of the gut

1009    microbiome following colonization with human feces determines colonic tumor burden.

1010    *Microbiome* **2**, 20 (2014). https://doi.org:10.1186/2049-2618-2-20

1011 60  Prasad, H. & Visweswariah, S. S. Impaired Intestinal Sodium Transport in

1012    Inflammatory Bowel Disease: From the Passenger to the Driver's Seat. *Cell Mol*

1013    *Gastroenterol Hepatol* **12**, 277-292 (2021).

1014    https://doi.org:10.1016/j.jcmgh.2021.03.005

1015 61  Harris, R. A. *et al.* Colonic Mucosal Epigenome and Microbiome Development in

1016    Children and Adolescents. *Journal of Immunology Research* **2016**, 9170162 (2016).

1017    https://doi.org:https://doi.org/10.1155/2016/9170162

1018 62  Larmonier, C. B. *et al.* Reduced colonic microbial diversity is associated with colitis in

1019    NHE3-deficient mice. *Am J Physiol Gastrointest Liver Physiol* **305**, G667-677 (2013).

1020    https://doi.org:10.1152/ajpgi.00189.2013

1021 63  Li, J. *et al.* Integrating microbial GWAS and single-cell transcriptomics reveals

1022    associations between host cell populations and the gut microbiome. *Nature*

1023    *Microbiology* **10**, 1210-1226 (2025). https://doi.org:10.1038/s41564-025-01978-w

1024 64  Astbury, S. *et al.* Lower gut microbiome diversity and higher abundance of

1025    proinflammatory genus Collinsella are associated with biopsy-proven nonalcoholic

1026    steatohepatitis. *Gut Microbes* **11**, 569-580 (2020).

1027    https://doi.org:10.1080/19490976.2019.1681861

1028 65  Shimizu, H. *et al.* Sucrose-preferring gut microbes prevent host obesity by producing

exopolysaccharides. *Nature Communications* **16**, 1145 (2025).

https://doi.org:10.1038/s41467-025-56470-0

66 Zhang, Y. X. *et al.* An exploratory study of the association between KCNB1 rs1051295

and type 2 diabetes and its related traits in Chinese Han population. *PLoS One* **8**,

e56365 (2013). https://doi.org:10.1371/journal.pone.0056365

67 Qiu, Q. W. *et al.* Evidence of recent natural selection on the Southeast Asian deletion

(--(SEA)) causing α-thalassemia in South China. *BMC Evol Biol* **13**, 63 (2013).

https://doi.org:10.1186/1471-2148-13-63

68 Gao, Y. *et al.* A pangenome reference of 36 Chinese populations. *Nature* **619**,

112-121 (2023). https://doi.org:10.1038/s41586-023-06173-7

69 Liu, X. *et al.* Mendelian randomization analyses support causal relationships between

blood metabolites and the gut microbiome. *Nature genetics* **54**, 52-61 (2022).

https://doi.org:10.1038/s41588-021-00968-y

70 Järvelä, I. *et al.* Biosynthesis and Intracellular Targeting of the CLN3 Protein Defective

in Batten Disease. *Human Molecular Genetics* **7**, 85-90 (1998).

https://doi.org:10.1093/hmg/7.1.85

71 Warkocki, Z., Liudkovska, V., Gewartowska, O., Mroczek, S. & Dziembowski, A.

Terminal nucleotidyl transferases (TENTs) in mammalian RNA metabolism. *Philos*

Trans R Soc Lond B Biol Sci **373** (2018). https://doi.org:10.1098/rstb.2018.0162

72 Katoh, T. *et al.* Selective stabilization of mammalian microRNAs by 3' adenylation

mediated by the cytoplasmic poly(A) polymerase GLD-2. *Genes Dev* **23**, 433-438

(2009). https://doi.org:10.1101/gad.1761509

73   Scher, J. U. *et al.* Decreased bacterial diversity characterizes the altered gut

microbiota in patients with psoriatic arthritis, resembling dysbiosis in inflammatory

bowel disease. *Arthritis Rheumatol* **67**, 128-139 (2015).

https://doi.org:10.1002/art.38892

74   Olejniczak-Staruch, I. *et al.* Alterations of the Skin and Gut Microbiome in Psoriasis

and Psoriatic Arthritis. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22083998

75   Zou, X., Zou, X., Gao, L. & Zhao, H. Gut microbiota and psoriasis: pathogenesis,

targeted therapy, and future directions. *Front Cell Infect Microbiol* **14**, 1430586 (2024).

https://doi.org:10.3389/fcimb.2024.1430586

76   Suzuki, T. A. & Ley, R. E. The role of the microbiota in human genetic adaptation.

Science **370** (2020). https://doi.org:10.1126/science.aaz6827

77   Ireland, J. H. *et al.* Detection of the (--SEA) double alpha-globin gene deletion by a

simple immunologic assay for embryonic zeta-globin chains. *Am J Hematol* **44**, 22-28

(1993). https://doi.org:10.1002/ajh.2830440106

78   Boettger, L. M. *et al.* Recurring exon deletions in the HP (haptoglobin) gene contribute

to lower blood cholesterol levels. *Nature genetics* **48**, 359-366 (2016).

https://doi.org:10.1038/ng.3510

79   Seo, H. *et al.* TOX and TOX2 transcription factors cooperate with NR4A transcription

factors to impose CD8[+] T cell exhaustion. *Proceedings of the National Academy of*

Sciences **116**, 12410-12415 (2019). https://doi.org:doi:10.1073/pnas.1905675116

80   Tall, A. R. & Yvan-Charvet, L. Cholesterol, inflammation and innate immunity. *Nat Rev*

Immunol **15**, 104-116 (2015). https://doi.org:10.1038/nri3793

81   Jukema, R. A., Ahmed, T. A. N. & Tardif, J.-C. Does low-density lipoprotein

cholesterol induce inflammation? If so, does it matter? Current insights and future

perspectives for novel therapies. *BMC Medicine* **17**, 197 (2019).

https://doi.org:10.1186/s12916-019-1433-3

82   Mizuguchi, T. *et al.* A 12-kb structural variation in progressive myoclonic epilepsy was

newly identified by long-read whole-genome sequencing. *Journal of Human Genetics*

64, 359-368 (2019). https://doi.org:10.1038/s10038-019-0569-5

83   Cen, Z. *et al.* Intronic pentanucleotide TTTCA repeat insertion in the SAMD12 gene

causes familial cortical myoclonic tremor with epilepsy type 1. *Brain* **141**, 2280-2288

(2018). https://doi.org:10.1093/brain/awy160

84   Kaivola, K. *et al.* Genome-wide structural variant analysis identifies risk loci for

non-Alzheimer's dementias. *Cell Genom* **3**, 100316 (2023).

https://doi.org:10.1016/j.xgen.2023.100316

85   Visconti, A. *et al.* Interplay between the human gut microbiome and host metabolism.

Nature Communications **10**, 4505 (2019).

https://doi.org:10.1038/s41467-019-12476-z

86   Kinross, J. M., Darzi, A. W. & Nicholson, J. K. Gut microbiome-host interactions in

health and disease. *Genome Medicine* **3**, 14 (2011). https://doi.org:10.1186/gm228

87   Dekkers, K. F. *et al.* An online atlas of human plasma metabolite signatures of gut

microbiome composition. *Nature Communications* **13**, 5370 (2022).

https://doi.org:10.1038/s41467-022-33050-0

88   Fang, C. *et al.* Assessment of the cPAS-based BGISEQ-500 platform for

metagenomic sequencing. *Gigascience* **7**, 1-8 (2018).

https://doi.org:10.1093/gigascience/gix133

89   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754-1760 (2009).

https://doi.org:10.1093/bioinformatics/btp324

90   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352

91   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297-1303

(2010). https://doi.org:10.1101/gr.107524.110

92   Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based

linkage analyses. *Am J Hum Genet* **81**, 559-575 (2007).

https://doi.org:10.1086/519795

93   Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline and

cancer sequencing applications. *Bioinformatics* **32**, 1220-1222 (2016).

https://doi.org:10.1093/bioinformatics/btv710

94   Kronenberg, Z. N. *et al.* Wham: Identifying Structural Variants of Biological

Consequence. *PLoS Comput Biol* **11**, e1004572 (2015).

https://doi.org:10.1371/journal.pcbi.1004572

95   Wala, J. A. *et al.* SvABA: genome-wide detection of structural variants and indels by

local assembly. *Genome research* **28**, 581-591 (2018).

https://doi.org:10.1101/gr.221028.117

96   Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end and

split-read   analysis.   *Bioinformatics*   **28**,   i333-i339   (2012).

https://doi.org:10.1093/bioinformatics/bts378

97   Chu, C. *et al.* Comprehensive identification of transposable element insertions using

multiple sequencing technologies. *Nature Communications* **12**, 3836 (2021).

https://doi.org:10.1038/s41467-021-24041-8

98   Klambauer, G. *et al.* cn.MOPS: mixture of Poissons for discovering copy number

variations in next-generation sequencing data with a low false discovery rate. *Nucleic*

acids research **40**, e69 (2012). https://doi.org:10.1093/nar/gks003

99   Van der Auwera, G. A. & O'Connor, B. D. *Genomics in the cloud: using Docker, GATK,*

and WDL in Terra.   (O'Reilly Media, 2020).

100   Kosugi, S. *et al.* Comprehensive evaluation of structural variation detection algorithms

for   whole   genome   sequencing.   *Genome*   *Biol*   **20**,   117   (2019).

https://doi.org:10.1186/s13059-019-1720-5

101   Li, R. *et al.* SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics*

25, 1966-1967 (2009). https://doi.org:10.1093/bioinformatics/btp336

102   Li, J. *et al.* An integrated catalog of reference genes in the human gut microbiome. *Nat*

Biotechnol **32**, 834-841 (2014). https://doi.org:10.1038/nbt.2942

103   Vieira-Silva, S. *et al.* Species-function relationships shape ecological properties of the

human   gut   microbiome.   *Nat*   *Microbiol*   **1**,   16088   (2016).

https://doi.org:10.1038/nmicrobiol.2016.88

104   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes and the

BioCyc collection of Pathway/Genome Databases. *Nucleic acids research* **42**,

D459-471 (2014). https://doi.org:10.1093/nar/gkt1103

1141

1142

Figure 1. Characteristics of the SVs discovered in this study (4D-SZ Cohort).

(**a**) Descriptive statistics of the SVs count for each SV type.

(**b**) The count of SVs per sample by each SV type.

(**c**) SV count distribution by minor allele frequency (MAF) and size.

(**d**) Length distribution of SVs. The peaks represent the Alu, SVA, and LINE1 events.

(**e**) The count of SVs according to the predicted function region annotation by each SV type.

(**f**) The count of SVs per sample according to the predicted function region annotation by each SV type.

(**g**) Venn diagram showing the overlap between SVs in this study and published SV datasets. Among the SVs, 49.57% are novel, while 50.43% have been previously reported.

(**h**) The difference in SV counts between novel and reported SVs according to the predicted function region annotation.

(**i**) The counts of affected genes per genome by novel and reported SVs according to the predicted function region annotation. Differences between novel and reported SVs were assessed with two-sided t-tests. CPXs were not included in this analysis due to a lack of accurate annotation of the genes they affected.

Abbreviations: SV, structural variant; DEL, deletions; DUP, duplications; INS, insertions; INV, inversions; CPX, complex variants; CNV, copy number variation.

Figure 2. Genome-wide associations of SVs with the gut microbiome identified by mbGWAS.

(**a**) Manhattan plot representing the GWAS results (619 microbial features; 22,519 common high-quality SVs with MAF $\geq$ 1%). The y-axis indicates the log10-transformed p-values, while the x-axis indicates the genomic positions of the SVs. The study-wide significance (p $< 3.59 \times 10^{-9}$) and genome-wide significance (p $< 2.2 \times 10^{-6}$) are indicated by horizontal lines colored blue and black, respectively. The annotations highlight the study-wide association of deletion variant at the intronic region of *TENT2* gene and the species *Coprobacillus sp. 29_1*.

(**b**)The distribution of genomic inflation factor (λ) for 616 mbGWAS tests, ranging from 0.84 to 1.33 with a median value of 0.98. The λ of GWAS for species *Coprobacillus sp. 29_1* is 0.87.

(**c**) The regional plot showing mbGWAS results of the *TENT2* deletion (purple diamond) and its nearby SNPs associated with the species *Coprobacillus sp. 29_1*.

(**d**) The log-transformed relative abundances of *Coprobacillus sp. 29_1* in carriers of *TENT*2 deletion and non-carriers.

Figure 3. The raw and conditional GWAS results for significant SVs and their nearby SNPs/indels with the gut microbiome.

(**a**) The raw associations of SVs and their nearby SNPs/indels (flanking 500kb of the SV) with gut microbiome.

(**b**) The associations of SVs and their nearby SNPs after conditioning on the tagging SNP/indel showing the strongest LD with the SV. The bar in the right indicates the significance of SV association after conditional analysis, including no longer significant (colored in grey), not reaching genome-wide significant (colored in dark red) and retaining genome-wide significant

1186 (colored in red).

1187 The x axis indicates the log10-transformed p values. The y axis indicates the associations of the

1188 SVs and nearby SNPs/indels with gut microbiome (N = 30). The red dots represent the p-values

1189 associated with SVs, and the blue dots represent the p-values associated with SNPs/indels.

1190

1191 Figure 4. Genome-wide associations of SVs with blood metabolites.

1192 (**a**) Manhattan plot depicting the GWAS results (121 blood metabolite traits; 22,519 common

1193 high-quality SVs with MAF $\geq$ 1%). The study-wide significance ($p < 1.84 \times 10^{-8}$) and

1194 genome-wide significance ($p < 2.2 \times 10^{-6}$) are indicated by horizontal lines colored blue and black,

1195 respectively. The annotations show the DEL variant on chromosome 16 (chr16: 165396-184700),

1196 corresponding to the *HBA1*, *HBA2*, *HBM*, and *HBQ1* genes.

1197 (**b**) Boxplots of nine blood metabolite levels associated with the *HBA1/HBA2/HBM/HBQ1* DEL

1198 that passed study-wide significance, comparing the metabolite levels between DEL carriers ("0/1":

1199 individuals with DEL) and non-carriers ("0/0": individuals without DEL).

1200 MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin; MCHC, mean corpuscular

1201 hemoglobin Concentration; RBC distribution, red blood cell distribution.

1202

1203 Figure 5. The enrichment analysis of significantly associated SVs in blood
1204 metabolites.

1205 (**a**) The distribution shows the number of SVs significantly associated with blood metabolites

1206 (N=48).

1207 (**b and c**) GO (b) and KEGG enrichment analysis (c) for genes harboring SVs significantly

1208 associated with the blood metabolites. The bar plot (b) only displays the biological process (BP)

1209 results from the GO analysis terms.

1210

1211 Figure 6. Causal relationships between the gut microbiome and blood
1212 metabolites supported by SVs.

1213 (**a**) Venn diagrams showing the number of overlapped SVs significantly associated with the gut

1214 microbiome (p $< 2.2 \times 10^{-6}$) that also linked to blood metabolites ($p < 0.05$).

1215 (**b**) Venn diagrams showing the number of overlapped SVs significantly associated with the blood

1216 metabolites (p $< 2.2 \times 10^{-6}$) that also linked to the gut microbiome ($p < 0.05$).

1217 (**c**) The deletion on chromosome 8 (SV length: 141 bp) was significantly associated with both

1218 *Clostridium ramosum* ($\beta = 0.131$, $p = 7.47 \times 10^{-7}$) and the blood metabolite serum uric acid ($\beta =$

1219 0.073, $p = 1.52 \times 10^{-3}$). A significant observational correlation and further causal effect of

1220 *Clostridium ramosum* on serum uric acid supported by MR.

1221 (**d**) A significant observational correlation and further causal effect of *Clostridium ramosum* on

1222 hyperuricemia supported by MR.

1223 (**e**) The causal effect of MCHC on MF0064: the triacylglycerol degradation supported by their

1224 overlapped genetic SV, such as the HBA1/HBA2/HBM/HBQ1 deletion

1225 (chr16:165396-184700_DEL).

1226 (**f**) The causal effect of zinc on MF0064: the triacylglycerol degradation supported by their

1227 overlapped genetic SV, such as the HBA1/HBA2/HBM/HBQ1 deletion

1228 (chr16:165396-184700_DEL).

1229   Obs., observational correlation; MR, Mendelian randomization. MCHC, mean corpuscular

1230   hemoglobin Concentration;

1231

Graphic abstract

Based on a multi-omics cohort of 2,002 individuals comprising high-depth whole genome and metagenomic sequencing data, blood metabolites, and detailed questionnaires , we identified 138,859 high-confidence SVs, uncovering 36 significant associations between SVs and microbial taxa/pathways, as well as 48 associations between SVs and blood metabolites. Mendelian randomization using SVs demonstrated causal relationships, such as *Clostridium ramosum*'s causal effect on serum uric acid and hyperuricemia. This study provides a detailed SV catalog for the Chinese population and highlights the significant role of SVs in the gut microbiome, blood metabolites, and their causal relationships.

Figure 1. Characteristics of the SVs discovered in this study (4D-SZ Cohort).

(**a**) Descriptive statistics of the SVs count for each SV type.

(**b**) The count of SVs per sample by each SV type.

(**c**) SV count distribution by minor allele frequency (MAF) and size.

(**d**) Length distribution of SVs. The peaks represent the Alu, SVA, and LINE1 events.

(**e**) The count of SVs according to the predicted function region annotation by each SV type.

(**f**) The count of SVs per sample according to the predicted function region annotation by each SV type.

(**g**) Venn diagram showing the overlap between SVs in this study and published SV datasets. Among the SVs, 49.57% are novel, while 50.43% have been previously reported.

(**h**) The difference in SV counts between novel and reported SVs according to the predicted function region annotation.

(**i**) The counts of affected genes per genome by novel and reported SVs according to the predicted function region annotation. Differences between novel and reported SVs were assessed with

two-sided t-tests. CPXs were not included in this analysis due to a lack of accurate annotation of the genes they affected.

Abbreviations: SV, structural variant; DEL, deletions; DUP, duplications; INS, insertions; INV, inversions; CPX, complex variants; CNV, copy number variation.

Figure 2. Genome-wide associations of SVs with the gut microbiome identified by mbGWAS.

(**a**) Manhattan plot representing the GWAS results (619 microbial features; 22,519 common high-quality SVs with MAF $\geq$ 1%). The y-axis indicates the log10-transformed p-values, while the x-axis indicates the genomic positions of the SVs. The study-wide significance (p $< 3.59 \times 10^{-9}$) and genome-wide significance (p $< 2.2 \times 10^{-6}$) are indicated by horizontal lines colored blue and black, respectively. The annotations highlight the study-wide association of deletion variant at the intronic region of *TENT2* gene and the species *Coprobacillus sp. 29_1*.

(**b**)The distribution of genomic inflation factor (λ) for 616 mbGWAS tests, ranging from 0.84 to 1.33 with a median value of 0.98. The λ of GWAS for species *Coprobacillus sp. 29_1* is 0.87.

(**c**) The regional plot showing mbGWAS results of the *TENT2* deletion (purple diamond) and its nearby SNPs associated with the species *Coprobacillus sp. 29_1*.

(**d**) The log-transformed relative abundances of *Coprobacillus sp. 29_1* in carriers of *TENT2* deletion and non-carriers.

Figure 3. The raw and conditional GWAS results for significant SVs and their nearby SNPs/indels with the gut microbiome.

(a) The raw associations of SVs and their nearby SNPs/indels (flanking 500kb of the SV) with gut microbiome.

(b) The associations of SVs and their nearby SNPs after conditioning on the tagging SNP/indel showing the strongest LD with the SV. The bar in the right indicates the significance of SV association after conditional analysis, including no longer significant (colored in grey), not reaching genome-wide significant (colored in dark red) and retaining genome-wide significant (colored in red).

The x axis indicates the log10-transformed p values. The y axis indicates the associations of the SVs and nearby SNPs/indels with gut microbiome (N = 30). The red dots represent the p-values associated with SVs, and the blue dots represent the p-values associated with SNPs/indels.

Figure 4. Genome-wide associations of SVs with blood metabolites.

(**a**) Manhattan plot depicting the GWAS results (121 blood metabolite traits; 22,519 common high-quality SVs with MAF ≥ 1%). The study-wide significance (p < 1.84×10^{-8}) and genome-wide significance (p < 2.2×10^{-6}) are indicated by horizontal lines colored blue and black, respectively. The annotations show the DEL variant on chromosome 16 (chr16: 165396-184700), corresponding to the *HBA1*, *HBA2*, *HBM*, and *HBQ1* genes.

(**b**)Boxplots of nine blood metabolite levels associated with the *HBA1/HBA2/HBM/HBQ1* DEL that passed study-wide significance, comparing the metabolite levels between DEL carriers ("0/1": individuals with DEL) and non-carriers ("0/0": individuals without DEL).

MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin; MCHC, mean corpuscular hemoglobin Concentration; RBC distribution, red blood cell distribution.

Figure 5. The enrichment analysis of significantly associated SVs in blood metabolites.

(**a**)The distribution shows the number of SVs significantly associated with blood metabolites (N=48).

(**b and c**) GO (b) and KEGG enrichment analysis (c) for genes harboring SVs significantly associated with the blood metabolites. The bar plot (b) only displays the biological process (BP) results from the GO analysis terms.

Figure 6. Interactions between the gut microbiome and blood metabolites supported by SVs.

(a) Venn diagrams showing the number of overlapped SVs significantly associated with the gut microbiome ($p < 2.2 \times 10^{-6}$) that also linked to blood metabolites ($p < 0.05$).

(b) Venn diagrams showing the number of overlapped SVs significantly associated with the blood metabolites ($p < 2.2 \times 10^{-6}$) that also linked to the gut microbiome ($p < 0.05$).

(c) The deletion on chromosome 8 (SV length: 141 bp) is significantly associated with both *Clostridium ramosum* ($\beta = 0.131$, $p = 7.47 \times 10^{-7}$) and the blood metabolite serum uric acid ($\beta = 0.073$, $p = 1.52 \times 10^{-3}$). A significant observational correlation and further causal effect of *Clostridium ramosum* on serum uric acid was supported by MR.

(d) The causal effect of MCHC on MF0064: the triacylglycerol degradation supported by their overlapped genetic SV, such as the HBA1/HBA2/HBM/HBQ1 deletion (chr16:165396-184700_DEL).

(e) The causal effect of zinc on MF0064: the triacylglycerol degradation supported by their overlapped genetic SV, such as the HBA1/HBA2/HBM/HBQ1 deletion (chr16:165396-184700_DEL).

(f) The causal effect of zinc on MF0064: the triacylglycerol degradation supported by their overlapped genetic SV, such as the HBA1/HBA2/HBM/HBQ1 deletion

(chr16:165396-184700_DEL).

Obs., observational correlation; MR, Mendelian randomization. MCHC, mean corpuscular hemoglobin Concentration;