1   Differential temperature adaptation mechanisms in the High Arctic-adapted *Cerastium regelii*
2   Ostenf. and the widespread *Stellaria longipes* Goldie.

Sarah L. Lane[1], Lauren A E Erland [1*]

[1] Agriculture, University of the Fraser Valley, Stó:lō Temexw, Abbotsford, BC, Canada, V2S 7M8

ORCID:
Erland - 0000-0003-3157-0535
Lane - 0009-0008-5507-2956

*Corresponding Author
Lauren Erland
Assistant Professor
University of the Fraser Valley | Agriculture, CEP Q109c
33844 King Rd, Abbotsford, BC, V2S 7M8
lauren.erland@ufv.ca | 604-504—7221 x 2485

Summary (200 words max)

- Climate change impacts arctic latitudes more acutely than other latitudes, resulting in arctic shrubification. How individual species in these climes respond to warming temperature is poorly understood. Understanding species resiliency to climate change will help us conserve plant species at risk.

- We performed a survey of plants in a permafrost anomaly in Resolute (Qausuittuq), Nunavut, Canada. Two identified species, *Stellaria longipes* Goldie and *Cerastium regelii* Ostenf., were investigated through modelled niche suitability under future climate scenarios, phenological analysis, and *in vitro* warming experiments to investigate growth and phytochemical profiles.

- 10 species including *Stellaria longipes* and *Cerastium regelii* were identified in the anomaly. Predicted niche suitability increased under SSP126 for *C. regelii*, with compressed and later flowering period since 1850. *In vitro, S. longipes* maximized growth at 24 °C with greater abundance of cytokinins than *C. regelii*, which increased growth at 28 °C.

- *Stellaria longipes* is self-limiting at higher temperatures, and is less temperature-dependent for its success, while *C. regelii* is more affected by warming temperatures, showing increases in growth and predicted niche suitability. Our work increases understanding of plant resiliency and vulnerability in Canada's High Arctic, and sheds light on the biology of an understudied arctic specialist in *C. regelii*.

Keywords (5-8, alphabetical order): chickweed; climate change; ecological niche model; long stalk starwort; metabolomics; Nunavut; plant environment resilience; temperature stress

Introduction

As climate change has shifted to become our global reality, high latitude and high altitude regions are experiencing the effects of these changes more acutely than other global regions (Rantanen *et al.*, 2022). Inuit Nunangat is a distinct geopolitical region, with Inuit self-governance, which is also referred to as Canada's High Arctic. While significant knowledge has been generated surrounding predicted impacts of climate change on terrestrial and marine ecosystems in this region, relatively little is still understood about the plants from a conservation perspective. For example, a review of the assessment status of medicinal plants in Canada in 2022 found that only 22 % of plant species have been assessed in Nunavut for a conservation ranking by NatureServe and related conservation data centres (Erland, Turi and Murch, 2022). Much of the available research has focused on woody plants examining the concept of shrubification of Arctic and tundra environments globally. Under these predicted future environments woody shrubs, including willows and perennial berry plants, are anticipated to grow taller with increasing growing degree days in their environments (Myers-Smith *et al.*, 2011; Kodl *et al.*, 2024). As primary producers, plants form the basis of ecosystems. Understanding how plants will respond to future environment scenarios can help to better predict and understand future environment ecosystem interactions.

Plant vulnerability and resiliency requires understanding how plants perceive and respond to environmental cues. Recent literature has proposed that vulnerability of plants to climate change is determined by three key factors: (a) exposure, i.e. extrinsic factors, (b) sensitivity, essentially the degree to which the plant is affected by the extrinsic factors, and (c) adaptive capacity, the ability for a plant to respond *in situ* to a changing climate and persist (Dawson *et al.*, 2011; Foden *et al.*, 2013). The first two factors have been widely investigated and the NatureServe Climate Change Vulnerability Index provides a foundation for identification of generalized adaptive traits, however, there is still a general lack of information on many of the traits mediating adaptive capacity. One of the ways that plants can interact with the world is through the production of specialized phytochemicals. Plant growth regulators or phytohormones are a set of phytochemicals produced by plants, which enable plants to perceive environmental changes and respond by redirecting growth, detoxifying stress metabolites and stabilizing physiological processes (Erland, Saxena and Murch, 2017).

In 2019 we conducted a botanical survey and collection in Resolute (Qausuittuq) Nunavut, which included 83 vouchers and 74 seed collections, with 46 vouchers submitted to the BOLD database and the Canadian Museum of Nature Herbarium. As part of this effort, we worked with local Inuk guide, Brandy Iqaluk, and we identified an area of interest, which was a permafrost anomaly with increased temperature and associated modified plant community composition. This site unlike surrounding permafrost had an abundance of herbaceous species present. We hypothesized that this region could serve as a model for how the plant community composition could shift under future climate scenarios. This anomaly was dominated by the presence of two species in the family Caryophyllaceae which were present, but not abundant, in the surrounding areas. Therefore, our overall objective was to integrate data from field survey, ecological niche modelling, in vitro temperature experiments and metabolomics to predict and understand the differential impacts of environmental warming on two related herbaceous species growing in the Canadian High Arctic: *Cerastium regelii* Ostenf. and *Stellaria longipes* Goldie.

Stellaria longipes, longstalk starwort or Goldie's starwort is a small herbaceous member of the Caryophyllaceae family, first identified in the woods of Ontario. It has a broad circumboreal distribution and has been found to grow in diverse ecosystem types including both arctic and alpine environments, though some debate exists in the literature about the potential existence of several ecotypes, species or subspecies regionally, with the species being referred to as a species complex (Porsild, Porsild and Cody, 1980; Chinnappa *et al.*, 2005; Sharples, 2019; *Stellaria longipes s. lat.*, no date). Extensive previous work has established *S. longipes* as a remarkably phenotypically and morphologically plastic species, which likely contributes significantly to this confusion. Alpine and arctic specimens tend to have compact or dwarf phenotypes (referred to as alpine ecotypes), while plants growing in lower elevation and altitude locations including prairie ecosystems are much taller with longer internodal lengths (prairie ecotype). While initially hypothesized to be an environmentally sensitive species, as it is very responsive to temperature and photoperiod cues, *S. longipes* has been found to be remarkably resilient through in field experimentation (Macdonald, Reid and Chinnappa, 1986; Emery, Chinnappa and Chmielewski, 1994; Maillette *et al.*, 2000; Li *et al.*, 2011) with the ecotypes have differential tolerance to stresses including drought stress (Sangtarash *et al.*, 2009). The role of plant hormones including abscisic acid, gibberellins,

ethylene, auxins and cytokinins in mediating these responses (Kathiresan, Reid and Chinnappa,

1996; Kurepin *et al.*, 2006, 2008; Li *et al.*, 2011), especially stem elongation in response to changes

in photoperiod and temperature (Macdonald, Chinnappa and Reid, 1984; Macdonald, Reid and

Chinnappa, 1986; Thuy, Yeung and Chinnappa, 2009) (Macdonald *et al.*, 1984, 1986; Thuy *et al.*,

2009). Together this makes *S. longipes* an interesting and well characterized system for the study

of environmental resilience (Chinnappa *et al.*, 2005).

Cerastium regelii or Regel's chickweed is a subarctic to arctic species found across the

circumpolar region, with populations restricted primarily to above 70 °N (Heide, Pedersen and

Dahl, 1990). In contrast very little literature has been published on *C. regelii*, with for example a

search for the species "*Cerastium regelii*" yielding only two documents on Web of Science, and

only 242 observations globally recorded on iNaturalist (*Regel's chickweed*, no date). Much of the

available information is in the form of occurrence records and presence in botanical surveys e.g.

(Hanssen, 1932; Fredskild, 1984; Matveyeva, 1994) or inclusion in phylogenetic studies e.g. (Yao

et al., 2021; Liu *et al.*, 2023). A study of plant community composition after glacial moraine retreat

in Svalbard found that *C. regelii* had higher occurrence in sites with higher amounts of fine texture

soil cover. Heide et al. 1990 describe photoperiod and flowering time requirements for this species,

noting that while it is a short day plant, it grows exclusively in a region where it would experience

long day lengths during the flowering season. The authors also identify *Cerastium jenisejense*

(tundra chickweed) as a synonym and high temperature morphotype of *C. regelii* (Heide, Pedersen

and Dahl, 1990; Kew Science, no date), though inclusion of the synonym in our Web of Science

search does not expand the literature available. Lastly, a cyclopeptide deemed 'regelin A' has been

isolated from *C. regelii* with the primary interest in cyclopeptides being as natural products (Ai-

Qun *et al.*, 2004; Zorzi, Deyle and Heinis, 2017).

We hypothesize that *S. longipes*, a generalist species with a pan-Canadian distribution

would have higher temperature resilience as defined by enhanced growth under a wider

temperature range, than *C. regelii*, and that this would be associated with modified phytohormone

metabolism profiles. Guided by our in-field observations, we use three main approaches to

investigate this hypothesis (1) in vitro germination studies, (2) phenology & ecological niche

modelling and (3) untargeted hormonomics studies. We generate new information on the little

studied arctic chickweed *C. regelii* and generate novel hypotheses for the role of phytohormones

in environmental resilience.

Materials and Methods

Botanical Collections and Ecological Surveys

Field samples of growing aerial tissues of *Cerastium regelii* Ostenf. and *Stellaria longipes*

Goldie were collected in July - August 2019 from sites just outside the Hamlet of Resolute,

Qausuittuq, Nunavut, near 3 Mile Lake (74.77011, -95.09212). Samples were collected under

Wildlife Research Permit #2019-047 granted by the Nunavut Department of Environment pursuant

to the Wildlife Act. Working with local Inuk guide Brandy Iqaluk, we asked "Are there areas that

you have noticed in which the plants are changing in recent years?". An area of interest was

identified for sampling due to its unusually high temperature compared to the surrounding areas

and unique plant community composition. This site showed no signs of animal activity such as a

lemming den or carcass which may explain the increased plant density and Ms. Iqaluk did not

know of any previous use of the site which could account for, for example, a greater nutrient

availability.

Thermal readings of the site were conducted using a FLIR ONE Pro thermal infrared

camera and the FLIR ONE app (Teledyne FLIR, USA). The feature was divided into three zones

for survey of the vegetation based on the differences in the plant community and "inside", "middle"

and "outside" region (Figure 1) which had distinct community composition. Detailed photos were

taken of each of the regions. The plants present were identified in the field and density was counted

based on the number of stems. While this does not provide an accurate representation of the

number of individual plants due to the dense matte forming growth form of the chickweeds, it

provides an approximate of the coverage stem density of the species.

For all species botanical vouchers were collected and were submitted for DNA barcoding

and confirmation of botanical identification to the Herbarium at the Canadian Museum of Nature

(BOLD Project ID: CCDB-25796).

Seed and aerial tissue samples were collected. Tissues were collected and transported in nutrient solution composed of one electrolyte tablet (Nuun Hydration Sport "Citrus Fruit", NPN 80089152) in 500 mL of water with 1% Head and Shoulders shampoo containing zinc pyrithione (Classic, Proctor & Gamble), prior to establishment of cultures at the University of British Columbia, Okanagan. Seed samples were isolated from receptacle tissue and stored in envelopes at -20 °C.

Establishment of an in vitro germplasm collection in controlled environments

Field-collected explants were transported from Nunavut to Kelowna, BC over 9 days. Upon arrival in the lab, explants were surface sterilized in a 10 % sodium hypochlorite (Chlorox ® Bleach) with 1 % Head and Shoulders shampoo for 15 minutes followed by three rinses with sterile distilled water. Explants were cultured onto Murashige and Skoog basal media (MSO) (Murashige and Skoog, 1962) containing full strength MS salts with Gamborg's B5 Vitamins (Gamborg, Miller and Ojima, 1968), 3% sucrose and 0.25 % phytagel, pH adjusted to 5.75 using NaOH. All media was sterilized by autoclave (Steris, Mentor, OH, USA; 121 °C at 18 psi for 20 min) prior to being poured across 100 mm x 15 mm petri dishes (Thermo Fisher Scientific, Canada). Sequential surface sterilization and re-subcultures of field-collected explants was necessary to establish axenic material for the germplasm collection and chemical studies. Cultures were maintained in a growth chamber at 21 °C under full spectrum white light (Sunblaster™ full spectrum daylight fluorescent bulbs; 40 μmol/m^2/s), with a 16 h photoperiod. Once established, species were subcultured as needed when media was spent or tissues had fully filled the plate.

Botanical Records, Species Distribution

Historical occurrence records for both species were downloaded from the Global Biodiversity Information Facility (GBIF):

Cerastium regelii (LAEE0038):https://doi.org/10.15468/dl.pcu6j5
Stellaria longipes (LAEE0051):https://doi.org/10.15468/dl.6uqxtz)

Data was cleaned to include only 'SPECIES' (and for *S. longipes* also 'SUBSPECIES = longipes') and country 'CA', duplicates were removed and the data was formatted in species

with data (SWD) format. Occurrences recorded during the 2019 field collection trip were then
added.

Changes in phenology in the historic record were screened. Occurrence data acquired
from the GBIF used in ecological niche models was manually screened for the presence of
flowers, flower buds and seeds. Data was grouped into the date ranges 1850-1950, 1951 – 1975,
1976-2000, 2000-2010 ad 2011-2019. The proportion of the observations with visible flowers or
flower buds was calculated by month and plotted in Prism (GraphPad, v10).

Ecological Niche Modelling

Ecological niche modelling using the maximum entropy model (Maxent) was performed
following procedures previously described (Hirabayashi, Murch and Erland, 2022) with
modification. In brief, ENMEvaluate, ENMTools and ecospat R packages were used for variable
selection and to optimize model settings (regularization multiplier and feature type) and variables
retained for assessment are detailed in Table S1 (Muscarella *et al.*, 2014; Kass *et al.*, 2021).
ENMEvaluate was run with spatial block partitions, testing regularization multiplier (RM, 1-5)
and feature type (linear (L), quadratic (Q), hinge (H), LQ or LQH), with parallel = TRUE,
algorithm = Maxent.jar, all other settings were left as default. Random background points were
generated in QGIS (v3.16.14 long term release, QGIS.org, 2021) using the Database of Global
Administration Areas (GADM) shapefile (Level 0) for Canada (www.gadm.org) as the extent.
Optimal models were selected based on 10 % omission rate, and where similar choosing the model
with the lower Akaike information criterion (AICc) value (Muscarella *et al.*, 2014; Kass *et al.*,
2021). Null models were constructed using the optimized parameters as an additional quality check
to ensure empirical > null (Table S2).

Environmental variables including landcover (Tuanmu and Jetz, 2014), topography
(Amatulli *et al.*, 2018), edaphic properties (Poggio *et al.*, 2021), habitat heterogeneity (Tuanmu
and Jetz, 2015) and bioclimatic variables (Fick and Hijmans, 2017) were assessed and selected to
avoid collinearity based on correlations as reported by the dataset authors original reports (Tuanmu
and Jetz, 2014, 2015; Fick and Hijmans, 2017; Amatulli *et al.*, 2018; Poggio *et al.*, 2021) and as
described in Hirabayashi et al. 2022 (Table S2). Soil and edaphic properties were downloaded
from the SoilGrids2 Web Coverage Service (WCS mean @ 0-5 cm) as both species have relatively
shallow root systems. Topographic, consensus landcover and habitat heterogeneity variables were

downloaded from www.earthenv.org at 30 arc seconds resolution based on GMTED2010 and

median aggregation where applicable (Amatulli *et al.*, 2018). Consensus land cover variables were

downloaded with DIScover (includes integration of Global Land Cover Characterization, GLCC

v2.0) and included: regularly flooded vegetation, cultivated and managed vegetation, snow and ice

cover (Tuanmu and Jetz, 2014). Habitat heterogeneity variables included: homogeneity, evenness,

entropy, dissimilarity, and coefficient of variation (coeffvar) (Tuanmu and Jetz, 2015). Historic

bioclimatic variables were downloaded from WorldClim v2.1 at 30 arc second resolution. Future

climate data (bioclimatic variables) were downloaded from www.worldclim.org for the Coupled

Model Intercomparison Project (CMIP) 6 multi-model ensemble under the least and most extreme

shared socioeconomic pathways (SSPs): 126 (lowest emissions, smallest increase in temperature)

and 585 (highest emissions, largest increase in temperature) averages; for projections to 2041-

2060 and 2061-2080 (referred to as 2060 and 2080). Three global climate models (GCMs):

CNRM-CM6-1 (CMIP6.CMIP.CNRM-CERFACS.CNRM-CM6-1; Voldoire *et al.*, 2019),

MIROC6 (CMIP6.ScenarioMIP.MIROC.MIROC6; Shiogama, Abe and Tatebe, 2019) and MRI-

ESM2-0 (CMIP6.CMIP.MRI.MRI-ESM2-0; Yukimoto *et al.*, 2019) were selected based on best

performance in GCMEval (gcm.met.no) using focus regions: Alaska/NW Canada and

Canada/Greenland/Iceland with equal weights for skill evaluation (Parding *et al.*, 2020; Shiogama

et al., 2021). All variables were resampled to a resolution of 30 arc seconds in RStudio.

Maxent was run in dismo in R with RM = 4 for *C. regelii* and = 5 for *S. longipes*. All other

parameters were as follows: feature class = linear quadratic hinge, autofeatures = FALSE, using

maximum of 10,000 background points, maximum of 5,000 iterations, random seed = TRUE,

allowing partial data using raster grids of selected environmental variables. Ten replicate cross-

validation runs, withholding one fifth of the data as testing data were used for training and Maxent

was then run using optimized settings for each species for historic and future climate scenarios.

Receiver operator area under the curve (AUC) and true skill statistic (TSS) were calculated using

the SDMtune R package using optimized Maxent settings for each species and withholding 20%

of data as a testing set and using k-fold cross validation (Vignali *et al.*, 2020). Scores are reported

for training, testing and heldback datasets (Table S3). The output for the GCMs (CNRM-CM6-1,

MIROC6 and MRI-ESM2-0) used under each SSP and at each timeframe were then averaged and

the mean model output (cloglog) was visualized in QGIS.

Temperature Response Experiments

Two node long cuttings were taken from *in vitro* cultured in established plants for each species, with four explants (pseudoreplicates) started per petri plate (12 °C, 24 °C, or 28 °C). Cultures were maintained under cool white fluorescent lights (~40 µmol m^{-2} s^{-1}, Sunblaster T4) with a 16 h photoperiod. Cultures grown for three weeks under test conditions at which point samples were destructively sampled and growth characteristics measured including: length of long shoot, length of longest root, number of shoots, number of roots, number of nodes, number of aerial shoots, total biomass accumulation. A total of five replicate plates per species (*C. regelii* and *S. longipes*) and per temperature treatment were used. Pseudoreplicates were averaged for each replicated prior to plotting and statistical analysis. Data were analyzed by analysis of variance (ANOVA) with post hoc multiple comparisons analysis using Tukey's honestly significant difference and plotted in GraphPad Prism v11, GraphPad,). Experiments were replicated twice.

Metabolomics and Hormonomics Analysis

Sample Preparation

Samples were harvested from the temperature treatment experiments, rinsed to remove any tissue culture medium, patted dry and frozen at -20 °C until analysis. Methods for metabolomics and hormonomics analysis followed previously published standard operating protocols (Giebelhaus, Erland and Murch, 2024). In brief, tissues were homogenized in microcentrifuge tubes using a disposable tissue grinder (Konte Pestle Grinder) under liquid nitrogen and extracted in 50 % methanol (Fisher Scientific, Optima Grade) with 4 % acetic acid (Sigma Aldrich, LC-MS Grade) with 30 s of vortexing followed by 15 minutes sonication in ice water. Samples were centrifuged (3 min, 13, 000 rpm) and supernatant diluted 5X with water. The diluted sample was centrifuge filtered (Millipore Ultra Free MC, 0.22 µm, 1 min @ 13,000 rpm) prior to injection. Extraction solvent and water blanks were included, as well as a mixed hormonomics sample, for compound identification and quality control (Giebelhaus, Erland and Murch, 2024).

Liquid Chromatography and Mass Spectrometry

Tissue extracts (10 μL) were separated by reverse phase chromatography (Phenomenex
Kinetex EVO C18 column (2.1 x 150 mm, 17. μm Torrance, CA, USA) using a Thermo Scientific
Vanquish ultra performance liquid chromatography (UHPLC) with column oven temperature of
30 °C and a flow rate of 0.250 mL/min. A gradient of A: 0.1 % aqueous formic acid B: acetonitrile
was used, 0 – 25 min, 95:5 – 5:95, 25 min – 30 min, 5:95 – 95:5 with an equilibration time of 5
min between runs. Features were detected by a Q-Exactive hybrid quadrupole-orbitrap mass
spectrometer with heated electrospray ionization (HESI) ionization (Thermo Scientific); settings
were as follows: spray voltage: 7kV; sheath gas pressure: 48; auxillary gas flow rate: 11; sweep
gas flow rate:1; capillary temperature: 250 °C; auxiliary gas heater temperature: 300 °C, S-lens RF
level: 50.0, scan mode - Full MS – SIM, resolution: 70,000; polarity: positive; automatic gain
control (AGC); injection time (IT): 200 ms; scan range: 100 – 1200 *m/z*; AGC target: 1E06; loop
count: 5; auxillary gas heater temp: 300 °C; dynamic exclusion: spray stabilization and collision
gas in the C-trap.

Data Processing and Statistical Analysis

Raw data files were exported and processed in MZmine 3.9 (Pluskal *et al.*, 2010; Schmid
et al., 2023). Mass detection used presets of MS1, all scan types and with noise level (baseline
peaks) established at 3E05, a retention time window of 1-19 min and mass range of 100 – 2000
m/z. The mass list was filtered and ADAP chromatogram algorithm use to generate the extracted
ion chromatograms (EIC). For deconvolution, the group intensity threshold was set to three times
noise level (9E05), and the minimum number of consecutive scans set to 5. Minimum absolute
height was set at 7 times noise level (2E06), with a scan to scan accuracy of 0.002 m/z or 10 ppm.
Savitsky Golay smoothing was perfomred using a window of 7 scans. Chromatograph resolving
was completed using the ADAP Chromatogram builder module, using the local minimum resolver
with a chromatographic threshold of 90 %, minimum search range of 0.05 min, minimum relative
height of 0 %, minimum absolute hiehgt of 2E06, minimum peak top/edge ratio of 1.7, minimum
number of data points as a group set to 5, and a peak duriation range of 0.8 min. Settings were
chosen in agreement with suggested settings for the Orbitrap detector for mzmine3. Features were
then aligned across samples using join aligner with settings: m/z tolerance 0.001 or 5 ppm, weight
for m/z 3, retention time tolerance 0.1 min and weight for retention time 1. Gap filling of the

aligned feature list was completed using peak finder, and was applied with settings matching those

used for join aligner with minimum data points 3, retention time tolerance 0.1 min and internsity

tolerance 20%. Features were initially annotated by retention time and accurate mass (precursor

mass) matching to the HormonomicsDB database with a 25 % retention time tolerance and mass

difference of 0.02 m/z. Because the database includes both predicted and observed retention times,

where retention time differed, all annotated features were compared to expected m/z and kept if

within 5 ppm error regardless of matched retention time.The feature list of both annotated and

unannotated features were exported with retention time, m/z ratio, and peak area to Microsoft

Excel for curation and further analysis.

For data curation, detected features were compared to solvent blanks, and removed if they

were also found in both samples and blanks. Features were normalized to mass dry weight (DW)

used during extractions. Features were then imported to Metabolanalyst 6.0 and processed using

the metadata and one-factor modules. During import, data were transformed using a square root

transformation, and scaled using Pareto scaling prior to statistical analysis. For untargeted

metabolomic analysis, a principal component analysis was performed, followed by a two-way

ANOVA using an adjusted p-value cutoff of 0.05 and with false discovery rate for multiple testing

correction. Significant features from this analysis were visualized as a heat map. The heat map was

prepared from the normalized feature list, where data were $\log_2(x+1)$ transformed and mean-

centered, then clustered by both sample and mass features using a Euclidean distance matrix and

Ward's D agglomerative clustering in Genesys. Annotated features were analyzed by two-way

ANOVA with post hoc multiple comparisons analysis using Tukey's honestly significant

difference and plotted using GraphPad Prism v.10.

Results

Field Collection and Species Identification

The permafrost anomaly itself had no signs of animal activity or otherwise that would

increase in nutrient availability leading to increased plant growth. It was warm to the touch which

was initially what brought our interest to the site. We hypothesize that this may be the result of

either permafrost thaw or geological activity. Field surveys identified a total of ten plant species

of vascular plants (excluding grasses and sedges) were present at or adjacent to the permafrost

anomaly including: *Draba subcapitata* Simmons, *Stellaria longipes*, *Papaver cornwallisense*

D.Löve, *Saxifraga cernua* L., *Micranthes nivalis* (L.) Small, *Saxifraga cespitosa* L., *Saxifraga*

oppositifolia L., *Salix arctica* Pall., *Cochlearia groendlandica* L. and *Cerastium regelii*. (Figure

1). Generally, the species found in the centre of the anomaly were herbaceous species, while those

in the surrounding areas had a higher proportion of woody and shrubby species (Figure 1A).

Stellaria longipes (Figure 1C & E) and *C. regelii* (Figure 1B & F) were selected for further

investigation and comparison due to their high population density at the anomaly and their close

phylogenetic relationship (Figure S1).

Figure 1. Plant community composition at the permafrost feature (a), density of (b) Cerastium regelii and (c) Stellaria longipes at

the sites (d) thermal camera readings overlaid with RGB (bottom) image of the anomaly and field photos from Resolute

(Qausuittuq), NU of: (e) S. longipes and (f) C. regelii.

In Vitro Germplasm Collection

Vegetative tissues of *C. regelii* and *S. longipes* were successfully established in culture.

Field collected tissues were plated in small tufts on MSO media and were maintained in perpetual

culture prior to conducting experiments. Both species form low growing mats across the surface

of the media in these conditions. We found that maintenance in petri dishes was more successful

than culture boxes as this helped to minimize contamination in *S. longipes* cultures, which

appeared to be due to the presence of an endophyte which shifted to external growth from plant

tissues when cultured in magenta boxes.

Phenology

Cerastium regelii (Figure 2A) showed a more significant shift in phenological patterns than

S. longipes (Figure 2B). Data from 1850-1950 shows a bimodal distribution in proportion of

observations in bloom with a smaller peak in July followed by a larger maxima in September

(Figure 2A, purple). The most recent observations from 2011 – 2019 show a shift towards a single

slightly skewed peak in July with the intervening decades representing a gradual transition between

the two distributions (Figure 2A, green). In contrast the distributions for *S. longipes* (Figure 2B)

do not have a clear shift in phenological distributions over time, though there are a greater

proportion of observation with flowers in later months in the year in 2011-2022 time frame

compared to 1850-1950. Note that *C. regelii* observations end in 2019 while *S. longipes* end in

2022 due to limitations in the number of observations recorded in the database.

Figure 2. Phenological shifts in flowering time for (A) Cerastium regelii and (B) Stellaria longipes from historic observations in

the Territories displayed as proportion of observations in bloom per month. More recent observations are lighter in colour.

Ecological Niche Modeling

A total of 2,471 occurrence points for *S. longipes* and 79 for *C. regelii* were included in the

final models. Differences in future ecological niche suitability in the Canadian High Arctic showed

a stark contrast between the two species. The models generated predict little change in niche

suitability for *S. longipes* between current and future climate scenarios (Figure 3B) but a significant

increase in future niche suitability for *C. regelii* (Figure 3C). National models align with current

distributions of *C. regelii* with suitable habitat occurring only in the High Arctic with no range

expansion predicted (Figure S2). *Stellaria longipes* national models (Figure S3) again show a more

moderated response with marginal improvements in niche suitability in Northern and alpine

regions. Under Shared Socioeconomic Pathway (SSP) 126, Resolute (Qausuittuq) is predicted to

experience an ~5 °C increase in mean annual temperature, with this increasing to almost 10°C

under SSP 585 (Table 1).

Table 1. Average annual temperature (BIO1) of Cornwallis Island (-96.554589, 75.64527, -93.1-

1039, 74.515367) sampled at 1,000 random points under historic and mean future climate

scenarios.

	ANNUAL MEAN (°C)[2]	ANNUAL DELTA (°C)
HISTORIC	-17.19	
SSP585[1]	-7.50	9.69
SSP126[1]	-12.11	5.08

[1]Average of CNRM, MRI & MIROC projections

[2]Calculated with no data values removed

Variable contributions to models for *C. regelii* and *S. longipes* differed strongly as reflected

in the magnitude of the change in niche suitability under future climates. The largest contributor

to models for *C. regelii* was a bioclimatic variable, bio5 (39%, maximum temperature of the

warmest month, a bioclimatic variable), followed by elevation and coefficient of variation of

habitat heterogeneity (Figure 3A). Habitat homogeneity and elevation were the two largest

contributors to the model for *S. longipes*, while the highest contributing bioclimatic variable was

bio6 (minimum temperature of the coldest month), with a contribution of only ~ 5% (Table S3).

Figure 3. (A) Variable contribution to ecological niche models for Stellaria longipes and Cerastium regelii and predicted change

in suitability based on maxent models for SSP 126 projected to 2080 for the region around Cornwallis Island for (B) Stellaria

longipes, yellow and (C) Cerastium regelii, blue where points indicate occurrence records included in the model, green and

increase in predicted niche suitability, white no change and pink a decrease in predicted niche suitability as compared to current

range predictions.

In Vitro Experimental Warming

Field grown tissues grown under three temperatures 12, 24 or 28 °C of *Stellaria longipes*

had a significant maxima in growth across all parameters determined at 24 °C (Figure 4). While

C. regelii showed a similar and significant pattern in growth for root growth and shoot height,

the number of shoots showed a linear increase as temperature increased, with the highest number

of shoots produced at 28 °C (Figure 4C).

Figure 4. Changes in selected growth parameters for Cerastium regelii and Stellaria longipes after 3 weeks at 12, 24, or 28 °C.

Both species had maximal increases to shoot height (A), root length (B), and number of roots (D) at 24 °C. S. longipes also had

maximal increase to number of shoots at this temperature, but C. regelii instead showed increasing number of shoots with

increasing temperature, with maximum number of shoots observed at 28 °C. Data were analysed with one-way ANOVA with

Tukey's test for multiple comparisons for each species, asterisks represent significant differences (α = 0.05) within a parameter

and species.

Metabolomics and Hormonomics

Following UPLC-MS analysis of plant extracts, a total of 345 mass features were

detected, and 11 of these were annotated as hormones, representing predominantly metabolites

from the cytokinin or tryptophan metabolism pathways. From the pool of both annotated and

unannotated mass features, a two-way ANOVA revealed 274 mass features that were

differentially abundant between species and across increasing temperature (Figure 5A). 18 of the

features identified via ANOVA were unique to species and 14 to temperature. 74 mass features

were significantly different when either species or temperature were considered, 71 of which had

a significant interaction between temperature and species. Principal component analysis showed

that the greatest variation within the dataset was due to species, as shown in principal component

1 representing 28 % of variation (Figure 5B). However, PC2 and PC3 (not shown) collectively

separated samples by temperature, representing a total of ~36 % of all variation.

Hierarchical clustering produced relatively distinct clusters for each condition tested,

with samples of a specific species and temperature being closely related (Figure 5C). *C. regelii*

clustered strongly together, along with *S. longipes* grown at low temperatures. A second large

cluster included *S. longipes* grown at 24 and 28 °C, with some intermixing between treatments,

indicating *S. longipes* significantly modifies metabolism in response to increased temperature.

Notably, this includes cytokinins topolin riboside, dihydrozeatin glucose and zeatin-9-glucoside.

These cytokins were not detected in *C. regelii* at 12 or 28 °C, only at 24 °C.

Figure 5. Untargeted metabolomic analysis of Stellaria longipes and Cerastium regelii extracts after growth at 12, 24, or 28 °C

for 3 weeks. A two-way ANOVA shows unique mass features related to species or temperature, and a large proportion of mass

features shared that have an interaction (A). Principal component analysis (B) separates samples by species along PC1,

representing the most variation within the dataset. A heat map prepared using all significant mass features clustered using

Euclidean distance and Ward's D method (C) highlights strong differences between species and temperature. A large group of

mass features are mostly unique to S. longipes grown at 24 or 28 °C, including some mass features annotated as cytokinins

involved in shoot growth and elongation.

Cytokinin Metabolism

Most annotated features were related to cytokinin biosynthesis. Zeatin increased with

increasing temperature in both *C. regelii* and *S. longipes* (Figure 6), but the normalized abundance

was similar between species at each temperature. In contrast, dihydrozeatin, dihydrozeatin

glucoside, and zeatin-9-glucoside were not detected in *C. regelii*, but significantly increased ($\alpha <$

0.05) with temperature in *S. longipes*, with the highest abundance at 28 °C, suggesting that *C.*

regelii and *S. longipes* utilize different signalling pathways to adapt to high temperature. Zeatin-

7-glucoside abundance in *C. regelii* mirrored growth patterns observed for shoot height, with

maximum abundance at 24 °C representing a significant increase ($p = 0.033$), but was absent in *S.*

longipes except at 28 °C. Topolin riboside increased with increasing temperature for both species,

but was in greater abundance *S. longipes* than *C. regelii*, to a significant extent ($p = 0.0046$).

Figure 6. Effects of temperature on cytokinin metabolism for zeatin, dihydrozeatin, dihydrozeatin glucoside, zeatin-7-glucoside,

zeatin-9-glucoside, and topolin-riboside in Stellaria longipes and Cerastium regelii. For clarity, only the structures for trans-

zeatin-7-glucoside, trans-zeatin-9-glucoside and ortho-topolin riboside are depicted, but this analysis did not differentiate between

isomers. Shoot growth patterns are shown below for reference.

4.0 Discussion

Following a field survey of a permafrost anomaly in Resolute (Qausuittuq), Nunavut, we

established cultures of *S. longipes* and *C. regelii* for *in vitro* studies, along with field-collected

seeds. We performed experimental warming to test the effects of rising temperature in comparison

to field observation and predicted ecological niche modelling for both *S. longipes* and *C. regelii,*

focusing on impacts to reproductive and vegetative growth. We hypothesized that between these

related species, as a more generalist species with a greater distribution, *S. longipes* would be more

resilient to temperature increases than *C. regelii*, as evidenced by increased growth and a modified

phytohormone profile.

Field observations of the permafrost anomaly showed that for the plant community

established there, an increase in temperature shifted species composition towards herbaceous

plants including *S. longipes* and *C. regelii*, suggesting that increased warming created a more

suitable habitat for both species in these conditions. As species in this area are not currently in

competition with heavy shrub cover yet, our observations may provide insight into early

succession stages prior to shrubification associated with arctic greening, following tundra fires,

or following browning events with high mortality (Box *et al.*, 2019; Chen *et al.*, 2024; Phoenix *et*

al., 2025). Increased herbification with warming was largely supported by ecological niche

modeling for these species. Under SSP126 Resolute (Qausuittuq) is predicted to experience an

average temperature increase of ~5 °C. In these models *S. longipes* shows very little change in

habitat suitability across its current range, with only a slight Northward expansion. In contrast,

habitat suitability in current regions for *C. regelii* is predicted to significantly increase, although

its range will remain similar.

Within these models, the main contributors of environmental and bioclimatic variables to

model outcomes were different between species. Variables relating to temperature did not strongly

contribute to niche suitability, while habitat homogeneity and elevation was highly predictive of

niche suitability for *S. longipes*. In contrast, the main contributor to niche suitability for *C. regelii*

was the maximum temperature of the warmest month, highlighting a temperature-dependent

relationship of niche suitability and range expansion for *C. regelii* not predicted for *S.* longipes.

This is consistent with our hypothesis that *S. longipes*, as a generalist, would be less sensitive to

temperature change in moderate scenarios such as those explored here. This also suggests that

within these scenarios, increased warming may have a positive, rather than negative influence on

C. regelii. Phenology is the measure of the biological life cycles over an annual cycle, and includes

important events such as timing of bud break, flowering, fruiting, and dormancy (Morales-Castilla

et al., 2024). As these events are closely linked to environmental conditions, phenology has been

increasingly used as an indicator of climate change, but it is important to investigate this across a

broad range of species and climes as both phylogenetics and environment play a role in

phenological shifts (Diepstraten *et al.*, 2018; Morales-Castilla *et al.*, 2024). In arctic climes, the

effects of climate change are accelerated, and early loss of sea ice, increased temperature and

radiation, and a longer growing period are contributing to changes in flowering time, fruit and bud

set, along with pollinator emergence (Stewart *et al.*, 2018; Box *et al.*, 2019). We evaluated the

phenology of *S. longipes* and *C. regelii* over the past 170 years to evaluate how warming

temperatures affected the flowering period. In most species, the flowering period is compressed,

with phenological shifts towards earlier flowering times such as in *Dryas integrifolia* Vahl where

early flowering time has been linked to monthly temperature just prior to flowering (Panchen and

Root, 2015). We found that neither *S. longipes* nor *C. regelii* flowered earlier. *S. longipes*

experienced no change in flowering time. The number of plants in bloom observed increased with

time, but may reflect increased access to the area or similar observational bias during this time

period. *C. regelii* did undergo contraction of the flowering period from a bimodal distribution with

two blooming periods in July and September, to a unimodal distribution over a shorter flowering

period, but this also resulted in a general shift to later blooming time in late July to early August.

Based on niche suitability modelling and historical phenology, *C. regelii* exhibits a more

pronounced change in characteristics due to temperature than *S. longipes,* but contrary to our

hypothesis, this may actually improve abundance and suitability for this species within this region.

Based on the modelled results, we followed up with *in vitro* experiments to test the effects

of temperature increases on growth and phytohormone profiles of these two species

experimentally. We grew cultured *S. longipes* and *C. regelii* at 12 °C (a typical summer

temperature in the High Arctic), 24 °C (a typical summer temperature in the southern-most

latitudes where *S. longipes* is found), and 28 °C to model increased warming for these areas. An

increase in temperature from 12 to 24 °C, significantly increased the number of shoots and roots

produced in *S. longipes*, accompanied by increases in shoot and root length. Further increases of

temperature to 28 °C had the opposite effect, reducing these parameters to values observed at 12

°C. These patterns were also observed in *C. regelii* for the number and length of roots, and the
length of shoots, but the number of shoots increased linearly with increasing temperature. G rowth
trends were similar between these two species for most conditions, which may reflect on their
similar abundance and suitability at the temperatures observed in the permafrost anomaly. An
opposite trend in *C. regelii* associated with number of shoots shows that temperature does affect
some aspects of growth differently for *C. regelii* and *S. longipes*, which may have downstream
implications for their individual adaptability. *C. regelii* shows consistent increases in the number
of shoots it produces even at the highest temperature included in this study, while *S. longipes*
shows reduced number of shoots at these temperatures.

Given that plant growth is regulated by cross-talk between a growing number of
phytohormone classes, we theorized that differences in growth, such as in the number of shoots,
could be resolved by exploring differences in phytohormone production (Banoriya *et al.*, 2024;
Mughal *et al.*, 2024). We used metabolomics to broadly explore changes to metabolomic profile
and phytohormone production. The primary driver of variability in metabolites was species, but
following hierarchical analysis, we found that warming temperature has a strong influence on both
species. We had expected to see the greatest differences in metabolites at the temperatures which
were associated with the biggest increases in growth, 24 °C for *S. longipes*, and 28 °C for *C. regelii*.
What we observed in our results, however, was that the largest differences for *S. longipes* was
when it was warmed past the optimal temperature. Given that growth patterns tended to be self-
limiting with a reduction back to normal growth patterns at 28 °C, additional changes to the
metabolite profile are thus likely indicative of an early stress response. These results support the
predictions generated by ecological niche suitability modeling and phenology analysis, where *S.*
longipes was not predicted to change substantially in suitability for the region studied. The results
for *C. regelii* are more challenging to interpret. Although the responsiveness of *C. regelii* was
highest at 24 °C, we continued to see increased growth in the number of shoots beyond this
temperature. Based on these results, we hypothesized that the production of specific
phytohormones required for growth is differentially synthesized or regulated between these two
species, and is more influenced by temperature in *C. regelii* that is reflected in niche suitability
modelling.

Following the complex growth response of *C. regelii* to temperature, we explored the
abundance of phytohormones annotated from our metabolomics analysis. The strong effect of

experimental conditions on shoot growth and number highlights a relationship between
temperature and the production of cytokinins. Cytokinins are strongly associated with shoot
growth, and are known to stimulate cellular proliferation, vascular development, and biomass
distribution in addition to abiotic stress responses like heat (Li *et al.*, 2021). Zeatin, a major
cytokinin, was similarly produced by both *S. longipes* and *C. regelii* as temperature increased.
Zeatin is a major isoprenoid cytokinin produced by most plants as *cis-* or *trans-*zeatin, and is one
of the bioactive forms with strong bioactivity (Sakakibara, 2006). Other cytokinin derivatives
strongly differ between species. Dihydrozeatin, a bioactive zeatin derivative steadily increased in
abundance in *S. longipes* as temperature increased, but this coincided with an increase in
dihydrozeatin glucoside especially at 28 °C, widely considered to be an inactive form. *S. longipes*
also produced detectable quantities of two other inactive zeatin forms, zeatin-7-glucoside and
zeatin-9-glucoside. Collectively, the production of increased inactivated zeatin derivatives at the
highest temperature is likely to have a dampening effect on growth, supporting the observed
decrease in growth for *S. longipes* at this temperature. In *C. regelii,* the two isoprenoid cytokinins
detected were zeatin and the inactivated zeatin-7-glucoside. Zeatin-7-glucoside was most abundant
at 24 °C. At 28 °C, zeatin-7-glucoside decreased alongside a corresponding increase in zeatin, and
this may help explain the increased growth observed at this temperature. Lastly, both *S. longipes*
and *C. regelii* produced an abundance of topolin riboside, an aromatic cytokinin found in a limited
number of plant species (Zhao *et al.*, 2024). It was increased in abundance at 28 °C, although in
significantly greater abundance in *S. longipes*, and may suggest an alternative cytokinin production
pathway under heat stress. These results show that differences in these species as they react to
temperature increases are governed by alternative hormone signaling pathways. Further
investigations are needed to determine the influence of temperature on other hormone classes, and
given the differences in cytokinin abundance, what cross-talk may exist between them.

Collectively, our results show two strikingly different outcomes for these two herbaceous
species of the Arctic tundra. *S. longipes* shows very little influence of temperature on its niche
suitability or phenology, and may maintain vigor and robustness under the moderate future climate
scenarios explored here, supporting our hypothesis that as a generalist species it would be robust
to change at increased temperatures. In direct contrast, although *C. regelii* is more affected by
temperature, but for the conditions evaluated in our study is generally positively influenced with
warming, which is surprising given that it is more specialized with a smaller distribution and

predicted lower tolerance to change. Our *in vitro* investigations supported these predictions, showcasing that *S. longipes* is self-limiting with temperature changes, while increased temperatures tend to favor increased growth in *C. regelii.* Despite steady increases in cytokinin production that typically induce increased growth, *S. longipes* is self-limiting within the temperature ranges evaluated. Yet our work highlights the importance of studying climate change in the Arctic not only as a function of ecosystems and biomes, but also as a function of species, location, and metabolism. It is clear that even related species such as *S. longipes* and *C. regelii* can have different and unexpected responses to both moderate and extreme temperature increases as explored here. We also show the benefit and importance of using techniques like metabolomics to investigate these questions, helping us to evaluate how warming temperatures of the future will impact growth patterns and other aspects of plant biology for a variety of species with different responses to climate change. Our study also expands our knowledge and understanding of *C. regelii*, an understudied and important herbaceous plant in the Canadian High Arctic.

6.0 Acknowledgements

Funding support was provided by the Alpine Club of Canada Environment Grant, the Explorer's Club Mamont Scholar Grant and the Natural Sciences and Engineering Council of Canada to Dr. Erland. Logistical support was provided by the Polar Continental Shelf Program (PCSP) of the National Research Council of Canada (Project: 63319 – "ARCTIC Change - Arctic Research and Conservation Team Investigating Climate Change"). Guidance for this project was provided by staff and researchers at the PCSP facility in Resolute (Qausuittuq). We are grateful to Brandy Iqaluk, our guide in Resolute (Qausuittuq). We would also like to acknowledge undergraduate students at the University of the Fraser Valley who contributed to visual assessment of the phenology results Jiadong Li, Sebastian Molina, Elisha Marks and Oscar Caunce. Ryland Giebelhaus who assisted in maintenance of the initial cultures, and Sarah Lyons at Supra RnD Labs who assisted in acquisition of the metabolomics data. We are grateful to Paul Sokoloff at the Canadian Museum of Nature for his guidance in, and identification of the botanical vouchers.

Competing Interests

The authors declare no competing interests

Author Contributions

L.A.E Erland performed experimental work, data analysis, writing, and editing of manuscript. S.L. Lane performed data analysis for metabolomics experiments, writing, and editing of manuscript.

Data Availability

Metabolomics data can be found on Metabolomics Workbench, Project ID (unassigned, submission under review).

Supplementary information below.

References

Ai-Qun, J.I.A. *et al.* (2004) 'Cyclopeptides from Three Arctic Caryophyllaceae Plants, Chemotaxonomy and Distribution Significance of Caryophyllaceae Cyclopeptides', *Journal of Integrative Plant Biology*, 46(5), p. 625.

Amatulli, G. *et al.* (2018) 'A suite of global, cross-scale topographic variables for environmental and biodiversity modeling', *Scientific Data*, 5(1), p. 180040. Available at: https://doi.org/10.1038/sdata.2018.40.

Banoriya, R. *et al.* (2024) 'Salicylic Acid: A Key Player in the Regulation of Interactions Between Heat Stress and Plant Defense', *Journal of Plant Growth Regulation*, pp. 1–25. Available at: https://doi.org/10.1007/s00344-024-11586-3.

Box, J.E. *et al.* (2019) 'Key indicators of Arctic climate change: 1971–2017', *Environmental Research Letters*, 14(4), p. 045010. Available at: https://doi.org/10.1088/1748-9326/aafc1b.

Chen, D. *et al.* (2024) 'Regional fire–greening positive feedback loops in Alaskan Arctic tundra', *Nature Plants*, 10(12), pp. 1886–1891. Available at: https://doi.org/10.1038/s41477-024-01850-5.

Chinnappa, C.C. *et al.* (2005) 'The biology of *Stellaria longipes* (Caryophyllaceae)', *Canadian Journal of Botany*, 83(11), pp. 1367–1383. Available at: https://doi.org/10.1139/b05-117.

Dawson, T.P. *et al.* (2011) 'Beyond Predictions: Biodiversity Conservation in a Changing Climate', *Science*, 332(6025), pp. 53–58. Available at: https://doi.org/10.1126/science.1200303.

Diepstraten, R.A.E. *et al.* (2018) 'Does climate change and plant phenology research neglect the Arctic tundra?', *Ecosphere*, 9(9), p. e02362. Available at: https://doi.org/10.1002/ecs2.2362.

Emery, R.J.N., Chinnappa, C.C. and Chmielewski, J.G. (1994) 'Specialization, Plant Strategies, and Phenotypic Plasticity in Populations of *Stellaria longipes* along an Elevational Gradient', *International Journal of Plant Sciences*, 155(2), pp. 203–219. Available at: https://doi.org/10.1086/297160.

Erland, L.A.E., Saxena, P.K. and Murch, S.J. (2017) 'Melatonin in plant signalling and behaviour', *Functional Plant Biology*, 45(2), pp. 58–69. Available at: https://doi.org/10.1071/FP16384.

Erland, L.A.E., Turi, C.E. and Murch, S.J. (2022) 'Preliminary assessment of the conservation status of medicinal plant species in Canada', *Botany*, 100(2), pp. 247–260. Available at: https://doi.org/10.1139/cjb-2021-0090.

Fick, S.E. and Hijmans, R.J. (2017) 'WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas', *International Journal of Climatology*, 37(12), pp. 4302–4315. Available at: https://doi.org/10.1002/joc.5086.

Foden, W.B. *et al.* (2013) 'Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and Corals', *PLOS ONE*, 8(6), p. e65427. Available at: https://doi.org/10.1371/journal.pone.0065427.

Fredskild, B. (1984) 'Distribution and occurrence of Onagraceae in Greenland', *Nordic Journal of Botany*, 4(4), pp. 475–480. Available at: https://doi.org/10.1111/j.1756-1051.1984.tb02051.x.

Gamborg, O.L., Miller, R.A. and Ojima, K. (1968) 'Nutrient requirements of suspension cultures of soybean root cells', *Experimental Cell Research*, 50(1), pp. 151–158. Available at: https://doi.org/10.1016/0014-4827(68)90403-5.

Giebelhaus, R.T., Erland, L.A.E. and Murch, S.J. (2024) 'HormonomicsDB: a novel workflow for the untargeted analysis of plant growth regulators and hormones', *F1000Research*, 11, p. 1191. Available at: https://doi.org/10.12688/f1000research.124194.2.

Hanssen, O. (1932) *Flowering plants of Franz Josef Land : collected on the Norwegian Scientific Expedition 1930*. Oslo: I kommisjon hos J. Dybwad. Available at: https://search.library.wisc.edu/catalog/999520306202121 (Accessed: 14 March 2025).

Heide, O.M., Pedersen, K. and Dahl, E. (1990) 'Environmental control of flowering and morphology in the high-arctic *Cerastium regelii*, and the taxonomic status of *C. jenisejense*', *Nordic Journal of Botany*, 10(2), pp. 141–147. Available at: https://doi.org/10.1111/j.1756-1051.1990.tb01761.x.

Hirabayashi, K., Murch, S.J. and Erland, L.A.E. (2022) 'Predicted impacts of climate change on wild and commercial berry habitats will have food security, conservation and agricultural implications', *Science of The Total Environment*, 845, p. 157341. Available at: https://doi.org/10.1016/j.scitotenv.2022.157341.

Kass, J.M. *et al.* (2021) 'ENMeval 2.0: Redesigned for customizable and reproducible modeling of species' niches and distributions', *Methods in Ecology and Evolution*, 12(9), pp. 1602–1608. Available at: https://doi.org/10.1111/2041-210X.13628.

Kathiresan, A., Reid, D.M. and Chinnappa, C.C. (1996) 'Light- and temperature-entrained circadian regulation of activity and mRNA accumulation of 1-aminocyclopropane-1-carboxylic

acid oxidase in Stellaria longipes', *Planta*, 199(3), pp. 329–335. Available at:
https://doi.org/10.1007/BF00195723.

Kew Science (no date) *Cerastium jenisejense Hultén*, *Plants of the World Online*. Available at:
http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:152328-1 (Accessed: 14 March
2025).

Kodl, G. *et al.* (2024) 'Arctic tundra shrubification can obscure increasing levels of soil erosion
in NDVI assessments of land cover derived from satellite imagery', *Remote Sensing of
Environment*, 301, p. 113935. Available at: https://doi.org/10.1016/j.rse.2023.113935.

Kurepin, L.V. *et al.* (2006) 'Involvement of gibberellins in the stem elongation of sun and shade
ecotypes of *Stellaria longipes* that is induced by low light irradiance', *Plant, Cell &
Environment*, 29(7), pp. 1319–1328. Available at: https://doi.org/10.1111/j.1365-
3040.2006.01512.x.

Kurepin, L.V. *et al.* (2008) 'Light irradiance differentially regulates endogenous levels of
cytokinins and auxin in alpine and prairie genotypes of *Stellaria longipes*', *Physiologia
Plantarum*, 134(4), pp. 624–635. Available at: https://doi.org/10.1111/j.1399-
3054.2008.01163.x.

Li, S.-M. *et al.* (2021) 'Cytokinins as central regulators during plant growth and stress response',
Plant Cell Reports, 40(2), pp. 271–282. Available at: https://doi.org/10.1007/s00299-020-02612-
1.

Li, W. *et al.* (2011) 'Effects of day length, light quality and ethylene on PHYTOCHROME B
expression during stem elongation in *Stellaria longipes*', *Plant Growth Regulation*, 63(3), pp.
291–300. Available at: https://doi.org/10.1007/s10725-010-9529-y.

Liu, C. *et al.* (2023) 'Phylogenetic relationships and diversification dynamics of *Cerastium*
(Alsineae: Caryophyllaceae): implications for biogeographical patterns of herbs in the Northern
Hemisphere', *Botanical Journal of the Linnean Society*, 203(1), pp. 67–77. Available at:
https://doi.org/10.1093/botlinnean/boad018.

Macdonald, S.E., Chinnappa, C.C. and Reid, D.M. (1984) 'Studies on the *Stellaria longipes*
complex: phenotypic plasticity. I. Response of stem elongation to temperature and photoperiod',
Canadian Journal of Botany, 62(3), pp. 414–419. Available at: https://doi.org/10.1139/b84-063.

Macdonald, S.E., Reid, D.M. and Chinnappa, C.C. (1986) 'Studies on the *Stellaria longipes*
complex: phenotypic plasticity. II. Gibberellins, abscisic acid, and stem elongation', *Canadian
Journal of Botany*, 64(11), pp. 2617–2621. Available at: https://doi.org/10.1139/b86-346.

Maillette, L. *et al.* (2000) 'Module demography does not mirror differentiation among
populations of *Stellaria longipes* along an elevational gradient', *Plant Ecology*, 149(2), pp. 143–
156. Available at: https://doi.org/10.1023/A:1026525106640.

Matveyeva, N.V. (1994) 'Floristic Classification and Ecology of Tundra Vegetation of the Taymyr Peninsula, Northern Siberia', *Journal of Vegetation Science*, 5(6), pp. 813–828. Available at: https://doi.org/10.2307/3236196.

Morales-Castilla, I. *et al.* (2024) 'Phylogenetic estimates of species-level phenology improve ecological forecasting', *Nature Climate Change*, 14(9), pp. 989–995. Available at: https://doi.org/10.1038/s41558-024-02102-2.

Mughal, N. *et al.* (2024) 'Adaptive roles of cytokinins in enhancing plant resilience and yield against environmental stressors', *Chemosphere*, 364, p. 143189. Available at: https://doi.org/10.1016/j.chemosphere.2024.143189.

Murashige, T. and Skoog, F. (1962) 'A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures', *Physiologia Plantarum*, 15(3), pp. 473–497. Available at: https://doi.org/10.1111/j.1399-3054.1962.tb08052.x.

Muscarella, R. *et al.* (2014) 'ENMeval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models', *Methods in Ecology and Evolution*, 5(11), pp. 1198–1205. Available at: https://doi.org/10.1111/2041-210X.12261.

Myers-Smith, I.H. *et al.* (2011) 'Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities', *Environmental Research Letters*, 6(4), p. 045509. Available at: https://doi.org/10.1088/1748-9326/6/4/045509.

Panchen, Z.A. and Root, G. (2015) 'Flowering and fruiting responses to climate change of two Arctic plant species, purple saxifrage (*Saxifraga oppositifolia*) and mountain avens (*Dryas integrifolia*)', *Arctic Science* [Preprint]. Available at: https://doi.org/10.1139/as-2015-0016.

Parding, K.M. *et al.* (2020) 'GCMeval – An interactive tool for evaluation and selection of climate model ensembles', *Climate Services*, 18, p. 100167. Available at: https://doi.org/10.1016/j.cliser.2020.100167.

Phoenix, G.K. *et al.* (2025) 'Browning events in Arctic ecosystems: Diverse causes with common consequences', *PLOS Climate*, 4(1), p. e0000570. Available at: https://doi.org/10.1371/journal.pclm.0000570.

Pluskal, T. *et al.* (2010) 'MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data', *BMC Bioinformatics*, 11(1), p. 395. Available at: https://doi.org/10.1186/1471-2105-11-395.

Poggio, L. *et al.* (2021) 'SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty', *SOIL*, 7(1), pp. 217–240. Available at: https://doi.org/10.5194/soil-7-217-2021.

Porsild, A.E., Porsild, A.E. and Cody, W.J. (1980) *Vascular plants of continental Northwest Territories, Canada*. Ottawa: National Museum of Natural Sciences, National Museums of Canada, pp. 1–690. Available at: https://doi.org/10.5962/bhl.title.70336.

Rantanen, M. *et al.* (2022) 'The Arctic has warmed nearly four times faster than the globe since 1979', *Communications Earth & Environment*, 3(1), pp. 1–10. Available at: https://doi.org/10.1038/s43247-022-00498-3.

Regel's chickweed (no date) *iNaturalist*.

Sakakibara, H. (2006) 'CYTOKININS: Activity, Biosynthesis, and Translocation', *Annual Review of Plant Biology*, 57(1), pp. 431–449. Available at: https://doi.org/10.1146/annurev.arplant.57.032905.105231.

Sangtarash, M.H. *et al.* (2009) 'Differential responses of two *Stellaria longipes* ecotypes to ultraviolet-B radiation and drought stress', *Flora - Morphology, Distribution, Functional Ecology of Plants*, 204(8), pp. 593–603. Available at: https://doi.org/10.1016/j.flora.2008.08.004.

Schmid, R. *et al.* (2023) 'Integrative analysis of multimodal mass spectrometry data in MZmine 3', *Nature Biotechnology*, 41(4), pp. 447–449. Available at: https://doi.org/10.1038/s41587-023-01690-2.

Sharples, M.T. (2019) 'Taxonomic Observations within *Stellaria* (Caryophyllaceae): Insights from Ecology, Geography, Morphology, and Phylogeny Suggest Widespread Parallelism in Starworts and Erode Previous Infrageneric Classifications', *Systematic Botany*, 44(4), pp. 877–886. Available at: https://doi.org/10.1600/036364419X15710776741459.

Shiogama, H. *et al.* (2021) 'Selecting CMIP6-Based Future Climate Scenarios for Impact and Adaptation Studies', *Sola*, 17, pp. 57–62. Available at: https://doi.org/10.2151/sola.2021-009.

Shiogama, H., Abe, M. and Tatebe, H. (2019) 'MIROC MIROC6 model output prepared for CMIP6 ScenarioMIP ssp245'. Earth System Grid Federation. Available at: https://doi.org/10.22033/esgf/cmip6.5746.

Stellaria longipes s. lat. (no date). Available at: https://www.svalbardflora.no/index.php/stellaria/stellaria-longipes-s-lat (Accessed: 14 March 2025).

Stewart, L. *et al.* (2018) 'Forecasted homogenization of high Arctic vegetation communities under climate change', *Journal of Biogeography*, 45(11), pp. 2576–2587. Available at: https://doi.org/10.1111/jbi.13434.

Thuy, D.T.T., Yeung, E.C. and Chinnappa, C.C. (2009) 'Differences in cellular structure regulate stem elongation of alpine and prairie ecotypes of *Stellaria longipes* grown under two contrasting temperature regimes', *Botany*, 87(12), pp. 1167–1176. Available at: https://doi.org/10.1139/B09-077.

Tuanmu, M.-N. and Jetz, W. (2014) 'A global 1-km consensus land-cover product for biodiversity and ecosystem modelling', *Global Ecology and Biogeography*, 23(9), pp. 1031–1045. Available at: https://doi.org/10.1111/geb.12182.

Tuanmu, M.-N. and Jetz, W. (2015) 'A global, remote sensing-based characterization of
terrestrial habitat heterogeneity for biodiversity and ecosystem modelling', *Global Ecology and*
Biogeography, 24(11), pp. 1329–1339. Available at: https://doi.org/10.1111/geb.12365.

Vignali, S. *et al.* (2020) 'SDMtune: An R package to tune and evaluate species distribution
models', *Ecology and Evolution*, 10(20), pp. 11488–11506. Available at:
https://doi.org/10.1002/ece3.6786.

Voldoire, A. *et al.* (2019) 'Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1',
Journal of Advances in Modeling Earth Systems, 11(7), pp. 2177–2213. Available at:
https://doi.org/10.1029/2019MS001683.

Yao, G. *et al.* (2021) 'Phylogenetic estimation and morphological evolution of Alsineae
(Caryophyllaceae) shed new insight into the taxonomic status of the genus *Pseudocerastium*',
Plant Diversity, 43(4), pp. 299–307. Available at: https://doi.org/10.1016/j.pld.2020.11.001.

Yukimoto, S. *et al.* (2019) 'MRI MRI-ESM2.0 model output prepared for CMIP6 CMIP'. World
Data Center for Climate (WDCC) at DKRZ. Available at: https://www.wdc-
climate.de/ui/entry?acronym=C6_5243961 (Accessed: 14 March 2025).

Zhao, J. *et al.* (2024) 'Spatially distributed cytokinins: Metabolism, signaling, and transport',
Plant Communications, 5(7), p. 100936. Available at:
https://doi.org/10.1016/j.xplc.2024.100936.

Zorzi, A., Deyle, K. and Heinis, C. (2017) 'Cyclic peptide therapeutics: past, present and future',
Current Opinion in Chemical Biology, 38, pp. 24–29. Available at:
https://doi.org/10.1016/j.cbpa.2017.02.006.

 Supporting Information

843   Table S1. Summary of historic climate variables assessment for inclusion in the model

Variable	Variable Shorthand	Source/Type
Annual Mean Temperature (°C)	bio1	WorldClim
Maximum Temperature Warmest Month (°C)	bio5	WorldClim
Minimum Temperature Coldest Month (°C)	bio6	WorldClim
Mean Temperature of Wettest Quarter (°C)	bio8	WorldClim
Mean Temperature of Driest Quarter (°C)	bio9	WorldClim
Annual Precipitation (mm)	bio12	WorldClim
Precipitation Wettest Quarter (mm)	bio16	WorldClim
Precipitation Driest Quarter (mm)	bio17	WorldClim
Precipitation Warmest Quarter (mm)	bio18	WorldClim
Precipitation Coldest Quarter (mm)	bio19	WorldClim
Aspect Sine	aspectsine	topography
Aspect Eastness	eastness	topography
Aspect Northness	northness	topography
Geomorphological landform (Shannon Index)	geom	topography
Slope	slope	topography
Cultivated & Managed Vegetation (% consensus prevalence)	cultivatedmanagedveg	landcover DIScover
Regularly Flooded Land (% consensus prevalence)	regflooded	landcover DIScover
Snow & Ice Cover (% consensus prevalence)	snowice	landcover DIScover

Homogeneity	homogeneity	habitat heterogeneity
Entropy	entropy	habitat heterogeneity
Dissimilarity	dissimilarity	habitat heterogeneity
Evenness	evenness	habitat heterogeneity
Co-efficient of Variation	coeffvar	habitat heterogeneity
Soil Organic Carbon Densities (hg/m3)	ocd	SoilGrids
Soil Organic Carbon Stock (t/ha)	ocs	SoilGrids
Soil Organic Carbon Content (dg/kg)	soc	SoilGrids
Silt (g/kg)	silt	SoilGrids
Sand (g/kg)	sand	SoilGrids
Nitrogen (cg/kg)	nitrogen	SoilGrids
Clay (g/kg)	clay	SoilGrids
Bulk Density of Fine Earth Fraction (cg/cm3)	bdod	SoilGrids
Volumetric fraction of coarse fragments > 2mm (cm2/dm3 / vol%)	cfvo	SoilGrids
Cation exchange capacity of the soil (mmol(c)/kg)	cec	SoilGrids
Soil pH (pH x 10)	phh2o	SoilGrids

846 *Table S2. Model performance statistics for the null and empirical models generated for ecological niche modelling of Cerastium*
847 *regelii and Stellaria longipes.*

Statistic	AUC		Minimum Training Presence Omission Rate		10 Percent Omission Rate	
	null	empirical	null	empirical	null	empirical
Cerastium regelii RM = 4, Features = LQH						
mean	0.558	0.804	0.000	0.000	0.105	0.103
standard deviation	0.032	0.054	0.001	0.000	0.030	0.091
zscore	7.604		-0.118		-0.060	

pvalue	<0.0001		0.453		0.476
Stellaria longipes RM = 5, Features = LQH					
mean	0.797	0.865	0.000	0.000	0.034	0.083
standard deviation	0.026	0.039	0.002	0.000	0.019	0.052
z-score	2.654		-0.188		2.526	
p-value	0.004		0.425		0.994	

Table S3. Maxent ecological niche model performance characteristics and variable contribution for variables retained in the model for Cerastium regelii and Stellaria longipes.

	Stellaria longipes		*Cerastium regelii*	
Variable	Percent Contribution	Permutation Importance	Percent Contribution	Permutation Importance
bdod	0.0193	0.9225	1.1442	2.8595
bio1	0.4998	0.1934	0	0
bio12	2.8942	0.5572	0	0
bio16	3.2335	0.6501	0.1314	0.0036
bio17	0.5264	7.1554	0	0
bio18	0.2939	4.4969	0	0
bio19	0.0088	0.0071	0.0522	0.0044
bio5	0.0295	0.0794	39.4382	42.7297
bio6	5.3681	0.9594	0	0
bio8	0.5832	0.917	0	0
bio9	0.5934	5.0889	0.0527	0.0285
cec	0.0199	0	0.0716	0.8538
coeffvar	4.98	0	14.9568	0
dissimilarity	0.0408	0	1.6561	0
elevation	30.906	25.3389	26.4285	16.868
entropy	0.8703	0	0.9293	0
evenness	1.4589	1.1872	0.4248	15.1271
geom	0.0111	0.0864	0	0
homogeneity	46.0241	38.5921	6.6315	19.4896
nitrogen	0.1316	4.3129	0.3514	0.5358
northness	0.0204	0.6623	4.00E-04	0
ocd	0.1348	5.6983	0.8418	0.781

ocs	0	0	0.0064	0.0372
openwater	0.3542	0	3.8697	0.1823
phh2o	0	0	0.2547	0
silt	0	0	0.2982	0
slope	0.6206	1.008	0.011	0.0617
snowice	0.3013	1.4626	2.4147	0.0176
soc	0.0211	0.447	0	0
TSS Train	0.528		0.777	
TSS test	0.511		0.73	
TSS heldback	0.546		0.81	
AICc	59554.01		31389.84	
AUC	0.782		0.929	

Tree scale:

Figure S1. Phylogenetic tree of field collections from 2019 based on DNA fingerprinting data uploaded to BOLD and generated

in PhyloT. Species of interest are highlighted in purple.

Figure S2. Predicted change in ecological niche suitability for *Cerastium regelii* under current and

future climates. Inset shows a closer view of the Canadian High Arctic where *C. regelii* is known

to occur.

Figure S3. Predicted change in ecological niche suitability for *Stellaria longipes* under current and

future climates

868