Synteny – a high throughput web tool to streamline causal gene prioritisation and provide insight into protein function

Harry B. Cutler[1,2], Jacqueline Stöckli[1,2], Søren Madsen[1,2], Stewart W.C. Masson[1,2], Oliver K. Fuller[1,2], Tom Buckingham Shum[2], David E. James[1,2,3,4]

[1] School of Life and Environmental Sciences, University of Sydney, Camperdown, New South Wales, Australia.
[2] Charles Perkins Centre, University of Sydney, Camperdown, New South Wales, Australia.
[3] School of Medical Sciences, University of Sydney, Camperdown, New South Wales, Australia.
[4] Correspondence: david.james@sydney.edu.au

Summary

Accounting for human genetic evidence can improve the outcomes and impact of basic research studies. However, current approaches are incompatible with the high volume of disease-associated genes that require mechanistic interrogation. *Synteny* responds to an urgent need in systems biology, scaling human genetic analysis to match the throughput of modern omics technologies. This approach prioritises candidates with the strongest human disease relevance and unearths functionally important proteins (https://bigproteomics.shinyapps.io/Synteny/).

Keywords

Disease-associated genes, candidate prioritisation, causal genes, human genetic data, proteomics, transcriptomics, systems biology

Introduction

Advances in omics-technologies have revolutionised the study and understanding of human disease. However, the complexity of the data generated by these techniques can obscure key biological insights.

A major challenge is that omics studies frequently identify hundreds of disease-associated genes, many of which remain poorly characterised. Even a comprehensive review of the literature often provides insight into just a small subset of these genes[1–3]. Combined with the fact that data analysis approaches like differential expression offer little insight into whether changes are causes or consequences of disease processes, researchers are often drawn to genes with known functions[3,4]. This 'streetlight' bias – searching for answers where the light is brightest rather than venturing into the relative unknown – impedes new breakthroughs and signals an unmet need for robust and unbiased tools to prioritise genes with the strongest causal links to human disease.

One approach that is becoming popular is to integrate omics data with findings from human genetics, since genetic variants are causal by definition. Genetic and phenotypic data are now available across large human cohorts and several studies using model systems have demonstrated success in applying such an integrated approach[5–7]. For example, to pinpoint causal genes contained within human GWAS loci, Li et al[6] integrated genetic associations from a GWAS of human lipid metabolism with gene co-expression networks from mouse liver samples. This identified 54 genes with conserved function in mice and humans, of which 25 had not previously been linked to lipid metabolism. It also motivated experiments to validate the role of Sestrin 1 in regulating cholesterol biosynthesis, identifying a previously unknown therapeutic target. Examples like this emphasise the importance of bridging the

gap between studies in humans and model systems. However, the complexity of current
approaches provides a considerable challenge to those without specialist bioinformatic
training. For example, Keller et al[5] devised an approach to determine whether loci identified
in their mouse GWAS of pancreatic islet function were conserved in humans by mapping
syntenic regions between mouse and human GWAS and then testing for overlap with human
GWAS loci. Although very rigorous, this approach is technically challenging and only
accessible to those with access to genetically diverse mouse populations.
62
To address these challenges, we developed *Synteny*, a user-friendly web tool that enables
anybody with proteomic or transcriptomic data to leverage large-scale human genetic
datasets to rank genes by human disease relevance. In contrast to current gene-
prioritisation platforms that are limited to searching genes one-at-a-time, such as
OpenTargets Genetics[8] and the Type 2 Diabetes Knowledge Portal[9], *Synteny* was built to
perform analyses at scale – data for several hundred genes can be acquired in minutes.
Synteny automatically maps human orthologs from 8 common model organisms using the
Alliance for Genome Resources database[10], before acquiring humans phenome-wide
association statistics from the Association to Function (A2F) database
(https://a2f.hugeamp.org/). Cross-species orthologs are typically highly conserved in terms
of structure and function[11] and so this approach enables findings in model organisms to be
leveraged to infer potential human disease mechanisms and gene function, and vice-versa.
As of June 2025, A2F reports summary statistics for 898 genetic studies and 1175
phenotypes, enabling unbiased assessment of gene function.
77
78 **Synteny powers improved access to human genetic evidence**
79
While technical barriers once limited the ability to integrate human genetic data into broader
biological analyses, recent approaches such as the Human Genetic Evidence (HuGE)
framework by Dornbos et al [12] have greatly expanded access to these insights. The main
benefit of HuGE scores is that they provide a single interpretable measure of the strength of
association between every gene and every phenotype. They are calculated using a
Bayesian model to account for the effects of rare and common genetic variation, the
proximity of a gene to the peak genetic association in genome wide association studies and
the impact of coding mutations detected in whole exome sequencing[12]. By aggregating
genetic data across many independent studies, HuGE scores provide an unbiased and
robust assessment of how strongly a gene of interest might relate to human physiology. For
example, HuGE scores for peroxisome proliferator activated receptor gamma (*PPARG*), a
master-regulator of adipogenesis, show strong associations (HuGE ≥ 30) to body fat
percentage and muscle fat infiltration (Fig 1A), consistent with its established role in
adipogenesis. While databases like A2F have made it straightforward to access HuGE
scores for individual genes, the restriction on searching genes one-at-a-time is incompatible
with high-throughput, hypothesis-generating studies where hundreds of disease-associated
genes can be identified in a single experiment (Fig 3B).
97
98 *Synteny* overcomes this limitation by enabling bulk querying of the A2F database (Fig 1C),
99 streamlining the process of integrating human genetic evidence with data from model
100 systems. Its operation is simple, requiring only uploading a list of genes. All results can be
101 exported for analysis in external software and data visualisation plots can be downloaded as
102 high-resolution vector PDFs.
103
To demonstrate the efficiency of this approach, we used *Synteny* to analyse a list of 52
genes that Dornbos et al[12] previously used to assess the use of human genetic evidence by
metabolism researchers working with model systems. Dornbos et al[12] selected these genes
from studies published in high profile journals between January 2017 and October 2020 that
contained any of "diabetes," "glucose," or "insulin" in their abstracts. Examining the HuGE
scores for these genes, Dornbos et al[12] identified a striking lack of genetic evidence to

support therapeutic targeting of these genes, highlighting an essential need for increased
use of human genetic data. However, without efficient tools to query human genetic
databases, it remains challenging to ensure that research is targeted towards genes with the
strongest links to human disease.
114
115   While manually querying each gene using the A2F database required more than an hour,
116   using *Synteny* this process was completed in just 80 seconds (Fig 1D). This boost in
117   efficiency makes this approach much more accessible than existing alternatives, enhancing
118   the potential for increased translatability.
119
Investigation of HuGE scores across genes frequently reveals well-studied associations,
however, the true value of this analysis lies in highlighting relationships that would otherwise
have gone unnoticed. For example, while the link between phosphatase and tensin homolog
(*PTEN*) and systemic glucose homeostasis is well-established[13,14], we were surprised by the
association between carnitine palmitoyltransferase I (*CPT1A*) and platelet number. CPT1A
plays a central role in transporting fatty acids into the mitochondria for oxidation[15,16] and,
consistent with human genetic evidence, recent studies have revealed that the contribution
of fatty acids to ATP production in platelets is likely greater than currently appreciated[17,18].
These insights suggest new avenues for exploring platelet pathology and showcase the
power of unbiased analyses.
130
The Human Phenotype Enrichment (HPE) tool in *Synteny* enables users to quantify the
extent to which gene sets of interest are supported by human genetic evidence. Similar to
existing over-representation analyses for molecular functions, HPE works by comparing
HuGE scores for a user-supplied gene set against those obtained by randomly sampling the
genome (or a user-supplied background gene set). Consistent with the findings of Dornbos
et al[12] , that their list of 52 genes was overall not supported by human genetic evidence,
HPE using *Synteny* failed to detect significant enrichment for glycaemic traits that would be
expected if these genes were supported by human genetic evidence (Fig 1E). Instead, the
52 genes were enriched for early life Body Mass Index (BMI), Alzheimer's disease (AD) and
age-related macular degeneration (AMD). This outcome supports the conclusions of
Dornbos et al[12], emphasising the importance of interrogating results in the context of human
genetics to maximise the extent to which advances in basic research can be translated into
human therapeutic breakthroughs. Many existing approaches using human genetic data are
designed to test single phenotypes and so here we report both unadjusted and FDR-
corrected p values to provide compatibility with these approaches.
146
To evaluate the sensitivity and robustness of HPE, we performed a series of benchmarking
experiments (Fig 2). To determine the degree of enrichment required to detect statistical
significance, power analyses were performed using gene sets of varying size between 10-
300 genes, obtained by weighted sampling of the HuGE score distribution (Fig 2A: i). Using
obesity as an example trait, 90% power was obtained at 4-fold enrichment with 100 genes
(Fig 2A: ii). To assess sensitivity more broadly, we extended this analysis to 15 randomly
sampled human traits from 16 trait categories (240 phenotypes in total). Except for
haematological traits that required a median 40-fold enrichment to reach 80% power, the
other 225 traits were broadly consistent, requiring a median enrichment of 1.66-fold (Fig 2B).
These results suggest that HPE is highly sensitive for phenome-wide enrichment analysis.
157
To assess the robustness of HPE against varying gene set compositions, we used HPE to
analyse several systematically curated HALLMARK pathways (Fig 2C). This identified
enrichments that are consistent with known phenotypic associations. For example, the
'Adipogenesis' pathway was enriched for adiponectin (an adipokine), body fat percentage
and plasma triglyceride (TG) levels; the 'Pancreatic β cells' pathway was enriched for fasting
glucose and traits related to insulin secretion, a key function of β cells; and the 'Inflammatory
response' pathway was enriched for Crohn's disease and coronary artery disease, both well-

studied inflammatory conditions. These results suggest that HPE can infer true biological
meaning from user-supplied gene sets.
167
HPE aggregates enrichment signals over entire gene sets, and it is therefore important to
determine whether findings are driven by a small number of genes or are distributed across
entire the gene set. *Synteny* facilitates this by automatically visualising the contributions of
user-supplied genes to an enriched phenotype. For example, while the enrichment for
adiponectin in the 'Adipogenesis' gene set seems to be primarily driven by the inclusion of
the adiponectin (ADIPOQ) gene, the Crohn's disease enrichment for the 'Inflammatory
response' pathway is more evenly distributed.
175
Enabling users to determine which genes contribute most strongly to human traits also
addresses a core challenge in omics analysis. Genes belonging to the same pathway are
often co-regulated[19], and this can obscure which genes causally influence the phenotype.
For example, although omics analysis might identify several glycolytic enzymes as hits, it
may be unclear which gene/s should be pursued experimentally. This is important since,
while there are 12 steps in glycolysis – from glucose import to lactate export – only 4 of
these appear to regulate total flux through the pathway[20]. Selecting genes that do not
regulate flux might result in a false negative result. *Synteny* addresses this challenge by
providing access to the effects of human genetic variation. Because the effects of genetic
variation are independent of pathway membership, this enables users to disentangle the
contributions of individual genes to human disease. Applied to the analysis of glycolytic
genes expressed in skeletal muscle, *Synteny* identified very strong genetic evidence (HuGE
≥ 30) for several enzymes that Tanner et al[20] previously showed to regulate total glycolytic
flux (Fig 2D). Thus, *Synteny* provides a powerful tool to assist users in identifying high-
priority genes for mechanistic follow-up.
191

192
193
Figure 1 – Synteny powers improved access to human genetic evidence. (A) Human
genetic evidence (HuGE) scores aggregate genetic data from many studies to summarise
the strength of association between genes and phenotypes[12], enabling gene function to be
rapidly assessed. Example of HuGE score plot linking *PPARG* to 1175 phenotypes. **(B)**
Schematic of the current implementation of the HuGE score framework via the Association to
Function web portal. **(C)** Schematic illustrating the high throughput implementation in
Synteny. API = application programming interface. Examples of data visualisation and
analysis in *Synteny* for 52 genes hypothesised to be linked to glucose metabolism[12] **(D & E)**.

(D) Heatmap showing associations between genes and human phenotypes, filtered for
genes with HuGE scores ≥ 30. **(E)** Results of HPE. Asterisks represent unadjusted p values.
204

205
206
Figure 2 – Benchmarking for human phenotype enrichment. (A) Example of sensitivity
analysis performed using Obesity as a model trait. Random weighted sampling (i) was used
to generate enriched gene sets with different numbers of genes that were then subjected to
HPE (ii). **(B)** Power simulations for 15 randomly sampled phenotypes from 16 trait
categories, with gene sets of size n = 100. Each dot is the average of 1000 permutations per
phenotype, showing the degree of enrichment required to reach 80% power. Median
enrichment for 80% power ($\tilde{x}$) = 1.66-fold. **(C)** HPE using gene sets from HALLMARK
pathways. Genes contributing to the top enrichments for 'Adipogenesis', 'Pancreatic β cells'
and 'Inflammatory response' are shown in i-iii, respectively. **(D)** Heatmap presenting HuGE

scores for glycolytic enzymes expressed in skeletal muscle for selected phenotypes.
Enzymes that regulate total glycolytic flux[20] are highlighted in red.
218
219 **Synteny leverages human genetic data for discovery**
220
221 Less than half of all human genes have been studied mechanistically[4,21,22]. To evaluate
222 whether *Synteny* could be used to infer protein function, we applied it to the solute carrier
223 (SLC) superfamily, containing over 400 genes. We chose this family since, despite its
224 members playing crucial roles in many aspects of biology, the substrates and biological
225 functions of many of these proteins remain unknown[23].
226
227 Consistent with their roles as transporters, HPE analysis identified enrichment of the SLC
228 family in regulating circulating levels of a variety of metabolites including creatinine, urate
229 and sodium (Fig 3A). Several human diseases related to dysregulated ion transport,
230 including chronic kidney disease and epilepsy with sclerosis were also enriched.
231
To identify putative functions for orphan SLC members – SLCs without known substrates –
we identified SLCs with similar associations to human traits by correlating HuGE scores
across genes. Such guilt-by-association strategies are implemented in several established
tools including WGCNA[24], GeneBridge[22] and QENIE[25], all of which assign putative functions
to genes by means of correlations or co-expression networks. This identified 6 clusters
containing more than 2 SLC proteins (Fig 3B). Each SLC cluster was enriched for at least
one human trait (Fig 3C). Strikingly, cluster 4 was enriched for genes associating with serum
urate levels and 3 of the 5 genes in this cluster (*SLC2A9*, *SLC22A12*, *SLC22A13*) have
previously been shown to actively transport urate[26–28]. This raised the intriguing possibility
that the remaining two 2 SLC members in this cluster (*SLC5A2* and *SLC5A9*) may also
function as urate transporters.
243
SLC5A2 encodes the SGLT2 protein, a sodium-dependent glucose transporter highly
expressed in the kidney and a well-established target for Type 2 Diabetes[29,30]. Both genetic
and pharmacological evidence support a potential role of *SLC5A2* in urate transport,
although this has not been directly tested yet. Specifically, a coding variant in *SLC5A2* has
been linked to uricosuria[31], a condition with high urate levels in the urine, and
pharmacological inhibition of SGLT2 decreases the re-absorption of urate from the urine[32].
While it has been suggested that this effect is mediated indirectly through the effects of
elevated urinary glucose on urate transport by GLUT9 (*SLC2A9*)[31] and URAT1 (*SLC22A12*)
[31,33], studies have shown that GLUT9 is dispensable for the hyperuricosuria[32] and the impact
of glucose on URAT1 remains unclear. This raises the possibility that SGLT2 may directly
mediate urate transport, a hypothesis worthy of future investigation.
255
SLC5A9 encodes the SGLT4 protein, another sodium-dependent sugar transporter
expressed in the small intestine[34,35], although little else is known about its function.
Combined with the fact that approximately 30% of uric acid is excreted into the small
intestine, the known overlap between urate and sugar transporters[26] supports the possibility
that *SLC5A9* could also transport urate. Enhancing intestinal urate disposal has been
proposed as a therapeutic target for chronic kidney disease[36] and so it will be important for
future studies to clarify whether this might be possible via targeting *SLC5A9*.
263
264 These examples highlight how human genetic data can be leveraged to gain insights into
265 protein function and emphasise the utility of *Synteny* in informing targeted validation
266 experiments
267

268
269
270 **Figure 3 – Synteny leverages human genetic data for discovery. (A)** Results of HPE for
271 413 members of the solute carrier (SLC) superfamily, only possible using Synteny. Asterisks
272 represent unadjusted p values. **(B)** Correlation heatmap of HuGE scores for SLC genes with
273 more than one R > 0.95 p < 0.01. Clusters contain SLC genes with similar phenotypic
274 associations. **(C)** Top scoring phenotype for each SLC cluster.

Discussion

Combined with existing bioinformatic tools for identifying genes of interest, *Synteny* provides a systematic and scalable pipeline to maximise the biological insight that can be drawn from omics studies (https://bigproteomics.shinyapps.io/Synteny/).

Synteny is not intended to replace existing knowledge portals. These databases contain vast amounts of information and facilitate targeted inspection of individual genes. Rather, *Synteny* complements these resources by helping researchers prioritise where to look, highlighting which genes or pathways are most strongly supported by human genetic evidence. To streamline downstream analysis, *Synteny* automatically generates hyperlinks to the A2F database for genes and phenotypes of interest. This functionality enables users to transition seamlessly from broad scale discovery to targeted experimental follow-up.

Several new analyses are possible using *Synteny*. By searching for genes with shared disease associations, it is possible to assign putative functions to proteins. In an analysis of the SLC family, this approach led us to propose SGLT2 and SGLT4 as putative urate transporters. Combined with the observation that both genetic and pharmacological manipulation of SGLT2 influences plasma urate levels[31,32], the fact that these genes clustered together with three known urate transporters provides considerable support for the use of human genetic evidence to group genes by their physiological functions.

Beyond functional annotation, *Synteny's* phenotype enrichment capabilities offer clinical applications. For rare diseases, HPE might reveal related phenotypes that could serve to improve diagnosis or treatment, highlighting the importance of comprehensive phenotype analysis.

As with all tools, however, *Synteny* is not without limitation. Firstly, analysis of genes from diverse tissues could limit mechanistic insight in cases where gene function is not well characterised. To mitigate this, we suggest users stratify genes by tissue expression prior to HPE. This issue does not apply to many common situations where genes sets are obtained by analysing omics data in a specific tissue. Secondly, *Synteny* does not currently incorporate information from Mendelian randomisation or protein quantitative trait loci (pQTL) studies linking proteins to disease. These datasets offer an additional form of evidence to aid candidate prioritisation. We encourage users to make use of data from proteogenomics consortia such as SCALLOP[37] and the UK BioBank[38] to perform these analyses once they have identified high-priority candidates using *Synteny*. Finally, *Synteny* does not currently support custom trait-gene mapping using local GWAS data.

Synteny addresses a fundamental challenge in modern biology by bridging the gap between studies in model systems and human genetic studies. While the benefits to those working in model systems were discussed above, this approach also offers considerable benefits to those working with human genetic data. Human genome-wide association studies (GWAS) have identified thousands of disease-associated loci. However, these loci often contain many genes and so pinpointing the causal gene/s within these loci remains a persistent challenge. By enabling users to cross-reference GWAS candidates with functional evidence from model systems, *Synteny* helps to prioritise which genes within complex loci are most likely to be causal and merit experimental validation.

Resource availability

Synteny is freely available at the following address:
https://bigproteomics.shinyapps.io/Synteny/. Code is open-source and available in the
supplementary materials. This can be run locally using RStudio.

Acknowledgements

This work was supported by an Australian Research Council Laureate Fellowship (to DEJ)
and an Australian Government Research Training Program Scholarship (to HBC). The
content is solely the responsibility of the authors and does not necessarily represent the
official views of the ARC.

We would like to thank several colleagues for their constructive feedback: Marcus Seldin for
suggestions relating to enabling broader use of the platform by incorporating the Alliance for
Genome Resources database; Alice Williamson for suggestions relating to heatmap
visualisation settings and improving the documentation for HuGE correlation analysis; Rob
Williams for suggestions relating to the inclusion of metadata in exported results; and Mark
Keller for suggestions relating to using *Synteny* with large (> 500) numbers of genes. These
conversations greatly improved the user experience of *Synteny.* We welcome future
suggestions from the community.

Author contributions

Conceptualisation, HBC, JS, SM, SWCM, OKF, DEJ; Methodology, HBC; Formal Analysis,
HBC; Data curation, HBC; Writing – Original Draft, HBC; Writing – Review & Editing, HBC,
JS, SM, SWCM, OKF, DEJ; Visualisation, HBC, TBS; Supervision, DEJ; Funding
Acquisition, DEJ.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Supplementary file 1: zip file containing source code required to run *Synteny* locally.

Figure legends

Figure 1 – Synteny powers improved access to human genetic evidence. (A) Human genetic
evidence (HuGE) scores aggregate genetic data from many studies to summarise the
strength of association between genes and phenotypes12, enabling gene function to be
rapidly assessed. Example of HuGE score plot linking PPARG to 1175 phenotypes. (B)
Schematic of the current implementation of the HuGE score framework via the Association to
Function web portal. (C) Schematic illustrating the high throughput implementation in
Synteny. API = application programming interface. Examples of data visualisation and
analysis in Synteny for 52 genes hypothesised to be linked to glucose metabolism12 (D &
E). (D) Heatmap showing associations between genes and human phenotypes, filtered for
genes with HuGE scores ≥ 30. (E) Results of HPE. Asterisks represent unadjusted p values.

Figure 2 – Benchmarking for human phenotype enrichment. (A) Example of sensitivity
analysis performed using Obesity as a model trait. Random weighted sampling (i) was used
to generate enriched gene sets with different numbers of genes that were then subjected to
HPE (ii). (B) Power simulations for 15 randomly sampled phenotypes from 16 trait

categories, with gene sets of size n = 100. Each dot is the average of 1000 permutations per
phenotype, showing the degree of enrichment required to reach 80% power. Median
enrichment for 80% power ($\tilde{x}$) = 1.66-fold. (C) HPE using gene sets from HALLMARK
pathways. Genes contributing to the top enrichments for 'Adipogenesis', 'Pancreatic β cells'
and 'Inflammatory response' are shown in i-iii, respectively. (D) Heatmap presenting HuGE
scores for glycolytic enzymes expressed in skeletal muscle for selected phenotypes.
Enzymes that regulate total glycolytic flux19 are highlighted in red.
384
385 Figure 3 – Synteny leverages human genetic data for discovery. (A) Results of HPE for 413
386 members of the solute carrier (SLC) superfamily, only possible using Synteny. Asterisks
387 represent unadjusted p values. (B) Correlation heatmap of HuGE scores for SLC genes with
388 more than one R > 0.95 p < 0.01. Clusters contain SLC genes with similar phenotypic
389 associations. (C) Top scoring phenotype for each SLC cluster.
390
391 **Methods**
392
393 Statistical analyses
394
395 All analysis and data visualisation were conducted using the R programming environment[39].
396 Unless otherwise stated, correlations were assessed by Pearson correlation analysis.
397 Significance is represented, with a p value < 0.05 by *, < 0.01 by **, < 0.001 by *** and <
398 0.0001 by ****.
399
400 Human phenotype enrichment
401
Human genetic evidence (HuGE) scores were downloaded for each gene. To assess
enrichment for a particular phenotype, the same number of genes were randomly selected
from the Association to Function database (for naïve enrichment) or randomly from a set of
background genes (for background-corrected enrichment). This was repeated for 10,000
permutations to enable the average HuGE score in the user-defined gene set to be
compared to the sampled distribution/s. An enrichment score was calculated as the mean
fold change of the average scores, and significance was determined by calculating the
proportion of sampled scores exceeding that of the user-defined set. Multiple corrections
were performed using a false-discovery rate (FDR) correction to account for the number of
phenotype tests.
412
413 Benchmarking
414
415 *Sensitivity analysis*
416
417 Gene sets with progressively greater enrichment were generated by weighted sampling. We
418 performed permutation testing to determine the power of HPE to detect enrichment for gene
419 sets with different levels of enrichment (n = 1000). Power is reported as the proportion of
420 gene sets where enrichment was detected with p < 0.05. Phenotypes in Figure 2B were
421 obtained by randomly sampling 15 phenotypes per trait category.
422
423 *HALLMARK analysis*
424
425 Human HALLMARK gene sets were downloaded from https://www.gsea-
426 msigdb.org/gsea/msigdb/human/genesets.jsp?collection=H and analysed using HPE.
427 Significantly enriched phenotypes are reported.
428
429 Synteny
430

431   *Web server implementation*
432
433   *Synteny* is implemented as a Shiny application using the R programming language and is
434   hosted on a shinyapps.io server.
435
436   *Privacy and security*
437
438   Gene lists are only uploaded if the user accesses *Synteny* via the web server. No user data
439   are retained following session termination. Secure HTTPS connections are used to transfer
440   data to and from the server. The country in which each user is located is logged using a
441   google analytics cookie. This data is used solely to monitor global adoption of the tool and
442   ensure server requirements are sufficient to meet demand. No other data is recorded.
443
444   *Local installation*
445
446   The code required to run *Synteny* locally is available in the supplementary materials. After
447   installing the required R packages, *Synteny* will still require an internet connection to connect
448   to the Association to Function application programming interface. No data will be uploaded to
449   the *Synteny* shinyapps.io server if running locally.
450
451   *Packages and databases*
452
453   *Synteny* maps gene names from any organism to human orthologs using the Alliance for
454   Genome Resources database[10]. Human genetic evidence (HuGE) scores are retrieved via
455   an application programming interface connected to the Association to Function database
456   using the jsonlite package[40]. Static heatmaps are created using the ComplexHeatmap
457   package[41] and interactive plots are created using a combination of ggplot2[42] and plotly[43].
458
459   *Exporting results*
460
461   Results and figures can be explored and visualised within *Synteny* itself or exported for
462   analysis in the users' software of choice. Data is exportable as excel sheets (.xlsx) and
463   figures are exportable as full resolution vector PDFs.

References

1. Müller, J.B., Geyer, P.E., Colaço, A.R., Treit, P. V., Strauss, M.T., Oroshi, M., Doll, S., Virreira Winter, S., Bader, J.M., Köhler, N., et al. (2020). The proteome landscape of the kingdoms of life. Nature 2020 582:7813 *582*, 592–596. https://doi.org/10.1038/s41586-020-2402-x.

2. Hutchison, C.A., Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman, M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L., et al. (2016). Design and synthesis of a minimal bacterial genome. Science (1979) *351*. https://doi.org/10.1126/SCIENCE.AAD6253/SUPPL_FILE/AAD6253-HUTCHISON-SM.PDF.

3. Kustatscher, G., Collins, T., Gingras, A.C., Guo, T., Hermjakob, H., Ideker, T., Lilley, K.S., Lundberg, E., Marcotte, E.M., Ralser, M., et al. (2022). Understudied proteins: opportunities and challenges for functional proteomics. Nat Methods *19*, 774–779. https://doi.org/10.1038/S41592-022-01454-X.

4. Edwards, A.M., Isserlin, R., Bader, G.D., Frye, S. V., Willson, T.M., and Yu, F.H. (2011). Too many roads not taken. Nature 2011 470:7333 *470*, 163–165. https://doi.org/10.1038/470163a.

5. Keller, M.P., Rabaglia, M.E., Schueler, K.L., Stapleton, D.S., Gatti, D.M., Vincent, M., Mitok, K.A., Wang, Z., Ishimura, T., Simonett, S.P., et al. (2019). Gene loci associated with insulin secretion in islets from nondiabetic mice. J Clin Invest *129*, 4419–4432. https://doi.org/10.1172/JCI129143.

6. Li, Z., Votava, J.A., Zajac, G.J.M., Lamming, D.W., Liang, C.-, Yen, E., and Parks Correspondence, B.W. (2020). Integrating Mouse and Human Genetic Data to Move beyond GWAS and Identify Causal Genes in Cholesterol Metabolism. Cell Metab *31*, 741-754.e5. https://doi.org/10.1016/j.cmet.2020.02.015.

7. Parker, B.L., Calkin, A.C., Seldin, M.M., Keating, M.F., Tarling, E.J., Yang, P., Moody, S.C., Liu, Y., Zerenturk, E.J., Needham, E.J., et al. (2019). An integrative systems genetic analysis of mammalian lipid metabolism. Nature *567*, 187. https://doi.org/10.1038/S41586-019-0984-Y.

8. Mountjoy, E., Schmidt, E.M., Carmona, M., Schwartzentruber, J., Peat, G., Miranda, A., Fumis, L., Hayhurst, J., Buniello, A., Karim, M.A., et al. (2021). An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated loci. Nat Genet *53*, 1527–1533. https://doi.org/10.1038/S41588-021-00945-5.

9. Costanzo, M.C., von Grotthuss, M., Massung, J., Jang, D., Caulkins, L., Koesterer, R., Gilbert, C., Welch, R.P., Kudtarkar, P., Hoang, Q., et al. (2023). The Type 2 Diabetes Knowledge Portal: An open access genetic resource dedicated to type 2 diabetes and related traits. Cell Metab *35*, 695-710.e6. https://doi.org/10.1016/J.CMET.2023.03.001.

10. Bult, C.J., and Sternberg, P.W. (2023). The alliance of genome resources: transforming comparative genomics. Mammalian Genome *34*, 531. https://doi.org/10.1007/S00335-023-10015-2.

11. Pers, T.H., Karjalainen, J.M., Chan, Y., Westra, H.J., Wood, A.R., Yang, J., Lui, J.C., Vedantam, S., Gustafsson, S., Esko, T., et al. (2015). Biological interpretation of genome-wide association studies using predicted gene functions. Nature Communications 2015 6:1 *6*, 1–9. https://doi.org/10.1038/ncomms6890.

12. Dornbos, P., Singh, P., Jang, D.K., Mahajan, A., Biddinger, S.B., Rotter, J.I., McCarthy, M.I., and Flannick, J. (2022). Evaluating human genetic support for hypothesized metabolic disease genes. Cell Metab *34*, 661–666. https://doi.org/10.1016/j.cmet.2022.03.011.

13. Pal, A., Barber, T.M., Van de Bunt, M., Rudge, S.A., Zhang, Q., Lachlan, K.L., Cooper, N.S., Linden, H., Levy, J.C., Wakelam, M.J.O., et al. (2012). PTEN Mutations as a Cause of Constitutive Insulin Sensitivity and Obesity . New England Journal of Medicine *367*, 1002–1011.

https://doi.org/10.1056/NEJMOA1113966/SUPPL_FILE/NEJMOA1113966_DISCLOS
URES.PDF.

14. Wijesekara, N., Konrad, D., Eweida, M., Jefferies, C., Liadis, N., Giacca, A.,
Crackower, M., Suzuki, A., Mak, T.W., Kahn, C.R., et al. (2005). Muscle-specific Pten
deletion protects against insulin resistance and diabetes. Mol Cell Biol 25, 1135–1145.
https://doi.org/10.1128/MCB.25.3.1135-1145.2005.

15. McGarry, J.D., Leatherman, G.F., and Foster, D.W. (1978). Carnitine
palmitoyltransferase I. The site of inhibition of hepatic fatty acid oxidation by malonyl-
CoA. Journal of Biological Chemistry 253, 4128–4136. https://doi.org/10.1016/S0021-
9258(17)34693-8.

16. Schlaepfer, I.R., and Joshi, M. (2020). CPT1A-mediated Fat Oxidation, Mechanisms,
and Therapeutic Potential. Endocrinology 161.
https://doi.org/10.1210/ENDOCR/BQZ046.

17. Kulkarni, P.P., Ekhlak, M., Singh, V., Kailashiya, V., Singh, N., and Dash, D. (2023).
Fatty acid oxidation fuels agonist-induced platelet activation and thrombus formation:
Targeting β-oxidation of fatty acids as an effective anti-platelet strategy. FASEB J 37.
https://doi.org/10.1096/FJ.202201321RR.

18. Kulkarni, P.P., Ekhlak, M., and Dash, D. (2023). Energy metabolism in platelets fuels
thrombus formation: Halting the thrombosis engine with small-molecule modulators of
platelet metabolism. Metabolism 145, 155596.
https://doi.org/10.1016/J.METABOL.2023.155596.

19. Öztürk, M., Freiwald, A., Cartano, J., Schmitt, R., Dejung, M., Luck, K., Al-Sady, B.,
Braun, S., Levin, M., and Butter, F. (2022). Proteome effects of genome-wide single
gene perturbations. Nature Communications 2022 13:1 13, 1–10.
https://doi.org/10.1038/s41467-022-33814-8.

20. Tanner, L.B., Goglia, A.G., Wei, M.H., Sehgal, T., Parsons, L.R., Park, J.O., White, E.,
Toettcher, J.E., and Rabinowitz, J.D. (2018). Four Key Steps Control Glycolytic Flux in
Mammalian Cells. Cell Syst 7, 49-62.e8. https://doi.org/10.1016/J.CELS.2018.06.003.

21. Stoeger, T., Gerlach, M., Morimoto, R.I., and Nunes Amaral, L.A. (2018). Large-scale
investigation of the reasons why potentially important genes are ignored. PLoS Biol
16, e2006643. https://doi.org/10.1371/JOURNAL.PBIO.2006643.

22. Li, H., Rukina, D., David, F.P.A., Yang Li, T., Oh, C.M., Gao, A.W., Katsyuba, E.,
Sleiman, M.B., Komljenovic, A., Huang, Q., et al. (2019). Identifying gene function and
module connections by the integration of multispecies expression compendia.
Genome Res 29, 2034–2045. https://doi.org/10.1101/GR.251983.119/-/DC1.

23. Superti-Furga, G., Lackner, D., Wiedmer, T., Ingles-Prieto, A., Barbosa, B., Girardi, E.,
Goldmann, U., Gürtl, B., Klavins, K., Klimek, C., et al. (2020). The RESOLUTE
consortium: unlocking SLC transporters for drug discovery. Nature Reviews Drug
Discovery 2020 19:7 19, 429–430. https://doi.org/10.1038/d41573-020-00056-6.

24. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted
correlation network analysis. BMC Bioinformatics 9, 559. https://doi.org/10.1186/1471-
2105-9-559.

25. Seldin, M.M., Koplev, S., Rajbhandari, P., Vergnes, L., Rosenberg, G.M., Meng, Y.,
Pan, C., Phuong, T.M.N., Gharakhanian, R., Che, N., et al. (2018). A Strategy for
Discovery of Endocrine Interactions with Application to Whole-Body Metabolism. Cell
Metab 27, 1138-1155.e6. https://doi.org/10.1016/J.CMET.2018.03.015.

26. Vitart, V., Rudan, I., Hayward, C., Gray, N.K., Floyd, J., Palmer, C.N.A., Knott, S.A.,
Kolcic, I., Polasek, O., Graessler, J., et al. (2008). SLC2A9 is a newly identified urate
transporter influencing serum urate concentration, urate excretion and gout. Nat
Genet 40, 437–442. https://doi.org/10.1038/NG.106.

27. Dai, Y., and Lee, C.H. (2024). Transport mechanism and structural pharmacology of
human urate transporter URAT1. Cell Research 2024 34:11 34, 776–787.
https://doi.org/10.1038/s41422-024-01023-1.

28.   Bahn, A., Hagos, Y., Reuter, S., Balen, D., Brzica, H., Krick, W., Burckhardt, B.C., and
Sabolic´, I. (2008). Identification of a New Urate and High Affinity Nicotinate
Transporter, hOAT10 (SLC22A13) *. https://doi.org/10.1074/jbc.M800737200.
29.   Ferrannini, E., and Solini, A. (2012). SGLT2 inhibition in diabetes mellitus: rationale
and clinical prospects. Nature Reviews Endocrinology 2012 8:8 *8*, 495–502.
https://doi.org/10.1038/nrendo.2011.243.
30.   Chao, E.C., and Henry, R.R. (2010). SGLT2 inhibition — a novel strategy for diabetes
treatment. Nature Reviews Drug Discovery 2010 9:7 *9*, 551–559.
https://doi.org/10.1038/nrd3180.
31.   Inthasot, S., Vanderhulst, J., Janssens, P., Daele, S. Van, Hoof, E. Van, Kint, C.,
Iconaru, L., and de Filette, J. (2024). A novel heterozygous likely pathogenic SLC5A2
variant in a diabetic patient with glucosuria and aminoaciduria. Endocrinol Diabetes
Metab Case Rep *2024*. https://doi.org/10.1530/EDM-24-0065.
32.   Novikov, A., Fu, Y., Huang, W., Freeman, B., Patel, R., van Ginkel, C., Koepsell, H.,
Busslinger, M., Onishi, A., Nespoux, J., et al. (2019). SGLT2 inhibition and renal urate
excretion: role of luminal glucose, GLUT9, and URAT1. Am J Physiol Renal Physiol
316, F173–F185. https://doi.org/10.1152/AJPRENAL.00462.2018.
33.   Dong, M., Chen, H., Wen, S., Yuan, Y., Yang, L., Xu, D., and Zhou, L. (2023). The
Mechanism of Sodium-Glucose Cotransporter-2 Inhibitors in Reducing Uric Acid in
Type 2 Diabetes Mellitus. Diabetes, Metabolic Syndrome and Obesity *16*, 437–445.
https://doi.org/10.2147/DMSO.S399343.
34.   Tazawa, S., Yamato, T., Fujikura, H., Hiratochi, M., Itoh, F., Tomae, M., Takemura, Y.,
Maruyama, H., Sugiyama, T., Wakamatsu, A., et al. (2005). SLC5A9/SGLT4, a new
Na+-dependent glucose transporter, is an essential transporter for mannose, 1,5-
anhydro-D-glucitol, and fructose. Life Sci *76*, 1039–1050.
https://doi.org/10.1016/J.LFS.2004.10.016.
35.   Sano, R., Shinozaki, Y., and Ohta, T. (2020). Sodium–glucose cotransporters:
Functional properties and pharmaceutical potential. J Diabetes Investig *11*, 770.
https://doi.org/10.1111/JDI.13255.
36.   Johnson, R.J. (2023). Intestinal Hyperuricemia as a Driving Mechanism for CKD.
American Journal of Kidney Diseases *81*, 127–130.
https://doi.org/10.1053/j.ajkd.2022.08.001.
37.   Zhao, J.H., Stacey, D., Eriksson, N., Macdonald-Dunlop, E., Hedman, Å.K.,
Kalnapenkis, A., Enroth, S., Cozzetto, D., Digby-Bell, J., Marten, J., et al. (2023).
Genetics of circulating inflammatory proteins identifies drivers of immune-mediated
disease risk and therapeutic targets. Nature Immunology 2023 24:9 *24*, 1540–1551.
https://doi.org/10.1038/s41590-023-01588-w.
38.   Sun, B.B., Chiou, J., Traylor, M., Benner, C., Hsu, Y.H., Richardson, T.G., Surendran,
P., Mahajan, A., Robins, C., Vasquez-Grinnell, S.G., et al. (2023). Plasma proteomic
associations with genetics and health in the UK Biobank. Nature 2023 622:7982 *622*,
329–338. https://doi.org/10.1038/s41586-023-06592-6.
39.   R Core Team (2021). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria., URL https://www.R-
project.org/.
40.   Ooms, J. (2024). A Simple and Robust JSON Parser and Generator for R [R package
jsonlite version 1.8.9]. CRAN: Contributed Packages.
https://doi.org/10.32614/CRAN.PACKAGE.JSONLITE.
41.   Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. Bioinformatics *32*, 2847–2849.
https://doi.org/10.1093/BIOINFORMATICS/BTW313.
42.   Wickham, H. (2016). ggplot2. https://doi.org/10.1007/978-3-319-24277-4.
43.   Sievert, C. (2021). Interactive Web-Based Data Visualization with R, Plotly, and Shiny.
J R Stat Soc Ser A Stat Soc *184*, 1150–1150. https://doi.org/10.1111/RSSA.12692.
625

A Human Genetic Evidence

PPARG

B Current implementation

ASSOCIATION TO FUNCTION
KNOWLEDGE PORTAL

Manual querying

Search gene, variant, region, phenotype, or tissue

Wait for results to load

Genetic support workflow

1 Find significant variants
2 Explore common & rare effects
3 Investigate HuGE scores

Repeat (2-5 min/gene)

C High throughput workflow

SYNTENY

Experimental candidates → API ← A2F

multi species support

1 Upload genes — 2 Acquire data — 3 Visualise interactively

D HuGE scores

log₂(HuGE)

A2F: 1hr   Synteny: 1 min 20 sec

E Human phenotype enrichment

AMD progression ***
Omega-3 fatty acids **
DHA
Late onset AD ***
Valine
Peptic Ulcers
TSH **
AD
BMI 6 months ***
BMI 1-5 yrs ***
BMI 3 months ***
BMI 1 yr ***
Asc. aorta distensibility *
Psychiatric *
Chylomicrons *
Glycoproteins *
Fat intake adj. BMI *
AMD *
Long QT syndrome *

p < 0.05
FDR < 0.05

log₂(HuGE) enrichment

15 min (299 phenotypes)

A Sensitivity analysis

i) Weighted sampling

increased HuGE score weighting

log₂(Enrichment): -2, 0, 2, 4

ii) Effect of sample size

Power (%): 0, 25, 50, 75, 100

Fold Enrichment: 1–10

Gene set size (n): 10, 20, 30, 40, 50, 100, 200, 300

B Robustness across phenotypes

- Aging and longevity
- Anthropometric
- Cancer
- Cardiovascular
- Glycemic
- Hematological
- Hepatic
- Immunological
- Lipids
- Lung
- Musculoskeletal
- Neurological
- Psychiatric
- Renal
- Reproductive traits
- Sleep and circadian

x̄ = 1.66-fold (n = 100)

log₂(Fold Enrichment) for 80% power: 1–8

C Robustness against gene set composition

Adipogenesis (i)

- Adiponectin
- TG-HDL ratio
- Body fat percentage
- HDL
- TG
- ALS
- Cholesterol
- LDL
- HDL-cholesterol ratio
- Maternal longevity
- Extreme longevity
- Paternal Alzheimer's disease
- Hip fracture
- AMD progression
- NAFLD
- TG in T2D
- Paternal longevity
- Carbohydrate intake
- Parental Lifespan
- TG in non-T2D
- Brain microbleeds

log₂(Enrichment): 0–6

Pancreatic β cells (ii)

- Fasting glucose
- Diabetic retinopathy
- Blood sugar adj. fasting
- Insulinogenic index
- Modified insulinogenic index
- Stumvoll
- Corrected insulin response
- DI (Kadowaki/Matsuda)
- Fetal genetic effect on body weight
- Smoking-T2D interaction

log₂(Enrichment): 0–6

Inflammatory response (iii)

- Crohn's disease
- Coronary artery disease
- RML mean attenuation

log₂(Enrichment): 0, 2, 4, 6

Significant phenotypes

Aging and longevity, Anthropometric, Cardiovascular, Cerebrovascular MRI traits, Diabetic complications, Glycemic, Hepatic, Inflammatory bowel, Lipids, Lung, Musculoskeletal, Neurological, Nutritional, Ocular, Reproductive traits, Stroke

i) Adiponectin
log₂(HuGE): 0–15; Rank: 0–6000
ADIPOQ, SLC1A5, ADIG, PPARG, ADIPOR2, RTN3, GPD2, RAB34

ii) Fasting Glucose
G6PC2, FOXA2, GCK, PCSK1, SLC2A2, PDX1, SRP14, PAX4; Rank: 0–8000

iii) Crohn's Disease
NOD2, NFKBIA, TAPBP, PTAFR, CALCRL, IRF1, GABBR1, CD40, IL10, IL12B, ICAM4, CCL7; Rank: 0–8000

D Disentangling pathways (glycolysis)

Import | Upper glycolysis | Lower glycolysis | Export

Rows: Fasting insulin adj. BMI, Heart rate, ALT, AST, Triglycerides, Plasma pyruvate, HBA1C, BMI, T2D

Genes: SLC2A1, SLC2A4, HK1, HK2, GPI, PFKM, PFKFB1, PFKFB2, PFKFB3, TIGAR, FBP1, FBP2, ALDOA, GAPDH, PGK1, BPGM, PGAM2, ENO1, ENO3, PKM, LDHA, SLC16A3

log₂(HuGE): 0–10

* regulates flux (Tanner et al. 2018)

A Solute carrier (SLC) enrichment

413 SLC Proteins

Chronic kidney disease ***
Creatinine ***
Phosphate *
Urate ***
Cystatin C *
Retinal vessel branching ***
Fasting glucose ***
Renal cortex stiffness ***
Liver age ***
Disposition index ***
Sodium ***
Pancreatic stiffness ***
COVID-19 ***
Alanine *
Low hand grip strength ***
Cerebellar volume *
T2D + neuro. outcome *
Corrected QT interval *
BMI adj. smoking *
Meconium ileus in CF ***
Minimum SpO₂ (sleep) ***
F'wd LV stroke volume **
Avg SpO₂ (hypopnea) **
Epilepsy + sclerosis *

p < 0.05
FDR < 0.05

$\log_2$(HuGE enrichment)

B Phenotypic correlations

R
-1 0 1

SLCO2B1
SLC35A3
SLC9A3P1
SLC28A3
SLC17A6
SLC25A43
SLC52A3
SLC2AXP1
SLC30A4
SLC36A4
SLC45A11
SLC36A3
SLC22A7
SLC47A1
SLC25A45
SLC34A1
SLC24A5
SLC4A5
SLC7A4
SLC18A3
SLC6A5
SLC35G3
SLC22A13
SLC22A9
SLC5A2
SLC22A12
SLC22A11
MFSD9
SLC39A5
SLC4A6
SLC30A2

C Top phenotype by cluster

Cluster 1 — Height
Cluster 2 — Serum creatinine
Cluster 3 — Fasting insulin adj. BMI
Cluster 4 — Serum urate
  SLC2A9 – known urate transporters
  SLC22A12
  SLC5A2
  SLC22A13
  SLC5A9
Cluster 5 — Testosterone
Cluster 6 — Alkaline phosphatase (ALP)

$\log_2$(HuGE)