Knocking out Arl13b completely abolishes primary ciliogenesis

Divyanshu Mahajan[1], Hui Min Chia[1], and Lei Lu[1*]

[1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,

Singapore 637551.

*Correspondence should be addressed to:

Lei Lu (PhD)

School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore

637551

Tel: 65-65922591

Email: lulei@ntu.edu.sg

Keywords: ARL13B, cilium, ciliogenesis, sonic hedgehog signaling, Joubert syndrome.

Abstract:

Small G protein ARL13B localizes to the cilium and plays essential roles in cilium biogenesis, organization, trafficking, and signaling. Here, we established multiple ARL13B knockout cell lines using the CRISPR/Cas9 system. Surprisingly, all our cell lines lost their cilia completely, in contrast to the reported short cilium and reduced ciliogenesis phenotype. We found that multiple regions of ARL13B are necessary for a complete rescue. Additionally, we found that ARL13B knockout cells also lost their response to sonic hedgehog stimulation. Our work demonstrates the absolute requirement of ARL13B for ciliogenesis and sonic hedgehog signaling, at least in cultured cells, and suggest that ARL13B plays a far more critical role in ciliary function than previously understood.

Introduction

The primary cilium (hereafter cilium) is a microtubule-based plasma membrane protrusion that functions as the cell's antenna. The cilium detects extracellular signals through cilium-localized receptors and triggers intracellular responses through various signaling pathways, such as sonic hedgehog, Wnt, and TGFβ signal transduction pathways (Anvarian, Mykytyn, Mukhopadhyay, Pedersen, & Christensen, 2019; Pala, Alomari, & Nauli, 2017; Wheway, Nazlamova, & Hancock, 2018). The critical role of the cilium in sensing and transducing signals for development and physiological function depends on its unique proteins, mutations of which can cause human disorders collectively termed ciliopathies (Goetz & Anderson, 2010; Hildebrandt, Benzing, & Katsanis, 2011). The ARF family small G protein ARL13B, the mutation of which causes the ciliopathy Joubert syndrome, has been recognized as one of the most crucial cilium regulators. ARL13B is highly enriched in the cilium, and such localization was mediated by the ciliary transport adaptor Rab8-TNPO1 or Tulp3 (Mahajan, Madugula, & Lu, 2023; Palicharla et al., 2023). Within the cilium, ARL13B recruits INPP5E to modulate the level of phosphoinositides (Chavez et al., 2015; Garcia-Gonzalo et al., 2015; Humbert et al., 2012; Nozaki et al., 2017). ARL13B can also function as a guanine nucleotide exchange factor to activate another cilium-localized ARF family small G protein, ARL3, to transport lipidated cargos to the cilium (Gotthardt et al., 2015; Ivanova et al., 2017; Q. Zhang et al., 2016).

Deletion of *ARL13B* in animals causes diverse cilium-associated physiological and cellular phenotypes. For example, *ARL13B* null zebrafish (*scorpion* mutant) is characterized by cystic kidney and body curvature with significantly reduced cilium number and abnormal cilium organization (Duldulao, Lee, & Sun, 2009). *ARL13B* null mice (*hennin* mutant) have defective neural tubes, abnormal eyes, and axial polydactyly. At the molecular and cellular level, *hennin* mutant cells display reduced cilium length and biogenesis, abnormal cilium organization, and defective sonic hedgehog signaling (Caspary, Larkins, & Anderson, 2007; Larkins, Aviles, East, Kahn, & Caspary, 2011). With the advent of CRISPR/Cas9 technology, the *ARL13B* gene has been targeted to generate knockout (KO) cell lines. *ARL13B*-KO hTert-RPE1 (hereafter RPE1) cells display short cilium with reduced ciliogenesis, consistent with observations in *ARL13B* null animals (Nozaki et al., 2017). In this work, we produced *ARL13B*-KO RPE1 and HEK293T cells and confirmed the KO of *ARL13B* by genotyping and Western blotting. Surprisingly, we found that cilia were completely lost in our *ARL13B*-KO cells. Concomitantly, the sonic hedgehog signaling was abolished in these cells. Therefore, our data showed the absolute requirement of ARL13B for cilium biogenesis and sonic hedgehog signaling.

Result

ARL13B-KO RPE1 cells completely lack cilia.

To study the cellular function of ARL13B, we employed CRISPR/Cas9-induced frameshift mutation to KO the *ARL13B* gene in RPE1 cells. We designed two sgRNAs, sg1 and 2, to target exons 1 and 5 of the human *ARL13B* gene, respectively (Fig. 1a). After lentivirus-mediated transduction of a single expression vector containing the sgRNA and Cas9, RPE1 cells were subjected to puromycin selection. We selected two stable cell lines, sg1#a and sg2#4, corresponding to sg1 and sg2, respectively. We sequenced the genomic DNA fragments containing the sgRNA target sites from sg1#a and sg2#4 cell lines. We found that the two cell lines have frameshift insertion or deletion of nucleotides in both alleles, which were expected to cause a premature translation termination of the transcribed mRNA and trigger the non-sense-mediated mRNA decay (Kurosaki, Popp, & Maquat, 2019). The open reading frames of frameshifted *ARL13B* in sg1#a and sg2#4 cell lines contain less than 13 and 169 AAs (Supplementary File 1), respectively, compared to wild type (WT) ARL13B, 428 AAs. Next, we assessed ARL13B protein levels in cell lysates prepared from sg1#a and sg2#4 cell lines using Western blotting. A single band at the expected molecular weight of ~ 48 kDa was observed in parental RPE1 cells. In contrast, no band was observed in the two KO cell lines (Fig. 1b), supporting that sg1#a and sg2#4 are *ARL13B*-KO cells.

Serum starvation dramatically increased the ciliogenesis rate of parental RPE1 cells from ~ 20% to 70%, as revealed by the immunostaining of endogenous ARL13B, acetylated tubulin, or glutamylated tubulin (Fig. 1c,d), which are conventional cilium markers. Our immunofluorescence staining confirmed that the endogenous ARL13B level was undetectable in sg1#a and sg2#4 cells, as expected for *ARL13B*-KO. Surprisingly, we found no cilia in our two *ARL13B*-KO cell lines by the acetylated or glutamylated tubulin immunostaining. Our measured ciliogenesis rates are 0.0 ± 0.0 % (*mean ± standard deviation* and same for the rest of the manuscript) (n = 3) in the two KO cell lines stained by the two tubulin cilium markers. Due to their possible short length and weak intensity, many acetylated or glutamylated tubulin-positive cilia might not be identified unambiguously against the cellular background in RPE1 cells. To facilitate the identification of cilia, we co-labeled the cilium and its base by immunostaining endogenous glutamylated tubulin and pericentrin (Jurczyk et al., 2004) (Fig. 1e). In parental RPE1 cells, the cilium marked by glutamylated tubulin always extended from the cilium base marked by pericentrin-positive puncta. In contrast, we never observed cilia around the pericentrin puncta in sg1#a and sg2#4 cells. We also employed another cilium marker, fibrocystin ciliary targeting signal (CTS)-fused CD8a (CD8a-fCTS) (Madugula & Lu, 2016) (Supplementary Fig. 1a,b), and we could not find cilia in our *ARL13B*-KO cell lines either. In summary, our data shows that *ARL13B*-KO RPE1 cells completely lack cilia.

To confirm that the no-cilium phenotype is not an off-target effect of gene editing, we performed rescue experiments by exogenously expressing CRISPR/Cas9-resistant ARL13B coding sequence (ARL13B[R]). ARL13B[R] was constructed by making multiple silent point mutations within sg1 and sg2 target sites in the coding sequence of ARL13B (Fig. 1f). After the lentivirus-transduction of ARL13B[R]-GFP, sg1#a and sg2#4 cells were subjected to serum starvation and immunostaining of endogenous acetylated tubulin (Fig. 1g). Among GFP-positive cells, we observed 65 ± 3 % (n = 3) and 67 ± 2 % (n = 3), respectively, had cilia (Fig. 1h; Supplementary Fig. 1c). The ciliogenesis rates of rescued cells had no significant difference from that of parental RPE1 cells (p = 0.001 by unpaired and two-tailed *t*-test). We previously introduced the cilium to

plasma membrane intensity ratio (*CPIR*) as an expression level-independent metric that quantitatively measures the cilium localization (Madugula & Lu, 2016). We found that the *CPIR*s of ARL13B^R-GFP and cilium lengths in rescued cells had no significant difference from those of parental RPE1 cells (Fig. 1i, j; Supplementary Fig. 1d, e). Therefore, our study demonstrates that ARL13B^R-GFP can rescue the no-cilium phenotype in *ARL13B*-KO cells, suggesting that the no-cilium phenotype should be specific to *ARL13B*-KO.

No cilia could be found in *ARL13B*-KO HEK293T cells either.

We employed HEK293T cell lines to perform similar KO and characterization experiments to confirm our findings in another cell line. We used the same sgRNAs, sg1 and sg2, to generate *ARL13B*-KO cell lines, sg1#5 and sg2#5, respectively. Their genotypes confirmed the frameshift mutations in both alleles (Fig. 2a). By immunofluorescence imaging and Western blotting, we verified that ARL13B protein levels were undetectable in both cell lines (Fig. 2b, c). After serum starvation, we observed 43 ± 4 % (n = 3) parental HEK293T cells acquired acetylated or glutamylated tubulin-positive cilia (Fig. 2d). However, under identical treatment, we found no cilia in our two *ARL13B*-KO HEK293T cell lines, in which all ciliogenesis rates were measured to be 0.0 ± 0.0 % (n = 3). We also demonstrated that the no-cilium phenotype in our *ARL13B*-KO HEK293T cell lines could be rescued by exogenously expressing ARL13B^R-GFP (Supplementary Fig. 2). Therefore, our data confirms the no-cilium phenotype upon *ARL13B*-KO in HEK293T cells and suggests that such an observation might apply to multiple types of cells.

ARL13B-KO by previously reported sgRNAs also displayed the no-cilium phenotype.

The no-cilium phenotype we observed in two *ARL13B*-KO cell lines differs from previous studies in *ARL13B* null mice (*hennin*) (Caspary et al., 2007; Larkins et al., 2011) and zebrafish (*scorpion*) (Duldulao et al., 2009). It is especially in stark contrast to a similar work by Nozaki *et al.*, who similarly produced *ARL13B*-KO RPE1 cells using CRISPR/Cas9 technique (Nozaki et al., 2017). Nozaki *et al*. reported that cilia were still clearly present in *ARL13B*-KO cells but with a reduced length. To resolve this discrepancy, we adopted the same two sgRNAs they designed, sg4 and sg5, corresponding to sgRNA #1 and #2 in their study, to knock out *ARL13B* gene in the same RPE1 cells. Our effort resulted in two clonal cell lines, sg4#2 and sg5#3, respectively (Fig. 3a). Their frameshift insertion and deletion genotypes and undetectable expression levels of ARL13B in Western blot and immunofluorescence confirmed their ARL13B null phenotypes in both cell lines (Fig. 1b; Fig. 3b). Importantly, consistent with our sg1#a, sg2#4, sg1#5, and sg2#5 cells, we still observed a complete loss of cilia by immunostaining of endogenous acetylated and glutamylated tubulin with a ciliogenesis rate of 0.0 ± 0.0 % (n = 3) for both cell lines (Fig. 3b,c). Similarly, we also demonstrated that the no-cilium phenotype was rescued by exogenously expressing ARL13B^R-GFP (Fig. 3d).

Gene KOs generated by the CRISPR/Cas9-induced frameshift mutation can still express residual target protein, resulting in a weak phenotype (Meraldi, 2019; Smits et al., 2019; G. Zhang et al., 2019). Since our KO cells exhibit a much stronger phenotype, no-cilium, than that reported by Nozaki *et al*. and others, which is reduced cilium length and ciliogenesis rate (Caspary et al., 2007; Duldulao et al., 2009; Nozaki et al., 2017), we believe our KO cells likely represent a more complete deletion of the *ARL13B* gene (see Discussion). In summary, using two sgRNAs in two cell lines and replicating a previous study, we demonstrated that *ARL13B*-KO cells lack cilia entirely rather than merely exhibiting reduced cilium length and biogenesis.

Multiple ARL13B regions or domains are essential for its ciliogenesis function.

To investigate the roles of different ARL13B regions or domains in ciliogenesis, we assessed the
rescue of the no-cilium phenotype of sg1#a cells using a series of ARL13B mutants. ARL13B
consists of the following regions or domains from the N to the C-terminus: an α-helix region with
palmitoylation lipid anchorage sites (PAL), a GTPase domain, a coiled-coil region (CC), an
intrinsically disordered region (IDR), the RVEP motif, and a proline-rich region (PRR) (Fig 4a).
The N-terminal PAL contains two palmitoylation sites at C8 and C9, which enable the membrane
anchorage of ARL13B. Mutation of both palmitoylation sites from cysteine to serine (C8S/C9S) in
PAL abolishes the cilium localization of ARL13B (Cevik et al., 2010; Cevik et al., 2013; Mahajan
et al., 2023; Roy et al., 2017). The GTPase domain is essential for the interaction with INPP5E
and can act as a guanine nucleotide exchange factor for ARL3 (Gotthardt et al., 2015; Humbert
et al., 2012; Ivanova et al., 2017; Nozaki et al., 2017; Q. Zhang et al., 2016). According to the
structural and functional analysis of Ras superfamily (including ARF family) small G proteins, the
T35N mutation of the GTPase domain's G1 box is expected to lock ARL13B in a GDP-binding
state (Feig, 1999; Sprang, 1997; Sztul et al., 2019). The cellular function of the CC remains
elusive, but it is essential for the cilium localization of ARL13B and INPP5E interaction (Mahajan
et al., 2023; Nozaki et al., 2017). While the function of the IDR is still unclear, the PRR is known
to interact with the IFT-B complex and INPP5E (Nozaki et al., 2017). Although both IDR and PRR
are dispensable for the cilium localization of ARL13B, the RVEP motif, a stretch of 17 AAs located
in between the IDR and PRR, serves as the CTS of ARL13B by interacting with ciliary transport
adaptors, TNPO1 and Rab8 (Mahajan et al., 2023).

We prepared the following mutants in the background of ARL13B^R-GFP: C8S/C9S, ΔGTPase
(deletion of GTPase domain), ΔCC (deletion of CC), ΔIDR (deletion of IDR), RVEP-4A (mutation
of RVEP to AAAA), ΔPRR (deletion of PRR), and AA1-245 (deletion of AAs from 246 to 428) (Fig.
4a). In parental RPE1 cells, we demonstrated that C8S/C9S and ΔCC do not localize to the cilium,
while ΔGTPase, ΔIDR, and ΔPRR show robust cilium localization and RVEP-4A and AA1-245
exhibit weak cilium localization (Fig. 4a; Fig. 4b, parental RPE1 cells), consistent with our previous
study (Mahajan et al., 2023). Additionally, we observed that T35N has a robust cilium localization
(Fig. 4a; Fig. 4b, parental RPE1 cells).

Next, we tested these ARL13B mutants for their ability to rescue the no-cilium phenotype of sg1#a
cells. Following lentivirus-mediated transduction and serum starvation, we analyzed ciliogenesis
by immunostaining endogenous acetylated tubulin and pericentrin (Fig. 4b,c). We found that
T35N, ΔIDR, and ΔPRR successfully rescued the no-cilium phenotype with a ciliogenesis
percentage of $\geq$ 70 %, showing no significant difference from ARL13B^R-GFP ($p > 0.05$)(Fig. 4c).
In contrast, no cilia were found in C8S/C9S, ΔGTPase, ΔCC, and AA1-245 expressing sg1#a
cells, all exhibiting a ciliogenesis percentage of 0.0 ± 0.0 % ($n = 3$) (Fig. 4c).

Interestingly, although RVEP-4A-expressing sg1#a cells did not show a significant difference (p
> 0.05) from WT-expressing cells in terms of the ciliogenesis percentage, 65 ± 3% vs. 67 ± 2% (n
$= 3$) (Fig. 4d), the resulting cilia were significantly ($p < 0.05$) shorter than WT, measuring 3.3 ± 0.6
vs 4.4 ± 0.3 μm, respectively ($n = 3$)(Fig. 4e). Our observation of the RVEP-4A mutant is
consistent with a previous study where ARL13B with V358A mutation did not rescue the cilium
length in *ARL13B* null *hennin* mouse embryonic fibroblasts (Mariani et al., 2016). The partial
rescue phenotype might be due to the insufficient amount of cilium-localized ARL13B, as revealed
by its significantly ($p < 0.05$) lower *CPIR* than WT in parental RPE1 cells, at 0.61 ± 0.04 vs. 8.5 ±
0.5 ($n = 3$), respectively (Fig. 4f).

In summary, our extensive mutagenesis demonstrated that the ciliogenesis function of ARL13B
requires the N-terminal palmitoylation, the GTPase domain, and the CC. We found that cilium
localization of ARL13B mutants is necessary but insufficient for ciliogenesis, as shown by
ΔGTPase and AA1-245. Furthermore, our data on the RVEP-4A mutant suggests that cilium
formation requires a minimal amount of ciliary ARL13B, while the dose of cilium-localized ARL13B
might be a limiting factor for the cilium length.

200 Joubert syndrome mutation, R200C, abolishes the ciliogenesis function of ARL13B

We focused on the three point mutations in the coding sequence of the *ARL13B* gene, which have
been associated with the Joubert syndrome (Cantagrel et al., 2008; Thomas et al., 2015). Among
them, R79Q, and Y86C are in the GTPase domain, while R200C is in the CC. To characterize
the impact of these point mutations on the ciliogenesis function of ARL13B, we made
corresponding mutations in the background of ARL13B^R-GFP (Fig. 5a). First, we examined their
cilium localization in parental RPE1 cells, similar to our approach with other mutants. We found
that R79Q and Y86C had robust cilium localization comparable to WT (Fig. 5b). The cilium
localization of R200C was significantly weaker than that of WT (Fig. 5b). Next, we tested if
ARL13B with these Joubert syndrome point mutations can rescue the no-cilium phenotype of
sg1#a cells (Fig. 5c). Following lentivirus-mediated transduction, serum starvation, and
immunostaining, we analyzed the ciliogenesis of these mutants. R79Q and Y86C displayed
ciliogenesis percentages of 71 ± 3 ($n = 3$) and 73 ± 1 ($n = 3$) in sg1#a cells (Fig. 5d), respectively,
similar to WT (Fig. 1h). In contrast, no cilia were observed in R200C-expressing sg1#a cells,
which exhibited a ciliogenesis percentage of 0.0 ± 0.0 ($n = 3$) (Fig. 5d). In summary, our data
revealed that R200C, but not R79Q and Y86C, completely abolishes the ciliogenesis function of
ARL13B, highlighting the essential role of the CC and suggesting a cellular mechanism of R200C
mutation for the Joubert syndrome pathology.

218 Sonic hedgehog signaling is abolished in *ARL13B*-KO cells.

It is known that the SHH signaling requires the cilium (Bangs & Anderson, 2017). The stimulation
of SHH recruits SMO to the cilium and upregulates the transcription of the SHH signaling target
genes, such as *SMO*, *PTCH1*, and *GLI1* genes. Previous studies established that the loss of
ARL13B disrupts the SHH signaling in mouse (*hennin* mutant) and zebrafish (*scorpion* mutant)
models (Caspary et al., 2007; Duldulao et al., 2009), which have the phenotypes of reduced cilia.
Therefore, we want to investigate the impact of a complete cilium loss on the SHH signaling
pathway in our *ARL13B*-KO cells by assessing the cilium localization of SMO and measuring the
transcription of *SMO*, *PTCH1*, and *GLI1* genes.

In serum-starved parental RPE1 cells, we immunostained the endogenous ARL13B and SMO.
We observed that only 11 ± 3 % ($n = 4$) of ciliated cells had the cilium localization of SMO (Fig.
6a,b control). In comparison, after 24 h treatment of 0.5 mM SAG, an SMO agonist, the cilium
localization of SMO significantly ($p < 0.005$) increased to 75 ± 5 % ($n = 4$) in ciliated cells (Fig.
6a,b). Using qPCR, we found that SAG treatment after serum starvation increased the normalized
transcription levels of *SMO*, *PTCH1*, and *GLI1* genes to 2.3 ± 0.9, 1.6 ± 0.3, and 4.5 ± 3.6,
respectively, ($n = 5$, normalized by control), respectively (Fig. 6c). Our results are consistent with
what we know about the SHH signaling in RPE1 cells, thereby validating our assay setup.

Since *ARL13B*-KO sg1#a cells have no cilia, SHH signaling components, including SMO, could
not localize to the cilium. To assess the SHH signaling upon the KO of *ARL13B*, we quantified

the SHH-dependent transcription of *SMO*, *PTCH1*, and *GLI1* genes. First, we found no significant
difference in the transcript levels of *SMO*, *PTCH1*, and *GLI1* genes between parental RPE1 and
sg1#a cells (Supplementary Fig. 3). Next, we treated sg1#a and parental RPE1 cells in parallel
with serum-starvation followed by SAG, as described in Figure 6c. In contrast to the parental
RPE1 cells, we found that SAG treatment did not increase the relative transcription levels of *SMO*,
PTCH1, and *GLI1* genes; instead, it slightly reduced their levels to 0.7 ± 0.1, 0.7 ± 0.5, and $0.7 \pm$
0.2, respectively, ($n = 5$, normalized by untreated), although the transcriptional change in *SMO*
and *GLI1*, but not *PTCH1* gene, was statistically significant. The exogenous expression of
ARL13B^R-GFP rescued the SAG-induced cilium localization of SMO in sg1#a cells, as we
observed an increase in the cilium localization of SMO among GFP-positive ciliated cells from 0
% to 56 ± 5 % ($n = 3$) following SAG treatment (Fig. 6d,e), demonstrating that the observed effect
on SHH signaling is specific to *ARL13B*-KO. In summary, our results revealed that SHH signaling
is severely blunted upon the KO of *ARL13B*, suggesting that *ARL13B*-KO cells might become
irresponsive to SHH.

Discussion

ARL13B null mice and zebrafish have been reported previously to display a reduced but still appreciable ciliogenesis percentage, cilium length, and SHH signaling. Additionally, with these reports, Nozaki *et al.* made *ARL13B*-KO RPE1 cells and observed the phenotype of a reduced cilium length (Nozaki et al., 2017). However, our *ARL13B*-KO cells, generated using four sgRNAs in two cell types, consistently exhibited much more extreme phenotypes — both cilia and SHH signaling appear to be completely lost. Despite using the same cell type (RPE1) and technique (CRISPR/Cas9), our findings are starkly different from those of Nozaki *et al.* as we observed a no-cilium phenotype even after adopting the same sgRNAs.

It has recently been reported that KO clones generated by CRISPR/Cas9-induced frameshift often express a residual amount of the target protein and, therefore, can produce much milder phenotypes than expected (Meraldi, 2019; Smits et al., 2019; G. Zhang et al., 2019). The low expression of the target protein in these "pseudo-KO" clones could result from multiple mechanisms, such as alternative splicing or usage of the downstream start codon of the target mRNA. Thus, it is tempting to speculate that *ARL13B*-KO clones and null mutants generated by Nozaki *et al.* and other labs could be such cases with a low level of ARL13B protein but still sufficient to support the ciliogenesis. Hence, their phenotypes are weaker than ours. Supporting this hypothesis, we found that reducing the ciliary localization of ARL13B by mutating its CTS, the RVEP motif, did not affect the ciliogenesis but significantly shortened the cilium length (Fig. 4e). To confirm such a hypothesis, we propose three possible experiments using these "pseudo-KO" cells. First, we can enrich the residual amount of the target protein and subject it to a more sensitive Western blotting assay. Second, we can employ reverse transcription PCR to detect alternative splicing events of the *ARL13B* mRNA. Last, we predict that further knockdown of ARL13B by RNAi in these cells should produce no-cilium phenotypes like ours.

Our data revealed the absolute requirement of ARL13B in ciliogenesis. Our rescue experiments in *ARL13B*-KO cells using ARL13B mutants further suggest the roles of different ARL13B regions and domains in ciliogenesis. The palmitoylation membrane anchorage, the GTPase domain, and the CC, but not the IDR and PRR, are required for ciliogenesis. In addition, the guanine nucleotide exchange of ARL13B might not be required, as suggested by the GDP-locked mutation, T35N. The RVEP motif is essential for the cilium growth, as its mutation, RVEP-4A, reduces the ciliary localization of the resulting ARL13B and causes the short-cilium phenotype (Fig. 4e). Additionally, we examined three Joubert syndrome point mutations and found that R200C, but not R79Q and Y86C, abolished the ciliogenesis. Therefore, our data also help us to understand the pathogenesis of Joubert syndrome with R200C mutation.

Material and methods:

DNA plasmids

Please see Supplementary Table 1 for all the plasmids used in this study. pLenti-CRISPR v2 was a gift from Feng Zhang (Addgene plasmid # 52961; http://n2t.net/addgene:52961; RRID:Addgene_52961).

Antibodies

Mouse monoclonal antibodies (mAbs) against acetylated α-tubulin (Merck, #T6793, 1:1000 for immunofluorescence or IF), polyglutamylated tubulin (Adipogen, #GT335, 1:1000 for IF), CD8a (Developmental Studies Hybridoma Bank, clone OKT8, 1:200 for IF), α-tubulin (Developmental Studies Hybridoma Bank, clone 12G10, 1:1000 for Western blotting or WB), GAPDH (Santa Cruz, #sc25778, 1:1000 for WB), and SMO (Santa Cruz, #sc-166685) Rabbit polyclonal antibodies (pAbs) against Arl13b (in-house made; 1:1000 for IF)(Madugula & Lu, 2016), ARL13B (ProteinTech, #17711-AP, 1:1000 for IF and 1:3000 for WB), and pericentrin (Abcam, #ab4448, 1:1000 for IF). Horseradish peroxidase (HRP) -conjugated goat anti-mouse and anti-rabbit IgG antibodies were purchased from Bio-Rad. Alexa Fluor-conjugated goat anti-mouse and anti-rabbit IgG antibodies (1:500 for IF) were purchased from Invitrogen.

Cell culture and transfection

hTert-RPE1 and HEK293T cells were from the American Type Culture Collection. 293FT cells were from Thermo Fisher Scientific. RPE1 cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) and Ham's F12 mixture medium supplemented with 10% fetal bovine serum (FBS). HEK293T cells were cultured in DMEM supplemented with 10% FBS. 293FT were cultured in DMEM supplemented with 10% FBS and 500 µg/ml Geneticin™ (Thermo Fisher Scientific, #10131035). Transfection of plasmid DNA was performed using polyethylenimine (Polysciences) or Lipofectamine 2000 (Thermo Fisher Scientific, #11668019), according to the manufacturer's protocol. Ciliogenesis was induced by incubating cells in the corresponding medium without the FBS for 48 h (serum starvation).

Lentivirus-transduced KO and rescue of Arl13b

sgRNA cloned in pLenti-CRISPRV2 vector or ARL13B[R] WT and mutants cloned in pLVX-puro vector was transiently transfected with packaging plasmids pLP1, pLP2, and pLP/VSVG (Thermo Fisher Scientific) to 293FT cells using Lipofectamine 2000 according to manufacturer's protocol. Lentiviruses were harvested 36 and 60 h post-transfection, filtered through 0.45 µm syringe-driven filter (Sartorius), and immediately used to infect WT RPE1, HEK293T, or *ARL3B*-KO cell lines.

To make *ARL13B*-KO cell lines, RPE1 or HEK293T cells seeded in 6-well plate were infected twice every 24 h with filtered lentiviruses using 8 µg/ml hexadimethrine bromide (Sigma-Aldrich, #H9268). 24 h after the second infection, the cells were then selected in media supplemented with 8 and 1 µg/ml puromycin (Sigma-Aldrich, #P8833) for 72 h for RPE1 and HEK293T cells, respectively. Subsequently, highly diluted cells were seeded into a 96-well plate so that each well had at most one colony. Colonies were harvested, expanded, and characterized for *ARL13B*-KO using immunofluorescence and Western blotting. At least one clonally selected cell line for each sgRNA was selected and sequenced around the sgRNA target site to confirm frameshift mutations in both.

326 In rescue experiments, *ARL13B*-KO cells seeded in a 6-well plate were infected twice every 24 h
327 with filtered lentiviruses using 8 μg/ml hexadimethrine bromide. The cells were then seeded into
328 a 24-well plate containing coverslips for immunofluorescence followed by serum starvation for 48
329 h to induce ciliogenesis before immunolabeling. For the SHH signaling assay, 24 h treatment with
330 0.5 mM SAG (Abcam, #ab142160) was performed before immunolabeling.

Immunolabeling and wide-field fluorescence microscopy

RPE1 and HEK293T cells were first seeded on No. 1.5 Φ 12-mm coverslips in a 24-well plate.
Next, cells were serum starved for 48 h to induce ciliogenesis, fixed with 4% paraformaldehyde
in PBS, and subsequently neutralized with ammonium chloride. Subsequently, the cells were
incubated with primary antibodies diluted in the antibody dilution buffer, which consists of PBS
supplemented with 5% FBS, 2% bovine serum albumin, and 0.1% Saponin (Sigma-Aldrich). After
several washings, cells were incubated with secondary antibodies diluted in the antibody dilution
buffer. The coverslips were mounted in Mowiol 4-88 (EMD Millipore) after extensive washing
using PBS. Finally, cells were imaged under an inverted wide-field Olympus IX83 microscope
system equipped with a Plan Apo oil objective lens (63✗, NA 1.40), a motorized stage, a focus
drift correction device, motorized filter cubes, a scientific complementary metal oxide
semiconductor camera (Neo; Andor Technology), and a 200 W metal halide excitation light source
(Lumen Pro 200; Prior Scientific). Dichroic mirrors and filters in filter turrets were optimized for
Alexa Fluor 488 (or GFP), Alexa Fluor 594, and Alexa Fluor 680. The microscope system was
controlled by MetaMorph software (Molecular Devices).

Image analysis for the ciliogenesis percentage, the cilium length, and the *CPIR*

Untransfected cells were serum-starved to induce ciliogenesis. They were subsequently fixed and
subjected to immunofluorescence labeling for endogenous cilium markers before wide-field
microscopy imaging. The ciliogenesis percentage was calculated by dividing the number of
ciliated cells by the total number of analyzed cells. Parental or *ARL13B*-KO cells expressing
ARL13B^R-GFP WT and mutants were similarly processed and imaged. Their ciliogenesis
percentages were calculated by dividing the number of ciliated GFP-positive cells by the total
number of analyzed GFP-positive cells. The cilium length and the *CPIR* of ARL13B^R-GFP WT or
mutant were measured in ciliated GFP-positive cells using Fiji (https://fiji.sc/). To calculate cilium
length, the cilium was traced by a line, and the length of the line was measured in Fiji. The
calculation of the *CPIR* was performed as described previously (Madugula & Lu, 2016). Briefly,
first, a ~1 μm thick line was drawn orthogonally across the cilium in Fiji. Next, the line intensity
profile of the cilium was determined, and the maximum intensity value (I_{max}) was obtained.
Similarly, by drawing a circular ROI (region of interest) on the PM and the background region, the
mean intensity of PM (I_{PM}) and the background value ($I_{background}$) were acquired, respectively.
Finally, the CPIR of ARL13B^R-GFP WT or mutant was calculated as $(I_{max} - I_{PM})/(I_{PM} - I_{background})$.

Data analysis and graphs were plotted using OriginPro 9 (OriginLab). The sample size *n* is
indicated wherever applicable in figures or corresponding legends. Data are presented as the
mean ± standard deviation (s.d.). The two-tailed unpaired *t*-test was conducted in Excel
(Microsoft). A *p*-value ≤ 0.05 was considered statistically significant.

References:

Anvarian, Z., Mykytyn, K., Mukhopadhyay, S., Pedersen, L. B., & Christensen, S. T. (2019). Cellular signalling by primary cilia in development, organ function and disease. *Nat Rev Nephrol, 15*(4), 199-219. doi:10.1038/s41581-019-0116-9

Bangs, F., & Anderson, K. V. (2017). Primary Cilia and Mammalian Hedgehog Signaling. *Cold Spring Harb Perspect Biol, 9*(5). doi:10.1101/cshperspect.a028175

Cantagrel, V., Silhavy, J. L., Bielas, S. L., Swistun, D., Marsh, S. E., Bertrand, J. Y., . . . Dobyns, W. B. (2008). Mutations in the cilia gene ARL13B lead to the classical form of Joubert syndrome. *The American Journal of Human Genetics, 83*(2), 170-179.

Caspary, T., Larkins, C. E., & Anderson, K. V. (2007). The graded response to Sonic Hedgehog depends on cilia architecture. *Dev Cell, 12*(5), 767-778. doi:10.1016/j.devcel.2007.03.004

Cevik, S., Hori, Y., Kaplan, O. I., Kida, K., Toivenon, T., Foley-Fisher, C., . . . Blacque, O. E. (2010). Joubert syndrome Arl13b functions at ciliary membranes and stabilizes protein transport in Caenorhabditis elegans. *J Cell Biol, 188*(6), 953-969. doi:10.1083/jcb.200908133

Cevik, S., Sanders, A. A., Van Wijk, E., Boldt, K., Clarke, L., van Reeuwijk, J., . . . Blacque, O. E. (2013). Active transport and diffusion barriers restrict Joubert Syndrome-associated ARL13B/ARL-13 to an Inv-like ciliary membrane subdomain. *PLoS Genet, 9*(12), e1003977. doi:10.1371/journal.pgen.1003977

Chavez, M., Ena, S., Van Sande, J., de Kerchove d'Exaerde, A., Schurmans, S., & Schiffmann, S. N. (2015). Modulation of Ciliary Phosphoinositide Content Regulates Trafficking and Sonic Hedgehog Signaling Output. *Dev Cell, 34*(3), 338-350. doi:10.1016/j.devcel.2015.06.016

Duldulao, N. A., Lee, S., & Sun, Z. (2009). Cilia localization is essential for in vivo functions of the Joubert syndrome protein Arl13b/Scorpion. *Development, 136*(23), 4033-4042. doi:10.1242/dev.036350

Feig, L. A. (1999). Tools of the trade: use of dominant-inhibitory mutants of Ras-family GTPases. *Nat Cell Biol, 1*(2), E25-27. doi:10.1038/10018

Garcia-Gonzalo, F. R., Phua, S. C., Roberson, E. C., Garcia, G., 3rd, Abedin, M., Schurmans, S., . . . Reiter, J. F. (2015). Phosphoinositides Regulate Ciliary Protein Trafficking to Modulate Hedgehog Signaling. *Dev Cell, 34*(4), 400-409. doi:10.1016/j.devcel.2015.08.001

Goetz, S. C., & Anderson, K. V. (2010). The primary cilium: a signalling centre during vertebrate development. *Nature Reviews Genetics, 11*(5), 331-344.

Gotthardt, K., Lokaj, M., Koerner, C., Falk, N., Giessl, A., & Wittinghofer, A. (2015). A G-protein activation cascade from Arl13B to Arl3 and implications for ciliary targeting of lipidated proteins. *Elife, 4*. doi:10.7554/eLife.11859

Hildebrandt, F., Benzing, T., & Katsanis, N. (2011). Ciliopathies. *New England Journal of Medicine, 364*(16), 1533-1543.

Humbert, M. C., Weihbrecht, K., Searby, C. C., Li, Y., Pope, R. M., Sheffield, V. C., & Seo, S. (2012). ARL13B, PDE6D, and CEP164 form a functional network for INPP5E ciliary targeting. *Proc Natl Acad Sci U S A, 109*(48), 19691-19696. doi:10.1073/pnas.1210916109

Ivanova, A. A., Caspary, T., Seyfried, N. T., Duong, D. M., West, A. B., Liu, Z., & Kahn, R. A. (2017). Biochemical characterization of purified mammalian ARL13B protein indicates that it is an atypical GTPase and ARL3 guanine nucleotide exchange factor (GEF). *J Biol Chem, 292*(26), 11091-11108. doi:10.1074/jbc.M117.784025

Jurczyk, A., Gromley, A., Redick, S., San Agustin, J., Witman, G., Pazour, G. J., . . . Doxsey, S. (2004). Pericentrin forms a complex with intraflagellar transport proteins and polycystin-2 and is required for primary cilia assembly. *J Cell Biol, 166*(5), 637-643. doi:10.1083/jcb.200405023

Kurosaki, T., Popp, M. W., & Maquat, L. E. (2019). Quality and quantity control of gene expression by nonsense-mediated mRNA decay. *Nat Rev Mol Cell Biol, 20*(7), 406-420. doi:10.1038/s41580-019-0126-2

Larkins, C. E., Aviles, G. D., East, M. P., Kahn, R. A., & Caspary, T. (2011). Arl13b regulates ciliogenesis and
the dynamic localization of Shh signaling proteins. *Mol Biol Cell, 22*(23), 4694-4703.
doi:10.1091/mbc.E10-12-0994
Madugula, V., & Lu, L. (2016). A ternary complex comprising transportin1, Rab8 and the ciliary targeting
signal directs proteins to ciliary membranes. *Journal of cell science, 129*(20), 3922-3934.
Mahajan, D., Madugula, V., & Lu, L. (2023). Rab8 and TNPO1 are ciliary transport adaptors for GTPase
Arl13b by interacting with its RVEP motif containing ciliary targeting sequence. *J Biol Chem, 299*(5),
104604. doi:10.1016/j.jbc.2023.104604
Mariani, L. E., Bijlsma, M. F., Ivanova, A. A., Suciu, S. K., Kahn, R. A., & Caspary, T. (2016). Arl13b
regulates Shh signaling from both inside and outside the cilium. *Mol Biol Cell*. doi:10.1091/mbc.E16-03-
0189
Meraldi, P. (2019). Bub1-the zombie protein that CRISPR cannot kill. *EMBO J, 38*(7).
doi:10.15252/embj.2019101912
Nozaki, S., Katoh, Y., Terada, M., Michisaka, S., Funabashi, T., Takahashi, S., . . . Nakayama, K. (2017).
Regulation of ciliary retrograde protein trafficking by the Joubert syndrome proteins ARL13B and
INPP5E. *J Cell Sci, 130*(3), 563-576. doi:10.1242/jcs.197004
Pala, R., Alomari, N., & Nauli, S. M. (2017). Primary cilium-dependent signaling mechanisms.
International journal of molecular sciences, 18(11), 2272.
Palicharla, V. R., Hwang, S. H., Somatilaka, B. N., Legue, E., Shimada, I. S., Familiari, N. E., . . .
Mukhopadhyay, S. (2023). Interactions between TULP3 tubby domain and ARL13B amphipathic helix
promote lipidated protein transport to cilia. *Mol Biol Cell, 34*(3), ar18. doi:10.1091/mbc.E22-10-0473
Roy, K., Jerman, S., Jozsef, L., McNamara, T., Onyekaba, G., Sun, Z., & Marin, E. P. (2017). Palmitoylation
of the ciliary GTPase ARL13b is necessary for its stability and its role in cilia formation. *J Biol Chem,
292*(43), 17703-17717. doi:10.1074/jbc.M117.792937
Smits, A. H., Ziebell, F., Joberty, G., Zinn, N., Mueller, W. F., Clauder-Munster, S., . . . Huber, W. (2019).
Biological plasticity rescues target activity in CRISPR knock outs. *Nat Methods, 16*(11), 1087-1093.
doi:10.1038/s41592-019-0614-5
Sprang, S. R. (1997). G protein mechanisms: insights from structural analysis. *Annu Rev Biochem, 66*,
639-678. doi:10.1146/annurev.biochem.66.1.639
Sztul, E., Chen, P. W., Casanova, J. E., Cherfils, J., Dacks, J. B., Lambright, D. G., . . . Kahn, R. A. (2019). ARF
GTPases and their GEFs and GAPs: concepts and challenges. *Mol Biol Cell, 30*(11), 1249-1271.
doi:10.1091/mbc.E18-12-0820
Thomas, S., Cantagrel, V., Mariani, L., Serre, V., Lee, J. E., Elkhartoufi, N., . . . Attie-Bitach, T. (2015).
Identification of a novel ARL13B variant in a Joubert syndrome-affected patient with retinal impairment
and obesity. *Eur J Hum Genet, 23*(5), 621-627. doi:10.1038/ejhg.2014.156
Wheway, G., Nazlamova, L., & Hancock, J. T. (2018). Signaling through the Primary Cilium. *Frontiers in
Cell and Developmental Biology, 6*(8). doi:10.3389/fcell.2018.00008
Zhang, G., Kruse, T., Guasch Boldu, C., Garvanska, D. H., Coscia, F., Mann, M., . . . Nilsson, J. (2019).
Efficient mitotic checkpoint signaling depends on integrated activities of Bub1 and the RZZ complex.
EMBO J, 38(7). doi:10.15252/embj.2018100977
Zhang, Q., Li, Y., Zhang, Y., Torres, V. E., Harris, P. C., Ling, K., & Hu, J. (2016). GTP-binding of ARL-3 is
activated by ARL-13 as a GEF and stabilized by UNC-119. *Sci Rep, 6*, 24534. doi:10.1038/srep24534

Acknowledgments

We would like to thank the following Feng Zhang (Massachusetts Institute of Technology, USA) for sharing the DNA plasmid (pLenti-CRISPR v2) with us.

Author contributions

L. L. and D. M. conceptualization; D. M., H. M. C., and L. L. methodology; D. M. and H. M. C. analysis, investigation, and validation; D. M. and L. L. writing; L. L. supervision, project administration, and funding acquisition.

Data availability

The datasets generated during the current study are available in the GenBank repository [PV240348 to 359] and are included in Supplementary File 2. The datasets are available from the corresponding author on request.

Supplementary Information

This article contains supporting information.

Competing interests

The authors declare that they have no conflicts of interest with the contents of this article.

Abbreviations

The abbreviations used are listed below: AA, amino acid; CC, coiled-coil region; *CPIR*, the cilium to plasma membrane intensity ratio; IDR, intrinsically disordered region; KO, knock out; NS, not significant; PAL, palmitoylation lipid anchorage site; PTCH1, patched1; PRR, proline-rich region; sgRNA, single guide RNA; SHH, sonic hedgehog; SMO, smoothened; WT, wild type.

Funding

This project is supported by the Ministry of Education, Singapore, under its Tier 1 RG 25/22.

Figure Legends

Figure 1

***ARL13B*-KO RPE1 cells have no cilia. (a)** A schematic diagram of CRISPR/Cas9 strategy to make frameshift mutations in human *ARL13B* gene. The scheme of the exon organization of the human *ARL13B* gene is illustrated at the top. The middle shows parts of the WT genomic sequences of exon 1 and exon 5, where sequences of two sgRNAs, sg1 and sg2, are highlighted in light blue. Underlined sequences are protospacer-adjacent motifs or PAMs. The bottom shows genomic sequences at sgRNA target sites of two alleles in sg1#a and sg2#4 cell lines. Sequences are aligned with the corresponding WT ones. Inserted and deleted bases are colored red and indicated by "-", respectively. "+2" and "-7" indicate insertion of 2 and deletion of 7 bases, respectively. **(b)** Western blotting of endogenous ARL13B in parental and KO RPE1 cells. Cell lysates from indicated cell lines were subjected to 12% SDS-PAGE followed by immunoblotting using anti-ARL13B and anti-α-tubulin (loading control) antibodies. Molecular weight markers are labeled to the right of the immunoblot. **(c)** Fluorescence imaging of endogenous ARL13B and acetylated tubulin in parental and KO RPE1 cells. Indicated cells were fixed and co-immunostained by antibodies against indicated cilium markers before being subjected to wide-field microscopy. (**d**) Quantification of the ciliogenesis percentage in parental and KO RPE1 cells. Images acquired as described in (c) were analyzed. $n = 3$ independent experiments with ≥ 100 cells analyzed in each case. (**e**) Fluorescence imaging of endogenous glutamylated tubulin and pericentrin in parental and KO RPE1 cells, similar to (c). (**f**) Silent mutations in ARL13B^R-GFP make it resistant to sgRNAs. Regions of the ARL13B^R coding sequence, corresponding to sg1, sg2, sg4, and sg5 (colored blue), are shown with introduced multiple silent mutations colored red. Protospacer-adjacent motifs, or PAMs, are underlined. Superscript numbers indicate base pair positions in the coding sequence. (**g**) ARL13B^R-GFP can rescue the no-cilium phenotype of *ARL13B*-KO cell lines. Parental RPE1, sg1#a, and sg2#4 cells expressing ARL13B^R-GFP were processed for immunofluorescence to label endogenous acetylated tubulin and pericentrin. Arrows indicate ciliated cells. (**h-j**) ARL13B^R-GFP expressing sg1#a cells exhibit normal cilia. Images acquired as described in (g) were used for quantification. Cilia were identified by acetylated tubulin. Panel h shows the ciliogenesis percentage in GFP-positive parental and sg1#a cells (indicated by "+"). It also displays the ciliogenesis percentage of GFP-negative sg1#a cells from the same coverslips (indicated by "-"). Quantification was from $n = 3$ independent experiments with ≥ 100 cells analyzed in each experiment. In (i), the cilium length was measured in ciliated GFP-positive cells. Quantification was from $n = 3$ independent experiments with ≥ 30 cells analyzed in each experiment. In (j), the *CPIR* of ARL13B^R-GFP was quantified in ciliated GFP-positive cells. Quantification was from $n = 3$ independent experiments with ≥ 30 cells analyzed in each experiment. In (d, h-j), the error bar represents the *mean ± standard deviation*. In (c, e, and g), the scale bar represents 10 μm.

Figure 2

***ARL13B*-KO HEK293T cells have no cilia. (a)** The genotypes of *ARL13B*-KO HEK293T cell lines, sg1#5 and sg2#5. See the legend of Figure 1a. (**b-e**) Endogenous ARL13B is undetectable by Western blot and immunofluorescence imaging and no cilia could be found in sg1#5 and sg2#5 cells. In (b), anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used for loading control. See the legend of Figure 1b-e.

Figure 3

***ARL13B*-KO RPE1 cells prepared using previously reported sgRNAs have no cilia either.**
(**a**) The genotypes of *ARL13B*-KO RPE1 cell lines, sg4#2 and sg5#3, using previously published
sgRNAs. See the legend of Figure 1a. (**b-c**) Endogenous ARL13B is undetectable by
immunofluorescence imaging and no cilia could be found in sg4#2 and sg5#3 cells. See the
legend of Figure 1c,e (**d**) Exogenously expressed ARL13B^R-GFP can rescue the no-cilium
phenotype of sg4#2 and sg5#3 cells. See the legend of Figure 1g.

Figure 4

Multiple ARL13B regions or domains are required for ciliogenesis. (**a**) The schematic
diagram illustrating the organization of WT and mutant ARL13B which were constructed in the
background of ARL13B^R-GFP. The top shows the WT ARL13B. Different functional regions and
domains are indicated with AA positions. Interacting proteins previously reported are labeled
above the corresponding regions or domains and shaded green. The AA sequences of PAL and
RVEP motifs are shown below. ARL13B mutants with point mutations or truncations are illustrated
below the WT. On the right is the qualitative summary of the cilium localization of corresponding
GFP-tagged WT and mutant ARL13B (see below for details). "+" indicates a positive cilium
localization; "-" indicates a negative cilium localization or no ciliogenesis; "W" indicates a weak
cilium localization. (**b**) The cellular distribution of GFP-tagged WT and mutant ARL13B^R-GFP.
Parental and sg1#a RPE1 cells expressing GFP-tagged WT and mutant ARL13B^R-GFP were
immunostained for indicated proteins. Scale bar, 10 μm. (**c**) The cilium localization percentage
(Parental RPE1 cells) and cilium biogenesis percentage (sg1#a RPE1 cells) of GFP-positive cells.
The images acquired as described in (b) were analyzed. n = 3 independent experiments with $\geq$
100 cells analyzed in each. (**d-f**) The RVEP motif is non-essential for ciliogenesis but essential
for the cilium length and the cilium enrichment of ARL13B^R-GFP in *ARL13B*-KO cells. Images
acquired as described in (b) were quantified. GFP-positive cells were analyzed for the ciliogenesis
percentage, the cilium length, and the *CPIR* of ARL13B^R-GFP. See legend of Figure 1h-j. p-values
are from unpaired, two-tailed t-tests. Not significant or NS, p > 0.05; ∗, $p \leq 0.05$; ∗∗, $p \leq 0.005$;
∗∗∗, $p \leq 0.0005$.

Figure 5

**Joubert syndrome mutation R200C, but not R79Q and Y86C, is essential for the
ciliogenesis function of ARL13B.** (**a**) The schematic diagram illustrates the organization of
ARL13B and its three Joubert syndrome mutants. See the legend of Figure 4a. (**b-d**) The cellular
distribution, cilium localization and ciliogenesis percentage of GFP-tagged WT and Joubert
syndrome mutant ARL13B^R. See the legend of Figure 4b,c.

Figure 6

Sonic hedgehog signaling is abolished in *ARL13B*-KO cells. (**a**) SAG treatment increased the cilium localization of SMO in RPE1 cells. RPE1 cells treated with or without (control) 0.5 mM SAG for 24 hours were fixed and processed for immunofluorescence labeling of endogenous SMO and ARL13B. Arrows and arrowheads indicate SMO-positive and -negative cilia, respectively. (**b**) Percentage of ciliated cells with SMO-positive cilia. Images acquired as described in (a) were analyzed. $n = 4$ independent experiments. (**c**) Normalized transcription levels of *SMO*, *PTCH1*, and *GLI1* genes in parental RPE1 or sg1#a cells with or without (Control) 0.5 mM SAG treatment for 24 h. The transcription was quantified by qPCR and normalized by the corresponding control. $n = 5$ independent experiments. (**d-e**) Exogenously expressed ARL13B^R-GFP can rescue the SAG-induced cilium localization of SMO in *ARL13B*-KO cells. Parental RPE1 or sg1#a cells expressing ARL13B^R-GFP were treated with or without (Control) 0.5 mM SAG for 24 h. Cells were subsequently fixed and processed for immunofluorescence labeling of endogenous SMO and pericentrin (d). Arrows indicate ciliated cells. The percentage of ARL13B^R-GFP expressing ciliated cells with SMO-positive cilia is quantified in (e). $n = 3$ independent experiments with ≥ 100 cells analyzed in each. In (a,d), scale bar, 10 μm. In (b,c,e), p-values were from unpaired, two-tailed t-tests. NS, $p > 0.05$; $*, p \leq 0.05$; $** p \leq 0.005$; $*** p \leq 0.0005$.

Figure 1

Figure 2

a

ARL13B gene

exon 1   2   3   4 5 6 7   8 9 10

HEK293T cells

sg1   PAM
AGTTGCTGCGGCTGGTTC-AAGCGGTGGCGGGAGC   WT

sg1#5 [
-AGTTGCTGCGGCTGGTT--------TGGCGGGAGC   allele1: -7
-AGTTGCTGCGGCTGGTTCCAAGCGGTGGCGGGAGC   allele2: +1
]

sg2   PAM
ATGTTCAGCAATCTCGGGGTATGGAAAGAAAATT   WT

sg2#5 [
-ATGTTCAGCAATCTCGGG--------AAAGAAAATT   allele1: -7
-ATGTTCAGCAATCTCGGG-------AAGAAAATT   allele2: -7
]

b

HEK293T

parental sg1#5 sg2#5   kDa

Blot: ARL13B
*   *   *   100
                           63
                           48
                           35
                           25

                           180
                           130

Blot: GAPDH   48
                         35

c

ARL13B   Acetylated tubulin   Merge

HEK293T

sg1#5

sg2#5

d

n = 3

Ciliogenesis percentage

■ Acetylated tubulin
■ Glutamylated tubulin

HEK293T   sg1#5   sg2#5

e

Glutamylated tubulin   pericentrin   Merge

HEK293T

sg1#5

sg2#5

Figure 3

Figure 4

Figure 5

a

b

c

d

Figure 6