The influence of heavy metal stress on the evolutionary transition

of teosinte to maize

Jonathan Acosta-Bayona[1], Miguel Vallebueno-Estrada[2],

and Jean-Philippe Vielle-Calzada[1]*

[1]Grupo de Desarrollo Reproductivo y Apomixis, Unidad de Genómica Avanzada

Langebio, Centro de Investigación y de Estudios Avanzados del IPN, Km 9.6

Libramiento Norte Carretera Irapuato-León, CP36825 Irapuato Guanajuato, México

[2]Swedish University of Agricultural Sciences, Umeå, Sweden

*Author for correspondence:

Jean-Philippe Vielle-Calzada

Tel: 421663000 ext. 3100

Email: vielle@cinvestav.mx

ORCIDs

Acosta-Bayona: https://orcid.org/0009-0005-7305-6245

Vallebueno-Estrada: https://orcid.org/0000-0001-8647-8758

Vielle-Calzada: https://orcid.org/0000-0002-3017-2490

Summary

Maize originated from teosinte *parviglumis* following a subspeciation event occurred in

volcanic regions of Mesoamerica. The elucidation of the phenotypic changes that gave rise

to maize have focused on the direct consequences of domestication, with no insights on

how environmental factors could have influenced specific gene function and human

selection. We explored the impact of heavy metal (HM) stress by exposing both subspecies

to sublethal concentrations of copper and cadmium. We also assessed the genetic diversity

of loci encompassing three HM response genes affected by domestication: *ZmHMA1*,

ZmHMA7 – encoding for heavy metal ATPases of the P1$_b$ family-, and *ZmSKUs5,* encoding

a multicopper oxidase. *ZmHMA1* and *ZmSKUs5* map in the short arm of chromosome five,

in a genomic region containing multiple linked QTLs with pleiotropic effects on

domestication. A genomic analysis of the full chromosome shows that their loci show

strong positive selection as compared to previously identified domestication genes.

Exposure of teosinte *parviglumis* to HM stress results in a plant architecture reminiscent

of extant maize, and upregulation of *Teosinte branched1* (*Tb1*) in the meristem. *ZmHMA1*

and *ZmHMA7* are expressed throughout development and respond to HM stress in both

subspecies. *ZmHMA1* is mainly involved in restricting plant height and optimizing the

number of female inflorescences and seminal roots. Our results suggest that HM stress

acted on specific ATPases involved in homeostasis, giving rise to phenotypic variants that

were identified and selected by humans during domestication.

Key words: maize, teosinte *parviglumis*, heavy metals, domestication, ATPases, seminal

roots.

Introduction.

Driven by either conscious or unconscious human selection, domestication can be considered a gradual evolutionary process resulting in the adaptation to agroecological environments and anthropogenic preferences (Ross-Ibarra et al. 2007). In the case of plants, the obvious consequences of domestication are represented by visually recognizable traits that include changes in photoperiod response, reduced capacity of seed dispersal, reduced lateral branching, larger fruits or grains, reduced duration of seed dormancy, and increase in the size of female inflorescences (Pickersgill 2007; Larson et al. 2014). Notably, research efforts to understand the phenotypic changes that gave rise to maize have mainly focused on the direct consequences of domestication in aerial organs and sometimes roots (Chen et al. 2022; Lopez-Valdivia et al. 2022; Ren et al. 2022), with no emphasis on the role of adaptive environmental responses that, by acting on specific gene function, could have given rise to phenotypic traits that were identified and selected by humans.

It is generally accepted that domestication of maize (*Zea mays* ssp. *mays*) from teosinte *parviglumis* (*Zea mays* ssp. *parviglumis*) initiated in Mesoamerica around 9,000 years before present (Matsuoka et al. 2002). The initial evolutionary transition that resulted in the event of subspeciation most likely occurred within the Balsas river drainage, in regions geographically converging with the Trans-Mexican Volcanic Belt, at the intersection of the States of Mexico, Guerrero, and Michoacan (Matsuoka et al. 2002). Although the nature of the initial phenotypic changes that allowed the emergence of primitive maize remains unclear, the most obvious architectural and morphological differences between both subspecies affect their vegetative and reproductive organs. Teosinte *parviglumis* generates numerous lateral branches ending in male inflorescences, while maize often has a few short or no lateral branches, not ending in male reproductive organs (Wills et al. 2013). Additionally, the seeds of teosinte *parviglumis* are covered by a hard glume that protects the kernel externally, while in maize glumes are reduced, soft and do not cover the kernel (Doebley and Stec 1993). Large scale efforts involving physical mapping and comparative genomic analysis have resulted in the identification of numerous regions showing differential nucleotide variability between teosinte *parviglumis* and the B73 reference

maize genome. In some cases, mutations in genes located within those genomic regions result in phenotypes that reveal their involvement in domestication traits. One of the major genes affected by maize domestication is *Teosinte branched1* (*Tb1*), a transcription factor of the TCP family that controls lateral branching, sexual determination, and glume formation (Doebley et al. 1997; Martín-Trillo and Cubas 2010). *Tb1* acts as a repressor of lateral branching (Studer et al. 2017), giving rise to maize plants usually composed of a single stem. Other genes involved in domestication include *Teosinte glume architecture1 (Tga1)*, *ramosa2 (ra2)* and *Zea floricaula leafy2 (zfl2),* among others (Bomblies and Doebley 2006; Sigmon and Vollbrecht 2010; Wang et al. 2015).

The importance of phenotypic plasticity is considered as likely relevant in the domestication history of crops (Ross-Ibarra et al. 2007; Lorant et al. 2020; Mueller et al., 2023); however, the role of environmental factors influencing their origins remains elusive and poorly investigated. A previous study showed that teosinte *parviglumis* grown under low temperatures and CO_2 concentrations - reminiscent of those prevailing in the late Pleistocene and early Holocene - exhibit some phenotypes related to the domestication syndrome, including short branches, a single main stalk, and partially open seed fruitcases (Piperno et al. 2015; Piperno et al. 2019); however, the genetic basis of these changes have not been investigated. Although these results suggest that phenotypic plasticity could have influenced the domestication process, to this date there are no hints on how the adaptation to local environmental conditions could have affected specific gene function and guided human selection.

Recent genomic evidence indicates that admixture of maize with native teosinte *mexicana* populations (*Zea mays* ssp. *mexicana*) contributed to maize adaptation across a wide range of altitudes and latitudes in the American continent (Yang et al. 2023); however, the earliest maize macro-specimens found to date lack teosinte *mexicana* admixture and exhibit morphological attributes of fully domesticated cobs, indicating that *mexicana* is not a direct ancestor of maize and that its introgression occurred after initial divergence from teosinte *parviglumis* (Vallebueno-Estrada et al. 2023; Acosta-Bayona et al. 2024). Consequently, the nature of the genetic factors that guided the emergence of maize remain largely

unknown, and the initial phenotypic changes that caused the original divergence of these

two subspecies have not been identified. It is also unknown if domestication initiated by

direct action of human selection on teosinte natural standing variation, or if the response to

specific environmental factors generated phenotypic changes that were identified and

selected by humans to act on native teosinte *parviglumis* populations.

Maize can adapt to abiotic factors such as heavy metal (HM) stress through numerous

mechanisms that prevail during development (AbdElgawad et al. 2020; Gallo-Franco et

al. 2020; Malenica et al. 2021; Gao et al. 2022). These include metal immobilization by

mycorrhizal association, metal sequestration, or complexation by exuding organic

compounds from roots, metal intracellular sequestration and compartmentalization in

vacuoles. HMs include essential elements that are necessary for maize growth through

enzymatic function, such as iron (Fe), manganese (Mn), zinc (Zn), magnesium (Mg),

molybdenum (Mo), and copper (Cu); and non-essential elements like Cadmium (Cd),

chromium (Cr), lead (Pb), aluminum (Al), and selenium (Se). In general, HM stress leads

to visibly reduced plant growth due to the reduced cell elongation and cell wall elasticity.

At the molecular level, metal toxicity can disturb the cellular redox balance and lead

directly or indirectly to oxidative stress and the formation of reactive oxygen species

(ROS), shifting the redox balance to the pro-oxidative side until the antioxidative defense

system allows a new balanced redox status (Smeets et al. 2009; Hall 2002).

The comparison of the *Palomero toluqueño* and B73 genomes resulted in the identification

of a large collection of identical sequence regions (IdSRs) with low nucleotide variability

and transcriptional evidence of expression (Vielle-Calzada et al. 2009). Putative

domestication loci included *ZmHMA1* and *ZmHMA7*, two genes encoding members of the

P1$_b$-ATPases protein superfamily, also known as Heavy Metal ATPases (P-type ATPases)

characterized by the generation of a phosphorylated intermediate during the transport

reaction cycle. Members of this superfamily contain 8 to 12 transmembrane domains as

well as conserved ATP binding, phosphorylation and dephosphorylation sites (Solioz and

Vulpe 1996; Axelsen and Palmgren 2001; Voskoboinik et al. 2003; Williams and Mills

2005). The superfamily can be divided in subfamilies that include H±ATPases (type 3$_A$) in

plants and fungi, Na+/K ATPases (type $2_{C/D}$) in animals, Ca2±ATPases (type $2_{A/B}$), and

heavy metal (HM) transporting ATPases (type 1_B) capable of transporting a large variety

of HM cations across cell membranes. *ZmHMA1* and *ZmHMA7* are type 1_B ATPases that

include a conserved Cys-Pro-Cys/His/Ser motif within a transmembrane domain necessary

for the translocation of HMs cations through the membrane channel. A third *Palomero*

toluqueño/B73 IdSR included *Skewed Root Growth Similar5* (*ZmSKUs5*), a gene encoding

a Skewed5 (SKU5)-similar (SKS) protein classified as a multi-copper oxidase containing

cuperdoxin domains (Sedbrook et al. 2002). All three genes map to chromosome five

(chr.5), with *ZmHMA1* and ZmSKUs*5* within a genomic region that fractionates into

multiple QTLs under selection during the *parviglumis* to maize transition (Doebley and

Stec 1993; Lemmon and Doebley, 2014). Contrary to major domestication loci showing a

single highly pleiotropic gene responsible for important domestication traits, in this chr.5

genomic region the smallest 1.5-LOD support interval contained more than 50 genes,

suggesting that phenotypic effects are due to multiple linked QTLs (Lemmon and Doebley,

2014). While the functional characterization of all three maize genes remains elusive, the

Arabidopsis and barley *ZmHMA1* homologs act during heavy metal transport in the

chloroplast (Boutigny et al. 2014), demonstrating their functional involvement in

homeostasis.

To investigate a possible involvement of heavy metal response in the origin and

domestication of maize, we compared the phenotypic changes that teosinte *parviglumis*

and maize undergo when grown in the presence of sublethal concentrations of Cu and Cd.

We show that teosinte *parviglumis* responds to HM stress by acquiring an aerial

architecture reminiscent of extant maize, and that *Tb1* expression is upregulated in teosinte

parviglumis meristems when grown under HM stress. A large-scale genomic analysis of

nucleotide variability across maize chr.5 indicates that loci encompassing *ZmHMA1*,

ZmHMA7, and *ZmSKUs5* show reduced nucleotide variability in maize as compared to

teosinte *parviglumis*. Both *ZmHMA1* and *ZmHMA7* are active throughout maize and

teosinte *parviglumis* development, and their expression responds to HM stress. In addition,

mutations in *ZmHMA1* show phenotypic defects related to the maize domestication

syndrome, some of which are only manifested under exposure to HMs. Overall, our results

indicate that the evolutionary transition that gave rise to extant maize was influenced by
heavy metal stress through the activity of P1b-ATPases.

Results

Teosinte parviglumis responds to heavy metal stress by acquiring an architecture
reminiscent of extant maize.

To compare HM response in teosinte *parviglumis* and maize, we exposed wild-type W22
maize inbred and CIMMYTMA 29791 teosinte *parviglumis* individuals to sublethal
concentrations of Cu (400 mg/kg of $CuSO_4$) and Cd (16mg/kg of $CdCl_2$). $CuSO_4$ and $CdCl_2$
were homogenized in soil before transplant and grown under controlled greenhouse
conditions. HM concentrations were based on preliminary dose responses that allowed
maize plants to reach reproductive stages and complete their life cycle (AbdElgawad et al.
2020). As previously reported (Atta et al. 2023), maize individuals grown in the presence
of HMs showed a significant reduction in aerial traits such as stem diameter, size of leaves,
size and number of roots, and size of female inflorescences (Figure 1; Tables 2 and S2).
On average, stem diameter showed a reduction of about 17% as compared to plants grown
in absence of HM stress (Figure 1f to 1j). Similarly, the length and width of developing
leaves showed average reductions of 12%, and mature leaves of 6%. Reproductive traits
such as the number of female inflorescences (NFI) and their average weight (WFI) showed
a significant reduction (NFI = 1.20 ± 0.42 in absence of HMs and NFI = 0.70 ± 0.67 in
presence of HMs; WFI = 40.06 ± 14.39 in absence of HMs and WFI = 37.75 ± 18.83 in
presence of HMs; Table 2). As expected, the most significant effects occurred in the root
system, with a significant increase in the number of brace roots under HMs (NBR = 12.90
± 4.33 in absence of HMs; NBR = 16.30 ± 4.99 in presence of HMs; Table 2), and a
significant decrease in their length (LBR = 16.15 ± 9.75 cm in absence of HMs; LBR =
4.00 ± 2.04 cm in presence of HMs; Table 2). The number and length of crown roots were
reduced on average by 33% and 18%, respectively (NCR = 11.30 ± 2.11 in absence of
HMs; NCR = 7.60 ± 0.97 in presence of HMs; LCR = 58.90 ± 21.87 cm in absence of
HMs; LCR = 48.85 ± 11.80 cm in presence of HMs; Table 2). Nodal roots also showed a

significant reduction in their number and length (NNR = 10.50 ± 2.27 in absence of HMs;

NNR = 6.70 ± 2.79 in presence of HMs; LNR = 87.20 ± 16.80 cm in absence of HMs; LNR

= 55.30 ± 21.10 cm in presence of HMs; Tables 2 and S2). By contrast, the number (NSR)

and length (LSR) of seminal roots was significantly increased in the presence of HMs (NSR

= 4.10 ± 1.45 in absence of HMs; NSR = 6.60 ± 1.35 in presence of HMs; LSR = 27.70 ±

13.38 cm in absence of HMs; LSR = 43.50 ± 14.01 cm in presence of HMs) when measured

50 days after transplant (Table 2).

Interestingly, teosinte *parviglumis* exhibited a maize-like phenotype when grown in the

presence of HMs, with complete absence of tillering in most plants (Figure 1; Tables 1, S1

and S3). From a total of 30 plants tested, 28 showed total absence of tillering, and two

initiated secondary stems during vegetative stages but arrested their growth early during

development, resulting in the presence of a single dominant stem at reproductive stages

(Figure 1a to 1d). Additionally, all 30 plants tested showed complete absence or residual

lateral branching, with unusual proliferation of female inflorescences arising from

independent axillary meristems in primary stems (Figure 1e). This significant increase in

the number of female inflorescences clustered within a single stem (NFI = 1.80 ± 0.63 in

absence of HMs; NFI = 3.90 ± 1.66 in presence of HMs; Figure 1e and Table 1) was

associated with shorter flowering time as compared with normal growing conditions in

absence of HMs (Figure 1d). In addition, teosinte *parviglumis* grown in the presence of

HMs showed significantly reduced leaf length (LIL = 71.30 ± 11.32 cm in absence of HMs;

56.25 ± 7.80 cm in presence of HMs; Table 1). The absence of lateral branches resulted in

lower values of aerial dry weight (DWAP = 133.49 ± 28.28 gr in the absence of HMs;

DWAP = 82.72 ± 11.67 gr in presence of HMs; Table 1). Teosinte *parviglumis* grown

under HMs showed a significant reduction in the length of brace roots (LBR = 20.00 ±

11.49 cm in absence of HMs; LBR = 2.88 ± 0.66 cm in presence of HMs; Table 1), but no

significant changes in their number. The number and length of crown and nodal roots was

not affected by HM stress, nor did the presence of HMs modify the absence of seminal

roots intrinsic to teosinte *parviglumis* accessions (Perkins and Lynch 2021). Taken

together, these results suggest that the response of maize and teosinte *parviglumis* to HM

stress can give rise to phenotypes previously shown to distinguish both subspecies at the
light of the domestication syndrome.

Genomic regions encompassing ZmHMA1, ZmHMA7 and ZmSKUs5 show reduced
nucleotide variability in extant maize as compared to teosinte parviglumis.

ZmHMA1 and *ZmSKU5* mapped 52.1 and 41.8 cM, respectively, in a genetic map of chr.5
calculated to be 86.64 cM, with an average megabase pair to centimorgan ratio of 1.873
Mbp/cM (Lemmon and Doebley, 2014; Figure S1). The corresponding mapping population
used isogenic recombinant inbred lines (NIRILs) with informative cross-over events
concentrated in the region of interest and replicated phenotypic assessments in block
experiments. Multiple QTLs were detected in a region spanning across 82 cM, none with
single large effect genes such as those found in other chromosomes (Wang et al., 2005;
Studer et al. 2011; Wills et al., 2013), suggesting that several associated domestication
traits are due to multiple linked QTLs of small effect (Lemmon and Doebley, 2014). The
location of *ZmSKU5* overlaps with a single wide 1.5-LOD support interval that spreads
across 50.6 cM (number of kernels per rank; Figure S1). By contrast, *ZmHMA1* overlaps
with a contiguous region containing no less than five QTL 1.5- LOD support intervals (ear
diameter; number of kernels per rank; percentage of staminate spikelets; tassel branch
number; and tillering index; Figure S1).

To determine if the regulatory and coding regions of *ZmHMA1, ZmHMA7,* and *ZmSKUs5*
were affected by human selection, we calculated their level of nucleotide variability based
on single nucleotide polymorphisms (SNPs) for all extant maize and teosinte *parviglumis*
accessions included in the diversity panel HapMap3, using the corresponding sequences of
Tripsacum dactyloides as an outgroup. The levels of the nucleotide variability index - π(m)
and π(t) for maize and teosinte *parviglumis*, respectively - were determined in bins of 100
nt using VCFtools, including alignments encompassing a genomic region that included 15
kb upstream of the START codon, the corresponding coding region for each gene, and 15
kb downstream of the STOP codon (Figure 2 and Table S4). In the case of *ZmHMA1*, the

upstream region values were $\pi(m)$= 0.00071 and $\pi(t)$= 0.00870, and for the downstream

region, $\pi(m)$=0.00097 and $\pi(t)$=0.00248. By contrast, the *ZmHMA1* coding sequence

showed low values in both subspecies: $\pi(m)$= 0.0050 and $\pi(t)$= 0.00106. These results

indicate that human selection in the *ZmHMA1* locus preferentially affected its upstream

and downstream regulatory regions (Figure 2a and Table S4). Interestingly, although maize

π values are at least one order of magnitude lower than π values in teosinte, $\pi(t)$ values are

also one order of magnitude lower than π values of *Tripsacum sp.*, suggesting that a genetic

bottleneck also affected teosinte *parviglumis* in these loci. *ZmHMA7* showed contrasting

values of $\pi(m)$ and $\pi(t)$ across the entire coding sequence and its regulatory regions (Figure

2b). For the upstream region, values were $\pi(m)$= 0.0009 and $\pi(t)$=0.00411; for the

ZmHMA7 coding region, $\pi(m)$= 0.00015 and $\pi(t)$= 0.00148; and for the downstream region,

$\pi(m)$=0.00066 and $\pi(t)$=0.00454. *ZmSKUs5* also showed contrasting nucleotide variability

in the entire coding and flanking regions (Figure 2c and Table S4). Values were $\pi(m)$=

0.0011 and $\pi(t)$= 0.00735 for the upstream region, $\pi(m)$= 0.00044 and $\pi(t)$= 0.00754 for

the coding sequence, and $\pi(m)$=0.00203 and $\pi(t)$=0.00626 for the downstream region

(Table S4). For all three genes, similar π values in the coding region of maize landraces

and improved lines indicates that the loss of nucleotide variability occurred during the

teosinte *parviglumis* to maize transition, and not during subsequent contemporary breeding

(Table S4). By contrast, the *ZmGLB*1 neutral control locus shows equivalent levels of

nucleotide variability in maize and teosinte *parviglumis* (Eyre-Walker et al. 1998).

We also compared the level of genetic variability in *ZmHMA1*, *ZmSKUs5*, and *ZmHMA7*

(and their neighboring regions) to all chr.5 genes categorized as likely affected by

domestication (Hufford et al., 2012), and to the full collection of neutral genes distributed

across the whole chromosome. The analysis was conducted using 100 bp bins and sliding

windows of 50 bp; for a total of 385,829 bins in the case of chr.5 domestication genes, and

2,373,010 bins for neutral genes. As shown in Figure S2, the levels of nucleotide variability in

the three candidate genes and their six neighboring loci are significantly reduced in maize

landraces as compared to teosinte *parviglumis* inbred lines. When compared to the full neutral

gene set of chr.5, this reduction is more severe than the reduction affecting 304 previously

identified domestication genes, confirming the exceptional selective pressure that was imposed

in those three loci during the evolutionary transition from teosinte *parviglumis* to extant

maize.

ZmHMA1 and ZmHMA7 are active throughout development and their expression is

influenced by the presence of heavy metals.

The gene structure of *ZmHMA1, ZmHMA7,* and *ZmSKUs5* is shown in Figure 3. The coding

sequence of *ZmHMA1* and *ZmHMA7* is approximately 21.6 Kb and 4.5 Kb, respectively,

while *ZmSKUs5* is approximately 4.2 Kb in length. Within the short arm of chromosome

5, *ZmHMA1* and *ZmSKUs5* are separated by 9.16 Mb, with *ZmHMA1* approximately 15

Mb away from the maximum likelihood prediction of a QTL involved in maize

domestication (Doebley and Stec 1993; Vielle-Calzada et al. 2009). *ZmHMA7* maps to the

centromeric region of chromosome 5. A detailed PlantCARE-based analysis identified

several conserved binding sites and regulatory elements in the promoter region of these

three genes (Lescot et al. 2002). *ZmHMA1* and *ZmHMA7* share binding sites to basic Helix-

Loop-Helix (bHLH) transcription factors, as well as abscisic acid response (ABRE),

ethylene responsive (ERF), and G-box (CACGTG) elements recognized by leucine zipper

family (bZIP) and bHLH transcription factors (Figure 3a). The regulatory region of both

ZmHMA1 and *ZmSKUs5* contain a binding site for MIKC-type MADS box transcription

factors (Figure 3a), whereas *ZmHMA7* and *ZmSKUs5* share a jasmonic acid-response

TGACG-motif (Figure 3a), as well as a Myb-binding (MBS) and GATA binding site. In

addition, *ZmHMA1* contains a binding site for TCP proteins such as Tb1, increasing the

number of families of transcription factors that could be involved in shared or individual

regulation of these three genes (Figure 3a).

To elucidate the expression patterns of *ZmHMA1* and *ZmHMA7*, we conducted reverse

transcriptase PCR (RT-PCR) in leaves, stems, roots, and flowers at two vegetative (V2 and

V6) and one reproductive (R1) developmental stage, in teosinte *parviglumis* and wild-type

W22 maize individuals (Figure 3b). In the case of maize, *ZmHMA1* was expressed in

leaves, stems, roots and flowers throughout development in either absence or presence of

HMs. *ZmHMA7* was also expressed in aerial organs throughout development in both

absence and presence of HMs but was not expressed in roots at reproductive stage R1. In

teosinte *parviglumis, ZmHMA1* was also expressed throughout development in either

absence or presence of HMs. By contrast, ZmHMA*7* was constitutively expressed in aerial

organs, but only expressed in roots under HM stress during V2 and V6 stages. To further

determine if the level of expression of these two genes is influenced by the presence of

HMs, we conducted quantitative real-time PCR (qPCR) using the primary root of teosinte

parviglumis and maize plantlets at four weeks after germination (Figures 3c and 3d). In

wild-type W22 maize, the expression of *ZmHMA1* was not significantly different in the

presence and absence of HMs. By contrast, in teosinte *parviglumis* the expression of

ZMHMA1 was significantly increased in presence of HMs (Figure 3c). In the case of

ZmHMA7, expression was significantly increased in both maize and teosinte *parviglumis*

when plants were grown in the presence of HMs (Figure 3d). Overall, these results indicate

that the activity of both *ZmHMA1* and *ZmHMA7* is influenced by HM stress in teosinte

parviglumis and maize.

ZmHMA1 null mutant individuals show phenotypic defects related to the domestication

syndrome.

Transposon *Mu* insertional lines have been widely used to elucidate gene function in maize.

To elucidate the function of *ZmHMA1*, we phenotypically analyzed *LANMu1029790*, a 4-

generation backcross of line *mu1029790* into the W22 background selected for single copy

insertions in *ZmHMA1* (McCarty et al. 2005). In *LANMu1029790,* a single *Mu* insertion is

in the first exon of *ZmHMA1*, 65 bp downstream of the START codon (Figure 3a). RT-

PCR analysis showed that expression of Z*mHMA1* is completely absent in homozygous

individuals for the *LANMu1029790* insertion (Figure S3), confirming that the

corresponding mutant is a null allele. Plants harboring a homozygous *LANMu1029790*

insertion are subsequently designated as *zmhma1-m1::MuDR* (abbreviated *zmhma1-m1*).

To determine the function of *ZmHMA1* in maize, we compared the phenotype of wild-type

W22 and homozygous *zmhma1-m1* individuals at the reproductive R1 stage (Table 2).

Under normal growing conditions, homozygous *zmhma1-m1* individuals generated

significantly less leaves than wild-type (NL = 13.4 ± 0.84 in wild-type, and NL = 11.4 ±

0.84 in *zmhma1-m1; P<0.001*). Whereas the length of mutant leaves was smaller than wild-

type (LML = 77.4 ± 6.64 cm in wild-type, and LML = 69.85 ± 9.18 cm in *zmhma1-m1*),

their width was significantly larger (WML = 6.75 ± 1.3 cm in wild-type, and WML = 9.75

± 1.4 cm in *zmhma1-m1*; P<0.001), resulting in an increase of the total dry weight of aerial

organs in mutant plants (DWAP = 18.43 ± 8.04 grams in wild-type, and DWAP = 28.46 ±

9.54 grams in *zmhma1-m1*). Homozygous *zmhma1-m1* individuals also produced

significantly more female inflorescences per plant (NFI = 1.2 ± 0.42 in wild-type, and NFI

= 1.9 ± 0.32 in *zmhma1*; P<0.001) a difference that reflected in the total dry weight of

female inflorescences per plant (WFI = 40.06 ± 14.39 gr in wild-type and WFI = 69.5 ±

11.2 gr in *zmhma1*; P<0.001); and significantly more spikes per male inflorescence (NSB

= 9 ± 1.63 in wild-type, and NSB = 13.4 ± 2.32 in *zmhma1-m1*; P<0.001). Interestingly,

the general length of roots was significantly smaller in homozygous *zmhma1-m1*

individuals, in particular nodal roots (LNR = 87.2 ± 16.8 cm in wild-type, and LNR = 50.9

± 12.83 cm in *zmhma1-m1*; P<0.001); however, mutant plants showed a significant

increase in the number of seminal roots per plant (NSR = 4.1 ± 1.45 in wild-type, and NSR

= 7.3 ± 1.16 in *zmhma1-m1*; P<0.001). When wild-type and mutant individuals were grown

under HM stress, differences in the size of leaves, number of female inflorescences per

plant, and number of spikes per male inflorescence were maintained. In addition, mutant

individuals exposed to HM stress were significantly taller than wild-type (HEI = 82.2 ±

34.94 cm in wild-type, and HEI = 125.9 ± 18.81 cm in *zmhma1-m1*; P<0.01). In addition,

total root and seminal root length were equivalent in wild-type and mutant individuals,

mainly because the length of wild-type roots was significantly reduced under HM stress;

however, the number of seminal roots was again significantly higher in mutant individuals

(NSR = 6.6 ± 1.35 in wild-type, and NSR = 8.2 ± 1.62 in *zmhma1-m1*; P<0.05). These

results indicate that in maize aerial organs, *ZmHMA1* is involved in promoting leaf growth,

restricting the number of female inflorescences per plant, and restricting the number of

spikes in male inflorescences. In underground organs, *ZmHMA1* is involved in promoting

root growth and restricting the number of seminal roots per individual. Under HM stress,

ZmHMA1 shows an additional role in restricting plant height, a function that was not

identified under normal growing conditions. Overall, these results suggest that *ZmHMA1*

is involved in the control of some of the aerial and underground phenotypic traits that

distinguish maize from teosinte *parviglumis,* especially under HM stress conditions*.*

Under heavy metal stress, Tb1 is overexpressed in the apical meristem of teosinte
parviglumis.

The absence of secondary ramifications in teosinte *parviglumis* grown under HM stress is

reminiscent of the aerial architecture of wild-type maize. Previous QTL and molecular

analysis suggested that *Tb1* contributed to the architectural difference between maize and

teosinte, as *tb1* maize mutants develop secondary stem ramifications ending in male

inflorescences, resembling teosinte in their overall architecture (Doebley et al. 1997)

Interestingly, the regulatory region of *ZmHMA1* contains a specific binding site for

members of the Teosinte-branched 1/Cycloidea/Proliferating (TCP) family of transcription

factors that includes Tb1 (Figure 3a). To investigate the possibility that the phenotypic

response of teosinte *parviglumis* under HM stress could be regulated by *Tb1*, we conducted

quantitative real-time PCR (qPCR) to determine the expression of *ZmHMA1* and *Tb1* in

the shoot apical meristem (SAM) of one month old W22 maize and teosinte *parviglumis*

plantlets (Figure 4a and 4b). The expression *ZmHMA1* was downregulated in W22 wild-

type maize meristems in the presence of HMs. By contrast, HM stress resulted in a

significant increase of *ZmHMA1* expression in teosinte *parviglumis*, confirming that in the

maize ancestor the level of *ZmHMA1* activity is influenced by the presence of HMs in the

soil. In addition, the expression of *Tb1* was significantly upregulated in teosinte

parviglumis meristems in the presence of HMs, suggesting that formation of secondary

ramifications in teosinte *parviglumis* is repressed by *Tb1* function under HM stress. Taken

together, these results open the possibility for a direct positive regulation of *ZmHMA1* by

transcriptions factor of the TCP family, including *Tb1*.

Discussion

Although the evolutionary transition from teosinte *parviglumis* to maize resulted in

important phenotypic divergence between landraces and their wild progenitor, the

relevance of teosinte *parviglumis* phenotypic plasticity and its genetic consequences during

domestication remain poorly understood. Our study offers the first evidence showing that

a specific environmental factor (heavy metal stress) influenced specific gene function

(heavy metal response ATPases) and human selection during the *parviglumis* to maize

transition. The comparison of HM responses between both subspecies indicates that

teosinte *parviglumis* phenotypically responds to HM stress acquiring phenotypes

reminiscent of maize. Whereas low temperatures and CO_2 concentrations also caused a

significant decrease in lateral branching, these environmental response has not been related

to specific gene function and human selection. The absence of tillering exhibited by

teosinte *parviglumi*s under HM stress was not reported under conditions simulating an

early Holocene environment (Piperno et al., 2015). Our results also show an increase in the

number of female inflorescences per plant. This HM stress-dependent proliferation of

several closely associated female inflorescences in a single teosinte *parviglumis* stem has

not been previously reported. Female inflorescences arose from individual axillary

meristems in multiple internodes, significantly increasing teosinte *parviglumis* seed yield.

Our evidence indicates that HM stress revealed a teosinte *parviglumis* environmental

plasticity that is directly related to the function of specific HM response genes that were

affected by domestication through human selection.

We also show that *ZmHMA1*, *ZmHMA7* and *ZmSKUs5* exhibit reduced genetic diversity in

extant maize as compared to teosinte *parviglumis*, suggesting that all three loci were

affected by the evolutionary transition that gave rise to maize. The maize regulatory region

and downstream 3' area of the *ZmHMA1* locus exhibit pronounced reduction in SNPs

compared to teosinte *parviglumis* lines, likely affecting the molecular scope of its

regulatory proteins, including transcription factors of the TCP family (Martín-Trillo and

Cubas 2010). By contrast, *ZmHMA7* and *ZmSKUs5* showed a selective sweep across the

entire maize locus. Although loss of genetic diversity is usually the result of human

selection during domestication, it can also represent a consequence of natural selective

pressures favoring fitness of specific teosinte *parviglumis* allelic variants better adapted to

environmental changes and subsequently affected by human selection during the

domestication process. The significant reduction of nucleotide variability in the *ZmHMA1*

locus of teosinte *parviglumis* suggests that indeed that a natural selection pressure

Examples of edaphological factors driving genetic divergence in the teosintes or maize

include local adaptation to phosphorus concentration in *mexicana* and *parviglumis*

(Aguirre-Liguori et al. 2019), and fast maize adaptation to changing iron availability

through the action of genes involved in its mobilization, uptake, and transport (Benke and

Stich 2011).

The maize genome contains 12 genes encoding for members of the P1b family of ATPases

(Cao et al. 2019). They are subdivided according to their preferential substrate affinity: the

Copper (Cu)/Silver (Ag) group includes *ZmHMA1* to *ZmHMA4*; and the Zinc (Zn)/Cobalt

(Co)/Cadmium (Cd)/Lead (Pb) includes *ZmHMA5* to *ZmHMA12*. Initial studies suggested

that *ZmHMA1* is expressed at low levels in vegetative, reproductive or root organs

throughout development, contrary to *ZmHMA7* that is strongly expressed in developing

seeds 10 days after pollination; however, both genes appeared downregulated in plants

grown under either Cu or Cd stress (Zhiguo et al. 2018). Additional studies by Cao et al

(2019) indicated low expression of *ZmHMA1* and high levels of *ZmHMA7* expression in

12 tissues: roots, nodes, shoots, leaves, tassel, shoot apical meristem and young stem,

internodes, anthers, silks, whole seedlings, endosperm, embryo, and pericarp. Under Cd

stress, expression of both *ZmHMA1* and *ZmHMA7* was significantly upregulated in roots

and downregulated in stems (Cao et al. 2019), contradicting previous results by Zhiguo et

al. (2018). In Arabidopsis and barley, the corresponding *ZmHMA1* structural homologs

(*AtHMA1* and *HvHMA1*, respectively) can transport a wide range of metals into the

chloroplast (Mikkelsen et al. 2012; Boutigny et al. 2014). In the case of the Arabidopsis

structural homolog of ZmHMA7, AtHMA7 (also called RAN1) is proposed to act as a

metallochaperone located in the Golgi apparatus, delivering Cu to the ethylene receptor

located in the plasma membrane (Hirayama et al. 1999). Our evidence complements

previous studies by showing that, in maize, *ZmHMA1* root expression is restricted to

vegetative developmental stages, and that *ZmHMA7* is expressed in roots in response to

HM stress. We also show through qPCR analysis that, under HM stress, the expression of

both *ZmHMA1* and *ZmHMA7* is upregulated in teosinte *parviglumis* primary roots of 4-

week-old plantlets. Overall, these results confirm that both *ZmHMA1* and *ZmHMA7* are

heavy metal response genes active throughout teosinte and maize development.

In maize, the phenotypic analysis of null mutant individuals indicates that *ZmHMA1* is

involved in the control of aerial and underground developmental traits, some of which are

related to important functional adaptations that distinguish maize from its wild ancestor. In

addition to changes in leaf number, leaf length, and number of spikes per male

inflorescence, *zmhma1-m1* mutant individuals grown under normal edaphological

conditions showed a significant increase in the number of female inflorescences per plant,

a significant reduction in the length of nodal roots, and a significant increase in the number

of seminal roots per plant. Whereas the promotion of nodal root growth is a distinctive trait

of maize phenotypic plasticity and adaptation, previous studies showed that contrary to

maize, plants of teosinte *parviglumis* lack seminal roots. This absence is also characteristic

of paleobotanical maize specimens dating more than 5,000 years before present,

confirming that the acquisition of seminal roots is an important trait acquired during

domestication (Lopez-Valdivia et al. 2022). The presence of seminal roots distinguishes

extant maize from the teosintes (Burton et al. 2013), suggesting a possible involvement in

domestication (Perkins and Lynch 2021; Lopez-Valdivia et al. 2022). Our results suggest

that the allelic variants of the *ZmHMA1* locus selected during domestication could have

contributed to promote seminal root formation in maize, optimizing their number in

accordance to seed endosperm availability and carbohydrate reserves (Perkins and Lynch

2021; Lopez-Valdivia et al. 2022). Interestingly, *ZmHMA1* plays an important role in

contributing to restricting plant height in response to HM stress, a phenotype that is also

an important distinction between teosinte *parviglumis* and maize. Taken together, these

results indicate that *ZmHMA1* function was involved in the phenotypic transition from

teosinte *parviglumis* to maize through the influence of HM stress; however, a better

definition of its role in the origin and domestication of maize will require a functional

analysis of the *ZmHMA1* teosinte *parviglumis* homolog, and a detailed molecular analysis

of the corresponding allelic variants and their phenotypic effects in both subspecies.

Ideas and Speculation.

Our results point to an evolutionary model in which environmental changes in the form of

edaphological HM stress imposed natural selective forces that acted on native teosinte

parviglumis populations, causing phenotypic responses and a severe genetic bottleneck

(Figure S4). Adapted plants developed phenotypic traits such as shorter height and

emergence of seminal roots, progressively giving raise to primitive maize individuals with

reduced genetic variation in HM response genes such as *ZmHMA1*. The subsequent action

of human selection caused a drastic reduction in allelic variation of heavy metal response

genes, and genetic fixation of the resulting domestication traits.

The most important source of HMs in ancient soils of Mesoamerica is volcanic activity

through short- and long-term effects related to ores, hydrothermal water flow and ash

deposits. The history of eruptive activity in the Trans Mexican Volcanic Belt during the

early Holocene coincides with the temporal estimation of initial stages in the evolutionary

transition that gave rise to maize. Intense and recurrent eruptions occurred 24,000 to 6,000

years ago near the region of the Balsas River basin presumed to be the cradle of maize

domestication (Matsuoka et al. 2002; Macias 2005). Located less than 100 Km from the

region that harbors the natural teosinte *parviglumis* populations most closely related to

maize (Matsuoka et al. 2002), the Nevado de Toluca volcano underwent four major

eruptions approximately 24100, 12100, 10500, and 8500 years before present (yr BP;

Macias 2005), causing in the region a period of climatic instability that lasted until 8500

yr BP (Ludlow-Wiechers et al., 2005). Could volcanic eruptions during the early Holocene

have modified soil composition and influence the initial evolutionary transition from

teosinte *parviglumis* to primitive maize? The possibility of environmental cues influencing

maize emergence was mentioned by Benz and Iltis (1992), but not in relation to volcanic

activity. Deciphering the ancient DNA sequence of HM response genes such as *ZmHMA1*

in paleobotanical maize specimens found in Central Mexico could help determine if its

genetic diversity was already reduced by 5,300 to 6,200 yr BP (Ramos-Madrigal et al.,

2016 ; Vallebueno-Estrada et al., 2016; Benz, 2001). Our results open the possibility for

environmental changes caused by volcanic activity acting as a driving force in the selection

of loci involved in adaptive responses such as heavy metal tolerance.

Materials and Methods

Plant materials and growth conditions.

Seeds from maize inbred line W22, teosinte *parviglumis* accession CIMMYTMA 29791,

and insertional line *LANMu1029790* were sterilized using a 1:1 distilled water and bleach

solution, rinsed three times in water, and germinated in MS solid medium. One week-old

seedings were transplanted into PVC tubes (length: 90 cm; diameter: 20 cm) or 20-liter

pots filled with a pre-sterilized mix composed of 30% peat moss, 10% vermiculite, 10%

perlite, and 50% ground litter, complemented with 15-9-12 (N-P-K) Osmocote^{TM} fertilizer.

For HM growth conditions, 16 mg/kg of cadmium chloride (CdCl$_2$) and 400 mg/kg of

copper sulfate (CuSO$_4$) were added to the substrate before seedling transplant. All plants

were grown under glasshouse conditions at 28°C during day and 24°C during night, with

relative humidity comprised between 40 and 50%.

Phenotypic analysis.

All plants were harvested at 15 days (V2 stage), 30 days (V6 stage), and 50 days (R1 stage)

after transplant. No less than ten plants were measured for each trait and stage. The width

and length of immature leaves was measured in the 5th leaf. The length and width of mature

leaves in the 3rd leaf. The total number of leaves, total number of stem internodes, plant

height, basal stem diameter, total number of stem ramifications, and total fresh and dry

weight of the aerial biomass were estimated at the R1 stage. Estimated root traits included

the number and length of brace roots (aerial roots), the number and length of crown roots

(underground roots originating within 2 cm below the basal node), number and length of

nodal roots (underground roots originating more than 2 cm below the basal node), the

number and length of seminal roots and maximum length of the root system.

Genetic diversity analysis.

The SNP collection of maize chromosome 5 was analyzed on the basis of the HapMap

V3.2.1 dataset (Bukowski et al. 2018), selecting individuals from maize landraces (LR)

(Table S6) and teosinte *parviglumis* inbred line accessions (Teo; Table S7). A sliding

window diversity (π) was calculated using VCFtools v0.1.13 (Danecek et al. 2011) with a

window size of 100 bp (--window-pi 100) and a sliding step of 50 bp (--window-pi-step

50). Regions under selection were extracted from a list of 304 genes with domestication-

related selection signatures across the whole chromosome (Hufford et al. 2012; Tables S8),

for a total of 385,829 sliding windows (Table S9 and S10). A second set of neutral gene

regions comprising 2,373,010 windows was also extracted across the entirety of

chromosome 5. These two collections were compared to *ZmHMA1* (GRMZM2G067853),

ZmHMA7 (GRMZM2G029951), and *ZmSKUs5* (GRMZM2G076985), encompassing

2,743 windows (Table S11 and S12). Additionally, we analyzed genetic diversity in the

neighboring regions of these candidate genes, which included six genes

(*GRMZM2G701784*, *GRMZM2G014508*, *GRMZM2G367857*, *GRMZM2G375607*,

GRMZM2G374375, and *GRMZM2G067883*), totaling 3,088 windows. We employed a

custom Perl script (Vallebueno GitHub repository) to calculate two diversity metrics: fold

change (FC= π(LR)/π(Teo)) and the proportional diversity (FP=π(LR)/(π(LR) + π(Teo)).

Data visualization was performed using violin plots generated in R v4.2.2.

For an arbitrary locus encompassing 30 Kbs up and downstream of the coding sequence of

ZmMA1, *ZmHMA7*, and *ZmSKUs5*, the number of segregating sites (S), the number of

unique sequences (haplotypes, h), and the nucleotide diversity index (θ) were estimated

using DnaSP version 6.0 for all accessions included in HapMap3 (Rozas et al. 2017;

Bukowski et al. 2018). DnaSP was also used to estimate Tajima's D, Fu and Li's D, and

Fu and Li's F indexes (Tajima 1989; Fu and Li 1993). The average proportion of pairwise

nucleotide differences per nucleotide site (π) was calculated with VCFtools using bins of

100 bp and steps of 25 bp (Tajima 1983; Weir and Cockerham 1984). Null hypothesis of

equality of parameters followed two-tailed tests as previously described (Rozas et al.

2017).

Gene expression analysis.

Leaves, stems, tassels, female inflorescences, as well as primary, nodal, and crown roots

were manually isolated, frozen in liquid nitrogen and stored at 4°C. Total RNA was

extracted using TRIZOL (Thermo Fisher Scientific, USA). For reverse-transcriptase PCR

(RT-PCR), RNA was converted to coding DNA using the One-step RT-PCR kit (QIAGEN,

USA). The full list of primers is presented in Table S8. *ZmCDK* was used as a positive

control. PCR amplification was conducted using the following parameters: at 94°C for 5

min.; 94°C 30 sec., 60°C 30 sec., 72°C 1 min. for 35 cycles; final stop after 72°C for 10

min. For quantitative real-time PCR (qPCR), total RNA was isolated from 4-week-old

meristems using TRIZOL. Complementary DNA (cDNA) was synthesized using 4 μg of

total RNA, 10 mM oligodTs and Superscript reverse transcriptase II (Invitrogen, USA).

Primers for PCR were designed using the online program Primer 3 Plus (v.0.4.0) and

verified with Oligocalculator (Sigma Chemical, St Louis MO) to discard dimer formation.

PCR efficiencies of target and reference gene were determined by generating standard

curves based on serial dilutions prepared for cDNA templates. PCR efficiency was

calculated according to the slope of the standard curve (primers with 100% efficiency, the

fold equals 2). Each quantitative real-time PCR (qPCR) reaction was performed in a 10 μl

volume consisting of 5 μl of 23 SYBR Green PCR reaction Mix (Applied Biosystems,

Foster City, CA). 3.5 μl of the DNA template (100 ng/ml), 0.5 μl of the forward primer (5

mM), 0.5 μl of the reverse primer (5 mM), and 0.5 μl of ultrapure water. The qPCR

reactions were performed using the CFX96 Touch Real-Time PCR detection System and

the data were analyzed using the Bio-Rad CFX Manager software v3.1. The thermal profile

consisted of 10 min at 95°C, 40 cycles of 15 sec at 95°C, and 1 min at 60°C. Amplification

results were collected at the end of the extension step. Primer sequences for qPCR

amplification are listed in Table S7. A comparative 2-ΔΔCT method was used for

determining a relative target quantity of gene expression (Schefe et al. 2006), and the

cyclin-dependent kinase gene *ZmCDK* (GRMZM2G149286) was used as a control (Hu et
al. 2019). Reproducibility of the results was evaluated for each sample by running three
technical and three biological replicates of each of the reactions and each sample.
642

643 *Statistical Analysis.*
644 Statistical analyses were conducted using RStudio, version 4.3.0 (Horton and Kleinman
645 2015). The data are presented as mean ± standard deviation and were tested by a paired
646 Student's T-Test. When the effects were significant according to T-Test, the treatments
647 were compared with the Welch two sample T-Test at $P < 0.05$.
648

Acknowledgments
We would like to thank Ueli Grossniklaus for critical comments of this manuscript; Juana
de la Cruz for supporting greenhouse activities; Ivan López-Valdivia, Eduardo González-
Orozco and Cristian Eduardo Martínez Guerrero for help with bioinformatic analysis; and
Enrique Pola for help with qPCR analysis. All members of the Apolab offered valuable
suggestions and support. This research was funded by Conahcyt grant (CB-256826) and
the INAH-Cinvestav Tehuacán Collaborative Initiative. J.A.-B. is the recipient of a
graduate scholarship from Consejo Nacional de Humanidades Ciencia y Tecnologia
(CONAHCyT).
658

659 **Competing interests**
660 None declared.
661

Author contributions
J. A-B and J.P. V-C designed the study. J. A-B and J.P. V-C coordinated and managed the
project. M.V.E. designed and performed computational analysis. J. gfrtcdA-B performed
the experiments, J. A-B, M.V.E and J.P. V-C analyzed and interpreted the data, J. A-B and
J.P. V-C wrote the article. All authors read and approved the final version of the
manuscript.
668

List of References

AbdElgawad H, Zinta G, Hamed BA, Selim S, Beemster G, Hozzein WN, Wadaan MAM, Asard H, Abuelsoud W. 2020. Maize roots and shoots show distinct profiles of oxidative stress and antioxidant defense under heavy metal toxicity. *Environmental Pollution* 258: 113705.

Acosta-Bayona J, León-Martínez G, Vielle-Calzada JP. 2024. Origin and diversification of maize: two teosintes but different contributions. *Molecular Plant* 17: 233–235.

Aguirre-Liguori JA, Gaut BS, Jaramillo-Correa JP, Tenaillon MI, Montes-Hernández S, García- Oliva F, Hearne SJ, Eguiarte LE. 2019. Divergence with gene flow is driven by local adaptation to temperature and soil phosphorus concentration in teosinte subspecies (*Zea mays parviglumi*s and *Zea mays mexicana*). *Molecular Ecology* 28: 2814–2830.

Atta MI, Zehra SS, Ali H, Ali B, Abbas SN, Aimen S, Sarwar S, Ahmad I, Hussain M, Al-Ashkar I, Elango D, Sabagh AE, et al. 2023. Assessing the effect of heavy metals on maize (*Zea mays L.*) growth and soil characteristics: Plants-implications for phytoremediation. *PeerJ* 11: e16067.

Axelsen KB, Palmgren MG. 2001. Inventory of the superfamily of P-Type ion pumps in *Arabidopsis*. *Plant Physiology* 126: 696-706.

Benke A, Stich B. 2011. An analysis of selection on candidate genes for regulation, mobilization, uptake, and transport of iron in maize. *Genome* 54: 674–683.

Benz B. 2001. Archaelogical evidence of teosinte domestication from Guilá Naquitz, Oaxaca. *Proceedings of the National Academy of Sciences USA* 98:2104-2106.

Bomblies K, Doebley JF. 2006. Pleiotropic effects of the duplicate maize *FLORICAULA/LEAFY* genes *zfl1* and *zfl2* on traits under selection during maize domestication. *Genetics* 172: 519–531.

Boutigny S, Sautron E, Finazzi G, Rivasseau C, Frelet-Barrand A, Pilon M, Rolland N, Seigneurin-Berny D. 2014. HMA1 and PAA1, two chloroplast-envelope PIB-ATPases, play distinct roles in chloroplast copper homeostasis. *Journal of Experimental Botany* 65: 1529–1540.

Bukowski R, Guo X, Lu Y, Zou C, He B, Rong Z, Wang B, Xu D, Yang B, Xie C, Fan L, Gao S, Xu X, Zhang G, Li Y, Jiao Y, Doebley JF, Ross-Ibarra J, Lorant A, Buffalo V, Cinta Romay M, Buckler ES, Ware D, Lai J, Sun Q, Xu Y. 2018. Construction of the third-generation *Zea mays* haplotype map. *GigaScience* 7: 1-12.

Burton AL, Brown KM, Lynch JP. 2013. Phenotypic diversity of root anatomical and architectural traits in *Zea* species. *Crop Science* 53: 1042–1055.

Cao Y, Zhao X, Liu Y, Wang Y, Wu W, Jiang Y, Liao C, Xu X, Gao S, Shen Y, Lan H, Zou C, Pan G, Lin H, et al. 2019. Genome-wide identification of *ZmHMAs* and association of natural variation in *ZmHMA2* and *ZmHMA3* with leaf cadmium accumulation in maize. *PeerJ* 7: e7877.

Chen Z, Sun J, Li D, Li P, He K, Ali F, Mi G, Chen F, Yuan L, Pan Q, et al. 2022. Plasticity of root anatomy during domestication of a maize-teosinte derived population. *Journal of Experimental Botany* 73: 139–153.

Danecek, Petr, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A. DePristo, Robert E. Handsaker, et al. 2011. The Variant Call Format and VCFtools. *Bioinformatics* 27(15): 2156–2158.

Doebley J, Stec A. 1993. Inheritance of the morphological differences between maize and
teosinte: comparison of results for two F2 populations. *Genetics* 134: 559-570.

Doebley J, Stec A, Hubbard L. 1997. The evolution of apical dominance in maize. *Nature*
6724: 236-239.

Eyre-Walker A, Gaut RL, Hilton H, Feldman DL, Gaut BS. 1998. Investigation of the
bottleneck leading to the domestication of maize. *PNAS* 95: 4441-4446.

Fu Y-X, Li W-H. 1993. Statistical tests of neutrality of mutations. *Genetics* 133: 693-709.

Gallo-Franco JJ, Sosa CC, Ghneim-Herrera T, Quimbaya M. 2020. Epigenetic control
of plant response to heavy metal stress: a new view on aluminum tolerance. *Frontiers*
in Plant Science 11: 602625.

Gao C, Gao K, Yang H, Ju T, Zhu J, Tang Z, Zhao L, Chen Q. 2022. Genome-wide
analysis of metallothionein gene family in maize to reveal its role in development and
stress resistance to heavy metal. *Biological Research* 55:1.

Gaudin ACM, McClymont SA, Raizada MN. 2011. The nitrogen adaptation strategy of
the wild teosinte ancestor of modern maize, *Zea mays* subsp. *parviglumis*. *Crop*
Science 51: 2780–2795.

Hall JL. 2002. Cellular mechanisms of heavy metal detoxification and tolerance. *Journal*
of Experimental Botany 53: 1-11.

Hirayama T, Kieber JJ, Hirayama N, Kogan M, Guzman P, Nourizadeh S, Alonso
JM, Dailey WP, Dancis A, Ecker JR, et al. 1999. RESPONSIVE-TO-
ANTAGONIST1, a Menkes/Wilson disease-related copper transporter, is required for
ethylene signaling in *Arabidopsis*. *Cell* 97: 383-393.

Horton NJ, Kleinman, K. 2015. Using R and RStudio for data management, statistical
analysis, and graphics second edition. New York, US: Chapman and Hall/CRC.

Hu B, Jiang Z, Wang W, Qiu Y, Zhang Z, Liu Y, Li A, Gao X, Liu L, Qian Y, Huang
X, Yu F, Kang S, Wang Y, Xie J, Cao S, Zhang L, Wang Y, Xie Q, Kopriva S,
Chu C, et al. 2019. Nitrate–NRT1.1B–SPX4 cascade integrates nitrogen and
phosphorus signaling networks in plants. *Nature Plants* 5: 401–413.

Hufford, Matthew B., Xun Xu, Joost van Heerwaarden, Tanja Pyhäjärvi, Jer-Ming
Chia, Reed A. Cartwright, Robert J. Elshire, et al. 2012. Comparative population
genomics of maize domestication and mprovement. *Nature Genetics* 44 (7): 808–11.

Benz BF, Iltis HH. 1992. Evolution of Female Sexuality in the Maize Ear (Zea Mays L.
subsp. Mays-Gramineae). *Economic Botany* 46: 212–22.

Larson G, Piperno DR, Allaby RG, Purugganan MD, Andersson L, Arroyo-Kalin M,
Barton L, Vigueira CC, Denham T, Dobney K, Doust AN, Gepts P, Thomas M
Gilbert P, Gremillion KJ, Lucas L, Lukens L, Marshall FB, Olsen KM, Pires
JC, Richerson PJ, Rubio De Casas R, Sanjur OI, Thomas MG, Fuller DQ, et al.
2014. Current perspectives and the future of domestication studies. *PNAS* 111: 6139–
6146.

Lescot M, Déhais P, Thijs G, Marchal K, Moreau Y, Van de Peer Y, Rouzé P,
Rombauts S. 2002. PlantCARE, a database of plant *cis*-acting regulatory elements
and a portal to tools for *in silico* analysis of promoter sequences. *Nucleic Acids*
Research 30: 325–327.

Lopez-Valdivia I, Perkins AC, Schneider HM, Vallebueno-Estrada M, Burridge JD,
Gonzalez-Orozco E, Montufar A, Montiel R, Lynch JP, Vielle-Calzada JP, et al.
2022. Gradual domestication of root traits in the earliest maize from Tehuacán. *PNAS*
119: e2110245119.

Lorant A, Ross-Ibarra J, Tenaillon M. 2020. Genomics of long- and short-term adaptation in maize and teosintes. *Methods in Molecular Biology* 2090: 289-311.

Ludlow-Wiechers B, Almeida-Leñero L, Isble G. 2005. Paleoecological and climatic changes of the Upper Lerma Basin, Central Mexico during the Holocene. *Quaternary Research* 64: 318-332.

Macías JL. 2005. Geología e historia eruptiva de algunos de los grandes volcanes activos de México. *Boletín De La Sociedad Geológica Mexicana*. 107: 379-424.

Malenica N, Dunić JA, Vukadinović L, Cesar V, Šimić D. 2021. Genetic approaches to enhance multiple stress tolerance in maize. *Genes* 12:1760.

Martín-Trillo M, Cubas P. 2010. TCP genes: a family snapshot ten years later. *Trends in Plant Science* 15: 31–39.

Matsuoka Y, Vigouroux Y, Goodman MM, Sanchez JG, Buckler E, Doebley J. 2002. A single domestication for maize shown by multilocus microsatellite genotyping. *PNAS* 99: 6080-6084.

McCarty DR, Mark Settles A, Suzuki M, Tan BC, Latshaw S, Porch T, Robin K, Baier J, Avigne W, Lai J, Messing J, Koch KE, Curtis Hannah L, et al. 2005. Steady-state transposon mutagenesis in inbred maize. *Plant Journal* 44: 52–61.

Mikkelsen MD, Pedas P, Schiller M, Vincze E, Mills RF, Borg S, Møller A, Schjoerring JK, Williams LE, Baekgaard L, Holm PB, Palmgren MG, et al. 2012. Barley *HvHMA1* is a heavy metal pump involved in mobilizing organellar Zn and Cu and plays a role in metal loading into grains. *PLoS ONE* 7: e49027.

Perkins AC, Lynch JP. 2021. Increased seminal root number associated with
domestication improves nitrogen and phosphorus acquisition in maize seedlings.
Annals of Botany 128: 453–468.

Pickersgill B. 2007. Domestication of plants in the Americas: insights from mendelian and
molecular genetics. *Annals of Botany* 100: 925–940.

Piperno DR, Holst I, Winter K, McMillan O. 2015. Teosinte before domestication:
experimental study of growth and phenotypic variability in late Pleistocene and early
Holocene environments. *Quaternary International* 363: 65–77.

Ramos-Madrigal J, Smith BD, Moreno-Mayar J, Gopalakrishnan S, Ross-Ibarra J,
Gilbert MTP, Wales N. 2016. Genome sequence of a 5,310-year-old maize cob
provides insights into the early stages of maize domestication. *Current Biology*
26:3195-3201

838
Ren W, Zhao L, Liang J, Wang L, Chen L, Li P, Liu Z, Li X, Zhang Z, Li J, He K,
Zhao Z, Ali F, Mi G, Yan J, Zhang F, Chen F, Yuan L, Pan Q, et al. 2022. Genome-
wide dissection of changes in maize root system architecture during modern breeding.
Nature Plants 8: 1408–1422.

Ross-Ibarra J, Morrell PL, Gaut BS. 2007. Plant domestication, a unique opportunity to
identify the genetic basis of adaptation. *PNAS* 104: 8641-8648.

Rozas J, Ferrer-Mata A, Sanchez-Del Barrio JC, Guirao-Rico S, Librado P, Ramos-
Onsins SE, Sanchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism
analysis of large data sets. *Molecular Biology and Evolution* 34: 3299–3302.

Schefe JH, Lehmann KE, Buschmann IR, Unger T, Funke-Kaiser H. 2006.
Quantitative real-time RT-PCR data analysis: current concepts and the novel "gene
expression's CT difference" formula. *Journal of Molecular Medicine* 84: 901–910.

Sedbrook JC, Carroll KL, Hung KF, Masson PH, Somerville CR. 2002. The *Arabidopsis SKU5* gene encodes an extracellular glycosyl phosphatidylinositol-anchored glycoprotein involved in directional root growth. *Plant Cell* 14: 1635–1648.

Sigmon B, Vollbrecht E. 2010. Evidence of selection at the *ramosa1* locus during maize domestication. *Molecular Ecology* 19: 1296–1311.

Smeets K, Opdenakker K, Remans T, Van Sanden S, Van Belleghem F, Semane B, Horemans N, Guisez Y, Vangronsveld J, Cuypers A, et al. 2009. Oxidative stress-related responses at transcriptional and enzymatic levels after exposure to Cd or Cu in a multipollution context. *Journal of Plant Physiology* 166: 1982–1992.

Solioz M, Vulpe C. 1996. CPx-type ATPases: a class of P-type ATPases that pump heavy metals. *Trends in Biochemical Sciences* 21: 237-241.

Studer AJ, Wang H, Doebley JF. 2017. Selection during maize domestication targeted a gene network controlling plant and inflorescence architecture. *Genetics* 207: 755–765.

Tajima F. 1983. Evolutionary relationship of DNA sequences in finite populations. *Genetics* 105: 437-460.

Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123: 585-595.

Vallebueno-Estrada M, Rodriguez-Arevalo I, Rougon-Cardoso A, Martinez Gonzalez J, Garcia Cook A, Montiel R, Vielle-Calzada JP. 2016. The earliest maize from San Marcos Tehuacán is a aprtial domesticate with genomic evidence of inbreeding. *Proceedings of the National Academy of Sciences USA* 113:14151-14156.

Vallebueno-Estrada M, Hernández-Robles G, González-Orozco E, López-Valdivia I,
Rosales-Tham T, Vásquez-Sánchez V, Swarts K, Dillehay T, Vielle-Calzada JP,
Montiel R, et al. 2023. Domestication and lowland adaptation of coastal preceramic
maize from Paredones, Peru. *Elife* 12: e83149.

Vielle-Calzada JP, De La Vega OM, Hernández-Guzmán G, Ibarra-LacLette E,
Alvarez-Mejía C, Vega-Arreguín JC, Jiménez-Moraila B, Fernández-Cortés A,
Corona-Armenta G, Herrera-Estrella L, Herrera-Estrella A. 2009. The Palomero
genome suggests metal effects on domestication. *Science* 326: 1078.

Voskoboinik I, Fernando R, Veldhuis N, Hannan KM, Marmy-Conus N, Pearson RB,
Camakaris J. 2003. Protein kinase-dependent phosphorylation of the Menkes copper
P-type ATPase. *Biochemical and Biophysical Research Communications* 303: 337–
342.

Wang H, Studer AJ, Zhao Q, Meeley R, Doebley JF. 2015. Evidence that the origin of
naked kernels during maize domestication was caused by a single amino acid
substitution in *tga1*. *Genetics* 200: 965–974.

Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the analysis of population
structure. *Evolution* 38: 1358-1370.

Williams LE, Mills RF. 2005. P1B-ATPases - an ancient family of transition metal pumps
with diverse functions in plants. *Trends in Plant Science* 10: 491–502.

Wills DM, Whipple CJ, Takuno S, Kursel LE, Shannon LM, Ross-Ibarra J, Doebley
JF. 2013. From many, one: genetic control of prolificacy during maize domestication.
PLoS Genetics 9: e1003604.

Yang N, Wang Y, Liu X, Jin M, Vallebueno-Estrada M, Calfee E, Chen L, Dilkes BP,
Gui S, Fan X, Harper TK, Kennett DJ, Li W, Lu Y, Ding J, Chen Z, Luo J,

Mambakkam S, Menon M, Snodgrass S, Veller C, Wu S, Wu S, Zhuo L, Xiao Y,

Yang X, Stitzer MC, Runcie D, Yan J, Ross-Ibarra J, et al. 2023. Two teosintes

made modern maize. *Science* 382: 1013–1021.

Zhiguo E, Tingting L, Chen C, Lei W. 2018. Genome-wide survey and expression

analysis of P1B-ATPases in rice, maize and sorghum. *Rice Science* 25: 208–217.

921
922

Legend of Figures.

Figure 1. Phenotypic response of teosinte *parviglumis* and wild-type W22 maize grown under heavy metal stress. (**a**) to (**e**), teosinte *parviglumis*; (**f**) to (**j**), W22 maize. (**a**) Teosinte *parviglumis* in absence of heavy metal stress at early flowering stage. (**b**) Teosinte *parviglumis* grown under heavy metal stress at early flowering stage. (**c**) Teosinte *parviglumis* in absence of heavy metal stress at late flowering stage. (**d**) Teosinte *parviglumis* grown under heavy metal stress at late flowering stage. (**e**) Teosinte *parviglumis* showing proliferation of female inflorescences arising from independent axillary meristems under heavy metal stress (detail of d). (**f**) W22 maize in absence of heavy metal stress at early flowering stage. (**g**) W22 maize grown under heavy metal stress at early flowering stage. (**h**) W22 maize in absence of heavy metal stress at late flowering stage. (**i**) W22 maize grown under heavy metal stress at late flowering stage. (**j**) W22 maize female inflorescences at late flowering stage (detail of i). Scale bars: 10 cm.

Figure 2. Genetic diversity of loci encompassing *ZmHMA1, ZmHMA7,* and *ZmSKUs5.* The nucleotide variability index (π) was calculated for all HapMap3 accessions of teosinte *parviglumis* (blue line), maize (red line), and *Tripsacum dactyloides* (orange line), taking in consideration the coding sequence (arrow) and an upstream and downstream region of 15 Kb encompassing each gene. (**a**) Nucleotide variability in the *ZmHMA1* locus. (**b**) Nucleotide variability in the ZmHMA7 locus. (**c**) Nucleotide variability in the *ZmSKUs5* locus. (**d**) Nucleotide variability in the *ZmGLB1* locus.

Figure 3. Structure and gene expression of *ZmHMA1* and *ZmHMA7* in teosinte *parviglumis* and W22 maize. (**a**) Structural analysis of *ZmHMA1*, *ZmHMA7*, and *ZMSKUs5*. Blue triangles indicate the location of motifs predicted in the regulatory region of each gene. The insertion site of *Mu* line *LANMu1029790* is indicated by a red triangle. (**b**) RT-PCR analysis of *ZmHMA1* and *ZMHMA7* expression throughout development of teosinte *parviglumis* and W22 maize in presence or absence of heavy metal stress; L=leaf; S=stem; PR=primary root; NR=nodal root; CR=crown root. V2: 15 days after transplant; V6: 30 days after transplant; R1: 50 days after transplant. Controls: (-) absence of DNA;

(+) genomic DNA. (**c**) qPCR analysis of *ZmHMA1* expression in 1-month old primary roots, in absence or presence of heavy metal stress. (**d**) qPCR analysis of *ZmHMA7* expression in 1-month old primary roots, in absence or presence of heavy metal stress. * P<0.01; ** P<0.001.

Figure 4. *Tb1* and *ZmHMA1* expression in the shoot apical meristem of plantlets grown under heavy metal stress. (a) qPCR analysis of *Tb1* expression in 1-month old teosinte *parviglumis* plantlets grown in absence or presence of heavy metal stress. (**b**) qPCR analysis of *ZmHMA1* expression in 1-month old teosinte *parviglumis* and W22 maize plantlets in absence or presence of heavy metal stress. * P<0.01.

Legend of Appendix and Supplementary Files.

Figure S1. Cumulative plot of QTL detected in a region of chromosome five that includes *ZmSKUs5* and *ZmHMA1* (modified from Lemmon and Doebley, 2024).

Figure S2. Comparison of chromosome 5 nucleotide variability between maize landraces and teosinte *parviglumis*.

Figure S3. Expression of *ZmHMA1* in wild-type and homozygous *zmhma1* maize individuals.

Figure S4. Speculative model illustrating the influence of heavy metal stress on the evolutionary transition of teosinte *parviglumis* to maize.

Table S1. Additional estimation of phenotypic traits in teosinte *parviglumis* grown in absence or presence of heavy metal stress.

Table S2. Additional estimation of phenotypic traits in wild-type and *zmhma1* maize grown in absence or presence of heavy metal stress.

Table S3. Full comparison of phenotypic values for wild-type W22, *zmhma1*, and teosinte *parviglumis* individuals grown under absence or presence of heavy metal stress

Table S4. Nucleotide variability across the *ZmSKUS5, ZmHMA*1 and *ZmHMA7* locus; and their upstream and downstream neighboring genes in maize landraces, maize improved lines, and teosinte *parviglumi*s accessions included in HapMap3.

Table S5. Collection of primers used in this study.

Table S6. Maize landrace accessions used in the genetic diversity analysis chr5.

Table S7. Teosinte *parviglumis* accessions used in the genetic diversity analysis chr5.

Supplementary File 1. List of domestication genes present on chr5 as defined by Hufford et al. 2012

Supplementary File 2. FC and FP values for all domestication genes present in chr5.

Supplementary File 3. FC and FP values of neutral regions included in chr5.

Supplementary File 4. FC and FP values of candidate genes *ZmHMA1*, *ZmHMA7* and *ZmSKUs5*.

Supplementary File 5. FC and FP values of candidate neighboring genes.

Figure 1

Figure 2.

Figure 3.

Figure 4.

Table 1. Estimation of phenotypic traits in teosinte *parviglumis* grown in absence and presence of heavy metal stress.

		Description	Cu/Cd(-)		Cu/Cd(+)		
			Mean	**SD**	**Mean**	**SD**	**P Value**
Aerial traits	LIL	Length of Immature Leaf (cm)	71.30	11.92	56.25	7.80	0.0570
	SB	Stem Branches	5.30	1.83	0.60	0.84	0.0002[***]
	DWAP	Dry Weight of Aerial Part (g)	133.49	28.28	82.72	11.67	0.7430
	NFI	Number of Female Inflorescences	1.80	0.63	3.90	1.66	0.0670
	LFI	Length of Female Inflorescences (cm)	15.40	1.35	12.50	2.12	0.0650
Root traits	NBR	Number of Brace Roots	10.60	6.13	11.60	4.55	0.1210
	LBR	Length of Brace Roots (cm)	20.00	11.49	2.88	0.66	0.0045[**]
	NCR	Number of Crown Roots	12.80	4.34	9.60	1.43	0.0002[***]
	LCR	Length of Crown Roots (cm)	60.70	16.95	62.80	14.10	0.2219
	NNR	Number of Nodal Roots	16.20	4.69	14.00	3.20	0.0039[**]
	LNR	Length of Nodal Roots (cm)	83.00	15.76	104.00	8.46	0.0016[***]

** $P < 0.01$; *** $P < 0.001$.

Table 2. Estimation of phenotypic traits in wild-type and z*mhma1* maize grown in
absence and presence of heavy metal stress.

	Description	Cu/Cd(-) W22 Mean	SD	zmhma1 Mean	SD	P value	Cu/Cd(+) W22 Mean	SD	zmhma1 Mean	SD	P value
Aerial traits											
NL	Number of Leaves	13.40	0.84	11.40	0.84	0.00005***	12.00	1.70	11.10	0.74	0.14194
LML	Length of Mature Leaf (cm)	77.40	6.64	69.85	9.18	0.04930*	72.35	8.81	63.85	7.31	0.03054*
WML	Width of Mature Leaf (cm)	6.75	1.30	9.75	1.40	0.00010***	7.20	1.42	9.95	0.86	0.00006***
LIL	Length of Immature Leaf (cm)	59.50	6.10	48.25	6.88	0.00112**	51.90	9.97	45.35	9.43	0.14868
WIL	Width of Immature Leaf (cm)	7.45	0.80	9.25	0.72	0.00005***	6.60	1.10	8.85	0.85	0.00007***
HEI	Height (cm)	99.60	32.61	103.15	10.76	0.74750	82.20	34.94	125.90	18.81	0.00266**
SD	Stem Diameter (cm)	3.24	0.52	3.02	0.51	0.34945	2.60	0.53	2.86	0.59	0.31207
DWAP	Dry Weight of Aerial Part (g)	28.43	8.04	28.46	9.54	0.02042*	25.22	8.26	31.44	18.85	0.00156**
NSB	Number of Spike Branches	9.00	1.63	13.40	2.32	0.00011***	7.20	3.16	13.60	1.96	0.00004***
NFI	Number of Female Inflorescences	1.20	0.42	1.90	0.32	0.00054***	0.70	0.67	1.60	0.52	0.00357***
WFI	Weight of Female Inflorescences (g)	40.06	14.39	69.53	11.20	0.00007***	37.75	18.83	76.06	25.04	0.00113**
Root traits											
NBR	Number of Brace Roots	12.90	4.33	10.60	6.29	0.35370	16.30	4.99	9.50	6.79	0.01998*
LBR	Length of Brace Roots (cm)	16.15	9.75	15.75	14.03	0.94179	4.00	2.04	2.67	2.01	0.15890
NCR	Number of Crown Roots	11.30	2.11	12.10	2.42	0.44153	7.60	0.97	9.00	2.58	0.12569
LCR	Length of Crown Roots (cm)	58.90	21.87	50.50	13.44	0.31442	48.85	11.80	44.40	20.75	0.56281
NNR	Number of Nodal Roots	10.50	2.27	8.50	2.42	0.07262	6.70	2.79	8.00	1.56	0.21505
LNR	Length of Nodal Roots (cm)	87.20	16.80	50.90	12.83	0.00004***	55.30	21.10	53.90	31.59	0.90850
NSR	Number of Seminal Roots	4.10	1.45	7.30	1.16	0.00004***	6.60	1.35	8.20	1.62	0.02743*
LSR	Length of Seminal Roots (cm)	27.70	13.38	21.33	10.31	0.24838	43.50	14.01	53.90	20.85	0.20694

* P<0.05; ** P<0.01; *** P<0.001.

Appendix

Article title: The influence of heavy metal stress on the evolutionary transition of teosinte to maize

Authors: Jonathan Acosta-Bayona, Miguel Vallebueno-Estrada, and and Jean-Philippe Vielle-Calzada

Figure S1. Cumulative plot of QTL detected in a region of chromosome five that includes *ZmSKUs5* and *ZmHMA1* (modified from Lemmon and Doebley, 2024).

Figure S2. Comparison of chromosome 5 nucleotide variability between maize landraces and teosinte *parviglumis*.

Figure S3. Expression of *ZmHMA1* in wild-type and homozygous *zmhma1* maize individuals.

Figure S4. Speculative model illustrating the influence of heavy metal stress on the evolutionary transition of teosinte *parviglumis* to maize.

Table S1. Additional estimation of phenotypic traits in teosinte *parviglumis* grown in absence or presence of heavy metal stress.

Table S2. Additional estimation of phenotypic traits in wild-type and *zmhma1* maize grown in absence or presence of heavy metal stress.

Table S3. Full comparison of phenotypic values for wild-type W22, *zmhma1*, and teosinte *parviglumis* individuals grown under absence or presence of heavy metal stress

Table S4. Nucleotide variability across the *ZmSKUS5, ZmHMA*1 and *ZmHMA7* locus; and their upstream and downstream neighboring genes in maize landraces, maize improved lines, and teosinte *parviglumi*s accessions included in HapMap3.

Table S5. Collection of primers used in this study.

Table S6. Maize landrace accessions used in the genetic diversity analysis chr5.

Table S7. Teosinte *parviglumis* accessions used in the genetic diversity analysis chr5.

Table S8. List of domestication genes present on chr5 as defined by Hufford et al. 2012

Supplementary Files.

Supplementary File 1. List of domestication genes present on chr5 as defined by Hufford et al. 2012

Supplementary File 2. FC and FP values for all domestication genes present in chr5.

Supplementary File 3. FC and FP values of neutral regions included in chr5.

Supplementary File 4. FC and FP values of candidate genes *ZmHMA1*, *ZmHMA7* and *ZmSKUs5*.

Supplementary File 5. FC and FP values of *ZmHMA1*, *ZmHMA7* and *ZmSKUs5* neighboring genes

Figure S1. Cumulative plot of QTL detected in a region of chromosome five that includes *ZmSKUs5* and *ZmHMA1* (modified from Lemmon and Doebley, 2024). Molecular marker positions and gene location of *ZmSKUs5* and *ZmHMA1* are in centimorgans. Abbreviated QTLs are in the left margin. The 1.5-LOD support intervals for QTLs are in black horizontal bars and peak LOD scores in vertical lines. Hatched bars indicate interacting QTLs. Gray vertical shadings indicate the presence of a QTL cluster.

culm: diameter of culm
dtp: days to pollen shed
eard: ear diameter
earl: ear length
kpr: kernels per rank
krn: kernel row number
lblh: primary lateral branch legth
plht: plant height
prol: prolificacy, ears on lateral branch
splh: spikelet length
stam: % of staminate spikelets
tbn: tassel branch number
till: tillering index

Fig. S2 Comparison of chromosome 5 nucleotide variability between maize landraces and teosinte *parviglumis*. Violin plots represent all neutral regions comprised chromosome 5 (blue), domestication genes previously reported in chromosome 5 (Hufford et al., 2012; green), the three candidate genes in this study (white), and their six neighboring genes (red). ANOVA-Tukey tests are represented by letters.

Fig. S3 Expression of *ZmHMA1* in wild-type and homozygous *zmhma1* maize individuals. A fragment of *ZmHMA1* was the target of RT-PCR amplification in fourteen W22 wild-type and fourteen *zmhma1* homozygous individuals. *ZmCDK* was used as a positive control in Wild-type individual number 1. A reaction without cDNA addition was used as a negative control. Corresponding primers are presented in Table S7.

Figure S4. Model illustrating the influence of heavy metal stress on the evolutionary transition of teosinte *parviglumis* to maize.

Table S1   **Additional estimation of phenotypic traits in teosinte**

parviglumis **grown in absence or presence of heavy metal stress**.　　* P<0.05; ** P<0.01.

		Cu/Cd(-)		Cu/Cd(+)		
	Description	Mean	SD	Mean	SD	P Value
NL	Number of Leaves	11.00	2.21	11.40	1.71	0.0360*
LML	Length of Mature Leaf	84.50	4.74	81.30	8.33	0.1662
WML	Width of Mature Leaf	5.45	0.50	6.00	1.25	0.4685
WIL	Width of Immature Leaf	7.10	1.02	6.90	0.61	0.0650
NN	Number of Nodes	9.30	0.95	9.80	1.55	0.0770
HEI	Height	127.30	27.33	134.56	18.51	0.2648
SD	Stem Diameter	2.94	0.66	3.12	0.46	0.0130**
FWAP	Fresh Weight of Aerial Part	667.04	141.02	358.94	49.84	0.7397
NSB	Number of Spike Branches	22.10	4.75	24.70	4.90	0.1323
LLBS	Length of Longest Branch in Spike	21.25	4.57	20.80	5.95	0.3917
TRL	Total Root Length	105.10	7.10	94.34	44.47	0.3021
FWR	Fresh Weight of Complete Root System	244.60	101.99	225.70	40.05	0.0892
DWR	Dry Weight of Complete Root System	34.21	14.29	31.84	3.57	0.0892

(Aerial traits: NL–LLBS; Root traits: TRL–DWR)

Table S2 Additional estimation of phenotypic traits in wild-type and *zmhma1* maize grown in absence or presence of heavy metal

stress.

| | | | Cu/Cd(-) | | | | | Cu/Cd(+) | | | | |
| | | | W22 | | zmhma1 | | | W22 | | zmhma1 | | |
		Description	Mean	SD	Mean	SD	P value	Mean	SD	Mean	SD	P value
Aerial traits	NN	Number of Nodes	11.00	2.36	10.50	1.35	0.56800	9.10	2.18	10.10	1.52	0.25038
	FWAP	Fresh Weight of Aerial Part	242.56	105.73	218.93	73.36	0.56867	226.39	108.55	318.78	145.03	0.12420
	LLBS	Length of Longest Branch in Spike	21.60	1.66	17.85	4.59	0.02596*	20.45	3.74	15.75	6.99	0.07720
	LFI	Length of Female Inflorescences	24.20	2.97	24.10	3.38	0.94479	21.50	3.17	25.50	4.60	0.03618*
Root traits	TRL	Total Root Length	91.90	17.12	72.29	9.51	0.00535**	83.98	16.19	82.33	17.27	0.82821
	FWR	Fresh Weight of Complete Root System	162.20	76.45	78.76	26.24	0.00431**	104.14	47.28	79.85	31.95	0.19497
	DWR	Dry Weight of Complete Root System	6.81	3.21	5.51	1.84	0.28136	4.37	1.98	5.59	2.24	0.21342

P<0.05; ** P<0.01

Table S3 Full comparison of phenotypic values for wild-type W22, *zmhma1*, and teosinte *parviglumis* individuals grown under

absence or presence of heavy metal stress.

| | | W22 | | | | zmhma1 | | | | Teosinte *parviglumis* | | | |
| | | Cu/Cd(-) | | Cu/Cd(+) | | Cu/Cd(-) | | Cu/Cd(+) | | Cu/Cd(-) | | Cu/Cd(+) | |
	Description	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
NL	Number of Leaves	13.40	0.84	12.00	1.70	11.40	0.84	11.10	0.74	11.00	2.21	11.40	1.71
LML	Length of Mature Leaf	77.40	6.64	72.35	8.81	69.85	9.18	63.85	7.31	84.50	4.74	81.30	8.33
WML	Width of Mature Leaf	6.75	1.30	7.20	1.42	9.75	1.40	9.95	0.86	5.45	0.50	6.00	1.25
LIL	Length of Immature Leaf	59.50	6.10	51.90	9.97	48.25	6.88	45.35	9.43	71.30	11.92	56.25	7.80
WIL	Width of Immature Leaf	7.45	0.80	6.60	1.10	9.25	0.72	8.85	0.85	7.10	1.02	6.90	0.61
NN	Number of Nodes	11.00	2.36	9.10	2.18	10.50	1.35	10.10	1.52	9.30	0.95	9.80	1.55
HEI	Height	99.60	32.61	82.20	34.94	103.15	10.76	125.90	18.81	127.30	27.33	134.56	18.51
SD	Stem Diameter	3.24	0.52	2.60	0.53	3.02	0.51	2.86	0.59	2.94	0.66	3.12	0.46
SB	Stem Branches	ND	ND	ND	ND	ND	ND	ND	ND	5.30	1.83	0.60	0.84
FWAP	Fresh Weight of Aerial Part	242.56	105.73	226.39	108.55	218.93	73.36	318.78	145.03	667.04	141.02	358.94	49.84
DWAP	Dry Weight of Aerial Part	18.43	8.04	17.22	8.26	28.46	9.54	41.44	18.85	133.49	28.28	82.72	11.67
NSB	Number of Spike Branches	9.00	1.63	7.20	3.16	13.40	2.32	13.60	1.96	22.10	4.75	24.70	4.90
LLBS	Length of Longest Branch in Spike	21.60	1.66	20.45	3.74	17.85	4.59	15.75	6.99	21.25	4.57	20.80	5.95
NFI	Number of Female Inflorescences	1.20	0.42	0.70	0.67	1.90	0.32	1.60	0.52	1.80	0.63	3.90	1.66
LFI	Length of Female Inflorescences	24.20	2.97	21.50	3.17	24.10	3.38	25.50	4.60	15.40	1.35	12.50	2.12
WFI	Weight of Female Inflorescences	40.06	14.39	37.75	18.83	69.53	11.20	76.06	25.04	1.29	0.74	1.29	0.56
NBR	Number of Brace Roots	12.90	4.33	16.30	4.99	10.60	6.29	9.50	6.79	10.60	6.13	11.60	4.55
LBR	Length of Brace Roots	16.15	9.75	4.00	2.04	15.75	14.03	2.67	2.01	20.00	11.49	2.88	0.66
NCR	Number of Crown Roots	11.30	2.11	7.60	0.97	12.10	2.42	9.00	2.58	12.80	4.34	9.60	1.43
LCR	Length of Crown Roots	58.90	21.87	48.85	11.80	50.50	13.44	44.40	20.75	60.70	16.95	62.80	14.10
NNR	Number of Nodal Roots	10.50	2.27	6.70	2.79	8.50	2.42	8.00	1.56	16.20	4.69	14.00	3.20
LNR	Length of Nodal Roots	87.20	16.80	55.30	21.10	50.90	12.83	53.90	31.59	83.00	15.76	104.00	8.46
NSR	Number of Seminal Roots	4.10	1.45	6.60	1.35	7.30	1.16	8.20	1.62	ND	ND	ND	ND
LSR	Length of Seminal Roots	27.70	13.38	43.50	14.01	21.33	10.31	53.90	20.85	ND	ND	ND	ND
TRL	Total Root Length	91.90	17.12	83.98	16.19	72.29	9.51	82.33	17.27	105.10	7.10	94.34	44.47
FWR	Fresh Weight of Complete Root System	162.20	76.45	104.14	47.28	78.76	26.24	79.85	31.95	244.60	101.99	225.70	40.05
DWR	Dry Weight of Complete Root System	6.81	3.21	4.37	1.98	5.51	1.84	5.59	2.24	34.21	14.29	31.84	3.57

Aerial traits (NL–WFI); Root traits (NBR–DWR)

Table S4 Nucleotide variability across the *ZmSKUS5, ZmHMA*1 and *ZmHMA7* locus; and their upstream and downstream neighboring genes in maize landraces, maize improved lines, and teosinte *parviglumi*s accessions included in HapMap3.

Gene ID	GeneLength(Kp)	DistanceOfGene(Kp)	Upstream(30Kb)			Coding Region			Downstream(30Kb)		
			π(Maize)	π(Teosinte)	**πM/πT**	π(Maize)	π(Teosinte)	**πM/πT**	π(Maize)	π(Teosinte)	**πM/πT**
GRMZM2G375607[a]	0.17	37.3	0.00105	0.000675	1.5556	0.037894	0.0304718	1.2436	0.00041	0.00064	0.6406
GRMZM2G076985[b]	4.17	-	0.011884529	0.017749593	0.669566282	0.000437979	0.007541557	0.058075436	0.002030405	0.006264344	0.324120984
GRMZM2G014508[c]	22.8	97.2	0.00844561	0.005282053	1.598925645	0.037149063	0.011257665	3.299890491	0.002433718	0.017648094	0.13790262
GRMZM2G367857[d]	8.72	25.7	0.005243813	0.006394847	0.820006014	0.008991137	0.011367691	0.790937843	0.002425106	0.006884682	0.352246578
GRMZM2G067853[e]	21.6	-	0.01329488	0.01772011	0.750270738	0.005748	0.0010578	5.433919455	0.00096874	0.0024874	0.389458873
GRMZM2G067883[f]	0.78	24.9	0.005048118	0.007825611	0.645076531	0.00119172	0.000102418	11.63584526	0.003030221	0.00681471	0.444658834
GRMZM2G374375[g]	1.84	228.6	0.0007481	0.000974	0.768069815	0.00974	0.01495	0.651505017	0.00054	0.00948	0.056962025
GRMZM2G029951[h]	4.5	-	0.000678	0.010784	0.06287092	0.00015	0.00148	0.101351351	0.00066	0.00454	0.145374449
GRMZM2G701784[i]	0.54	56	0.0005974	0.009748	0.061284366	0.00074	0.01784	0.041479821	0.00017	0.00009	1.888888889
GRMZM2G067919[j]	2.4	-	0.00611	0.00809	0.755253399	0.01035	0.01058	0.97826087	0.01417	0.01254	1.129984051

[a]Uncharacterized gene; [b]*ZmSKUS5* (Multicopper oxidase); [c]*ZmDds_sub1* (Decaprenyl-diphosphate synthase subunit 1); [d]Uncharacterized gene; [e]*ZmHMA1;* [f]Uncharacterized gene; [g]*Zmpgl20* (Exopolygalacturonase-like); [h]*ZmHMA7*; [i]Uncharacterized gene; [j]*ZmGLB1* (Globulin1)

Table S5. Collection of primers used in this study.

Primer Name	Gene ID	Forward	Reverse
*ZmHMA1*_PCR/qPCR	GRMZM2G067853_T02	TGAACACTCCGACATGGTCC	GGTGTAGCTTGAGGGAGAGC
*ZmHMA7*_PCR/qPCR	GRMZM2G029951_T01	ATCTCCTGCGCTTCTTGTGC	CCTCTGGCCTGTACTGAACC
*ZmTb1*_qPCR	Zm00001eb054440	CCCATGGACTTACCGCTTTA	GGGAGCATGGGTAGCAGTAG
*ZmCDK*_PCR	Zm00001eb215750_T001	GGAAGGTATGCACAGGACAGAT	TTCAGCACAATCTTGGCAAAAC
*ZmCDK*_qPCR	Zm00001eb215750_T001	CCGTCATCGCCTCACGAAGAG	AGAGCCTGCCTTACGGAATTGG
ZmHMA1 genotyping	GRMZM2G067853_T02	GACGACACAAGCCAAAACAG	AGGGAGATGCAGCAGAGGT

Table S6. Maize landrace accessions used in the genetic diversity analysis of chr5.

1	BKN009
2	BKN010
3	BKN011
4	BKN014
5	BKN015
6	BKN016
7	BKN017
8	BKN018
9	BKN019
10	BKN020
11	BKN022
12	BKN023
13	BKN025
14	BKN026
15	BKN027
16	BKN029
17	BKN030
18	BKN031
19	BKN032
20	BKN033
21	BKN034
22	BKN035
23	BKN040

Table S7. Teosinte *parviglumis* accessions used in the genetic diversity analysis of chr5.

1	TIL01
2	TIL01-JD
3	TIL02
4	TIL03
5	TIL04-TIP285
6	TIL04-TIP454
7	TIL05
8	TIL06-TIP260
9	TIL06-TIP496
10	TIL07
11	TIL09
12	TIL10
13	TIL11
14	TIL12
15	TIL14-TIP498
16	TIL15
17	TIL16
18	TIL17