Neural Correlates of Social Withdrawal and Preference for Solitude in

Adolescence

Matthew Risner, BS[1], Catherine Stamoulis, PhD[1,2*]

[1]Department of Pediatrics, Division of Adolescent and Young Adult Medicine, Boston

Children's Hospital, Boston, MA, 02115, USA

[2]Department of Pediatrics, Harvard Medical School, Boston, MA, 02115, USA

*To whom correspondence should be addressed:

caterina.stamoulis@childrens.harvard.edu

Abstract

Social isolation during development, especially in adolescence, has detrimental but incompletely understood effects on the brain. This study investigated the neural correlates of preference for solitude and social withdrawal in a sample of 2,809 youth (median (IQR) age = 12.0 (1.1) years, 1440 (51.26%) females) from the Adolescent Brain Cognitive Development study. Older youth whose parents had mental health issues more frequently preferred solitude and/or were socially withdrawn (β=0.04–0.14, CI=[0.002,0.19], p<0.05), both of which were associated with internalizing and externalizing behaviors, depression, and anxiety (β=0.25-0.45, CI=[0.20,0.49], p<0.05). Youth who preferred solitude and/or were socially withdrawn had lower cortical thickness in regions supporting social function (cuneus, insula, anterior cingulate and superior temporal gyri) and/or mental health (β=-0.09 to -0.02, CI=[-0.14,-0.003], p<0.05), and higher amygdala, entorhinal cortex, parahippocampal gyrus, and basal ganglia volume (β=2.62-668.10, CI=[0.13,668.10], p<0.05). Youth preferring solitude had more topologically segregated dorsal attention, temporoparietal, and social networks (β=0.07-0.10, CI=[0.02,0.14], p≤0.03). Socially withdrawn youth had a less topologically robustness and efficient (β=-0.05 to -0.80, CI=[-1.34,-0.01], p<0.03), and more fragile cerebellum (β=0.04, CI=[0.01,0.07], p<0.05). These findings suggest that social isolation in adolescence may be a risk factor for widespread alterations in brain regions supporting social function and mental health.

Keywords

• Adolescent brain

• Social withdrawal

• Preference for solitude

• Resting-state brain networks

• Brain Structures

• Mental Health

Word count: 4991

1. INTRODUCTION

Adolescence is a period of extensive physical, cognitive, and social changes (Best & Ban, 2021; Blakemore & Mills, 2014a; Klimstra et al., 2010; Luna, 2009; Özdemir et al., 2016; Pfeifer & Allen, 2021; Rogol et al., 2002; Siervogel et al., 2004; Steinberg & Morris, 2001; Susman & Rogol, 2004; Wigfield et al., 2006). It is also a period of heightened neural maturation and circuit reorganization (Andrews et al., 2021; Ashtari et al., 2007; Blakemore, 2008, 2012; Goddings et al., 2014; Lenroot et al., 2007; Mills et al., 2016; Peper et al., 2011; Sturman & Moghaddam, 2011; Vijayakumar et al., 2018; Wierenga et al., 2014; Yurgelun-Todd, 2007). Concurrently, the youth social world expands and changes, in part due to social reorientation –a shift in the relative importance of peer relationships compared to those with family (Albert et al., 2013; Giordano, 2003; Jaccard et al., 2005; Lerner & Steinberg, 2009; Maxwell, 2002; Sawyer et al., 2018; Wang et al., 1995). Changes in the youth social environment can be overwhelming, sometimes leading to social isolation (Barzeva et al., 2019; Biggs et al., 2012; Oh et al., 2008; Porcelli et al., 2019; Rubin et al., 2009; Wood et al., 2022). Although spending time alone may be motivated by a desire for self-reflection or creative endeavors, habitual withdrawal driven by social stressors, for example persistent peer rejection (Borg & Willoughby, 2022; Vasey, 2001), can lead to social isolation, which can have profound adverse effects on social development (Almeida et al., 2021; Orben et al., 2020; Thompson et al., 2022). The adolescent brain is vulnerable to social stressors, which increase the risk of circuit miswiring, and mental

health and behavioral issues (Blakemore, 2019; Bor et al., 2014; Costello et al., 2011;

Reiss, 2013).

Negative factors in the youth's immediate environment increase the risk for social

withdrawal, isolation, and loneliness. Children whose parents have mental health issues

may be at higher risk of behavioral and mental health problems, including social

isolation (Manning & Gregoire, 2009; Matthews et al., 2015; Van Loon et al., 2014).

Stigma surrounding parental mental illness can cause children to face bullying,

embarrassment, and guilt. This often leads to concealing their family circumstances and

avoiding support, further exacerbating isolation (Bosch et al., 2017; Cogan et al., 2005;

Dam et al., 2018; Reupert et al., 2021). Relationships with peers are similarly influential,

and peer rejection, victimization, lacking friends, and low friendship quality also

contribute to loneliness (Asher & Paquette, 2003; Cassidy & Asher, 1992; Cheng &

Furnham, 2002; Lodder et al., 2017; Schwartz-Mette et al., 2020; Vanhalst et al., 2014;

Woodhouse et al., 2012). These relationships are, however, bidirectional since social

withdrawal may worsen the quality and/or depth of friendships (Barzeva et al., 2022;

Biggs et al., 2012; Coplan et al., 2018; Rubin et al., 2006). Social exclusion at school

may lead to social withdrawal and loneliness (G. Arslan et al., 2023; Ph. D. Arslan

Gökmen, 2021; Ford et al., 2018; Patton et al., 2006). In contrast, a sense of school

connectedness may mitigate the adverse psychological effects of social isolation

(Foster et al., 2017; Hall-Lande et al., 2007; London & Ingram, 2018; Marraccini & Brier,

2017; Preston & Rew, 2022). These associations are bidirectional, as socially withdrawn

children often encounter difficulties in school, including poor relationships with teachers,

academic struggles, and may avoid school altogether. These challenges further

reinforce their social isolation and limit opportunities for meaningful social engagement

(Coplan et al., 2018; Rubin et al., 2009; Stenseng et al., 2022).

Lack of social interaction increases the risk of physical, cognitive, and mental health

problems (Christiansen et al., 2021; Fuhrmann et al., 2019; Hall-Lande et al., 2007;

Houghton et al., 2022; Rubin et al., 2009; Stenseng et al., 2022). Social isolation and

loneliness have been associated with increased risk for cancer, obesity, diabetes,

asthma, migraine, hypertension, and back pain in adolescents and young adults

(Almeida et al., 2021; Christiansen et al., 2021; Goosby et al., 2013; Mahon et al., 1993;

Mushtaq et al., 2014; von Soest et al., 2020). They have also been associated with

longer-term physical issues, including obesity, cardiovascular problems, inflammation,

and impaired immunoregulation in adulthood (Caspi et al., 2006; Danese et al., 2009;

Goosby et al., 2013; Hawkley et al., 2010; Hawkley & Cacioppo, 2010; Lacey et al.,

2014; Matthews et al., 2024; Patterson & Veenstra, 2010).

Social isolation may also significantly increase the risk of mental health issues, In

adolescence, during which many common mental health disorders emerge (Blakemore,

2019; de Girolamo et al., 2012; Paus et al., 2008), it has been linked to anxiety,

depression, internalizing problems, and suicidal ideation and self-harm (Blakemore,

2019; Bor et al., 2014; Cacioppo et al., 2006; Costello et al., 2011; Endo et al., 2017;

Gallagher et al., 2014; Harman et al., 2021; Jones et al., 2011; McClelland et al., 2020;

Reiss, 2013; Rubin et al., 2009; Schinka et al., 2012). Social isolation has also been

associated with cognitive deficits, including worse academic performance and impaired

reward processing and language skills (Almeida et al., 2021; Jefferson et al., 2023;

Matthews et al., 2023; Rubin et al., 2009; Tomova et al., 2022). In addition, recent

studies focusing on social isolation of youth during the COVID-19 pandemic have

identified associations between pandemic-related social isolation and deficits in

executive function, attention, and memory (Lavigne-Cerván et al., 2021; Murtaza et al.,

2023).

The neural correlates of social isolation (resulting from social withdrawal and/or

preference for solitude) in the adolescent brain remain poorly understood. This is a

significant gap in knowledge given that adolescence represents a formative period for

the establishment of social identity and mental health outcomes (Blakemore, 2019;

Blakemore & Mills, 2014b; Branje et al., 2021; Fergusson et al., 2005; Kessler et al.,

2005; Kroger, 2004; Meeus, 1996; Paus et al., 2008; Schlack et al., 2021). Adult studies

have shown that loneliness and social withdrawal are associated with lower gray matter

volume in the left posterior superior temporal, the amygdala, hippocampus and

cerebellum (Düzel et al., 2019; Kanai et al., 2012; Koolschijn et al., 2013; Lind et al., 2020),

disrupted white matter integrity in the salience network (Tian et al., 2014), and lower

white matter density in the inferior parietal lobule and anterior insula (Lam et al., 2021;

Nakagawa et al., 2015). It has also been associated with changes in gray matter volume

and lower white matter density in the dorsolateral prefrontal cortex, likely leading to

emotional dysregulation (Kong et al., 2015; Liu et al., 2016; Zheng et al., 2023). Social

isolation may also affect emotional processing in adolescents through its impact on

anatomical connections between the amygdala and the orbitofrontal cortex (Goetschius

et al., 2020). Together, these findings suggest that social isolation adversely impacts

the structural integrity of brain regions that play critical roles in social perception,

cognitive control, and emotional processing.

Beyond its impact on brain structure, social withdrawal, isolation and loneliness in

young adults has been linked to higher resting-state functional connectivity in the right

central operculum, supramarginal gyrus, and circuits involved in sustained attention

(Layden et al., 2017), and weaker connections between dorsal attention and salience

networks, as well within and between temporal, limbic, and prefrontal networks (Feng et

al., 2019; McIver et al., 2019; Tian et al., 2017). In adolescents, social withdrawal has

been associated with aberrant stronger connections between the left amygdala and

regions of the dorsal attention, frontoparietal control, and auditory networks, and

between the prefrontal and limbic networks (Thomas et al., 2024). Loneliness been

associated with lower activation of the ventromedial prefrontal cortex (Golde et al.,

2019) and higher functional connectivity in visual attention networks (Brilliant T. et al.,

2022).

Despite valuable insights provided by prior studies, adolescent brain correlates of social

isolation (as a result of withdrawal and/or preference for solitude) remain incompletely

understood and have not been systematically characterized. A significant barrier to

robust investigations is the heterogeneity of adolescent brain development, and the

complexity of associations between social behaviors, environmental factors, and the

brain. However, the availability of multimodal (including neuroimaging) data from the

Adolescent Brain Cognitive Development (ABCD) study, a historically large (~12,000

youth) investigation of adolescent brain development (Casey et al., 2018) provides a

unique opportunity to bridge this gap in knowledge, and investigate relationships

between social isolation and brain development during this complex period.

The present study leveraged the ABCD structural MRI and resting-state fMRI data,

mental health assessments and survey data on social behaviors and environmental

factors, in order to investigate associations between preference for solitude and social

withdrawal (both leading to social isolation), and the organization of intrinsically

coordinated brain networks, and their structural correlates in almost 3,000 adolescents

(ages 11-12 years), who at this age may be at increased risk of mental health disorders

and maladaptive social behaviors (Kessler et al., 2005). The study hypothesized that

preference for solitude and social withdrawal are in part driven by negative factors in the

youth social environment and are associated with widespread structural and functional

brain changes, especially in underdeveloped regions involved in social function.

175 2. MATERIALS AND METHODS

176 This study analyzed publicly available and thus anonymized data, and was approved by

177 the Institutional Review Board.

2.1 Participants

A sample of n = 2809 youth from the ABCD study, measured at the two-year follow-up, was analyzed. The primary criterion for inclusion was availability of at least one high-quality (≤10% of frames censored for motion) 5-min resting-state fMRI run that had been collected prior to the onset of the COVID-19 pandemic (assumed to be March 11, 2020, when the World Health Organization declared it as such). Youth measured during the pandemic were excluded to eliminate its potential confounding effects on the brain and social behaviors. Although studying the effects of social isolation during the pandemic on the brain is critical, it was outside the scope of the present study. In addition, participants with clinical MRI findings and neuropsychiatric or neurodevelopmental disorders were also excluded, since they may independently affect brain morphology and the organization of neural circuits (Brooks et al., 2021, 2022, 2023). The final analytic sample included 1366 biological boys (48.63%) and 1440 girls (51.26%), with a median age of 12.0 years (Interquartile Range (IQR) = 1.1 years). Over half of youth were white non-Hispanic (1488 (53.0%)), 1069 (38.1%) were non-Hispanic racial minorities, and 243 (8.7%) were Hispanic. Socioeconomic status was assessed based on annual household income, with a median income $75,000-$99,999. The sample included 515 (18.3%) youth in pre-puberty, 609 (21.7%) in early puberty, 918 (32.7%) in mid-puberty, and 622 (22.1%) in late/post-puberty. Detailed demographic and other participant characteristics are provided in Table 1.

2.2 Measures of social withdrawal

Two measures directly assessing social withdrawal and preference for solitude were extracted from the Child Behavior Checklist (CBCL), completed by parents: *"withdrawn, doesn't get involved with others"* and *"would rather be alone than with others."* Both were rated on a Likert scale from not true (0) to very/often true (2).

2.3 Measures of mental health

Data on common mental health problems that were adequately represented in the study sample, including anxiety, depression, internalizing and externalizing behaviors (all t-scores) were extracted from the CBCL. Additional information on social anxiety disorder and anhedonia was extracted from the Kiddie Schedule for Affective Disorders and Schizophrenia (K-SADS), also completed by parents. For each mental health factor from the K-SADS, responses to two yes/no questions about past and present diagnoses or symptoms were combined into a single binary variable based on any positive responses.

2.4 Environmental factors

Multiple aspects of the youth social environment were investigated. Parental mental health was based on the Adult Behavior Checklist (ABCL), and was measured as a t-score of critical items related to depression, self-harm, substance use, mood swings, and suicidal thoughts. Family dynamics were assessed based on the parent-reported Family Environment Scale. Conflict and cohesion were measured on continuous scales by aggregating responses from multiple survey items designed to assess specific

aspects of the family's organizational structure. In addition, parent responses to one

question from the Mexican American Cultural Values Scale were extracted, providing a

measure for how strongly they believed that family members should show love and

affection to one another. Responses were coded on a Likert scale from not at all (1) to

completely (5). Prior work has identified associations between this parental belief youth

prosocial behaviors (Smith & Stamoulis, 2023).

Questions on peer relations were: 1) number of close friends (from the Other Resilience

survey); 2) regular group of friends (binary variable, from the KSADS); 3) bullying,

represented as a binary variable based on questions on whether a child had been

bullied in any social setting (from a parent-reported K-SADS question) or had been

cyberbullied (from the youth-reported Cyber-bullying instrument). The Discrimination

Measure, reported by youth, assessed experiences of perceived discrimination in four

domains: race/ethnicity/color, being from another country, sexual orientation, and/or

weight. Given small samples with positive responses in each domain, responses were

combined into a single binary variable for analysis, indicating any type of discrimination.

Youth responses on whether they got along with their teacher and liked school (from the

School Risk & Protective Factors survey) were also examined. Responses were coded

on a Likert scale from definitely not true (1) to definitely true (4).

2.5 Additional characteristics

Other assessments of youth social behaviors were extracted from The Early Adolescent

Temperament Questionnaire (completed by the parents) and included *"Is energized by*

being in large crowds of people," "Wants to have close relationships with other people,"

"Is shy," and *"Feels shy about meeting new people"*, all on a 5-point scale (almost

always untrue (1) to almost always true (5)). Another question from the same instrument

asked if the child *"Likes meeting new people"*, and was reverse-coded by the ABCD,

with higher values indicating less frequent social engagement. Two questions on past or

present symptoms of a fear of social situations were extracted from the KSAD-S and

combined into a binary variable with 1 corresponding to a 'true' response to either or

both questions, and 0 otherwise. Another question from the CBCL asked parents if their

child was *"self-conscious or easily embarrassed"* and was coded on a Likert scale from

not true (0) to very/often true (2).

2.6 Neuroimaging data analysis

2.6.1. Resting-state fMRI processing and topological property estimation

Individual structural MRI and resting-state fMRI (rs-fMRI; participants completed up to

four 5-minute runs), underwent initial preprocessing by the ABCD study's Data Analysis,

Informatics & Resource Center (DAIRC), to correct for bias field, B0 distortion, grad

warp and motion in the scanner (Hagler et al., 2019)). Additional processing was

necessary to further suppress motion-related and other artifacts, harmonize the data

across scanners (data were collected with a 3T GE, Siemens or Philips scanner), and

downsample the voxel-level time series. The custom Next Generation Neural Data

Analysis (NGNDA) pipeline was used for this purpose (Next-Generation Neural Data

Analysis (NGNDA), 2020). Following segmentation of each participant's structural MRI

(sMRI), coregistration of their fMRI to the sMRI, and normalization to common MNI152

space, fMRI signals were further processed to improve their quality. This process

included removal of initial frames, correction for motion, and denoising using signal

decomposition to eliminate signal components likely associated with cardiorespiratory

and other artifacts (Brooks et al., 2021). Voxel-level time series were then downsampled

using three atlases for parcellating cortical and subcortical structures and the

cerebellum, resulting in 1088 parcel signals (Diedrichsen et al., 2009; Schaefer et al.,

2018; Tian et al., 2020). Only fMRI runs with ≤10% motion-censored frames were

retained for further analysis. Each participant's best-quality run (based on lowest

median connectivity, given that the brain at rest is overall weakly coordinated at rest)

was first analyzed. Often this run also had the lowest number of frames censored for

motion. For replication purposes, the second best-quality run was selected from a

subsample of participants (n = 2160 (~77%) of the sample) who had more than one run

of adequate quality for analysis. Median (IQR) percent of frames censored for motion

was 1.07% (IQR = 4.00%) in the best-quality run and 1.07% (IQR = 3.73%) in the

second run. Associations between resting-state topological properties and measures of

social withdrawal that were consistent across runs are reported in primary analyses and

tables. Additional associations based on the best-quality run, and also larger sample,

are reported in secondary analyses and supplemental tables (see Supplementary

Materials).

Resting-state connectivity was calculated as the peak cross-correlation between pairs of

fMRI signals, resulting in symmetric 1088-by-1088 matrices. All methods used to

estimate connectivity yield full matrices that include spurious correlations that need to

be eliminated prior to the estimation of topological properties. In this study, statistical

thresholds were estimated from the entire cohort and all available runs, and

bootstrapping with replacement was used to derive thresholds for different levels of

connection strength. A conservative threshold, equal to the moderate outlying value

(median + 1.5*IQR) of peak cross-correlation was then used to obtain adjacency

matrices. Topological properties were estimated from these matrices at multiple spatial

scales: a) brain regions (network nodes), b) large-scale networks that included the

reward and social networks (Yeo et al, 2011; Haber & Knudson, 2010, Blakemore,

2008), c) entire brain (the connectome). Topological properties at the network and

connectome levels included modularity, global clustering, global efficiency, natural

connectivity (robustness), median connectivity (within and between networks), fragility,

stability, and segregation. The latter was calculated as the ratio of within-network over

out-of-network median connectivity. Node properties included degree, local clustering,

and eigenvector centrality. Detailed descriptions of each of these properties are

provided elsewhere (Pasqualetti et al., 2020; Rubinov & Sporns, 2010; Wu et al., 2011).

2.6.2. Structural parameters

Morphometric brain properties (provided by the ABCD) were analyzed. These were

estimated using a parcellation based on the Desikan-Killiany atlas, which segments the

cortex into 68 regions based on anatomical landmarks (Desikan et al., 2006).

Subcortical voxels were parcellated into 30 regions using a probabilistic method (Fischl

et al., 2002). Properties of 98 regions were analyzed, including cortical thickness and

white matter intensity, and cortical and subcortical volume (Hagler et al., 2019). White

matter intensity was was also examined as a potentially sensitive cortical marker of

brain development (Hagler et al., 2019; Lewis et al., 2018; Westlye et al., 2009).

2.7 Statistical analysis

All relationships of interest were investigated using linear mixed-effects regression

models that included a random intercept and slope for each of the 21 ABCD sites to

account for potential site effects. All models included age, pubertal stage, sex, race-

ethnicity, family income, and BMI z-score stratified by sex (which has been previously

linked to differences in topological properties in the ABCD cohort (Brooks et al., 2023).

All analyses accounted for potential bias associated with sampling differences at the 21

sites, using the ABCD-provided propensity scores. Given limited statistical power to

compare individual race/ethnic groups, race-ethnicity was coded as a binary variable

representing white non-Hispanic participants (0) and racioethnic minorities (1). Pubertal

stage was coded as an ordinal variable in the range 1 (pre-puberty) to 4 (late/post-

puberty).

In models that included brain parameters (the dependent variables), social withdrawal

and preference for solitude were the independent variables of interest. These models

were also adjusted for internalizing behavior scores, a variable that was consistently

significant across analyses. In models that examined association between social

isolation and environmental, mental health, and other behavioral factors, social

withdrawal and preference for solitude were the dependent variables. Finally, models

with resting-state network parameters included percent of frames censored for motion

and scan time of day participants (0-23 h) as additional variables (Brooks et al., 2021;

Hu et al., 2023; Vaisvilaite et al., 2022). Variations of models were also developed, in

order to assess pairwise comparisons between specific response categories of the

social withdrawal measures, focusing on two categories at a time, or treating these

measures as binary variables.

The significance level was set at α = 0.05. All parameter p-values were adjusted for the

False Discovery Rate (FDR), using a well-established method (Benjamini & Hochberg,

1995). In analyses focusing on topological properties of the entire brain or individual

networks, the FDR correction was done over properties. In analyses focusing on

regional (node) properties, the correction was done over properties of all nodes within a

particular network. P-values for parameters in structural models were adjusted over

regional morphometric properties. Associations were considered significant if the model

intercept, parameter of interest, and model p-value were all significant. Model cross-

validation was performed by randomly dividing the data into training and testing sets

(75:25) 100 times. The coefficient of variation of the root-mean-squared error

(CV[RMSE]) between predicted and observed responses was used to assess each

model's predictive power. All data were analyzed in the Harvard Medical School high-

performance cluster using the software MATLAB (release R2023a, Mathworks, Inc).

3. RESULTS

Across the cohort, 455 (17.9%) youth sometimes or often preferred solitude, and 170

(6.0%) were sometimes or often socially withdrawn. Less than 10% reported past or

present anhedonia (264 (9.4%)) and 40 (1.4%) were diagnosed with social anxiety

disorder. Median (IQR) t-scores for other mental health measures were 51.0 (2.0) for

anxiety, 50.0 (4.0) for depression, 41.0 (16.0) for externalizing behaviors, and 46.0

(14.0) for internalizing behaviors.

Median (IQR) parent mental health score was 51.0 (4.0). Almost 90% of parents said

that it was important for the family to show love and affection towards each other.

Median family cohesion was 8.0 (2.0) whereas median family conflict was 2.0 (2.0).

Median (IQR) family size was 4 (1). Participants had on average 5 close friends, and

almost 90% had a regular group of kids to hang out with. About 10% reported being

discriminated against (302 (10.7%)), and 519 (18.5%) had been bullied. Almost 95%

reported having good relationships with teachers, and ~70% said that they liked school

a lot (1991 (70.9%)). Summary statistics for social environmental factors are provided in

Table 2.

Almost 70% of parents reported that their child wanted close relationships with other

people (1938 (69.0%)), ~60% of youth were not self-conscious or easily embarrassed

(1717 (61.%), and liked to meet new people (1654 (58.9%)), and almost 30% were

energized by large crowds (813 (28.9%)). About a quarter were shy (552 (26.4%)),

fewer were specifically shy about meeting new people (466 (16.6%)), and even fewer

were afraid of social situations (150 (5.3%)). Detailed distributions of responses are

provided in Table S1.

3.1 Associations between social isolation, behaviors and mental health outcomes

Older participants and at more advanced pubertal stages were more frequently socially isolated (β = 0.04 to 0.07, CI = [0.002, 0.13], p < 0.05). Preferring solitude was associated with higher internalizing behaviors (β = 0.45, CI = [0.42, 0.49], p < 0.01), externalizing behaviors (β = 0.23, CI = [0.20, 0.27], p < 0.01), depression (β = 0.39, CI = [0.35, 0.42], p < 0.01), and anxiety (β = 0.29, CI = [0.26, 0.33], p < 0.01). Similarly, being socially withdrawn was associated with higher internalizing behavior (β = 0.41, CI = [0.37, 0.45], p < 0.01), externalizing behaviors (β = 0.25, CI = [0.21, 0.29], p < 0.01), depression (β = 0.45, CI = [0.42, 0.49], p < 0.01), and anxiety (β = 0.35, CI = [0.31, 0.39], p < 0.01).

Parental mental health issues were positively associated with youth preferring solitude and social withdrawal (β = 0.14, CI = [0.10, 0.19], p < 0.01, and β = 0.13, CI = [0.09, 0.17], p< 0.01), respectively). In contrast, being energized by large crowds of people was negatively associated with preferring solitude (β = -0.23, CI = [-0.27, -0.19], p < 0.01). When participants who often preferred solitude were excluded (due to their small sample size (n = 48)), and only participants who sometimes preferred solitude were compared to those who did not, additional positive associations were identified with being shy (β = 0.45, CI = [0.34, 0.55], p < 0.01), and not liking to meet new people (β = 0.58, CI = [0.46, 0.70], p < 0.01). Social withdrawal was also associated with being shy (β = 0.21, CI = [0.16, 0.25], p < 0.01), not liking to meet new people (β = 0.21, CI = [0.17, 0.25], p < 0.01), and feeling shy about meeting new people (β = 0.19, CI = [0.15, 0.23], p < 0.01).

3.2 Associations between social isolation and resting-state network properties

3.2.1 Whole-brain topology

In the full cohort, there were no consistent associations across both fMRI runs.

However, within-group comparisons of those who preferred solitude identified negative

correlations between preference for solitude and connectome efficiency and global

clustering (β = -0.03 to -0.02, CI = [-0.05, -0.01], p ≤ 0.04). Additional associations were

estimated based on only the larger sample with the best-quality fMRI run. In the full

cohort, preferring solitude was negatively associated with global efficiency (β = -0.07, CI

= [-0.11, -0.03], p = 0.01), and similarly for pairwise group comparisons. Also, negative

associations between preferring solitude and topological robustness, and similarly for

stability, were estimated in group comparisons, and positive associations with

modularity and segregation. There were no significant associations between brain-wide

topological properties and being withdrawn. Statistics of models based only on the best

run are summarized in Table S2.

3.2.2 Network topology

Preferring solitude was associated with higher segregation (more connected local

communities) of the right dorsal attention, temporoparietal, and social networks (β =

0.07 to 0.10, CI = [0.02, 0.14], p < 0.04). Within-group comparisons identified

associations between preference for solitude and higher segregation of the right social

network (β = 0.01, CI = [0.003, 0.01], p = 0.04) but lower global efficiency of the left

cerebellum (β = -0.09, CI = [-0.13, -0.05], p < 0.01). In addition, social withdrawal was

associated with lower topological robustness and stability (β = -0.78 to -0.67, CI = [-

1.34, -0.19], p < 0.03) and higher fragility (β = 0.04, CI = [0.01, 0.07], p < 0.05) of the left

cerebellum. Model statistics are summarized in Table 3.

Additional associations were identified in analyses of only the best-quality fMRI.

Preferring solitude was associated with higher segregation, modularity and/or fragility of

multiple networks, including the basal ganglia, dorsal attention, prerontal, reward,

salience, somatomotor, social and temporoparietal networks (β = 0.06 - 0.10, CI = [0.01,

0.14], p < 0.05). It was also associated with lower connectivity, topological robustness,

stability and efficiency of most of the same networks, as well as the frontoparietal

control network (β = -0.06 to -0.05, CI = [-0.10, -0.01], p < 0.05). Being withdrawn was

also associated with higher segregation of the right temporoparietal network (β = -0.08,

CI = [-0.03, -0.12], p < 0.01). Across networks, models on associations between

preference for solitude and topological properties (including within-network median

connectivity, fragility and global clustering) had good predictive power (CV[RMSE] ≤

0.20). Similar results were obtained across pairwise group comparisons of those who

strongly preferred solitude vs those who did not, within-group comparisons of those who

preferred it, and those who somewhat preferred solitude vs those who did not, and

similarly for social withdrawal. Detailed results and model statistics are provided in

Table S3.

Finally, there were no consistent associations between social isolation and node

properties across both fMRI runs. For the best-quality run, preferring solitude was

associated with lower connectedness (degree) in nodes of the left dorsal attention

network (β = -0.08 to -0.06, CI = [-0.12, -0.02], p <= 0.04). There were no other

significant associations.

3.3 Associations between social withdrawal and structural properties

Preferring solitude was associated with lower thickness of the left superior temporal,

right caudal anterior cingulate, and right insular gyri and the left cuneus and pars

opercularis (β = -0.09 to -0.05, CI = [-0.14, -0.01], p < 0.05), and lower white matter

intensity but higher volume of the right parahippocampal gyrus (β = -0.02, CI = [-0.04, -

0.002], p < 0.05, and β = 0.06, CI = [0.01, 0.10], p < 0.05 respectively). Social

withdrawal was associated with higher white matter intensity in the right isthmus of the

cingulate gyrus (β = 0.03, CI = [0.01, 0.05], p = 0.04). These associations were based

on models with good predictive power (CV[RMSE] ≤0.20).

Comparisons of those who strongly preferred solitude to those who did not, identified

associations between preference for solitude and lower thickness of the left cuneus, and

left superior temporal, right caudal anterior cingulate and right insular gyri (β = -0.04 to -

0.02, CI = [-0.07, -0.003], p < 0.05), lower volume of the left cuneus and right fusiform

gyrus (β = -264.31 to -109.54, CI = [-474.51, -11.40], p = 0.04), and lower white matter

intensity of the right parahippocampal gyrus (β = -0.23, CI = [-0.42, -0.05], p = 0.04).

Social withdrawal was associated with higher volume of the right pallidum (β = 81.06, CI

= [12.50, 149.62], p = 0.02), lower thickness of the right cuneus and right insular and

lingual gyri (β = -0.06, CI = [-0.09, -0.01], p $\leq$ 0.04), and lower volume of the right

cuneus (β = -333.56, CI = [-529.69, -137.44], p < 0.01) and right lingual gyrus (β = -

492.48, CI = [-848.47, -136.49], p = 0.02).

In within group comparisons, preference for solitude was associated with lower white

matter intensity in the right inferior temporal gyrus (β = -0.39, CI = [-0.68, -0.10], p =

0.03), and social withdrawal was associated with higher volume of left caudate nucleus,

right pallidum and the putamen (β = 231.75 - 369.01, CI = [2.68, 668.10], p < 0.05),

lower thickness of the right cuneus (β = -0.15, CI = [-0.23, -0.06], p < 0.01) and right

lingual gyrus (β = -0.09, CI = [-0.17, -0.01], p = 0.04), and lower volume of the right

cuneus (β = -723.26, CI = [-1185.41, -261.12], p < 0.01).

Comparisons of those who somewhat preferred solitude versus those who did not,

identified associations between preference for solitude and higher volume of the left

amygdala, and right entorhinal and parahippocampal gyri (β = 25.72 - 61.62, CI = [0.13,

106.57], p < 0.05), and lower thickness of the right pars opercularis and right caudal

anterior cinguate gyrus (β = -0.03, CI = [-0.06, -0.003], p < 0.05). Similar comparisons of

those who were somewhat socially withdrawn to those who were not, identified a

negative association between social withdrawal and volume in of the right caudal middle

frontal gyrus (β = -319.88, CI = [-579.79, -59.97], p < 0.05). Model statistics are

summarized in Table 4. Negative associations between preference for solitude and

cortical thickness, which were more consistent across group comparisons (especially in

the caudal anterior cingulate cortex) are shown in Figure 1.

4. DISCUSSION

In a large cohort of almost 3,000 adolescents, we have systematically investigated the

neural correlates of preference for solitude and social withdrawal, especially their

impacts on hallmark structural and topological characteristics of large scale networks

that play critical roles in cognitive (including social) function. Social isolation in

adolescence, a sensitive period of brain and social development but also vulnerable

period for mental health, may lead to brain circuit miswiring, and increased risk of

cognitive delays and deficits, and maladaptive behaviors across the lifespan. In this

study's cohort, about 18% of youth sometimes or often preferred solitude, and 6% were

socially withdrawn. Older youth whose parents had mental health issues were more

frequently preferred solitude and/or were socially withdrawn. These findings are in

agreement with prior studies that have shown that parental behaviors and their mental

health issues are significant risk factors for youth loneliness, social withdrawal and

isolation (Matthews et al, 2015; Almeida et al, 2021), as early as infancy (Mantymaa et

al, 2008).

As their social world expands beyond the family, and changes as a function of age,

adolescents may become increasingly exposed to additional social stressors, such as

peer rejection and victimization, which may, in turn, lead to social withdrawal and

isolation. In this cohort, almost 90% had a regular group of friends, only ~10% had

experienced discrimination (or any form), and ~20% had been bullied. Neither overall

discrimination nor bullying were statistically associated with an increased likelihood of

social isolation. However, unmeasured (by the ABCD) granular aspects of peer

relationships and their quality and other latent factors in the youth environment that

change with age could contribute to social isolation. Preference for solitude and social

withdrawal was also associated with youth depression, anxiety, and internalizing and

externalizing behaviors, in agreement with prior work that has identified extensive links

between social isolation and mental health issues in youth (Rubin et al, 2009; Rubin &

Lollis, 2015; Orben et al, 2020; Christiansen et al, 2021).

Preferring solitude and social withdrawal had overlapping structural but distinct

topological brain correlates. Both preference for solitude and social withdrawal were

associated with lower thickness of the cuneus and insular gyrus, while preference for

solitude was also associated with lower thickness of the superior temporal and caudal

anterior cingulate gyri and the pars opercularis, and social withdrawal with lower

thickness of the lingual gyrus. The cuneus is a structural brain hub (Oldham and

Fornito, 2019), i.e., a brain region where multimodal information is integrated, and has

been implicated both in emotional processing and social function (Riedel et al 2018;

Eslinger et al, 2021), but also loneliness (Lam et al, 2021; Wong et al, 2022). The

insular cortex is another structural hub involved in information integration and supports

interoception, but is also involved in (social) emotional processing and maladaptive

social behaviors (Lamm & Singer, 2010; Terasawa et al, 2013; Emmerling et al,

2016;Gogolla, 2017). The superior temporal gyrus is considered a 'hub' of the social

brain and is extensively involved in social perception and processing (Blakemore, 2008;

Pelphrey & Carter, 2008; Lahnakoski et et al, 2012; Beauchamp, 2015). Decreased

thickness of the cuneus, and insular and superiot temporal gyri has been associated

with cognitive deficits but also mental health issues, not only in adults but also

adolescents (von Tol et al, 2014; Pan, 2015; Sheffield et al, 2021). The anterior

cingulate cortex has distributed connections with frontal and limbic regions and plays a

ubiquitous role in cognitive function, including decision-making, cognitive control, and

emotion and reward processing and regulation, and is also involved in social cognition

(Devinsky et al, 1995; Vogt, 2005; Apps et al, 2016). The caudal part of the anterior

cingulate cortex is involved in cognitive control, and decreased thickness of this region

has been associated with depression, and developmental disorders affecting social

function, such as Autism Spectrum Disorders (ASD) (Laidi et al, 2019; Mertse et al,

2022). Finally, the pars opercularis and lingual gyrus are both involved in speech and

language processing (Palejwala et al, 2021; Zaccarella et al, 2015), and morphological

anomalies in these regions have been associated both with linguistic deficits but more

broadly impaired social communication, and mental health issues, including depression

and internalizing symptoms (Jensen et al, 2015; Rosada et al, 2023).

Preference for solitude and social withdrawal were also associated with other structural

differences in some of the same areas, including lower volume of the cuneus and the

lingual gurys, but also higher volume of the entorhinal and parahippocampal gyri,

amygdala and basal ganglia structures, including the caudate, pallidum and putamen.

These findings are in agreement with those of prior studies that have associated social

isolation and loneliness with higher amygdala volume (Xiong et al, 2023), potentially as

a result of higher emotional sensitivity, but also distress in social settings (Lam et al,

2021; Vitale et al, 2022) and mental health issues (Espinoza et al, 2020; Suor et al,

2024). Specifically in children and adolescents, it has been associated with anxiety,

negative affect and behavioral issues (Merz et al, 2018; Liu et al, 2024; Pereira et al,

2024). In addition, higher putamen volume has been associated with disorders affecting

social function, such as ASD (Sato et al, 2014), and higher caudate volume with

behavioral disregulation, impulsivity, and risk for developmental disorders such as ASD

(Voelbel et al, 2006). Together these findings suggest that social isolation is associated

with extensive morphological alterations in the adolescent brain in regions supporting

social function but that have also been implicated in mental health disorders and social

dysfunction, and developmental disorders impacting social communication. In addition,

cortical thinning and increased structural volume are hallmark characteristics of brain

development. It is possible that social isolation may be associated with accelerated

neural maturation in selective regions, which, in turn, increases the risk for mental

health and behavioral problems. A recent study on adolescents during the COVID-19

pandemic reported accelerated brain maturation partly as a result of social isolation due

to lockdowns and social distancing (Corrigan et al, 2024).

Preference for solitude was also associated with higher topological segregation

(modularity) of the dorsal attention and temporoparietal networks, and an overlapping

distributed network representing the social brain (Blakemore, 2008). Increased

structural and functional network modularity is also a fundamental characteristic of brain

and cognitive development (Baum et al, 2017; Tooley et al, 2022), however in socially

isolated youth it may reflect aberrantly accelerated neural maturation. Higher modularity

has also been associated with mental health issues (Gao et al, 2023). Furthermore,

social withdrawal was associated with topological changes in the cerebellum, including

lower robustness and efficiency and higher fragility. A number of studies have shown

that the cerebellum, which has extensive cortical and subcortical connections with brain

regions supporting social function, plays an important role in social development,

perception and prediction, and emotional processing) (Van Overwalle, 2020, 2024;

Turrini & Avenanti, 2024). Abnormal structural changes in the cerebellum have been

associated with developmental and mental health disorders, including those directing

affecting social function (Fett et al, 2014; Jack & Morris, 2014; Phillips et al, 2015; Zhu &

Qiu, 2022; Olivito et al, 2023; Wang et al, 2023). Together, these findings suggest that

social isolation during adolescence may have profound detrimental impacts on large-

scale networks and their constituent structures that play ubiquitous roles in cognitive,

and specifically social function, and are abnormally modulated by mental health

disorders.

Despite its strengths, including the large sample that captures the heterogeneity of

adolescent brain development, comprehensive assessment of topological properties of

resting-state networks, concurrent investigation of morphological and functional brain

characteristics, and examination of environmental correlates of youth social isolation,

this study also had some limitations. First, assessments of social withdrawal and

preference for solitude were based on parent reports. When possible, youth surveys

were analyzed, but across surveys measuring the youth environment and mental

health/behavioral assessments, parent reports were in general more complete and

reliable, or the only available reports (corresponding youth ones were not available).

Second, as is the case with any retrospective investigation, analyses were limited by the

experimental decisions made by the ABCD investigators. Thus, more granular

assessments of the youth social environment and related behaviors were not available.

Nevertheless, the ABCD is the only investigation that extensively samples the youth

social world, and relationships with parents, family, teachers, and peers. The present

study leveraged data on these relationships in order to examine their associations with

social isolation.

This study makes a significant scientific contribution and provides novel insights into the

detrimental effects of social isolation on the developing adolescent brain and its

maturating, and vulnerable to miswiring, neural circuitry. It has identified widespread

morphological and topological brain alterations in youth who prefer to be alone and/or

are socially withdrawn, several of which have not been previously related directly to

social isolation, but to disorders affecting social communication. It has also associated

preference for solitude and social withdrawal with parental mental health but also

common youth mental health issues. Adolescence is a sensitive period for mental

health. Many identified brain alterations were in regions that support social function but

have also been implicated in mental health disorders. In addition, lower cortical

thickness and higher subcortical volume and increased modularity of several large-scale

networks also suggest potential accelerated maturation of the socially isolated brain,

which has also been associated with mental health issues. The ABCD follows youth

longitudinally, thus future investigations could specifically examine the hypothesis of

accelerated neural maturation and track the emergence of mental health issues as a

function of age in socially withdrawn youth.

Funding

This study was supported by the National Science Foundation, Awards #2116707,

2207733, 1940096

Data Availability Statement

The data underlying this article are available in National Institute of Mental Health Data

Archive (NDA). https://nda.nih.gov/.

Declaration of Interest Statement

The authors have no interests to declare.

5. REFERENCES

Albert, D., Chein, J., & Steinberg, L. (2013). The Teenage Brain: Peer Influences on

Adolescent Decision Making. *Current Directions in Psychological Science*, *22*(2),

114–120. https://doi.org/10.1177/0963721412471347

Almeida, I. L. de L., Rego, J. F., Teixeira, A. C. G., & Moreira, M. R. (2021). Social

isolation and its impact on child and adolescent development: A systematic

review. *Revista Paulista de Pediatria*, *40*, e2020385.

https://doi.org/10.1590/1984-0462/2022/40/2020385

Andrews, J. L., Ahmed, S. P., & Blakemore, S.-J. (2021). Navigating the Social

Environment in Adolescence: The Role of Social Brain Development. *Biological*

Psychiatry, *89*(2), 109–118. https://doi.org/10.1016/j.biopsych.2020.09.012

Apps MA, Rushworth MF, Chang SW. The Anterior Cingulate Gyrus and Social

Cognition: Tracking the Motivation of Others. Neuron. 2016 May 18;90(4):692-

707.

Arslan, G., Yıldırım, M., Tanhan, A., & Kılınç, M. (2023). Social Inclusion to Promote

Mental Health and Well-Being of Youths in Schools. In C. Boyle & K.-A. Allen

(Eds.), *Research for Inclusive Quality Education: Leveraging Belonging,*

Inclusion, and Equity (pp. 113–122). Springer Nature.

https://doi.org/10.1007/978-981-16-5908-9_9

Arslan, Ph. D., Gökmen. (2021). School belongingness, well-being, and mental health

among adolescents: Exploring the role of loneliness. *Australian Journal of*

669   *Psychology*, *73*(1), 70–80. https://doi.org/10.1080/00049530.2021.1904499

670 Asher, S. R., & Paquette, J. A. (2003). Loneliness and Peer Relations in Childhood.

671   *Current Directions in Psychological Science*, *12*(3), 75–78.

672   https://doi.org/10.1111/1467-8721.01233

673 Ashtari, M., Cervellione, K. L., Hasan, K. M., Wu, J., McIlree, C., Kester, H., Ardekani,

674   B. A., Roofeh, D., Szeszko, P. R., & Kumra, S. (2007). White matter development

675   during late adolescence in healthy males: A cross-sectional diffusion tensor

676   imaging study. *NeuroImage*, *35*(2), 501–510.

677   https://doi.org/10.1016/j.neuroimage.2006.10.047

678 Barzeva, S. A., Meeus, W. H. J., & Oldehinkel, A. J. (2019). Social Withdrawal in

679   Adolescence and Early Adulthood: Measurement Issues, Normative

680   Development, and Distinct Trajectories. *Journal of Abnormal Child Psychology*,

681   *47*(5), 865–879. https://doi.org/10.1007/s10802-018-0497-4

682 Barzeva, S. A., Richards, J. S., Veenstra, R., Meeus, W. H. J., & Oldehinkel, A. J.

683   (2022). Quality over quantity: A transactional model of social withdrawal and

684   friendship development in late adolescence. *Social Development*, *31*(1), 126–

685   146. https://doi.org/10.1111/sode.12530

686 Baum GL, Ciric R, Roalf DR, Betzel RF, Moore TM, Shinohara RT, Kahn AE, Vandekar

687   SN, Rupert PE, Quarmley M, Cook PA, Elliott MA, Ruparel K, Gur RE, Gur RC,

688   Bassett DS, Satterthwaite TD. Modular Segregation of Structural Brain Networks

689   Supports the Development of Executive Function in Youth. Curr Biol. 2017 Jun

690   5;27(11):1561-1572.e8.

691 Beauchamp MS. The social mysteries of the superior temporal sulcus. Trends Cogn

Sci. 2015 Sep;19(9):489-90.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical

and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical*

Society: Series B (Methodological), *57*(1), 289–300.

https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Best, O., & Ban, S. (2021). Adolescence: Physical changes and neurological

development. *British Journal of Nursing*, *30*(5), 272–275.

https://doi.org/10.12968/bjon.2021.30.5.272

Biggs, B. K., Vernberg, E. M., & Wu, Y. P. (2012). Social Anxiety and Adolescents'

Friendships: The Role of Social Withdrawal. *The Journal of Early Adolescence*,

32(6), 802–823. https://doi.org/10.1177/0272431611426145

Blakemore, S.-J. (2008). The social brain in adolescence. *Nature Reviews*

Neuroscience, *9*(4), 267–277. https://doi.org/10.1038/nrn2353

Blakemore, S.-J. (2012). Development of the social brain in adolescence. *Journal of the*

Royal Society of Medicine, *105*(3), 111–116.

https://doi.org/10.1258/jrsm.2011.110221

Blakemore, S.-J. (2019). Adolescence and mental health. *The Lancet*, *393*(10185),

2030–2031. https://doi.org/10.1016/S0140-6736(19)31013-X

Blakemore, S.-J., & Mills, K. L. (2014a). Is Adolescence a Sensitive Period for

Sociocultural Processing? *Annual Review of Psychology*, *65*(Volume 65, 2014),

187–207. https://doi.org/10.1146/annurev-psych-010213-115202

Blakemore, S.-J., & Mills, K. L. (2014b). Is Adolescence a Sensitive Period for

Sociocultural Processing? *Annual Review of Psychology*, *65*(Volume 65, 2014),

187–207. https://doi.org/10.1146/annurev-psych-010213-115202

Bor, W., Dean, A. J., Najman, J., & Hayatbakhsh, R. (2014). Are child and adolescent
   mental health problems increasing in the 21st century? A systematic review.
   Australian & New Zealand Journal of Psychiatry, *48*(7), 606–616.
   https://doi.org/10.1177/0004867414533834

Borg, M. E., & Willoughby, T. (2022). Affinity for Solitude and Motivations for Spending
   Time Alone Among Early and Mid- Adolescents. *Journal of Youth and
   Adolescence*, *51*(1), 156–168. https://doi.org/10.1007/s10964-021-01520-1

Bosch, A., Riebschleger, J., & van Loon, L. (2017). Dutch youth of parents with a mental
   illness reflect upon their feelings of guilt and shame. *International Journal of
   Mental Health Promotion*, *19*(3), 159–172.
   https://doi.org/10.1080/14623730.2017.1315955

Branje, S., de Moor, E. L., Spitzer, J., & Becht, A. I. (2021). Dynamics of Identity
   Development in Adolescence: A Decade in Review. *Journal of Research on
   Adolescence*, *31*(4), 908–927. https://doi.org/10.1111/jora.12678

Brilliant T., D., Takeuchi, H., Nouchi, R., Yokoyama, R., Kotozaki, Y., Nakagawa, S.,
   Hanawa, S., Sekiguchi, A., Ikeda, S., Sakaki, K., Kawata, K. H. dos S., Nozawa,
   T., Yokota, S., Magistro, D., & Kawashima, R. (2022). Loneliness inside of the
   brain: Evidence from a large dataset of resting-state fMRI in young adult.
   Scientific Reports, *12*(1), 7856. https://doi.org/10.1038/s41598-022-11724-5

Brooks, S. J., Katz, E. S., & Stamoulis, C. (2022). Shorter Duration and Lower Quality
   Sleep Have Widespread Detrimental Effects on Developing Functional Brain
   Networks in Early Adolescence. *Cerebral Cortex Communications*, *3*(1), tgab062.

https://doi.org/10.1093/texcom/tgab062

Brooks, S. J., Parks, S. M., & Stamoulis, C. (2021). Widespread positive direct and

indirect effects of regular physical activity on the developing functional

connectome in early adolescence. *Cerebral Cortex*, *31*(10), 4840–4852.

https://doi.org/10.1093/cercor/bhab126

Brooks, S. J., Smith, C., & Stamoulis, C. (2023). Excess BMI in early adolescence

adversely impacts maturating functional circuits supporting high-level cognition

and their structural correlates. *International Journal of Obesity (2005)*, *47*(7),

590–605. https://doi.org/10.1038/s41366-023-01303-7

Cacioppo, J. T., Hughes, M. E., Waite, L. J., Hawkley, L. C., & Thisted, R. A. (2006).

Loneliness as a specific risk factor for depressive symptoms: Cross-sectional and

longitudinal analyses. *Psychology and Aging*, *21*(1), 140–151.

https://doi.org/10.1037/0882-7974.21.1.140

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M.,

Soules, M. E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C. A., Wager, T.

D., Banich, M. T., Speer, N. K., Sutherland, M. T., Riedel, M. C., Dick, A. S.,

Bjork, J. M., Thomas, K. M., … ABCD Imaging Acquisition Workgroup. (2018).

The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition

across 21 sites. *Developmental Cognitive Neuroscience*, *32*, 43–54.

https://doi.org/10.1016/j.dcn.2018.03.001

Caspi, A., Harrington, H., Moffitt, T. E., Milne, B. J., & Poulton, R. (2006). Socially

isolated children 20 years later: Risk of cardiovascular disease. *Archives of*

Pediatrics & Adolescent Medicine, *160*(8), 805–811.

https://doi.org/10.1001/archpedi.160.8.805

Cassidy, J., & Asher, S. R. (1992). Loneliness and Peer Relations in Young Children.

Child Development, *63*(2), 350–365. https://doi.org/10.1111/j.1467-

8624.1992.tb01632.x

Cheng, H., & Furnham, A. (2002). Personality, peer relations, and self-confidence as

predictors of happiness and loneliness. *Journal of Adolescence*, *25*(3), 327–339.

https://doi.org/10.1006/jado.2002.0475

Christiansen, J., Qualter, P., Friis, K., Pedersen, S., Lund, R., Andersen, C., Bekker-

Jeppesen, M., & Lasgaard, M. (2021). Associations of loneliness and social

isolation with physical and mental health among adolescents and young adults.

Perspectives in Public Health, *141*(4), 226–236.

https://doi.org/10.1177/17579139211016077

Cogan, N., Riddell, S., & Mayes, G. (2005). Children living with an affectively ill parent:

How do they cope? *Educational and Child Psychology*, *22*(4), 16–28.

Coplan, R. J., Ooi, L. L., Xiao, B., & Rose-Krasnor, L. (2018). Assessment and

implications of social withdrawal in early childhood: A first look at social

avoidance. *Social Development*, *27*(1), 125–139.

https://doi.org/10.1111/sode.12258

Corrigan NM, Rokem A, Kuhl PK. COVID-19 lockdown effects on adolescent brain

structure suggest accelerated maturation that is more pronounced in females

than in males. Proc Natl Acad Sci U S A. 2024 Sep 17;121(38):e2403200121.

Costello, E. J., Copeland, W., & Angold, A. (2011). Trends in psychopathology across

the adolescent years: What changes when children become adolescents, and

when adolescents become adults? *Journal of Child Psychology and Psychiatry*,

52(10), 1015–1025. https://doi.org/10.1111/j.1469-7610.2011.02446.x

Dam, K., Joensen, D. G., & Hall, E. O. C. (2018). Experiences of adults who as children

lived with a parent experiencing mental illness in a small-scale society: A

Qualitative study. *Journal of Psychiatric and Mental Health Nursing*, *25*(2), 78–

87. https://doi.org/10.1111/jpm.12446

Danese, A., Moffitt, T. E., Harrington, H., Milne, B. J., Polanczyk, G., Pariante, C. M.,

Poulton, R., & Caspi, A. (2009). Adverse childhood experiences and adult risk

factors for age-related disease: Depression, inflammation, and clustering of

metabolic risk markers. *Archives of Pediatrics & Adolescent Medicine*, *163*(12),

1135–1143. https://doi.org/10.1001/archpediatrics.2009.214

de Girolamo, G., Dagani, J., Purcell, R., Cocchi, A., & McGorry, P. D. (2012). Age of

onset of mental disorders and use of mental health services: Needs,

opportunities and obstacles. *Epidemiology and Psychiatric Sciences*, *21*(1), 47–

57. https://doi.org/10.1017/s2045796011000746

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D.,

Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., &

Killiany, R. J. (2006). An automated labeling system for subdividing the human

cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*,

31(3), 968–980. https://doi.org/10.1016/j.neuroimage.2006.01.021

Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to

behaviour. Brain. 1995 Feb;118 (Pt 1):279-306.

Diedrichsen, J., Balsters, J. H., Flavell, J., Cussans, E., & Ramnani, N. (2009). A

probabilistic MR atlas of the human cerebellum. *NeuroImage*, *46*(1), 39–46.

https://doi.org/10.1016/j.neuroimage.2009.01.045

Düzel, S., Drewelies, J., Gerstorf, D., Demuth, I., Steinhagen-Thiessen, E.,

Lindenberger, U., & Kühn, S. (2019). Structural Brain Correlates of Loneliness

among Older Adults. *Scientific Reports*, *9*(1), 13569.

https://doi.org/10.1038/s41598-019-49888-2

Emmerling, F., Schuhmann, T., Lobbestael, J., Arntz, A., Brugman, S., & Sack, A. T.

(2016). The role of the insular cortex in retaliation. PloS one, 11(4), e0152000.

Endo, K., Ando, S., Shimodera, S., Yamasaki, S., Usami, S., Okazaki, Y., Sasaki, T.,

Richards, M., Hatch, S., & Nishida, A. (2017). Preference for Solitude, Social

Isolation, Suicidal Ideation, and Self-Harm in Adolescents. *Journal of Adolescent*

Health, *61*(2), 187–191. https://doi.org/10.1016/j.jadohealth.2017.02.018

Eslinger, P. J., Anders, S., Ballarini, T., Boutros, S., Krach, S., Mayer, A. V., ... & Zahn,

R. (2021). The neuroscience of social feelings: mechanisms of adaptive social

functioning. Neuroscience & Biobehavioral Reviews, 128, 592-620.

Feng, C., Wang, L., Li, T., & Xu, P. (2019). Connectome-based individualized prediction

of loneliness. *Social Cognitive and Affective Neuroscience*, *14*(4), 353–365.

https://doi.org/10.1093/scan/nsz020

Fergusson, D. M., Horwood, L. J., Ridder, E. M., & Beautrais, A. L. (2005).

Subthreshold Depression in Adolescence and Mental Health Outcomes in

Adulthood. *Archives of General Psychiatry*, *62*(1), 66–72.

https://doi.org/10.1001/archpsyc.62.1.66

Fett, A. K., Shergill, S. S., & Krabbendam, L. (2015). Social neuroscience in psychiatry:

unravelling the neural mechanisms of social dysfunction. Psychological Medicine,

45(6), 1145-1165.

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der

Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N.,

Rosen, B., & Dale, A. M. (2002). Whole brain segmentation: Automated labeling

of neuroanatomical structures in the human brain. *Neuron*, *33*(3), 341–355.

https://doi.org/10.1016/s0896-6273(02)00569-x

Ford, T., Parker, C., Salim, J., Goodman, R., Logan, S., & Henley, W. (2018). The

relationship between exclusion from school and mental health: A secondary

analysis of the British Child and Adolescent Mental Health Surveys 2004 and

2007. *Psychological Medicine*, *48*(4), 629–641.

https://doi.org/10.1017/S003329171700215X

Foster, C. E., Horwitz, A., Thomas, A., Opperman, K., Gipson, P., Burnside, A., Stone,

D. M., & King, C. A. (2017). Connectedness to family, school, peers, and

community in socially vulnerable adolescents. *Children and Youth Services*

Review, *81*, 321–331. https://doi.org/10.1016/j.childyouth.2017.08.011

Fuhrmann, D., Casey, C. S., Speekenbrink, M., & Blakemore, S.-J. (2019). Social

exclusion affects working memory performance in young adolescent girls.

Developmental Cognitive Neuroscience, *40*, 100718.

https://doi.org/10.1016/j.dcn.2019.100718

Gallagher, M., Prinstein, M. J., Simon, V., & Spirito, A. (2014). Social Anxiety Symptoms

and Suicidal Ideation in a Clinical Sample of Early Adolescents: Examining

Loneliness and Social Support as Longitudinal Mediators. *Journal of Abnormal*

Child Psychology, *42*(6), 871–883. https://doi.org/10.1007/s10802-013-9844-7

Gao Y, Guo X, Zhong Y, Liu X, Tian S, Deng J, Lin X, Bao Y, Lu L, Wang G. Decreased

dorsal attention network homogeneity as a potential neuroimaging biomarker for

major depressive disorder. J Affect Disord. 2023 Jul 1;332:136-142.

Giordano, P. C. (2003). Relationships in Adolescence. *Annual Review of Sociology*,

29(Volume 29, 2003), 257–281.

https://doi.org/10.1146/annurev.soc.29.010202.100047

Goddings, A.-L., Mills, K. L., Clasen, L. S., Giedd, J. N., Viner, R. M., & Blakemore, S.-J.

(2014). The influence of puberty on subcortical brain development. *NeuroImage*,

88, 242–251. https://doi.org/10.1016/j.neuroimage.2013.09.073

Goetschius, L. G., Hein, T. C., Mitchell, C., Lopez-Duran, N. L., McLoyd, V. C., Brooks-

Gunn, J., McLanahan, S. S., Hyde, L. W., & Monk, C. S. (2020). Childhood

violence exposure and social deprivation predict adolescent amygdala-

orbitofrontal cortex white matter connectivity. *Developmental Cognitive*

Neuroscience, *45*, 100849. https://doi.org/10.1016/j.dcn.2020.100849

Gogolla, N. (2017). The insular cortex. Current Biology, 27(12), R580-R586.

Golde, S., Romund, L., Lorenz, R. C., Pelz, P., Gleich, T., Beck, A., & Raufelder, D.

(2019). Loneliness and Adolescents' Neural Processing of Self, Friends, and

Teachers: Consequences for the School Self-Concept. *Journal of Research on*

Adolescence, *29*(4), 938–952. https://doi.org/10.1111/jora.12433

Goosby, B. J., Bellatorre, A., Walsemann, K. M., & Cheadle, J. E. (2013). Adolescent

Loneliness and Health in Early Adulthood. *Sociological Inquiry*, *83*(4), 505–536.

https://doi.org/10.1111/soin.12018

Haber SN, Knutson B. The reward circuit: linking primate anatomy and human imaging.

Neuropsychopharmacology. 2010 Jan;35(1):4-26.

Hagler, D. J., Hatton, S., Cornejo, M. D., Makowski, C., Fair, D. A., Dick, A. S.,

Sutherland, M. T., Casey, B. J., Barch, D. M., Harms, M. P., Watts, R., Bjork, J.

M., Garavan, H. P., Hilmer, L., Pung, C. J., Sicat, C. S., Kuperman, J., Bartsch,

H., Xue, F., … Dale, A. M. (2019). Image processing and analysis methods for

the Adolescent Brain Cognitive Development Study. *NeuroImage*, *202*, 116091.

https://doi.org/10.1016/j.neuroimage.2019.116091

Hall-Lande, J. A., Eisenberg, M. E., Christenson, S. L., & Neumark-Sztainer, D. (2007).

Social Isolation, Psychological Health, and Protective Factors in Adolescence.

Adolescence, *42*(166), 265–286.

Harman, G., Kliamovich, D., Morales, A. M., Gilbert, S., Barch, D. M., Mooney, M. A.,

Feldstein Ewing, S. W., Fair, D. A., & Nagel, B. J. (2021). Prediction of suicidal

ideation and attempt in 9 and 10 year-old children using transdiagnostic risk

features. *PloS One*, *16*(5), e0252114.

https://doi.org/10.1371/journal.pone.0252114

Hawkley, L. C., & Cacioppo, J. T. (2010). Loneliness Matters: A Theoretical and

Empirical Review of Consequences and Mechanisms. *Annals of Behavioral*

Medicine, *40*(2), 218–227. https://doi.org/10.1007/s12160-010-9210-8

Hawkley, L. C., Thisted, R. A., Masi, C. M., & Cacioppo, J. T. (2010). Loneliness

predicts increased blood pressure: 5-year cross-lagged analyses in middle-aged

and older adults. *Psychology and Aging*, *25*(1), 132–141.

https://doi.org/10.1037/a0017805

Houghton, S., Kyron, M., Hunter, S. C., Lawrence, D., Hattie, J., Carroll, A., & Zadow,

C. (2022). Adolescents' longitudinal trajectories of mental health and loneliness:

The impact of COVID-19 school closures. *Journal of Adolescence*, *94*(2), 191–

205. https://doi.org/10.1002/jad.12017

Hu, L., Katz, E. S., & Stamoulis, C. (2023). Modulatory effects of fMRI acquisition time

of day, week and year on adolescent functional connectomes across spatial

scales: Implications for inference. *NeuroImage*, *284*, 120459.

https://doi.org/10.1016/j.neuroimage.2023.120459

Jaccard, J., Blanton, H., & Dodge, T. (2005). Peer influences on risk behavior: An

analysis of the effects of a close friend. *Developmental Psychology*, *41*(1), 135–

147. https://doi.org/10.1037/0012-1649.41.1.135

Jack, A., & Morris, J. P. (2014). Neocerebellar contributions to social perception in

adolescents with autism spectrum disorder. Developmental cognitive

neuroscience, 10, 77-92.

Jefferson, R., Barreto, M., Jones, F., Conway, J., Chohan, A., Madsen, K. R., Verity, L.,

Petersen, K. J., & Qualter, P. (2023). Adolescent loneliness across the world and

its relation to school climate, national culture and academic performance. *British*

Journal of Educational Psychology, *93*(4), 997–1016.

https://doi.org/10.1111/bjep.12616

Jefferson, R., Barreto, M., Verity, L., & Qualter, P. (2023). Loneliness During the School

Years: How It Affects Learning and How Schools Can Help. *Journal of School*

Health, *93*(5), 428–435. https://doi.org/10.1111/josh.13306

Jensen SK, Dickie EW, Schwartz DH, Evans CJ, Dumontheil I, Paus T, Barker ED.

Effect of Early Adversity and Childhood Internalizing Symptoms on Brain

Structure in Young Men. JAMA Pediatr. 2015 Oct;169(10):938-46

Jones, A. C., Schinka, K. C., van Dulmen, M. H. M., Bossarte, R. M., & Swahn, M. H.

(2011). Changes in Loneliness during Middle Childhood Predict Risk for

Adolescent Suicidality Indirectly through Mental Health Problems. *Journal of*

Clinical Child & Adolescent Psychology, *40*(6), 818–824.

https://doi.org/10.1080/15374416.2011.614585

Kanai, R., Bahrami, B., Duchaine, B., Janik, A., Banissy, M. J., & Rees, G. (2012). Brain

Structure Links Loneliness to Social Perception. *Current Biology*, *22*(20), 1975–

1979. https://doi.org/10.1016/j.cub.2012.08.045

Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E.

(2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in

the National Comorbidity Survey Replication. *Archives of General Psychiatry*,

62(6), 593–602. https://doi.org/10.1001/archpsyc.62.6.593

Klimstra, T. A., Hale III, W. W., Raaijmakers, Q. A. W., Branje, S. J. T., & Meeus, W. H.

J. (2010). Identity Formation in Adolescence: Change or Stability? *Journal of*

Youth and Adolescence, *39*(2), 150–162. https://doi.org/10.1007/s10964-009-

9401-4

Kong, X., Wei, D., Li, W., Cun, L., Xue, S., Zhang, Q., & Qiu, J. (2015). Neuroticism and

extraversion mediate the association between loneliness and the dorsolateral

prefrontal cortex. *Experimental Brain Research*, *233*(1), 157–164.

https://doi.org/10.1007/s00221-014-4097-4

Koolschijn, P. C. M. P., van IJzendoorn, M. H., Bakermans-Kranenburg, M. J., & Crone,

E. A. (2013). Hippocampal volume and internalizing behavior problems in

adolescence. *European Neuropsychopharmacology*, *23*(7), 622–628.

https://doi.org/10.1016/j.euroneuro.2012.07.001

Kroger, J. (2004). *Identity In Adolescence: The balance between self and other*.

Routledge. https://doi.org/10.4324/9780203346860

Lacey, R. E., Kumari, M., & Bartley, M. (2014). Social isolation in childhood and adult

inflammation: Evidence from the National Child Development Study,

Psychoneuroendocrinology, *50*, 85–94.

https://doi.org/10.1016/j.psyneuen.2014.08.007

Laidi C, Boisgontier J, de Pierrefeu A, Decreased Cortical Thickness in the Anterior

Cingulate Cortex in Adults with Autism. J Autism Dev Disord. 2019

Apr;49(4):1402-1409

Lam, J. A., Murray, E. R., Yu, K. E., Ramsey, M., Nguyen, T. T., Mishra, J., Martis, B.,

Thomas, M. L., & Lee, E. E. (2021). Neurobiology of loneliness: A systematic

review. *Neuropsychopharmacology*, *46*(11), 1873–1887.

https://doi.org/10.1038/s41386-021-01058-7

Lamm, C., & Singer, T. (2010). The role of anterior insular cortex in social emotions.

Brain structure and function, 214, 579-591.

Lahnakoski JM, Glerean E, Salmi J, Jääskeläinen IP, Sams M, Hari R, Nummenmaa L.

Naturalistic FMRI mapping reveals superior temporal sulcus as the hub for the

distributed brain network for social perception. Front Hum Neurosci. 2012 Aug

13;6:233.

Lavigne-Cerván, R., Costa-López, B., Juárez-Ruiz de Mier, R., Real-Fernández, M.,

Sánchez-Muñoz de León, M., & Navarro-Soria, I. (2021). Consequences of

COVID-19 Confinement on Anxiety, Sleep and Executive Functions of Children

and Adolescents in Spain. *Frontiers in Psychology*, *12*, 565516.

https://doi.org/10.3389/fpsyg.2021.565516

Layden, E. A., Cacioppo, J. T., Cacioppo, S., Cappa, S. F., Dodich, A., Falini, A., &

Canessa, N. (2017). Perceived social isolation is associated with altered

functional connectivity in neural networks associated with tonic alertness and

executive control. *NeuroImage*, *145*(Pt A), 58–73.

https://doi.org/10.1016/j.neuroimage.2016.09.050

Lenroot, R. K., Gogtay, N., Greenstein, D. K., Wells, E. M., Wallace, G. L., Clasen, L.

S., Blumenthal, J. D., Lerch, J., Zijdenbos, A. P., Evans, A. C., Thompson, P. M.,

& Giedd, J. N. (2007). Sexual dimorphism of brain developmental trajectories

during childhood and adolescence. *NeuroImage*, *36*(4), 1065–1073.

https://doi.org/10.1016/j.neuroimage.2007.03.053

Lerner, R. M., & Steinberg, L. (2009). *Handbook of Adolescent Psychology, Volume 2:*

Contextual Influences on Adolescent Development. John Wiley & Sons.

Lewis, J. D., Evans, A. C., Tohka, J., Brain Development Cooperative Group, &

Pediatric Imaging, Neurocognition, and Genetics Study. (2018). T1 white/gray

contrast as a predictor of chronological age, and an index of cognitive

performance. *NeuroImage*, *173*, 341–350.

https://doi.org/10.1016/j.neuroimage.2018.02.050

Lind, A., Salomäki, S., Parkkola, R., Haataja, L., Rautava, P., Junttila, N., Koikkalainen,

J., Lötjönen, J., Saunavaara, V., Korja, R., & Group, P. S. (2020). Brain volumes

in relation to loneliness and social competence in preadolescents born very

preterm. *Brain and Behavior*, *10*(6), e01640. https://doi.org/10.1002/brb3.1640

Liu, H., Wang, Y., Liu, W., Wei, D., Yang, J., Du, X., Tian, X., & Qiu, J. (2016).

Neuroanatomical correlates of attitudes toward suicide in a large healthy sample:

A voxel-based morphometric analysis. *Neuropsychologia*, *80*, 185–193.

https://doi.org/10.1016/j.neuropsychologia.2015.11.012

Liu J, Wang L, Zhang L, Ding Y, Zhang X, Hu Z, Zhao X. Abnormal amygdala volume

moderates parenting and anxiety symptoms in children and adolescents with

anxiety disorder. J Psychiatr Res. 2024 Jul;175:316-322.

Lodder, G. M. A., Scholte, R. H. J., Goossens, L., & Verhagen, M. (2017). Loneliness in

Early Adolescence: Friendship Quantity, Friendship Quality, and Dyadic

Processes. *Journal of Clinical Child & Adolescent Psychology*, *46*(5), 709–720.

https://doi.org/10.1080/15374416.2015.1070352

London, R., & Ingram, D. (2018). Social Isolation in Middle School. *School Community*

Journal, *28*(1), 107–127.

Luna, B. (2009). Developmental Changes in Cognitive Control through Adolescence. In

P. Bauer (Ed.), *Advances in Child Development and Behavior* (Vol. 37, pp. 233–

278). JAI. https://doi.org/10.1016/S0065-2407(09)03706-9

Mahon, N. E., Yarcheski, A., & Yarcheski, T. J. (1993). Health consequences of

loneliness in adolescents. *Research in Nursing & Health*, *16*(1), 23–31.

https://doi.org/10.1002/nur.4770160105

Manning, C., & Gregoire, A. (2009). Effects of parental mental illness on children.

Psychiatry, *8*(1), 7–9. https://doi.org/10.1016/j.mppsy.2008.10.012

Mäntymaa, M., Puura, K., Luoma, I., Kaukonen, P., Salmelin, R. K., & Tamminen, T. (2008). Infants' social withdrawal and parents' mental health. *Infant Behavior and Development*, *31*(4), 606-613.

Marraccini, M. E., & Brier, Z. M. F. (2017). School connectedness and suicidal thoughts and behaviors: A systematic meta-analysis. *School Psychology Quarterly*, *32*(1), 5–21. https://doi.org/10.1037/spq0000192

Matthews, T., Danese, A., Wertz, J., Ambler, A., Kelly, M., Diver, A., Caspi, A., Moffitt, T. E., & Arseneault, L. (2015). Social Isolation and Mental Health at Primary and Secondary School Entry: A Longitudinal Cohort Study. *Journal of the American Academy of Child & Adolescent Psychiatry*, *54*(3), 225–232. https://doi.org/10.1016/j.jaac.2014.12.008

Matthews, T., Qualter, P., Bryan, B. T., Caspi, A., Danese, A., Moffitt, T. E., Odgers, C. L., Strange, L., & Arseneault, L. (2023). The developmental course of loneliness in adolescence: Implications for mental health, educational attainment, and psychosocial functioning. *Development and Psychopathology*, *35*(2), 537–546. https://doi.org/10.1017/S0954579421001632

Matthews, T., Rasmussen, L. J. H., Ambler, A., Danese, A., Eugen-Olsen, J., Fancourt, D., Fisher, H. L., Iversen, K. K., Schultz, M., Sugden, K., Williams, B., Caspi, A., & Moffitt, T. E. (2024). Social isolation, loneliness, and inflammation: A multi-cohort investigation in early and mid-adulthood. *Brain, Behavior, and Immunity*, *115*, 727–736. https://doi.org/10.1016/j.bbi.2023.11.022

Maxwell, K. A. (2002). Friends: The Role of Peer Influence Across Adolescent Risk Behaviors. *Journal of Youth and Adolescence*, *31*(4), 267–277.

https://doi.org/10.1023/A:1015493316865

McClelland, H., Evans, J. J., Nowland, R., Ferguson, E., & O'Connor, R. C. (2020).

Loneliness as a predictor of suicidal ideation and behaviour: A systematic review

and meta-analysis of prospective studies. *Journal of Affective Disorders*, *274*,

880–896. https://doi.org/10.1016/j.jad.2020.05.004

McIver, T. A., Bosma, R. L., Goegan, S., Sandre, A., Klassen, J., Chiarella, J., Booij, L.,

& Craig, W. (2019). Functional connectivity across social inclusion and exclusion

is related to peer victimization and depressive symptoms in young adults. *Journal*

of Affective Disorders, *253*, 366–375. https://doi.org/10.1016/j.jad.2019.04.085

Meeus, W. (1996). Studies on identity development in adolescence: An overview of

research and some new data. *Journal of Youth and Adolescence*, *25*(5), 569–

598. https://doi.org/10.1007/BF01537355

Mertse N, Denier N, Walther S, Breit S, Grosskurth E, Federspiel A, Wiest R, Bracht T.

Associations between anterior cingulate thickness, cingulum bundle

microstructure, melancholia and depression severity in unipolar depression. J

Affect Disord. 2022 Mar 15;301:437-444.

Merz EC, Tottenham N, Noble KG. Socioeconomic Status, Amygdala Volume, and

Internalizing Symptoms in Children and Adolescents. J Clin Child Adolesc

Psychol. 2018 Mar-Apr;47(2):312-323

Mills, K. L., Goddings, A.-L., Herting, M. M., Meuwese, R., Blakemore, S.-J., Crone, E.

A., Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., & Tamnes, C. K.

(2016). Structural brain development between childhood and adulthood:

Convergence across four longitudinal samples. *NeuroImage*, *141*, 273–281.

https://doi.org/10.1016/j.neuroimage.2016.07.044

Murtaza, G., Sultana, R., Abualait, T., Fatima, M., & Bashir, S. (2023). Social isolation

during the COVID-19 pandemic is associated with the decline in cognitive

functioning in young adults. *PeerJ*, *11*, e16532.

https://doi.org/10.7717/peerj.16532

Mushtaq, R., Shoib, S., Shah, T., & Mushtaq, S. (2014). Relationship Between

Loneliness, Psychiatric Disorders and Physical Health ? A Review on the

Psychological Aspects of Loneliness. *Journal of Clinical and Diagnostic*

Research : JCDR, *8*(9), WE01–WE04.

https://doi.org/10.7860/JCDR/2014/10077.4828

Nakagawa, S., Takeuchi, H., Taki, Y., Nouchi, R., Sekiguchi, A., Kotozaki, Y., Miyauchi,

C. M., Iizuka, K., Yokoyama, R., Shinada, T., Yamamoto, Y., Hanawa, S., Araki,

T., Hashizume, H., Kunitoki, K., Sassa, Y., & Kawashima, R. (2015). White

matter structures associated with loneliness in young adults. *Scientific Reports*,

5, 17001. https://doi.org/10.1038/srep17001

Next-Generation Neural Data Analysis (NGNDA) (Version 1.0). (2020). [Computer

software]. Boston Childrens Hospital/Harvard Medical School.

https://github.com/cstamoulis1/Next-Generation-Neural-Data-Analysis-NGNDA-

Oh, W., Rubin, K. H., Bowker, J. C., Booth-LaForce, C., Rose-Krasnor, L., & Laursen,

B. (2008). Trajectories of Social Withdrawal from Middle Childhood to Early

Adolescence. *Journal of Abnormal Child Psychology*, *36*(4), 553–566.

https://doi.org/10.1007/s10802-007-9199-z

Oldham S, Fornito A. The development of brain network hubs. Dev Cogn Neurosci.

2019 Apr;36:100607.

Olivito G, Siciliano L, Clausi S, Lupo M, Baiocco R, Gragnani A, Saettoni M, Delle

Chiaie R, Laghi F, Leggio M. The Cerebellum Gets Social: Evidence from an

Exploratory Study of Cerebellar, Neurodevelopmental, and Psychiatric Disorders.

Biomedicines. 2023; 11(2):309. https://doi.org/10.3390/biomedicines11020309

Orben, A., Tomova, L., & Blakemore, S.-J. (2020). The effects of social deprivation on

adolescent development and mental health. *The Lancet. Child & Adolescent*

Health, *4*(8), 634–640. https://doi.org/10.1016/S2352-4642(20)30186-3

1091

Espinoza Oyarce DA, Shaw ME, Alateeq K, Cherbuin N. Volumetric brain differences in

clinical depression in association with anxiety: a systematic review with meta-

analysis. J Psychiatry Neurosci. 2020 Nov 1;45(6):406-429.

Özdemir, A., Utkualp, N., & Palloş, A. (2016). *Physical and Psychosocial Effects of the*

Changes in Adolescence Period.

Pan, L. A., Ramos, L., Segreti, A., Brent, D. A., & Phillips, M. L. (2015). Right superior

temporal gyrus volume in adolescents with a history of suicide attempt. The

British Journal of Psychiatry, 206(4), 339-340.

Palejwala AH, Dadario NB, Young IM, O'Connor K, Briggs RG, Conner AK,

O'Donoghue DL, Sughrue ME. Anatomy and White Matter Connections of the

Lingual Gyrus and Cuneus. World Neurosurg. 2021 Jul;151:e426-e437.

Pasqualetti, F., Zhao, S., Favaretto, C., & Zampieri, S. (2020). Fragility Limits

Performance in Complex Networks. *Scientific Reports*, *10*(1), 1774.

https://doi.org/10.1038/s41598-020-58440-6

Patterson, A. C., & Veenstra, G. (2010). Loneliness and risk of mortality: A longitudinal

investigation in Alameda County, California. *Social Science & Medicine (1982)*,

71(1), 181–186. https://doi.org/10.1016/j.socscimed.2010.03.024

Patton, G. C., Bond, L., Carlin, J. B., Thomas, L., Butler, H., Glover, S., Catalano, R., &

Bowes, G. (2006). Promoting Social Inclusion in Schools: A Group-Randomized

Trial of Effects on Student Health Risk Behavior and Well-Being. *American*

Journal of Public Health, *96*(9), 1582–1587.

https://doi.org/10.2105/AJPH.2004.047399

Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders

emerge during adolescence? *Nature Reviews. Neuroscience*, *9*(12), 947–957.

https://doi.org/10.1038/nrn2513

Pelphrey, K. A., & Carter, E. J. (2008). Brain mechanisms for social perception: lessons

from autism and typical development. *Annals of the New York Academy of*

Sciences, *1145*(1), 283-299.

Peper, J. S., Hulshoff Pol, H. E., Crone, E. A., & van Honk, J. (2011). Sex steroids and

brain structure in pubertal boys and girls: A mini-review of neuroimaging studies.

Neuroscience, *191*, 28–37. https://doi.org/10.1016/j.neuroscience.2011.02.014

Pereira Camejo, M., Escobar Saade, L., Liverani, M.C. et al. Amygdala volumes and

associations with socio-emotional competencies in preterm youth: cross-

sectional and longitudinal data. Pediatr Res 96, 1868–1877 (2024)

Pfeifer, J. H., & Allen, N. B. (2021). Puberty Initiates Cascading Relationships Between

Neurodevelopmental, Social, and Internalizing Processes Across Adolescence.

Biological Psychiatry, *89*(2), 99–108.

https://doi.org/10.1016/j.biopsych.2020.09.002

Phillips, J. R., Hewedi, D. H., Eissa, A. M., & Moustafa, A. A. (2015). The cerebellum

and psychiatric disorders. Frontiers in public health, 3, 66.

Porcelli, S., Van Der Wee, N., van der Werff, S., Aghajani, M., Glennon, J. C., van

Heukelum, S., Mogavero, F., Lobo, A., Olivera, F. J., Lobo, E., Posadas, M.,

Dukart, J., Kozak, R., Arce, E., Ikram, A., Vorstman, J., Bilderbeck, A., Saris, I.,

Kas, M. J., & Serretti, A. (2019). Social brain, social dysfunction and social

withdrawal. *Neuroscience & Biobehavioral Reviews*, *97*, 10–33.

https://doi.org/10.1016/j.neubiorev.2018.09.012

Preston, Angela J., & Rew, L. (2022). *Connectedness, Self-Esteem, and Prosocial*

Behaviors Protect Adolescent Mental Health Following Social Isolation: A

Systematic Review.

Reiss, F. (2013). Socioeconomic inequalities and mental health problems in children

and adolescents: A systematic review. *Social Science & Medicine*, *90*, 24–31.

https://doi.org/10.1016/j.socscimed.2013.04.026

Reupert, A., Gladstone, B., Helena Hine, R., Yates, S., McGaw, V., Charles, G., Drost,

L., & Foster, K. (2021). Stigma in relation to families living with parental mental

illness: An integrative review. *International Journal of Mental Health Nursing*,

30(1), 6–26. https://doi.org/10.1111/inm.12820

Riedel, M. C., Yanes, J. A., Ray, K. L., Eickhoff, S. B., Fox, P. T., Sutherland, M. T., &

Laird, A. R. (2018). Dissociable meta-analytic brain networks contribute to

coordinated emotional processing. Human brain mapping, 39(6), 2514-2531.

Rogol, A. D., Roemmich, J. N., & Clark, P. A. (2002). Growth at puberty. *Journal of*

Adolescent Health, *31*(6, Supplement), 192–200. https://doi.org/10.1016/S1054-

139X(02)00485-8

Rosada C, Bauer M, Golde S, Metz S, Roepke S, Otte C, Buss C, Wingenfeld K.

Childhood trauma and cortical thickness in healthy women, women with post-

traumatic stress disorder, and women with borderline personality disorder.

Psychoneuroendocrinology. 2023 Jul;153:106118.

Rubin, K. H., Coplan, R. J., & Bowker, J. C. (2009). Social Withdrawal in Childhood.

Annual Review of Psychology, *60*(Volume 60, 2009), 141–171.

https://doi.org/10.1146/annurev.psych.60.110707.163642

Rubin, K. H., Wojslawowicz, J. C., Rose-Krasnor, L., Booth-LaForce, C., & Burgess, K.

B. (2006). The Best Friendships of Shy/Withdrawn Children: Prevalence,

Stability, and Relationship Quality. *Journal of Abnormal Child Psychology*, *34*(2),

139–153. https://doi.org/10.1007/s10802-005-9017-4

Rubin, K. H., & Lollis, S. P. (2015). Origins and consequences of social withdrawal.

In *Clinical implications of attachment* (pp. 219-252). Routledge.

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity:

Uses and interpretations. *NeuroImage*, *52*(3), 1059–1069.

https://doi.org/10.1016/j.neuroimage.2009.10.003

Sato W, Kubota Y, Kochiyama T, Uono S, Yoshimura S, Sawada R, Sakihama M,

Toichi M. Increased putamen volume in adults with autism spectrum disorder.

Front Hum Neurosci. 2014 Nov 25;8:957

Sawyer, S. M., Azzopardi, P. S., Wickremarathne, D., & Patton, G. C. (2018). The age

of adolescence. *The Lancet Child & Adolescent Health*, *2*(3), 223–228.

https://doi.org/10.1016/S2352-4642(18)30022-1

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J.,

Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human

Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*

(New York, N.Y.: 1991), *28*(9), 3095–3114. https://doi.org/10.1093/cercor/bhx179

Schinka, K. C., VanDulmen, M. H. M., Bossarte, R., & Swahn, M. (2012). Association

Between Loneliness and Suicidality During Middle Childhood and Adolescence:

Longitudinal Effects and the Role of Demographic Characteristics. *The Journal of*

Psychology, *146*(1–2), 105–118. https://doi.org/10.1080/00223980.2011.584084

Schlack, R., Peerenboom, N., Neuperdt, L., Junker, S., & Beyer, A.-K. (2021). The

effects of mental health problems in childhood and adolescence in young adults:

Results of the KiGGS cohort. *Journal of Health Monitoring*, *6*(4), 3–19.

https://doi.org/10.25646/8863

Schwartz-Mette, R. A., Shankman, J., Dueweke, A. R., Borowski, S., & Rose, A. J.

(2020). Relations of friendship experiences with depressive symptoms and

loneliness in childhood and adolescence: A meta-analytic review. *Psychological*

Bulletin, *146*(8), 664–700. https://doi.org/10.1037/bul0000239

Sheffield, J.M., Huang, A.S., Rogers, B.P. et al. Insula sub-regions across the psychosis

spectrum: morphology and clinical correlates. Transl Psychiatry 11, 346 (2021).

Siervogel, R. M., Demerath, E. W., Schubert, C., Remsberg, K. E., Chumlea, W. C.,

Sun, S., Czerwinski, S. A., & Towne, B. (2004). Puberty and Body Composition.

Hormone Research, *60*(Suppl. 1), 36–45. https://doi.org/10.1159/000071224

Smith, C., & Stamoulis, C. (2023). Effects of multidomain environmental and mental

health factors on the development of empathetic behaviors and emotions in

adolescence. *PLoS ONE*, *18*(11). https://doi.org/10.1371/journal.pone.0293473

Steinberg, L., & Morris, A. S. (2001). Adolescent development. *Annual Review of*

Psychology, *52*, 83–110. https://doi.org/10.1146/annurev.psych.52.1.83

Stenseng, F., Tingstad, E. B., Wichstrøm, L., & Skalicka, V. (2022). Social withdrawal

and academic achievement, intertwined over years? Bidirectional effects from

primary to upper secondary school. *British Journal of Educational Psychology*,

92(4), 1354–1365. https://doi.org/10.1111/bjep.12504

Sturman, D. A., & Moghaddam, B. (2011). The neurobiology of adolescence: Changes

in brain architecture, functional dynamics, and behavioral tendencies.

Neuroscience & Biobehavioral Reviews, *35*(8), 1704–1712.

https://doi.org/10.1016/j.neubiorev.2011.04.003

Suor JH, Jimmy J, Monk CS, Phan KL, Burkhouse KL. Parsing differences in amygdala

volume among individuals with and without social and generalized anxiety

disorders across the lifespan. J Psychiatr Res. 2020 Sep;128:83-89.

Susman, E. J., & Rogol, A. (2004). Puberty and Psychological Development. In

Handbook of Adolescent Psychology (pp. 15–44). John Wiley & Sons, Ltd.

https://doi.org/10.1002/9780471726746.ch2

Terasawa, Y., Shibata, M., Moriguchi, Y., & Umeda, S. (2013). Anterior insular cortex

mediates bodily sensibility and social anxiety. Social cognitive and affective

neuroscience, 8(3), 259-266.

Thomas, E., Juliano, A., Owens, M., Cupertino, R. B., Mackey, S., Hermosillo, R.,

Miranda-Dominguez, O., Conan, G., Ahmed, M., Fair, D. A., Graham, A. M.,

Goode, N. J., Kandjoze, U. P., Potter, A., Garavan, H., & Albaugh, M. D. (2024).

Amygdala connectivity is associated with withdrawn/depressed behavior in a

large sample of children from the Adolescent Brain Cognitive Development

(ABCD) Study®. *Psychiatry Research. Neuroimaging*, *344*, 111877.

https://doi.org/10.1016/j.pscychresns.2024.111877

Thompson, K. N., Odgers, C. L., Bryan, B. T., Danese, A., Milne, B. J., Strange, L.,

Matthews, T., & Arseneault, L. (2022). Trajectories of childhood social isolation in

a nationally representative cohort: Associations with antecedents and early

adulthood outcomes. *JCPP Advances*, *2*(2), e12073.

https://doi.org/10.1002/jcv2.12073

Tian, Y., Liang, S., Yuan, Z., Chen, S., Xu, P., & Yao, D. (2014). White matter structure

in loneliness: Preliminary findings from diffusion tensor imaging. *Neuroreport*,

25(11), 843–847. https://doi.org/10.1097/WNR.0000000000000197

Tian, Y., Margulies, D. S., Breakspear, M., & Zalesky, A. (2020). Topographic

organization of the human subcortex unveiled with functional connectivity

gradients. *Nature Neuroscience*, *23*(11), 1421–1432.

https://doi.org/10.1038/s41593-020-00711-6

Tian, Y., Yang, L., Chen, S., Guo, D., Ding, Z., Tam, K. Y., & Yao, D. (2017). Causal

interactions in resting-state networks predict perceived loneliness. *PloS One*,

12(5), e0177443. https://doi.org/10.1371/journal.pone.0177443

Tomova, L., Towner, E., Thomas, K., Zhang, L., Palminteri, S., & Blakemore, S.-J.

(2022). *Acute isolation is associated with increased reward responsiveness in*

human adolescents. OSF. https://doi.org/10.31234/osf.io/ndt9k

Tooley UA, Park AT, Leonard JA, Boroshok AL, McDermott CL, Tisdall MD, Bassett DS, Mackey AP. The Age of Reason: Functional Brain Network Development during Childhood. J Neurosci. 2022 Nov 2;42(44):8237-8251.

Turrini S, Avenanti A. Cerebellum function: The chronometry of social perception. Curr Biol. 2024 May 6;34(9):R340-R343.

Vaisvilaite, L., Hushagen, V., Grønli, J., & Specht, K. (2022). Time-of-Day Effects in Resting-State Functional Magnetic Resonance Imaging: Changes in Effective Connectivity and Blood Oxygenation Level Dependent Signal. *Brain Connectivity*, *12*(6), 515–523. https://doi.org/10.1089/brain.2021.0129

Van Loon, L. M. A., Van de Ven, M. O. M., Van Doesum, K. T. M., Witteman, C. L. M., & Hosman, C. M. H. (2014). The Relation Between Parental Mental Illness and Adolescent Mental Health: The Role of Family Factors. *Journal of Child and Family Studies*, *23*(7), 1201–1214. https://doi.org/10.1007/s10826-013-9781-7

Van Overwalle, F., Manto, M., Cattaneo, Z., Clausi, S., Ferrari, C., Gabrieli, J. D., ... & Leggio, M. (2020). Consensus paper: cerebellum and social cognition. *The Cerebellum*, *19*, 833-868.

Van Overwalle, F. Social and emotional learning in the cerebellum. Nat. Rev. Neurosci. 25, 776–791 (2024)

van Tol MJ, Li M, Metzger CD, Hailla N, Horn DI, Li W, Heinze HJ, Bogerts B, Steiner J, He H, Walter M. Local cortical thinning links to resting-state disconnectivity in major depressive disorder. Psychol Med. 2014 Jul;44(10):2053-65.

Vanhalst, J., Luyckx, K., & Goossens, L. (2014). Experiencing Loneliness in Adolescence: A Matter of Individual Characteristics, Negative Peer Experiences,

or Both? *Social Development*, *23*(1), 100–118.

https://doi.org/10.1111/sode.12019

Vasey, M. W. (2001). *The Developmental Psychopathology of Anxiety*. Oxford

University Press, USA.

Vijayakumar, N., Op de Macks, Z., Shirtcliff, E. A., & Pfeifer, J. H. (2018). Puberty and

the human brain: Insights into adolescent development. *Neuroscience &*

Biobehavioral Reviews, *92*, 417–436.

https://doi.org/10.1016/j.neubiorev.2018.06.004

Vitale EM, Smith AS. Neurobiology of Loneliness, Isolation, and Loss: Integrating

Human and Animal Perspectives. Front Behav Neurosci. 2022 Apr 8;16:846315.

Voelbel GT, Bates ME, Buckman JF, Pandina G, Hendren RL. Caudate nucleus volume

and cognitive performance: Are they related in childhood psychopathology? Biol

Psychiatry. 2006 Nov 1;60(9):942-50.

Vogt, Pain and Emotion Interactions in Subregions of the Cingulate Gyrus, Nat Rev

Neurosci. 2005 July ; 6(7): 533–544.

von Soest, T., Luhmann, M., & Gerstorf, D. (2020). The development of loneliness

through adolescence and young adulthood: Its nature, correlates, and midlife

outcomes. *Developmental Psychology*, *56*(10), 1919–1934.

https://doi.org/10.1037/dev0001102

Wang, M. Q., Fitzhugh, E. C., Westerfield, R. C., & Eddy, J. M. (1995). Family and peer

influences on smoking behavior among American adolescents: An age trend.

Journal of Adolescent Health, *16*(3), 200–203. https://doi.org/10.1016/1054-

139X(94)00097-X

Wang, L., Zhao, P., Zhang, J., Zhang, R., Liu, J., Duan, J., ... & Wang, F. (2023).

Functional connectivity between the cerebellar vermis and cerebrum

distinguishes early treatment response for major depressive episodes in

adolescents. Journal of Affective Disorders, 339, 256-263.

Westlye, L. T., Walhovd, K. B., Dale, A. M., Espeseth, T., Reinvang, I., Raz, N., Agartz,

I., Greve, D. N., Fischl, B., & Fjell, A. M. (2009). Increased sensitivity to effects of

normal aging and Alzheimer's disease on cortical thickness by adjustment for

local variability in gray/white contrast: A multi-sample MRI study. *NeuroImage*,

47(4), 1545–1557. https://doi.org/10.1016/j.neuroimage.2009.05.084

Wierenga, L., Langen, M., Ambrosino, S., van Dijk, S., Oranje, B., & Durston, S. (2014).

Typical development of basal ganglia, hippocampus, amygdala and cerebellum

from age 7 to 24. *NeuroImage*, *96*, 67–72.

https://doi.org/10.1016/j.neuroimage.2014.03.072

Wigfield, A., Byrnes, J. P., & Eccles, J. S. (2006). Development During Early and Middle

Adolescence. In *Handbook of educational psychology* (pp. 87–113). Lawrence

Erlbaum Associates Publishers.

Wong NML, Mabel-Kenzie STST, Lin C, Huang CM, Liu HL, Lee SH, Lee TMC. Meta-

analytic evidence for the cognitive control model of loneliness in emotion

processing. Neurosci Biobehav Rev. 2022 Jul;138:104686.

Wood, K. R., Coplan, R. J., Hipson, W. E., & Bowker, J. C. (2022). Normative Beliefs

about Social Withdrawal in Adolescence. *Journal of Research on Adolescence*,

32(1), 372–381. https://doi.org/10.1111/jora.12617

Woodhouse, S. S., Dykas, M. J., & Cassidy, J. (2012). Loneliness and Peer Relations in

Adolescence. *Social Development*, *21*(2), 273–293.

https://doi.org/10.1111/j.1467-9507.2011.00611.x

Wu, J., Barahona, M., Tan, Y.-J., & Deng, H.-Z. (2011). Spectral Measure of Structural

Robustness in Complex Networks. *IEEE Transactions on Systems, Man, and*

Cybernetics - Part A: Systems and Humans, *41*(6), 1244–1252. IEEE

Transactions on Systems, Man, and Cybernetics - Part A: Systems and Humans.

https://doi.org/10.1109/TSMCA.2011.2116117

Xiong, Y., Hong, H., Liu, C. *et al.* Social isolation and the brain: effects and

mechanisms. *Mol Psychiatry* 28, 191–201 (2023).

Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, Roffman JL,

Smoller JW, Zöllei L, Polimeni JR, Fischl B, Liu H, Buckner RL. The organization

of the human cerebral cortex estimated by intrinsic functional connectivity. J

Neurophysiol. 2011 Sep;106(3):1125-65

Yurgelun-Todd, D. (2007). Emotional and cognitive changes during adolescence.

Current Opinion in Neurobiology, *17*(2), 251–257.

https://doi.org/10.1016/j.conb.2007.03.009

Zaccarella, E., & Friederici, A. D. (2015). Merge in the human brain: A sub-region based

functional investigation in the left pars opercularis. Frontiers in psychology, 6,

1818.

Zheng, S., Chen, X., Liu, W., Li, Z., Xiao, M., Liu, Y., & Chen, H. (2023). Association of

loneliness and grey matter volume in the dorsolateral prefrontal cortex: The

mediating role of interpersonal self-support traits. *Brain Imaging and Behavior*,

17(5), 481–493. https://doi.org/10.1007/s11682-023-00776-4

Zhu J, Qiu A. Interindividual variability in functional connectivity discovers differential

development of cognition and transdiagnostic dimensions of psychopathology in

youth. Neuroimage. 2022 Oct 15;260:119482.

1339

1340

1341   **Table 1**. Demographic and other individual characteristics (n = 2809).

Age (years)	Median (IQR)	12.00 (1.08)
Sex	Male	1366 (48.63%)
	Female	1440 (51.26%)
	Missing	3 (0.11%)
Race/Ethnicity	White Non-Hispanic	1488 (52.97%)
	Black Non-Hispanic	363 (12.92%)
	Asian Non-Hispanic	645 (22.96%)
	Other (including mixed race) Non-Hispanic	61 (2.17%)
	Hispanic	243 (8.65%)
	Missing	9 (0.32%)
Family Income ($)	<25,000	266 (9.47%)
	25,000-49,999	331 (11.78%)
	50,000 - 74,999	355 (12.64%)
	75,000 - 99,999	361 (12.85%)
	100,000-199,999	883 (31.43%)
	>=200,000	405 (14.42%)
	Missing	208 (7.40%)
BMI Z-Score	Median (IQR)	-0.31 (1.138)
	Missing	17 (0.61%)
Pubertal Stage	Pre-Puberty	515 (18.33%)
	Early Puberty	609 (21.68%)
	Mid-Puberty	918 (32.68%)
	Late/Post-Puberty	622 (22.14%)
	Missing	145 (5.16%)

1342

1343   **Table 2**. Distribution of social environmental factors in the cohort.

PARENTAL FACTORS		
Parental mental health (T-score)	**Median (IQR)**	51 (4)
	Missing	378 (13.46%)
Parental belief that family members should show love and affection to one another	**Not at all**	15 (0.53%)
	A little	23 (0.82%)
	Somewhat	254 (9.04%)
	Very much	1237 (44.04%)
	Completely	1279 (45.53%)
	Missing	1 (0.04%)
FAMILY ENVIRONMENT		
Family conflict Scale: 0-9	**Median (IQR)**	2 (2)
	Missing	1 (0.04%)
Family cohesion Scale: 0-9	**Median (IQR)**	8 (2)
	Missing	1 (0.04%)
Family size	**Median (IQR)**	4 (1)
	Missing	36 (1.28%)
PEER RELATIONSHIPS		
Number of close friends	**Median (IQR)**	5 (5)
	Missing	6 (0.21%)
Has a regular group of kids to hang out with at school/neighborhood	**No**	276 (9.83%)
	Yes	2427 (86.40%)
	Missing	106 (3.77%)
Faced discrimination in the past 12 months	**No**	2507 (89.25%)
	Yes	302 (10.75%)
	Missing	0 (0%)
Problems with bullying at school/neighborhood OR cyberbullying	**No**	2276 (81.03%)
	Yes	519 (18.48%)
	Missing	14 (0.50%)
SCHOOL		
Gets along with their teachers	**Strongly disagree**	25 (0.89%)
	Disagree	120 (4.27%)
	Agree	1188 (42.29%)
	Strongly agree	1471 (52.37%)
	Missing	5 (0.18%)
Likes school a lot	**Strongly disagree**	259 (9.22%)
	Disagree	554 (19.72%)
	Agree	1204 (42.86%)
	Strongly agree	787 (28.02%)
	Missing	5 (0.18%)

1344

1345 **Table 3.** Statistics of mixed-effect models testing associations between preferring
1346 solitude, social withdrawal, and topological properties of individual networks. All
1347 reported p-values have been adjusted for the False Discovery Rate. Regression
1348 coefficients in models comparing pairs of groups were not standardized. *-:
1349 Nonsignificant; CI: confidence interval.

Would rather be alone than with others						
Property	Network	Sample comparison	Beta	95th % CI*	P-value	Beta
			Left hemisphere			Right he
Segregation	Dorsal attention	Entire sample	-*			0.066
	Social	Entire sample	-			0.085
		Somewhat vs Very True	-			0.008
	Temporoparietal	Entire sample	-			0.095
Global efficiency	Cerebellum	Somewhat vs Very True	-0.089	[-0.133, -0.045]	<0.001	-
Withdrawn, does not get involved with others						
Property	Network	Sample comparison	Beta	95th % CI	P-value	Beta
			Left hemisphere			Right he
Topological robustness	Cerebellum	Somewhat vs Very True	-0.781	[-1.342, -0.220]	0.021	-
Topological stability	Cerebellum	Somewhat vs Very True	-0.670	[-1.151, -0.189]	0.021	-
Fragility	Cerebellum	Somewhat vs Very True	0.039	[0.005, 0.073]	0.047	-

1350

1351 **Table 4.** Statistics of mixed-effect models testing associations between preferring
1352 solitude and morphological properties. All reported p-values have been adjusted for the
1353 False Discovery Rate. Regression coefficients in models comparing only 2 response
1354 options were not standardized. *-: Nonsignificant *CI: confidence interval.

			Would rather be alone than with others				
Property	Structure	Sample comparison	Beta	95th % CI*	P-value	Beta	
			Left hemisphere			Right her	
Cortical Thickness	Cuneus	Entire sample	-0.054	[-0.096, -0.011]	0.039	-*	
		Not vs Very True	-0.024	[-0.044, -0.004]	0.043	-	
	Superior temporal gyrus	Entire sample	-0.058	[-0.100, -0.016]	0.019	-	
		Not vs Very True	-0.024	[-0.045, -0.003]	0.0496	-	
	Caudal anterior cingulate gyrus	Entire sample	-			-0.086	[
		Not vs Somewhat True	-			-0.032	[
		Not vs Very True	-			-0.038	[
	Pars opercularis	Entire sample	-			-0.053	[
		Not vs Somewhat True	-			-0.018	[
	Insular gyrus	Entire sample	-			-0.062	[
		Not vs Very True	-			-0.035	[
Volume	Amygdala	Not vs Somewhat True	25.718	[0.128, 51.308]	0.049	-	
	Cuneus	Not vs Very True	-109.540	[-207.7, -11.4]	0.043	-	
	Entorhinal	Not vs Somewhat True	-			61.62	[
	Fusiform	Not vs Very True	-			-264.3	[
	Parahippocampal gyrus	Entire sample	-			0.055	[
		Not vs Somewhat True	-			53.95	[
	Supramarginal gyrus	Somewhat vs Very True	-1044.6	[-1851, -248]	0.031	-	
Cortical white matter intensity	Parahippocampal gyrus	Entire sample	-			-0.020	[
		Not vs Very True	-			-0.234	[
	Inferior temporal gyrus	Somewhat vs Very True	-			-0.389	[
			Withdrawn, does not get involved with others				

Property	Structure	Sample comparison	Statistic	Value	Statistic
			Left hemisphere		**Right hen**
Cortical thickness	**Cuneus**	**Not vs Very True**	-		-0.055 [
		Somewhat vs Very True	-		-0.146 [
	Insular gyrus	**Not vs Very True**	-		-0.057 [
	Lingual gyrus	**Not vs Very True**	-		-0.040 [
		Somewhat vs Very True	-		-0.090 [
Cortical volume	**Caudal middle frontal gyrus**	**Not vs Somewhat True**	-		-319.9 [
	Caudate nucleus	**Somewhat vs Very True**	308.498 [2.68, 614.31]	0.048	-
	Cuneus	**Not vs Very True**	-		-333.6 [
		Somewhat vs Very True	-		-723.3 [
	Lingual gyrus	**Not vs Very True**	-		-492.5 [
	Pallidum	**Not vs Very True**	-		81.06 [
		Somewhat vs Very True	-		231.7 [
	Putamen	**Somewhat vs Very True**	-		369.0 [
White matter intensity	**Isthmus of cingulate cortex**	**Entire sample**	-		0.026 [

1355

1356

1357 **FIGURES**

1359

1360 **Figure 1.** Associations between preferring solitude and cortical thickness. The color
1361 map represents the regression coefficients corresponding to cortical thickness.