original article to Primates

Analysis of individual identification and age-class classification of wild female

macaque vocalizations without pitch- and formant-based acoustic parameter

measurements

Rentaro Kimpara[1*], Fumiya Kakuta[2], Hiroki Koda[3,4], Ikki Matsuda[1,5,6], and Goro

Hanya[2]

[1] Wildlife Research Center, Kyoto University, Kyoto, Japan

[2] Center for Ecological Research, Kyoto University, Inuyama, Japan

[3] Graduate School of Arts and Sciences, The University of Tokyo, Japan

[4] Center of Evolutionary Cognitive Science, The University of Tokyo, Japan

[5] Chubu Institute for Advanced Studies, Chubu University, Kasugai, Japan

[6] Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Kota

Kinabalu, Malaysia

Correspondence to RK (kimpara.rentaro.75c@st.kyoto-u.ac.jp)

Abstract

In recent years, deep learning has achieved high performance in bioacoustic

classification tasks by leveraging automatically processed acoustic features for large

datasets. However, few performance evaluations of automatically processed acoustic

features have been conducted on small-scale data because deep learning requires large

datasets. To test whether mel spectrograms (an automatically processed acoustic

features) are effective for classifying relatively small acoustic data, we evaluated the

performance of two classification machines (random forest and support vector machine)

using mel-spectrograms of 651 coo calls of six wild female Japanese macaques on two

tasks: 1) individual identification and 2) age-class classification between younger (<10

yrs) and the older animals (>20 yrs). For the individual identification task, the mean

balanced accuracy was 0.81 for random forest and 0.82 for support vector machine. For

the age-class classification task, the mean balanced accuracy was 0.91 for random forest

and 0.93 for support vector machine. Considering that of all the calls were recorded in

the wild, methods using automatically processed acoustic features, such as mel

spectrogram, are effective in classifying small acoustic data for the individual

identification task. The high performance in the age-class classification task might be

attributable to the potential of mel spectrograms to capture the characteristics of older

individuals (e.g. harshness).

Introduction

To elucidate the vocal communication of animals, bioacoustic studies have

implemented tasks to classify calls from acoustic features for such purposes as

identifying individuals making vocalizations and classifying vocal repertoires. This

work has been pursued for more than half a century (Arnaud et al., 2023; Fukushima et

al., 2015; Marler, 1956; Marler and Hobbett, 1975; Mitani, 1986; Rendall et al., 1998;

Winter et al., 1966). Historically, conventional acoustic analysis has measured

numerous acoustic features like pitch and formant and has examined classification

performance using dimensionality reduction and multivariate analysis (Ceugniet and

Izumi, 2004; Oyakawa et al., 2007; Rendall et al., 1998). Reported classification

accuracies were 83.7-92.3%, 90.5-98.4%, and 63.6-90.9%, respectively. Although these

conventional approaches are still in use, they have suffered from problems such as the

arbitrary nature of selecting acoustic features and the lack of versatility due to the

definition of species-specific acoustic features.

Recent machine learning, especially deep learning, can automatically process

acoustic signals for classification tasks without explicitly measuring specific features,

which solves the problems associated with the conventional method (Arnaud et al.,

2023; Lakdari et al., 2024). Nevertheless, current deep-learning techniques have a

methodological shortcoming in that they typically require a large amount of supervised

data, often ranging from thousands to millions of annotated samples in animal sound

classification tasks (Nanni et al. 2020; Heinrich et al. 2025). This is an obstacle to

building models because preparing a large amount of supervised data requires a huge

amount of time or is nearly impossible due to difficulties such as collecting sufficient

data in the field. Therefore, developing classification models for relatively small

amounts of data without measuring specific acoustic features could alleviate the

difficulty of model building and facilitate bioacoustic studies of many species.

However, no such models have so far been built.

Coo call is a contact call of macaques and accounts for a large part of Japanese

macaque vocalizations (Koda and Sugiura, 2010). Acoustically, the characteristic of coo

call is a tonal sound, that is to say, the fundamental frequency of the harmonics is clear

(Green, 1975). Concerning individual identification from coo calls of Japanese macaques,

Ceugniet and Izumi suggest that call duration and start and end fundamental frequency

are crucial (Ceugniet and Izumi, 2004), and Mitani pointed out that fundamental

frequency is less variable in the same individual compared to call duration (Mitani, 1986).

Likewise, formant frequencies—which are generally determined by vocal tract length and

shape—have been examined to describe acoustic properties relevant to vocal identity. A

study in rhesus macaques, a closely related species, demonstrated that vocal tract

characteristics contribute to individual discrimination (Rendall et al., 1998), and this

finding has been supported by experimental work in Japanese macaques (Furuyama et al.,

2016). These studies attempted to identify individuals based on vocalizations by

measuring specific acoustic characteristics. However, to date, no research has applied

automatically extracted acoustic features to distinguish Japanese macaques based on their

vocalizations.

Acoustic features are relatively unstable due to age-related changes. Aging in

animals is associated with physical and cognitive changes (Hindle et al., 2009; Manrique

and Call, 2015). In humans, aging physio-morphologically affects acoustic features such

as fundamental frequency and signal-to-noise ratio (Abitbol et al., 1999; Ferrand, 2002;

Pontes et al., 2006; Ryan and Burk, 1974; Stathopoulos et al., 2011), suggesting that the

age of a human could be identified from the voice (Harnsberger et al., 2008; Huntley et

al., 1987). Interestingly, some studies suggest that impressions of the speaker are

influenced by age-related acoustic features such as pitch (Feinberg et al., 2005; Klofstad

et al., 2015; Tigue et al., 2012). While fewer studies have explored senescence-related

acoustic changes in non-human animals, some research suggests that such changes

influence the listener's perceptual estimation of caller age based on vocalization

(Lemasson et al., 2010; Zipple et al., 2020). However, bioacoustics research has largely

focused on early life stages, with limited attention to senescence (reviewed by Ey et al.

2007).

Individual identification of vocalizations or classification of vocal repertoires,

notably in group-living primates, has been attempted repeatedly over the past half-

century (Arnaud et al., 2023; Fukushima et al., 2015; Marler and Hobbett, 1975; Mitani,

1986; Rendall et al., 1998). However, few studies have attempted to identify vocalizers

or classify age classes in wild primates without explicitly measuring specific acoustic

features such as pitch or resonance frequency. In this study, therefore, we address the

tasks of vocalizer identification and age-class classification (younger: <10 yrs or older:

20 > yrs) in wild Japanese macaques (*Macaca fuscata yakui*) without measuring

specific acoustic features.

Materials and methods

Recordings of vocalization

The Japanese macaques (*Macaca fuscata yakui*) of Yakushima Island in

southern Japan (30.41N, 130.41E), have been studied intensively since the 1970s, and

multiple groups have been identified (Yamagiwa and Hill, 1998). We investigated one

of the identified groups, named the Petit group, from October to December 2023. At that

time, the Petit Group was composed of about 33 individuals, including 10 adult females

(>=6 years), 5 or more adult males (>=6 years), 8 juvenile females (1–5 years), 7

juvenile males (1–5 years), and 3 infants. We selected 10 adult females as focal animals

and collected vocalization data using the focal animal sampling method, where RK and

FK followed the focal animal. The reason we only focused on adult females is that we

concerned that male individuals vocalize infrequently (Mitani, 1986), making it difficult

to obtain sufficient vocalizations in the wild. One observation session lasted two hours,

and we randomly changed focal individuals from those with the least observation time.

We maintained a distance of 3 meters or less from the focal individual. We did not

follow estrous females although the observation period included mating season. We

recorded vocalizations with annotations of the vocalizer using an audio recorder with

two channels and 32-bit-float/48 kHz sampling rate formats (48 kHz), which was

connected to a directional microphone (Sennheiser MKE600) for recording the animal's

vocalizations and a lavalier microphone (COMICA CVM-V02O) for recording the

observer's annotations, such as recording vocalizers. The total observation time was 112

hours over 20 days. We created classifiers for six adult females (Kapa, 9 yrs; Rine, 8

yrs; Sasa, 7 yrs; Sazae, estimated 21-22 yrs; Taiko, estimated 23-25 yrs; Taiwu, 9 yrs),

for which we obtained sufficient high-quality vocal data. The other four adult females

were excluded because they had fewer than 80 high-quality calls. The genetic

relationship of the six adult females used for machine learning is not fully known, but

two pairs (Sazae and Sasa; Taiko and Taiwu) are mother-daughter pairs.

Analysis

Data selection and preprocessing

138   We used the acoustic analysis software Praat (Boersma, 2024) for annotation

to select data in the recordings for which we could reliably identify the vocalizer. After

the annotation, we selected the calls that met all five of the following criteria for further

analysis: 1) Little background noise, 2) coo call (see Koda and Sugiura, 2010 for its

common definition and function), 3) no overlap with other individuals, and 4) certainty

about the vocalizer, and 5) less than 1 second, based on the information by Mitani

(1986) and Sugiura (1998). From 1181 audio recordings, we finally selected 116 calls

for Kapa (102 calls excluded), 89 for Rine (149 calls excluded), 138 for Sasa (84 calls

excluded), 122 for Sazae (84 calls excluded), 84 for Taiko (77 calls excluded), and 102

for Taiwu (34 calls excluded), and we used them in subsequent models for classification

tasks. Noted that even though the recordings were made continuously and the time of

recording is the same among individuals, thereby minimizing bias, it cannot be denied

that bias may have been introduced during the data preprocessing. Situations prone to

noise, such as those with strong winds and proximity to streams, probably tended to be

removed.

Generation of Mel Spectrograms

155   A mel spectrogram is a set of acoustic features derived through logarithmic

scale filters designed using the frequency-sensitivity characteristics of human auditory

perception, (Stevens et al., 1937) and it is often used values in machine learning

(Baowaly et al., 2024; Dossou and Gbenou, 2021; Walsh et al., 2023; Xie et al., 2024).

We also used mel-spectrograms in this study, as previous work showed that UMAP

successfully classified animal call types, such as those of meerkats (Thomas et al.,

2022).

162   We generated mel spectrograms of the vocal region with Python's audio

processing module *librosa* (McFee et al., 2015). The original code for the generation of

mel spectrograms was based on a previous bioacoustics study on meerkat calls

including seven call types (Thomas et al., 2022), and we set the key parameters of

generating mel spectrograms generations as follows: N_MELS=40 (number of Mel

bands), FFT_WIN=0.03 (30-ms window size for STFT), FFT_HOP=0.00375 (hop size,

set to 1/8th of the window size), WINDOW="hann" (Hann window function), FMIN=0,

and FMAX=24000 (frequency range in Hz). To handle varying call durations, zero-

padding was applied to match the length of the longest call, ensuring consistent input

dimensions for the classifiers. Each spectrogram was also z-transformed to normalize

intensity values across calls, minimizing the effects of amplitude variation. These

parameter settings, along with zero-padding and z-transformation, were all based on the

suggestions of previous study (Thomas et al., 2022). This process resulted in 10,360

dimensions for each call.

Data structure analysis with UMAP

To assess the separability of individual classes and age classes, we performed

dimensionality reduction using Uniform Manifold Approximation and Projection

(UMAP) from *umap-learn* (McInnes et al., 2018). We repeated supervised UMAP and

unsupervised UMAP 100 times each using all calls (651 calls). Supervised UMAP aims

to create a latent space where labels (e.g., individual identity) help guide the projection,

making it easier to see how well the classes separate. In contrast, unsupervised UMAP

does not use labels and reveals the natural structure of the data. In this study, we used

two types of labels: individual and age-class (older (over 20 years) and younger (less

than 10 years)). Both methods used the same parameters as those in the previous

research (Thomas et al., 2022) except for the n_components used for comparison in the

previous research: n_components=2, metric='euclidean'; min_dist=0. We calculated

silhouette scores and distances among calls using each UMAP projection to evaluate

cohesion within the same class and separation between different classes. Silhouette

scores range from -1 to 1, indicating how tightly data points are grouped within a cluster

and how well clusters are separated here; higher positive values mean better separation,

while higher negative values indicate overlap between clusters.

Train-test split

Given the unbalanced sample sizes among individuals, we randomly selected

80 coo calls per individual in each classifier learning iteration. As with the 80 calls in

each individual, we allocated 64 calls to the training data and the remainder (16 calls) to

the test data. This process was repeated 1000 times, resulting in 1000 pairs of training

and test datasets to account for variability and provide a robust estimate of classifier

performance.

Classifier learning

In this study, we tackled two tasks: One was to identify individuals from the

calls of six adult females, and the other was to classify age classes (older = over 20

years old and younger = less than 10 years old). For individual and age-class

classification, we trained random forests (RF) and support vector machines (SVM)

using training datasets and then predicted test datasets 1000 times using *scikit-learn,* a

commonly used Python module (Pedregosa et al., 2011). In each iteration, to estimate

the performance of classifiers, we calculated balanced accuracy, which is defined as the

average of recall in each class and reduced the biases due to the differences in data size

between classes compared to the standard accuracy. By using the prediction based on

the test data, we calculated the accuracy per call defined as follows: the number of times

the predicted label in the test data matched the actual label divided by the number of

times the call was used as the test data.

Hyperparameter tuning

218   Since it is time-consuming to tune the hyperparameters of the classifiers in

each iteration, we explored the best parameters using 5-repeated 5-fold cross-validation

in the grid search method. In other words, we repeated the random selection of 80 calls

from each individual and 5-fold cross-validation 5 times, and then we searched for the

combinations of hyperparameters with the highest average balanced accuracy,

commonly known as the grid search method. We searched for the best combination of

hyperparameters with high average balanced accuracy for RF and SVM in the age-class

classification and individual identification tasks, respectively. The grids explored by RF

and SVM are as follows: for RF i) max_depth (10, 20, 50, None), ii) max_features (sqrt,

log2, None), iii) criterion (gini, entropy, log_loss), iv) min_samples_split (2, 4, 6, 8), v)

min_samples_leaf (1, 2, 5); for SVM i) the kernel type (linear, poly, rbf, sigmoid), ii) C,

which is the regularization parameter (0.1, 1, 10, 100), iii) the kernel coefficient in poly,

rbf and sigmoid (scale, 0.001, 0.01, 0.1, 1), iv) decision_function_shape (ovo, ovr), v)

for poly or sigmoid, coef0 (0.0, 0.1, 0.5, 1), and vi) for poly, degree (1, 2, 3, 4). To

reduce the computational costs, we set n_estimator =100 in the RF. The best

hyperparameters were as follows: In the individual identification task, for RF best

hyperparameter combination was n_estimators=100, max_depth=20,

max_features=sqrt, criterion=entropy, min_samples_split=2, and min_samples_leaf=1;

for SVM C=10, kernel=rbf, gamma=scale, and decision_function_shape=ovo. In the

age-class classification task, for RF the best hyperparameter combination was

n_estimators=100, max_depth=20, max_features=sqrt, criterion=gini,

min_samples_split=2, and min_samples_leaf=2; for SVM C=0.1, kernel=poly,

degree=3, coef0=0.5, gamma=0.001, and decision_function_shape=ovo.

Results

Individual identification

Coo calls were generally clustered by individual in the supervised UMAP,

while the unsupervised UMAP did not produce clear clusters (Fig. 1). This was also

reflected in the results of the silhouette score. The mean silhouette score for all six

individuals of the supervised UMAP (0.60) was much higher compared to the

unsupervised UMAP (0.05) (Fig. 2). The mean silhouette score in each individual was

positive (0.21-0.90) in the supervised UMAP; however, the silhouette scores of Kapa

and Rine were negative in the unsupervised UMAP. The all-average distances within

individuals were smaller than between individuals in the supervised UMAP; on the

other hand, the unsupervised UMAP did not show much difference in the within-

individual and between-individual distances (Fig. 2).

In the individual identification task, the mean balanced accuracy of RF was

81% and that of SVM was 82% (Table 1). The mean accuracy varied across individuals,

with values ranging from 0.68 to 0.94 in RF and from 0.72 to 0.94 in SVM. Both RF

and SVM had the highest accuracy in Sasa and the lowest in Kapa. The main factor

lowering the accuracy for both RF and SVM was the misidentification of Kapa, Rine,

and Taiwu among them.

The distribution of the accuracies of all coo calls was divided into high-

accuracy and low accuracy calls (Fig. 4). When we defined the calls with accuracy

below 1/6 (lower than expected) as low-accuracy calls and over 0.9 as high-accuracy

calls, 55 coo calls were classified as low accuracy (Table 2) and 447 as high accuracy

(Table 3) in both RF and SVM. The proportion of low-accuracy calls in each individual

varied, with the highest for Kapa at 0.15 and the lowest for Sasa at 0.02. On the other

hand, the proportion of high-accuracy calls ranged from 0.54 to 0.89.

Age-class classification

In the supervised UMAP, coo calls were clustered tightly by age-class, while

the unsupervised UMAP showed a looser separation between coo calls of younger and

older individuals (Fig. 5). As with the supervised UMAP, the averages of silhouette

scores were 0.86 for the younger and 0.80 for the older, respectively, which were much

higher than the unsupervised UMAP with 0.18 for the younger and 0.01 for the older

(Fig. 6). Nevertheless, only some calls of the older individuals had negative silhouette

scores in the supervised UMAP. Compared these calls with calls with high silhouette

score, but there was no consistent difference between low silhouette calls and high

silhouette calls. In terms of distances, the mean pairwise distance within age classes

(3.05 for the younger and 3.76 for the older) was smaller than between age classes

(22.66) in the supervised UMAP, whereas no differences were observed within and

between age classes in the unsupervised UMAP (Fig. 6).

In the age classification task, the balanced accuracy for RF was 91% and that

for SVM was 93% (Table 4), thus exceeding 90%. The accuracy for young individuals

exceeded 98% in both RF and SVM, while that for older individuals was less than 87%.

Therefore, misclassification was caused by classifying the older calls as younger ones.

In the age classification task, the accuracy scores of each call were sharply

divided between high and low values (Fig. 8). When we defined the calls with accuracy

below 0.5 (lower than expected) as low accuracy calls, the common low-accuracy calls

between RF and SVM were 23, comprising of only Sazae and Taiko (Table 5). Here

12.3% of the Sazae calls and 9.5% of Taiko calls were low accuracy.

Discussion

Individual identification

In this study, RF and SVM achieved 81% and 82%, respectively, in identifying the individual from coo calls. The previous research on individual identification from contact calls in macaques achieved variable accuracy (0.32-0.92), although the datasets were obtained in laboratory settings (Ceugniet and Izumi, 2004; Fukushima et al., 2015; Trawicki, 2024). Although it is difficult to simply compare accuracy because of the datasets are different, the dataset used by Fukushima et al. whose study achieved the highest accuracy (0.92) and Trawicki more clustered individually in the UMAP latent space than this study (Sainburg et al., 2020). Therefore, the dataset is probably easier to classify than this study dataset. In addition, considering the difficulty of collecting a balanced set of vocalizations of sufficient and well-controlled sound quality from a social group of wild macaques, the performance of our approach seemed not to be inferior to classifiers that relied solely on highly controlled calls in a laboratory setting. In previous studies, the performance of classifiers varied greatly depending on the acoustic features used (Fukushima et al., 2015; Rendall et al., 1998). This means that the way of selecting acoustic features may distort the performance evaluation. Our approach using mel spectrograms would be more stable due to eliminating the variance of selecting specific acoustic features.

The method using mel spectrograms for individual identification has the potential to facilitate density estimation in the wild. Recently, vocal individual identification has received attention for its use in mark-and-recapture methods due to its non-invasive nature and ability to collect data even at night (Longden et al., 2020). While the average accuracy of individual identification in this study was 80%, more

than half of calls achieved over 90% accuracy for any individual with either the RF or

SVM models. Highly accurate discrimination is possible by collecting enough data for

each individual, identifying calls that are easier to discriminate in advance, and focusing

on those calls. Additionally, employing supervised UMAP to pre-check the data

structure, as recommended in a previous study (Arnaud et al., 2023), would be

beneficial, since supervised UMAP can capture structures that unsupervised UMAP

cannot clarify, such as higher silhouette scores and longer inter-class distances (Figs. 1,

2, 5, and 6). Although this study demonstrates the effectiveness of mel spectrograms in

small groups, further research is needed to verify whether similar accuracy can be

achieved with larger numbers of individuals including not only females but also males

with various age-class.

326   In the individual identification task, the accuracy of each call was sharply

divided and skewed into high and the low scores, and low accuracy calls were made by

all individuals. These results might be caused by mislabeling or data sampling bias,

such as the collection of only a small number of specific types of coo calls in an

individual. Otherwise, the results suggest that some of the individuals' calls might be

distinctive. Such differences in individuality might differ depending on the context. For

example, in some primates, the call tends to be less individualistic when the distance

from other individuals is sufficiently close (Leliveld et al., 2011; Mitani et al., 1996). In

addition, the coo call in response tends to resemble the call of the individual whose call

was responded to (Sugiura, 1998). These factors may affect the individuality of each

coo call. Further study of the relationship between individuality and the context of the

calls would help to elucidate which factor affects the difference in individuality.

Age-class classification

This study achieved 91-93% accuracy in the age-class classification task, i.e.

young (<10 yrs) and old (20> yrs) individuals. One possible reason for this high

accuracy is the use of the entire mel spectrogram, which preserves global spectral

patterns rather than relying on a limited set of predefined acoustic features. However,

whether age-class classification based on such representations can be generalized across

a broader range of age groups remains to be examined. Because mel spectrograms

encode the overall distribution and temporal structure of acoustic energy, they may

implicitly reflect complex spectral characteristics. For example, harshness, previously

reported to increase with age in Japanese macaques (Inoue, 1988), is difficult to capture

using specific acoustic features such as pitch or resonance frequency. Thus, the current

results do not demonstrate that the classifiers explicitly relied on harshness or other

specific age-related acoustic features, nor do they allow identification of which acoustic

properties may have implicitly contributed to the classification.

A key limitation of the present study is that the classifiers may have

discriminated age classes based on individual-specific characteristics of a small number

of older females rather than generalizable age-related vocal features. In this study, only

two older females were included, and although we did not observe an obvious

difference in body size between older and younger individuals, we cannot rule out the

possibility that subtle individual differences influenced classification. The limited

sample size of older individuals therefore represents an important constraint when

interpreting the results. Classification accuracy was lower for older individuals than for

younger individuals. One possible explanation is that older individuals may not

consistently produce vocalizations with age-typical acoustic characteristics, and may, in

certain contexts, emit calls that are acoustically similar to those of younger individuals.

This variability could reduce classification accuracy for older individuals. However,

further investigation is required to evaluate whether such tendencies systematically exist

and whether classifiers can robustly capture age-class-specific vocal characteristics.

Taken together, the present findings demonstrate that age-class classification

of wild Japanese macaque vocalizations is feasible without explicitly measuring specific

acoustic features. At the same time, clarifying which acoustic properties underlie this

classification will require future studies incorporating a larger number of older

individuals, the use of independent test data from novel individuals, and analytical

approaches that allow for greater interpretability of model decisions.

Acknowledgements

We appreciate the residents and researchers on Yakushima Island for supporting

our field study, as well as the Yakushima Forest Ecosystem Conservation Center and the

Kagoshima Prefectural Government for permission to perform the research. We also Drs.

Hideki Sugiura and Mariko Suzuki for giving thoughtful comments and Sarah Katherine

Carnes for preprocessing the vocal data. The study was supported by the Japan Society

for the Promotion of Science, Grant-in-Aid for Scientific Research (No. 23H02563 to GH,

No. 24H00774 to IM, and No. 23K25168 and No. 23H05428 to HK), JSPS Core-to-Core

Program, Asia–Africa Science Platforms (JPJSCCB20250006) and the Collaborative

Research Program of Wildlife Research Center, Kyoto University.

Author contributions

Convinced procedures: RK(lead), FK(supporting), HK(equally), IM

(supporting), GH (supporting); Data collection: RK (lead), FK(equally); Software: RK

(lead), HK(supporting); Grant obtained: HK(supporting); IM (supporting); GH (lead);

Supervision; HK(supporting), IM (supporting), GH (lead); Formal analysis: RK(lead),

HK(supporting); Manuscript writing: RK(equally), IM(supporting), HK(lead), GH

(supporting)

Conflict of interests

The authors declare no conflicts of interest.

Figs.

Fig. 1 Projections of all coo calls into two-dimensional space through UMAP using

individual labels (651 calls; each dot = 1 call; different colors indicate different

individuals). a: unsupervised UMAP. b: supervised UMAP.

Fig. 2 Silhouette plots and distances between individuals using projections of UMAPs and individual labels. Top: silhouette plots of unsupervised (a) and supervised (b) UMAP with averages and standard deviations per individual (black line) and in all individuals (red line). Bottom: distance matrix of unsupervised (c) and supervised (d) UMAP.

410 **Table 1 Balanced accuracy of six individual identifications**

Classifier	Mean	std	min	25%	50%	75%	max
RF	0.81	0.04	0.69	0.78	0.81	0.83	0.92
SVM	0.82	0.04	0.71	0.79	0.82	0.84	0.93

Fig. 3 Average confusion matrix for 1000 iterations in identifying six females. a:

RF. b: SVM.

Fig. 4 Frequency of accuracy for coo calls in individual identification. Blue bar and

orange bar represent SVM results and RF results, respectively.

422   **Table 2 Low-and-high accuracy calls in individual identification**

Individual	Number of low-accuracy calls in RF	Number of low-accuracy calls in SVM	Number of low-accuracy calls in common between RF and SVM	Number of calls
Kapa	26	20	17	116
Rine	12	10	9	89
Sasa	6	6	3	138
Sazae	18	16	13	122
Taiko	7	7	4	84
Taiwu	12	9	9	102

425   **Table 3 Low-and-high accuracy calls in individual identification**

Individual	Number of high accuracy calls in RF	Number of high accuracy calls in SVM	Number of high-accuracy calls in common between RF and SVM	Number of calls
Kapa	67	74	63	116
Rine	58	54	49	89
Sasa	126	127	124	138
Sazae	94	94	92	122
Taiko	67	64	60	84
Taiwu	63	72	59	102

Fig. 5 Projections of all coo calls into two-dimensional space through UMAP using

age labels (651 calls; each dot = 1 call; different colors indicate different age classes).

a: Unsupervised UMAP. b: Supervised UMAP

Fig. 6 Silhouette plots and distances between individuals using projections of

UMAPs and age class labels. Top: silhouette plots of unsupervised (a) and

supervised (b) UMAP with averages and standard deviations per age class (black

line) and in all classes (red line). Bottom: distance matrix of unsupervised (c) and

supervised (d) UMAP.

441   **Table 4 Balanced accuracy of age classification**

Classifier	Mean	std	min	25%	50%	75%	max
RF	0.91	0.03	0.78	0.88	0.91	0.93	0.98
SVM	0.93	0.03	0.81	0.91	0.93	0.95	1.00

445 **Fig. 7 The Confusion matrix for classifying age-class between six females. a: RF. b: SVM.**

Fig. 8 Frequency of accuracy for coo calls in age-class classification. Blue bar and

orange bar represent SVM results and RF results, respectively

451   **Table 5 Low-accuracy calls in age classification**

Individual	Number of low-accuracy calls in RF	Number of low-accuracy calls in SVM	Number of low-accuracy calls in common between RF and SVM	Number of calls
Kapa	0	2	0	116
Rine	0	0	0	89
Sasa	1	0	0	138
Sazae	23	16	15	122
Taiko	14	10	8	84
Taiwu	0	1	0	102

Reference

Abitbol, J., P. Abitbol, and B. Abitbol, 1999, Sex hormones and the female voice:

JOURNAL OF VOICE, v. 13, p. 424-446.

Arnaud, V., F. Pellegrino, S. Keenan, X. St-Gelais, N. Mathevon, F. Levréro, and C.

Coupé, 2023, Improving the workflow to crack Small, Unbalanced, Noisy, but

Genuine (SUNG) datasets in bioacoustics: The case of bonobo calls: PLOS

Computational Biology, v. 19, p. e1010325.

Baowaly, M. K., B. C. Sarkar, M. A. A. Walid, M. M. Ahamad, B. C. Singh, E. S.

Alvarado, I. Ashraf, and M. A. Samad, 2024, Deep transfer learning-based bird

species classification using mel spectrogram images: PLOS ONE, v. 19.

Boersma, P. W., David., 2024, Praat: doing phonetics by computer [Computer program].

Version 6.4.12, retrieved 8 May 2024 from   http://www.praat.org/.

Ceugniet, M., and A. Izumi, 2004, Individual vocal differences of the coo call in

Japanese monkeys: Comptes Rendus. Biologies, v. 327.

Dossou, B. F., and Y. K. Gbenou, 2021, FSER: Deep convolutional neural networks for

speech emotion recognition: Proceedings of the IEEE/CVF International

Conference on Computer Vision, p. 3533-3538.

Ey, E., D. Pfefferle, and J. Fischer, 2007, Do age- and sex-related variations reliably

reflect body size in non-human primate vocalizations? A review: Primates, v. 48,

p. 253-267.

Feinberg, D. R., B. C. Jones, A. C. Little, D. M. Burt, and D. I. Perrett, 2005,

Manipulations of fundamental and formant frequencies influence the

attractiveness of human male voices: Animal Behaviour, v. 69.

Ferrand, C. T., 2002, Harmonics-to-Noise Ratio: An Index of Vocal Aging: Journal of

Voice, v. 16.

Fukushima, M., A. M. Doyle, M. P. Mullarkey, M. Mishkin, and B. B. Averbeck, 2015,

481   Distributed acoustic cues for caller identity in macaque vocalization: Royal

482   Society Open Science, v. 2, p. 150432.

483   Furuyama, T., Kobayasi, K. & Riquimaroux, H, 2016, Role of vocal tract characteristics

484   in individual discrimination by Japanese macaques (*Macaca fuscata*): Scientific

485   Repports, v 6, p. 32042.

486   Green S, 1975, Variation of vocal pattern with social situation in the Japanese monkey

487   (Macaca fuscata): a field study. In: Rosenblum LA (ed) Primate behavior, vol 4.

488   Academic Press, New York, pp 1–102

489   Harnsberger, J. D., R. Shrivastav, W. S. Brown, H. Rothman, and H. Hollien, 2008,

490   Speaking Rate and Fundamental Frequency as Speech Cues to Perceived Age:

491   Journal of Voice, v. 22.

492   Heinrich, R., L. Rauch, B. Sick, and C. Scholz, 2025, AudioProtoPNet: An interpretable

493   deep learning model for bird sound classification: Ecological Informatics, v. 87,

494   p. 103081.

495   Hindle, A. G., M. Horning, J.-A. E. Mellish, and J. M. Lawler, 2009, Diving into old

496   age: muscular senescence in a large-bodied, long-lived mammal, the Weddell

497   seal (*Leptonychotes weddellii*): Journal of Experimental Biology, v. 212.

498   Huntley, R., H. Hollien, and T. Shipp, 1987, Influences of listener characteristics on

499   perceived age estimations: Journal of Voice, v. 1.

500   Inoue, M., 1988, Age Gradations in Vocalization and Body Weight in Japanese Monkeys

501   (*Macaca fuscata*): Folia Primatologica, v. 51.

502   Klofstad, C. A., R. C. Anderson, and S. Nowicki, 2015, Perceptions of Competence,

503   Strength, and Age Influence Voters to Select Leaders with Lower-Pitched

504   Voices: PLOS ONE, v. 10.

Koda, H., and H. Sugiura, 2010, The ecological design of the affiliative vocal

communication style in wild Japanese macaques: behavioral adjustments to

social contexts and environments, The Japanese Macaques, Springer, p. 165-190.

Lakdari, M. W., A. H. Ahmad, S. Sethi, G. A. Bohn, and D. J. Clink, 2024, Mel-

frequency cepstral coefficients outperform embeddings from pre-trained

convolutional neural networks under noisy conditions for discrimination tasks of

individual gibbons: Ecological Informatics, v. 80.

Leliveld, L. M. C., M. Scheumann, and E. Zimmermann, 2011, Acoustic correlates of

individuality in the vocal repertoire of a nocturnal primate (*Microcebus*

murinus): The Journal of the Acoustical Society of America, v. 129.

Lemasson, A., E. Gandon, and M. Hausberger, 2010, Attention to elders' voice in non-

human primates: Biology Letters, v. 6.

Longden, E. G., S. H. Elwen, B. McGovern, B. S. James, C. B. Embling, and T. Gridley,

2020, Mark–recapture of individually distinctive calls—a case study with

signature whistles of bottlenose dolphins (*Tursiops truncatus*): Journal of

Mammalogy, v. 101.

Manrique, H. M., and J. Call, 2015, Age-dependent cognitive inflexibility in great apes:

Animal Behaviour, v. 102.

Marler, P., 1956, The voice of the chaffinch and its function as a language: Ibis.

Marler, P., and L. Hobbett, 1975, Individuality in a Long-Range Vocalization of Wild

Chimpanzees: Zeitschrift für Tierpsychologie, v. 38.

McFee, B., C. Raffel, D. Liang, D. Ellis, M. McVicar, E. Battenberg, and O. Nieto,

2015, librosa: Audio and Music Signal Analysis in Python: SciPy.

McInnes, L., J. Healy, and J. Melville, 2018, Umap: Uniform manifold approximation

and projection for dimension reduction: arXiv preprint arXiv:1802.03426.

Mitani, J. C., J. Gros-Louis, and J. M. Macedonia, 1996, Selection for acoustic

individuality within the vocal repertoire of wild chimpanzees: International

Journal of Primatology, v. 17, p. 569-583.

Mitani, M., 1986, Voiceprint identification and its application to sociological studies of

wild Japanese monkeys (*Macaca fuscata yakui*): Primates, v. 27, p. 397-412.

Nanni, L., S. Brahnam, A. Lumini, and G. Maguolo, 2020, Animal sound classification

using dissimilarity spaces: Applied Sciences, v. 10, p. 8578.

Oyakawa, C., H. Koda, and H. Sugiura, 2007, Acoustic features contributing to the

individuality of wild agile gibbon (*Hylobates agilis agilis*) songs: American

Journal of Primatology, v. 69.

Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel,

P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau,

M. Brucher, M. Perrot, and E. Duchesnay, 2011, Scikit-learn: Machine Learning

in Python: JOURNAL OF MACHINE LEARNING RESEARCH, v. 12, p.

2825-2830.

Pontes, P., R. Yamasaki, and M. Behlau, 2006, Morphological and functional aspects of

the senile larynx: FOLIA PHONIATRICA ET LOGOPAEDICA, v. 58, p. 151-

158.

Rendall, D., M. J. Owren, and P. S. Rodman, 1998, The role of vocal tract filtering in

identity cueing in rhesus monkey (*Macaca mulatta*) vocalizations: The Journal

of the Acoustical Society of America, v. 103.

Ryan, W. J., and K. W. Burk, 1974, Perceptual and acoustic correlates of aging in the

speech of males: Journal of Communication Disorders, v. 7.

Sainburg, T., M. Thielk, and T. Q. Gentner, 2020, Finding, visualizing, and quantifying

latent structure across diverse animal vocal repertoires: PLOS Computational

Biology, v. 16, p. e1008228.

Stathopoulos, E. T., J. E. Huber, and J. E. Sussman, 2011, Changes in Acoustic

Characteristics of the Voice Across the Life Span: Measures From Individuals 4–

93 Years of Age: Journal of Speech, Language, and Hearing Research, v. 54.

Stevens, S. S., J. Volkmann, and E. B. Newman, 1937, A Scale for the Measurement of

the Psychological Magnitude Pitch: The Journal of the Acoustical Society of

America, v. 8.

Sugiura, H., 1998, Matching of acoustic features during the vocal exchange of coo calls

by Japanese macaques: Animal Behaviour, v. 55, p. 673-687.

Thomas, M., F. H. Jensen, B. Averly, V. Demartsev, M. B. Manser, T. Sainburg, M. A.

Roch, and A. Strandburg-Peshkin, 2022, A practical guide for generating

unsupervised, spectrogram-based latent space representations of animal

vocalizations: Journal of Animal Ecology, v. 91, p. 1567-1581.

Tigue, C. C., D. J. Borak, J. J. M. O'Connor, C. Schandl, and D. R. Feinberg, 2012,

Voice pitch influences voting behavior: Evolution and Human Behavior, v. 33.

Trawicki, M. B., 2024, Automatic gender recognition and speaker identification of

Rhesus Macaques (*Macaca mulatta*) using hidden Markov models (HMMs):

International Journal of Speech Technology, v. 27, p. 179-186.

Walsh, S. L., S. Engesser, S. W. Townsend, and A. R. Ridley, 2023, Multi-level

combinatoriality in magpie non-song vocalizations: Journal of the Royal Society

Interface, v. 20, p. 20220679.

Winter, P., D. Ploog, and J. Latta, 1966, Vocal repertoire of the squirrel monkey (*Saimiri*

sciureus), its analysis and significance: Experimental Brain Research, v. 1, p.

359-384.

Xie, B., V. Daunay, T. C. Petersen, and E. F. Briefer, 2024, Vocal repertoire and

individuality in the plains zebra (*Equus quagga*): Royal Society Open Science,

v. 11.

Yamagiwa, J., and D. A. Hill, 1998, Intraspecific variation in the social organization of

Japanese macaques: Past and present scope of field studies in natural habitats:

Primates, v. 39, p. 257-273.

Zipple, M. N., S. Peters, W. A. Searcy, and S. Nowicki, 2020, Sounds of senescence:

male swamp sparrows respond less aggressively to the songs of older

individuals: Behavioral Ecology, v. 31.

a

b