Optimising 7T-fMRI for imaging regions of magnetic susceptibility

Saskia L. Frisby[1], Marta M. Correia[1], Minghao Zhang[2], Christopher T. Rodgers[2], Timothy T.

Rogers[3], Matthew A. Lambon Ralph[1], and Ajay D. Halai[1]

[1]MRC Cognition and Brain Sciences Unit, 15 Chaucer Road, Cambridge CB2 7EF

[2]Wolfson Brain Imaging Centre, Cambridge Biomedical Campus, Cambridge CB2 0QQ

[3]Department of Psychology, University of Wisconsin-Madison, 1202 W Johnson Street,

Madison 53706

Correspondence to:

Dr. Saskia L. Frisby

saskia.frisby@mrc-cbu.cam.ac.uk

+44 (0)1223 769703

and

Dr. Ajay D. Halai

ajay.halai@mrc-cbu.cam.ac.uk

+44 (0)1223 767655

MRC Cognition & Brain Sciences Unit

University of Cambridge

15 Chaucer Road

Cambridge

CB2 7EF

Abstract

The temporal signal-to-noise ratio (tSNR) of functional magnetic resonance imaging (fMRI) is particularly poor in ventral anterior temporal and orbitofrontal regions because of magnetic field inhomogeneity, a problem that is exacerbated at higher field strengths. In this 7T-fMRI study we compared three methods of improving sensitivity in these areas: parallel transmit, which uses multiple transmit elements, controlled independently, to homogenise the flip angle experienced by the tissue; multi-echo, which entails collection of multiple volumes at different echo times following a single radiofrequency pulse; and multiband, in which multiple slices are acquired simultaneously. We found that parallel transmit and multi-echo increased the magnitude of the BOLD signal change, but only multi-echo increased BOLD magnitude in areas prone to susceptibility artefacts. Multiband and denoising of multi-echo data with independent components analysis (ICA) both improved precision of GLM fit. Exploratory results suggested that multi-echo and ICA denoising can both benefit multivariate analyses. In conclusion, a multi-echo, multiband sequence improved fMRI quality in areas prone to susceptibility artefacts while maintaining sensitivity across the whole brain. We recommend this approach for studies investigating the functional roles of ventral temporal and orbitofrontal regions with 7T fMRI.

Keywords: 7T-fMRI, parallel transmit, multi-echo, multiband, temporal lobe, orbitofrontal cortex

1. Introduction

The temporal signal-to-noise ratio (tSNR) of functional magnetic resonance imaging (fMRI) varies across the brain. The ventral anterior temporal cortex and orbitofrontal cortex, for example, are located next to air-filled sinuses and so are affected by magnetic field inhomogeneity that causes signal dropout and geometric distortions (Devlin et al., 2000; Halai et al., 2014, 2015, 2024). This makes it challenging to use fMRI to investigate the roles that these regions may play in a myriad of cognitive processes – including vision (Devereux et al., 2018), language (Borghesani et al., 2016), multimodal semantic cognition (Lambon Ralph et al., 2017), emotion (Fernandez et al., 2017), social cognition (Binney et al., 2016; Zahn et al., 2007), theory of mind (DuPre et al., 2016), and executive function (Duncan, 2010). Signal dropout and geometric distortions are more severe at higher field strengths, implying that it may be especially difficult to measure task-related activity in susceptible regions with ultra-high-field fMRI (e.g., 7T-fMRI). However, 7T-fMRI also has many advantages in regions unaffected by magnetic susceptibility artefacts: 7T-fMRI offers improved tSNR relative to 3T-fMRI (Morris et al., 2019), which can be used to reduce voxel size and enable applications such as laminar fMRI (Koopmans et al., 2011) or to reduce acquisition times and enable shorter scan times for special populations such as patients with neurodegenerative diseases (Cope et al., 2023). 7T-fMRI also benefits from improved spatial specificity relative to 3T-fMRI because the signal from cortical microvasculature is enhanced while the signal from large veins is reduced (Marques & Norris, 2018). Improving signal homogeneity would allow researchers studying the whole brain (or focusing on regions prone to susceptibility artefacts) to take full advantage of 7T-fMRI; therefore, in this study we compared three methods of doing so.

One possible method for counteracting signal dropout is parallel transmit (pTx), which uses multiple transmit elements, controlled independently, to homogenise the flip angle pattern experienced by the tissue (Deniz et al., 2019). A recent study used pTx to counteract signal dropout in ventral anterior temporal regions for echo-planar imaging (EPI) 7T fMRI (Ding et al., 2022). pTx improved tSNR across the brain compared to a standard sequence, particularly in the temporal lobes. However, there was no improvement in functional contrast during a semantic association task that is known to recruit the anterior temporal lobes in 3T-fMRI studies (Jung et al., 2017).

A second method of recovering signal in these regions is multi-echo (ME) imaging

(Kundu et al., 2017; Poser et al., 2006; Posse, 2012). T_2^* is known to vary across the brain

(Hagberg et al., 2002); in areas prone to magnetic susceptibility artefacts, T_2^* is particularly

short due to increased intravoxel dephasing. A single echo provides sensitivity to a narrow

range of T_2^* values; the echo time (TE) is therefore selected to provide the best compromise

of sensitivity to T_2^* across the whole brain. Combining data from multiple echoes increases

the range of T_2^* that can be imaged with high fidelity. ME has been shown to improve

functional contrast (Poser & Norris, 2009) and spatial specificity (Boyacioğlu et al., 2015) at

7T and 3T (Fernandez et al., 2017; Halai et al., 2024; Kirilina et al., 2016; Lynch et al., 2020).

Having multiple echoes also facilitates the separation of signal and noise because signals

decay in a well-characterised way across echoes, whereas noise does not. This principle

underpins multi-echo independent components analysis (ME-ICA), via which ICA

components that are TE-independent, and thus are likely to be noise rather than blood-

oxygen-level-dependent (BOLD) signal, can be removed (Dipasquale et al., 2017; Kundu et

al., 2011, 2013, 2015, 2017). This method may enhance signal detection in areas prone to

susceptibility artefacts on top of the advantage offered by ME alone (e.g. Lombardo et al.,

2016). ME sequences have some potential disadvantages. For example, multi-echo can

lengthen repetition time (TR). In-plane acceleration is frequently needed to achieve a

sufficiently short first TE, which reduces tSNR (Yun & Shah, 2017). In turn, a short first TE,

combined with hardware constraints, often limits the minimum voxel size (Koopmans et al.,

2011). Critically, in many previous studies examining the benefits of ME sequences

compared to single-echo (SE), the "single-echo" data were extracted from the ME dataset

(Amemiya et al., 2019; Bhavsar et al., 2014; Caballero-Gaudes et al., 2019; Cohen et al.,

2017, 2017, 2018; Dipasquale et al., 2017; Evans et al., 2015; Fernandez et al., 2017; Gilmore

et al., 2022; Kovářová et al., 2022). This means that the "single-echo" data will inherit

suboptimal parameters that are ME-specific, making the comparison unfair.

A third strategy for improving acquisition is multiband (MB) imaging, also known as

simultaneous multi-slice, in which multiple slices are acquired simultaneously (Barth et al.,

2016; Moeller et al., 2010; Setsompop et al., 2012). Typically MB has been applied to

increase temporal resolution – multiple studies have shown that MB can reduce noise

aliasing, increase statistical power and counteract the increase in TR associated with ME

(Feinberg et al., 2010; Griffanti et al., 2014; Halai et al., 2024; Puckett et al., 2018; Smith et

al., 2013). Benefits specific to task fMRI have been less clear (Demetriou et al., 2018; Todd et al., 2016) since some noise features are unlikely to be correlated with task regressors and can be removed with high-pass filtering. Other possible disadvantages of MB include a reduction in tSNR due to increases in g-factor effects (Demetriou et al., 2018; Risk et al., 2021; Setsompop et al., 2012) and leakage of signal into the simultaneously-excited slices (Todd et al., 2016).

There is therefore a need to evaluate 7T-fMRI sequences to determine which parameters are important for counteracting signal dropout and improving sensitivity in regions prone to susceptibility artefacts without compromising signal quality elsewhere. We tested five possible sequences. These consisted of pTx, plus a 2 x 2 factorial design varying number of echoes and multiband factor: single-echo single band (SESB), single-echo multiband (SEMB), multi-echo single band (MESB), and multi-echo multiband (MEMB). We used a semantic judgment task that is known to evoke activity across the semantic network, including areas severely affected and those relatively unaffected by susceptibility artefacts (Jung et al., 2017).

We had two univariate effects of interest – activation magnitude and activation precision (Halai et al., 2024). Activation magnitude is the magnitude of the BOLD signal change, operationalised as the 1^{st}-level beta values extracted from each voxel. We hypothesised that both pTx and ME would recover signal and hence increase activation magnitude relative to the SESB sequence (which we used as a baseline). Activation precision is the reliability of the BOLD signal change, analogous to the functional contrast-to-noise ratio (fCNR) and operationalised as the 1^{st}-level t-values extracted from each voxel. We hypothesised that MB sequences, with greater effective degrees of freedom, would increase activation precision relative to single band (SB) sequences. We also had two secondary hypotheses: first, since ME-ICA denoising removes TE-independent noise, greater activation precision would be observed in multi-echo denoised data (MEdn) relative to ME data without denoising; second, that any MB advantage would be due to the increase in the number of volumes.

Our study focused primarily on univariate effects. However, multivariate analysis techniques, which exploit variance and covariance between voxels and can accommodate participant-specific differences in activation patterns (Coutanche, 2013; Davis et al., 2014; Davis & Poldrack, 2013) are rapidly gaining popularity. These methods frequently rely on the

assumption of good-quality signal across the whole brain (Frisby et al., 2023) and so we

conducted an exploratory analysis, following the method of Haxby et al. (2001), to decode

task condition from the data acquired with each sequence. Finally, we tested for the

presence of slice leakage artefacts in our MB data.

To summarise, our study aimed to compare pTx, ME and MB as methods for

improving sensitivity in ventral temporal and orbitofrontal regions while maintaining image

quality across the rest of the brain.

2.   Methods

2.1. Participants

20 healthy native speakers of British English (age range 18-50, mean age 33.45 years, 12

female, 8 male) participated in the study. All were right-handed, had normal or corrected-

to-normal vision, and had no neurological or sensory disorders. All participants gave written

informed consent. The research was approved by a local National Health Service (NHS)

ethics committee (04/Q105/66).

2.2. Stimuli and task

All participants performed a semantic association task and a visual pattern matching task

(hereafter called the "control task") adapted from a previous study (Jung et al., 2017). Each

stimulus consisted of three pictures presented simultaneously (Figure 1). Some pictures

were line drawings taken from the Pyramids and Palm Trees Test (Howard & Patterson,

1992) and some were colour cartoons or photographs taken from the Camel and Cactus Test

(Bozeat et al., 2000). In the semantic task, participants were instructed to indicate which of

the two pictures at the bottom of the screen had the closest semantic relationship to the

picture at the top (hereafter the "probe picture"). In the control task, participants were

instructed to indicate which of two scrambled pictures (generated from the pictures used in

the semantic task) at the bottom of the screen was identical to a scrambled probe. There

were 248 unique picture triplets, so some stimuli were repeated between runs, but no

stimulus was repeated within a run. E-Prime software (Psychology Software Tools Inc.,

Pittsburgh, USA) was used to display stimuli and record responses. Stimuli were rear-

projected onto a screen at the back of the MRI scanner, and observers viewed stimuli

through a mirror mounted to the head coil directly above the eyes.

Figure 1: One block of the semantic association task (left of the arrow) and one block of the

control task (right of the arrow). Each trial consisted of a fixation cross (500 ms) followed by

three pictures presented simultaneously. Participants had to select which of the two

pictures at the bottom was more semantically similar to (in the semantic task) or visually

identical to (in the control task) the picture at the top (the probe picture). Each block lasted

16 s in total. Reprinted with permission from (Halai et al., 2024).

The study had a block design with three types of block: semantic, control and rest.

Each task block consisted of four trials. Each trial consisted of a fixation cross presented for

500 ms followed by a stimulus presented for 3500 ms. Each rest block consisted of a fixation

cross presented for 16 s. Each run began 16 s after the start of the MR sequence and then

consisted of 30 blocks presented in the order: semantic, control, semantic, control, rest.

There were five runs per participant, collected in a single session. Accuracy and reaction

time were measured for each trial. Since reaction time is not normally distributed, both

metrics were compared across tasks using Wilcoxon's signed-rank tests and across

sequences using Friedman's nonparametric ANOVA.

2.3. Image acquisition

All images were acquired on a whole-body MAGNETOM Terra 7T MRI (Siemens

Healthcare, Germany). An 8Tx32Rx head coil (Nova Medical, USA) was used to run all

structural and functional imaging sequences.

An MP2RAGE anatomical scan was acquired with the following parameters: 224

sagittal slices (interleaved acquisition), FOV 240 x 225.12 x 240 mm^3, voxel size 0.75 mm

isotropic, TR 4300 ms, inversion times 840 and 2370 ms, TE 1.99 ms, nominal flip angles 5°

and 6°, GRAPPA acceleration factor 3, and duration 8 minutes 50 seconds.

Next, manual B_0-shimming was performed over the volume to be used for EPI. The

aim was to reduce the water linewidth, defined as full width of the spectrum at half height,

to below 40 Hz. However, for some participants the adjustments proved time-consuming

and so adjustment time was capped at 30 minutes from the start of scanning after which

the best shim parameters were adopted. The actual average water linewidth was 42 Hz

(standard deviation = 9 Hz; missing data for 4 participants). Then a dummy pTx-EPI scan of

one volume was acquired to trigger the acquisition of subject-specific B_0 and per-channel B_1^+

field maps. The brain was divided into 5 slabs along the slice direction and slab-specific 2-

spoke pTx excitation pulses were designed offline. Variable-rate selective excitation (VERSE;

Hargreaves et al., 2004) was applied to reduce specific absorption rate (SAR) for the pTx

sequence only.

There were five functional runs of EPI, one run of each sequence. Key parameters

are given in Table 1. The order of sequences was counterbalanced across participants. The

following parameters were held constant across sequences: 48 axial slices (interleaved

acquisition), FOV 210 x 210 x 210 mm^3 to cover the whole brain in most participants (visual

inspection ensured that the ventral anterior temporal lobe was included in the FOV for all

participants and the FOV was tilted up at the nose to avoid ghosting of the eyes into the

temporal lobe), voxel size 2.5 mm isotropic (no gap), and A-P phase encoding direction.

After each run, 5 further volumes were acquired with the phase encoding direction changed

to P-A to facilitate distortion correction during preprocessing.

	SESB	SEMB	MESB	MEMB	pTx
No. echoes	1	1	3	3	1
Multiband factor	1	2	1	2	1
TR (ms)	3020	1510	3020	1510	3000
TE_1 (ms)	25.00	25.00	11.80	11.80	25.00
TE_2 (ms)	-	-	27.05	27.05	-
TE_3 (ms)	-	-	42.30	42.30	-
iPAT type	Off	Off	GRAPPA	GRAPPA	GRAPPA
iPAT factor	Off	Off	3	3	2
Phase partial Fourier	7/8	7/8	7/8	7/8	Off

Nominal flip angle (°)	50	50	78	63	40
Bandwidth (Hz/Px)	2204	2204	2204	2204	1804
Number of volumes acquired	171	340	171	340	172
Number of pulses	1	1	1	1	5 (VERSE 2-spoke pTx pulses)

Table 1: parameters of each sequence.

2.4. Data analysis

All analysis code is available at https://github.com/slfrisby/7TOptimisation/.

2.4.1. Preprocessing

For ease of data sharing we converted all DICOMs to BIDS format (Gorgolewski et al.,

2016) using *heudiconv* v1.0.0 (Halchenko et al., 2024).

Standard reproducible preprocessing pipelines designed for 3T-fMRI, such as

fMRIprep (Esteban et al., 2019; Gorgolewski et al., 2011; Markiewicz et al., 2024) perform

poorly on 7T-fMRI EPI data. Therefore, the analysis pipeline was split into stages using

different software packages.

Since it is notoriously difficult to perform a good-quality brain extraction on

MP2RAGE data (because of salt-and-pepper noise in the background and cavities), two

pipelines were used. The MP2RAGE T1w (combined) image first had its background noise

removed using O'Brien regularisation (O'Brien et al., 2014) and was then submitted to the

CAT12 pipeline for segmentation (Gaser et al., 2023; in SPM12;

https://www.fil.ion.ucl.ac.uk/spm/). The bias- and global-intensity corrected T1w image

produced was provided to as input to the anatomy pipeline in *fMRIPrep* 21.0.1. TheT1w

image was skull-stripped with a Nipype implementation of the *antsBrainExtraction.sh*

workflow (ANTs 2.3.3; Avants et al., 2009, 2011; https://github.com/ANTsX/ANTs/). Volume-

based spatial normalisation to standard space (MNI152NLin2009cAsym) was performed

through nonlinear registration with *antsRegistration.sh* (ANTs).

Functional preprocessing was performed using in-house code, composed of

functions from AFNI (v.18.3.03; Cox, 1996; Cox & Hyde, 1997; https://afni.nimh.nih.gov/),

FSL (v.5.0; Andersson et al., 2003; Jenkinson et al., 2012; Smith, 2002; Smith et al., 2004;

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/), tedana (v.23.0.1; DuPre et al., 2021; Kundu et al.,

2011, 2013; The tedana Community et al., 2023;

https://tedana.readthedocs.io/en/stable/index.html) and ANTS (v. 2.2.0; Avants et al., 2009,

2011; https://github.com/ANTsX/ANTs/). EPIs were despiked using *3dDespike* (AFNI), slice

timing was corrected to the middle slice using *3dTshift* (AFNI), motion was corrected with

3dvolreg and *3dAllineate* (AFNI) using the first volume of each run as a reference (for ME

datasets, the TE_1 image was aligned and the resulting transform was applied to the TE_2 and

TE_3 images), and skull-stripped using *BET* (FSL) to create a participant-specific brain mask.

For ME datasets only, *tedana* was used to create two timeseries – one in which the

echoes were optimally combined based on T_2^* weighting (Posse, 2012), and one in which

the T_2^* optimally-combined data were denoised using ICA. *tedana* conducts denoising by

decomposing data using PCA and ICA, classifying components according to whether the

signal scales linearly with TE (as the BOLD signal does), and reconstructing the data using

only BOLD-like components. The brain mask created with *BET* was used as the mask for this

stage.

Next, for all datasets, unwarping was conducted using *topup* and *applytopup* (FSL).

Field displacement maps were calculated using ten volumes (five with A-P phase encoding

direction, extracted from the start of each functional run, and five with P-A phase encoding

direction, collected separately after each run) and the resulting correction was applied to all

images. Finally, the mean EPI for each run was coregistered to the skull-stripped native

structural image using a rigid-body registration with *AntsRegistrationSyN.sh* (ANTS). EPIs

were then transformed into standard space (MNI152NLin2009cAsym) by combining the

transforms from native EPI to native T1 and the transforms from native T1 to standard space

and applying those transforms to the EPIs using *antsApplyTransforms* (ANTS). Images were

smoothed with a 6 mm FWHM Gaussian filter in SPM12 (https://www.fil.ion.ucl.ac.uk/spm/)

for GLM analysis.

A separate functional preprocessing pipeline was used to create images for slice

leakage analysis. This pipeline differed from the main pipeline in the following ways.

Despiking was omitted to avoid the removal of noise. For ME datasets only, rather than run

the full *tedana* workflow, we conducted optimal combination of data from multiple echoes

(but no denoising) using the *t2smap* command using all voxels in the volume (i.e. no brain

mask). All coregistration steps were omitted (the images remained in native EPI space) and
no smoothing was applied.

2.4.2. 1st-level (within-participant) GLM

Data were analysed using the general linear model (GLM) approach implemented in
SPM12 in MATLAB r2019a. We had 5 timeseries of primary interest – standard single-echo
single band (SESB), parallel transmit (pTx), single-echo multiband (SEMB), multi-echo single
band (MESB) and multi-echo multiband (MEMB). For the latter two sequences, data from all
echoes were optimally combined but were not ICA-denoised. We also generated two
timeseries with ME-ICA denoising - multi-echo single band with ICA denoising (MESBdn) and
multi-echo multiband with ICA denoising (MEMBdn) - and two downsampled MB timeseries
created by extracting odd-numbered volumes to match the number of volumes in the single
band timeseries - odd-numbered volumes of single-echo multiband (SEMBodd) and odd-
volumes of multi-echo multiband (MEMBodd).

At the individual subject level, each block of the semantic and control task was
modelled as a boxcar function (resting blocks were modelled implicitly) and these boxcar
functions were subsequently convolved with SPM's difference of gammas haemodynamic
response function. The six motion parameters extracted during preprocessing were used as
regressors of no interest. The micro-time resolution was set as the number of slices (n = 48),
the micro-time onset was set as the reference slice for slice-time correction (n = 24), and the
high-pass filter cutoff was 128 seconds. The same MNI template used during preprocessing
was used as a mask for the analysis. The parameter estimation method was restricted
maximum likelihood estimation (ReML) and serial correlations were accounted for using an
autoregressive AR(1) model during estimation. For univariate analyses the contrast of
interest was greater activation for the semantic task than the control task (S > C). For
exploratory multivariate pattern analysis (MVPA) each block (12 semantic and 12 control)
was modelled individually in order to obtain one beta image per block.

Finally, for the slice leakage analysis, the modelling was rerun on the minimally-
preprocessed timeseries without any brain mask. We obtained both univariate contrasts of
interest (S > C) and beta images per block (for MVPA).

2.4.3. 2nd-level (across-participant) GLM

2.4.3.1. Region-of-interest (ROI) analysis

Regions of interest (ROIs) were defined based on a large-scale distortion-corrected 3T-fMRI study of the semantic network (Humphreys et al., 2015). For each comparison, ROIs were analysed only if they overlapped by at least one voxel with the whole-brain contrast of interest (S > C) summed over all sequences in the comparison.

2.4.3.1.1. Univariate analysis

Activation magnitude (1st-level beta values) and activation precision (1st-level t-values) were extracted from each ROI using a publicly-available script, *roi_extract.m* (https://github.com/MRC-CBU/riksneurotools/blob/master/Util/roi_extract.m). There were two planned t-tests for activation magnitude (pTx > SESB, ME > SE) and four planned t-tests for activation precision (MB > SB, MEdn > ME, MBodd > SB, SB > MBodd). For each planned t-test, results were Bonferroni-corrected for the number of ROIs included.

2.4.3.1.2. Exploratory multivariate pattern analysis (MVPA)

The input to this analysis was the activation magnitude values extracted from each block individually (12 semantic and 12 control). For each block and each ROI, a vector of beta values within that ROI was created. The cosine dissimilarity between every possible pair of blocks was calculated. MVPA performance was operationalised as the mean between-task dissimilarity minus the mean within-task dissimilarity and paired t-tests were used to compare the metric across sequences (all planned t-tests described in the univariate ROI analysis were conducted; Haxby et al., 2001).

2.4.3.2. Whole-brain analysis

All the above contrasts were assessed at the whole-brain level using t-tests (for comparing pTx and SESB) or using random-effects ANOVAs with one-sample t-tests on the summary statistic (for the three factorial designs: varying echo and band (SESB, SEMB, MESB and MEMB); varying denoising and band (MESBdn, MEMBdn, MESB and MEMB); and varying echo and downsampled band (SEMBodd, MEMBodd, SEMB and MEMB)). The ANOVAs were conducted using a publicly-available script, *batch_spm_anova.m* (https://github.com/MRC-CBU/riksneurotools/blob/master/SPM/batch_spm_anova.m). The group t-maps were assessed for significance by using a voxel-height threshold of p < 0.001

350   to define clusters and then a cluster-defining family-wise-error corrected threshold of p <

351   0.05 for statistical inference.

353   2.4.3.3.   Slice leakage analysis

The group-level, whole-brain contrast of interest (S > C) in standard space was

inspected and the coordinates of the peak t-values of the top 5 clusters were identified.

These coordinates were back-projected to obtain 5 sets of corresponding coordinates in

each participant's native EPI space (hereafter "seeds", labelled A). Next, voxels to which

signal might be warped were identified as artefact locations based on phase shift (FOV/2,

labelled B) and, in the MEMB data, GRAPPA (labelled Ag) and phase shift plus GRAPPA

(labelled Bg). A spherical ROI, 4 voxels in radius, was defined around each seed location and

possible artefact location using a modified version of the scripts developed for McNabb et

al. (2020; https://github.com/DrMichaelLindner/MAP4SL/; our version available at

https://github.com/slfrisby/7TOptimisation/).

Both univariate (McNabb et al., 2020) and multivariate (Halai et al., 2024) slice

leakage tests were conducted. Activation magnitude was extracted from minimally-

preprocessed data, and, for the multivariate analysis, the difference between mean within-

task similarity and mean between-task similarity was calculated as in the ROI analysis.

Paired t-tests were conducted for each seed and artefact location between each sequence

of interest (SEMB and MEMB) and a control sequence. For artefact locations based on phase

shift (B), the control sequence was the corresponding SB sequence (SESB for SEMB and

MESB for MEMB). For artefact locations based on GRAPPA, the control sequence was the

corresponding SE sequence, because SE sequences were collected without GRAPPA (SEMB

for MEMB). Results were Bonferroni-corrected for the number of peaks (n = 5) and the

number of possible artefact regions (n = 1 for SEMB and n = 3 for MEMB).

3. Results

3.1. Excluded participants

Two participants were excluded because of excessive head motion (this was defined

by calculating, for each participant, the percentage of volumes per run with absolute

translation values of over 2 mm or absolute rotation values over 1°, averaging these

percentages over runs, and excluding any participant whose mean percentage was greater

than 2 standard deviations above the mean percentage across participants). One participant was excluded because of technical problems during data acquisition which meant that the pTx run failed. All subsequent analyses were conducted on the remaining 17 participants.

3.2. Behavioural results

The 17 participants had good performance on both tasks and, importantly, there were no significant differences between the 5 sequences in terms of accuracy (Friedman's χ^2 = 5.26, p = 0.26) or reaction time (Friedman's χ^2 = 7.20, p = 0.16). The semantic and control tasks did not differ reliably in accuracy (Wilcoxon's z = 0; p = 0.06) or reaction time (Wilcoxon's z = 7; p = 1.00).

3.3. Region-of-interest analysis

ROIs that overlapped with the whole-brain contrast of interest (S > C) for at least one of the five sequences are shown in Figure 2.

Figure 2: Regions of interest, taken from a meta-analysis of semantic tasks by (Humphreys et al., 2015). All spheres are 8 mm in radius. Regions of interest are overlaid on the MNI152NLin2009cAsym template. LTP = left temporal pole; LvATL = left ventral anterior temporal lobe; RITG = right inferior temporal gyrus; LFP = left frontal pole; LmMTG = left medial middle temporal gyrus; LpMTG = left posterior middle temporal gyrus; LIFGpt = left inferior temporal gyrus pars triangularis.

	LTP	LvATL	RITG	LFP	LmMTG	LpMTG	LIFGpt
Activation magnitude							
pTx > SESB	-	0.2488	0.0240*	-	-	0.0042**	0.4114
ME > SE	0.0502	0.0001**	0.0333*	0.2951	0.0092*	0.0003*	0.4155
Activation precision							
MB > SB	0.4569	0.0007**	0.0019**	0.0989	0.2187	0.0266*	0.0016**
MEdn > ME	0.3117	<0.0001**	<0.0001**	0.0014**	0.1363	<0.0001**	<0.0001**
MBodd > SB	0.4225	0.1057	0.0481*	-	0.5697	0.1879	0.3300

MVPA							
pTx > SESB	-	0.8569	0.3280	-	-	0.1529	0.9179
ME > SE	0.0096*	0.0063**	0.1612	0.0655	0.0465*	0.0024**	0.2508
MB > SB	0.2455	0.1417	0.0667	0.3021	0.3949	0.2693	0.5207
MEdn > ME	0.0027**	<0.0001**	<0.0001**	0.0023**	0.0001**	0.0012**	0.0007**
SB > MB	0.7416	0.4570	0.7222	-	0.3803	0.6716	0.3242
MBodd > SB	0.2584	0.5430	0.2778	-	0.6197	0.3284	0.6758

Table 2: p-values for all t-tests within regions of interest (ROIs) based on the semantic network. * = $p < 0.05$, ** = $p < 0.05$ (Bonferroni-corrected for the number of ROIs), - = ROI does not overlap by at least one voxel with the whole-brain contrast of interest (S > C) summed over all sequences in the comparison, LTP = left temporal pole, LvATL = left ventral anterior temporal lobe, RITG = right inferior temporal gyrus, LFP = left frontal pole, LmMTG = left medial middle temporal gyrus, LpMTG = left posterior middle temporal gyrus, LIFGpt = left inferior temporal gyrus pars triangularis.

3.3.1. Univariate analysis

Table 2 shows the p-values for all planned t-contrasts. pTx provided significantly better activation magnitude than SESB in the left posterior middle temporal gyrus (LpMTG). ME sequences provided significantly better activation magnitude than SE sequences in the left ventral anterior temporal lobe (LvATL).

MB sequences offered significantly better activation precision than SB sequences in the LvATL, right inferior temporal gyrus (RITG) and left inferior frontal gyrus pars triangularis (LIFGpt). MEdn sequences offered significantly better activation precision than ME in the LvATL, RITG, LpMTG, LIFGpt and left frontal pole (LFP). There was no significant difference in either direction between MBodd sequences and SB sequences (see Supplementary Table 1 for detailed results, plus reverse contrasts). We also found no significant interaction between multi-echo and multi-band.

3.3.2. Exploratory MVPA

Table 2 also shows the p-values for t-tests comparing our MVPA dissimilarity metric between sequences. The ME sequences produced significantly better performance than SE sequences in the LvATL and LpMTG. MEdn sequences produced better performance than ME sequences in every ROI. All other comparisons failed to reach significance.

3.4. Whole-brain analysis

Figure 3 shows the results for activation magnitude. Figure 3A shows a single cluster,

within the right fusiform gyrus and lateral inferior occipital cortex, that showed greater

activation magnitude with pTx than with SESB (pTx > SESB). Figure 3B shows the main effect

of ME over SE, featuring clusters in the fusiform and inferior temporal gyri bilaterally plus

the left orbitofrontal cortex (cluster and peak information is provided in Supplementary

Table 2).

Figure 3: Effects on activation magnitude: (A) effect of parallel transmit (pTx > SESB); (B)
effect of echo (ME > SE). Results are cluster-corrected at p < 0.05 based on an uncorrected
voxel threshold of p < 0.001 and are overlaid on the MNI152NLin2009cAsym template.
445

446   Figure 4 shows the results for activation precision. Both the main effect of MB over

447   SB and the main effect of ME-ICA denoising over ME without ICA denoising extended down

448   the temporal lobe and included frontal regions (Supplementary Table 2). There was no

449   significant difference between downsampled MB sequences and SB sequences in either

450   direction.

Figure 4: Effects on activation precision: (A) effect of multiband (MB > SB); (B) effect of ME-ICA denoising (MEdn > ME). Results are cluster-corrected at p < 0.05 based on an uncorrected voxel threshold of p < 0.001 and are overlaid on the MNI152NLin2009cAsym template.

To summarise, these results aligned with results from our ROI analyses – ME increased activation magnitude and both MB and ME-ICA denoising increased activation precision (see Supplementary Figures 1-6 for detailed results, plus reverse contrasts).

3.5. Slice leakage analysis

Seed and possible artefact locations for an example participant are shown in Figure 5A. Figure 5B shows activation magnitude for an example seed and its corresponding possible artefact locations for all participants in each MB sequence and its corresponding sequences. Figure 5C shows MVPA performance. Violin plots for all other peaks are shown in Supplementary Figure 7. t-tests were conducted at each location, but no comparison reached statistical significance (p-values are shown in Supplementary Table 3). We therefore concluded that there was no evidence for slice leakage in either of our MB datasets.

Figure 5: Slice leakage analysis. (A) Seed and possible artefact locations for a single
participant. The ROIs (radius 4 voxels) indicate the seed location (blue), possible artefact
location based on phase shift (green), possible artefact location based on GRAPPA (red), and
possible artefact location based on phase shift and GRAPPA (yellow) in native EPI space.
Informed consent was obtained from the participant for this image to be published. (B)
Mean activation magnitude within each sphere for each MB sequence and the

477 corresponding control sequences for the first seed location (plots for other seeds are shown
478 in Supplementary Figure 7; statistics for all seeds are shown in Supplementary Table 3). (C)
479 Mean MVPA dissimilarity within each sphere for each MB sequence and the corresponding
480 control sequences.

4. Discussion

The tSNR of fMRI varies across the brain. This is especially evident in ventral anterior temporal and orbitofrontal regions, which are located next to the air-filled sinuses and are therefore affected by signal dropout and geometric distortions (Devlin et al., 2000; Halai et al., 2014, 2015, 2024). This study is the first 7T-fMRI study systematically comparing pTx, ME and MB as methods for improving sensitivity in these regions while maintaining sensitivity across the brain. We found that pTx improved activation magnitude in posterior temporal and occipital regions. ME, however, resulted in improved activation magnitude extending down the temporal lobe and including inferior frontal regions. Both MB and ME-ICA denoising resulted in improved activation precision in the same areas. In an exploratory analysis we found that ME and ME-ICA improved MVPA performance but MB did not. No slice leakage artefacts were associated with our multiband sequences.

Although parallel transmit produced better activation magnitude in posterior temporal and occipital regions than the baseline (SESB) sequence without compromising activation magnitude elsewhere, activation magnitude in anterior temporal areas was comparable for both sequences. These results replicated those of Ding et al. (2022) who failed to find improved task contrast in anterior temporal regions with pTx in spite of improved tSNR in the resting state. Note that, despite visible signal dropout on the EPIs, even the baseline sequence was able to identify semantic activity extending into ventral anterior temporal regions (Figure 6). It is possible that this finding reflects the increased sensitivity of 7T-fMRI compared to 3T-fMRI, where semantic activity is rarely observed without using a method such as multi-echo (Halai et al., 2014, 2015, 2024) or spin-echo (Binney et al., 2010; Embleton et al., 2010). That a standard sequence can detect signal in these regions might come as a surprise to many neuroimaging researchers as 7T-fMRI is not only associated with increased sensitivity, but also with exacerbated magnetic susceptibility artefacts. This finding should therefore embolden researchers to use 7T fMRI for experimental designs that are difficult to conduct at a lower field strength - for example, studies of special populations, who may benefit from shorter scan times (Cope et al., 2023),

511   or studies requiring high spatial specificity (Marques & Norris, 2018). We do acknowledge

512   that the way we have utilised 7T fMRI in this study will not be suitable for, for example,

513   studies that require ultra-high resolution (in our case, scanner hardware constrained voxel

514   size when combined with a short first echo time).

516
517

Figure 6: Contrast despite signal dropout and geometric distortions. (A) Mean EPI across all
subjects for the baseline sequence (SESB) registered to MNI152NLin2009cAsym space. (B)
Selected regions of interest overlaid on the mean EPI image. (C) Main effect of activation
magnitude for the contrast of interest (S > C) for the baseline sequence. (D) Main effect of
activation magnitude for the contrast of interest (S > C) for the MEMB sequence. Results are
cluster-corrected at $p < 0.05$ based on an uncorrected voxel threshold of $p < 0.001$.
524

The factorial design enabled us to disentangle the effects of ME and MB. As

hypothesised, ME improved activation magnitude and these effects were localised to

inferior temporal and orbitofrontal areas. Adverse effects on activation magnitude were

negligible - although ME reduced activation precision in two very small clusters

(Supplementary Figure 4), these were not within areas that the semantic task is known to

recruit (Jung et al., 2017). These results demonstrated that having one very short echo is a

successful method of increasing activation magnitude in areas where T_2* is particularly short

(Halai et al., 2014, 2015, 2024; Jung et al., 2017). ME also opens up the opportunity for

sophisticated denoising such as *tedana* (DuPre et al., 2021; Kundu et al., 2011, 2013; The

tedana Community et al., 2023), which improved activation precision in agreement with

previous findings (Amemiya et al., 2019; Gonzalez-Castillo et al., 2016). Small clusters in

which ME-ICA increased activation magnitude (Supplementary Figure 5) or decreased

activation precision (Supplementary Figure 6) were not within regions recruited by the task

(Jung et al., 2017).

As hypothesised, multiband improved activation precision in areas of inhomogeneity

and in agreement with other work (Puckett et al., 2018). Supplementary Figure 4 shows two

very small clusters in which multiband reduced activation precision, but which lie outside

regions recruited by the task (Jung et al., 2017). After downsampling the MB timeseries to

match the number of volumes in the SB timeseries, there was no longer any difference

between the SB and MB sequences; this suggests that it is the increase in the number of

volumes that accounts for the benefits of multiband. Additionally, there was no evidence for

signal leakage into the simultaneously-acquired slice location (McNabb et al., 2020; Todd et

al., 2016), at least for our modest multiband factor of 2. We note that MB and ME seem to

have independent effects as the interaction was not significant within any ROI. However,

ME seems to offer larger activation magnitude (Figure 6) and MB offers better activation

precision. MB can help reduce the longer TR associated with ME and may decrease noise

aliasing in studies using short TRs (~1s).

Although this study focused primarily on univariate effects, we also identified

improvements to multivariate metrics that will be of interest to researchers using

increasingly-popular MVPA tools (Frisby et al., 2023). ME was found to improve MVPA

performance compared to SE sequences and, in turn, MEdn sequences performed

significantly better than ME sequences without denoising in every region of interest (with

no detrimental effects). Future studies employing multivariate methods should strongly

consider using both ME acquisitions and ME-ICA preprocessing.

This study provides evidence supporting use of ME and/or MB sequences to detect

signal in regions prone to susceptibility artefacts. However, it is important to note that

individual 7T MRI scanners have their own hardware and software limitations; this means

that each site must optimise and test feasible parameters locally. For example, our system

did not allow us to take advantage of the higher spatial resolution offered by 7T-fMRI while

retaining an adequately short first echo for our multi-echo sequences. Puckett et al. (2018)

used comparable voxel sizes but had a shorter first TE than us (9.9 ms vs our 11.8 ms);

Miletić et al. (2020) had higher resolution (1.6 mm vs our 2.5 mm isotropic) and a shorter

first echo of 9.66 ms. Additionally, our 7T Terra pTx system imposes very conservative SAR

limits (1W per channel, 8W total) for pTx scans but permits 20W total (i.e. 2.5W per

channel) for circularly polarised mode (CP-mode or "TrueForm") scans (equivalent to single

transmit). We therefore had to use VERSE-modified pTx pulses which are known to be more

sensitive to B_0 inhomogeneity. We also acknowledge that pTx can be combined with multi-

echo or multiband (Ding et al., 2023; Wu et al., 2016). Although we used the pTx head coil

for all scans, the combination of sequences was not implemented on our scanner at the

time of running this study and therefore needs to be tested empirically in future. Thus,

rather than identifying the limits of what is possible, we offer an accessible framework that

will enable researchers to leverage 7T-fMRI for investigating regions affected by

susceptibility artefacts despite the hardware and software constraints of individual

scanners.

5. Conclusion

In this study we compared pTx, ME and MB as methods of improving sensitivity in

temporal and frontal areas prone to magnetic susceptibility artefacts. Both pTx and ME

improved activation magnitude, but only ME showed improvements in artefact-prone

regions. MB and ME-ICA improved activation precision in these areas. Exploratory results

suggested that both ME and ME-ICA may also benefit MVPA. We demonstrated that a multi-

echo, multiband sequence running on the 8Tx32Rx pTx head coil (in CP mode) can detect

signal in susceptible regions while maintaining sensitivity across the whole brain and is

therefore a versatile choice for future studies using high field strength and investigating the

functional roles of ventral temporal and/or orbitofrontal cortex (Binney et al., 2016; Borghesani et al., 2016; Devereux et al., 2018; Duncan, 2010; DuPre et al., 2016; Fernandez et al., 2017; Lambon Ralph et al., 2017; Zahn et al., 2007).

Data and code availability: Data will be made publicly available upon peer review and acceptance. Code is publicly available at: https://github.com/slfrisby/7TOptimisation/.

CRediT statement: Saskia L. Frisby: Conceptualisation, Methodology, Investigation, Formal Analysis, Writing – Original Draft, Writing – Review & Editing, Visualisation; Marta M. Correia: Conceptualisation, Methodology, Writing – Review & Editing, Funding Acquisition; Minghao Zhang: Methodology, Writing – Review & Editing; Christopher T. Rodgers: Methodology, Writing – Review & Editing, Supervision; Timothy T. Rogers: Writing – Review & Editing, Supervision; Matthew A. Lambon Ralph: Writing – Review & Editing, Supervision, Funding Acquisition; Ajay D. Halai: Conceptualisation, Methodology, Formal Analysis, Writing – Review & Editing, Supervision, Funding Acquisition.

Funding: This work was supported by an MRC Unit grant (SUAG/019 G116768) to M.M.C., an MRC PhD studentship (MR N013433-1) to M.Z., an MRC programme grant (MR/R023883/1) and intramural funding (MC_UU_00005/18) to M.A.L.R, and an MRC Career Development Award (MR/V031481/1) to A.D.H. This work was also supported by the NIHR Cambridge Biomedical Research Centre (NIHR203312) and an MRC Clinical Research Infrastructure Award for 7T (MR/M008983/1). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Competing interests: C.T.R. receives research support from Siemens.

Acknowledgements: We thank the participants and the Wolfson Brain Imaging Centre radiographers for their assistance.

References

Amemiya, S., Yamashita, H., Takao, H., & Abe, O. (2019). Integrated multi-echo denoising strategy improves identification of inherent language laterality. *Magnetic Resonance in Medicine*, *81*(5), 3262–3271. https://doi.org/10.1002/mrm.27620

Andersson, J. L. R., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-echo echo-planar images: Application to diffusion tensor imaging. *NeuroImage*, *20*(2), 870–888. https://doi.org/10.1016/S1053-8119(03)00336-7

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage*, *54*(3), 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Avants, B. B., Tustison, N., & Johnson, H. (2009). *Advanced Normalization Tools (ANTS)*.

Barth, M., Breuer, F., Koopmans, P. J., Norris, D. G., & Poser, B. A. (2016). Simultaneous multislice (SMS) imaging techniques. *Magnetic Resonance in Medicine*, *75*(1), 63–81. https://doi.org/10.1002/mrm.25897

Bhavsar, S., Zvyagintsev, M., & Mathiak, K. (2014). BOLD sensitivity and SNR characteristics of parallel imaging-accelerated single-shot multi-echo EPI for fMRI. *NeuroImage*, *84*, 65–75. https://doi.org/10.1016/j.neuroimage.2013.08.007

Binney, R. J., Embleton, K. V., Jefferies, E., Parker, G. J. M., & Lambon Ralph, M. A. (2010). The ventral and inferolateral aspects of the anterior temporal lobe are crucial in semantic memory: Evidence from a novel direct comparison of distortion-corrected fMRI, rTMS, and semantic dementia. *Cerebral Cortex*, *20*(11), 2728–2738. https://doi.org/10.1093/cercor/bhq019

Binney, R. J., Hoffman, P., & Lambon Ralph, M. A. (2016). Mapping the Multiple Graded Contributions of the Anterior Temporal Lobe Representational Hub to Abstract and Social Concepts: Evidence from Distortion-corrected fMRI. *Cerebral Cortex*, *26*(11), 4227–4241. https://doi.org/10.1093/cercor/bhw260

Borghesani, V., Pedregosa, F., Buiatti, M., Amadon, A., Eger, E., & Piazza, M. (2016). Word meaning in the ventral visual path: A perceptual to conceptual gradient of semantic coding. *NeuroImage*, *143*, 128–140. https://doi.org/10.1016/j.neuroimage.2016.08.068

Boyacioğlu, R., Schulz, J., Koopmans, P. J., Barth, M., & Norris, D. G. (2015). Improved sensitivity and specificity for resting state and task fMRI with multiband multi-echo EPI compared to multi-echo EPI at 7 T. *NeuroImage*, *119*, 352–361. https://doi.org/10.1016/j.neuroimage.2015.06.089

Bozeat, S., Lambon Ralph, M. A., Patterson, K., Garrard, P., & Hodges, J. R. (2000). Non-verbal semantic impairment in semantic dementia. *Neuropsychologia*, *38*(9), 1207–1215. https://doi.org/10.1016/S0028-3932(00)00034-8

Caballero-Gaudes, C., Moia, S., Panwar, P., Bandettini, P. A., & Gonzalez-Castillo, J. (2019). A deconvolution algorithm for multi-echo functional MRI: Multi-echo Sparse Paradigm Free Mapping. *NeuroImage*, *202*, 116081. https://doi.org/10.1016/j.neuroimage.2019.116081

Cohen, A. D., Nencka, A. S., Lebel, R. M., & Wang, Y. (2017). Multiband multi-echo imaging of simultaneous oxygenation and flow timeseries for resting state connectivity. *PLOS ONE*, *12*(3), e0169253. https://doi.org/10.1371/journal.pone.0169253

Cohen, A. D., Nencka, A. S., & Wang, Y. (2018). Multiband multi-echo simultaneous ASL/BOLD for task-induced functional MRI. *PLOS ONE*, *13*(2), e0190427. https://doi.org/10.1371/journal.pone.0190427

Cope, T. E., Sohoglu, E., Peterson, K. A., Jones, P. S., Rua, C., Passamonti, L., Sedley, W., Post, B., Coebergh, J., Butler, C. R., Garrard, P., Abdel-Aziz, K., Husain, M., Griffiths, T. D., Patterson, K., Davis, M. H., & Rowe, J. B. (2023). Temporal lobe perceptual predictions for speech are instantiated in motor cortex and reconciled by inferior frontal cortex. *Cell Reports*, *42*(5). https://doi.org/10.1016/j.celrep.2023.112422

Coutanche, M. N. (2013). Distinguishing multi-voxel patterns and mean activation: Why, how, and what does it tell us? *Cognitive, Affective, & Behavioral Neuroscience*, *13*(3), 667–673. https://doi.org/10.3758/s13415-013-0186-2

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173. https://doi.org/10.1006/cbmr.1996.0014

Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data. *NMR in Biomedicine*, *10*(4–5), 171–178. https://doi.org/10.1002/(SICI)1099-1492(199706/08)10:4/5<171::AID-NBM453>3.0.CO;2-L

Davis, T., LaRocque, K. F., Mumford, J. A., Norman, K. A., Wagner, A. D., & Poldrack, R. A. (2014). What do differences between multi-voxel and univariate analysis mean? How subject-, voxel-, and trial-level variance impact fMRI analysis. *NeuroImage*, *97*, 271–283. https://doi.org/10.1016/j.neuroimage.2014.04.037

Davis, T., & Poldrack, R. A. (2013). Measuring neural representations with fMRI: Practices and pitfalls. *Annals of the New York Academy of Sciences*, *1296*(1), 108–134. https://doi.org/10.1111/nyas.12156

Demetriou, L., Kowalczyk, O. S., Tyson, G., Bello, T., Newbould, R. D., & Wall, M. B. (2018). A comprehensive evaluation of increasing temporal resolution with multiband-accelerated protocols and effects on statistical outcome measures in fMRI. *NeuroImage*, *176*, 404–416. https://doi.org/10.1016/j.neuroimage.2018.05.011

Deniz, F., Nunez-Elizalde, A. O., Huth, A. G., & Gallant, J. L. (2019). The Representation of Semantic Information Across Human Cerebral Cortex During Listening Versus Reading Is Invariant to Stimulus Modality. *The Journal of Neuroscience*, *39*(39), 7722–7736. https://doi.org/10.1523/JNEUROSCI.0675-19.2019

Devereux, B. J., Clarke, A., & Tyler, L. K. (2018). Integrated deep visual and semantic attractor neural networks predict fMRI pattern-information along the ventral object processing pathway. *Scientific Reports*, *8*(1), 10636. https://doi.org/10.1038/s41598-018-28865-1

Devlin, J. T., Russell, R. P., Davis, M. H., Price, C. J., Wilson, J., Moss, H. E., Matthews, P. M., & Tyler, L. K. (2000). Susceptibility-induced loss of signal: Comparing PET and fMRI on a semantic task. *NeuroImage*, *11*(6), 589–600. https://doi.org/10.1006/nimg.2000.0595

Ding, B., Dragonu, I., Rua, C., Carlin, J. D., Halai, A. D., Liebig, P., Heidemann, R., Correia, M. M., & Rodgers, C. T. (2022). Parallel transmit (pTx) with online pulse design for task-based fMRI at 7T. *Magnetic Resonance Imaging*, *93*, 163–174. https://doi.org/10.1016/j.mri.2022.07.003

Ding, B., Williams, S. N., Zhang, M., Herrler, J., Liebig, P., Dragonu, I., Neji, R., Rodgers, C. T., & Porter, D. A. (2023). *Optimisation of pulse-specific phase schedule reduces peak RF amplitude in multiband parallel-transmit pulses*. ISMRM & ISMRT Annual Meeting & Exhibition, Toronto, Canada.

Dipasquale, O., Sethi, A., Laganà, M. M., Baglio, F., Baselli, G., Kundu, P., Harrison, N. A., & Cercignani, M. (2017). Comparing resting state fMRI de-noising approaches using

multi- and single-echo acquisitions. *PLOS ONE*, *12*(3), e0173289.
https://doi.org/10.1371/journal.pone.0173289

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: Mental programs
for intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179.
https://doi.org/10.1016/j.tics.2010.01.004

DuPre, E., Luh, W.-M., & Spreng, R. N. (2016). Multi-echo fMRI replication sample of
autobiographical memory, prospection and theory of mind reasoning tasks. *Scientific
Data*, *3*(1), 160116. https://doi.org/10.1038/sdata.2016.116

DuPre, E., Salo, T., Ahmed, Z., Bandettini, P. A., Bottenhorn, K. L., Caballero-Gaudes, C.,
Dowdle, L. T., Gonzalez-Castillo, J., Heunis, S., Kundu, P., Laird, A. R., Markello, R.,
Markiewicz, C. J., Moia, S., Staden, I., Teves, J. B., Uruñuela, E., Vaziri-Pashkam, M.,
Whitaker, K., & Handwerker, D. A. (2021). TE-dependent analysis of multi-echo fMRI
with tedana. *Journal of Open Source Software*, *6*(66), 3669.
https://doi.org/10.21105/joss.03669

Embleton, K. V., Haroon, H. A., Morris, D. M., Lambon Ralph, M. A. L., & Parker, G. J. M.
(2010). Distortion correction for diffusion-weighted MRI tractography and fMRI in
the temporal lobes. *Human Brain Mapping*, *31*(10), 1570–1587.
https://doi.org/10.1002/hbm.20959

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D.,
Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J.,
Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: A robust preprocessing
pipeline for functional MRI. *Nature Methods*, *16*(1), 111–116.
https://doi.org/10.1038/s41592-018-0235-4

Evans, J. W., Kundu, P., Horovitz, S. G., & Bandettini, P. A. (2015). Separating slow BOLD
from non-BOLD baseline drifts using multi-echo fMRI. *NeuroImage*, *105*, 189–197.
https://doi.org/10.1016/j.neuroimage.2014.10.051

Feinberg, D. A., Moeller, S., Smith, S. M., Auerbach, E., Ramanna, S., Glasser, M. F., Miller, K.
L., Ugurbil, K., & Yacoub, E. (2010). Multiplexed Echo Planar Imaging for Sub-Second
Whole Brain FMRI and Fast Diffusion Imaging. *PLOS ONE*, *5*(12), e15710.
https://doi.org/10.1371/journal.pone.0015710

Fernandez, B., Leuchs, L., Sämann, P. G., Czisch, M., & Spoormaker, V. I. (2017). Multi-echo
EPI of human fear conditioning reveals improved BOLD detection in ventromedial
prefrontal cortex. *NeuroImage*, *156*, 65–77.
https://doi.org/10.1016/j.neuroimage.2017.05.005

Frisby, S. L., Halai, A. D., Cox, C. R., Lambon Ralph, M. A., & Rogers, T. T. (2023). Decoding
semantic representations in mind and brain. *Trends in Cognitive Sciences*, *27*(3),
258–281. https://doi.org/10.1016/j.tics.2022.12.006

Gaser, C., Dahnke, R., Thompson, P. M., Kurth, F., Luders, E., & Alzheimer's Disease
Neuroimaging Initiative. (2023). *CAT − A computational anatomy toolbox for the
analysis of structural MRI data* (p. 2022.06.11.495736). bioRxiv.
https://doi.org/10.1101/2022.06.11.495736

Gilmore, A. W., Agron, A. M., González-Araya, E. I., Gotts, S. J., & Martin, A. (2022). A
comparison of single- and multi-echo processing of functional MRI data during overt
autobiographical recall. *Frontiers in Neuroscience*, *16*, 854387.
https://doi.org/10.3389/fnins.2022.854387

Gonzalez-Castillo, J., Panwar, P., Buchanan, L. C., Caballero-Gaudes, C., Handwerker, D. A.,
Jangraw, D. C., Zachariou, V., Inati, S., Roopchansingh, V., Derbyshire, J. A., &

Bandettini, P. A. (2016). Evaluation of multi-echo ICA denoising for task based fMRI studies: Block designs, rapid event-related designs, and cardiac-gated fMRI. *NeuroImage*, *141*, 452–468. https://doi.org/10.1016/j.neuroimage.2016.07.049

Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P., Flandin, G., Ghosh, S. S., Glatard, T., Halchenko, Y. O., Handwerker, D. A., Hanke, M., Keator, D., Li, X., Michael, Z., Maumet, C., Nichols, B. N., Nichols, T. E., Pellman, J., … Poldrack, R. A. (2016). The brain imaging data structure, a format for organizing and describing outputs of neuroimaging experiments. *Scientific Data*, *3*(1), 160044. https://doi.org/10.1038/sdata.2016.44

Gorgolewski, K. J., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., & Ghosh, S. S. (2011). Nipype: A flexible, lightweight and extensible neuroimaging data processing framework in python. *Frontiers in Neuroinformatics*, *5*. https://doi.org/10.3389/fninf.2011.00013

Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C. E., Zsoldos, E., Ebmeier, K. P., Filippini, N., Mackay, C. E., Moeller, S., Xu, J., Yacoub, E., Baselli, G., Ugurbil, K., Miller, K. L., & Smith, S. M. (2014). ICA-based artefact removal and accelerated fMRI acquisition for improved resting state network imaging. *NeuroImage*, *95*, 232–247. https://doi.org/10.1016/j.neuroimage.2014.03.034

Hagberg, G. E., Indovina, I., Sanes, J. N., & Posse, S. (2002). Real-time quantification of T2* changes using multiecho planar imaging and numerical methods. *Magnetic Resonance in Medicine*, *48*(5), 877–882. https://doi.org/10.1002/mrm.10283

Halai, A. D., Henson, R. N., Finoia, P., & Correia, M. M. (2024). *Comparing the effect of multi gradient echo and multi band fMRI during a semantic task* (p. 2024.03.20.585909). bioRxiv. https://doi.org/10.1101/2024.03.20.585909

Halai, A. D., Parkes, L. M., & Welbourne, S. R. (2015). Dual-echo fMRI can detect activations in inferior temporal lobe during intelligible speech comprehension. *NeuroImage*, *122*, 214–221. https://doi.org/10.1016/j.neuroimage.2015.05.067

Halai, A. D., Welbourne, S. R., Embleton, K., & Parkes, L. M. (2014). A comparison of dual gradient-echo and spin-echo fMRI of the inferior temporal lobe. *Human Brain Mapping*, *35*(8), 4118–4128. https://doi.org/10.1002/hbm.22463

Halchenko, Y. O., Goncalves, M., Ghosh, S., Velasco, P., Castello, M. V. di O., Salo, T., Wodder, J. T., Hanke, M., Sadil, P., Gorgolewski, K. J., Ioanas, H.-I., Rorden, C., Hendrickson, T. J., Dayan, M., Houlihan, S. D., Kent, J., Strauss, T., Lee, J., To, I., … Kennedy, D. N. (2024). HeuDiConv—Flexible DICOM conversion into structured directory layouts. *Journal of Open Source Software*, *9*(99), 5839. https://doi.org/10.21105/joss.05839

Hargreaves, B. A., Cunningham, C. H., Nishimura, D. G., & Conolly, S. M. (2004). Variable-rate selective excitation for rapid MRI sequences. *Magnetic Resonance in Medicine*, *52*(3), 590–597. https://doi.org/10.1002/mrm.20168

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and overlapping representations of faces and objects in ventral temporal cortex. *Science*, *293*(5539), 2425–2430. https://doi.org/10.1126/science.1063736

Howard, D., & Patterson, K. E. (1992). The pyramids and palm trees test. *Https://Eprints.Ncl.Ac.Uk*. https://eprints.ncl.ac.uk

Humphreys, G. F., Hoffman, P., Visser, M., Binney, R. J., & Lambon Ralph, M. A. (2015). Establishing task- and modality-dependent dissociations between the semantic and

default mode networks. *Proceedings of the National Academy of Sciences*, *112*(25), 7857–7862. https://doi.org/10.1073/pnas.1422760112

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL. *NeuroImage*, *62*(2), 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jung, J., Williams, S. R., Sanaei Nezhad, F., & Lambon Ralph, M. A. (2017). GABA concentrations in the anterior temporal lobe predict human semantic processing. *Scientific Reports*, *7*(1), 15748. https://doi.org/10.1038/s41598-017-15981-7

Kirilina, E., Lutti, A., Poser, B. A., Blankenburg, F., & Weiskopf, N. (2016). The quest for the best: The impact of different EPI sequences on the sensitivity of random effect fMRI group analyses. *NeuroImage*, *126*, 49–59. https://doi.org/10.1016/j.neuroimage.2015.10.071

Koopmans, P. J., Barth, M., Orzada, S., & Norris, D. G. (2011). Multi-echo fMRI of the cortical laminae in humans at 7T. *NeuroImage*, *56*(3), 1276–1285. https://doi.org/10.1016/j.neuroimage.2011.02.042

Kovářová, A., Gajdoš, M., Rektor, I., & Mikl, M. (2022). Contribution of the multi-echo approach in accelerated functional magnetic resonance imaging multiband acquisition. *Human Brain Mapping*, *43*(3), 955–973. https://doi.org/10.1002/hbm.25698

Kundu, P., Benson, B. E., Baldwin, K. L., Rosen, D., Luh, W.-M., Bandettini, P. A., Pine, D. S., & Ernst, M. (2015). Robust resting state fMRI processing for studies on typical brain development based on multi-echo EPI acquisition. *Brain Imaging and Behavior*, *9*(1), 56–73. https://doi.org/10.1007/s11682-014-9346-4

Kundu, P., Brenowitz, N. D., Voon, V., Worbe, Y., Vértes, P. E., Inati, S. J., Saad, Z. S., Bandettini, P. A., & Bullmore, E. T. (2013). Integrated strategy for improving functional connectivity mapping using multiecho fMRI. *Proceedings of the National Academy of Sciences*, *110*(40), 16187–16192. https://doi.org/10.1073/pnas.1301725110

Kundu, P., Inati, S. J., Evans, J. W., Luh, W.-M., & Bandettini, P. A. (2011). Differentiating BOLD and non-BOLD signals in fMRI time series using multi-echo EPI. *NeuroImage*, *60*(3), 1759–1770. https://doi.org/10.1016/j.neuroimage.2011.12.028

Kundu, P., Voon, V., Balchandani, P., Lombardo, M. V., Poser, B. A., & Bandettini, P. A. (2017). Multi-echo fMRI: A review of applications in fMRI denoising and analysis of BOLD signals. *NeuroImage*, *154*, 59–80. https://doi.org/10.1016/j.neuroimage.2017.03.033

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and computational bases of semantic cognition. *Nature Reviews Neuroscience*, *18*(1), 42–55. https://doi.org/10.1038/nrn.2016.150

Lombardo, M. V., Auyeung, B., Holt, R. J., Waldman, J., Ruigrok, A. N. V., Mooney, N., Bullmore, E. T., Baron-Cohen, S., & Kundu, P. (2016). Improving effect size estimation and statistical power with multi-echo fMRI and its impact on understanding the neural systems supporting mentalizing. *NeuroImage*, *142*, 55–66. https://doi.org/10.1016/j.neuroimage.2016.07.022

Lynch, C. J., Power, J. D., Scult, M. A., Dubin, M., Gunning, F. M., & Liston, C. (2020). Rapid Precision Functional Mapping of Individuals Using Multi-Echo fMRI. *Cell Reports*, *33*(12), 108540. https://doi.org/10.1016/j.celrep.2020.108540

Markiewicz, C. J., Esteban, O., Goncalves, M., Provins, C., Salo, T., Kent, J. D., DuPre, E., Ciric, R., Pinsard, B., Blair, R. W., Poldrack, R. A., & Gorgolewski, K. J. (2024). *fMRIPrep: A*

855   *robust preprocessing pipeline for functional MRI* (Version 24.0.1) [Computer
856   software]. Zenodo. https://doi.org/10.5281/zenodo.12774221

857 Marques, J. P., & Norris, D. G. (2018). How to choose the right MR sequence for your
858   research question at 7 T and above? *NeuroImage*, *168*, 119–140.
859   https://doi.org/10.1016/j.neuroimage.2017.04.044

860 McNabb, C. B., Lindner, M., Shen, S., Burgess, L. G., Murayama, K., & Johnstone, T. (2020).
861   Inter-slice leakage and intra-slice aliasing in simultaneous multi-slice echo-planar
862   images. *Brain Structure and Function*, *225*(3), 1153–1158.
863   https://doi.org/10.1007/s00429-020-02053-2

864 Miletić, S., Bazin, P.-L., Weiskopf, N., van der Zwaag, W., Forstmann, B. U., & Trampel, R.
865   (2020). FMRI protocol optimization for simultaneously studying small subcortical and
866   cortical areas at 7 T. *NeuroImage*, *219*, 116992.
867   https://doi.org/10.1016/j.neuroimage.2020.116992

868 Moeller, S., Yacoub, E., Olman, C. A., Auerbach, E., Strupp, J., Harel, N., & Uğurbil, K. (2010).
869   Multiband multislice GE-EPI at 7 Tesla, with 16-fold acceleration using partial parallel
870   imaging with application to high spatial and temporal whole-brain fMRI. *Magnetic
871   Resonance in Medicine*, *63*(5), 1144–1153. https://doi.org/10.1002/mrm.22361

872 Morris, L. S., Kundu, P., Costi, S., Collins, A., Schneider, M., Verma, G., Balchandani, P., &
873   Murrough, J. W. (2019). Ultra-high field MRI reveals mood-related circuit
874   disturbances in depression: A comparison between 3-Tesla and 7-Tesla. *Translational
875   Psychiatry*, *9*(1), 94. https://doi.org/10.1038/s41398-019-0425-6

876 O'Brien, K. R., Kober, T., Hagmann, P., Maeder, P., Marques, J., Lazeyras, F., Krueger, G., &
877   Roche, A. (2014). Robust T1-weighted structural brain imaging and morphometry at
878   7T using MP2RAGE. *PLOS ONE*, *9*(6), e99676.
879   https://doi.org/10.1371/journal.pone.0099676

880 Poser, B. A., & Norris, D. G. (2009). Investigating the benefits of multi-echo EPI for fMRI at 7
881   T. *NeuroImage*, *45*(4), 1162–1172.
882   https://doi.org/10.1016/j.neuroimage.2009.01.007

883 Poser, B. A., Versluis, M. J., Hoogduin, J. M., & Norris, D. G. (2006). BOLD contrast sensitivity
884   enhancement and artifact reduction with multiecho EPI: Parallel-acquired
885   inhomogeneity-desensitized fMRI. *Magnetic Resonance in Medicine*, *55*(6), 1227–
886   1235. https://doi.org/10.1002/mrm.20900

887 Posse, S. (2012). Multi-echo acquisition. *NeuroImage*, *62*(2), 665–671.
888   https://doi.org/10.1016/j.neuroimage.2011.10.057

889 Puckett, A. M., Bollmann, S., Poser, B. A., Palmer, J., Barth, M., & Cunnington, R. (2018).
890   Using multi-echo simultaneous multi-slice (SMS) EPI to improve functional MRI of
891   the subcortical nuclei of the basal ganglia at ultra-high field (7T). *NeuroImage*, *172*,
892   886–895. https://doi.org/10.1016/j.neuroimage.2017.12.005

893 Risk, B. B., Murden, R. J., Wu, J., Nebel, M. B., Venkataraman, A., Zhang, Z., & Qiu, D. (2021).
894   Which multiband factor should you choose for your resting-state fMRI study?
895   *NeuroImage*, *234*, 117965. https://doi.org/10.1016/j.neuroimage.2021.117965

896 Setsompop, K., Gagoski, B. A., Polimeni, J. R., Witzel, T., Wedeen, V. J., & Wald, L. L. (2012).
897   Blipped-controlled aliasing in parallel imaging for simultaneous multislice echo
898   planar imaging with reduced *g* -factor penalty. *Magnetic Resonance in Medicine*,
899   *67*(5), 1210–1224. https://doi.org/10.1002/mrm.23097

900 Smith, S. M. (2002). Fast robust automated brain extraction. *Human Brain Mapping*, *17*(3),
901   143–155. https://doi.org/10.1002/hbm.10062

Smith, S. M., Beckmann, C. F., Andersson, J., Auerbach, E. J., Bijsterbosch, J., Douaud, G.,
    Duff, E., Feinberg, D. A., Griffanti, L., Harms, M. P., Kelly, M., Laumann, T., Miller, K.
    L., Moeller, S., Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A. Z., Vu, A. T., …
    Glasser, M. F. (2013). Resting-state fMRI in the Human Connectome Project.
    NeuroImage, *80*, 144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-
    Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K.,
    Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M.
    (2004). Advances in functional and structural MR image analysis and implementation
    as FSL. *NeuroImage*, *23*, S208–S219.
    https://doi.org/10.1016/j.neuroimage.2004.07.051

The tedana Community, Ahmed, Z., Bandettini, P. A., Bottenhorn, K. L., Caballero-Gaudes,
    C., Dowdle, L. T., DuPre, E., Gonzalez-Castillo, J., Handwerker, D., Heunis, S., Kundu,
    P., Laird, A. R., Markello, R., Markiewicz, C. J., Maullin-Sapey, T., Moia, S., Molfese,
    P., Salo, T., Staden, I., … Whitaker, K. (2023). *ME-ICA/tedana: 23.0.1* (Version 23.0.1)
    [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.7926293

Todd, N., Moeller, S., Auerbach, E. J., Yacoub, E., Flandin, G., & Weiskopf, N. (2016).
    Evaluation of 2D multiband EPI imaging for high-resolution, whole-brain, task-based
    fMRI studies at 3T: Sensitivity and slice leakage artifacts. *NeuroImage*, *124*, 32–42.
    https://doi.org/10.1016/j.neuroimage.2015.08.056

Wu, X., Schmitter, S., Auerbach, E. J., Uğurbil, K., & Van de Moortele, P.-F. (2016). A
    generalized slab-wise framework for parallel transmit multiband RF pulse design.
    Magnetic Resonance in Medicine, *75*(4), 1444–1456.
    https://doi.org/10.1002/mrm.25689

Yun, S. D., & Shah, N. J. (2017). Whole-brain high in-plane resolution fMRI using accelerated
    EPIK for enhanced characterisation of functional areas at 3T. *PLOS ONE*, *12*(9),
    e0184759. https://doi.org/10.1371/journal.pone.0184759

Zahn, R., Moll, J., Krueger, F., Huey, E. D., Garrido, G., & Grafman, J. (2007). Social concepts
    are represented in the superior anterior temporal cortex. *Proceedings of the
    National Academy of Sciences*, *104*(15), 6430–6435.
    https://doi.org/10.1073/pnas.0607061104