Optimising 7T-fMRI for imaging regions of magnetic susceptibility

Saskia L. Frisby[1], Marta M. Correia[1], Minghao Zhang[2], Christopher T. Rodgers[2], Timothy T.

Rogers[3], Matthew A. Lambon Ralph[1], and Ajay D. Halai[1]

[1]MRC Cognition and Brain Sciences Unit, 15 Chaucer Road, Cambridge CB2 7EF

[2]Wolfson Brain Imaging Centre, Cambridge Biomedical Campus, Cambridge CB2 0QQ

[3]Department of Psychology, University of Wisconsin-Madison, 1202 W Johnson Street,

Madison 53706

Correspondence to:

Dr. Saskia L. Frisby

saskia.frisby@mrc-cbu.cam.ac.uk

+44 (0)1223 769703

and

Dr. Ajay D. Halai

ajay.halai@mrc-cbu.cam.ac.uk

+44 (0)1223 767655

MRC Cognition & Brain Sciences Unit

University of Cambridge

15 Chaucer Road

Cambridge

CB2 7EF

28 **Acknowledgements:** We thank the participants and the Wolfson Brain Imaging Centre

29 radiographers for their assistance.

Abstract

The temporal signal-to-noise ratio (tSNR) of functional magnetic resonance imaging (fMRI) is

particularly poor in ventral anterior temporal and orbitofrontal regions because of B_0 and

B_1^+ magnetic field inhomogeneity, a problem that is exacerbated at higher field strengths. In

this 7T-fMRI study we compared three methods of improving sensitivity in these areas:

parallel transmit, which uses multiple transmit elements, controlled independently, to

homogenise the flip angle experienced by the tissue; multi-echo, which entails collection of

multiple volumes at different echo times following a single radiofrequency pulse; and

multiband, in which multiple slices are acquired simultaneously. We found that parallel

transmit and multi-echo increased the magnitude of the BOLD signal change, but only multi-

echo increased BOLD magnitude in areas prone to susceptibility artefacts. Multiband and

denoising of multi-echo data with independent components analysis (ICA) both improved

precision of GLM fit. Exploratory results suggested that multi-echo and ICA denoising can

both benefit multivariate analyses. In conclusion, a multi-echo, multiband sequence

improved fMRI quality in areas prone to susceptibility artefacts while maintaining sensitivity

across the whole brain. We recommend this approach for studies investigating the

functional roles of ventral temporal and orbitofrontal regions with 7T fMRI.

Keywords: 7T-fMRI, parallel transmit, multi-echo, multiband, temporal lobe, orbitofrontal

cortex

1. Introduction

The temporal signal-to-noise ratio (tSNR) of functional magnetic resonance imaging

(fMRI) varies across the brain. The ventral anterior temporal cortex and orbitofrontal cortex,

for example, are affected by two kinds of magnetic field inhomogeneity. First, these regions

are located next to air-filled sinuses and so are affected by B_0 inhomogeneity that causes

signal dropout and geometric distortions (Devlin et al., 2000; Halai et al., 2014, 2015, 2024;

Jezzard & Clare, 1999). The effects are more severe at higher field strengths, implying that it

may be especially difficult to measure task-related activity in susceptible regions with ultra-

high-field fMRI (e.g., 7T-fMRI). Second, at 7T and above, the radiofrequency wavelength is

similar to the size of the head, resulting in standing wave effects that cause B_1^+

inhomogeneity in the temporal lobes (Gras et al., 2019; van der Kolk et al., 2013). This in

turn causes spatial variation in flip angle and therefore in image intensity (Uğurbil, 2018; Wu

et al., 2018). B_0 and B_1^+ inhomogeneity make it challenging to use fMRI to investigate the

roles that these regions may play in a myriad of cognitive processes – including vision

(Devereux et al., 2018), language (Borghesani et al., 2016), multimodal semantic cognition

(Lambon Ralph et al., 2017), emotion (Fernandez et al., 2017), social cognition (Binney et al.,

2016; Zahn et al., 2007), theory of mind (DuPre et al., 2016), and executive function

(Duncan, 2010). However, 7T-fMRI also has many advantages in regions unaffected by

inhomogeneity: 7T-fMRI offers improved tSNR relative to 3T-fMRI (Morris et al., 2019),

which can be used to reduce voxel size and enable applications such as laminar fMRI

(Koopmans et al., 2011) or to reduce acquisition times and enable shorter scan times for

special populations such as patients with neurodegenerative diseases (Cope et al., 2023).

7T-fMRI also benefits from improved spatial specificity relative to 3T-fMRI because the

signal from cortical microvasculature is enhanced while the signal from large veins is

reduced (Marques & Norris, 2018). Improving signal homogeneity would allow researchers

studying the whole brain (or focusing on regions prone to susceptibility artefacts) to take

full advantage of 7T-fMRI; therefore, in this study we compared three methods of doing so.

One possible method for improving image quality is parallel transmit (pTx), which

uses multiple transmit elements, controlled independently, to homogenise the flip angle

pattern experienced by the tissue (Adriany et al., 2005; Deniz et al., 2019; Roemer et al.,

1990; Uğurbil, 2018; Van de Moortele et al., 2005). In this way, pTx directly counteracts the

effect of B_1^+ inhomogeneity – however, since the design of spoke pTx pulses incorporates B_0

field maps, pTx is also capable of counteracting B_0 inhomogeneity to an extent (Zhang et al.,

2024). A recent study used pTx to improve image quality in ventral anterior temporal

regions for echo-planar imaging (EPI) 7T fMRI (Ding et al., 2022). pTx improved tSNR across

the brain compared to a standard sequence, particularly in the temporal lobes. However,

there was no improvement in functional contrast during a semantic association task that is

known to recruit the anterior temporal lobes in 3T-fMRI studies (Jung et al., 2017).

A second method of recovering signal in these regions is multi-echo (ME) imaging

(Kundu et al., 2017; Poser et al., 2006; Posse, 2012), which aims to counteract the effects of

B_0 inhomogeneity. T_2* is known to vary across the brain (Hagberg et al., 2002); in areas

affected by B_0 inhomogeneity, T_2* is particularly short due to increased intravoxel

dephasing. A single echo provides sensitivity to a narrow range of T_2* values; the echo time

(TE) is therefore selected to provide the best compromise of sensitivity to T_2* across the

whole brain. Combining data from multiple echoes increases the range of T_2* that can be

imaged with high fidelity. ME has been shown to improve functional contrast (Poser &

Norris, 2009) and spatial specificity (Boyacioğlu et al., 2015) at 7T and 3T (Fernandez et al.,

2017; Halai et al., 2024; Kirilina et al., 2016; Lynch et al., 2020). Having multiple echoes also

facilitates the separation of signal and noise because signals decay in a well-characterised

way across echoes, whereas noise does not. This principle underpins multi-echo

independent components analysis (ME-ICA), via which ICA components that are TE-

independent, and thus are likely to be noise rather than blood-oxygen-level-dependent

(BOLD) signal, can be removed (Dipasquale et al., 2017; Kundu et al., 2011, 2013, 2015,

2017). This method may enhance signal detection in areas prone to susceptibility artefacts

on top of the advantage offered by ME alone (e.g. Lombardo et al., 2016). ME sequences

have some potential disadvantages. For example, ME can lengthen repetition time (TR). In-

plane acceleration is frequently needed to achieve a sufficiently short first TE, which reduces

tSNR (Yun & Shah, 2017). In turn, a short first TE, combined with hardware constraints,

often limits the minimum voxel size (Koopmans et al., 2011). Critically, in many previous

studies examining the benefits of ME sequences compared to single-echo (SE), the "single-

echo" data were extracted from the ME dataset (Amemiya et al., 2019; Bhavsar et al., 2014;

Caballero-Gaudes et al., 2019; Cohen et al., 2017, 2017, 2018; Dipasquale et al., 2017; Evans

et al., 2015; Fernandez et al., 2017; Gilmore et al., 2022; Kovářová et al., 2022). This means

that the "single-echo" data will inherit suboptimal parameters that are ME-specific, making

the comparison unfair.

A third strategy for improving acquisition is multiband (MB) imaging, also known as

simultaneous multi-slice, in which multiple slices are acquired simultaneously (Barth et al.,

2016; Moeller et al., 2010; Setsompop et al., 2012). Rather than counteracting B_0 or B_1^+

inhomogeneity directly, MB has typically been applied to increase temporal resolution –

multiple studies have shown that MB can reduce noise aliasing, increase statistical power

and counteract the increase in TR associated with ME (Feinberg et al., 2010; Griffanti et al.,

2014; Halai et al., 2024; Puckett et al., 2018; Smith et al., 2013). Benefits specific to task

fMRI have been less clear (Demetriou et al., 2018; Todd et al., 2016) since some noise

features are unlikely to be correlated with task regressors and can be removed with high-

pass filtering. Other possible disadvantages of MB include a reduction in tSNR due to

increases in g-factor effects (Demetriou et al., 2018; Risk et al., 2021; Setsompop et al.,

2012) and leakage of signal into the simultaneously-excited slices (Todd et al., 2016).

Previous work therefore makes clear predictions about which 7T-fMRI sequences are

capable of improving sensitivity in regions affected by B_0 and B_1^+ inhomogeneity without

compromising signal quality elsewhere. In this study we tested five possible sequences with

the goal of demonstrating their *relative* impacts on functional effects of interest. These

consisted of a 2 x 2 factorial design varying number of echoes and multiband factor – single-

echo single band (SESB), single-echo multiband (SEMB), multi-echo single band (MESB), and

multi-echo multiband (MEMB) – plus a fifth sequence that used pTx. Rather than attempting

to map the limits of each parameter, our aim was to generate sequences that would be of

practical benefit to neuroscientists seeking to answer a range of cognitive questions that

implicate regions affected by inhomogeneity.

We used a semantic judgment task that is known to evoke activity across the

semantic network, including areas severely affected and those relatively unaffected by

susceptibility artefacts (Jung et al., 2017).

We had two univariate effects of interest – activation magnitude and activation

precision (Halai et al., 2024). Activation magnitude is the magnitude of the BOLD signal

change, operationalised as the 1st-level beta values extracted from each voxel. We

hypothesised that both pTx and ME would recover signal and hence increase activation

magnitude relative to the SESB sequence (which we used as a baseline). Activation precision

is the reliability of the BOLD signal change, analogous to the functional contrast-to-noise

ratio (fCNR) and operationalised as the 1st-level t-values extracted from each voxel. We

hypothesised that MB sequences, with greater effective degrees of freedom, would increase

activation precision relative to single band (SB) sequences. We also had two secondary

hypotheses: first, since ME-ICA denoising removes TE-independent noise, greater activation

precision would be observed in multi-echo denoised data (MEdn) relative to ME data

without denoising; second, that any MB advantage would be due to the increase in the

number of volumes.

Our study focused primarily on univariate effects. However, multivariate analysis

techniques, which exploit variance and covariance between voxels and can accommodate

participant-specific differences in activation patterns (Coutanche, 2013; Davis et al., 2014;

Davis & Poldrack, 2013) are rapidly gaining popularity. These methods frequently rely on the

assumption of good-quality signal across the whole brain (Frisby et al., 2023) and so we

conducted an exploratory analysis, following the method of Haxby et al. (2001), to decode

task condition from the data acquired with each sequence. Finally, we tested for the

presence of slice leakage artefacts in our MB data.

To summarise, our study aimed to compare pTx, ME and MB as methods for

improving sensitivity in ventral temporal and orbitofrontal regions while maintaining image

quality across the rest of the brain.

2. Methods

2.1. Participants

20 healthy native speakers of British English (age range 18-50, mean age 33.45 years, 12

female, 8 male) participated in the study. All were right-handed, had normal or corrected-

to-normal vision, and had no neurological or sensory disorders. All participants gave written

informed consent. The research was approved by a local National Health Service (NHS)

ethics committee (04/Q105/66).

2.2. Stimuli and task

All participants performed a semantic association task and a visual pattern matching task

(hereafter called the "control task") adapted from a previous study (Jung et al., 2017). Each

stimulus consisted of three pictures presented simultaneously (Figure 1). Some pictures

were line drawings taken from the Pyramids and Palm Trees Test (Howard & Patterson,

1992) and some were colour cartoons or photographs taken from the Camel and Cactus Test

(Bozeat et al., 2000). In the semantic task, participants were instructed to indicate which of

the two pictures at the bottom of the screen had the closest semantic relationship to the

picture at the top (hereafter the "probe picture"). In the control task, participants were

instructed to indicate which of two scrambled pictures (generated from the pictures used in

the semantic task) at the bottom of the screen was identical to a scrambled probe. There

were 248 unique picture triplets, so some stimuli were repeated between runs, but no

stimulus was repeated within a run. E-Prime software (Psychology Software Tools Inc.,

Pittsburgh, USA) was used to display stimuli and record responses. Stimuli were rear-

projected onto a screen at the back of the MRI scanner, and observers viewed stimuli

through a mirror mounted to the head coil directly above the eyes.

Figure 1: One block of the semantic association task (left of the arrow) and one block of the
control task (right of the arrow). Each trial consisted of a fixation cross (500 ms) followed by
three pictures presented simultaneously. Participants had to select which of the two
pictures at the bottom was more semantically similar to (in the semantic task) or visually
identical to (in the control task) the picture at the top (the probe picture). Each block lasted
16 s in total. Reprinted with permission from (Halai et al., 2024).
201

The study had a block design with three types of block: semantic, control and rest.

Each task block consisted of four trials. Each trial consisted of a fixation cross presented for

500 ms followed by a stimulus presented for 3500 ms. Each rest block consisted of a fixation

cross presented for 16 s. Each run began 16 s after the start of the MR sequence and then

consisted of 30 blocks presented in the order: semantic, control, semantic, control, rest.

There were five runs per participant, collected in a single session. Accuracy and reaction

time were measured for each trial. Since reaction time is not normally distributed, both

metrics were compared across tasks using Wilcoxon's signed-rank tests and across

sequences using Friedman's nonparametric ANOVA.

2.3. Image acquisition

All images were acquired on a whole-body MAGNETOM Terra 7T MRI (Siemens

Healthcare, Germany). An 8Tx32Rx head coil (Nova Medical, USA) was used to run all

structural and functional imaging sequences.

An MP2RAGE anatomical scan was acquired with the following parameters: 224

sagittal slices (interleaved acquisition), FOV 240 x 225.12 x 240 mm^3, voxel size 0.75 mm

isotropic, TR 4300 ms, inversion times 840 and 2370 ms, TE 1.99 ms, nominal flip angles 5°

and 6°, GRAPPA acceleration factor 3, and duration 8 minutes 50 seconds.

Next, manual B_0-shimming was performed over the volume to be used for EPI. This

was necessary because of a known bug in the scanner software which renders automatic

shims inaccurate. A double-echo steady state (DESS) sequence was used to generate a field

map and the radiographer adjusted shim parameters manually with the aim of reducing the

water linewidth, defined as full width of the spectrum at half height, to below 40 Hz.

However, for some participants the adjustments proved time-consuming and so adjustment

time was capped at 30 minutes from the start of scanning after which the best shim

parameters were adopted. The actual average water linewidth was 42 Hz (standard

deviation = 9 Hz; missing data for 4 participants). Then a dummy pTx-EPI scan was acquired

to trigger the acquisition of subject-specific B_0 and per-channel B_1^+ field maps. Parameters

of this dummy scan were the same as the pTx functional sequence (see Table 1 and below)

except that only one volume was acquired. The brain was divided into 5 slabs along the slice

direction and slab-specific 2-spoke pTx excitation pulses were designed offline. Variable-rate

selective excitation (VERSE; Hargreaves et al., 2004) was applied to reduce specific

absorption rate (SAR) for the pTx sequence only.

There were five functional runs of EPI, one run of each sequence. Key parameters

are given in Table 1. To ensure that our results would be of use to future cognitive

neuroimaging studies, the sequences were designed to represent the way that each

parameter would be used in a typical study. Three echoes were used because this is the

minimum number needed for ME-ICA and because our scanner hardware prevented us from

using four (Halai et al., 2024). An MB factor of 2 was used because previous work suggested

that this MB factor, combined with a GRAPPA factor of 3, would be unlikely to cause slice

leakage (Todd et al., 2016). GRAPPA and phase partial Fourier were used to ensure a

reasonable TE and TR for ME sequences and to ensure that the SE and pTx sequences would

have comparable TRs. However, in-plane acceleration reduces tSNR (Yun & Shah, 2017) and

using *both* GRAPPA and phase partial Fourier is not needed to achieve a reasonable TE for a

single-echo sequence. To facilitate a fair comparison between ME and SE sequences as they

would be used in typical studies, no GRAPPA was used for SE sequences. 2 spokes were used

because, as the number of spokes increases, the pulse length increases and the bandwidth

decreases, and using 2 spokes keeps both those parameters within a reasonable range. 5

slabs were used because, as the number of slabs increases, the time needed for pulse

calculation also increases, and using 5 slabs keeps the calculation time short (~ 5 minutes;

Ding et al., 2022). Due to experimenter error, the flip angle was not set to the Ernst angle

for SE sequences. The possible impact of this error on results is explored in Supplementary

Material 2. Bandwidth of the pTx sequence was selected to match previous work (Ding et

al., 2022). It was necessary to increase the bandwidth for our other sequences in order to

control TE, but doing so reduces tSNR. To facilitate a fair comparison of pTx to the other

sequences as they would be used in typical studies, we did not lower the bandwidth of the

pTx sequence to match the others.

The order of sequences was counterbalanced across participants. The following

parameters were held constant across sequences: 48 axial slices (interleaved acquisition),

FOV 210 x 210 x 210 mm^3 to cover the whole brain in most participants (visual inspection

ensured that the ventral anterior temporal lobe was included in the FOV for all participants

and the FOV was tilted up at the nose to avoid ghosting of the eyes into the temporal lobe),

voxel size 2.5 mm isotropic (no gap), and A-P phase encoding direction. After each run, 5

further volumes were acquired with the phase encoding direction changed to P-A to

facilitate distortion correction during preprocessing.

	SESB	SEMB	MESB	MEMB	pTx

No. echoes	1	1	3	3	1
Multiband factor	1	2	1	2	1
TR (ms)	3020	1510	3020	1510	3000
TE_1 (ms)	25.00	25.00	11.80	11.80	25.00
TE_2 (ms)	-	-	27.05	27.05	-
TE_3 (ms)	-	-	42.30	42.30	-
iPAT type	Off	Off	GRAPPA	GRAPPA	GRAPPA
iPAT factor	Off	Off	3	3	2
Phase partial Fourier	7/8	7/8	7/8	7/8	Off
Nominal flip angle (°)	50	50	78	63	40
Bandwidth (Hz/Px)	2204	2204	2204	2204	1804
Number of volumes acquired	171	340	171	340	172
Number of pulses	1	1	1	1	5 (VERSE 2-spoke pTx pulses; duration 5.12 ms)

Table 1: parameters of each sequence.

2.4. Data analysis

All analysis code is available at https://github.com/slfrisby/7TOptimisation/.

2.4.1. Preprocessing

For ease of data sharing we converted all DICOMs to BIDS format (Gorgolewski et al.,

2016) using *heudiconv* v1.0.0 (Halchenko et al., 2024).

Standard reproducible preprocessing pipelines designed for 3T-fMRI, such as

fMRIprep (Esteban et al., 2019; Gorgolewski et al., 2011; Markiewicz et al., 2024) perform

poorly on 7T-fMRI EPI data. Therefore, the analysis pipeline was split into stages using

different software packages.

Since it is notoriously difficult to perform a good-quality brain extraction on

MP2RAGE data (because of salt-and-pepper noise in the background and cavities), two

pipelines were used. The MP2RAGE T1w (combined) image first had its background noise

removed using O'Brien regularisation (O'Brien et al., 2014) and was then submitted to the

CAT12 pipeline for segmentation (Gaser et al., 2023; in SPM12;

https://www.fil.ion.ucl.ac.uk/spm/). The bias- and global-intensity corrected T1w image

produced was provided to as input to the anatomy pipeline in *fMRIPrep* 21.0.1. TheT1w

image was skull-stripped with a Nipype implementation of the *antsBrainExtraction.sh*

workflow (ANTs 2.3.3; Avants et al., 2009, 2011; https://github.com/ANTsX/ANTs/). Volume-

based spatial normalisation to standard space (MNI152NLin2009cAsym) was performed

through nonlinear registration with *antsRegistration.sh* (ANTs).

Functional preprocessing was performed using in-house code, composed of

functions from AFNI (v.18.3.03; Cox, 1996; Cox & Hyde, 1997; https://afni.nimh.nih.gov/),

FSL (v.5.0; Andersson et al., 2003; Jenkinson et al., 2012; Smith, 2002; Smith et al., 2004;

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/), tedana (v.23.0.1; DuPre et al., 2021; Kundu et al.,

2011, 2013; The tedana Community et al., 2023;

https://tedana.readthedocs.io/en/stable/index.html) and ANTS (v. 2.2.0; Avants et al., 2009,

2011; https://github.com/ANTsX/ANTs/). EPIs were despiked using *3dDespike* (AFNI), slice

timing was corrected to the middle slice using *3dTshift* (AFNI), motion was corrected with

3dvolreg and *3dAllineate* (AFNI) using the first volume of each run as a reference (for ME

datasets, the TE_1 image was aligned and the resulting transform was applied to the TE_2 and

TE_3 images), and skull-stripped using *BET* (FSL) to create a participant-specific brain mask.

For ME datasets only, *tedana* was used to create two timeseries – one in which the

echoes were optimally combined based on T_2* weighting (Posse, 2012), and one in which

the T_2* optimally-combined data were denoised using ICA. *tedana* conducts denoising by

decomposing data using PCA and ICA, classifying components according to whether the

signal scales linearly with TE (as the BOLD signal does), and reconstructing the data using

only BOLD-like components. The brain mask created with *BET* was used as the mask for this

stage.

Next, for all datasets, unwarping was conducted using *topup* and *applytopup* (FSL).

Field displacement maps were calculated using ten volumes (five with A-P phase encoding

direction, extracted from the start of each functional run, and five with P-A phase encoding

direction, collected separately after each run) and the resulting correction was applied to all

images. Finally, the mean EPI for each run was coregistered to the skull-stripped native

structural image using a rigid-body registration with *AntsRegistrationSyN.sh* (ANTS). EPIs

were then transformed into standard space (MNI152NLin2009cAsym) by combining the

transforms from native EPI to native T1 and the transforms from native T1 to standard space

and applying those transforms to the EPIs using *antsApplyTransforms* (ANTS). Images were

smoothed with a 6 mm FWHM Gaussian filter in SPM12 (https://www.fil.ion.ucl.ac.uk/spm/)

for GLM analysis.

A separate functional preprocessing pipeline was used to create images for slice

leakage analysis. This pipeline differed from the main pipeline in the following ways.

Despiking was omitted to avoid the removal of noise. For ME datasets only, rather than run

the full *tedana* workflow, we conducted optimal combination of data from multiple echoes

(but no denoising) using the *t2smap* command using all voxels in the volume (i.e. no brain

mask). All coregistration steps were omitted (the images remained in native EPI space) and

no smoothing was applied.

2.4.2.   1st-level (within-participant) GLM

Data were analysed using the general linear model (GLM) approach implemented in

SPM12 in MATLAB r2019a. We had 5 timeseries of primary interest – standard single-echo

single band (SESB), parallel transmit (pTx), single-echo multiband (SEMB), multi-echo single

band (MESB) and multi-echo multiband (MEMB). For the latter two sequences, data from all

echoes were optimally combined but were not ICA-denoised. We also generated two

timeseries with ME-ICA denoising - multi-echo single band with ICA denoising (MESBdn) and

multi-echo multiband with ICA denoising (MEMBdn) - and two downsampled MB timeseries

created by extracting odd-numbered volumes to match the number of volumes in the single

band timeseries - odd-numbered volumes of single-echo multiband (SEMBodd) and odd-

volumes of multi-echo multiband (MEMBodd).

At the individual subject level, each block of the semantic and control task was

modelled as a boxcar function (resting blocks were modelled implicitly) and these boxcar

functions were subsequently convolved with SPM's difference of gammas haemodynamic

response function. The six motion parameters extracted during preprocessing were used as

regressors of no interest. The micro-time resolution was set as the number of slices (n = 48),

the micro-time onset was set as the reference slice for slice-time correction (n = 24), and the

high-pass filter cutoff was 128 seconds. The same MNI template used during preprocessing

was used as a mask for the analysis. The parameter estimation method was restricted

maximum likelihood estimation (ReML) and serial correlations were accounted for using an

autoregressive AR(1) model during estimation. For univariate analyses the contrast of

interest was greater activation for the semantic task than the control task (S > C). For

exploratory multivariate pattern analysis (MVPA) each block (12 semantic and 12 control)

was modelled individually in order to obtain one beta image per block.

Finally, for the slice leakage analysis, the modelling was rerun on the minimally-

preprocessed timeseries without any brain mask. We obtained both univariate contrasts of

interest (S > C) and beta images per block (for MVPA).

2.4.3.   2nd-level (across-participant) GLM

2.4.3.1.   Region-of-interest (ROI) analysis

Regions of interest (ROIs) were defined based on a large-scale distortion-corrected

3T-fMRI study of the semantic network (Humphreys et al., 2015). For each comparison, ROIs

were analysed only if they overlapped by at least one voxel with the whole-brain contrast of

interest (S > C) summed over all sequences in the comparison.

2.4.3.1.1.   Univariate analysis

Activation magnitude (1st-level beta values) and activation precision (1st-level t-

values) were extracted from each ROI using a publicly-available script, *roi_extract.m*

(https://github.com/MRC-CBU/riksneurotools/blob/master/Util/roi_extract.m). There were

two planned t-tests for activation magnitude (pTx > SESB, ME > SE) and four planned t-tests

for activation precision (MB > SB, MEdn > ME, MBodd > SB, SB > MBodd). For each planned

t-test, results were Bonferroni-corrected for the number of ROIs included.

2.4.3.1.2.   Exploratory multivariate pattern analysis (MVPA)

The input to this analysis was beta images, one image per block (12 semantic and 12

control). For each block and each ROI, a vector v of beta values was created (voxels within

the ROI are numbered arbitrarily):

$$v_{block\ n} = [\beta_{voxel\ 1}, \beta_{voxel\ 2}, \dots \beta_{voxel\ n}]$$

For each ROI, the cosine distance D (1 – cosine similarity; (Halai et al., 2024)) between every

possible pair of blocks was calculated. For example:

$$D_{block\ 1, block\ 2} = 1 - \left(\frac{v_{block\ 1} \cdot v_{block\ 2}}{\|v_{block\ 1}\| \, \|v_{block\ 2}\|} \right)$$

For each ROI, MVPA performance was operationalised as the mean between-task distance

(over every possible pair of blocks of different tasks) minus the mean within-task distance

(over every possible pair of blocks of the same task). Where block S is a block of the

semantic task and block C is a block of the control task, and n_{blocks} is the total number of

blocks of each task (12):

$$MVPA\ metric$$

386
$$= \left(\frac{\sum D_{block\ S1,\ C1},\ D_{block\ S1,\ C2},\ ...,\ D_{block\ Sn,\ block\ Cn}}{n_{blocks}{}^2} \right)$$
$$- \left(\frac{\sum D_{block\ S1,\ block\ S2},\ ...,\ D_{block\ Sn-1,\ block\ Sn},\ D_{block\ C1,\ block\ C2},\ ...,\ D_{block\ Cn-1,\ block\ Cn}}{2(n_{blocks}{}^2)} \right)$$

If a region is encoding information about the task, we would expect mean within-task

similarity to be large and mean between-task similarity to be small, so the MVPA metric

would take a large negative value. If a region is not encoding information about the task, we

would expect both within-task and between-task similarity to be zero and the difference

between them to be negligible (Halai et al., 2024).

Paired t-tests were used to compare the MVPA metric across sequences (all planned

t-tests described in the univariate ROI analysis were conducted; Haxby et al., 2001).

2.4.3.2.   Whole-brain analysis

All the above contrasts were assessed at the whole-brain level using t-tests (for

comparing pTx and SESB) or using random-effects ANOVAs with one-sample t-tests on the

summary statistic (for the three factorial designs: varying echo and band (SESB, SEMB, MESB

and MEMB); varying denoising and band (MESBdn, MEMBdn, MESB and MEMB); and

varying echo and downsampled band (SEMBodd, MEMBodd, SEMB and MEMB)). The

ANOVAs were conducted using a publicly-available script, *batch_spm_anova.m*

(https://github.com/MRC-CBU/riksneurotools/blob/master/SPM/batch_spm_anova.m). The

group t-maps were assessed for significance by using a voxel-height threshold of p < 0.001

to define clusters and then a cluster-defining family-wise-error corrected threshold of p <

0.05 for statistical inference.

2.4.3.3.   Slice leakage analysis

The group-level, whole-brain contrast of interest (S > C) in standard space was

inspected and the coordinates of the peak t-values of the top 5 clusters were identified.

These coordinates were back-projected to obtain 5 sets of corresponding coordinates in

each participant's native EPI space (hereafter "seeds", labelled A). Next, voxels to which

signal might be warped were identified as artefact locations based on phase shift (FOV/2,

labelled B) and, in the MEMB data, GRAPPA (labelled Ag) and phase shift plus GRAPPA

(labelled Bg). A spherical ROI, 4 voxels in radius, was defined around each seed location and

possible artefact location using a modified version of the scripts developed for McNabb et

al. (2020; https://github.com/DrMichaelLindner/MAP4SL/; our version available at

https://github.com/slfrisby/7TOptimisation/).

Both univariate (McNabb et al., 2020) and multivariate (Halai et al., 2024) slice

leakage tests were conducted. Activation magnitude was extracted from minimally-

preprocessed data, and, for the multivariate analysis, the difference between mean within-

task similarity and mean between-task similarity was calculated as in the ROI analysis.

Paired t-tests were conducted for each seed and artefact location between each sequence

of interest (SEMB and MEMB) and a control sequence. For artefact locations based on phase

shift (B), the control sequence was the corresponding SB sequence (SESB for SEMB and

MESB for MEMB). For artefact locations based on GRAPPA, the control sequence was the

corresponding SE sequence, because SE sequences were collected without GRAPPA (SEMB

for MEMB). Results were Bonferroni-corrected for the number of peaks (n = 5) and the

number of possible artefact regions (n = 1 for SEMB and n = 3 for MEMB).

3. Results

3.1. Excluded participants

Two participants were excluded because of excessive head motion (this was defined

by calculating, for each participant, the percentage of volumes per run with absolute

translation values of over 2 mm or absolute rotation values over 1°, averaging these

percentages over runs, and excluding any participant whose mean percentage was greater

than 2 standard deviations above the mean percentage across participants). One participant

was excluded because of technical problems during data acquisition which meant that the

pTx run failed. All subsequent analyses were conducted on the remaining 17 participants.

3.2. Behavioural results

The 17 participants had good performance on both tasks and, importantly, there

were no significant differences between the 5 sequences in terms of accuracy (Friedman's χ^2

= 5.26, p = 0.26) or reaction time (Friedman's χ^2 = 7.20, p = 0.16). The semantic and control

tasks did not differ reliably in accuracy (Wilcoxon's z = 0; p = 0.06) or reaction time

(Wilcoxon's z = 7; p = 1.00).

3.3. Region-of-interest analysis

ROIs that overlapped with the whole-brain contrast of interest (S > C) for at least

one of the five sequences are shown in Figure 2.

Figure 2: Regions of interest, taken from a meta-analysis of semantic tasks by (Humphreys et
al., 2015). All spheres are 8 mm in radius. Regions of interest are overlaid on the
MNI152NLin2009cAsym template. LTP = left temporal pole; LvATL = left ventral anterior
temporal lobe; RITG = right inferior temporal gyrus; LFP = left frontal pole; LmMTG = left
medial middle temporal gyrus; LpMTG = left posterior middle temporal gyrus; LIFGpt = left
inferior temporal gyrus pars triangularis.
457
458

	LTP	LvATL	RITG	LFP	LmMTG	LpMTG	LIFGpt
Activation magnitude							
pTx > SESB	-	0.2488	0.0240*	-	-	0.0042**	0.4114
ME > SE	0.0502	0.0001**	0.0333*	0.2951	0.0092*	0.0003*	0.4155
Activation precision							
MB > SB	0.4569	0.0007**	0.0019**	0.0989	0.2187	0.0266*	0.0016**
MEdn > ME	0.3117	<0.0001**	<0.0001**	0.0014**	0.1363	<0.0001**	<0.0001**
MBodd > SB	0.4225	0.1057	0.0481*	-	0.5697	0.1879	0.3300
MVPA							
pTx > SESB	-	0.8569	0.3280	-	-	0.1529	0.9179
ME > SE	0.0096*	0.0063**	0.1612	0.0655	0.0465*	0.0024**	0.2508
MB > SB	0.2455	0.1417	0.0667	0.3021	0.3949	0.2693	0.5207
MEdn > ME	0.0027**	<0.0001**	<0.0001**	0.0023**	0.0001**	0.0012**	0.0007**
SB > MB	0.7416	0.4570	0.7222	-	0.3803	0.6716	0.3242
MBodd > SB	0.2584	0.5430	0.2778	-	0.6197	0.3284	0.6758

Table 2: p-values for all t-tests within regions of interest (ROIs) based on the semantic

network. * = p < 0.05, ** = p < 0.05 (Bonferroni-corrected for the number of ROIs), - = ROI

does not overlap by at least one voxel with the whole-brain contrast of interest (S > C)

summed over all sequences in the comparison, LTP = left temporal pole, LvATL = left ventral

anterior temporal lobe, RITG = right inferior temporal gyrus, LFP = left frontal pole, LmMTG

= left medial middle temporal gyrus, LpMTG = left posterior middle temporal gyrus, LIFGpt =

left inferior temporal gyrus pars triangularis.

3.3.1. Univariate analysis

Table 2 shows the p-values for all planned t-contrasts. pTx provided significantly

better activation magnitude than SESB in the left posterior middle temporal gyrus (LpMTG).

ME sequences provided significantly better activation magnitude than SE sequences in the

left ventral anterior temporal lobe (LvATL).

MB sequences offered significantly better activation precision than SB sequences in

the LvATL, right inferior temporal gyrus (RITG) and left inferior frontal gyrus pars triangularis

(LIFGpt). MEdn sequences offered significantly better activation precision than ME in the

LvATL, RITG, LpMTG, LIFGpt and left frontal pole (LFP). There was no significant difference in

either direction between MBodd sequences and SB sequences (see Supplementary Table 1

for detailed results, plus reverse contrasts). We also found no significant interaction

between ME and MB.

3.3.2. Exploratory MVPA

Table 2 also shows the p-values for t-tests comparing our MVPA metric between

sequences. The ME sequences produced significantly better performance than SE sequences

in the LvATL and LpMTG. MEdn sequences produced better performance than ME

sequences in every ROI. All other comparisons failed to reach significance.

3.4. Whole-brain analysis

Figure 3 shows the results for activation magnitude. Figure 3A shows a single cluster,

within the right fusiform gyrus and lateral inferior occipital cortex, that showed greater

activation magnitude with pTx than with SESB (pTx > SESB). Figure 3B shows the main effect

of ME over SE, featuring clusters in the fusiform and inferior temporal gyri bilaterally plus

the left orbitofrontal cortex (cluster and peak information is provided in Supplementary

Table 2).

Figure 3: Effects on activation magnitude: (A) effect of parallel transmit (pTx > SESB); (B)
effect of echo (ME > SE). Results are cluster-corrected at p < 0.05 based on an uncorrected
voxel threshold of p < 0.001 and are overlaid on the MNI152NLin2009cAsym template.
499

Figure 4 shows the results for activation precision. Both the main effect of MB over

SB and the main effect of ME-ICA denoising over ME without ICA denoising extended down

the temporal lobe and included frontal regions (Supplementary Table 2). There was no

significant difference between downsampled MB sequences and SB sequences in either

direction.

Figure 4: Effects on activation precision: (A) effect of multiband (MB > SB); (B) effect of ME-
ICA denoising (MEdn > ME). Results are cluster-corrected at p < 0.05 based on an

uncorrected voxel threshold of p < 0.001 and are overlaid on the MNI152NLin2009cAsym
template.
510

511   To summarise, these results aligned with results from our ROI analyses – ME

512 increased activation magnitude and both MB and ME-ICA denoising increased activation

513 precision (see Supplementary Figures 1-6 for detailed results, plus reverse contrasts).

514
515
3.5. Slice leakage analysis

Seed and possible artefact locations for an example participant are shown in Figure

5A. Figure 5B shows activation magnitude for possible artefact locations associated with one

peak for all participants in each MB sequence and its corresponding sequences. Figure 5C

shows MVPA performance. Violin plots for all other peaks are shown in Supplementary

Figure 7. t-tests were conducted at each artefact location, but no comparison reached

statistical significance (p-values are shown in Supplementary Table 3). t-tests were not

conducted at seed locations – if a difference were discovered, this would be attributable to

the MB sequence rather than to leakage of signal from the seed into a different location.

We therefore concluded that there was no evidence for slice leakage in either of our

MB datasets.

Figure 5: Slice leakage analysis. (A) Seed and possible artefact locations for a single
participant. The ROIs (radius 4 voxels) indicate the seed location (blue), possible artefact
location based on phase shift (green), possible artefact location based on GRAPPA (red), and
possible artefact location based on phase shift and GRAPPA (yellow) in native EPI space.
Informed consent was obtained from the participant for this image to be published. (B)
Group mean activation magnitude within each possible artefact location corresponding to

534   the first seed for each MB sequence and the corresponding control sequences (plots for
535   other seeds are shown in Supplementary Figure 7; statistics for all seeds are shown in
536   Supplementary Table 3). Each point within a violin plot indicates the mean for one
537   participant (n = 17). (C) Group mean MVPA distance within each sphere for each MB
538   sequence and the corresponding control sequences.
539
540
4. Discussion

The tSNR of fMRI varies across the brain. This is especially evident in ventral anterior

temporal and orbitofrontal regions, which are located next to the air-filled sinuses and are

therefore affected by both B_0 and B_1^+ magnetic field inhomogeneity (Devlin et al., 2000;

Gras et al., 2019; Halai et al., 2014, 2015, 2024; Uğurbil, 2018; van der Kolk et al., 2013; Wu

et al., 2018). This study is the first 7T-fMRI study systematically comparing pTx, ME and MB

as practical methods for improving sensitivity in these regions while maintaining sensitivity

across the brain. We found that pTx improved activation magnitude in posterior temporal

and occipital regions. ME, however, resulted in improved activation magnitude extending

down the temporal lobe and including inferior frontal regions. Both MB and ME-ICA

denoising resulted in improved activation precision in the same areas. In an exploratory

analysis we found that ME and ME-ICA improved MVPA performance but MB did not. No

slice leakage artefacts were associated with our MB sequences.

Although pTx produced better activation magnitude in posterior temporal and

occipital regions than the baseline (SESB) sequence without compromising activation

magnitude elsewhere, activation magnitude in anterior temporal areas was comparable for

both sequences. These results replicated those of Ding et al. (2022) who failed to find

improved task contrast in anterior temporal regions with pTx in spite of improved tSNR in

the resting state. Note that, despite visible effects of magnetic field inhomogeneity on the

EPIs, even the baseline sequence was able to identify semantic activity extending into

ventral anterior temporal regions. This is shown in Figure 6 in which the contour of the

whole-brain contrast of interest is overlaid on the mean EPI across all participants (detailed

contrast maps are shown in Supplementary Figures 1 and 3). It is possible that this finding

reflects the increased sensitivity of 7T-fMRI compared to 3T-fMRI, where semantic activity is

rarely observed without using a method such as ME (Halai et al., 2014, 2015, 2024) or spin-

echo (Binney et al., 2010; Embleton et al., 2010). That a standard sequence can detect signal

in these regions might come as a surprise to many neuroimaging researchers as 7T-fMRI is

not only associated with increased sensitivity, but also with exacerbated magnetic

susceptibility artefacts. This finding should therefore embolden researchers to use 7T fMRI

for experimental designs that are difficult to conduct at a lower field strength - for example,

studies of special populations, who may benefit from shorter scan times (Cope et al., 2023),

or studies requiring high spatial specificity (Marques & Norris, 2018). We do acknowledge

that the way we have utilised 7T fMRI in this study will not be suitable for, for example,

studies that require ultra-high resolution (in our case, scanner hardware constrained voxel

size when combined with a short first echo time).

577
578

Figure 6: Contrast despite magnetic field inhomogeneity. (A) Mean EPI across all participants

for the baseline sequence (SESB) registered to MNI152NLin2009cAsym space. (B) Selected

regions of interest overlaid on the mean EPI image. (C) Contour of the contrast of interest (S
> C) for the baseline sequence. (D) Contour of the contrast of interest (S > C) for the MEMB
sequence. Results are cluster-corrected at $p < 0.05$ based on an uncorrected voxel threshold
of $p < 0.001$.
585

The factorial design enabled us to disentangle the effects of ME and MB. As
hypothesised, ME improved activation magnitude and these effects were localised to
inferior temporal and orbitofrontal areas. Adverse effects on activation magnitude were
negligible - although ME reduced activation precision in two very small clusters
(Supplementary Figure 4), these were not within areas that the semantic task is known to
recruit (Jung et al., 2017). These results demonstrated that having one very short echo is a
successful method of increasing activation magnitude in areas where T_2* is particularly short
(Halai et al., 2014, 2015, 2024; Jung et al., 2017). ME also opens up the opportunity for
sophisticated denoising such as *tedana* (DuPre et al., 2021; Kundu et al., 2011, 2013; The
tedana Community et al., 2023), which improved activation precision in agreement with
previous findings (Amemiya et al., 2019; Gonzalez-Castillo et al., 2016). Small clusters in
which ME-ICA increased activation magnitude (Supplementary Figure 5) or decreased
activation precision (Supplementary Figure 6) were not within regions recruited by the task
(Jung et al., 2017).

As hypothesised, MB improved activation precision in areas of inhomogeneity and in
agreement with other work (Puckett et al., 2018). Supplementary Figure 4 shows two very
small clusters in which MB reduced activation precision, but which lie outside regions
recruited by the task (Jung et al., 2017). After downsampling the MB timeseries to match
the number of volumes in the SB timeseries, there was no longer any difference between
the SB and MB sequences; this suggests that it is the increase in the number of volumes that
accounts for the benefits of MB. Additionally, there was no evidence for signal leakage into
the simultaneously-acquired slice location (McNabb et al., 2020; Todd et al., 2016), at least
for our modest multiband factor of 2. We note that MB and ME seem to have independent
effects as the interaction was not significant within any ROI. However, ME seems to offer
larger activation magnitude (Figure 6) and MB offers better activation precision. MB can
help reduce the longer TR associated with ME and may decrease noise aliasing in studies
using short TRs (~1s).

Although this study focused primarily on univariate effects, we also identified

improvements to multivariate metrics that will be of interest to researchers using

increasingly-popular MVPA tools (Frisby et al., 2023). ME was found to improve MVPA

performance compared to SE sequences and, in turn, MEdn sequences performed

significantly better than ME sequences without denoising in every region of interest (with

no detrimental effects). Future studies employing multivariate methods should strongly

consider using both ME acquisitions and ME-ICA preprocessing.

This study provides evidence supporting use of ME and/or MB sequences to detect

signal in regions prone to susceptibility artefacts. It should be noted that, due to

experimenter error, the flip angles for the SE sequences were not set to the Ernst angle,

which could in theory have accounted for the ME benefit that we observed. However,

acquisition of additional SE data with the flip angle set to the Ernst angle indicated that,

although use of a suboptimal flip angle compromised tSNR, there was no consistent effect

on any of our univariate or multivariate effects of interest (methods and results of this

investigation are described in detail in Supplementary Material 2). From this we conclude

that the difference in flip angle is not sufficient to explain the ME advantage, although

future studies should seek to validate these results using the Ernst angle for all sequences.

It is also important to note that individual 7T MRI scanners have their own hardware

and software limitations; this means that each site must optimise and test feasible

parameters locally. For example, our system did not allow us to take advantage of the

higher spatial resolution offered by 7T-fMRI while retaining an adequately short first echo

for our ME sequences. Puckett et al. (2018) used comparable voxel sizes but had a shorter

first TE than us (9.9 ms vs our 11.8 ms); Miletić et al. (2020) had higher resolution (1.6 mm

vs our 2.5 mm isotropic) and a shorter first echo of 9.66 ms. While we could in principle

have achieved a smaller voxel size for our SE sequences, this would have been accompanied

either by a narrower FOV or, if we were to maintain the FOV by increasing the number of

slices, by a longer TR (which we wished to keep constant across sequences). Additionally,

our 7T Terra pTx system imposes very conservative SAR limits (1W per channel, 8W total)

for pTx scans but permits 20W total (i.e. 2.5W per channel) for circularly polarised mode

(CP-mode or "TrueForm") scans (equivalent to single transmit). We therefore had to use

VERSE-modified pTx pulses which are known to be more sensitive to B_0 inhomogeneity. We

also acknowledge that pTx can be combined with ME or MB (Ding et al., 2023; Wu et al.,

2016). Although we used the pTx head coil for all scans, we were not able to create

sequences that combined pTx with ME or MB because the software to support this was not

implemented on our scanner at the time of running this study. Future studies should seek to

test the potential benefits of these combinations. It is possible that studies seeking to test

more extreme parameter manipulations (numbers of echoes, multiband factors, or pTx

approaches) than we have employed may reach different conclusions to us. For example,

Todd et al. (2016) demonstrated that, although multiband factors of 3 or greater can

sometimes be associated with greater BOLD sensitivity than the multiband factor of 2 that

we have used, higher multiband factors are also associated with a risk of slice leakage. In

summary, rather than identifying the limits of what is possible (as others have done), we

aim to offer an accessible framework that will enable researchers to leverage 7T-fMRI for

investigating the functional roles of regions affected by susceptibility artefacts despite the

hardware and software constraints of individual scanners.

5. Conclusion

In this study we compared pTx, ME and MB as methods of improving sensitivity in

temporal and frontal areas prone to magnetic susceptibility artefacts. Both pTx and ME

improved activation magnitude, but only ME showed improvements in artefact-prone

regions. MB and ME-ICA improved activation precision in these areas. Exploratory results

suggested that both ME and ME-ICA may also benefit MVPA. We demonstrated that a multi-

echo, multiband sequence running on the 8Tx32Rx pTx head coil (in CP mode) can detect

signal in susceptible regions while maintaining sensitivity across the whole brain and is

therefore a versatile choice for future studies using high field strength and investigating the

functional roles of ventral temporal and/or orbitofrontal cortex (Binney et al., 2016;

Borghesani et al., 2016; Devereux et al., 2018; Duncan, 2010; DuPre et al., 2016; Fernandez

et al., 2017; Lambon Ralph et al., 2017; Zahn et al., 2007).

Data and code availability: Data will be made publicly available upon peer review and

acceptance. Code is publicly available at: https://github.com/slfrisby/7TOptimisation/.

CRediT statement: Saskia L. Frisby: Conceptualisation, Methodology, Investigation, Formal

Analysis, Writing – Original Draft, Writing – Review & Editing, Visualisation; Marta M.

Correia: Conceptualisation, Methodology, Writing – Review & Editing, Funding Acquisition;

Minghao Zhang: Methodology, Writing – Review & Editing; Christopher T. Rodgers:

Methodology, Writing – Review & Editing, Supervision; Timothy T. Rogers: Writing – Review

& Editing, Supervision; Matthew A. Lambon Ralph: Writing – Review & Editing, Supervision,

Funding Acquisition; Ajay D. Halai: Conceptualisation, Methodology, Formal Analysis,

Writing – Review & Editing, Supervision, Funding Acquisition.

Funding: This work was supported by an MRC Unit grant (SUAG/019 G116768) to M.M.C.,

an MRC PhD studentship (MR N013433-1) to M.Z., an MRC programme grant

(MR/R023883/1) and intramural funding (MC_UU_00005/18) to M.A.L.R, and an MRC

Career Development Award (MR/V031481/1) to A.D.H. This work was also supported by the

NIHR Cambridge Biomedical Research Centre (NIHR203312) and an MRC Clinical Research

Infrastructure Award for 7T (MR/M008983/1). The views expressed are those of the authors

and not necessarily those of the NIHR or the Department of Health and Social Care.

Competing interests: C.T.R. receives research support from Siemens.

References

Adriany, G., Van de Moortele, P.-F., Wiesinger, F., Moeller, S., Strupp, J. P., Andersen, P., Snyder, C.,
Zhang, X., Chen, W., Pruessmann, K. P., Boesiger, P., Vaughan, T., & Uğurbil, K. (2005).
Transmit and receive transmission line arrays for 7 Tesla parallel imaging. *Magnetic*
Resonance in Medicine, *53*(2), 434–445. https://doi.org/10.1002/mrm.20321
Amemiya, S., Yamashita, H., Takao, H., & Abe, O. (2019). Integrated multi-echo denoising strategy
improves identification of inherent language laterality. *Magnetic Resonance in Medicine*,
81(5), 3262–3271. https://doi.org/10.1002/mrm.27620
Andersson, J. L. R., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-
echo echo-planar images: Application to diffusion tensor imaging. *NeuroImage*, *20*(2), 870–
888. https://doi.org/10.1016/S1053-8119(03)00336-7
Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible
evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage*,
54(3), 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025
Avants, B. B., Tustison, N., & Johnson, H. (2009). *Advanced Normalization Tools (ANTS)*.
Barth, M., Breuer, F., Koopmans, P. J., Norris, D. G., & Poser, B. A. (2016). Simultaneous multislice
(SMS) imaging techniques. *Magnetic Resonance in Medicine*, *75*(1), 63–81.
https://doi.org/10.1002/mrm.25897
Bhavsar, S., Zvyagintsev, M., & Mathiak, K. (2014). BOLD sensitivity and SNR characteristics of
parallel imaging-accelerated single-shot multi-echo EPI for fMRI. *NeuroImage*, *84*, 65–75.
https://doi.org/10.1016/j.neuroimage.2013.08.007

Binney, R. J., Embleton, K. V., Jefferies, E., Parker, G. J. M., & Lambon Ralph, M. A. (2010). The ventral
and inferolateral aspects of the anterior temporal lobe are crucial in semantic memory:
Evidence from a novel direct comparison of distortion-corrected fMRI, rTMS, and semantic
dementia. *Cerebral Cortex*, *20*(11), 2728–2738. https://doi.org/10.1093/cercor/bhq019
Binney, R. J., Hoffman, P., & Lambon Ralph, M. A. (2016). Mapping the Multiple Graded
Contributions of the Anterior Temporal Lobe Representational Hub to Abstract and Social
Concepts: Evidence from Distortion-corrected fMRI. *Cerebral Cortex*, *26*(11), 4227–4241.
https://doi.org/10.1093/cercor/bhw260
Borghesani, V., Pedregosa, F., Buiatti, M., Amadon, A., Eger, E., & Piazza, M. (2016). Word meaning in
the ventral visual path: A perceptual to conceptual gradient of semantic coding.
NeuroImage, *143*, 128–140. https://doi.org/10.1016/j.neuroimage.2016.08.068
Boyacioğlu, R., Schulz, J., Koopmans, P. J., Barth, M., & Norris, D. G. (2015). Improved sensitivity and
specificity for resting state and task fMRI with multiband multi-echo EPI compared to multi-
echo EPI at 7 T. *NeuroImage*, *119*, 352–361.
https://doi.org/10.1016/j.neuroimage.2015.06.089
Bozeat, S., Lambon Ralph, M. A., Patterson, K., Garrard, P., & Hodges, J. R. (2000). Non-verbal
semantic impairment in semantic dementia. *Neuropsychologia*, *38*(9), 1207–1215.
https://doi.org/10.1016/S0028-3932(00)00034-8
Caballero-Gaudes, C., Moia, S., Panwar, P., Bandettini, P. A., & Gonzalez-Castillo, J. (2019). A
deconvolution algorithm for multi-echo functional MRI: Multi-echo Sparse Paradigm Free
Mapping. *NeuroImage*, *202*, 116081. https://doi.org/10.1016/j.neuroimage.2019.116081
Cohen, A. D., Nencka, A. S., Lebel, R. M., & Wang, Y. (2017). Multiband multi-echo imaging of
simultaneous oxygenation and flow timeseries for resting state connectivity. *PLOS ONE*,
12(3), e0169253. https://doi.org/10.1371/journal.pone.0169253
Cohen, A. D., Nencka, A. S., & Wang, Y. (2018). Multiband multi-echo simultaneous ASL/BOLD for
task-induced functional MRI. *PLOS ONE*, *13*(2), e0190427.
https://doi.org/10.1371/journal.pone.0190427
Cope, T. E., Sohoglu, E., Peterson, K. A., Jones, P. S., Rua, C., Passamonti, L., Sedley, W., Post, B.,
Coebergh, J., Butler, C. R., Garrard, P., Abdel-Aziz, K., Husain, M., Griffiths, T. D., Patterson,
K., Davis, M. H., & Rowe, J. B. (2023). Temporal lobe perceptual predictions for speech are
instantiated in motor cortex and reconciled by inferior frontal cortex. *Cell Reports*, *42*(5).
https://doi.org/10.1016/j.celrep.2023.112422
Coutanche, M. N. (2013). Distinguishing multi-voxel patterns and mean activation: Why, how, and
what does it tell us? *Cognitive, Affective, & Behavioral Neuroscience*, *13*(3), 667–673.
https://doi.org/10.3758/s13415-013-0186-2
Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance
neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173.
https://doi.org/10.1006/cbmr.1996.0014
Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data. *NMR in
Biomedicine*, *10*(4–5), 171–178. https://doi.org/10.1002/(SICI)1099-
1492(199706/08)10:4/5<171::AID-NBM453>3.0.CO;2-L
Davis, T., LaRocque, K. F., Mumford, J. A., Norman, K. A., Wagner, A. D., & Poldrack, R. A. (2014).
What do differences between multi-voxel and univariate analysis mean? How subject-,
voxel-, and trial-level variance impact fMRI analysis. *NeuroImage*, *97*, 271–283.
https://doi.org/10.1016/j.neuroimage.2014.04.037
Davis, T., & Poldrack, R. A. (2013). Measuring neural representations with fMRI: Practices and
pitfalls. *Annals of the New York Academy of Sciences*, *1296*(1), 108–134.
https://doi.org/10.1111/nyas.12156
Demetriou, L., Kowalczyk, O. S., Tyson, G., Bello, T., Newbould, R. D., & Wall, M. B. (2018). A
comprehensive evaluation of increasing temporal resolution with multiband-accelerated

766 protocols and effects on statistical outcome measures in fMRI. *NeuroImage*, *176*, 404–416.
767 https://doi.org/10.1016/j.neuroimage.2018.05.011
768 Deniz, F., Nunez-Elizalde, A. O., Huth, A. G., & Gallant, J. L. (2019). The Representation of Semantic
769 Information Across Human Cerebral Cortex During Listening Versus Reading Is Invariant to
770 Stimulus Modality. *The Journal of Neuroscience*, *39*(39), 7722–7736.
771 https://doi.org/10.1523/JNEUROSCI.0675-19.2019
772 Devereux, B. J., Clarke, A., & Tyler, L. K. (2018). Integrated deep visual and semantic attractor neural
773 networks predict fMRI pattern-information along the ventral object processing pathway.
774 *Scientific Reports*, *8*(1), 10636. https://doi.org/10.1038/s41598-018-28865-1
775 Devlin, J. T., Russell, R. P., Davis, M. H., Price, C. J., Wilson, J., Moss, H. E., Matthews, P. M., & Tyler, L.
776 K. (2000). Susceptibility-induced loss of signal: Comparing PET and fMRI on a semantic task.
777 *NeuroImage*, *11*(6), 589–600. https://doi.org/10.1006/nimg.2000.0595
778 Ding, B., Dragonu, I., Rua, C., Carlin, J. D., Halai, A. D., Liebig, P., Heidemann, R., Correia, M. M., &
779 Rodgers, C. T. (2022). Parallel transmit (pTx) with online pulse design for task-based fMRI at
780 7T. *Magnetic Resonance Imaging*, *93*, 163–174. https://doi.org/10.1016/j.mri.2022.07.003
781 Ding, B., Williams, S. N., Zhang, M., Herrler, J., Liebig, P., Dragonu, I., Neji, R., Rodgers, C. T., & Porter,
782 D. A. (2023). *Optimisation of pulse-specific phase schedule reduces peak RF amplitude in*
783 *multiband parallel-transmit pulses*. ISMRM & ISMRT Annual Meeting & Exhibition, Toronto,
784 Canada.
785 Dipasquale, O., Sethi, A., Laganà, M. M., Baglio, F., Baselli, G., Kundu, P., Harrison, N. A., &
786 Cercignani, M. (2017). Comparing resting state fMRI de-noising approaches using multi- and
787 single-echo acquisitions. *PLOS ONE*, *12*(3), e0173289.
788 https://doi.org/10.1371/journal.pone.0173289
789 Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: Mental programs for
790 intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179.
791 https://doi.org/10.1016/j.tics.2010.01.004
792 DuPre, E., Luh, W.-M., & Spreng, R. N. (2016). Multi-echo fMRI replication sample of
793 autobiographical memory, prospection and theory of mind reasoning tasks. *Scientific Data*,
794 *3*(1), 160116. https://doi.org/10.1038/sdata.2016.116
795 DuPre, E., Salo, T., Ahmed, Z., Bandettini, P. A., Bottenhorn, K. L., Caballero-Gaudes, C., Dowdle, L. T.,
796 Gonzalez-Castillo, J., Heunis, S., Kundu, P., Laird, A. R., Markello, R., Markiewicz, C. J., Moia,
797 S., Staden, I., Teves, J. B., Uruñuela, E., Vaziri-Pashkam, M., Whitaker, K., & Handwerker, D.
798 A. (2021). TE-dependent analysis of multi-echo fMRI with tedana. *Journal of Open Source*
799 *Software*, *6*(66), 3669. https://doi.org/10.21105/joss.03669
800 Embleton, K. V., Haroon, H. A., Morris, D. M., Lambon Ralph, M. A. L., & Parker, G. J. M. (2010).
801 Distortion correction for diffusion-weighted MRI tractography and fMRI in the temporal
802 lobes. *Human Brain Mapping*, *31*(10), 1570–1587. https://doi.org/10.1002/hbm.20959
803 Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D.,
804 Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R.
805 A., & Gorgolewski, K. J. (2019). fMRIPrep: A robust preprocessing pipeline for functional MRI.
806 *Nature Methods*, *16*(1), 111–116. https://doi.org/10.1038/s41592-018-0235-4
807 Evans, J. W., Kundu, P., Horovitz, S. G., & Bandettini, P. A. (2015). Separating slow BOLD from non-
808 BOLD baseline drifts using multi-echo fMRI. *NeuroImage*, *105*, 189–197.
809 https://doi.org/10.1016/j.neuroimage.2014.10.051
810 Feinberg, D. A., Moeller, S., Smith, S. M., Auerbach, E., Ramanna, S., Glasser, M. F., Miller, K. L.,
811 Ugurbil, K., & Yacoub, E. (2010). Multiplexed Echo Planar Imaging for Sub-Second Whole
812 Brain FMRI and Fast Diffusion Imaging. *PLOS ONE*, *5*(12), e15710.
813 https://doi.org/10.1371/journal.pone.0015710
814 Fernandez, B., Leuchs, L., Sämann, P. G., Czisch, M., & Spoormaker, V. I. (2017). Multi-echo EPI of
815 human fear conditioning reveals improved BOLD detection in ventromedial prefrontal
816 cortex. *NeuroImage*, *156*, 65–77. https://doi.org/10.1016/j.neuroimage.2017.05.005

Frisby, S. L., Halai, A. D., Cox, C. R., Lambon Ralph, M. A., & Rogers, T. T. (2023). Decoding semantic
representations in mind and brain. *Trends in Cognitive Sciences*, *27*(3), 258–281.
https://doi.org/10.1016/j.tics.2022.12.006
Gaser, C., Dahnke, R., Thompson, P. M., Kurth, F., Luders, E., & Alzheimer's Disease Neuroimaging
Initiative. (2023). *CAT − A computational anatomy toolbox for the analysis of structural MRI*
data (p. 2022.06.11.495736). bioRxiv. https://doi.org/10.1101/2022.06.11.495736
Gilmore, A. W., Agron, A. M., González-Araya, E. I., Gotts, S. J., & Martin, A. (2022). A comparison of
single- and multi-echo processing of functional MRI data during overt autobiographical
recall. *Frontiers in Neuroscience*, *16*, 854387. https://doi.org/10.3389/fnins.2022.854387
Gonzalez-Castillo, J., Panwar, P., Buchanan, L. C., Caballero-Gaudes, C., Handwerker, D. A., Jangraw,
D. C., Zachariou, V., Inati, S., Roopchansingh, V., Derbyshire, J. A., & Bandettini, P. A. (2016).
Evaluation of multi-echo ICA denoising for task based fMRI studies: Block designs, rapid
event-related designs, and cardiac-gated fMRI. *NeuroImage*, *141*, 452–468.
https://doi.org/10.1016/j.neuroimage.2016.07.049
Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P., Flandin, G., Ghosh, S.
S., Glatard, T., Halchenko, Y. O., Handwerker, D. A., Hanke, M., Keator, D., Li, X., Michael, Z.,
Maumet, C., Nichols, B. N., Nichols, T. E., Pellman, J., … Poldrack, R. A. (2016). The brain
imaging data structure, a format for organizing and describing outputs of neuroimaging
experiments. *Scientific Data*, *3*(1), 160044. https://doi.org/10.1038/sdata.2016.44
Gorgolewski, K. J., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., & Ghosh, S.
S. (2011). Nipype: A flexible, lightweight and extensible neuroimaging data processing
framework in python. *Frontiers in Neuroinformatics*, *5*.
https://doi.org/10.3389/fninf.2011.00013
Gras, V., Poser, B. A., Wu, X., Tomi-Tricot, R., & Boulant, N. (2019). Optimizing BOLD sensitivity in the
7T Human Connectome Project resting-state fMRI protocol using plug-and-play parallel
transmission. *NeuroImage*, *195*, 1–10. https://doi.org/10.1016/j.neuroimage.2019.03.040
Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C. E., Zsoldos,
E., Ebmeier, K. P., Filippini, N., Mackay, C. E., Moeller, S., Xu, J., Yacoub, E., Baselli, G.,
Ugurbil, K., Miller, K. L., & Smith, S. M. (2014). ICA-based artefact removal and accelerated
fMRI acquisition for improved resting state network imaging. *NeuroImage*, *95*, 232–247.
https://doi.org/10.1016/j.neuroimage.2014.03.034
Hagberg, G. E., Indovina, I., Sanes, J. N., & Posse, S. (2002). Real-time quantification of T2* changes
using multiecho planar imaging and numerical methods. *Magnetic Resonance in Medicine*,
48(5), 877–882. https://doi.org/10.1002/mrm.10283
Halai, A. D., Henson, R. N., Finoia, P., & Correia, M. M. (2024). *Comparing the effect of multi gradient*
echo and multi band fMRI during a semantic task (p. 2024.03.20.585909). bioRxiv.
https://doi.org/10.1101/2024.03.20.585909
Halai, A. D., Parkes, L. M., & Welbourne, S. R. (2015). Dual-echo fMRI can detect activations in
inferior temporal lobe during intelligible speech comprehension. *NeuroImage*, *122*, 214–221.
https://doi.org/10.1016/j.neuroimage.2015.05.067
Halai, A. D., Welbourne, S. R., Embleton, K., & Parkes, L. M. (2014). A comparison of dual
gradient-echo and spin-echo fMRI of the inferior temporal lobe. *Human Brain Mapping*,
35(8), 4118–4128. https://doi.org/10.1002/hbm.22463
Halchenko, Y. O., Goncalves, M., Ghosh, S., Velasco, P., Castello, M. V. di O., Salo, T., Wodder, J. T.,
Hanke, M., Sadil, P., Gorgolewski, K. J., Ioanas, H.-I., Rorden, C., Hendrickson, T. J., Dayan,
M., Houlihan, S. D., Kent, J., Strauss, T., Lee, J., To, I., … Kennedy, D. N. (2024). HeuDiConv—
Flexible DICOM conversion into structured directory layouts. *Journal of Open Source*
Software, *9*(99), 5839. https://doi.org/10.21105/joss.05839
Hargreaves, B. A., Cunningham, C. H., Nishimura, D. G., & Conolly, S. M. (2004). Variable-rate
selective excitation for rapid MRI sequences. *Magnetic Resonance in Medicine*, *52*(3), 590–
597. https://doi.org/10.1002/mrm.20168

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and
overlapping representations of faces and objects in ventral temporal cortex. *Science*,
293(5539), 2425–2430. https://doi.org/10.1126/science.1063736

Howard, D., & Patterson, K. E. (1992). The pyramids and palm trees test. *Https://Eprints.Ncl.Ac.Uk*.
https://eprints.ncl.ac.uk

Humphreys, G. F., Hoffman, P., Visser, M., Binney, R. J., & Lambon Ralph, M. A. (2015). Establishing
task- and modality-dependent dissociations between the semantic and default mode
networks. *Proceedings of the National Academy of Sciences*, *112*(25), 7857–7862.
https://doi.org/10.1073/pnas.1422760112

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL.
NeuroImage, *62*(2), 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jezzard, P., & Clare, S. (1999). Sources of distortion in functional MRI data. *Human Brain Mapping*,
8(2–3), 80–85. https://doi.org/10.1002/(SICI)1097-0193(1999)8:2/3<80::AID-
HBM2>3.0.CO;2-C

Jung, J., Williams, S. R., Sanaei Nezhad, F., & Lambon Ralph, M. A. (2017). GABA concentrations in
the anterior temporal lobe predict human semantic processing. *Scientific Reports*, *7*(1),
15748. https://doi.org/10.1038/s41598-017-15981-7

Kirilina, E., Lutti, A., Poser, B. A., Blankenburg, F., & Weiskopf, N. (2016). The quest for the best: The
impact of different EPI sequences on the sensitivity of random effect fMRI group analyses.
NeuroImage, *126*, 49–59. https://doi.org/10.1016/j.neuroimage.2015.10.071

Koopmans, P. J., Barth, M., Orzada, S., & Norris, D. G. (2011). Multi-echo fMRI of the cortical laminae
in humans at 7T. *NeuroImage*, *56*(3), 1276–1285.
https://doi.org/10.1016/j.neuroimage.2011.02.042

Kovářová, A., Gajdoš, M., Rektor, I., & Mikl, M. (2022). Contribution of the multi-echo approach in
accelerated functional magnetic resonance imaging multiband acquisition. *Human Brain
Mapping*, *43*(3), 955–973. https://doi.org/10.1002/hbm.25698

Kundu, P., Benson, B. E., Baldwin, K. L., Rosen, D., Luh, W.-M., Bandettini, P. A., Pine, D. S., & Ernst,
M. (2015). Robust resting state fMRI processing for studies on typical brain development
based on multi-echo EPI acquisition. *Brain Imaging and Behavior*, *9*(1), 56–73.
https://doi.org/10.1007/s11682-014-9346-4

Kundu, P., Brenowitz, N. D., Voon, V., Worbe, Y., Vértes, P. E., Inati, S. J., Saad, Z. S., Bandettini, P. A.,
& Bullmore, E. T. (2013). Integrated strategy for improving functional connectivity mapping
using multiecho fMRI. *Proceedings of the National Academy of Sciences*, *110*(40), 16187–
16192. https://doi.org/10.1073/pnas.1301725110

Kundu, P., Inati, S. J., Evans, J. W., Luh, W.-M., & Bandettini, P. A. (2011). Differentiating BOLD and
non-BOLD signals in fMRI time series using multi-echo EPI. *NeuroImage*, *60*(3), 1759–1770.
https://doi.org/10.1016/j.neuroimage.2011.12.028

Kundu, P., Voon, V., Balchandani, P., Lombardo, M. V., Poser, B. A., & Bandettini, P. A. (2017). Multi-
echo fMRI: A review of applications in fMRI denoising and analysis of BOLD signals.
NeuroImage, *154*, 59–80. https://doi.org/10.1016/j.neuroimage.2017.03.033

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and
computational bases of semantic cognition. *Nature Reviews Neuroscience*, *18*(1), 42–55.
https://doi.org/10.1038/nrn.2016.150

Lombardo, M. V., Auyeung, B., Holt, R. J., Waldman, J., Ruigrok, A. N. V., Mooney, N., Bullmore, E. T.,
Baron-Cohen, S., & Kundu, P. (2016). Improving effect size estimation and statistical power
with multi-echo fMRI and its impact on understanding the neural systems supporting
mentalizing. *NeuroImage*, *142*, 55–66. https://doi.org/10.1016/j.neuroimage.2016.07.022

Lynch, C. J., Power, J. D., Scult, M. A., Dubin, M., Gunning, F. M., & Liston, C. (2020). Rapid Precision
Functional Mapping of Individuals Using Multi-Echo fMRI. *Cell Reports*, *33*(12), 108540.
https://doi.org/10.1016/j.celrep.2020.108540

Markiewicz, C. J., Esteban, O., Goncalves, M., Provins, C., Salo, T., Kent, J. D., DuPre, E., Ciric, R.,
Pinsard, B., Blair, R. W., Poldrack, R. A., & Gorgolewski, K. J. (2024). *fMRIPrep: A robust*
preprocessing pipeline for functional MRI (Version 24.0.1) [Computer software]. Zenodo.
https://doi.org/10.5281/zenodo.12774221

Marques, J. P., & Norris, D. G. (2018). How to choose the right MR sequence for your research
question at 7 T and above? *NeuroImage*, *168*, 119–140.
https://doi.org/10.1016/j.neuroimage.2017.04.044

McNabb, C. B., Lindner, M., Shen, S., Burgess, L. G., Murayama, K., & Johnstone, T. (2020). Inter-slice
leakage and intra-slice aliasing in simultaneous multi-slice echo-planar images. *Brain*
Structure and Function, *225*(3), 1153–1158. https://doi.org/10.1007/s00429-020-02053-2

Miletić, S., Bazin, P.-L., Weiskopf, N., van der Zwaag, W., Forstmann, B. U., & Trampel, R. (2020).
fMRI protocol optimization for simultaneously studying small subcortical and cortical areas
at 7 T. *NeuroImage*, *219*, 116992. https://doi.org/10.1016/j.neuroimage.2020.116992

Moeller, S., Yacoub, E., Olman, C. A., Auerbach, E., Strupp, J., Harel, N., & Uğurbil, K. (2010).
Multiband multislice GE-EPI at 7 Tesla, with 16-fold acceleration using partial parallel
imaging with application to high spatial and temporal whole-brain fMRI. *Magnetic*
Resonance in Medicine, *63*(5), 1144–1153. https://doi.org/10.1002/mrm.22361

Morris, L. S., Kundu, P., Costi, S., Collins, A., Schneider, M., Verma, G., Balchandani, P., & Murrough,
J. W. (2019). Ultra-high field MRI reveals mood-related circuit disturbances in depression: A
comparison between 3-Tesla and 7-Tesla. *Translational Psychiatry*, *9*(1), 94.
https://doi.org/10.1038/s41398-019-0425-6

O'Brien, K. R., Kober, T., Hagmann, P., Maeder, P., Marques, J., Lazeyras, F., Krueger, G., & Roche, A.
(2014). Robust T1-weighted structural brain imaging and morphometry at 7T using
MP2RAGE. *PLOS ONE*, *9*(6), e99676. https://doi.org/10.1371/journal.pone.0099676

Poser, B. A., & Norris, D. G. (2009). Investigating the benefits of multi-echo EPI for fMRI at 7 T.
NeuroImage, *45*(4), 1162–1172. https://doi.org/10.1016/j.neuroimage.2009.01.007

Poser, B. A., Versluis, M. J., Hoogduin, J. M., & Norris, D. G. (2006). BOLD contrast sensitivity
enhancement and artifact reduction with multiecho EPI: Parallel-acquired inhomogeneity-
desensitized fMRI. *Magnetic Resonance in Medicine*, *55*(6), 1227–1235.
https://doi.org/10.1002/mrm.20900

Posse, S. (2012). Multi-echo acquisition. *NeuroImage*, *62*(2), 665–671.
https://doi.org/10.1016/j.neuroimage.2011.10.057

Puckett, A. M., Bollmann, S., Poser, B. A., Palmer, J., Barth, M., & Cunnington, R. (2018). Using multi-
echo simultaneous multi-slice (SMS) EPI to improve functional MRI of the subcortical nuclei
of the basal ganglia at ultra-high field (7T). *NeuroImage*, *172*, 886–895.
https://doi.org/10.1016/j.neuroimage.2017.12.005

Risk, B. B., Murden, R. J., Wu, J., Nebel, M. B., Venkataraman, A., Zhang, Z., & Qiu, D. (2021). Which
multiband factor should you choose for your resting-state fMRI study? *NeuroImage*, *234*,
117965. https://doi.org/10.1016/j.neuroimage.2021.117965

Roemer, P. B., Edelstein, W. A., Hayes, C. E., Souza, S. P., & Mueller, O. M. (1990). The NMR phased
array. *Magnetic Resonance in Medicine*, *16*(2), 192–225.
https://doi.org/10.1002/mrm.1910160203

Setsompop, K., Gagoski, B. A., Polimeni, J. R., Witzel, T., Wedeen, V. J., & Wald, L. L. (2012).
Blipped-controlled aliasing in parallel imaging for simultaneous multislice echo planar
imaging with reduced *g* -factor penalty. *Magnetic Resonance in Medicine*, *67*(5), 1210–1224.
https://doi.org/10.1002/mrm.23097

Smith, S. M. (2002). Fast robust automated brain extraction. *Human Brain Mapping*, *17*(3), 143–155.
https://doi.org/10.1002/hbm.10062

Smith, S. M., Beckmann, C. F., Andersson, J., Auerbach, E. J., Bijsterbosch, J., Douaud, G., Duff, E.,
Feinberg, D. A., Griffanti, L., Harms, M. P., Kelly, M., Laumann, T., Miller, K. L., Moeller, S.,
Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A. Z., Vu, A. T., … Glasser, M. F. (2013).

Resting-state fMRI in the Human Connectome Project. *NeuroImage*, *80*, 144–168.
https://doi.org/10.1016/j.neuroimage.2013.05.039
Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H.,
Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J.,
Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and
structural MR image analysis and implementation as FSL. *NeuroImage*, *23*, S208–S219.
https://doi.org/10.1016/j.neuroimage.2004.07.051
The tedana Community, Ahmed, Z., Bandettini, P. A., Bottenhorn, K. L., Caballero-Gaudes, C.,
Dowdle, L. T., DuPre, E., Gonzalez-Castillo, J., Handwerker, D., Heunis, S., Kundu, P., Laird, A.
R., Markello, R., Markiewicz, C. J., Maullin-Sapey, T., Moia, S., Molfese, P., Salo, T., Staden, I.,
… Whitaker, K. (2023). *ME-ICA/tedana: 23.0.1* (Version 23.0.1) [Computer software].
Zenodo. https://doi.org/10.5281/zenodo.7926293
Todd, N., Moeller, S., Auerbach, E. J., Yacoub, E., Flandin, G., & Weiskopf, N. (2016). Evaluation of 2D
multiband EPI imaging for high-resolution, whole-brain, task-based fMRI studies at 3T:
Sensitivity and slice leakage artifacts. *NeuroImage*, *124*, 32–42.
https://doi.org/10.1016/j.neuroimage.2015.08.056
Uğurbil, K. (2018). Imaging at ultrahigh magnetic fields: History, challenges, and solutions.
NeuroImage, *168*, 7–32. https://doi.org/10.1016/j.neuroimage.2017.07.007
Van de Moortele, P.-F., Akgun, C., Adriany, G., Moeller, S., Ritter, J., Collins, C. M., Smith, M. B.,
Vaughan, J. T., & Uğurbil, K. (2005). B1 destructive interferences and spatial phase patterns
at 7 T with a head transceiver array coil. *Magnetic Resonance in Medicine*, *54*(6), 1503–1518.
https://doi.org/10.1002/mrm.20708
van der Kolk, A. G., Hendrikse, J., Zwanenburg, J. J. M., Visser, F., & Luijten, P. R. (2013). Clinical
applications of 7 T MRI in the brain. *European Journal of Radiology*, *82*(5), 708–718.
https://doi.org/10.1016/j.ejrad.2011.07.007
Wu, X., Auerbach, E. J., Vu, A. T., Moeller, S., Lenglet, C., Schmitter, S., Van de Moortele, P.-F.,
Yacoub, E., & Uğurbil, K. (2018). High-resolution whole-brain diffusion MRI at 7T using
radiofrequency parallel transmission. *Magnetic Resonance in Medicine*, *80*(5), 1857–1870.
https://doi.org/10.1002/mrm.27189
Wu, X., Schmitter, S., Auerbach, E. J., Uğurbil, K., & Van de Moortele, P.-F. (2016). A generalized slab-
wise framework for parallel transmit multiband RF pulse design. *Magnetic Resonance in*
Medicine, *75*(4), 1444–1456. https://doi.org/10.1002/mrm.25689
Yun, S. D., & Shah, N. J. (2017). Whole-brain high in-plane resolution fMRI using accelerated EPIK for
enhanced characterisation of functional areas at 3T. *PLOS ONE*, *12*(9), e0184759.
https://doi.org/10.1371/journal.pone.0184759
Zahn, R., Moll, J., Krueger, F., Huey, E. D., Garrido, G., & Grafman, J. (2007). Social concepts are
represented in the superior anterior temporal cortex. *Proceedings of the National Academy*
of Sciences, *104*(15), 6430–6435. https://doi.org/10.1073/pnas.0607061104
Zhang, M., Ding, B., Dragonu, I., Liebig, P., & Rodgers, C. T. (2024). Dynamic parallel transmit
diffusion MRI at 7T. *Magnetic Resonance Imaging*, *111*, 35–46.
https://doi.org/10.1016/j.mri.2024.03.037
1010