Utility of the Recombinase Driver CX3CR1::Cre Rat Strain to Evaluate Microglial Contributions to Nicotine Addiction

Ashley White[1], Ashley J. Craig[1], Daryl Richie[1], Christa L. Corley[1], Emma O. Bondy[1], Percell T. Kendrick, Jr., Michael D. Scofield[2], and Cassandra D. Gipson[1]*

[1]Department of Pharmacology and Nutritional Sciences, University of Kentucky, Lexington KY

[2]Department of Anesthesiology, Medical University of South Carolina, Charleston, SC

*Correspondence to:

Cassandra D. Gipson, Ph.D.
Department of Pharmacology and Nutritional Sciences
University of Kentucky
900 S. Limestone
CTW Room 535
Lexington, KY, 40536, USA
Email: cdgips2@uky.edu
859-216-0625

Conflicts of interest: None
Funding Sources: This work was supported by the National Institutes of Health Grant DA058933, and DA055879, DA044479, DA046526, DA061626, and DA049130 (to CDG). All authors have no disclosures to declare.

Author Note: We would like to thank Mei Hong for technical assistance.

Abstract

Smoking remains a leading preventable cause of death, and nicotine is the primary substance responsible for maintaining use of tobacco products. Preclinical rodent models have shown that neuroimmune signaling is dysregulated by nicotine self-administration (SA) within the nucleus accumbens core (NAcore), which is a key region within the mesolimbic brain reward pathway. Microglia are the resident brain immune cell and prior studies have shown that they play an important role in nicotine-related behaviors. However, while there are transgenic mouse lines that allow for specific evaluations of microglia to neurobiology and behavior, there are fewer tools available for rats as a model species thus limiting our ability to evaluate specific contributions of microglia to nicotine SA. Using transgenic rats expressing Cre under the control of the CX3CR1 promoter bred on a Long Evans (LE) background, we show that NAcore microglia can be specifically transduced with Designer Receptors Exclusively Activated by Designer Drugs (DREADDs). We further show that CX3CR1::Cre rats readily self-administer nicotine, and display a characteristic extinction curve that is not different from outbred LE or Cre-negative littermates. Together, these validation studies lay the foundation for future use of this transgenic rat line to evaluate the specific contributions of microglia in the brain to neurobehavioral underpinnings of nicotine addiction.

-208 words-

KEYWORDS: Microglia, Nicotine, Self-administration, Behavior, DREADDs

Introduction

The immunomodulatory effects of nicotine have been clinically (Sher et al. 1999) and preclinically (Li et al. 2021; Namba et al. 2019; Shi et al. 2009) documented, however, the impacts of chronic, volitional nicotine use on neuroimmune signaling is a field in its infancy. Microglia are resident brain immune cells which regulate adaptive and innate immune responses (Paolicelli et al. 2022), regulate synaptic plasticity (Aramideh et al. 2021; Bessis et al. 2007; Paolicelli and Ferretti 2017), and are a part of the neurovascular unit (Liu et al. 2020). Recent studies have implicated microglia in nicotine addiction processes as microglia within the reward pathway and specifically the nucleus accumbens (NA) play a large role in regulating nicotine-related neurobehavior (Kumar et al. 2024; Adeluyi et al. 2019; Soares and Picciotto 2024). However, our ability to specifically manipulate microglia to determine causal roles of these brain immune cells in nicotine addiction processes remains limited.

The current neuroscience toolbox for evaluating microglia includes viruses, chemicals to deplete them, and transgenic rodent models. Viruses that have been utilized in prior studies include Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) which incorporate adeno-associated viral vectors (AAVs) with promoters that are thought to be microglia-specific (e.g., CD68; (Grace et al. 2016), and AAV variants that allow *in vitro* and *in vivo* microglial transduction through directed evolution of the AAV capsid protein to increase efficiency of genetic payload into microglia (Lin et al. 2022). Notably, the pAAV-CD68-hM4D-mCherry (Gi) and pAAV-CD68-EGFP DREADD utilized in (Grace et al. 2016) are commercially available but untested in their ability to penetrate microglia within the brain as that study infused DREADDs intrathecally. The AAV variants which were discovered a few years later to infect microglia were also found to infect neurons (Lin et al. 2022), thus making it difficult to tease apart microglial versus neuronal contributions to neurobiology and behavior.

As mentioned above, other methodologies to evaluate contributions of microglia to brain function include depletion methodologies with either a Cx3cr1-Dtr rat, which allows for short-

term microglia depletion (De Luca et al. 2019), or with colony-stimulating factor–1 receptor

(CSF1R) inhibitors (e.g., PLX5622 or PLX3397), which deplete a proportion of microglia as well

as other affect peripheral cells (Green et al. 2020) when administered in chow. An advantage of

this technique is that microglia can repopulate (Wickel et al. 2024), allowing for bidirectional

evaluations of microglia to neurobiology and behavioral outcomes. However, microglia depletion

is non-specific and can have off target effects, limiting the cell-type specificity of this approach.

An additional methodology includes frequently utilized transgenic mouse strains which have

been used primarily in other fields of research. Transgenic mice, including the recombinase

driver CX3CR1-Cre strains (Zhao et al. 2019), have been used to visualize or manipulate

microglia in the context of other pathologies (Wieghofer and Prinz 2016). Endogenous CX3CR1

is a microglia-specific marker, and it is a receptor with the endogenous ligand fractalkine

(Nishiyori et al. 1998). However, there has been some debate regarding specificity of CX3CR1

to microglia, as there are reports that CX3CR1 expression occurs in other cell types depending

on the tissue type (Lee et al. 2018), or co-localization of CX3CR1 with glial fibrillary acidic

protein (GFAP; (Peng et al. 2017), a marker of astrocyte function (Middeldorp and Hol 2011),

although this has not been evaluated within the brain reward pathway to our knowledge. Some

studies have also reported that CX3CR1::cre in mice is not specific to microglia, with significant

leakage into neurons (Haimon et al. 2018; Hwang et al. 2017; Zhang et al. 2018). While there

are two constitutive and two tamoxifen-inducible CX3CR1::cre mouse lines (Gong et al. 2007;

Parkhurst et al. 2013; Yona et al. 2013), there is only one CX3CR1::cre rat line which has never

been utilized to evaluate microglia contributions to addiction neuroscience and has never been

demonstrated to allow for microglia-specific infection.

A large majority of studies modeling nicotine addiction utilize self-administration (SA), which

allows for evaluations of volitional nicotine use and is most frequently implemented in rat models

(Corley et al. 2024). Studies also frequently incorporate extinction training following SA to

evaluate the ability to learn to withhold the prepotent response to receive the drug and is

typically conducted under withdrawal conditions (Namba, et al., 2019). While some studies

incorporate mouse SA to model aspects of nicotine addiction (Fowler et al. 2011; Fowler and

Kenny 2011), use of mice in nicotine SA is limited to a small number of researchers and can

present challenges with catheter patency among other issues. While all currently available

methodologies to evaluate microglia have limitations with some noted above, the microglia

addiction field has a significant barrier in that there are no validated rat strains for use in SA

procedures. Here, we utilized a commercially available CX3CR1::cre transgenic rat strain (LE-

Tg(OTTC1005) CX3CR1-Cre-ERT2), which is a tamoxifen-inducible Cre recombinase strain

based on a Long Evans background. The goal of the current study was therefore to determine if

this rat strain (1) displays microglia-specific expression of viral vectors within the NAcore without

infecting surrounding neurons, and (2) can acquire nicotine SA and extinction (3) similar to

CX3CR1::cre (-) littermates and wildtype Long Evans rats.

Methods

<u>Subjects.</u>

Breeder male LE-Tg(OTTC1005) CX3CR1-Cre-ERT2 rats were purchased from the Rat

Resource and Research Center (RRRC) and bred in-house with Long Evans females purchased

from Charles River. Rats were housed in a temperature and humidity-controlled animal facility

within the Division of Laboratory Animal Resources (DLAR) at the University of Kentucky. The

colony room was on a 12:12 hr light:dark reverse light cycle and rats in the breeding colony had

ad libitum access to food and water. Rats in nicotine SA were either CX3CR1::Cre (+; N=50),

CX3CR1::Cre (-; N=15), or purchased from Charles River as an outbred strain control (N=37).

Male and female rats were 60-90 days old at the beginning of the study and were food restricted

to 85% of free feed weight throughout SA experimentation. All animal use practices were

approved by the Institutional Animal Care and Use Committee (IACUC) of the University of

Kentucky (UK; Protocol #2020-3438). 19 CX3CR1::Cre (+), 3 CX3CR1::Cre (-), and 2 Long Evans

rats were excluded from the study due to catheter patency issues.

<u>CX3CR1::Cre Genotyping.</u>

Ear punches were collected from all animals on post-natal day 21. In separate 1.5 mL

microcentrifuge tubes, ear punch tissues were incubated in lysis buffer and Proteinase K

(Invitrogen, ~2 mg/ml, 24 µl total volume) at 55°C for 30 minutes, vortexed, and pelleted (18,000

x g). Samples were incubated for an additional 30 min at 55°C, vortexed, then resuspended in

distilled water up to 200 µl total volume, and pelleted (18,000 x g). The samples were heated at

100°C for 10 minutes, cooled to room temperature, and pelleted at 18,000 x g for 20 minutes at

4°C. Next, 100 µl of the supernatant containing the genomic DNA was transferred to a new 1.5

ml microcentrifuge tube, and standard PCR amplification was used for genotyping.

Subjects were genotyped using the following primers: CX3CR1 F103289 (5'- TCAGGGTGG

CCCATAACCAC-3′) and CreETT2 R713 (5'-AGGTAGTTATTCGGATCATCAGCTTAC AC-3')

which produce a 1017 bp amplicon spanning the 5' end of the CX3CR1-Cre-ERT2. Primers

CX3CR1 F103289 (5'-TCAGGGTGGCCCATAACCAC-3′) and CREERT2 R246 (5'-CAGTGA

AGCTTCGATGATGGG C-3'), which produce a 550 bp amplicon spanning the 5' end of the

CX3CR1-Cre-ERT2. Primers CreERT2 F1206 (5'-CAGTGAAGCTTCGATGATGGGC-3') and

CX3CR1 R104157 – 5' – GCCAGATTTCCTGCAGGACCT- 3'), which produce a 1553 bp

amplicon spanning the 3' end of the CX3CR1-Cre-ERT2. All animals used were confirmed

CX3CR1::cre transgene-positive.

<u>Surgical Procedures.</u>

Rats included in the immunohistochemistry study underwent surgery for viral infusions

targeting the NAcore as previously described (Leyrer-Jackson et al. 2021). Rats included in the

nicotine SA study underwent jugular catheter surgery and virus infusion surgery. All rats were

anesthetized using ketamine hydrochloride (80–100 mg/kg, i.m.) and xylazine (8 mg/kg, i.m.) and

underwent surgical implantation of intravenous jugular catheters as well as stereotaxically infused

viral delivery within the NAcore. Intravenous jugular catheters (made from polyurethane tubing;

BTPU-040; Instech) were inserted 2.5–3 cm into the right jugular vein and were threaded

subcutaneously to the posterior side of the animal where it was connected to an indwelling back

port (Instech). Dental cement (SNAP or Ortho-Jet) was used to adhere the catheter to the port.

The indwelling port was sutured using 4–0 vicryl braided suture (Ethicons) subcutaneously ~2 cm

caudal from the shoulder blades. Following jugular vein catheterization, animals were immediately

transferred to a rat stereotaxic frame and NAcore guide cannulae were bilaterally aimed at the

NAcore according to a stereotaxic atlas (Paxinos and Watson 2007). Viral vectors encoding

DREADDs for microglia infection were infused into the NAcore of CX3Cr1::cre animals: pAAV-

EF1a-DIO-hM4D(Gi)-mCherry (inhibitory; titer: 1.2×10^{13} vg/ml; Addgene, #50461), pAAV-EF1a-

DIO-hM3D(Gq)-mCherry (excitatory; titer: 1.3×10^{13} vg/ml; Addgene, #50460; packaged into an

AAV5 vector by UNC Vector Core), or pAAV5-EF1a-DIO-mCherry (control; i.e., only mCherry-

expressing; titer: 1.5×10^{13} vg/ml; Addgene, #44362) at a volume of 0.5 µl per hemisphere. Ef1a

was selected as the promoter because it is abundantly expressed in all cell types (Wakabayashi-

Ito and Nagata 1994), it is double floxed, and both Gq and Gi coupled-versions are commercially

available. Rats were immediately administered cefazolin (100 mg/kg, i.v.) and heparin (10 U/ml,

i.v.) and for seven consecutive days during the recovery period. Meloxicam (1 mg/kg, i.v.) was

given immediately and for the first 3 d of the recovery period. For rats in the SA study, heparin (10

U/ml, i.v.) was administered daily. All rats received 4-hydroxytamoxifen (20 mg/kg, I.P.; (Hughes

et al. 2024) 3 weeks following virus infusion and 2 days prior to experimentation.

<u>Food Training Procedures</u>

All rats in the nicotine SA study (N = 30) underwent food training on the sixth day of

postoperative care. Food restriction to maintain 85% of free feed weight was implemented a

minimum of 2 h before food training initiation and was maintained for the duration of the

experiment. Food training sessions were 15 h duration, and one active lever press resulted in the

delivery of one food pellet [fixed-ratio-1 (FR1), schedule of reinforcement; Bio-Serv, 45 mg/pellet)

Concurrently, one food pellet was delivered every 20 min regardless of response. Light and tone

stimuli were not paired with pellet administration. Food training criteria was set to 2:1 active to

inactive lever presses throughout the session and a minimum of 200 active lever presses. Rats

that did not pass the initial food training session completed a second session before beginning

SA procedures (N=10). Rats unable to pass food training procedures were excluded from the

study (N = 8). All food training, SA, and extinction sessions were conducted in modular operant

conditioning chambers (13 ENV-008, 15 ENV-007; Med Associates), which have been previously

described in detail (Sadek et al. 2024; Maher et al. 2023; Leyrer-Jackson et al. 2021).

<u>Nicotine SA Procedures.</u>

Nicotine infusions (0.06 mg/kg/infusion) were paired with a compound stimulus

(light+tone) and was followed by a 20-s timeout period. Nicotine was delivered across a 5.9-s

duration, at a total volume of 0.1 ml of nicotine was administered per infusion. Infusions were

paired with cue lights, located above each lever, and a 2900-Hz tone was presented throughout

the duration of the nicotine infusion. Active lever presses during the timeout period were recorded,

but did not result in additional nicotine infusions. An inactive lever was extended at all times and

responses were recorded; however, presses yielded no programmed consequences or rewards.

Session duration was 2 h in length and a FR1 schedule of reinforcement was used. All animals

were required to complete a minimum of 10 sessions before moving into extinction with the

following criteria: ≥10 nicotine infusions obtained and ≥2:1 active/inactive lever press ratio.

Additionally, extinction is defined either by (1) the lack of delivery of a reinforcer that was

previously delivered after a response (e.g., responses on the active lever no longer result in

nicotine infusions), or (2) as the absence of a contingency between response and reinforcer (e.g.,

the nicotine infusion occurs regardless of an animals' response; see Rescorla and Skucy,

1969; Lattal and Lattal, 2012). Given that the reinforcer here (the nicotine infusion) is not delivered

following a time-out active lever press, the latter definition applies to these types of responses

and are therefore distinct from active lever presses that directly result in the initiation of a nicotine

infusion. Thus, these two types of lever presses were analyzed separately in the current study.

Following 10 non-consecutive criteria-making sessions, rats were moved to the extinction

phase which took place in the same boxes as self-administration. During the extinction phase,

active lever presses no longer resulted in nicotine infusions or associated cues. A minimum of 14

2-h extinction sessions was required for each animal. Extinction criteria were set at <30 active

lever presses on the last day of extinction. Eight animals were removed from the current study

because of not meeting self-administration criteria ($N = 5$) or lack of DREADD

expression/improper cannula placement ($N = 3$).

<u>Immunohistochemistry.</u>

Rats were anesthetized with an overdose of euthasol (excess of 0.25 ml/kg i.p.) and

transcardially perfused with 0.01 M PBS followed by 4% paraformaldehyde. Brains were then

coronally sliced on a vibratome at 60 μm for immunohistochemistry staining. All antibodies and

dilutions are listed in **Table 1**. To determine if the virus infected microglia cells in the NAcore,

NAcore sections were first incubated in anti-Iba-1 primary antibody that was diluted in 1% BSA,

0.3% Triton, .05% azide, and 0.01 M PBS for 24 hours at room temperature on a shaker. Sections

were then rinsed with 0.01 M PBS three times for 30 minutes each then incubated in a secondary

antibody conjugated to Alexa Fluor 488 for 2 hours. Sections were mounted onto glass slides and

coverslipped using an anti-fade mounting medium. Next, to determine if the virus infected

microglia that expressed the cre (+) genotype, NAcore sections were simultaneously stained with

2 primary antibodies (anti-Iba1, and anti-CX3CR1) that were diluted in 1% BSA, 0.3% Triton, .05%

azide, and 0.01 M PBS for 24 hours at room temperature on a shaker. Sections were then rinsed

and incubated in secondary antibodies conjugated to Alexa Fluor 488 and Alexa Fluor 594 for 2

hours. Sections were mounted onto glass slides and coverslipped using an anti-fade mounting

medium containing DAPI to visualize cell bodies. Next to show colocalization of infected microglia

without infected neurons, NACore sections were incubated in 2 primary antibodies (anti-Iba1, and

anti-NeuN) that were diluted in 1% BSA, 0.3% Triton, .05% azide, and 0.01 M PBS for 24 hours

at room temperature on a shaker. Sections were then rinsed and incubated in secondary

antibodies conjugated to Alexa Fluor 488 and Alexa Fluor 594 for 2 hours. All confocal

micrographs were captured using a Nikon A1R Inverted Confocal Microscope.

<u>Statistical Analysis.</u>

Food training lever press data, drug infusions, and lever press discrimination (active,

inactive) across sessions were analyzed using linear mixed effects (LME) modeling (JMP

software from SAS). Tukey's HSD was used as post-hoc pairwise comparisons on nominal

variables where appropriate ($\alpha = 0.05$). Subject was treated as a random factor and session

was treated as a continuous factor whereas group was evaluated as a nominal variable in all

relevant analyses. Model fits were conducted to determine the best model fit for the datasets

where appropriate, and the models with the lowest Akaike information criterion (AICc) were

selected. Lever discrimination was assessed as the ratio of active/active+inactive lever presses

(including infusion-earning and timeout active lever presses, as well as all inactive lever

presses).

Results

<u>**Cre-Dependent Viruses Express Within NAcore Microglia**</u>

Immunohistochemistry was first conducted to determine if the fluorescent mCherry tag

from the viruses expressed in NAcore microglia following intracranial viral delivery. Images were

taken using a Nikon A1R Confocal Microscope. **Figure 1A** shows that mCherry from the Gi viral

construct was successfully observed within NAcore Iba1-(+) cells. Similarly, we found co-

localization of mCherry with NAcore Iba1-(+) cells with the Gq construct (**Figure 1B**). To

determine specificity to CX3CR1-expressing cells, we next evaluated co-localization of mCherry

from the Gi construct with CX3CR1 and Iba-1. Notably, cells that were mCherry-(+) expressed

both microglia markers (**Figure 1C**), demonstrating that the CX3CR1::cre transgenic rat strain

allowed for AAV infection of microglia. Finally, we confirmed somatic expression of mCherry with

co-localization of the fluorescent tag with DAPI, which visualizes nuclear DNA (Tarnowski et al.

1991); **Figure 1D**). Next, we determined if mCherry from the Gi construct co-localized with the

neuronal marker NeuN. As shown in **Figure 1E**, there was no co-localization of mCherry with

NeuN, demonstrating specificity of mCherry to microglia within the NAcore.

CX3CR1::cre Rats Acquire Nicotine SA and Extinction

First, baseline nicotine consumption was compared during nicotine SA acquisition as a

function of virus groups following viral administration (see **Figure 2A** for an experimental timeline).

The best model fit excluded sex in the model (ΔAICc = 31.9, relative to the next best model fit,

which included both sex and group as a variable). LME revealed no significant main effects group

($F_{2,2}$ = 0.6081, p>0.05), session ($F_{1,21}$ = 1.0336, p>0.05), or no group x session interactions ($F_{2,21}$

= 0.4445, p>0.05) (**Figure 3B**). These results indicate that virus administration did not change

nicotine SA acquisition rates regardless of the construct type. When nicotine consumption was

summed across experimentation, there were no group differences in the total amount of nicotine

consumption across the 15-day SA experiment ($F_{2,21}$ = 1.2943, p>0.05).

Next, data were collapsed across virus type and compared to CX3CR1::cre (-) littermates,

as well as purchased Long Evans rats. The best model fit excluded sex in the model (ΔAICc =

44.2, relative to the next best model fit, which included both sex and group as a variable). First

LME was conducted on lever discrimination during food training as a function of strain. No main

effect of strain ($F_{1,68}$ = 0.44, p>0.05 was found, indicating that all rats acquired food training

similarly with no strain differences in this effect. Baseline nicotine consumption was compared

during nicotine SA acquisition as a function of strain groups (see **Figure 3A** for an experimental

timeline). LME revealed no significant main effects of group ($F_{2, 64}$ = 1.1342, p>0.05), or session

$(F_{1, 64} = 3.0377, p>0.05)$ or group x session interactions $(F_{2, 64} = 2.2673, p>0.05)$ (**Figure 3B**).

These results indicate that the strain did not change nicotine SA acquisition rates regardless of

the genetic strain used. LME was conducted on lever discrimination during nicotine acquisition as

a function of strain. A significant main effect of strain $(F_{2,64.02} = 5.3284, p<0.05)$ was found,

however, all groups met the 2:1 active to inactive lever pressing ratio. When nicotine consumption

was summed across experimentation, there were no group $(F_{2,77} = 1.7039, p>0.05)$ differences in

the total amount of nicotine consumption across the 10-day SA experiment.

Discussion

Here we show that the CX3CR1::cre transgenic rat strain can be used to specifically

transduce DREADDs in microglia within a key region of the brain reward pathway, the NAcore,

and these rats can acquire intravenous nicotine SA and also undergo successful nicotine

extinction training. We compared acquisition curves across virus type and across strain with no

differences in either analyses. As noted previously, evaluations of microglia in addiction

neurobiology have been limited due to the tools available to study this cell type in rat models. Our

current data reported here support the use of this transgenic rat strain in the addiction

neuroscience field to evaluate microglia contributions to substance use, and particularly, in the

nicotine addiction field.

While the current study demonstrates the utility of this rat strain for studying aspects of

addiction neuroscience, there are notable limitations that must be outlined. It is important to note

that the current study did not functionally evaluate the impacts of chemogenetic activation or

inhibition on NAcore microglia, including evaluations of transcriptomic, epigenetic, metabolomic,

or proteomic changes induced by activation of the DREADD receptors. This is important because

the microglia field has recently shifted from a dichotomous thinking of "activated" versus "resting"

microglial states to a more complex multidimensional approach to microglial states (Paolicelli et

al. 2022). It is also unknown how muscarinic DREADD receptors expressed on CX3CR1(+)

microglia may shift these dynamic aspects of these immune cells. One study did find that

activation of Gi DREADDs expressed in microglia of CX3CR1$^{creER/+}$:R26$^{LSL-hM4Di/+}$ transgenic mice

inhibited chronic pain and spinal nerve transection-induced microglial reactivity (Yi et al. 2021).

This study also found that the Gi DREADD reduced interferon regulatory factor 8 (IRF8) and

interleukin (IL)-1β expression in Iba1(+) cells, although this study did not confirm that all microglia

were DREADD(+). Further, while this rat strain allows for infection of CX3CR1(+) microglia, this

methodology requires these cells to be virally infected with the construct. Here we did not find that

all microglia within the NAcore were mCherry(+), indicating that some microglia within the area of

viral delivery were not infected with the constructs. This could either be due to (1) too little infused

volume of the virus to perfuse through the visualized area of the NAcore, (2) some microglia did

not express cre, or (3) some microglia were able to mount an immune response from the viral

constructs. It is also important to note that this rat strain requires tamoxifen injections. Given that

tamoxifen is a selective estrogen receptor modulator and may have sex-specific antiestrogenic

effects (Rochefort and Borgna 1981) that could interfere with experimental outcomes, future

studies incorporating CX3CR1::cre rats should take this into account in experimental timelines.

Here, rats were administered tamoxifen several weeks prior to quantification of microglia

morphology, and during the early part of nicotine SA acquisition. Importantly, tamoxifen has a

half-life of 10.3 hrs in rats (Robinson et al. 1991), and five elimination half-lives are required for

approximately 97% of the bioavailable dose of a drug to be eliminated from the body (here, 51.5

hrs would be required; (Nnane 2005). Given this and that no sex differences were found in the

nicotine SA acquisition curves, it is unlikely that tamoxifen impacted behavioral outcomes here.

As well, another study administered 4-OHT immediately prior to cocaine conditioning and

observed measurable cocaine place preference in mice (Hughes et al. 2024). Finally, it is not

clear if CX3CR1::cre rat line has altered constitutive CX3CL1/CX3CR1 signaling, which is

important in neuron/microglia crosstalk in brain injury and inflammation (Pawelec et al. 2020).

Despite these caveats, this rat strain may allow for more granular *in vivo* evaluations of microglia

to glutamate dyshomeostasis induced by addictive drugs (Scofield et al. 2016) including nicotine

(Gipson et al. 2013), as well as cell type-specific dissections of glial contributions to the

neurobehavior of nicotine addiction.

Conclusions

Here, we have shown that the inducible recombinase driver CX3CR1::Cre rat strain can

be successfully used in a model of nicotine SA. CX3CR1::Cre rats readily self-administer nicotine,

while the viral constructs successfully infected microglia without neuronal co-infection within the

NAcore, eliminating one of the main issues with other methodologies utilizing viral constructs.

While mouse nicotine SA can be conducted successfully (Souter, et al., 2022), the majority of SA

studies utilize rats as the model rodent species, thus this rat will provide researchers with the

toolbox to readily manipulate microglia in rats undergoing SA of addictive drugs. Understanding

these contributions will provide insights on the neuroimmune landscape of nicotine addiction and

how immune modulation in different areas of the brain affect behavior.

-- 3854 words --

Table 1. Antibodies for Immunohistochemistry
347

Target	Company	Catalogue #	RRID	Dilution
Mouse Anti-NeuN antibody	Abcam	ab104224	RRID:AB_10711040	1:500
AlexaFluor 488 goat anti-chicken IgY	Abcam	ab150173	RRID:AB_2636995	1:500
Anti Iba1, Rabbit	Fujifilm	019-19741	RRID:AB_839504	1:500
Anti CX3CR1, Mouse	Abcam	ab167571	RRID:AB_2916288	1:500
DAPI antibody	Thermo Fisher	D1306	RRID:AB_2629482	1:1000

Figure Legends.

Figure 1. NAcore Microglia but not Neurons Are Transduced with AAV Constructs.
Representative immunohistochemical images from animals receiving viral infusions are shown.
Slices were stained with either whole cell microglial marker, Iba1; Cre-recombinase marker,
mCherry; microglial receptor, CX3CR1; cell soma marker, DAPI, and/or neuronal cell marker,
NeuN. Merged images show overlap of microglial and viral markers, indicating infection of
microglia with viral constructs without neuronal infection. AC = anterior commissure, denoted by
a dotted line.

Figure 2. CX3CR1::cre Rats Acquire Nicotine SA and Extinction. (A) Experimental timeline
for CX3CR1::cre rats. (B) Food training where animals met a 2:1 active to inactive lever
pressing ratio compared across virus groups with no significant differences *(p>0.05)*. (C$_{1-2}$)
CX3CR1::cre rats acquired lever responding to the active lever to receive IV infusions of
nicotine (0.06mg/kg/infusion) with no differences in lever discrimination. (D) CX3CR1::cre rats
had no significant differences in total nicotine consumption across the 15 day timeline. (E)
CX3CR1::cre rats showed a decline in lever pressing behavior during extinction training to < 30
lever presses per session.

**Figure 3. CX3CR1::cre Rats Acquire Nicotine SA similarly to Cre-negative Littermates and
Wild-Type rats.** (A) Experimental timeline for cre-negative littermates and ordered wild-type
rats from Charles River. (B) Food training where animals met a 2:1 active to inactive lever
pressing ratio compared across strain groups with no significant differences *(p>0.05)*. (C) Rats
had no significant differences in total nicotine consumption across the 10 day timeline across
strains. (D$_{1-2}$) CX3CR1::cre rats acquired lever responding to the active lever to receive IV
infusions of nicotine (0.06mg/kg/infusion) compared to cre-negative littermates and wild-type
ordered animals with a significant difference in lever discrimination between strains (p<0.05).

Figure 1.

Figure 2.

384 **Figure 3.**

REFERENCES

Adeluyi A, Guerin L, Fisher ML, Galloway A, Cole RD, Chan SSL, Wyatt MD, Davis SW, Freeman LR, Ortinski PI, Turner JR (2019) Microglia morphology and proinflammatory signaling in the nucleus accumbens during nicotine withdrawal. Sci Adv 5 (10):eaax7031. doi:10.1126/sciadv.aax7031

Aramideh JA, Vidal-Itriago A, Morsch M, Graeber MB (2021) Cytokine Signalling at the Microglial Penta-Partite Synapse. Int J Mol Sci 22 (24). doi:10.3390/ijms222413186

Bessis A, Bechade C, Bernard D, Roumier A (2007) Microglial control of neuronal death and synaptic properties. Glia 55 (3):233-238. doi:10.1002/glia.20459

Corley C, Craig A, Sadek S, Marusich JA, Chehimi SN, White AM, Holdiness LJ, Reiner BC, Gipson CD (2024) Enhancing translation: A need to leverage complex preclinical models of addictive drugs to accelerate substance use treatment options. Pharmacol Biochem Behav 243:173836. doi:10.1016/j.pbb.2024.173836

De Luca SN, Sominsky L, Soch A, Wang H, Ziko I, Rank MM, Spencer SJ (2019) Conditional microglial depletion in rats leads to reversible anorexia and weight loss by disrupting gustatory circuitry. Brain Behav Immun 77:77-91. doi:10.1016/j.bbi.2018.12.008

Fowler CD, Kenny PJ (2011) Intravenous nicotine self-administration and cue-induced reinstatement in mice: Effects of nicotine dose, rate of drug infusion and prior instrumental training. Neuropharmacology 61 (4):687-698. doi:10.1016/j.neuropharm.2011.05.012

Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ (2011) Habenular alpha5 nicotinic receptor subunit signalling controls nicotine intake. Nature 471 (7340):597-601. doi:10.1038/nature09797

Gipson CD, Reissner KJ, Kupchik YM, Smith AC, Stankeviciute N, Hensley-Simon ME, Kalivas PW (2013) Reinstatement of nicotine seeking is mediated by glutamatergic plasticity. Proc Natl Acad Sci U S A 110 (22):9124-9129. doi:10.1073/pnas.1220591110

Gong S, Doughty M, Harbaugh CR, Cummins A, Hatten ME, Heintz N, Gerfen CR (2007) Targeting Cre recombinase to specific neuron populations with bacterial artificial chromosome constructs. J Neurosci 27 (37):9817-9823. doi:10.1523/jneurosci.2707-07.2007

Grace PM, Strand KA, Galer EL, Urban DJ, Wang X, Baratta MV, Fabisiak TJ, Anderson ND, Cheng K, Greene LI, Berkelhammer D, Zhang Y, Ellis AL, Yin HH, Campeau S, Rice KC, Roth BL, Maier SF, Watkins LR (2016) Morphine paradoxically prolongs neuropathic pain in rats by amplifying spinal NLRP3 inflammasome activation. Proc Natl Acad Sci U S A 113 (24):E3441-3450. doi:10.1073/pnas.1602070113

Green KN, Crapser JD, Hohsfield LA (2020) To Kill a Microglia: A Case for CSF1R Inhibitors. Trends Immunol 41 (9):771-784. doi:10.1016/j.it.2020.07.001

Haimon Z, Volaski A, Orthgiess J, Boura-Halfon S, Varol D, Shemer A, Yona S, Zuckerman B, David E, Chappell-Maor L, Bechmann I, Gericke M, Ulitsky I, Jung S (2018) Re-evaluating microglia expression profiles using RiboTag and cell isolation strategies. Nat Immunol 19 (6):636-644. doi:10.1038/s41590-018-0110-6

Hughes BW, Huebschman JL, Tsvetkov E, Siemsen BM, Snyder KK, Akiki RM, Wood DJ, Penrod RD, Scofield MD, Berto S, Taniguchi M, Cowan CW (2024) NPAS4 supports cocaine-conditioned cues in rodents by controlling the cell type-specific activation balance in the nucleus accumbens. Nature communications 15 (1):5971. doi:10.1038/s41467-024-50099-1

Hwang HW, Saito Y, Park CY, Blachère NE, Tajima Y, Fak JJ, Zucker-Scharff I, Darnell RB (2017) cTag-PAPERCLIP Reveals Alternative Polyadenylation Promotes Cell-Type Specific Protein Diversity and Shifts Araf Isoforms with Microglia Activation. Neuron 95 (6):1334-1349.e1335. doi:10.1016/j.neuron.2017.08.024

Kumar M, Keady J, Aryal SP, Hessing M, Richards CI, Turner JR (2024) The Role of Microglia in Sex- and Region-Specific Blood-Brain Barrier Integrity During Nicotine Withdrawal. Biol Psychiatry Glob Open Sci 4 (1):182-193. doi:10.1016/j.bpsgos.2023.08.019

Lee M, Lee Y, Song J, Lee J, Chang SY (2018) Tissue-specific Role of CX(3)CR1 Expressing Immune Cells and Their Relationships with Human Disease. Immune Netw 18 (1):e5. doi:10.4110/in.2018.18.e5

Leyrer-Jackson JM, Holter M, Overby PF, Newbern JM, Scofield MD, Olive MF, Gipson CD (2021) Accumbens Cholinergic Interneurons Mediate Cue-Induced Nicotine Seeking and Associated Glutamatergic Plasticity. eNeuro 8 (1). doi:10.1523/ENEURO.0276-20.2020

Li H, Peng Y, Lin C, Zhang X, Zhang T, Wang Y, Li Y, Wu S, Wang H, Hutchinson MR, Watkins LR, Wang X (2021) Nicotine and its metabolite cotinine target MD2 and inhibit TLR4 signaling. Innovation (Camb) 2 (2):100111. doi:10.1016/j.xinn.2021.100111

Lin R, Zhou Y, Yan T, Wang R, Li H, Wu Z, Zhang X, Zhou X, Zhao F, Zhang L, Li Y, Luo M (2022) Directed evolution of adeno-associated virus for efficient gene delivery to microglia. Nat Methods 19 (8):976-985. doi:10.1038/s41592-022-01547-7

Liu LR, Liu JC, Bao JS, Bai QQ, Wang GQ (2020) Interaction of Microglia and Astrocytes in the Neurovascular Unit. Front Immunol 11:1024. doi:10.3389/fimmu.2020.01024

Maher EE, White AM, Craig A, Khatri S, Kendrick PT, Jr., Matocha ME, Bondy EO, Pallem N, Breakfield G, Botkins M, Sweatt O, Griffin WC, Kaplan B, Weafer JJ, Beckmann JS, Gipson CD (2023) Synthetic contraceptive hormones occlude the ability of nicotine to reduce ethanol consumption in ovary-intact female rats. Drug Alcohol Depend 252:110983. doi:10.1016/j.drugalcdep.2023.110983

Middeldorp J, Hol EM (2011) GFAP in health and disease. Prog Neurobiol 93 (3):421-443. doi:10.1016/j.pneurobio.2011.01.005

Namba MD, Kupchik YM, Spencer SM, Garcia-Keller C, Goenaga JG, Powell GL, Vicino IA, Hogue IB, Gipson CD (2019) Accumbens neuroimmune signaling and dysregulation of astrocytic glutamate transport underlie conditioned nicotine-seeking behavior. Addict Biol:e12797. doi:10.1111/adb.12797

Nishiyori A, Minami M, Ohtani Y, Takami S, Yamamoto J, Kawaguchi N, Kume T, Akaike A, Satoh M (1998) Localization of fractalkine and CX3CR1 mRNAs in rat brain: does fractalkine play a role in signaling from neuron to microglia? FEBS Lett 429 (2):167-172. doi:10.1016/s0014-5793(98)00583-3

Nnane IP (2005) PHARMACOKINETICS: Absorption, Distribution, and Elimination. In: Paul Worsfold AT, Colin Poole (ed) Encyclopedia of Analytical Science. 2nd edn. Elsevier, pp 126-133. doi:https://doi.org/10.1016/B0-12-369397-7/00458-1.

Paolicelli RC, Ferretti MT (2017) Function and Dysfunction of Microglia during Brain Development: Consequences for Synapses and Neural Circuits. Front Synaptic Neurosci 9:9. doi:10.3389/fnsyn.2017.00009

Paolicelli RC, Sierra A, Stevens B, Tremblay ME, Aguzzi A, Ajami B, Amit I, Audinat E, Bechmann I, Bennett M, Bennett F, Bessis A, Biber K, Bilbo S, Blurton-Jones M, Boddeke E, Brites D, Brône B, Brown GC, Butovsky O, Carson MJ, Castellano B, Colonna M, Cowley SA, Cunningham C, Davalos D, De Jager PL, de Strooper B, Denes A, Eggen BJL, Eyo U, Galea E, Garel S, Ginhoux F, Glass CK, Gokce O, Gomez-Nicola D, González B, Gordon S, Graeber MB, Greenhalgh AD, Gressens P, Greter M, Gutmann DH, Haass C, Heneka MT, Heppner FL, Hong S, Hume DA, Jung S, Kettenmann H, Kipnis J, Koyama R, Lemke G, Lynch M, Majewska A, Malcangio M, Malm T, Mancuso R, Masuda T, Matteoli M, McColl BW, Miron VE, Molofsky AV, Monje M, Mracsko E, Nadjar A, Neher JJ, Neniskyte U, Neumann H, Noda M, Peng B, Peri F, Perry VH, Popovich PG, Pridans C, Priller J, Prinz M, Ragozzino D, Ransohoff RM, Salter MW, Schaefer A, Schafer DP, Schwartz M, Simons M, Smith CJ, Streit WJ, Tay TL, Tsai LH, Verkhratsky A, von Bernhardi R, Wake H, Wittamer V, Wolf SA, Wu LJ,

Wyss-Coray T (2022) Microglia states and nomenclature: A field at its crossroads.
Neuron 110 (21):3458-3483. doi:10.1016/j.neuron.2022.10.020

Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR, 3rd, Lafaille JJ, Hempstead BL, Littman
DR, Gan WB (2013) Microglia promote learning-dependent synapse formation through
brain-derived neurotrophic factor. Cell 155 (7):1596-1609. doi:10.1016/j.cell.2013.11.030

Pawelec P, Ziemka-Nalecz M, Sypecka J, Zalewska T (2020) The Impact of the
CX3CL1/CX3CR1 Axis in Neurological Disorders. Cells 9 (10). doi:10.3390/cells9102277

Paxinos G, Watson C (2007) The rat brain in stereotaxic coordinates. 6th edn. Academic
Press/Elsevier, Amsterdam ; Boston ;

Peng Y, Guo G, Shu B, Liu D, Su P, Zhang X, Gao F (2017) Spinal CX3CL1/CX3CR1 May Not
Directly Participate in the Development of Morphine Tolerance in Rats. Neurochem Res
42 (11):3254-3267. doi:10.1007/s11064-017-2364-z

Robinson SP, Langan-Fahey SM, Johnson DA, Jordan VC (1991) Metabolites,
pharmacodynamics, and pharmacokinetics of tamoxifen in rats and mice compared to
the breast cancer patient. Drug Metab Dispos 19 (1):36-43

Rochefort H, Borgna JL (1981) Differences between oestrogen receptor activation by oestrogen
and antioestrogen. Nature 292 (5820):257-259. doi:10.1038/292257a0

Sadek SM, Khatri SN, Kipp Z, Dunn KE, Beckmann JS, Stoops WW, Hinds TD, Jr., Gipson CD
(2024) Impacts of xylazine on fentanyl demand, body weight, and acute withdrawal in
rats: A comparison to lofexidine. Neuropharmacology 245:109816.
doi:10.1016/j.neuropharm.2023.109816

Scofield MD, Heinsbroek JA, Gipson CD, Kupchik YM, Spencer S, Smith AC, Roberts-Wolfe D,
Kalivas PW (2016) The Nucleus Accumbens: Mechanisms of Addiction across Drug
Classes Reflect the Importance of Glutamate Homeostasis. Pharmacol Rev 68 (3):816-
871. doi:10.1124/pr.116.012484

Sher ME, Bank S, Greenberg R, Sardinha TC, Weissman S, Bailey B, Gilliland R, Wexner SD
(1999) The influence of cigarette smoking on cytokine levels in patients with
inflammatory bowel disease. Inflamm Bowel Dis 5 (2):73-78. doi:10.1097/00054725-
199905000-00001

Shi FD, Piao WH, Kuo YP, Campagnolo DI, Vollmer TL, Lukas RJ (2009) Nicotinic attenuation
of central nervous system inflammation and autoimmunity. J Immunol 182 (3):1730-
1739. doi:10.4049/jimmunol.182.3.1730

Soares AR, Picciotto MR (2024) Nicotinic regulation of microglia: potential contributions to
addiction. J Neural Transm (Vienna) 131 (5):425-435. doi:10.1007/s00702-023-02703-
94

Souter, E. A., Chen, Y. C., Zell, V., Lallai, V., Steinkellner, T., Conrad, W. S., Wisden, W.,
Harris, K. D., Fowler, C. D., & Hnasko, T. S. (2022). Disruption of VGLUT1 in Cholinergic
Medial Habenula Projections Increases Nicotine Self-Administration. eNeuro, 9(1),
ENEURO.0481-21.2021. https://doi.org/10.1523/ENEURO.0481-21.2021

Tarnowski BI, Spinale FG, Nicholson JH (1991) DAPI as a useful stain for nuclear quantitation.
Biotech Histochem 66 (6):297-302

Wakabayashi-Ito N, Nagata S (1994) Characterization of the regulatory elements in the
promoter of the human elongation factor-1 alpha gene. J Biol Chem 269 (47):29831-
29837

Wickel J, Chung HY, Ceanga M, von Stackelberg N, Hahn N, Candemir Ö, Baade-Büttner C,
Mein N, Tomasini P, Woldeyesus DM, Andreas N, Baumgarten P, Koch P, Groth M,
Wang ZQ, Geis C (2024) Repopulated microglia after pharmacological depletion
decrease dendritic spine density in adult mouse brain. Glia 72 (8):1484-1500.
doi:10.1002/glia.24541

Wieghofer P, Prinz M (2016) Genetic manipulation of microglia during brain development and
disease. Biochim Biophys Acta 1862 (3):299-309. doi:10.1016/j.bbadis.2015.09.019

Yi MH, Liu YU, Liu K, Chen T, Bosco DB, Zheng J, Xie M, Zhou L, Qu W, Wu LJ (2021)
Chemogenetic manipulation of microglia inhibits neuroinflammation and neuropathic
pain in mice. Brain Behav Immun 92:78-89. doi:10.1016/j.bbi.2020.11.030
Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, Strauss-Ayali D, Viukov S, Guilliams M,
Misharin A, Hume DA, Perlman H, Malissen B, Zelzer E, Jung S (2013) Fate mapping
reveals origins and dynamics of monocytes and tissue macrophages under
homeostasis. Immunity 38 (1):79-91. doi:10.1016/j.immuni.2012.12.001
Zhang B, Zou J, Han L, Beeler B, Friedman JL, Griffin E, Piao YS, Rensing NR, Wong M (2018)
The specificity and role of microglia in epileptogenesis in mouse models of tuberous
sclerosis complex. Epilepsia 59 (9):1796-1806. doi:10.1111/epi.14526
Zhao XF, Alam MM, Liao Y, Huang T, Mathur R, Zhu X, Huang Y (2019) Targeting Microglia
Using Cx3cr1-Cre Lines: Revisiting the Specificity. eNeuro 6 (4).
doi:10.1523/ENEURO.0114-19.2019