Title:

Pore-C sequencing identifies episome-driven chromosome conformation perturbations differentiating pneumococcal epigenetic variants

Author list:

Tze Yee Lim[1], Samuel T. Horsfield[1,2], Catherine M. Troman[1], Stephen D. Bentley[3], Min Jung Kwun[1,+] & Nicholas J. Croucher[1,+,*]

Affiliations:

[1] MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London, W12 0BZ, UK

[2] European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, CB10 1SA, UK

[3] Parasites & Microbes, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, UK

[+] These authors contributed equally

[*] Correspondence to n.croucher@imperial.ac.uk

<u>Abstract</u>

Streptococcus pneumoniae (pneumococcus) is a genetically diverse opportunistic bacterial pathogen that expresses two phase-variable loci encoding restriction-modification systems. Comparisons of two genetically-distinct pairs of epigenetically-distinct variants, each distinguished by a stabilised arrangement of one of these phase-variable loci, found the consequent changes in genome-wide DNA methylation patterns were associated with differential expression of mobile genetic elements (MGEs). This relationship was hypothesised to be mediated through changes in xenogenic silencing (XS) or nucleoid organisation. Therefore the chromosomal conformation of the both variants of each isolate were characterised using Illumina Hi-C, and Nanopore Pore-C, sequencing. Both methods concurred that the organisation of the pneumococcal chromosome was dominated by small-scale structures, with most pairwise interactions between loci <25 kb apart. Neither found substantial evidence for higher-order structure or XS in the pneumococcal genome, with more complex contact patterns only evident around the replication origin. Comparisons between the variants identified phage-related chromosomal islands (PRCIs) as the foci of differential contact densities between the variants. This was driven by copy number variation, resulting from variable excision and replication of the episomal PRCIs. However, the methods were discordant in their identification of the variant in which the PRCI was more actively replicating in both pairs. Validatory experiments demonstrated that the prevalence of circular PRCIs was not determined by DNA modification, but instead varied stochastically between colonies in both backgrounds, and was metastable during vegetative growth. PRCI excision was inducible by mitomycin C, but independent of the presence of a phage. Yet transcriptional activation of these elements was affected by both signals, indicating

1   transcription and replication are separately regulated. Therefore pneumococcal

2   MGEs do not appear to be subject to XS, resulting in heterogeneity being generated

3   within these bacterial populations through the frequent local disruption of

4   chromosome conformation resulting from the stochastic excision and reintegration of

5   episomal elements.

Author summary

The pneumococcus is a bacterium with a circular chromosome that is organised by

DNA-binding proteins and often contains mobile genetic elements (MGEs), genes

able to transmit between bacteria. All pneumococci encode defences against MGEs,

some of which create epigenetic modifications (typically methylation) genome-wide

at particular sequence motifs. Changes in these epigenetic patterns are associated

with altered MGE gene expression and replication in otherwise genetically-identical

bacteria. To test whether this was the result of methylation remodelling the

organisation of the chromosome, we compared the contact patterns across the

chromosome using two sequencing technologies. Both methods concurred that the

pneumococcal genome is generally folded into small structures, with the biggest

differences between the variants caused by the replication of MGEs. However, the

methods disagreed on the variant in which the MGEs replicated fastest. Further

experiments showed that MGE replication was stable over the course of culturing

over hours, but would randomly change level between days, explaining the

inconsistent observations. MGE replication was found to rise in response to DNA

damage, whereas gene expression also depended on the presence of other signals,

explaining the discrepancies in these activities between variants. Hence MGEs

significantly contribute to the heterogeneity that rapidly accumulates within

pneumococcal populations.

Introduction

Streptococcus pneumoniae (the pneumococcus) is a gram-positive commensal bacterium and opportunistic pathogen. Pneumococci are a common cause of diseases such as conjunctivitis, pneumonia, sepsis and meningitis [1]. The species has an extensive pangenome [2–4] as a consequence of exchange through transformation [5] and the frequent integration of mobile genetic elements (MGEs) into the pneumococcal chromosome [4]. However, the acquisition of genes can be inhibited by restriction modification systems (RMSs) [6,7]. Almost all pneumococci share two phase-variable Type I RMSs: the *Spn*III RMS encoded by the inverting variable restriction (*ivr*) locus [8,9], and the *Spn*IV RMS encoded by the translocating variable restriction (*tvr*) locus [4,9]. The rapid variation of these loci can inhibit the spread of prophage within clonally-related populations [10,11]. Yet the changes also cause global alterations to the methylation of the genome. Hence the sensitivity of pneumococcal transcription to epigenomic changes means these RMSs also cause phenotypic heterogeneity between isolates [9,12].

Variation in the *Spn*III RMS has been associated with changes in capsule expression [9,13,14], human carriage [15], adhesion to human cells [16], and virulence in a mouse model [9,17]. Variation at the *tvr* locus is instead associated with variation in transformation efficiency [12,18]. However, it has not been possible to identify methylation sites that can explain the transcriptional changes through proximal effects at gene regulatory loci [9,12]. Yet methylation has broader effects on the biophysical properties of DNA that means a global change in methylation patterns could affect the conformation of the bacterial nucleoid, and alter the interactions between regulatory proteins and DNA loci [19]. Hence it is plausible that the

epigenetic effects are instead mediated through longer-distance, larger-scale

changes in chromosomal conformation.

This is supported by analysis of two pneumococcal isolates, RMV7 and RMV8.

Mutations disrupting the recombinase that catalyses rearrangements at the *tvr* locus

enabled "locked" phase variants of each to be isolated: one representing the

common, dominant arrangement (RMV7$_{domi}$ and RMV8$_{domi}$), and one expressing a

rare arrangement (RMV7$_{rare}$ and RMV8$_{rare}$) [7]. These variants differ in the

methylation motif targeted by the *Spn*IV RMS, but are otherwise isogenic outside the

tvr loci. Both rare variants were more transformable than their dominant variant

counterparts, and in both cases this was at least partly attributable to the altered

activity of MGEs integrated into the chromosome [12,18]. RMV7 contains a cryptic

integrative and conjugative element next to *rpsI* (ICE$_{rpsI}$) and the antibiotic-resistance

transposon Tn*916*, as well as two phage-related chromosomal islands (PRCIs;

alternatively known as phage-inducible chromosomal islands, or PICIs) integrated

next to *dnaN* (PRCI$_{dnaN}$) and *uvrA* (PRCI$_{uvrA}$). PRCI$_{dnaN}$ was more transcriptionally

active in RMV7$_{domi}$, suppressing the induction of competence for transformation

through inducing stress responses [12]. RMV8 encodes a prophage (ɸRMV8), a

PRCI integrated within the *mal* operon (PRCI$_{malA}$), and a ClPhR prophage remnant

(previously annotated as PRCI$_{tadA}$) [18,20]. In RMV8$_{domi}$, reduced excision of ɸRMV8

resulted in greater expression of a prophage-modified RNA that suppressed

competence induction [18]. However, no specific methylation sites have been

identified that could explain this difference in MGE activity [12].

Hence host DNA conformation is a plausible mechanism by which epigenetic

modification could modulate the activity of MGEs through xenogeneic silencing (XS)

[21]. This process depends on nucleoid-associated proteins (NAPs) that

preferentially bind to the AT-rich DNA that is characteristic of MGEs [22,23], and are

capable of silencing their gene expression. As well as affecting MGE gene

transcription [23–25], these NAPs can modulate the excision and re-integration of

MGEs [26]. Such differences in XS activity could explain the inter-variant differences

observed in RMV7 and RMV8. At least four families of NAPs have been identified

[21,27–30]. Two are common in gram-negative *Proteobacteria*: H-NS, identified in

Escherichia coli [31], and MvaT, characterised in *Pseudomonas* [30]. By contrast,

Lsr2 has only been found in gram-positive *Actinobacteria* [29], and the Rok protein

has only been characterised in *Bacillus subtilis* [32]. Hence no XS system has yet

been characterised in *S. pneumoniae*. Nevertheless, the species encodes

orthologues of Rok, as well as other proteins that preferentially bind AT-rich DNA

[33]. Therefore the differential activity of MGEs in RMV7 and RMV8 may be

attributable to changes in the patterns of NAPs associated with the genome.

The conformation of the genome, and therefore the distribution of NAPs and regions

subject to XS, can be characterised using chromosome conformation capture (3C)

[34]. This approach typically involves crosslinking DNA to proteins with

formaldehyde, digesting the DNA with a frequently-cutting restriction enzyme, then

facilitating proximity ligation to link DNA fragments that were spatially close within the

cell (Fig. 1A). The subsequent use of paired-end Illumina data enables such contact

patterns to be assayed at a genome-wide scale using high-throughput 3C (Hi-C),

which infers instances of digestion and re-ligation within the DNA separating the two

reads based on their mapping positions and orientations relative to a reference

genome [35]. More recently, long-read data generated by Nanopore sequencing has

enabled 3C analyses to use information from concatemers consisting of multiple

DNA fragments ligated together [36,37]. Hence this Pore-C method makes it

possible to infer higher-order multiway chromatin interactions, and represents an

efficient method for generating data on the scale of a bacterial genome.

Therefore this study compared Hi-C and Pore-C approaches to inferring the

chromosomal conformation of the dominant and rare variants of *S. pneumoniae*

RMV7 and RMV8. These assays were employed to determine whether it is feasible

that the phenotypic heterogeneity associated with epigenetic variation can be

attributed to changes in XS, or other alterations in the distribution of NAPs across the

genome.

Results

Short-range interactions dominate pneumococcal Hi-C and Pore-C data

Three independent 3C preparations were generated for each of *S. pneumoniae*

RMV7$_{domi}$, RMV7$_{rare}$, RMV8$_{domi}$ and RMV8$_{rare}$ (Table S1, S2). These were digested with

*Nla*III, which can cut at sites found in the pneumococcal chromosome at intervals of

~300 bp (Table S3), and used to generate a 12-plex library for Nanopore sequencing.

However, this yielded few reads, as the activity of the pores declined rapidly during

the sequencing run (Fig. S1). Residual DNA from these samples was instead

processed for Illumina sequencing, and successfully generated Hi-C data (Table S2).

A further three independent 3C preparations were generated for each of the four

variants using *Mlu*CI, which can cut at sites found in the pneumococcal chromosome

at intervals of ~125 bp (Table S3). Previous applications of the Pore-C approach to

human cells found an additional protease treatment with pronase increased the

efficacy of de-cross-linking, thereby slowing the rate of nanopore blocking by proteins

attached to DNA [38]. Correspondingly, including this step in the sample preparation

protocol enabled more efficient data generation from individual flow cells (Fig. S1;

Table S2).

The three biological replicates for both methods for each of the variants were mapped

to the appropriate dominant variant reference genome (Fig. S2, S3; Table S1). The

necessity of mapping short read fragments meant the Pore-C data, which contains a

higher rate of sequencing errors, mapped less efficiently than the Hi-C data.

Nevertheless, a high proportion of the Nanopore data could be uniquely mapped to

locations within the pneumococcal genome (Fig. S3). To establish the highest

resolution at which analyses could be reliably conducted, the proportions of inferred

relative mapping orientations of sequence fragments were plotted against the

separation between their mapped locations if they came from the same read pair, for

Illumina Hi-C data, or from the same read, for Pore-C data. This found that sequence

fragments separated by less than 5 kb were typically mapped to opposite strands of

the genome for Illumina Hi-C data (Fig. S4), or to the same strand of the genome, for

Pore-C data (Fig. S5). These are the orientations expected for undigested DNA

sequenced with these technologies, consistent with only a subset of restriction sites

being cleaved during sample preparation. This is typical of 3C preparations, requiring

a convergence distance to be defined as the separation at which sufficient digestion

and re-ligation occurred to ensure all sequence fragment mapping orientations

become equally frequent [39]. This was estimated to be between 5 and 10 kb for both

technologies (Fig. S4-5).

Plotting the overall frequency of pairs against their separation highlighted particularly

strong interactions at an interval of ~5 kb (Fig. 1B), which may indicate a particular set

of atypical interactions. However, plotting the locations of these high-level short-range

interactions found they were distributed across the chromosome, and therefore may

represent false positive interactions resulting from incompletely-digested DNA (Fig.

S6). Therefore data were subsequently analysed at a resolution of 10 kb, to mitigate

any effects of DNA fragments that had not undergone digestion and religation.

This locus size was used to analyse the contact frequencies between all loci across

the chromosome for each replicate (Fig. S7-8), and combined across each set of

replicates (Fig. 1C). A strong diagonal was evident across all of the symmetrical

contact frequency matrices. This suggested most interactions occurred over distances

<25 kb, consistent with a simple genome organisation, dominated by short-range

structures. There were relatively few signals of higher-level chromosomal

organisation, apart from an elevated density of off-diagonal interactions evident

around the origin of replication (*ori*; Fig. 1D, S8). This was positioned ~1.6 Mb from

the beginning of the RMV7 reference, resulting in vertical and horizontal stripes across

the matrix (Fig. 1C). The RMV8 *ori* was at the start of the reference genome, resulting

instead in a framing effect, with elevated intensities in each corner of the matrix.

Focussing on the origin of replication suggested the presence of boundaries to

interactions that were consistent across technologies at ~40 kb or ~20 kb downstream

of *ori* in RMV7 and RMV8, respectively (Fig. 1D). These coincide with the location of

the only conserved *parS* site in this downstream replichore [40], which is more distant

from *ori* in RMV7 as a consequence of the integration of PRCI*dnaN* in the intervening

sequence [41]. The *parS* site is recognised by the ParB chromosome segregation

machinery, which is integral to organising the origin domain of the chromosomes of

the Bacillota species *B. subtilis* [42] and *S. pneumoniae* [40]. Hence this motif is likely

to underlie the observed disruption to chromosomal contacts in this region. Although

two *parS* sites are proximal to *ori* in the upstream replichore, these do not appear to

affect contact patterns in the same way, consistent with previous observations of *parS*

sites having distinct effects depending on their location [40,42]. Hence both Hi-C and

Pore-C data produced a consistent view of the contacts that dominate the

pneumococcal genome's conformation.

Differential interactions between variants span PRCIs

A genome-wide search for differences in chromosome conformation between the

variants was undertaken using contact matrices that summarised contact counts within

non-overlapping 10 bp loci. The DiffHiC (Fig. S9-S12) and multiHiCcompare (Fig. S13-

S16) methods were used, both of which fit generalised linear models to independent

biological replicate datasets to identify differences in contact frequencies [43,44].

However, they differ in their normalisation of the data [45]. Clustering of the biological

replicates by the diffHiC pipeline suggested the variants could be resolved from one

another by the genome-wide patterns of contacts in each comparison, except in the

Hi-C analysis of the RMV7 variants (Fig. S17). However, these separations were not

always associated with high bootstrap values, suggesting the differences were

attributable to specific loci, rather than broad changes in genome-wide contact

frequencies. Correspondingly, specific sites exhibiting significantly different contact frequencies between the genotypes, following Benjamini-Hochberg corrections for multiple testing, could be identified using Manhattan plots (Fig. 2A).

For RMV8, the only loci that were consistently identified as significantly differing between the two variants by all four comparisons were within, or adjacent to, PRCI$_{malA}$ [18]. The interacting loci, and the fold differences in contact frequency, were plotted against the annotation of this MGE (Fig. 2B). The hits were either between the flanking region and the element, or between the two flanks. However, while the spatial distribution and magnitude of the effects were similar across all comparisons, the Hi-C data suggested higher contact frequencies in RMV8$_{rare}$, whereas the Pore-C data suggested higher contact frequencies in RMV8$_{domi}$. To rule out sample misidentification as a cause of this discrepancy, it was verified that the patterns of contact frequencies at the *tvr* locus matched the arrangements characteristic of the variants, and that a simple analysis of contact frequency ratios within PRCI$_{malA}$ supported the genome-wide analysis results (Fig. S18). Detailed analysis of the Nanopore sequencing read lengths mapping to this region did not suggest this was an artefact of the Pore-C methodology preferentially sequencing shorter DNA fragments originating from the MGE (Fig. S19). Therefore the Hi-C and Pore-C analyses concurred that there was variation in the contact frequency within PRCI$_{malA}$, but disagreed on the variant in which the contact density was highest.

The comparison of the RMV7 variants provided an opportunity to test whether this observation could be independently replicated, as the analyses of Hi-C data were consistent in identifying PRCI$_{dnaN}$ [12] as the region differing most significantly

between RMV7$_{domi}$ and RMV7$_{rare}$ (Fig. 2A). This MGE was also the location of the most

significant difference in contact frequencies between the variants in the diffHiC

comparison of Pore-C datasets, in addition to being the site of multiple significant

differences in the multiHiCcompare analysis of the same data (Fig. 1D). The detailed

plot of these points relative to the PRCI$_{dnaN}$ annotation demonstrated these hits were

either within the element, or between the element and its flanking regions (Fig. 2B).

While the differences were again of similar magnitude and distribution across the two

methods, the Hi-C and Pore-C data were also discordant in identifying the variant in

which the contact frequency was highest. In this comparison, the Hi-C data suggested

the contact densities were higher in RMV7$_{domi}$, whereas the Pore-C data implied they

were higher in RMV7$_{rare}$. Therefore multiple significant differences in contact

frequencies between both pairs of variants were within or flanking PRCIs, although the

methods did not reach a consensus on the nature of the altered pattern of interactions.

Changes in PRCI contact frequency do not reflect changes in XS

These discrepancies between the datasets for each variant could represent either a

biological difference between the samples, or an artefact of the differing

methodologies used to generate the sequence reads. Consequently, a generalised

linear mixed-effects model was constructed that could resolve the effect of different

genetic and biological variables. This model was fitted to the four combinations of

sequencing methodology and genotype at a resolution of 10 kb using a zero-inflated

Poisson distribution with a logarithmic link function (see Methods; Fig. S20-4). The

fitted models were able to explain a substantial fraction of the variation in contact

frequencies across loci and datasets (Fig. S25). The effect of distance between loci

was estimated to be consistent between the genotypes, but contact frequencies

declined more rapidly with distance in the Hi-C data, suggesting the Pore-C data were

more efficient at detecting long-range interactions (Fig. 3A).

To test whether any of the differential contact frequency results were artefacts of the

restriction enzymes used to generate the samples, the model also estimated the

sensitivity of the inferred contact frequencies to the density of sites targeted by these

endonucleases (Fig. 3B). With the Pore-C data, there was no evidence of any strong

relationship between restriction sites and inferred contact density. With Hi-C data, a

higher density of *Nla*III sites was associated with an increased sensitivity for detecting

contacts, which is consistent with the experimental protocol. However, there was an

inverse relationship between the inferred contact density and the density of *Mlu*CI

sites, despite this enzyme not being involved in sample preparation. Given the AT-rich

nature of the *Mlu*CI sequence motif (AATT), such an observation may be attributable

to Illumina sequencing's bias towards DNA with a neutral GC content, which is not

shared with Nanopore data [46]. However, contact frequencies did not exhibit an

overall inverse relationship with GC content. Rather, contact frequencies increased

with rising GC content in RMV7, but showed the opposite relationship in RMV8 (Fig.

3C). Both trends were more extreme for the Hi-C data than the Pore-C data. This

suggested the Illumina data may be more affected by localised regions of extremely

AT-rich sequence, correlating with the presence of *Mlu*CI sites, rather than the mean

GC content averaged over a 10 kb locus. Nevertheless, no consistent evidence could

be found of a pneumococcal XS system that preferentially condensed AT-rich

sequences [47].

A more specific test for XS was quantifying the contact density across the MGEs within each of the genomes, relative to the rest of the chromosome. For RMV7, neither the Hi-C or Pore-C data found any evidence of elevated contact densities across the identified MGEs, with the confidence intervals of the level of contact density relative to the rest of the genome always spanning one (Fig. 3D). When comparing the variants, differences in the density of contacts were only inferred for PRCI$_{dnaN}$. In further agreement with the differential contact analyses (Fig. 2B), the contact frequencies were higher for PRCI$_{dnaN}$ in RMV7$_{rare}$ for the Pore-C data, but lower in the Hi-C data. For RMV8, the Illumina Hi-C data again found no evidence of the φRMV8 or PRCI$_{malA}$ MGEs being associated with elevated contact densities, although there was some evidence of higher contact frequencies within the latter in the Pore-C data. However, the Pore-C data also suggested that contact frequencies within PRCI$_{malA}$ were reduced in the rare variant, suggesting there was no consistent evidence for increased contact densities across both variants. By contrast, the Illumina Hi-C data suggested that PRCI$_{malA}$ was associated with higher contact densities in RMV8$_{rare}$. Therefore the model outputs were consistent with the differential contact analysis in both genotypes, and there was no evidence of elevated contact densities within the pneumococcal MGEs that could provide evidence of XS.

To test whether there was evidence of XS at any location within MGEs, locus-specific effects on contact frequencies were estimated for the sites within and flanking PRCI$_{dnaN}$ and PRCI$_{malA}$ (Fig. 3E). There was no evidence of heterogeneity within the elements, although the size of PRCI$_{malA}$ meant it contained only a single locus, making the elevated contact density inferred for this element in the Pore-C data highly susceptible to confounding by locus-level effects (Fig. 3D). There was also little

variation between the variants at these sites (Fig. 3F), which is consistent with the differences being captured by the MGE-level variables (FIg. 3D). However, variation was evident in the flanking regions, the direction of which mirrored the alterations in the elements themselves (Fig. 3D). Therefore the differential contact densities at the PRCIs was not caused by changing levels of XS-driven condensation, but instead reflected altered interactions between the MGEs and the proximal regions of the host chromosome.

Contact frequency changes are associated with altered PRCI activity

Such an observation suggested the differences between the variants may reflect changes in excision, replication and activation of the PRCIs. Given the conflicting data between the Hi-C and Pore-C methods, previously-collected RNA-seq data were used to identify the variants in which the MGEs were more active. PRCI$_{dnaN}$ had previously been identified as being transcribed at a significantly higher level in RMV7$_{domi}$ than RMV7$_{rare}$ during early exponential phase using both RNA-seq (Fig. 4A) and quantitative reverse transcriptase PCR (qRT-PCR) data [12]. Analogously, the PRCI$_{malA}$ genes were generally more highly expressed in RMV8$_{rare}$ than RMV8$_{domi}$ (Fig. 4B), although these differences were not consistent enough to reach genome-wide significance [18]. However, no difference was observed when the *tvr* loci of the variants were removed (Fig. 4C), indicating the differences might be associated with epigenetic modifications [18].

To validate whether transcription of PRCI$_{malA}$ substantially differed between the variants, qRT-PCR was used to assay the expression of *int*, encoding the element's integrase, and *dnaC*, encoding a DNA replication protein, during early exponential

(OD$_{600}$ = 0.2) and mid-exponential (OD$_{600}$ = 0.5) phase growth (Fig. 4D). Although no substantial difference was evident in early exponential phase, the transcription of the replication gene rose to be ~4-fold higher in RMV8$_{rare}$ than RMV8$_{domi}$ during mid-exponential phase, consistent with the higher expression of these genes in the same variant in the RNA-seq data (Fig. 4B). This suggested that PRCI$_{dnaN}$ and PRCI$_{malA}$ were less frequently transcribed in RMV7$_{rare}$ and RMV8$_{domi}$, respectively. Hence transcription, as well as contact densities, differed between epigenetic variants at these PRCIs.

The readily detectable levels of gene expression, which changed in response to growth, were also consistent with an absence of XS. Instead, these transcriptional signals indicated that the PRCIs were likely to be excising and replicating, which would explain the differences in contacts within these elements, and between them and their flanking regions [48]. Therefore qPCR assays were used to measure the ratios of excised to integrated elements, and the copy number of circularised elements within the cell (Fig. 4E,F). In RMV7, the ratio of excised to integrated PRCI$_{dnaN}$ was significantly higher in RMV7$_{rare}$, although there was no substantial difference in the number of circularized elements within the cell. In RMV8, both the ratio of excised to integrated elements, and the frequency of circularised PRCI$_{malA}$ genomes, were significantly higher in RMV8$_{domi}$. Therefore the altered contact frequency patterns between variants likely represented changes in the topological configuration of the elements.

For such a topological change to fully explain the significant differences of opposing magnitude in the Hi-C and Pore-C comparisons, it would need to be possible for

differences in PRCI circularization to be metastable [49]. Under such circumstances,

a variant may exhibit levels of PRCI circularization and replication that are stable over

enough bacterial generations such that separate biological cultures grown in parallel

have consistent properties. Yet the conflicting results between analyses demands that

it is also possible that re-isolation of the variant produces a genotype with a different

level of PRCI activity that is similarly persistent.

To test this, three independent colonies were cultured for each variant of RMV7 and

RMV8. Each of these cultures was frozen, used to inoculate overnight growth, then

diluted into three parallel tubes of fresh media until they reached exponential phase

growth. Measurements of the level of excision, and extracellular circularization, were

generally consistent across the parallel triplicate cultures inoculated from the same

isolate (Fig. 5A). However, there was substantial variation between cultures derived

from different colonies for RMV8. These results also contrasted with the previous

qPCR assays, which were performed on the same isolate used for the Pore-C analysis

(Fig. 4F). For RMV7, the results were more consistent across the colonies

representing each variant (Fig. 5A). However, RMV7$_{domi}$ was generally associated with

the higher median copy number across the replicates in this experiment, whereas this

value had been higher for RMV7$_{rare}$ in the previous assay (Fig. 4F). Therefore PRCI

copy number levels varied stochastically between even isogenic isolates, but are

metastable during culturing in liquid media.

PRCI excision and expression are separately regulated

The variation in contact frequency and copy number between isolates of the same

variant contrasted with the consistent differences in PRCI expression between RNA-

seq and qRT-PCR data experiments (Fig. 4B,D) [12]. Correspondingly, previous

passages of RMV7 genotypes, in which *tvr* loci with differing arrangements were

independently introduced, found phenotypic and expression differences were stable

over multiple days [12]. The topological arrangement of PRCI$_{dnaN}$ was assayed in the

same samples (Fig. 5B). This demonstrated substantial variation in the integration-

excision dynamics of this element across isolates of both variants. As these samples

were directly transferred between liquid cultures, this demonstrated the changes in

PRCI excision did not depend on culturing on solid media. Hence differences in PRCI

expression may be stable, despite stochastic variation in the topology of the element.

This implied the excision and transcription of PRCIs may be differentially regulated,

consistent with PRCI gene expression in at least some samples being highest in the

variants in which these elements were most frequently integrated into the chromosome

(Fig. 4D,F). This was tested using RMV8, which carries both the ɸRMV8 prophage

and PRCI$_{malA}$ [18]. RMV8$_{rare}$ cells were exposed to competence stimulating peptide

(CSP) and mitomycin C (MMC), both of which activate the ɸRMV8 prophage [20].

These assays were conducted with RMV8$_{rare}$ and a mutant in which ɸRMV8 had been

replaced with a chloramphenicol acetyltransferase (*cat*) resistance marker, to

ascertain whether the PRCI's activity was affected by the prophage (Fig. 5C). Both the

level of excision, and the concentration of circularised PRCI elements, rose in

response to MMC, but not CSP. This behaviour was independent of the presence of

a prophage within the same cell. This suggests the excision of PRCI$_{malA}$ is controlled

by the MGE itself in response to similar stimuli that activate prophage.

A qRT-PCR assay was then used to test whether MMC also activated PRCI$_{malA}$ *int* and

dnaC gene expression in both RMV8$_{rare}$ and RMV8$_{rare}$ ɸRMV8::*cat* (Fig. 5D). An

inconsistent rise in the transcription of both genes was recorded in MMC

supplemented cultures, correlating with the elevated copy number of the element (Fig.

5C). However, independently of treatment with MMC, transcription levels were

consistently lower in the absence of the prophage (Fig. 5D). This effect was greatest

for the *dnaC* replication gene, the median expression of which was 4.8-fold lower in

the absence of ɸRMV8 in unsupplemented media, and 3.4-fold lower in the absence

of ɸRMV8 in MMC-supplemented cultures. This is consistent with the PRCI's

dependence on the intact prophage for its ability to transmit between host cells [50].

Therefore expression of the PRCI is regulated by additional factors, associated with

the presence of prophage, that do not affect excision.

PRCIs do not cause a substantial fitness cost to their host cells

To test whether this regulation of PRCI activity limited their cost to their host cell, the

growth of mutants in which PRCIs had been replaced with a Janus cassette [51] was

compared to the corresponding unmodified genotypes in unsupplemented media (Fig.

6A). Neither RMV7$_{rare}$ PRCI$_{dnaN}$::Janus, nor RMV8$_{rare}$ PRCI$_{malA}$::Janus, exhibited

detectable changes in growth in unsupplemented media relative to the variant from

which they were derived. By contrast, removal of the prophage ɸRMV8 substantially

increased the growth of RMV8$_{rare}$ (Fig. 6B), consistent with the behaviour of a similar

RMV8 prophage null mutant [20]. The deletion of PRCI$_{malA}$ from this non-lysogenic

mutant caused no additional change. Therefore PRCIs cause a substantially smaller

impairment of host pneumococcus growth than prophage in the absence of an

activating stimulus.

Growth assays were then used to test whether this changed when PRCIs were

induced to excise from the chromosome by MMC during early exponential growth.

Both RMV7$_{rare}$ and the corresponding PRCI$_{dnaN}$::Janus grew similarly in the presence

of MMC, consistent with the MGE not reducing host cell fitness upon excision (Fig.

6C). Although MMC did cause a dose-dependent reduction in the density of RMV8$_{rare}$

cultures, this sensitivity was substantially reduced in a non-lysogenic φRMV8::*cat*

mutant (Fig. 6D). By contrast, removal of PRCI$_{malA}$ did not affect MMC sensitivity in

either the wild type or non-lysogenic RMV8 cells. Therefore stimulating the excision

and circularisation of PRCIs neither adversely affect pneumococcal host cell growth,

nor interfered with the lysis of cells following prophage induction.

<u>Discussion</u>

This application of both Hi-C and Pore-C methods to epigenetic variants of two *S.*

pneumoniae genotypes provides an overview of the pneumococcal chromosome's

organisation. Both methods concurred that the chromosomal contacts were dominated

by short-range interactions, consistent with a simple, locally-condensed structure.

However, the resolution at which these could be studied was limited by the density of

digestion and proximity re-ligation events. Future studies, using refined sample

preparation protocols, may enable higher-resolution analyses of bacterial

chromosomes.

At larger scales, some similarities were observed with data from *B. subtilis*, another

species within the Bacillota phylum [52], and a recently-published study of *S.*

pneumoniae R6 [53]. Both methods found evidence of an origin domain, indicative of

this region having a globular conformation, which included constrained contact

patterns (Fig. 1D) that likely reflected the positioning of *parS* sites, as seen in *B. subtilis*

[52] and *S. pneumoniae* [53]. The Pore-C data also identified a secondary diagonal

signal of long-range interactions that was orthogonal to that formed by the contacts

between neighbouring loci, previously determined as being indicative of a close

association of the chromosome arms (Fig. 1C) [52]. However, there was no evidence

of the chromosome interaction domains that have been noted not only in *B. subtilis*

[52], but also *E. coli* [54,55], *Caulobacter crescentus* [56] and *Mycoplasma*

pneumoniae [57]. This absence of discrete domains concurs with the recent study of

S. pneumoniae R6 [53]. It may be that the relatively small pneumococcal chromosome,

with strong coding biases on each replichore [58], is capable of orchestrating

concurrent transcription and regulation without substantial higher-order chromosomal

folding.

This comparatively simple structure may reflect the apparent paucity of NAPs in

pneumococci relative to *E. coli* and *B. subtilis*, with only the histone-like protein HU (or

HlpA) characterised in *S. pneumoniae* [59–62], and the relatively small number of

MGEs in a typically pneumococcal genome [4]. The lack of characterised orthologues

of many of the proteins involved in XS is consistent with the absence of condensed

regions that form boundaries between large domains in other species [52], and the

extensive disruption to pneumococcal chromosome conformation associated with the

depletion of HlpA [53]. Nevertheless, the epigenetic variants of RMV7 and RMV8 were

each distinguished by variable contact density at the sites of PRCI integration (Fig.

2A). However, the approximately two-fold increases in contact density at these

elements (Fig. 2B) was consistent with the *attP* locus of the circularised form having a

similar copy number to the chromosome itself (Fig. 4E, Fig. 5A), effectively doubling

the presence of the PRCI sequences. Hence the alternative explanation of

condensation by XS was unlikely. While it is possible that the high density of short-

range contacts identified chromosome-wide in our analyses prevents the identification

of greater condensation of loci, the absence of XS also explains the absence of any

detectable increase in contact density in AT-rich regions, or within other MGEs (Fig.

3C, D).

Furthermore, a lack of XS is consistent with the continued expression, excision and

activation of these PRCIs (Fig. 4, 5), and the larger φRMV8 prophage (Fig. 6), even in

the absence of inducing stimuli [18,20]. Such a difference in genome biology from *B.*

subtilis and *E. coli* suggests a fundamentally distinct interaction between MGEs and

host cells in these species. One possible explanation is that XS proteins have an

important role in controlling the activity of plasmids [63]. Such elements are common

in *E. coli* [64], found in ~10% of *B. subtilis* isolates [65], but rarer in *S. pneumoniae* [4].

Hence pneumococci may have adopted a different strategy for reducing the fitness

cost or transmissibility of their primarily chromosomally-integrated MGEs.

That the only consistent significant difference in contact frequencies between the

variants of RMV8 was caused by PRCI$_{malA}$ (Fig. 2A) contrasted with the previous RNA-

seq analysis, which only identified *comC* and φRMV8 as sites of significantly-differing

transcriptional variation between RMV8$_{domi}$ and RMV8$_{rare}$ [18]. However, the excision

of φRMV8 is linked to its increased expression, typically resulting in cell lysis [20]. By

contrast the excision of PRCI$_{malA}$ is reversible, without substantial fitness cost to the

host (Fig. 6), meaning these episomes are always likely to be hotspots of variation in

copy number even in cells growing in the same culture, thereby resulting in differential

contact densities. Additional significant differences in contact density were also

identified across the genome in the Pore-C data, with greater variation in RMV7 than

RMV8. This is consistent with the more widespread differences in transcription

between RMV7$_{domi}$ and RMV7$_{rare}$ than between the RMV8 variants [12,18], although it

may also reflect the higher and more even coverage across the RMV7 Pore-C

datasets (Fig. S8) providing greater power for detecting such variation. Hence

alterations in chromosomal conformation remains a feasible mechanism to explain

how distinct patterns of epigenetic DNA modification can affect transcription. However,

this observed correlation cannot establish causation without higher-resolution 3C data,

and further experimental work that can change both the DNA methylation patterns,

and the organisation of the genome.

However, the DNA modifications could not account for the differences in excision

dynamics, which were variable between different isolates of the same variant (Fig. 5A).

This diversity likely represents the PRCIs engaging in "bet hedging" behaviour [12,66],

as integrated elements are stably inherited following chromosomal replication, but are

at risk of deletion by homologous recombination [67]. Hence the vertical inheritance of

PRCIs is likely to be maximised by existing as both chromosomally-integrated and

episomal forms. That the *attB*/*attL* ratios were below one suggested most *attB* sites

contained a chromosomally-integrated form (Fig. 4D), ensuring they will be passed to

daughter cells in clonally-replicating populations. However, the *attP* levels were in the

range 0.3-5, implying that circular forms were also widespread, meaning the PRCI can

surivive even if the chromosomally-integrated copy is deleted by recombination.

Therefore these episomal dynamics can help explain the stable inheritance of PRCIs

by strains over decades, in contrast to the transient associations between prophage and host genotypes [4].

This accounts for why it is adaptive for the PRCIs to increase their excision in response to the detection of RecA-coated nucleoprotein filaments (Fig. 5C), which are generated by MMC, and mediate homologous recombination [20]. Yet PRCI excision does not cause the same fitness cost to the cell as prophage activation, because it cannot lyse the host in the absence of an intact "helper" phage (Fig. 5D). Therefore it is also adaptive for expression to be regulated in response to prophage activity, as was apparent in RMV8. This observed interdependence between the MGEs explains the rise in PRCI$_{malA}$ transcription in mid-exponential phase (Fig. 4F), as this is when ϕRMV8 is most active [68]. This distinction between the regulation of excision and activation means the former can occur stochastically, even in the absence of a prophage, as in RMV7. The observed epigenetic inheritance of a variable state suggests that the levels of excision are determined by the concentration of regulatory molecules being passed on from mother to daughter cells [69], although no mechanistic details were analysed in this study. Hence the discrepancy between the PRCI copy numbers detected in the Hi-C and Pore-C data is indicative of these genomes being dynamic systems, the behaviour of which is determined by metastable regulatory switches, which cannot by fully described by a single sequence alone.

Hence Pore-C will reveal additional heterogeneity within species, strains, and even clonal bacterial populations, with identical genomes being differentiated by their topology and copy number of individual elements. Although our analysis used data from two flow cells, one was sufficient to identify the key differences between the

studied genotypes, meaning this method can be implemented by any laboratory with

the capacity to use sequencing platforms on the scale of a MinION. Hence future

applications of this method to species with a greater diversity of replicons, NAPs and

XS systems promises to aid our understanding of the diversity of bacterial genome

biology across species.

Methods

Culturing of *S. pneumoniae*

All *S. pneumoniae* strains (Table S1) were cultured at 35°C in a 5% CO_2 atmosphere.

Culturing on solid media used Todd-Hewitt broth (Sigma-Aldrich) supplemented with

0.5% yeast extract (Sigma-Aldrich), 1.5% agar (Sigma-Aldrich) and 200 U ml^{-1}

catalase (Sigma-Aldrich). To select desired mutant genotypes, media were

additionally supplemented with antibiotics: streptomycin (Sigma-Aldrich) or kanamycin

(Sigma-Aldrich) at 400 µg ml^{-1}, or chloramphenicol (Sigma-Aldrich) at 4 µg ml^{-1}.

Unless stated otherwise, liquid culturing utilised a 'mixed' media consisting of Todd-

Hewitt broth supplemented with 0.5% yeast extract and Brain Heart Infusion broth

(Sigma-Aldrich) mixed at a 2:3 ratio [12]. For comparing PRCI dynamics between

variants, samples were collected during mid-exponential growth, at an optical density

at 600 nm (OD_{600}) of ~0.6. For measuring the PRCI responses to stimuli, either

unsupplemented media, MMC (850 ng), or competence stimulating peptide (CSP; 8.75

µg) was added to 7 ml of bacteria grown to an OD_{600} of ~0.2. The added CSP was

appropriate for inducing competence in each genotype: CSP1 for RMV7, and CSP2

for RMV8. The cultures were then incubated at 35ºC with 5% CO_2 for 2 h.

Subsequently, 5 ml of culture was used for RNA extraction, and 2 ml was centrifuged at 3220 g for 10 min to produce a cell pellet for DNA extraction.

To measure growth curves, 2×10^4 cells from titrated frozen glycerol stocks were grown in mixed liquid media in 96-well microtiter plates at 35ºC with 5% CO_2 for 20 h. For all experiments measuring bacterial sensitivity to inducing stimuli, 200 µl of bacterial cultures with an OD_{600} between 0.15 and 0.25 were mixed with MMC (10 ng), CSP (75 ng), or H_2O_2 (2 ng), then grown in microtiter plates at 35ºC with 5% CO_2 for 20 h. The OD_{600} was measured at 30 min intervals using a FLUOstar Omega microplate reader (BMG LABTECH). Three or four replicate wells were assayed for each tested genotype and condition in each experiment.

DNA purification and PCR amplification

Following overnight incubation, bacteria were pelleted by centrifugation at 3220 g for 10 min. Supernatants were discarded, and bacterial pellets were then resuspended in 480 µl of lysis buffer (Promega) and 120 µl of 30 mg ml^{-1} lysozyme (Sigma-Aldrich). Samples were incubated at 35ºC for 45 min and centrifuged for 2 min at 8000 g. Bacterial genomic DNA was then extracted using Wizard Genomic DNA Purification Kit (Promega), following the manufacturer's instructions. For experiments requiring subsequent PCR product purification, 1000 ng of genomic DNA, 25 µl of 2x DreamTaq Master Mix (Thermo Fisher), nuclease-free water (Thermo Fisher) and 2 µl each of the 10 µM forward and reverse primers listed in Table S4 were used for PCR amplification in a total reaction volume of 50 µl. Otherwise, 500 ng of genomic DNA, 7.5 µl of 2x DreamTaq Master Mix (Thermo Fisher), 1 µl each of a 10 µM forward and

reverse primers (Table S2), and nuclease-free water were combined in a total reaction

volume of 15 µl.

Construction of the *S. pneumoniae* RMV8 PRCI*malA* mutant

To generate the PRCI*malA*::Janus construct, the ~1 kb region flanking upstream (UP)

and downstream (DOWN) of PRCI*malA* were amplified by PCR using primers that

inserted restriction enzyme sites on the internal sides of each region. The UP region

was generated with primers malA_KO_UP_For and malA_KO_UP_ApaI, whereas the

DOWN region was generated with primers malA_KO_DOWN_For_BamHI and

malA_KO_DOWN_Rev. The Janus cassette was amplified with the corresponding

primers with added restriction sites (Table S4). Purified PCR amplicons were then

obtained via gel electrophoresis with a 1% agarose gel dyed with SYBR Safe (Thermo

Fisher) in Tris/Borate/EDTA buffer, followed by gel extraction with a GenElute Gel

Extraction Kit (Sigma-Aldrich).

Following extraction, purified PCR amplicons were digested at 37ºC for 2 h with the

appropriate restriction enzyme: ApaI (Promega) for UP, and BamHI (Promega) for

DOWN. Digested UP and DOWN products were then mixed with the digested Janus

cassette at a 2:2:1 ratio and incubated with T4 DNA ligase (New England Biolabs) at

room temperature (RT) overnight. The ligation mix was then used as the template for

a touchdown PCR, and the amplified PRCI*malA*::Janus construct was purified by

separation through electrophoresis on a 1% agarose gel, and extraction using a

QIAquick Gel & PCR Cleanup kit (Qiagen). To transform recipient bacteria (RMV8$_{rare}$

or RMV8$_{rare}$ ɸRMV8::*cat*) with the PRCI*malA*::Janus construct, a 1 ml sample of a

bacterial culture at an OD$_{600}$ of ~0.2 was incubated with 5 µl of 500 mM CaCl$_2$ (Sigma-

Aldrich), 1250 ng of CSP2, and 1 μg of the PRCI$_{malA}$::Janus construct at 35°C for 2 h.

Samples were then spread onto solid media supplemented with antibiotics (400 μg ml$^-$

^{1} of either kanamycin or streptomycin) to select for resistant transformants that grew

after incubation at 35ºC with 5% CO_2 for at least 16 h.

RNA purification and reverse transcription

A 5 ml sample of a bacterial culture was mixed with an equal volume of RNAprotect

(Qiagen) and incubated at room temperature for 5 min. RNA samples were extracted

using the SV Total Isolation System (Promega) according to the manufacturer's

instructions, with the exception that the initial extraction of RNA required treating cell

cultures with 480 μL 30 mg mL^{-1} lysozyme (Promega) for 30 min at 35 °C. The

purified RNA was treated with amplification-grade DNase I (Invitrogen), then used in

reverse transcription reactions using the First-Strand III cDNA synthesis kit

(Invitrogen). Each reaction used 0.2 μg RNA, 100 units of the SuperScript III reverse

transcriptase, 1 μL of 100 μM random hexamer primers (Thermo Fisher Scientific)

and 1 μL of 10 mM dNTP mix (Bioline). After a 5 min annealing period at 25 °C,

reverse transcription was conducted at 50 °C for 30 min, then 55°C for 30 min. The

reverse transcriptase was then heat inactivated at 70 °C for 15 min.

Quantitative PCR

The oligonucleotides used for qPCR were designed to generate amplicons that were

150-200 bp in length, and are listed in Table S4. Template cDNA was diluted in a

1:25 ratio with nuclease-free water (Qiagen). Each 15 μL reaction used 3.75 μL of

DNA template (either genomic DNA or cDNA), 0.75 μL of 10mM forward and reverse

primer solutions (Invitrogen), 7.5 μL of PowerUp SYBR Green Master Mix (Thermo

Fisher Scientific) and 2.25 µL of DNase-free and RNase-free water. Reactions were

run using MicroAmp Optical 96-well reaction plates (Thermo Fisher Scientific) and

the QuantStudio 7 Flex Real-Time PCR System (Applied Biosystems). The mixtures

were initially heated to 50 °C for 2 min. Subsequently, 40 amplification cycles were

run, each consisting of denaturation at 95 °C for 15 s, followed by annealing and

elongation at 60 °C for 1 min. The purity of the amplicon at the end of the reaction

was assessed through a melt curve analysis. In all experiments, three technical

replicate measurements were made for each of three biological replicates.

The expression of $PRCI_{malA}$ genes was quantified using the $\Delta\Delta Ct$ method. The $rpoA$

gene was used as the reference gene, against which target gene expression levels

were normalised to yield a ΔCt value [12]. The ΔCt values were calculated as the

difference between the mean Ct of the target gene and the reference gene across

technical replicates within the same biological replicate. The $\Delta\Delta Ct$ values were then

calculated relative to the reference ΔCt, which was that of $RMV8_{rare}$ at an OD_{600} of 0.2.

The fold gene expression difference between genotypes was then quantified as $2^{-\Delta\Delta Ct}$.

For the quantification of PRCI dynamics, the amplicons produced across the

integration site, and the $rpoA$ reference gene, were generated from genomic DNA

(Table S2). The concentration of each purified PCR product was measured using a

Qubit Broad Range Kit and Qubit 4 Fluorometer (Thermo Fisher). The Ct values for

copy numbers of 3×10^3, 3×10^4, 3×10^5 and 3×10^6 of these products were used to

generate a standard curve to convert the Ct values from experiments into absolute

copy numbers of DNA molecules.

Construction of Hi-C and Pore-C sequencing libraries

Three independent cultures of each of the four variants (*S. pneumoniae* RMV7$_{domi}$, RMV7$_{rare}$, RMV8$_{domi}$ and RMV8$_{rare}$) were grown to mid-exponential phase, corresponding to an OD$_{600}$ of ~0.60. The samples processed for Illumina Hi-C and Pore-C analyses were cultured independently. To crosslink DNA and proteins, cells were mixed with 1% v/v formaldehyde (Sigma-Aldrich) for 30 min at RT, followed by a 30 min incubation at 4ºC. The formaldehyde was then quenched through the addition of 1 ml 2.5 M glycine (Sigma-Aldrich), which was incubated at RT for 5 min, then at 4ºC for 15 min. Fixed cells were then collected by centrifugation at 3220 *g* for 10 min at 4ºC, flash frozen on dry ice, and stored at -80 ºC, to permit consistent processing of all samples in parallel.

Frozen pellets were thawed on ice, washed with 5 ml water, and centrifuged at 3220 *g* for 10 min. Pellets were then resuspended in 50 µl 30 mg ml^{-1} lysozyme, followed by incubation at 37ºC for 30 min. Both 10% v/v sodium dodecyl sulphate (SDS; Fisher Bioreagents) and 10% v/v Triton-X (Sigma-Aldrich) were added, followed by 10 min RT shaking incubation, then 10 min incubation at 55ºC. A 400 µl sample of the lysed cells was then combined with 130 µl of digestion mix, comprising NEB CutSmart buffer (NEB), 10% v/v Triton-X, and 1 U µl^{-1} of either *Nla*III (NEB; Illumina Hi-C library preparation), or *Mlu*CI (NEB; Pore-C library preparation). The chromatin was digested at 37ºC for 3 h. The *Nla*III enzyme was then heat inactivated through a 65ºC incubation for 20 min. The less stable *Mlu*CI enzyme did not require inactivation.

Digested chromatin was then ligated through the addition of 460 µL ligation mix, comprising ligation buffer (NEB), 0.1 µg µl^{-1} recombinant albumin (Promega), and 20

U µl^{-1} T4 DNA ligase (NEB). This mixture was incubated at RT for 1 h, followed by a

4ºC incubation for 48 h. After ligation, the DNA-protein crosslinks were reversed

through incubation of the mixture with 500 µl 20% Tween-20 (Sigma-Aldrich), 100 µl

10% v/v SDS, and 100 µl 20 µg µl^{-1} proteinase K (Sigma-Aldrich) at 65ºC overnight.

For the Pore-C library preparation, another round of protein denaturation was

performed by adding 118 µl 20 mg ml^{-1} pronase (Sigma-Aldrich) and 235 µl 10% v/v

SDS, followed by a 63ºC incubation for 1 h.

Following de-cross-linking, DNA was purified with 2.5 ml of a chilled 25:24:1 ratio

mixture of phenol:chloroform:isoamyl alcohol (Sigma-Aldrich), followed by separation

by centrifugation at 3220 g for 30 min at 4ºC. DNA was then precipitated from the

aqueous phase of each sample using 20 µl 5 M NaCl (Sigma Aldrich), 200 µl 3 M

sodium acetate (EMD Millipore Corporation) and 6 ml isopropanol (Sigma-Aldrich).

These mixtures were incubated at -80ºC overnight, then centrifuged at 3220 g for 30

min at 4ºC. The resulting pellets were washed with 1 ml of chilled 70% ethanol (VWR

Chemicals BDH), followed by a second centrifugation (3220 g, 15 min, 4ºC). All

remaining ethanol was removed through pipetting and drying at 65ºC in a heat block

for 15 min. The DNA was the resuspended in 100 µl nuclease-free water.

To enrich for longer molecules likely to contain multiple ligated DNA fragments, a 100

µl sample of each preparation was mixed with 50 µl of KAPA Pure Beads (Roche)

through vortexing, followed by incubation together at RT for 10 min. The beads were

then magnetically captured, and the supernatant discarded. The beads were washed

twice using 200 µl 80% ethanol and dried at RT for 4 min. DNA was then eluted from

the beads through a 10 min incubation in 100 µl of pre-heated nuclease-free water at

37ºC. The concentration and size distribution of the DNA molecules was checked

using an Agilent 2200 Tape Station (Agilent Technologies) and NanoDrop 1000

(Thermo Fisher).

DNA sequencing

Illumina Hi-C data were generated at the Wellcome Sanger Institute. Libraries were

prepared using standard approaches [70], enriching for a fragment size of

approximately 450 bp. Libraries corresponding to the three biological replicates for

each of the four variants were then combined into a 12-plex library, which was

sequenced on a single lane of the Illumina Novaseq 6000 platform. This generated

152 nt paired end reads.

Pore-C data were generated at Imperial College London using the native 12-plex

barcoding library preparation kit (Oxford Nanopore Technologies, ONT; product SQK-

NBD114.24) to process ~400 ng of DNA from each of the three biological replicates

for each of the four variants. Each DNA sample was individually treated to ensure

fragments were blunt ended with the NEBNext FFPE DNA Repair Mix (NEB) and the

NEBNext Ultra II End Repair/dA-tailing Module (NEB). Samples then underwent native

barcode ligation with NEB Blunt/TA Ligase Master Mix, before being pooled together.

The NEBNext Quick Ligation Module (NEB) was then used to attach sequencing

adapters. The completed libraries were then purified from the mixture using the short

fragment buffer to obtain DNA fragments of all sizes. The concentration of the final

DNA library was measured on a Qubit 4 fluorometer (Thermo Fisher). Two flow cells

(ONT product FLO-MIN114) were each used to sequence 15 fmol of the final library

using the MinION platform, recording data with a minimum quality score of 9 using high-accuracy basecalling with Guppy version 6.4.6 (ONT).

Bioinformatic processing

The Illumina Hi-C data were processed using version 2.1.0 of the hic workflow [71] implemented using Nextflow version 24.02.0 [72]. This workflow filtered the Illumina reads using FastQC [73], then mapped reads to the appropriate reference genome (Table S1) using bowtie2 [74], and generated raw contact maps using iced, as implemented within the HiC-Pro pipeline [75]. Genome-wide contact maps between loci were then generated at resolutions of 100, 500, 1000, 5000, 10000 and 50000 bp using cooler [76]. A quality control report was then generated with MultiQC [77], with additional visualisation files produced with juicer [78].

The Pore-C data were processed using version 1.1.0 of the wf-pore-c workflow [79] implemented using Nextflow version 23.10.1 [72]. This workflow mapped reads to the appropriate reference genome (Table S1) using minimap2 [80] with the non-default settings "-x map-ont -k 7 -w 5". These alignments were then processed with Pore-c-py [79] and pairtools [39] to generate raw contact maps, which were processed with cooler [76] and juicer [78], as for the Hi-C data.

Differential analysis of contact matrices

Analyses of differential contacts used matrices calculated at a resolution of 10 kb, with all self interactions along the matrix diagonals removed, and the *ivr* and *tvr* loci excluded, as they contained known rearrangements.

The diffHiC package [43] was applied using the recommended workflow [81], with modifications to parameter settings as appropriate for these datasets. Loci with fewer than five contacts across all replicates in each comparison were first removed, as there was low power to identify differential contacts at such positions. The counts across the filtered dataset were then normalised across replicates within each comparison between variants using a cyclic LOESS process, run for ten iterations. The biological coefficient of variation was then calculated across different contact densities, which informed the dispersion of the negative binomial generalised linear model that was fitted to the data. This used a tagwise, rather than trended, dispersion calculation, as there was not a strong relationship between variance and contact densities in some datasets. Differential contact densities between loci in the two variants were then identified from this model using quasi F-tests. The resulting *p* values were subject to a Benjamini-Hochberg correction for multiple testing to generate *q* values.

Similarly, analysis with the multiHiCcompare package [44] used a modified version of the recommended workflow [82]. This started by first both removing any loci for which the minimum frequency across replicates was zero. The remaining data were normalised using a cyclic LOESS process run for ten iterations. Significant differences were inferred through using a negative binomial exact test, as implemented in edgeR [83], with a Benjamini-Hochberg correction for multiple testing.

Statistical modelling of contact densities

The 10 kb resolution contact matrices were analysed using a generalised linear mixed effects model with the structure:

$$1 \quad \text{Contact frequency} \sim \text{Distance category} \; +$$

$$2 \quad \text{Distance from origin} + \text{GC} \; + \; Mlu\text{CI} \; + \; Nla\text{III} +$$

$$3 \quad \text{Locus classification} * \text{Variant} +$$

$$4 \quad (1|\text{Replicate}) \; + \; (\text{Variant}|\text{Position})$$

The terms all represent fixed effects, except those with the '|' notation, which denotes

a grouping variable for a random effect. Asterisks indicate an interaction between fixed

effects. A zero-inflated logarithmic Poisson link function was used to fit the model with

the glmmTMB package [84]. Parameter estimates reproducibly converged using the

default settings. However, attempts analyse the data using an equivalent model with

a negative binomial link function, which would account for any overdispersion in the

data, failed to converge on robust parameter estimates. Analysis of parameter

estimates used the sjPlot package [85], model fits were assessed with the

performance package [86], and data processing used the tidyverse packages [87].

Each interaction was categorised by the distance between the loci: <12.5 kb (ie.,

neighbouring 10 kb loci); 12.5-25 kb; 25- 50 kb, and >50 kb. Only loci within 100 kb of

each other were considered, to ensure parameter estimation was feasible, and avoid

the model fitting primarily to the noisy, sparse interactions between distant loci.

Loci were also classified based on the annotation of the corresponding reference

genomes. In both genomes, the four rRNA operons were annotated, as the difficulty

of mapping to these relatively GC-rich repeats reduces the detection of contacts

involving these loci. In RMV7, four MGEs were annotated: PRCI_{dnaN}, PRCI_{uvrA}, ICE_{rpsI},

and Tn*916* [12]. In RMV8, three MGEs were annotated: the prophage remnant CIPhR,

prophage ϕRMV8 and PRCI$_{malA}$ [18]. These classifications were fitted separately for each variant, to enable intervariant differences to be inferred at these loci.

As the DNA-protein crosslinks were fixed in mid-exponential phase, the detected contact frequencies were higher near the origin of replication, as some cells would have initiated, but not completed, chromosomal replication. Hence the number of detected contacts was expected to change with distance from the origin.

The additional fixed effects analysed the properties of each DNA locus, which were calculated using biopython [88]. The effect of the distribution of *Nla*III and *Mlu*CI sites was accounted for through calculating the mean number of restriction sites across the two interacting loci, and dividing this value by ten. The GC content was calculated as the mean proportion of bases in the two interacting loci, which was scaled upwards by a factor of ten. Similarly, the distance to the origin was calculated as the proportion of the genome between the site and the origin of replication, and this value was scaled upwards by a factor of ten when included in the model. These adjustments were necessary for numerical stability of the model fitting process.

The identity of the biological replicate was included as a random effect, to correct for differences in the amount of sequence data generated for each dataset. A second random effect estimated the corrected level of contact frequency for each locus in the dominant variant, and the difference in binding to each of these loci in the rare variant.

Data Availability

All raw genetic data used in this study are available from the ENA with the accession codes listed in Table S1 and S2. Processed files generated by bioinformatic pipelines are available from FigShare (https://figshare.com/projects/Hi-C_and_Pore-C_data_for_pneumococcal_epigenetic_variants/227616). Experimental results and sample information are available from Github (https://github.com/nickjcroucher/RMV_PoreC).

Code Availability

All code used in this analysis is available from Github (https://github.com/nickjcroucher/RMV_PoreC).

Acknowledgements

We thank Prof. Juanma Vaquerizas for helpful discussions about Hi-C data generation and analysis, and the Bespoke team at the Wellcome Sanger Institute for generating the Hi-C sequencing libraries.

Competing interests

I have read the journal's policy and the authors of this manuscript have the following competing interests: NJC has consulted for Antigen Discovery Inc., Merck and Pfizer, and has received an investigator-initiated award from GlaxoSmithKline; S.D.B. has consulted for Pfizer and Merck.

1   **References**

2   1.   Croucher NJ, Løchen A, Bentley SD. Pneumococcal Vaccines: Host Interactions,

3   Population Dynamics, and Design Principles. Annu Rev Microbiol. 2018;72: 521–549.

4   doi:10.1146/annurev-micro-090817-062338

5   2.   Hiller NL, Janto B, Hogg JS, Boissy R, Yu S, Powell E, et al. Comparative Genomic

6   Analyses of Seventeen *Streptococcus pneumoniae* Strains: Insights into the

7   Pneumococcal Supragenome. J Bacteriol. 2007;189: 8186 LP – 8195.

8   doi:10.1128/JB.00690-07

9   3.   Donati C, Hiller NL, Tettelin H, Muzzi A, Croucher NJ, Angiuoli S V, et al. Structure

10   and dynamics of the pan-genome of *Streptococcus pneumoniae* and closely related

11   species. Genome Biol. 2010;11: R107–R107. doi:10.1186/gb-2010-11-10-r107

12   4.   Croucher N, Coupland PG, Stevenson AE, Callendrello A, Bentley SD, Hanage WP.

13   Diversification of bacterial genome content through distinct mechanisms over different

14   timescales. Nat Commun. 2014;5: 5471. doi:10.1038/ncomms6471

15   5.   D'Aeth JC, van der Linden MPG, McGee L, de Lencastre H, Turner P, Song J-H, et al.

16   The role of interspecies recombination in the evolution of antibiotic-resistant

17   pneumococci. Elife. 2021;10: e67113. doi:10.7554/eLife.67113

18   6.   Johnston C, Martin B, Granadel C, Polard P, Claverys JP. Programmed Protection of

19   Foreign DNA from Restriction Allows Pathogenicity Island Exchange during

20   Pneumococcal Transformation. PLoS Pathog. 2013;9: e1003178.

21   doi:10.1371/journal.ppat.1003178

22   7.   Kwun MJ, Oggioni MR, De Ste Croix M, Bentley SD, Croucher NJ. Excision-

23   reintegration at a pneumococcal phase-variable restriction-modification locus drives

24   within- and between-strain epigenetic differentiation and inhibits gene acquisition.

25   Nucleic Acids Res. 2018;46: 11438–11453. doi:10.1093/nar/gky906

8. Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, et al. Complete

genome sequence of a virulent isolate of *Streptococcus pneumoniae*. Science (1979).

2001;293: 498–506. doi:10.1126/science.1061217

9. Manso AS, Chai MH, Atack JM, Furi L, De Ste Croix M, Haigh R, et al. A random six-

phase switch regulates pneumococcal virulence via global epigenetic changes. Nat

Commun. 2014;5: 5055. doi:10.1038/ncomms6055

10. Kwun MJ, Oggioni MR, Bentley SD, Fraser C, Croucher NJ. Synergistic Activity of

Mobile Genetic Element Defences in *Streptococcus pneumoniae*. Genes (Basel).

2019;10: 707.

11. Furi L, Crawford LA, Rangel-Pineros G, Manso AS, De Ste Croix M, Haigh RD, et al.

Methylation warfare: interaction of pneumococcal bacteriophages with their host. J

Bacteriol. 2019; JB.00370-19. doi:10.1128/JB.00370-19

12. Kwun MJ, Ion A V, Oggioni MR, Bentley SD, Croucher NJ. Diverse regulatory

pathways modulate bet hedging of competence induction in epigenetically-

differentiated phase variants of *Streptococcus pneumoniae*. Nucleic Acids Res.

2023;51: 10375–10394. doi:10.1093/nar/gkad760

13. Li J, Li JW, Feng Z, Wang J, An H, Liu Y, et al. Epigenetic Switch Driven by DNA

Inversions Dictates Phase Variation in *Streptococcus pneumoniae*. PLoS Pathog.

2016;12: e1005762. doi:10.1371/journal.ppat.1005762

14. Durmort C, Ercoli G, Ramos-Sevillano E, Chimalapati S, Haigh RD, Croix MDS, et al.

Deletion of the Zinc Transporter Lipoprotein AdcAII Causes Hyperencapsulation of

Streptococcus pneumoniae Associated with Distinct Alleles of the Type I Restriction-

Modification System. mBio. 2020;11: 10.1128/mbio.00445-20.

doi:10.1128/mbio.00445-20

15. De Ste Croix M, Mitsi E, Morozov A, Glenn S, Andrew PW, Ferreira DM, et al. Phase

variation in pneumococcal populations during carriage in the human nasopharynx. Sci

Rep. 2020;10: 1803. doi:10.1038/s41598-020-58684-2

16. Phillips ZN, Trappetti C, Van Den Bergh A, Martin G, Calcutt A, Ozberk V, et al. Pneumococcal Phasevarions Control Multiple Virulence Traits, Including Vaccine Candidate Expression. Microbiol Spectr. 2022;10: e0091622. doi:10.1128/spectrum.00916-22

17. Agnew HN, Atack JM, Fernando ARD, Waters SN, van der Linden M, Smith E, et al. Uncovering the link between the SpnIII restriction modification system and LuxS in *Streptococcus pneumoniae* meningitis isolates. Front Cell Infect Microbiol. 2023;13: 1177857. doi:10.3389/fcimb.2023.1177857

18. Kwun MJ, Ion A V, Cheng H-C, D'Aeth JC, Dougan S, Oggioni MR, et al. Post-vaccine epidemiology of serotype 3 pneumococci identifies transformation inhibition through prophage-driven alteration of a non-coding RNA. Genome Med. 2022;14: 144. doi:10.1101/2022.09.21.508813

19. Buitrago D, Labrador M, Arcon JP, Lema R, Flores O, Esteve-Codina A, et al. Impact of DNA methylation on 3D genome structure. Nat Commun. 2021;12: 3243. doi:10.1038/s41467-021-23142-8

20. Kwun MJ, Ion A V, Apagyi KJ, Croucher NJ. Chromosomal curing drives prophage evolution and bacterial epidemiology. bioRxiv. 2024; 2024.08.13.607808. doi:10.1101/2024.08.13.607808

21. Duan B, Ding P, Navarre WW, Liu J, Xia B. Xenogeneic Silencing and Bacterial Genome Evolution: Mechanisms for DNA Recognition Imply Multifaceted Roles of Xenogeneic Silencers. Mol Biol Evol. 2021;38: 4135–4148. doi:10.1093/molbev/msab136

22. Dorman CJ. H-NS, the genome sentinel. Nat Rev Microbiol. 2006;5: 157–161. doi:10.1038/nrmicro1598

23. Navarre WW. The Impact of Gene Silencing on Horizontal Gene Transfer and Bacterial Evolution. Advances in Microbial Physiology. Elsevier; 2016. pp. 157–186. doi:10.1016/bs.ampbs.2016.07.004

24. Wiechert J, Filipchyk A, Hünnefeld M, Gätgens C, Brehm J, Heermann R, et al.

Deciphering the Rules Underlying Xenogeneic Silencing and Counter-Silencing of

Lsr2-like Proteins Using CgpS of *Corynebacterium glutamicum* as a Model. mBio.

2020;11: e02273-19. doi:10.1128/mBio.02273-19

25. Navarre WW, McClelland M, Libby SJ, Fang FC. Silencing of xenogeneic DNA by H-

NS—facilitation of lateral gene transfer in bacteria by a defense system that

recognizes foreign DNA. Genes & Development . 2007;21: 1456–1471.

doi:10.1101/gad.1543107

26. Seah NE, Warren D, Tong W, Laxmikanthan G, Van Duyne GD, Landy A.

Nucleoprotein architectures regulating the directionality of viral integration and

excision. Proc Natl Acad Sci U S A. 2014/08/11. 2014;111: 12372–12377.

doi:10.1073/pnas.1413019111

27. Perez-Rueda E, Ibarra JA. Distribution of putative xenogeneic silencers in prokaryote

genomes. Comput Biol Chem. 2015;58: 167–172.

doi:10.1016/j.compbiolchem.2015.06.007

28. Winardhi RS, Fu W, Castang S, Li Y, Dove SL, Yan J. Higher order oligomerization is

required for H-NS family member MvaT to form gene-silencing nucleoprotein filament.

Nucleic Acids Res. 2012/07/13. 2012;40: 8942–8952. doi:10.1093/nar/gks669

29. Qu Y, Lim CJ, Whang YR, Liu J, Yan J. Mechanism of DNA organization by

Mycobacterium tuberculosis protein Lsr2. Nucleic Acids Res. 2013/04/10. 2013;41:

5263–5272. doi:10.1093/nar/gkt249

30. Tendeng C, Soutourina OA, Danchin A, Bertin PN. MvaT proteins in Pseudomonas

spp.: a novel class of H-NS-like proteins. Microbiology (N Y). 2003;149: 3047–3050.

doi:10.1099/mic.0.c0125-0

31. Wang W, Li G-W, Chen C, Xie XS, Zhuang X. Chromosome organization by a

nucleoid-associated protein in live bacteria. Science. 2011;333: 1445–1449.

doi:10.1126/science.1204697

32. Seid CA, Smith JL, Grossman AD. Genetic and biochemical interactions between the

bacterial replication initiator DnaA and the nucleoid-associated protein Rok in *Bacillus*

subtilis. Mol Microbiol. 2017/01/03. 2017;103: 798–817. doi:10.1111/mmi.13590

33. Solano-Collado V, Hüttener M, Espinosa M, Juárez A, Bravo A. Mga*Spn* and H-NS:

Two Unrelated Global Regulators with Similar DNA-Binding Properties. Front Mol

Biosci. 2016;3: 60. doi:10.3389/fmolb.2016.00060

34. Dekker J, Rippe K, Dekker M, Kleckner N. Capturing Chromosome Conformation.

Science (1979). 2002;295: 1306–1311. doi:10.1126/science.1067799

35. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, et

al. Comprehensive mapping of long-range interactions reveals folding principles of the

human genome. Science. 2009;326: 289–293. doi:10.1126/science.1181369

36. Ulahannan N, Pendleton M, Deshpande A, Schwenk S, Behr JM, Dai X, et al.

Nanopore sequencing of DNA concatemers reveals higher-order features of

chromatin structure. bioRxiv. 2019. doi:10.1101/833590

37. Li Z, Long Y, Yu Y, Zhang F, Zhang H, Liu Z, et al. Pore-C simultaneously captures

genome-wide multi-way chromatin interaction and associated DNA methylation status

in Arabidopsis. Plant Biotechnol J. 2022;20: 1009–1011. doi:10.1111/pbi.13811

38. Zhong J-Y, Niu L, Lin Z-B, Bai X, Chen Y, Luo F, et al. High-throughput Pore-C

reveals the single-allele topology and cell type-specificity of 3D genome folding. Nat

Commun. 2023;14: 1250. doi:10.1038/s41467-023-36899-x

39. Open2C, Abdennur N, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, et al.

Pairtools: From sequencing data to chromosome contacts. PLoS Comput Biol.

2024;20: e1012164.

40. Attaiech L, Minnen A, Kjos M, Gruber S, Veening JW. The ParB-parS chromosome

segregation system modulates competence development in Streptococcus

pneumoniae. mBio. 2015;6. doi:10.1128/mBio.00662-15

41. Kwun MJ, Ion A V, Oggioni MR, Bentley SD, Croucher NJ. Diverse regulatory

pathways modulate bet hedging of competence induction in epigenetically-

differentiated phase variants of *Streptococcus pneumoniae*. Nucleic Acids Res. 2023;51: 10375–10394. doi:10.1093/nar/gkad760

42. Marbouty M, Le Gall A, Cattoni DI, Cournac A, Koh A, Fiche J-B, et al. Condensin- and Replication-Mediated Bacterial Chromosome Folding and Origin Condensation Revealed by Hi-C and Super-resolution Imaging. Mol Cell. 2015;59: 588–602. doi:10.1016/j.molcel.2015.07.020

43. Lun ATL, Smyth GK. diffHic: a Bioconductor package to detect differential genomic interactions in Hi-C data. BMC Bioinformatics. 2015;16: 258. doi:10.1186/s12859-015-0683-0

44. Stansfield JC, Cresswell KG, Dozmorov MG. multiHiCcompare: joint normalization and comparative analysis of complex Hi-C experiments. Bioinformatics. 2019;35: 2916–2923. doi:10.1093/bioinformatics/btz048

45. Stansfield JC, Tran D, Nguyen T, Dozmorov MG. R Tutorial: Detection of Differentially Interacting Chromatin Regions From Multiple Hi-C Datasets. Curr Protoc Bioinformatics. 2019;66: e76. doi:10.1002/cpbi.76

46. Browne PD, Nielsen TK, Kot W, Aggerholm A, Gilbert MTP, Puetz L, et al. GC bias affects genomic and metagenomic reconstructions, underrepresenting GC-poor organisms. Gigascience. 2020;9: giaa008. doi:10.1093/gigascience/giaa008

47. Ali SS, Xia B, Liu J, Navarre WW. Silencing of foreign DNA in bacteria. Curr Opin Microbiol. 2012;15: 175–181. doi:https://doi.org/10.1016/j.mib.2011.12.014

48. Wang S, Lee S, Chu C, Jain D, Kerpedjiev P, Nelson GM, et al. HiNT: a computational method for detecting copy number variations and translocations from Hi-C data. Genome Biol. 2020;21: 73. doi:10.1186/s13059-020-01986-5

49. Newby J, Chapman J. Metastable behavior in Markov processes with internal states. J Math Biol. 2014;69: 941–976. doi:10.1007/s00285-013-0723-1

50. Martínez-Rubio R, Quiles-Puchalt N, Martí M, Humphrey S, Ram G, Smyth D, et al. Phage-inducible islands in the Gram-positive cocci. ISME J. 2017;11: 1029–1042. doi:10.1038/ismej.2016.163

51. Sung CK, Li H, Claverys JP, Morrison DA. An *rpsL* cassette, Janus, for gene replacement through negative selection in *Streptococcus pneumoniae*. Appl Environ Microbiol. 2001;67: 5190–5196.

52. Marbouty M, Le Gall A, Cattoni DI, Cournac A, Koh A, Fiche J-B, et al. Condensin- and Replication-Mediated Bacterial Chromosome Folding and Origin Condensation Revealed by Hi-C and Super-resolution Imaging. Mol Cell. 2015;59: 588–602. doi:10.1016/j.molcel.2015.07.020

53. Mazzuoli MV, Van Raaphorst R, Martin LS, Bock FP, Thierry A, Marbouty M, et al. HU promotes higher order chromosome organization and influences DNA replication rates in *Streptococcus pneumoniae*. Nucleic Acids Res. 2025;53. doi:10.1093/nar/gkaf312

54. Espeli O, Mercier R, Boccard F. DNA dynamics vary according to macrodomain topography in the *E.   coli* chromosome. Mol Microbiol. 2008;68: 1418–1427. doi:10.1111/j.1365-2958.2008.06239.x

55. Lioy VS, Cournac A, Marbouty M, Duigou S, Mozziconacci J, Espéli O, et al. Multiscale Structuring of the *E. coli* Chromosome by Nucleoid-Associated and Condensin Proteins. Cell. 2018;172: 771-783.e18. doi:10.1016/j.cell.2017.12.027

56. Le TBK, Imakaev M V, Mirny LA, Laub MT. High-Resolution Mapping of the Spatial Organization of a Bacterial Chromosome. Science (1979). 2013;342: 731–734. doi:10.1126/science.1242059

57. Trussart M, Yus E, Martinez S, Baù D, Tahara YO, Pengo T, et al. Defined chromosome structure in the genome-reduced bacterium *Mycoplasma pneumoniae*. Nat Commun. 2017;8: 14665. doi:10.1038/ncomms14665

58. Croucher NJ, Walker D, Romero P, Lennard N, Paterson GK, Bason NC, et al. Role of conjugative elements in the evolution of the multidrug-resistant pandemic clone *Streptococcus pneumoniae*[Spain23F] ST81. J Bacteriol. 2008/12/31. 2009;191: 1480–1489. doi:10.1128/jb.01343-08

59. Rode CK, Melkerson-Watson LJ, Johnson AT, Bloch CA. Type-specific contributions to chromosome size differences in *Escherichia coli*. Infect Immun. 1999;67: 230–236. doi:10.1128/IAI.67.1.230-236.1999

60. Wasim A, Gupta A, Bera P, Mondal J. Interpretation of organizational role of proteins on *E. coli* nucleoid via Hi-C integrated model. Biophys J. 2022/11/25. 2023;122: 63–81. doi:10.1016/j.bpj.2022.11.2938

61. Swetha RG, Sekar DKK, Devi ED, Ahmed ZZ, Ramaiah S, Anbarasu A, et al. *Streptococcus pneumoniae* Genome Database (SPGDB): A database for strain specific comparative analysis of *Streptococcus pneumoniae* genes and proteins. Genomics. 2014;104: 582–586. doi:10.1016/j.ygeno.2014.09.012

62. Ferrándiz M-J, Carreño D, Ayora S, de la Campa AG. HU of *Streptococcus pneumoniae* Is Essential for the Preservation of DNA Supercoiling. Front Microbiol. 2018;9: 493. doi:10.3389/fmicb.2018.00493

63. Dorman CJ. H-NS-like nucleoid-associated proteins, mobile genetic elements and horizontal gene transfer in bacteria. Plasmid. 2014;75: 1–11. doi:https://doi.org/10.1016/j.plasmid.2014.06.004

64. Denamur E, Clermont O, Bonacorsi S, Gordon D. The population genetics of pathogenic *Escherichia coli*. Nat Rev Microbiol. 2021;19: 37–54. doi:10.1038/s41579-020-0416-x

65. Zawadzki P, Riley MA, Cohan FM. Homology among nearly all plasmids infecting three Bacillus species. J Bacteriol. 1996;178: 191–198. doi:10.1128/jb.178.1.191-198.1996

66. Veening J-W, Smits WK, Kuipers OP. Bistability, epigenetics, and bet-hedging in bacteria. Annu Rev Microbiol. 2008;62: 193–210. doi:10.1146/annurev.micro.62.081307.163002

67. Croucher NJ, Mostowy R, Wymant C, Turner P, Bentley SD, Fraser C. Horizontal DNA Transfer Mechanisms of Bacteria as Weapons of Intragenomic Conflict. Barton NH, editor. PLoS Biol. 2016;14: e1002394. doi:10.1371/journal.pbio.1002394

68. Kwun MJ, Ion A V, Cheng H-C, D'Aeth JC, Dougan S, Oggioni MR, et al. Post-

vaccine epidemiology of serotype 3 pneumococci identifies transformation inhibition

through prophage-driven alteration of a non-coding RNA. Genome Med. 2022;14:

144. doi:10.1101/2022.09.21.508813

69. Martins BMC, Locke JCW. Microbial individuality: how single-cell heterogeneity

enables population level strategies. Curr Opin Microbiol. 2015;24: 104–112.

doi:https://doi.org/10.1016/j.mib.2015.01.003

70. Bronner IF, Quail MA. Best Practices for Illumina Library Preparation. Curr Protoc

Hum Genet. 2019;102: e86. doi:10.1002/cphg.86

71. Servant N, Ewels P, Garcia MU, Talbot A, Peltzer A, Miller E, et al. nf-core/hic: nf-

core/hic v2.1.0. Zenodo; 2023. doi:10.5281/zenodo.7994878

72. Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C.

Nextflow enables reproducible computational workflows. Nat Biotechnol. 2017;35:

316–319. doi:10.1038/nbt.3820

73. Andrews S. FastQC. 2023.

74. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9: 357–359. doi:10.1038/nmeth.1923

75. Servant N, Varoquaux N, Lajoie BR, Viara E, Chen C-J, Vert J-P, et al. HiC-Pro: an

optimized and flexible pipeline for Hi-C data processing. Genome Biol. 2015;16: 259.

doi:10.1186/s13059-015-0831-x

76. Abdennur N, Mirny LA. Cooler: scalable storage for Hi-C data and other genomically

labeled arrays. Bioinformatics. 2020;36: 311–316. doi:10.1093/bioinformatics/btz540

77. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048.

doi:10.1093/bioinformatics/btw354

78. Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, et al. Juicer

Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell

Syst. 2016;3: 95–98. doi:10.1016/j.cels.2016.07.002

79. EPI2ME. wf-pore-c. 2024.

80. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34: 3094–3100. doi:10.1093/bioinformatics/bty191

81. Lun ATL. DiffHiC: Differential analysis of Hi-C data User's Guide. 2017.

82. Stansfield JC, Dozmorov MG. multiHiCcompare Vignette. 2024.

83. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140. doi:10.1093/bioinformatics/btp616

84. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. {glmmTMB} Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. R J. 2017;9: 378–400. doi:10.32614/RJ-2017-066

85. Lüdecke D. sjPlot: Data Visualization for Statistics in Social Science, R package version 2.4.0. R package version 2.4.0. 2017.

86. Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. {performance}: An {R} Package for Assessment, Comparison and Testing of Statistical Models. J Open Source Softw. 2021;6: 3139. doi:10.21105/joss.03139

87. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

88. Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely available Python tools for computational molecular biology and bioinformatics. Bioinformatics. 2009;25: 1422–1423. doi:10.1093/bioinformatics/btp163

1　**Supporting Information Legends**

2　　　　　　　　　　　　　　　　　　　　　　　　　　- 49 -

3　**Supporting Information File 1** This file contains Figures S1-S25 and Tables S1-S4,

4　and their associated legends.

1   **<u>Figure Legends</u>**

Figure 1 Chromosomal conformation analysis of the epigenetic variants of two pneumococcal genotypes. (A) Diagram describing the methodology underpinning the use of high-throughput sequencing to capture chromatin conformation (3C). The final sequencing step distinguishes the Illumina (Hi-C) and Oxford Nanopore Technology (Pore-C) methods. **(B)** Relationship between the frequency of contacts between loci and their separation in the genome. Each panel represents a unique combination of sequencing technology and epigenetic variant. Each point corresponds to a detected pairwise interaction between loci in a contact matrix calculated at a 5 kb resolution for both technologies. The red dashed lines represent the thresholds used to define the closely-spaced loci between which particularly high contact densities were inferred. **(C)** Genome-wide Hi-C and Pore-C contact matrices shown at a 10 kb resolution. Each panel displays the cumulative number of contacts identified in three biological replicates for each unique combination of sequencing technology and epigenetic variant. The frequency of detected interactions between loci across the genome is represented by the colour of the corresponding cell in these symmetrical matrices. The black dashed lines indicate the position of the origin of replication. **(D)** Matrix detailing the cumulative number of contacts identified in loci surrounding the origin of replication, the position of which is marked by black dashed lines. The coloured dashed lines indicate the position of the *parS* site downstream of *ori*; the dotted lines show the position of the *parS* sites upstream of *ori*. Each cell in the matrix represents data from a 10 kb locus.

Figure 2 Genome-wide analysis of differential contact frequencies. (A)

Manhattan plots comparing the contact densities between epigenetic variants. For

each genotype, two different statistical methods (diffHiC and multiHiCCompare) were

used to compare the three biological replicates generated using each technology (Hi-

C and Pore-C) for both epigenetic variants. The negative logarithmic q values, corresponding to false discovery rates following a Benjamini-Hochberg correction for multiple testing, are shown for each locus across the genome. The threshold for genome-wide significance is shown by the red dashed horizontal line. The vertical black dashed lines show the boundaries of the variable PRCIs (PRCI$_{dnaN}$ in RMV7, and PRCI$_{malA}$ in RMV8) within the chromosomes. **(B)** Sites of differential contact frequencies identified by all four statistical comparisons within PRCI$_{dnaN}$ in RMV7, and PRCI$_{malA}$ in RMV8. Each point represents an interaction that was found to be significantly different between the corresponding dominant and rare variants following a correction for multiple testing. Its position on the horizontal axis represents the site of the interacting locus with the lower numerical position. The dashed line connects the point to the interacting locus with the higher numerical position. The vertical position of the point represents the base two logarithmic fold difference in the rare variant relative to the dominant variant, such that values greater than zero represent a higher contact frequency in the rare variant. The shapes of the points represent the method used to generate the data.

2 **Figure 3 Statistical modelling of contact frequencies.** A zero-inflated Poisson

3 generalised linear mixed effects model was fitted to the four combinations of

4 genotype and sequencing technology using contact matrices calculated at a

5 resolution of 10 kb. The two genotypes are distinguished by colours, and the two

6 technologies are distinguished by line and point styles. **(A)** Coefficients estimating

the effect of separation between loci on contact frequencies, normalised to the

expected contact frequency between neighbouring loci, shown by the horizontal line

at a value of one. **(B)** Coefficients estimating the effect of *Mlu*CI and *Nla*III site

densities on the detection of contacts; these coefficients reflect the impact of the

number of sites per 1 kb. The horizontal line at one represents the coefficient value if

the density of sites had no effect on the number of inferred contacts. **(C)** Coefficients

estimating the effect of the fraction of bases (scaled upwards 10-fold to aid model

fitting) that are guanine or cytosine (GC) on contact frequencies. **(D)** Coefficients

estimating the contact frequencies within mobile genetic elements. The top panel

shows the coefficients quantifying the relative contact frequencies in the MGEs,

normalised to the frequencies observed across the rest of the chromosome,

corresponding to the horizontal line at a value of one. The bottom panel shows the

coefficients estimating the contact frequencies at the same loci in the rare variant

relative to the dominant variant. **(E)** Locus-specific variation in the contact

frequencies across the loci encompassing the two PRCIs highlighted in Fig. 2:

PRCI$_{dnaN}$ within RMV7, and PRCI$_{malA}$ within RMV8. Data are plotted at 10 kb

intervals. The black vertical dashed lines indicate the boundaries of the PRCIs. **(F)**

Differences between the variants at the loci shown in (E).

Figure 4 Differential activity of PRCIs between variants. **(A)** Quantification of

PRCI*dnaN* expression in RMV7domi and RMV7rare using previously-published RNA-seq

data [12]. The annotated protein coding sequences (CDSs) within the PRCIs are shown by the arrows at the bottom of the panel, with their orientation indicating the direction in which they are transcribed. The lines in the main panel show the level of expression estimated for each CDS for each of the three independent biological replicates as normalised transcripts per million reads. The solid line indicates the median value for each variant, and the shaded area indicates the range between the maximum and minimum values. **(B)** Quantification of $PRCI_{malA}$ expression in $RMV7_{domi}$ and $RMV7_{rare}$ using previously-published RNA-seq data [18], as displayed in panel (A). **(C)** Quantification of $PRCI_{malA}$ expression in $RMV7_{domi}$ *tvr*::*cat* and $RMV7_{rare}$ *tvr*::*cat* using previously-published RNA-seq data [18], as displayed in panel (A). These two mutants do not have functional *Spn*IV RMSs. **(D)** Quantification of $PRCI_{malA}$ expression in $RMV7_{domi}$ and $RMV7_{rare}$ using qRT-PCR assays of transcription of an integrase (*int*) and replication (*dnaC*) CDS. Three technical replicates are shown for each of three biological replicates, as indicated by the shapes of the points. The violin plot summarises their distribution, with the median indicated by the horizontal line. Measurements were made at both early and mid-exponential timepoints. **(E)** Diagram showing the primers used to quantify the integration, excision and replication of the PRCIs. **(F)** Differences in integration, excision and replication of the PRCIs between variants. Data are shown as in panel (D). For both $PRCI_{dnaN}$, in RMV7, and $PRCI_{malA}$, in RMV8, the level of excision was quantified as the ratio of *attB* sites to *attL* sites A higher ratio indicates a greater level of excision. Similarly, the levels of circularised PRCIs were quantified by measuring the concentration of *attP*.

2 **Figure 5 Metastability and inducibility of PRCI activity**. **(A)** Violin plot showing

3 the variation in PRCI topology between cultures inoculated with independently-

isolated colonies of the four variants. Three biological replicate cultures were grown

from each of three independently-isolates colonies for each variant. Each violin plot

represents a single colony, and describes the distribution of three technical replicate

measurements for an individual biological replicate, with the shape of each point

indicating the biological replicate to which they relate. The horizontal line across

each violin represents the median value. **(B)** Violin plot showing the variation in

PRCI$_{dnaN}$ excision in a six-day passage of RMV7 genotypes. Four independent

mutants, in which the RMV8$_{domi}$ or RMV8$_{rare}$ *tvr* locus had been introduced into a *tvr*-

null RMV8 genotype, were passaged in parallel [12]. DNA was sampled from a

single biological replicate of each mutant at the start of the passage, and after two,

four and six days. Each violin summarises three technical replicate measurements of

a single biological replicate. **(C)** Violin plot showing the effect of inducing stimuli on

the topology of PRCI$_{malA}$. The excision and replication of PRCI$_{malA}$ following exposure

to competence stimulating peptide and mitomycin C were assayed two hours after

exposure in early exponential growth. These experiments used both RMV8$_{rare}$, and a

mutant derivative lacking the ϕRMV8 prophage. Data are shown as in panel (A). **(D)**

Violin plot quantifying the effect of mitomycin C on PRCI gene expression. The

expression of the *int* and *dnaC* genes of PRCI$_{dnaN}$ were assayed after two hours of

exposure to mitomycin C, in both RMV8$_{rare}$ and a mutant derivative lacking the

ϕRMV8 prophage. Data are shown as in Fig. 4.

Figure 6 The impact of induced and uninduced MGEs on the growth of RMV7

and RMV8 genotypes. Each panel shows the 20 h growth curve of a set of

pneumococcal genotypes, measured through assaying the optical density (OD_{600}) at

30 min intervals. In each plot, the solid line shows the median, and the shaded

region shows the full range observed across three biological replicates. **(A)** A

comparison of the effect of removing $PRCI_{dnaN}$ from $RMV7_{rare}$ and removing $PRCI_{malA}$

from $RMV8_{rare}$. **(B)** A comparison of the effect of removing ɸRMV8 from $RMV8_{rare}$

and $RMV8_{rare}$ $PRCI_{dnaN}$::Janus. **(C)** A comparison of the effect of exogenous stimuli

on the growth of $RMV7_{rare}$ and $RMV7_{rare}$ $PRCI_{dnaN}$::Janus. **(D)** The effect of

increasing concentrations of mitomycin C on $RMV8_{rare}$ and mutant derivatives lacking

ɸRMV8 or $PRCI_{malA}$.