Aging-associated Increase of GATA4 levels in Articular Cartilage is Linked to Impaired

Regenerative Capacity of Chondrocytes and Osteoarthritis

Meagan J. Makarczyk[1,2,#], Yiqian Zhang[1,3,#], Alyssa Aguglia[2], Olivia Bartholomew[2], Sophie

Hines[1,2], Suyash Sinkar[2], Silvia Liu[4,5], Craig Duvall[6], Hang Lin[1,2,7,*]

[1] Department of Orthopaedic Surgery, University of Pittsburgh School of Medicine, Pittsburgh,

PA, USA

[2] Department of Bioengineering, University of Pittsburgh Swanson School of Engineering,

Pittsburgh, PA, USA

[3] Xiangya Hospital, Central South University, Changsha, Hunan, China

[4] Department of Pharmacology and Chemical Biology, University of Pittsburgh School of

Medicine, Pittsburgh, PA

[5] Organ Pathobiology and Therapeutics Institute, University of Pittsburgh School of Medicine,

Pittsburgh, PA

[6] Department of Biomedical Engineering, Vanderbilt University, Nashville, TN, USA

[7] Bethel Family Musculoskeletal Research Center (BMRC), University of Pittsburgh School of

Medicine, Pittsburgh, PA, USA

[#] These authors contributed equally

*Correspondence:

Hang Lin, PhD

350 Technology Drive, Room 417, Pittsburgh, PA, 15219, USA

Phone: 412-624-5503; Fax: 412-624-5544

Email: hal46@pitt.edu

Abstract

Although the causal association between aging and osteoarthritis (OA) has been documented, our understanding of the underlying mechanism remains incomplete. To define the regulatory molecules governing chondrocyte aging, we performed transcriptomic analysis of young and old human chondrocytes from healthy donors. The data predicted that GATA binding protein 4 (GATA4) may play a key role in mediating the difference between young and old chondrocytes. Results from immunostaining and western blot showed significantly higher GATA4 levels in old human or mouse chondrocytes when compared to young cells. Moreover, overexpressing *GATA4* in young chondrocytes remarkably reduced their cartilage-forming capacity *in vitro* and induced the upregulation of proinflammatory cytokines. Conversely, suppressing *GATA4* expression in old chondrocytes, through either siRNA or a small-molecule inhibitor NSC140905, increased the production of aggrecan and collagen type II, and also decreased levels of matrix-degrading enzymes. In OA mice induced by surgical destabilization of the medial meniscus, intraarticular injection of lentiviral vectors carrying mouse *Gata4* resulted in a higher OA severity, synovial inflammation, and pain level when compared to control vectors. Mechanistically, we found that overexpressing GATA4 significantly increased the phosphorylation of SMAD1/5. Our work demonstrates that the aging-associated increase of GATA4 in chondrocytes plays a vital role in OA progression, which may also serve as a target to reduce osteoarthritis in the older population.

Keywords

 GATA4, chondrocyte, aging, osteoarthritis, DNA damage

Introduction

Aging is an inevitable phenomenon resulting in limited functionality, loss of structural integrity, and inability to effectively resist injury and disease.[1] Osteoarthritis (OA), the world's most common form of degenerative disease, has been closely associated with the advancement of age.[2] Specifically, OA is estimated to affect 32.5 million Americans, with most cases occurring in adults over the age of 45.[3] In fact, 1 in 3 people over the age of 65 are suffering from OA.[4] There are many contributors linked to the onset and progression of aging, such as organelle dysfunction and telomere shortening,[1] but the molecular mechanism underlying age-related OA has not been fully understood. Prior studies have demonstrated the role of DNA damage on chondrocyte aging due to oxidative stresses, resulting in an aged phenotype. [5,6] Aged cells can also undergo a state similar to permanent cell cycle arrest known as cellular senescence, which leads to a low-grade state of chronic inflammation and contributes to OA onset and progression in aged individuals.[6-8]

OA is now considered a whole joint disease, but cartilage degradation still represents the central feature.[3] The physiological role of articular cartilage is to support and protect the bones of diarthrodial joints through absorbing mechanical loads and facilitating frictionless movements. The extracellular matrix (ECM) of the cartilage is composed primarily of collagen type II and glycosaminoglycans (GAGs), and the constant breakdown and rebuilding of ECM components in the cartilage is common in healthy adults.[3,9] As we age, the regenerative response of chondrocytes begins to decline.[2] The whole joint nature of OA and the limited regenerative capacity of chondrocytes has contributed to the difficulty in developing disease-modifying osteoarthritis drugs (DMOADs). To date, no DMOADs have reached FDA approval.[3]

Currently, OA-associated changes in chondrocytes have been widely examined, which have significantly enhanced our understanding of this disease and promoted the development of potential treatments. However, OA is a combined consequence of different physiological stressors, including aging, injuries, obesity, etc. Therefore, aged chondrocytes in healthy humans without OA do not necessarily exhibit all the features of OA chondrocytes. In this study, to understand key regulators that contribute to an "aged" state in chondrocytes prior to OA onset, we first compared the transcriptome of young and old human chondrocytes isolated from healthy donors without joint diseases and employed unbiased analysis to define the key regulatory molecules that mediate chondrocyte aging. Assessment of the chondrocyte genome demonstrated the upregulation of several factors in aged chondrocytes, and transcription factor GATA binding protein 4 (GATA4)

specifically drew our attention since it is associated with DNA damage and cellular senescence.[5,10]

Upregulation of GATA4 in OA chondrocytes was also reported.[11] We thus hypothesized that the

increased GATA4 level contributes to chondrocyte aging and accelerated OA progression upon

injuries. We first tested the hypothesis by increasing or suppressing *GATA4* expression in healthy

chondrocytes and examining their cartilage-forming capacity. We also assessed the role of GATA4

in vivo by injecting lentiviral vectors carrying control or mouse *Gata4* genes into the knee joints

of OA mice induced by surgically created destabilization of the medial meniscus (DMM). Lastly,

a mechanistic study was conducted to explore how GATA4 impacts chondrocyte phenotypes and

functions.

Materials and Methods

Cell isolation and expansion

Healthy human knee cartilage tissues were harvested from arthritis-free donors through an

established protocol with the National Disease Research Interchange (NDRI). This study was

approved by the University of Pittsburgh Committee for Oversight of Research and Clinical

Training Involving Decedents. Cartilage was diced into ~ 1 mm^3 pieces with a scalpel and

incubated with a dissociation medium that was composed of high glucose Dulbecco's Modified

Eagle Medium (DMEM, Gibco/Thermo Fisher Scientific, Waltham, MA, United States), 2%

Antibiotics-Antimycotics (Life Technologies, Carlsbad, CA, United States), and collagenase type

II (1 mg/mL(w/v), Worthington Biochemical Corporation, Lakewood, NJ, United States). 10 mL

medium was used for 1g of cartilage, and the treatment lasted for 16 hours in a shaker at 37°C.

The mixture was then passed through a 70 μm strainer to collect single chondrocytes. Isolated

chondrocytes were seeded in tissue culture flasks at 1×10^4 cells/cm^2 and maintained in growth

medium (GM, DMEM containing 10% fetal bovine serum (FBS, Life Technologies) and 1%

Antibiotics-Antimycotics). After cells were fully attached to the culture substrate, the medium was

changed every three days until cells reached 70-80% confluency. Cells were detached with

Trypsin/EDTA (Gibco/ Thermo Fisher Scientific) and passaged.

Individual Chondrocyte Monolayer Culture

Chondrocytes isolated from healthy human cartilage were expanded to passage 1 (P1) and plated

in tissue culture 6 well plates at P2, treated with GM, and cultured until cells reached 80%

confluency. Media changes occurred every two days until cell collection. **Table S1** lists the
chondrocyte ages and genders used for different experiments.

Young and Old Chondrocyte Pools
Chondrocytes at passage 0 (P0) were pooled with three to four other young chondrocyte donors
and grown in tissue culture flasks with GM at a cell seeding density of $1\text{x}10^6$ cells per flask. Media
was changed weekly. Once cells reached 80% confluency, cells were collected using
Trypsin/EDTA. Some cells were frozen using Recovery Cell Culture Freezing Medium (Gibco/
Thermo Fisher Scientific), and other cells were used for subsequent passaging. The same methods
were used for the old (> 45 years) chondrocyte pool. **Table S1** lists the chondrocyte ages and
genders used in each pool. Due to the extensive amount of work conducted using these pools, two
separate chondrocyte pools were made.

RNA sequencing (RNA-Seq) and bioinformatics analysis
The transcriptomic differences among three young and three old chondrocyte donors were assessed
through RNA-Seq. Individual chondrocyte cultures (P0) were lysed with the QIAzol reagent
(Qiagen, German Town, MD, United States), and RNA was isolated from the lysate using an RNA
Easy Plus Universal Kit (Qiagen). Extracted RNA were quantitated Qubit™ RNA BR Assay Kit
(Thermo Fisher Scientific) followed by the RNA quality check using Fragment Analyzer (Agilent
Technologies, Santa Clara, CA). For each sample, RNA libraries were prepared from 500ng RNA
using the KAPA mRNA HyperPrep Kit (Roche, Indianapolis, IN) according to manufacturer's
protocol, followed by quality check using Fragment Analyzer (Agilent Technologies) and
fluorescent quantification on the Infinite F Nano + (Tecan, Männedorf, Switzerland). The libraries
were normalized and pooled, and then sequenced using NovaSeq6000 platform (Illumina, San
Diego, CA,) to an average of 50M 100PE reads.
Quality control was first applied to raw RNA sequencing reads by tool FastQC. Low-
quality reads and adapter sequences were filtered out by the Trimmomatic tool. Surviving reads
were then aligned to human reference genome hg38 using STAR aligner, and gene counts were
quantified. Differential expression analysis was performed based on gene counts by R package
"DESeq2" and DEGs were selected by adjusted p-value $< = 0.05$ and fold change $> = 1.5$. These
DEGs were then applied to Ingenuity Pathway Analysis (IPA) to detect enriched pathways. This

software employs databases of prebuilt pathways with known genes summarized from previous

studies, checks the overlap between the DEG list and known pathways and performs statistical

tests to determine the enrichment. Significant pathways were defined by p-value $<=0.05$.

Statistically stringently, FDR = 5% cutoff should be applied to control the false discovery rate. To

encourage more gene candidates, this study went by p-value$<=0.05$ and fold-change$>=1.5$ cutoff.

All the tools were run by default parameter settings.

Immunohistochemistry (IHC) to examine the GATA4 levels in native cartilage tissues.

Human cartilage tissues were fixed in 10% buffered formalin (Fisher Chemical, Fair Lawn, NJ) at

4°C overnight and then subjected to a graded ethanol dehydration series, starting from 20% ethanol

and progressing to 100% ethanol. Subsequently, they were embedded in Paraplast X-tra (Leica

Biosystems Inc. Richmond, IL). The Paraplast-embedded samples were sectioned at a thickness

of 6 µm using a rotary Leica microtome (Leica Microsystems Inc., Deerfield, IL, Model RM 2255).

Young and old mouse knee joints were gifts from Dr. Ana Mario Cuervo's lab at Albert Einstein

College of Medicine, which otherwise were wastes. After the specimens were fixed and decalcified

in a formic acid-based bone decalcifier (StatLab, Mckinney, TX, USA) for a period of two weeks,

they were then embedded and sectioned as described above.

For immunohistochemistry (IHC), the formalin-fixed paraffin-embedded sections first

underwent antigen retrieval based on different antibodies. Slides were then blocked with 10% goat

or horse serum (Abcam, Cambridge, MA) in phosphate buffered saline (PBS, Thermo Fisher

Scientific) for 1h, incubated at $4^{\circ C}$ overnight with the primary antibody against GATA4, then

incubated with a biotinylated anti-mouse/rabbit immunoglobulin G(IgG) secondary antibody for

1h, with signal detection via DAB substrate kit (Abcam). The Nikon Eclipse E800 upright

microscope (Melivile, NY, United States) was used to image the stained sections. Antibody

specifications can be found in **Table S2**.

Western blot to examine the GATA4 levels in human chondrocytes

P1 young and old chondrocytes were used for individual chondrocyte analysis of GATA4. Cells

were washed in pre-cooled PBS (Thermo Fisher Scientific) three times. Using the RIPA buffer

(Sigma-Aldrich) supplemented with the protease and phosphatase Inhibitor Single-Use Cocktail

(Gibco/Thermo Fisher Scientific) and a cell scraper, monolayer culture samples were collected. A

pestle was used to homogenize pellets in the RIPA cocktail for pellet culture samples. The protein

concentration of the supernatant was determined by the Pierce™ BCA Protein Assay Kit (Thermo

Scientific). Proteins were fractioned electrophoretically on the NuPAGE 4-12%, Bis-Tris Mini

Protein Gel (Gibco/Thermo Fisher Scientific) and then transferred to a polyvinylidene fluoride

(PVDF) membrane using the iBlot Dry Blotting System (Invitrogen, Waltham, MA, United States).

The membrane was blocked with 3% non-fat milk (Bio-Rad, Hercules, CA, USA), diluted with 1×

Tris-buffered saline (TBS, Gibco/ Thermo Fisher Scientific) and 0.1% Tween 20 (Sigma-Aldrich)

(TBST) at room temperature for 1.5 h, washed, and incubated with the primary antibody at 4°C

overnight on a rotating shaker. The membrane was washed 7 times for 3 min with TBST buffer

and incubated with horseradish peroxidase (HRP)-linked secondary antibodies (GE Healthcare

Life Sciences, Malborough, MA, United States) for 1.5 h at room temperature. After being washed

5 times with TBST, the membrane was incubated in the chemiluminescence substrate SuperSignal

West Dura Extended Duration Substrate (Thermo Fisher Scientific). Images were acquired using

the ChemiDocTM Touch Imaging System (Bio-Rad). Images were quantified using ImageJ.

Antibody information is included in **Table S2**.

Overexpression of *GATA4* in young human chondrocytes

P2 young, pooled chondrocytes or young, individual chondrocytes were transduced with the

lentiviral vector containing *GATA4* fused with dTomato gene or the control lentivirus carrying

EGFP for 10 hours. After that, flasks were rinsed with PBS for 2 times and the medium was

replaced by fresh GM. Both vectors were created and packed by VectorBuilder ($>10^8$ TU/mL,

VectorBuilder, Chicago, IL, United States). To detect the number of cells transduced, cultures were

imaged with an EVOS M5000 microscope (Thermo Fisher Scientific) after 72 h of initial

transduction. After transduction, western blot, RT-qPCR, and IHC were used to verify the stable

expression of GATA4 in cells.

RNA isolation and quantitative real-time polymerase chain reaction (RT-qPCR)

For pellet culture, samples were first rinsed with PBS twice and a pestle and electric pulverizer

were used to crush pellets. Cells were homogenized in Qiazol (Qiagen). Total RNA was isolated

and purified using the RNAeasy Plus Universal Mini Kit (Qiagen, Cat. NO. 74104) according to

the manufacturer's protocol. The reverse transcription to the complementary DNA (cDNA) was

accomplished using the SuperScript IV VILO Master Mix (Invitrogen). RT-qPCR was performed

on a real-time PCR instrument (QuantStudio 3, Applied Biosystems, Foster City, CA, United

States) using the SYBR Green Reaction Mix (Applied Biosystems) with custom primers ordered

from Integrated DNA Technologies (IDT, Newark, NJ, United States). Relative gene expression

levels were calculated through the $2^{-\Delta\Delta Ct}$ method. Ribosomal protein L13A (*RPL13A*) was used as

the housekeeping gene. Full names and abbreviations of genes and their corresponding proteins

are listed in **Table S3,** and primer sequences are listed in **Table S4**.

Pellet culture and chondrogenesis of young human Chondrocytes overexpressing *GATA4*

Following the lentiviral transduction of *GATA4* gene in young, pooled chondrocytes, the cells were

collected and formed into pellets at a cell seeding density of 3×10^5 cells per pellet. Pellets were

treated with chondrogenic medium (CM, DMEM with 1% v/v Insulin-Transferrin-Selenium-

Ethanolamine (ITS, Gibco/ Thermo Fisher Scientific), 1% antibiotic-antimycotics, 10^{-7} M

dexamethasone (Sigma-Aldrich), 40 µg/mL L-proline (Sigma-Aldrich), supplemented with 10

ng/mL transforming growth factor- β3 (TGF- β3, Peprotech, Rocky Hill, NJ, United States), and

50 µg/mL ascorbic acid-2-phosphate (Sigma-Aldrich)). Medium was changed daily for seven days.

RT-qPCR, histology, IHC, and western blot were used to characterize the tissues.

Safranin O/Fast green staining

Pellet samples were fixed in 10% buffered formalin (Fisher Chemical) for 2 h at room temperature

and then rinsed with PBS. Pellets then underwent serial dehydration in 30,50,70,95, and 100%

ethanol for one hour each. The 100% ethanol was refreshed once for an additional hour prior to

sample clearing in Xylene. Pellets were cleared in Xylene (Fisher Chemical) for two hours. Pellets

were then placed in Paraplast X-tra (Leica Biosystems Inc.) overnight. The next day, pellets were

embedded in Paraplast X-tra blocks and sectioned at a 6-µm thickness using a Leica microtome

(Leica Microsystems Inc., Model RM 2255).

Slides were stained using Safranin O (0.5%, Catalog number: 50240, Sigma-Aldrich), in 1%

acetic acid (Catalog number: A6283, Sigma-Aldrich), 0.005% fast green (0.05 g, Catalog number:

104022, Sigma-Aldrich) in 100 mL distilled water (Invitrogen) and counterstained with

Hematoxylin QS solution (Catalog number: H3404, Vector Laboratories INC). Imaging was

conducted using a Nikon Eclipse E800 upright microscope.

LUMINEX multiplex assays

Upon 7 days of chondrogenesis, condition medium from pellets was collected and flash frozen in liquid nitrogen and immediately stored in -80°C. LUMINEX assays were accomplished using the Bio-Plex 200 system (Bio-Rad). Data collection and analysis were conducted using the Bio-Plex Manager 6.1 software as established in previous studies.[12] LUMINEX kit information can be found in **Table S5**.

Knockdown of *GATA4* in old human chondrocytes

The siRNA targeting human *GATA4* (ON-TARGET plus Human GATA4 (2626) siRNA, J-008244-06-0005, Horizon Discovery Biosciences Limited, Cambridge, UK) was used in this study with a scrambled siRNA (ON-TARGET plus non-targeting siRNA #1, Catalog number: D-001810-01-05, Horizon Discovery Biosciences Limited) as the control. Old, pooled chondrocytes were transfected with the siRNA using Lipofectamine RNAiMAX reagent (Thermo Fisher Scientific). After 24 hours of incubation, the transfection medium (Opti-MEM Reduced Serum Medium, Thermo Fisher Scientific) was changed to GM. Transfected cells were collected after 48 hours for RT-qPCR to confirm the knockdown efficiency.

Pellet culture and chondrogenesis of old human chondrocytes with *GATA4* Knockdown

Following the 48h transfection of scrambled control siRNA or siRNA targeting *GATA4* , cells were collected using Trypsin/EDTA (Thermo Fisher Scientific) and pellets were made at a cell density of $3x10^5$ cells per pellet. Pellets were treated with CM supplemented with 10 ng/mL TGF- β3 and 50 µg/mL ascorbic acid-2-phosphate for 7 days. Upon day 7, pellets were collected for RT-qPCR, western blot, and IHC.

GATA4 Small Molecule NSC140905

GATA4 small molecule, NSC140905, also known as HCA 42027(Biosynth Ltd, Compton, United Kingdom), was reconstituted to 7.5 mM stock solution using UltraPure™ DNase/RNase-Free Distilled Water (Invitrogen) on a shaker at 37° until completely dissolved. Old, pooled chondrocytes were pelleted and treated with CM supplemented with 10 ng/mL TGF- β3 and 50

279 µg/mL ascorbic acid-2-phosphate with 100 µM NSC140905 for 14 days. Pellets were collected for

280 IHC, western blot, and RT-qPCR.

Animal model

All animal experiments were approved by the University of Pittsburgh Institutional Animal Care

and Use Committee (IACUC). Young (8 weeks) male C57BL/6 mice were purchased from Jackson

Laboratory (Bar Harbor, ME, USA) and maintained in pathogen-free conditions, with no more

than five mice per cage. Mice were provided ad libitum access to food and water, and a 12-hour

light/dark cycle to simulate natural circadian rhythms. To minimize bias, mice were randomly

assigned to either control or GATA4 groups, with 8 mice in each group.

Intraarticular injection

Intraarticular injections were administered under general anesthesia to safeguard the well-being of

the animals and to minimize procedural discomfort. Under general anesthesia with 2% isoflurane

in an oxygen mixture, the mice were placed in a supine position, and the right knee joint was

positioned at a 90-degree flexion to facilitate accurate injection into the joint space. The injection

site was meticulously identified medial to the patellar tendon. Using a 29-gauge needle, a volume

of 10 µL of lentiviral particles encoding either GATA4 or a control vector (at a concentration $>10^8$

TU/mL, VectorBuilder) was precisely administered into the intra-articular space of the right knee.

The precision of the injection was ensured by employing a consistent technique across all animals,

thereby reducing variability in the delivery of the viral vectors.

Destabilization of the medial meniscus (DMM) surgery

One week after viral vector injection, DMM surgery was performed to induce the OA model.[13,14]

Briefly, a medial parapatellar incision was made to expose the right knee joint, followed by a

careful opening of the joint capsule. The anterior medial menisco-tibial ligament was identified

and transected using micro scissors. The joint capsule and skin were subsequently sutured with 6-

0 silk thread. For experimental controls, a sham operation was performed in which the joint capsule

was exposed as in the DMM surgery, but the medial menisco-tibial ligament was left intact.

Knee hyperalgesia

We expected accelerated OA development after *Gata4* overexpression. To observe these differences, we used six weeks as the time point at which control mice had begun displaying mild OA symptoms. Knee hyperalgesia was evaluated using a Pressure Application Measurement (PAM) device (Ugo Basile, Varese, Italy).[15,16] Mechanical stimuli were applied to the mouse's knee joint to assess the degree of hyperalgesia based on the applied pressure and the mouse's response. Six weeks after either DMM or sham surgery, the mice were carefully removed from their cages to ensure they remained calm and unstimulated. The mouse was held securely by its back to maintain a straight posture, with the right knee joint flexed at approximately 90 degrees. The PAM device was placed on the index finger of the right hand of testers, which, along with the thumb, was used to apply pressure to the medial side of the knee joint, ensuring proper contact. Pressure was gradually applied at a constant rate of 30 g/s while monitoring the pressure curve displayed on the computer. The sensor was released immediately when the mouse exhibited a response to the applied stress, such as head movement, vocalization, or knee withdrawal. The pressure value displayed by the software, and the maximum pressures that mice can withstand were recorded. Two measurements were taken per knee, one on the medial side, and one on the lateral side the average was calculated for accuracy.

Synovial inflammation score

After 6 weeks, knee joints were collected and sectioned as described above. Histology staining and IHC were used to assess OA severity. Synovial inflammation score was evaluated according to a scoring protocol outlined in previous studies[17], which relies on the hematoxylin and eosin (H&E)-stained tissue sections.

Statistical Analysis

Each experiment was carried out with at least three biological replicates. Data is presented as mean ± standard deviation unless otherwise specified. Detailed information of sample size, pre-processing, and statistical methods have been specified in each figure legend. Prism 10 (GraphPad, San Diego, CA) was used for statistical analysis. The significance level was set at 0.05 and indicated by * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$), and **** ($p < 0.0001$).

Results

GATA4 is predicted to regulate chondrocyte aging

To limit the influence of *in vitro* expansion on cell phenotype, P0 human chondrocytes were used for RNA sequencing (**Figure 1A**). The volcano plot showed that 303 genes are upregulated and 163 genes are downregulated in aged chondrocytes compared to young cells (**Figure 1B**). The top 50 most changed genes are listed in **Figure 1C**. Interestingly, the osteoarthritis pathway was identified to be activated in old chondrocytes (**Figure 1E**).

Through Ingenuity Pathway Analysis (IPA) Upstream Regulator Analysis, transcriptional regulators that may mediate the difference between young and old cells were predicted, and the top 20 of them are shown in **Figure 1D**. Given that GATA4 was previously shown to regulate chondrocyte senescence[10] and be involved in the activation of Nuclear Factor Kappa B (NF-κB) pathway in tissues like the nucleus pulposus[18] and synovium,[19,20] it was selected for further investigation in this study. Initial data analysis predicted that GATA4 was upregulated in aged chondrocytes compared to young donors (**Figure 1D**). Immunohistochemistry (IHC) results indicated that GATA4 was more abundant in articular cartilage harvested from aged individuals (**Figure 1F**) (**Supplementary Figure S1**). Analyzing knee joints collected from young and old mice also demonstrated that GATA4 levels were higher in aged cartilage tissues compared to young (**Supplementary Figure S2**). To further confirm the findings, western blot was used to examine GATA4 levels in isolated chondrocytes, and the results indicated that aging correlates with the increase of GATA4 in chondrocytes (**Figure 1G**).

Figure 1: Bulk RNA sequencing to compare the transcriptome differences between young and old chondrocytes. *(A) Healthy chondrocytes were isolated from knee joint cartilage from young and old donors without osteoarthritis (assessed by experienced surgeons). P0 cells were used for RNA sequencing analysis. (B) Volcano plot demonstrating 303 upregulated and 163 downregulated genes in old chondrocytes when compared to young cells. (C) Top 50 genes that are significantly differently expressed in young and old chondrocytes. (D) Activation Z score of top 10 transcription regulators that are activated (positive) or inhibited (negative) in aged versus young chondrocytes. A comprehensive list of gene names can be found in **Table S3**. (E) Ingenuity Pathway Analysis (IPA) of young versus old cells. The gray bars indicate that no activity pattern is identified in IPA despite the highly significant association of the genes within the pathway. Orange, positive z-score; white, zero z-score. OBs=osteoblasts; OCs=osteoclasts. (F) GATA4 IHC of healthy human cartilage tissue from young and aged donors. Bar=50 μm. (G) Relative protein levels of GATA4 in P1 chondrocytes from individual human donors were analyzed by western blot.*

Overexpressing GATA4 impairs the hyaline cartilage formation capacity of young chondrocytes

To examine the functions of GATA4 in chondrocytes, we first overexpressed it in healthy young human chondrocytes (**Figure 2A**). Real-time quantitative PCR (RT-qPCR), immunostaining, and western blot confirmed the significantly increased GATA4 expression after infection, which did not impact the expression levels of collagen type II (COLII)-α1 (*COL2A1*) and aggrecan (*ACAN*), but significantly upregulated the expression of Collagen type X (COLX)-α1 (*COL10A1*) and India hedgehog (IHH), two representative chondrocytic hypertrophy markers (**Supplementary Figure S3**). We then examined the cartilage formation capacity of cells by culturing cells in chondrogenic medium for 7 days. IHC, RT-qPCR, and western blot demonstrated the continuous overexpression of *GATA4* in newly formed tissues (**Figure 2B-D**). Interestingly, significantly reduced GAG and COLII production was found in the overexpression *GATA4* group (GATA4 group) compared to the GFP control group (**Figure 2E-G**). Similar results were observed in the study testing chondrocytes from individual donors (**Supplementary Figure S4**). Interestingly, *GATA4* overexpression did not impact the expression of *COL10A1* and *IHH* in the cartilage pellets **(Figure 2G)**. Additionally, tissues from the GATA4 group secreted more proinflammatory cytokines, including IL-6, IL-8, TNF-α, and chemokine CCL2 (**Figure 2H&I**), as well as enzymes that can break down cartilage, including matrix metalloproteinases *(MMP)-1, 2, 12*, and a disintegrin and metalloproteinases *(ADAMTS) 5* (**Figure 2J&K**). Although the expression levels of *MMP-13* gene decreased in GATA4 group, the protein levels showed no difference between the two groups (**Supplementary Figure S5**).

Figure 2: **Impact of GATA4 overexpression on in vitro cartilage formation of young chondrocytes.** *(A) Timeline depicting the study. (B) IHC to assess GATA4 protein levels in young*

cells overexpressing GFP control or GATA4.   Scale Bar=50 µm. (C) RT-qPCR analysis of GATA4

gene expression in two groups. (D) Western blot to measure GATA4 protein levels. (E) Safranin O

staining and (F) Collagen type II (COLII) IHC to examine the production of cartilage matrix.

Scale Bar=50 µm. (G) RT-qPCR analysis of gene expression of cartilage matrix proteins aggrecan

(ACAN) and collagen type II-α1 (COL2A1) and hypertrophy markers collagen type X- α1

(COL10A1) and India hedgehog (IHH). (H) RT-qPCR analysis of gene expression of

proinflammatory cytokines, including interleukin (IL)-6, IL-8, and tumor necrosis factor- α (TNF-

α) (n=6). (I) Concentrations of IL-6, IL-8, and chemokine (C-C motif) ligand 2 (CCL2) in condition

medium (n=3). (J) RT-qPCR analysis of relative gene expression of matrix-degrading enzymes,

including matrix metalloproteinases (MMP)-1,2,3,12, and 13, and a disintegrin and

metalloproteinase (ADAMTS) 4 and 5 (n=6). (K) MMP-1 concentration in condition medium

(n=3). Student's two-tailed t-test with Welch's correction for standard deviation and a p-value of

0.05 was used for all statistical analysis.

GATA4 overexpression activates SMAD1/5

Increased SMAD1/5 phosphorylation represents a key feature of aged chondrocytes [21]. We

thus examined whether increased GATA4 levels are associated with SMAD1/5 activation (**Figure**

3A). In the experiment testing chondrocytes from individual donors (**Figure 3B, D-G**),

overexpression of GATA4, even without the stimulation of TGF-β3, was sufficient to activate

SMAD1/5, but not SMAD2/3. In addition, the group that was co-treated with GATA4

overexpression and TGF-β3 displayed the highest phosphorylated SMAD1/5 (pSMAD1/5) levels

in all tested groups. Of note, activation of SMAD2/3 was not impacted by GATA levels. A similar

trend was also observed in the study using pooled chondrocytes (**Figure 3C**).

Figure 3: Assessment of activation of SMADs following GATA4 overexpression. (A) Schematic showing the experiment. (B, C) Western blot to assess protein levels of phosphorylated SMAD1/5 (pSMAD1/5), phosphorylated SMAD2/3 (pSMAD2/3), and GATA4 in pellets derived from young (B)individual(Y1-3) or (C) pooled chondrocytes, which were infected with lentiviral vectors carrying GATA4 or control genes and then stimulated with (+) or without (-) TGF- β3 for two hours. Relative protein levels of (D)GATA4, (E)pSMAD1/5, and (F)pSMAD2/3 were semi-quantified using ImageJ (n=3). (G) The ratio of pSMAD1/5 compared to pSMAD2/3 was also calculated. Statistics were conducted using one-way Analysis of Variance (ANOVA) with Dunnett's post hoc analysis.

Suppressing GATA4 in old chondrocytes promotes ECM formation and lowers proinflammatory
cytokines

We then tested the potential of suppressing GATA4 in reversing chondrocyte aging (**Figure**
4A). Several GATA4 siRNAs were tested, and the one with the highest efficiency was selected
(**Supplementary Figure S6**). *GATA4* knockdown resulted in increased cartilage formation from
old chondrocytes, which did not influence the expression of hypertrophy marker *COL10A1*
(**Figure 4B-D**). Moreover, although we did not see a difference between scrambled control and
GATA4 siRNA groups regarding the expression of *IL-6* and *IL-8*, the protein level of IL-8 was
higher in the GATA4 group. Interestingly, the protein level of CCL2 was significantly decreased
after GATA4 knockdown (**Figure 4E, F**). We also tested the expression levels of MMPs and
ADAMTSs (**Figure 4H-I**). In general, suppressing GATA4 either decreased or caused no
significant changes to the levels of these enzymes. In particular, MMP-13 levels were reduced, in
both gene and protein levels (in condition medium), after GATA4 knockdown. Mechanistically,
GATA4 siRNA treatment also lowered the phosphorylation of SMAD1/5 and p-P65 (**Figure 4G**).

In a separate study, we used a small-molecule GATA4 inhibitor NSC140905, which was
shown to significantly promote cartilage formation from old chondrocytes and reduce the
expression of proinflammatory cytokines (**Supplementary Figure S7**). Taken together,
suppressing GATA4 partially restored the capacity of old chondrocytes to create new cartilage.

Figure 4: Influence of GATA4 knockdown on in vitro cartilage formation of old chondrocytes.
(A) Schematic showing the study. (B) Safranin O staining and (C) COLII IHC to examine the
production of cartilage matrix in the Scrambled Control or GATA4 siRNA group. Scale Bar=50
µm. (D) RT-qPCR analysis of relative gene expression of cartilage matrix proteins ACAN and
COL2A1 and hypertrophy marker COL10A1(n=6). *(E) RT-qPCR analysis of relative gene*

expression of proinflammatory cytokines IL-6 and IL-8 (n=6). (*F*) *Concentrations of IL-6, IL-8,*

and chemokine (CCL2) in condition medium (n=3). (*G*) *The relative protein levels of pSMAD1/5,*

pSMAD2/3, and phosphorylated p65 (p-P65) in two groups. (H) RT-qPCR analysis of relative gene

expression of matrix-degrading enzymes, including MMP-1,2,3,12, and 13, and ADAMTS 4 and 5

(n=6). *(I) MMP-1,2 and 13 concentrations in condition medium* (n=3). *Student's two-tailed t-test*

with Welch's correction for standard deviation and a p-value of 0.05 was used for all statistical

analysis.

Gata4 overexpression in the knee joints accelerates OA progression in mice

The physiological functions of GATA4 were further examined using a mouse model

(**Figure 5A**). Specifically, lentiviral vectors carrying *Gata4* or *mCherry* genes were injected into

the knee joints of young mice. One week after the injection, DMM surgery was conducted to

induce OA. Since we expected that Gata4 overexpression would accelerate OA progression, we

harvested samples for analysis 6 weeks after DMM.

Results from IHC indicated a successful Gata4 overexpression in hyaline cartilage even 6

weeks after injection (**Figure 5B, C**). Interestingly, mice overexpressing Gata4 in the knee joint

resulted in the elevation of p-P65 (**Figure 5D, E**), suggesting increased inflammation. Moreover,

the mice in the Gata4 group displayed a more severe OA and higher knee hyperalgesia than the

control group, revealed by a lower withdrawal threshold (**Figure 5F-H**). Interestingly, mice from

the Gata4 group displayed higher synovial inflammation (**Figure S8**).

Figure 5: Assessment of the physiological roles of Gata4 through intraarticular overexpression.
(A) schematic of the study. Mice received one intraarticular injection of lentiviral vectors that
carried mCherry or Gata4 gene one week before DMM surgery was performed. Knee joints were
harvested 6 weeks post-surgery. Levels of Gata4 (B, C) and p-P65 (D, E) were assessed with
IHC(B&D), and the staining was semi-quantitated with Image J (C&E). Cartilage degradation
was assessed with (F) Safranin O/ fast green (FG) staining, and (G)OARSI score was calculated.
(H)Knee hyperalgesia 6 weeks post-surgery. 507 g was the threshold baseline for non-surgery mice
(dashed line). Student's two-tailed t-test with Welch's correction for standard deviation and a p-
value of 0.05 was used for all statistical analysis.

Discussion

The causal relationship between aging and OA has been documented, and understanding
the molecular mechanisms is essential for the development of treatment methods. In this study, we
discovered the critical roles of aging-associated increases in GATA4 levels. Specifically,
overexpressing GATA4 in young chondrocytes impaired their capacity to form normal hyaline

cartilage, while suppressing GATA4 in old chondrocytes restored their chondrogenic potential. We also demonstrated that GATA4 functions partially by promoting the activation of SMAD1/5. Our *in vivo* work further confirmed that high *Gata4* expression accelerated OA progression in mice. Lastly, we defined a small-molecule GATA4 inhibitor that can partially restore the capacity of old chondrocytes to create healthy cartilage, representing a potential DMOAD for further validation in the future.

Given the recognized challenges in harvesting healthy cartilage tissues from donors without arthritis, there are limited reports investigating chondrocyte aging per se. The current findings include proliferation and post-expansion chondrogenic capacity reduction with aging,[22] increased MMP-13 production in response to catabolic stimuli,[23] and altered response to TGF-β.[24] One of our recent studies again demonstrated that aged chondrocytes displayed a reduced proliferation potential compared to young cells. Additionally, cartilage tissues generated by old chondrocytes contained more senescent cells than those from young cells.[25] However, to the best of our knowledge, there were no publications describing the transcriptomic comparisons between young and old chondrocytes, which would be informative in defining targets to stop or reverse chondrocyte aging.

Through transcriptomic analysis, we were able to assess the expression of different genes and genetic pathways that occur as chondrocytes age and found that GATA4 was predicted to have a higher z score in aged chondrocytes. The GATA family consists of several proteins (GATA1-6) with different variations of DNA-binding domains composed of zinc finger structures.[18] Particularly, GATA4, 5, and 6 are involved in the development of the mesoderm and endoderm tissues,[26] in which GATA4 has been detected in the development of structures such as the heart,[27,28] pancreas,[29] lung, and liver.[30,31] In addition, GATA4 and GATA6 are the only members that are associated with aging.[32] Particularly, GATA4 is unique in that it regulates tissues in a context-dependent manner and adopts a multifaceted role in the body, contributing to other age-related diseases such as atherosclerosis[33] and heart failure.[34]

In 2015, GATA4 was first introduced as a senescence regulator in a study conducted by Kang et al.[10] This work demonstrated the role of GATA4 in human fibroblasts and determined that the inhibition of autophagy caused GATA4 accumulation following DNA damage. They examined the presence of GATA4 in the human brain and found that there was an increase of GATA4 in the prefrontal cortex of aged human samples.[10] In fact, previous studies demonstrated that increased

GATA4 levels are associated with chondrocyte senescence,[11,35,36] an important change often
observed in OA chondrocytes. Moreover, suppressing GATA4 with siRNA was shown to
effectively abolish ionizing radiation-induced senescent phenotype in chondrocytes.[11] Of note, all
these studies relevant to chondrocytes were conducted *in vitro.* Herein, we, for the first time,
demonstrate the role of GATA4 in regulating chondrocyte aging.

A theory of chondrocyte aging proposed by van der Kramm suggests that alterations to the
TGF-β pathway induce chondrocyte hypertrophy and result in articular cartilage that is prone to
OA development.[24] While investigating the function of GATA4 in chondrocytes, we assessed how
its levels contribute to TGF-β alterations in chondrocytes and found that GATA4 levels negatively
correlated with the anabolic potential of chondrocytes (**Figures 2 and 4**). To maintain healthy
cartilage homeostasis, numerous pathways are involved. In particular, TGF-β is a crucial cytokine
involved in cartilage homeostasis during OA,[37,38] and aged chondrocytes respond differently to
TGF-β compared to their young counterparts.[39] Mechanistically, in the TGF-β pathway, TGF-β
binds to the heterotetrameric receptor complex, which can be grouped into three receptor types
(type I, type II, and type III). When TGF-β binds to its corresponding receptor, the activin-receptor-
like kinases (ALKs) are activated.[40] Typically, the anabolic binding of TGF-β to ALK4/5 results
in the phosphorylation of SMAD2/3, protecting chondrocytes from hypertrophy.[37] However, there
can also be catabolic effects associated with the expression of SMAD1/5, typically expressed when
TGF-β binds to ALK1/2/3/6.[37,40] The phosphorylation of SMAD1/5 results in the promotion of
ECM degrading proteins such MMP13.[21,37] As mentioned previously, research has demonstrated
that aged chondrocytes respond differently to TGF-β compared to young chondrocytes.[39,41]
Chondrocyte aging has been linked to the increase of pSMAD1/5 signaling.[21] These previous
studies inspired us to explore the potential association between GATA4 levels and the activation
of SMAD1/5. Our results found that GATA4 resulted in the phosphorylation of SMAD1/5 even
without TGF-β stimulation, which however did not alter the phosphorylation of SMAD2/3.
Although there are no current publications specifying the complex relationship of GATA4 and
SMAD1/5 in chondrocytes, a prior study reported that GATA4 was regulated by SMAD1/5.
Specifically, they can bind together to promote IL-6 expression in macrophages.[42]

While TGF-β pathway is closely associated with cartilage matrix remodeling, we also found
increases in the levels of multiple proinflammatory cytokines after GATA4 overexpression. We
investigated the NF-κB pathway since it is also activated during aging. [43,44] During the aging

process, NF-κB is thought to amplify and potentially propagate cellular senescence through the senescence-associated secretory phenotype (SASP).[45] A study by Qiao et al. indicated that GATA4 regulates NF-κB in dental pulp cells[46] and fibroblasts[47]. Specifically, using siRNAs, they determined that the knockdown of GATA4 decreased p65 production induced by lipopolysaccharide (LPS).[46] Other studies have investigated the role of GATA4 in the synovium and the progression of the disease state of rheumatoid arthritis and OA. Jia et al. showed that the knockdown of GATA4 attenuated synovial inflammation and joint damage in a collagenase-induced arthritis mouse model.[47] More recently, a study conducted by Chen et al. investigated the roles of GATA4 in fibroblast-like synoviocytes (FLS) and determined that GATA4 induced cellular senescence in FLSs and their role in OA progression.[20] Our study also discovered that the siRNA knockdown of GATA4 decreased the phosphorylation of p65 in aged chondrocytes (**Figure 4**). However, we also noticed that MMP-13 expression levels decreased in both *GATA4* overexpression and knockdown experiments. It should be noted that MMP-13 is constitutively produced in human chondrocytes but is only activated under pathological conditions. Given that MMP-13 is regulated by different transcriptional factors and cytokines, as well as RNAs,[21] its associations with GATA4 require further investigation in the future. Collectively, these findings further indicate that GATA4 might regulate aging partially through TGFβ and NF-κB pathways.

Our current study has not fully explored why aging promotes GATA4 expression. The study from Kang et al. indicated that DNA damage contributed to GATA4 accumulation.[10] It is known that DNA damage response (DDR) is regulated by ataxia telangiectasia mutated (ATM) and ataxia telangiectasia and Rad3–related (ATR) signaling.[10,48,49] The activation of these signaling pathways inhibits autophagy-related protein p62.[50] In addition, Copp et al. reported increased DNA damage in old chondrocytes.[5] These studies implied that DNA damage may be a reason for GATA4 upregulation. Our preliminary data support this possibility. Specifically, a DNA damage-inducing agent, doxorubicin, promoted the upregulation of GATA4 in chondrocytes (**Supplementary Figure S9**). To further link DNA damage to GATA4 accumulation, a study by Chung et al. used Tributyltin (TBT), a well-known endocrine-disrupting chemical, to induce DNA damage in articular chondrocytes. After 24 hours of TBT treatment, there was a significant increase in *GATA4* expression and expression of senescence markers.[35] Therefore, we believe that DNA damage due to aging is a key driver of the upregulation of GATA4 in old chondrocytes.

In conclusion, we have determined that GATA4 is increased in aged chondrocytes compared to young in both humans and mice, which may be induced by increased DNA damage observed in aged cells. We also demonstrated that GATA4 overexpression impaired the quality and quantity of cartilage created by chondrocytes and accelerated OA progression in mice. Conversely, suppressing GATA4 with siRNA or small-molecule inhibitors partially restores the capacity of old chondrocytes to form cartilage. Additionally, our study found that GATA4 can activate SMAD1/5 and change chondrocyte response to TGF-β. Overall, our study indicated that GATA4 could be a contributor to OA onset and progression in aged individuals, which can also serve as a potential target to prevent aging-associated OA.

There are some limitations of this study that can be further addressed in future studies. First, this study has allowed us to examine the potential mechanisms of GATA4 in chondrocyte aging. However, the relationship between GATA4 overexpression and SAMD1/5 activation should be further investigated to elucidate specific signaling mechanisms. Second, during our *in vivo* work, the intraarticular injection of GATA4 lentivirus was not tissue-specific. Therefore, the injection allowed for other cell types to overexpress GATA4. Future work should be conducted with cartilage-specific overexpression or depletion of GATA4 to clarify findings. Furthermore, studies using GATA4 knockdown do not demonstrate a complete reversal of aging in chondrocytes, and additional *in vivo* assessment needs to be conducted to verify whether GATA4 could be a therapeutic target for chondrocyte aging, in particular, the potential use of small molecule GATA4 inhibitor studied here. Lastly, our work indicates that GATA4 has a role in chondrocyte aging, but that does not negate the other aging factors and pathways. Combining current findings and investigating the prevalence of GATA4 in association with other aging molecules will further contribute to our understanding of the role of GATA4 in aging and possible OA onset.

Acknowledgments

This work was supported by the Department of Orthopaedic Surgery and the Orland Bethel Family Musculoskeletal Research Center (BMRC) at the University of Pittsburgh, as well as the University of Pittsburgh Center for Research Computing through the resources provided. This study used the Luminex Core Facility of UPCI. M.J.M was supported by the NIH T32 EB001026 - Cellular Approaches to Tissue Engineering and Regeneration, Cardiovascular Bioengineering Training Program T32-HL076124, and Ruth L. Kirschstein National Research Service Award

619   (NRSA) Individual Predoctoral Fellowship (Parent F31) 1F31AR083814 - 01A1. The authors

620   thank the funding support from the NIH (P30AG038072). The Chicago Center on Musculoskeletal

621   Pain Research Core Center, supported by NIH P30AR079206, provided training on pain-assessing

622   methods. Schematic figures were created at https://BioRender.com.

Data Availability

625   The Raw and processed RNA-seq data was uploaded to Gene Expression Omnibus (GEO) with

626   accession ID: GSE287861. The data will be publicly available when the manuscript is accepted.

References

1. Xiong, H., Hua, F., Dong, Y., Lin, Y., Ying, J., Liu, J., Wang, X., Zhang, L., and Zhang, J. (2022). DNA damage response and GATA4 signaling in cellular senescence and aging-related pathology. Front Aging Neurosci *14*, 933015. 10.3389/fnagi.2022.933015.
2. Rahmati, M., Nalesso, G., Mobasheri, A., and Mozafari, M. (2017). Aging and osteoarthritis: Central role of the extracellular matrix. Ageing Res Rev *40*, 20-30. 10.1016/j.arr.2017.07.004.
3. Makarczyk, M.J., Gao, Q., He, Y., Li, Z., Gold, M.S., Hochberg, M.C., Bunnell, B.A., Tuan, R.S., Goodman, S.B., and Lin, H. (2021). Current Models for Development of Disease-Modifying Osteoarthritis Drugs. Tissue Eng Part C Methods *27*, 124-138. 10.1089/ten.TEC.2020.0309.
4. Hawker, G.A., and King, L.K. (2022). The Burden of Osteoarthritis in Older Adults. Clin Geriatr Med *38*, 181-192. 10.1016/j.cger.2021.11.005.
5. Copp, M.E., Chubinskaya, S., Bracey, D.N., Shine, J., Sessions, G., Loeser, R.F., and Diekman, B.O. (2022). Comet assay for quantification of the increased DNA damage burden in primary human chondrocytes with aging and osteoarthritis. Aging Cell *21*, e13698. 10.1111/acel.13698.
6. Loeser, R.F., Collins, J.A., and Diekman, B.O. (2016). Ageing and the pathogenesis of osteoarthritis. Nat Rev Rheumatol *12*, 412-420. 10.1038/nrrheum.2016.65.
7. Ji, M.L., Jiang, H., Li, Z., Geng, R., Hu, J.Z., Lin, Y.C., and Lu, J. (2022). Sirt6 attenuates chondrocyte senescence and osteoarthritis progression. Nat Commun *13*, 7658. 10.1038/s41467-022-35424-w.
8. Franceschi, C., Bonafe, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E., and De Benedictis, G. (2000). Inflamm-aging. An evolutionary perspective on immunosenescence. Ann N Y Acad Sci *908*, 244-254. 10.1111/j.1749-6632.2000.tb06651.x.
9. Makarczyk, M.J. (2023). Cell Therapy Approaches for Articular Cartilage Regeneration. Organogenesis *19*, 2278235. 10.1080/15476278.2023.2278235.
10. Kang, C., Xu, Q., Martin, T.D., Li, M.Z., Demaria, M., Aron, L., Lu, T., Yankner, B.A., Campisi, J., and Elledge, S.J. (2015). The DNA damage response induces inflammation and senescence by inhibiting autophagy of GATA4. Science *349*, aaa5612. 10.1126/science.aaa5612.
11. Kang, D., Shin, J., Cho, Y., Kim, H.S., Gu, Y.R., Kim, H., You, K.T., Chang, M.J., Chang, C.B., Kang, S.B., et al. (2019). Stress-activated miR-204 governs senescent phenotypes of chondrocytes to promote osteoarthritis development. Sci Transl Med *11*. 10.1126/scitranslmed.aar6659.
12. Li, Z., Lin, Z., Liu, S., Yagi, H., Zhang, X., Yocum, L., Romero-Lopez, M., Rhee, C., Makarcyzk, M.J., Yu, I., et al. (2022). Human Mesenchymal Stem Cell-Derived Miniature Joint System for Disease Modeling and Drug Testing. Adv Sci (Weinh) *9*, e2105909. 10.1002/advs.202105909.
13. Glasson, S.S., Blanchet, T.J., and Morris, E.A. (2007). The surgical destabilization of the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse. Osteoarthritis Cartilage *15*, 1061-1069. 10.1016/j.joca.2007.03.006.

14.   Chen, D., Shen, J., Zhao, W., Wang, T., Han, L., Hamilton, J.L., and Im, H.J. (2017).
Osteoarthritis: toward a comprehensive understanding of pathological mechanism. Bone
Res *5*, 16044. 10.1038/boneres.2016.44.

15.   Barton, N.J., Strickland, I.T., Bond, S.M., Brash, H.M., Bate, S.T., Wilson, A.W.,
Chessell, I.P., Reeve, A.J., and McQueen, D.S. (2007). Pressure application measurement
(PAM): a novel behavioural technique for measuring hypersensitivity in a rat model of
joint pain. J Neurosci Methods *163*, 67-75. 10.1016/j.jneumeth.2007.02.012.

16.   Leuchtweis, J., Imhof, A.K., Montechiaro, F., Schaible, H.G., and Boettger, M.K. (2010).
Validation of the digital pressure application measurement (PAM) device for detection of
primary mechanical hyperalgesia in rat and mouse antigen-induced knee joint arthritis.
Methods Find Exp Clin Pharmacol *32*, 575-583. 10.1358/mf.2010.32.8.1532102.

17.   Hayer, S., Vervoordeldonk, M.J., Denis, M.C., Armaka, M., Hoffmann, M., Bäcklund, J.,
Nandakumar, K.S., Niederreiter, B., Geka, C., Fischer, A., et al. (2021). 'SMASH'
recommendations for standardised microscopic arthritis scoring of histological sections
from inflammatory arthritis animal models. Ann Rheum Dis *80*, 714-726.
10.1136/annrheumdis-2020-219247.

18.   Wang, X.H., Gao, J.W., Bao, J.P., Zhu, L., Xie, Z.Y., Chen, L., Peng, X., Zhang, C., and
Wu, X.T. (2022). GATA4 promotes the senescence of nucleus pulposus cells via NF-
kappaB pathway. Arch Gerontol Geriatr *101*, 104676. 10.1016/j.archger.2022.104676.

19.   Shi, W., Zheng, Y., Luo, S., Li, X., Zhang, Y., Meng, X., Huang, C., and Li, J. (2021).
METTL3 Promotes Activation and Inflammation of FLSs Through the NF-kappaB
Signaling Pathway in Rheumatoid Arthritis. Front Med (Lausanne) *8*, 607585.
10.3389/fmed.2021.607585.

20.   Chen, X., Gong, W., Shao, X., Shi, T., Zhang, L., Dong, J., Shi, Y., Shen, S., Qin, J.,
Jiang, Q., and Guo, B. (2022). METTL3-mediated m(6)A modification of ATG7 regulates
autophagy-GATA4 axis to promote cellular senescence and osteoarthritis progression.
Ann Rheum Dis *81*, 87-99. 10.1136/annrheumdis-2021-221091.

21.   Garcia de Vinuesa, A., Sanchez-Duffhues, G., Blaney-Davidson, E., van Caam, A.,
Lodder, K., Ramos, Y., Kloppenburg, M., Meulenbelt, I., van der Kraan, P., Goumans,
M.J., and Ten Dijke, P. (2021). Cripto favors chondrocyte hypertrophy via TGF-beta
SMAD1/5 signaling during development of osteoarthritis. J Pathol *255*, 330-342.
10.1002/path.5774.

22.   Barbero, A., Grogan, S., Schafer, D., Heberer, M., Mainil-Varlet, P., and Martin, I. (2004).
Age related changes in human articular chondrocyte yield, proliferation and post-
expansion chondrogenic capacity. Osteoarthritis Cartilage *12*, 476-484.
10.1016/j.joca.2004.02.010.

23.   Forsyth, C.B., Cole, A., Murphy, G., Bienias, J.L., Im, H.J., and Loeser, R.F., Jr. (2005).
Increased matrix metalloproteinase-13 production with aging by human articular
chondrocytes in response to catabolic stimuli. J Gerontol A Biol Sci Med Sci *60*, 1118-
1124. 10.1093/gerona/60.9.1118.

24.   van der Kraan, P.M. (2017). The changing role of TGFbeta in healthy, ageing and
osteoarthritic joints. Nat Rev Rheumatol *13*, 155-163. 10.1038/nrrheum.2016.219.

25.   Shen, H., He, Y., Wang, N., Fritch, M.R., Li, X., Lin, H., and Tuan, R.S. (2021).
Enhancing the potential of aged human articular chondrocytes for high-quality cartilage
regeneration. FASEB J *35*, e21410. 10.1096/fj.202002386R.

26.   Romano, O., and Miccio, A. (2020). GATA factor transcriptional activity: Insights from
717   genome-wide binding profiles. IUBMB Life *72*, 10-26. 10.1002/iub.2169.
718   27.   Dobrzycki, T., Lalwani, M., Telfer, C., Monteiro, R., and Patient, R. (2020). The roles
719   and controls of GATA factors in blood and cardiac development. IUBMB Life *72*, 39-44.
720   10.1002/iub.2178.
721   28.   Ang, Y.S., Rivas, R.N., Ribeiro, A.J.S., Srivas, R., Rivera, J., Stone, N.R., Pratt, K.,
722   Mohamed, T.M.A., Fu, J.D., Spencer, C.I., et al. (2016). Disease Model of GATA4
723   Mutation Reveals Transcription Factor Cooperativity in Human Cardiogenesis. Cell *167*,
724   1734-1749 e1722. 10.1016/j.cell.2016.11.033.
725   29.   Villamayor, L., Cano, D.A., and Rojas, A. (2020). GATA factors in pancreas development
726   and disease. IUBMB Life *72*, 80-88. 10.1002/iub.2170.
727   30.   Tremblay, M., Sanchez-Ferras, O., and Bouchard, M. (2018). GATA transcription factors
728   in development and disease. Development *145*. 10.1242/dev.164384.
729   31.   Molkentin, J.D. (2000). The zinc finger-containing transcription factors GATA-4, -5, and
730   -6. Ubiquitously expressed regulators of tissue-specific gene expression. J Biol Chem
731   *275*, 38949-38952. 10.1074/jbc.R000029200.
732   32.   Jiao, H., Walczak, B.E., Lee, M.S., Lemieux, M.E., and Li, W.J. (2021). GATA6 regulates
733   aging of human mesenchymal stem/stromal cells. Stem Cells *39*, 62-77.
734   10.1002/stem.3297.
735   33.   Mahmoud, M., Souilhol, C., Serbanovic-Canic, J., and Evans, P. (2019). GATA4-Twist1
736   Signalling in Disturbed Flow-Induced Atherosclerosis. Cardiovasc Drugs Ther *33*, 231-
737   237. 10.1007/s10557-019-06863-3.
738   34.   Katanasaka, Y., Suzuki, H., Sunagawa, Y., Hasegawa, K., and Morimoto, T. (2016).
739   Regulation of Cardiac Transcription Factor GATA4 by Post-Translational Modification in
740   Cardiomyocyte Hypertrophy and Heart Failure. Int Heart J *57*, 672-675. 10.1536/ihj.16-
741   404.
742   35.   Chung, Y.P., Weng, T.I., Chan, D.C., Yang, R.S., and Liu, S.H. (2023). Low-dose
743   tributyltin triggers human chondrocyte senescence and mouse articular cartilage aging.
744   Arch Toxicol *97*, 547-559. 10.1007/s00204-022-03407-x.
745   36.   Chung, Y.P., Chen, Y.W., Weng, T.I., Yang, R.S., and Liu, S.H. (2020). Arsenic induces
746   human chondrocyte senescence and accelerates rat articular cartilage aging. Arch Toxicol
747   *94*, 89-101. 10.1007/s00204-019-02607-2.
748   37.   Thielen, N.G.M., van Caam, A.P.M., HM, V.B., Vitters, E.L., van den Bosch, M.H.J.,
749   Koenders, M.I., van de Loo, F.A.J., Blaney Davidson, E.N., and van der Kraan, P.M.
750   (2023). Separating friend from foe: Inhibition of TGF-beta-induced detrimental
751   SMAD1/5/9 phosphorylation while maintaining protective SMAD2/3 signaling in OA
752   chondrocytes. Osteoarthritis Cartilage *31*, 1481-1490. 10.1016/j.joca.2023.08.004.
753   38.   van der Kraan, P.M., Goumans, M.J., Blaney Davidson, E., and ten Dijke, P. (2012). Age-
754   dependent alteration of TGF-beta signalling in osteoarthritis. Cell Tissue Res *347*, 257-
755   265. 10.1007/s00441-011-1194-6.
756   39.   Bauge, C., Girard, N., Lhuissier, E., Bazille, C., and Boumediene, K. (2014). Regulation
757   and Role of TGFbeta Signaling Pathway in Aging and Osteoarthritis Joints. Aging Dis *5*,
758   394-405. 10.14336/AD.2014.0500394.
759   40.   van Caam, A., Madej, W., Garcia de Vinuesa, A., Goumans, M.J., Ten Dijke, P., Blaney
760   Davidson, E., and van der Kraan, P. (2017). TGFbeta1-induced SMAD2/3 and SMAD1/5

phosphorylation are both ALK5-kinase-dependent in primary chondrocytes and mediated
by TAK1 kinase activity. Arthritis Res Ther *19*, 112. 10.1186/s13075-017-1302-4.

41. Wiegertjes, R., Thielen, N.G.M., van Caam, A.P.M., van Laar, M., van Beuningen, H.M.,
Koenders, M.I., van Lent, P., van der Kraan, P.M., van de Loo, F.A.J., and Blaney
Davidson, E.N. (2021). Increased IL-6 receptor expression and signaling in ageing
cartilage can be explained by loss of TGF-beta-mediated IL-6 receptor suppression.
Osteoarthritis Cartilage *29*, 773-782. 10.1016/j.joca.2021.01.008.

42. Lee, G.T., Kwon, S.J., Lee, J.H., Jeon, S.S., Jang, K.T., Choi, H.Y., Lee, H.M., Kim, W.J.,
Kim, S.J., and Kim, I.Y. (2010). Induction of interleukin-6 expression by bone
morphogenetic protein-6 in macrophages requires both SMAD and p38 signaling
pathways. J Biol Chem *285*, 39401-39408. 10.1074/jbc.M110.103705.

43. Yao, Q., Wu, X., Tao, C., Gong, W., Chen, M., Qu, M., Zhong, Y., He, T., Chen, S., and
Xiao, G. (2023). Osteoarthritis: pathogenic signaling pathways and therapeutic targets.
Signal Transduct Target Ther *8*, 56. 10.1038/s41392-023-01330-w.

44. Adler, A.S., Sinha, S., Kawahara, T.L., Zhang, J.Y., Segal, E., and Chang, H.Y. (2007).
Motif module map reveals enforcement of aging by continual NF-kappaB activity. Genes
Dev *21*, 3244-3257. 10.1101/gad.1588507.

45. Rahman, S.M.T., Aqdas, M., Martin, E.W., Tomassoni Ardori, F., Songkiatisak, P., Oh,
K.S., Uderhardt, S., Yun, S., Claybourne, Q.C., McDevitt, R.A., et al. (2022). Double
knockin mice show NF-kappaB trajectories in immune signaling and aging. Cell Rep *41*,
111682. 10.1016/j.celrep.2022.111682.

46. Qiao, W., Huang, Y., Bian, Z., Sun, X., Wang, X., Gao, Q., Peng, Y., and Meng, L.
(2019). Lipopolysaccharide-induced DNA damage response activates nuclear factor
kappaB signalling pathway via GATA4 in dental pulp cells. Int Endod J *52*, 1704-1715.
10.1111/iej.13180.

47. Jia, W., Wu, W., Yang, D., Xiao, C., Huang, M., Long, F., Su, Z., Qin, M., Liu, X., and
Zhu, Y.Z. (2018). GATA4 regulates angiogenesis and persistence of inflammation in
rheumatoid arthritis. Cell Death Dis *9*, 503. 10.1038/s41419-018-0570-5.

48. Marechal, A., and Zou, L. (2013). DNA damage sensing by the ATM and ATR kinases.
Cold Spring Harb Perspect Biol *5*. 10.1101/cshperspect.a012716.

49. Menolfi, D., and Zha, S. (2020). ATM, ATR and DNA-PKcs kinases-the lessons from the
mouse models: inhibition not equal deletion. Cell Biosci *10*, 8. 10.1186/s13578-020-
0376-x.

50. Wang, Y., Zhang, N., Zhang, L., Li, R., Fu, W., Ma, K., Li, X., Wang, L., Wang, J.,
Zhang, H., et al. (2016). Autophagy Regulates Chromatin Ubiquitination in DNA
Damage Response through Elimination of SQSTM1/p62. Mol Cell *63*, 34-48.
10.1016/j.molcel.2016.05.027.

Figure S1

Figure S1. *GATA4 IHC of healthy human cartilage tissue from 3 young (Y1,Y2) and aged (O1,O2) donors. Scale Bar=50 µm.*

Figure S2

Figure S2. *GATA4 IHC of healthy mouse cartilage tissue from young (left) and old (right) mice. Scale Bar=50 µm.*

Figure S3

Figure S3. (A) GATA4 overexpression of young, pooled chondrocytes in monolayer culture 48 hours after infection. The lentiviral control contained the EF1A promoter-driven expression of Green Fluorescent Protein (GFP), and the GATA4 lentivirus contained the EF1A promoter-driven overexpression of GATA4 with dTomato fluorescent protein. Scale Bar=100 μm. (B) Western blot to measure GATA4 protein levels. (C) RT-qPCR analysis of relative gene expression of cartilage matrix proteins aggrecan (ACAN) and collagen type II-α1 (COL2A1), and hypertrophy markers collagen type X- α1 (COL10A1) and India hedgehog (IHH).

859 **Figure S4**

861 ***Figure S4.*** *Safranin-O staining for pellets derived from young, individual chondrocyte (Y1-3)*
862 *transduced with GFP control lentivirus or GATA4 lentivirus. Pellets were cultured in chondrogenic*
863 *medium for 7 days. Scale Bar=50 μm.*

869 ***Figure S5***
870

871
872
873 ***Figure S5.*** *Western blot to examine MMP-13 protein levels in pellets derived from two young*
874 *chondrocyte lines (Y1 and Y2) overexpressing GFP control (GFP C) or GATA4.*

875
876
877
878

879 *Figure S6*

A

B

GATA4

Supplementary Figure 6. Assessment of GATA4 siRNAs in monolayer*. (A) Schematic of the four*
different GATA4 siRNAs with corresponding target sequences assessed for GATA4 knockdown in
monolayer culture of old pooled chondrocytes. (B) RT-qPCR assessing GATA4 levels after siRNA
treatment (n=3).

Figure S7

Supplementary Figure 7. Assessment of GATA4 small molecule inhibitor, NSC140905. *Pooled*
old human chondrocytes were pelleted and treated with the chondrogenic medium with or without
supplanting NSC140905 for 14 days. (A) RT-qPCR analysis of relative gene expression of matrix-
degrading enzymes, including MMP-1,13 and ADAMTS 4 and 5 (n=3). Student's two-tailed t-test
with Welch's correction for standard deviation and a p-value of 0.05 was used for all statistical
analysis. (B) Safranin-O/Fast green staining. Bar=50 μM.

Figure S8

898
899   ***Supplementary Figure S8.*** *(A) H&E staining and (B) p-P65 IHC to assess synovial inflammation*
900   *in mice treated with lentiviral vectors carrying mCherry Control or Gata4. C. p-P65 IHC staining*
901   *was also semi-quantitated.*

907   **Figure S9**

908
909   ***Supplementary Figure S9.*** *Wester blot to examine protein levels in chondrocytes treated with*
910   *doxorubicin (100nM) or vehicle control for 3 days.*

919 **Table S1. Information of chondrocyte donors**
920 RNA sequencing

Age	Gender
24	Female
36	Female
38	Female
70	Female
73	Female
74	Female

921
922 Western Blot

Age	Gender
22	Male
26	Male
29	Male
69	Male
70	Male
73	Female

923
924
925 Human Cartilage Tissue IHC

Age	Gender
24	Male
25	Female
27	Female
33	Male
67	Female
76	Female
78-1	Female
78-2	Male

926
927 Young Chondrocyte Pool

Age	Gender
22	Male
29	Male
38	Female
42	Female

928
929 Old Chondrocyte Pool #1

Age	Gender
66	Male
70	Male
73	Female

930
931 Old Chondrocyte Pool #2

Age	Gender
66-1	Male
66-2	Male
70-1	Male
70-2	Male

933 Monolayer young individual chondrocytes

Age	Gender
29	Male
36	Female
38	Female

935 Pellet young individual chondrocytes

Age	Gender
21	Male
35	Male
38	Female

938 **Table S2. Antibodies used for immunofluorescence (IF), Immunohistochemistry (IHC), or**
939 **Western blot (WB)**

Antibody	Origin	Cat. No	Species	Assay	Dilution
Anti-Human Collagen Type II	MP Biomedicals	SKU 0863171	Mouse	IHC	1:200
Biotinylated Goat Anti-Rabbit IgG	Vector Laboratories	PK-6101	Rabbit	IHC	1:50
Goat Anti-Rabbit IgG H&L (HRP)	Abcam	Ab6721	Goat	WB	1:5000
GAPDH (D16H11) XP® Rabbit mAb	Cell Signaling Technology	5174s	Rabbit	WB	1:2000
Smad1 (D59D7) XP® Rabbit mAb	Cell Signaling Technology	6944s	Rabbit	WB	1:500
Anti-GATA4 antibody (ab84593)- Discontinued	Abcam	ab84593	Rabbit	IHC/WB	1:500
GATA-4 (D3A3M) Rabbit mAb	Cell Signaling Technology	369366s	Rabbit	IHC/WB	1:1000
Phospho-Smad1/5 (Ser463/465) (41D10) Rabbit mAb	Cell Signaling Technology	9516s	Rabbit	WB	1:500
Phospho-Smad2 (Ser465/467) (138D4) Rabbit mAb	Cell Signaling Technology	3108s	Rabbit	WB	1:500
Smad2/3 (D7G7) XP® Rabbit mAb	Cell Signaling Technology	8685s	Rabbit	WB	1:500
Anti-NF-kB p65 (phospho S536) antibody [EP2294Y]	Abcam	ab76302	Rabbit	WB	1:500
Anti-NF-kB p65 antibody [E379]	Abcam	ab32536	Rabbit	WB	1:500
Anti-p21 antibody [EPR362] - BSA and Azide free	Abcam	ab218311	Rabbit	WB	1:1000
Phospho-Histone H2A.X (Ser139) (20E3) Rabbit mAb	Cell Signaling Technology	9718s	Rabbit	WB	1:500

 Table S3: Full names of genes shown in Figure 1

Transcription Regulator Name	Acronym
Hypoxia-inducible factor 1-alpha	HIF1A
GATA binding protein 4	GATA4
MAF bZIP transcription factor B	MAFB
Homeobox D10	HOXD10
CCAAT enhancer binding protein alpha	CEBPA
Forkhead box L2	FOXL2
Caudal type homeobox 2	CDX2
Early Growth Response 1	EGR1
Signal Transducer and Activator of Transcription 3	STAT3
Paired Box 1	PAX1
Sequestosome 1	SQSTM1
Myocardin Related Transcription Factor B	MRTFB
WW Domain Binding Protein 2	WBP2
Scleraxis bHLH Transcription Factor	SCX
Twist Family bHLH Transcription Factor 1	TWIST1
Myocyte Enhancer Factor 2D	MEF2D
Lysine Methyltransferase 2D	KMT2D
Achaete-scute Family bHLH Transcription Factor 1	ASCL1
Recombination Signal Binding Protein for Immunoglobulin kappa J region	RBPJ
SIX homeobox 1	SIX1
FERM domain containing 3	FRMPD3
Ceruloplasmin	CP
SPARC related modular calcium binding 2	SMOC2
ATP binding cassette subfamily A member 13	ABCA13
Inositol 1,4,5-triphosphate receptor type 1	ITPR1
NIM1 serine/threonine protein kinase	NIM1K
Fibroblast growth factor 13	FGF13
Ankyrin repeat domain 12	ANKRD12
GTP binding protein overexpressed in skeletal muscle	GEM
erythroferrone	FAM132B
Pappalysin 1	PAPPA
PPARG coactivator 1 alpha	PPARGC1A
Synaptotagmin like 2	SYTL2
Leucine rich repeat transmembrane neuronal 2	LRRTM2
G protein-coupled receptor class C group 5 member B	GPRC5B
Polypeptide N-acetylgalactosaminyltransferase 15	GALNT15

Growth associated protein 43	GAP43
Fibroblast growth factor binding protein 1	FGFBP1
ATRX chromatin remodeler	ATRX
Protocadherin gamma subfamily B, 2	PCDHGB2
Calsyntenin 2	CLSTN2
Fibroblast growth factor binding protein 2	FGFBP2
Desmocollin 2	DSC2
TNF receptor superfamily member 21	TNFRSF21
Kinesin family member 13B	KIF13B
Plexin A2	PLXNA2
Cysteine rich protein 2	CRIP2
Cysteine rich protein 1	CRIP1
KLF transcription factor 2	KLF2
Serum/glucocorticoid regulated kinase 1	SGK1
LIM zinc finger domain containing 2	LIMS2
Long intergenic non-protein coding RNA 1133	LINC01133
Adipogenesis regulatory factor	ADIRF
Ornithine decarboxylase 1	ODC1
Kazrin, periplakin interacting protein	KAZN
Chondroitin polymerizing factor	CHPF
RAB23, member RAS oncogene family	RAB23
Alkaline phosphatase, biomineralization associated	ALPL
Signal transducer and activator of transcription 4	STAT4
G protein-coupled receptor 1	GPR1
Spectrin alpha, non-erythrocytic 1	SPTAN1
Very low density lipoprotein receptor	VLDLR
Fatty acid desaturase 3	FADS3
Cancer susceptibility candidate 4	CASC4
NAD(P)H quinone dehydrogenase 1	NQO1
Pyrroline-5-carboxylate reductase 1	PYCR1
ERBB receptor feedback inhibitor 1	ERRFI1
NmrA-like family domain containing 1 pseudogene	LOC344887
Desumoylating isopeptidase 2	DESI2
Basonuclin zinc finger protein 1	BNC1

949   **Table S4. Primers for qRT-PCR**

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
RPL13A	GCCATCGTGGCTAAACAGGTA	GTTGGTGTTCATCCGCTTGC
GATA4	CAGTCTACGTGCCCACACC	TCCCGCCTGGCTCCAT
ACAN	AGTCACACCTGAGCAGCATC	AGTTCTCAAATTGCATGGGGTGTC
COL2A1	GGATGGCTGCACGAAACATACCGG	CAAGAAGCAGACCGGCCCTATG
COL10A1	CCCTCTTGTTAGTGCCAACC	AGATTCCAGTCCTTGGGTCA
IHH	AACTCGCTGGCTATCTCGGT	GCCCTCATAATGCAGGGACT
IL-6	ACTCACCTCTTCAGAACGAATTG	CCATCTTTGGAAGGTTCAGGTTG
IL-8	TTTTGCCAAGGAGTGCTAAAGA	AACCCTCTGCACCCAGTTTTC
TNF-α	CCTCTCTCTAATCAGCCCTCTG	GAGGACCTGGGAGTAGATGAG
MMP-1	AAAATTACACGCCAGATTTGCC	GGTGTGACATTACTCCAGAGTTG
MMP-2	GGTCACATCGCTCCAGACT	TACAGGATCATTGGCTACACACC
MMP-3	CGGTTCCGCCTGTCTCAAG	CGCCAAAAGTGCCTGTCTT
MMP-12	GGAATCCTAGCCCATGCTTTT	CATTACGGCCTTTGGATCACT
MMP-13	ACTGAGAGGCTCCGAGAAATG	GAACCCCGCATCTTGGCTT
ADAMTS4	GAGGAGGAGATCGTGTTTCCA	CCAGCTCTAGTAGCAGCGTC
ADAMTS5	GAACATCGACCAACTCTACTCCG	CAATGCCCACCGAACCATCT

953   **Table S5. Information of Luminex assay kit**

Target Molecule	Method	Catalog number & supplier
IL-8	Luminex® assay	HAGP1MAG-12K, EMD Millipore
IL-6, CCL2, MMP-1, MMP-2, MMP-13.	Luminex® assay	LXSAHM-18, R&D Systems

958