Addressing cultural and knowledge barriers to enable preclinical sex inclusive research.

Authors: Brianna N. Gaskill[1*], Benjamin Phillips[2*], Jonathan Ho[3], Holly Rafferty[4], Oladele Olajide Onada[5], Andrew Rooney[6], Amrita Ahluwalia[3*], Natasha A. Karp[2*+]

Affiliations:

1: Novartis Biomedical Research, Cambridge Massachusetts, USA

2: Data Sciences & Quantitative Biology, Discovery Sciences, R&D, AstraZeneca, Cambridge, United Kingdom

3: Faculty of Medicine & Dentistry, Queen Mary University of London, United Kingdom

4: University of Glasgow, United Kingdom

5: Obafemi Awolowo University and Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, Nigeria.

6. Institute of Pharmacy and Biomedical Science, University of Strathclyde, Glasgow, United Kingdom

* These authors contributed equally

+ Author for correspondence: natasha.karp@astrazeneca.com

Abbreviations:

SIRF: Sex Inclusive Research Framework

Key words:

Sex inclusive research

In-vivo

Theory of planned behaviour

25   Behaviour change

26   Preclinical research

27 Abstract

For over thirty years, research has highlighted a sex bias in early research, risking

the validity of biological knowledge. The first step towards change is effectively

challenging misconceptions allowing researchers to perceive sex inclusive research

as do-able. Utilising the theory of planned behaviour, we quantified researchers'

intention as a proxy measure for conducting sex inclusive research and explored

attitude (value of the behaviour), subjective norm (perceived social pressure) and

behavioural control (ability to conduct the behaviour). Additionally, we quantified the

knowledge gap, prevalence of misconceptions, and assessed perceived benefits and

barriers. We tested a workshop intervention that directly challenges the cultural

embedded barriers. The data shows researcher's intentions were high, but they had

weak statistical knowledge and misunderstandings leading to a perception that

inclusive research is prohibitive due to cost and animal use. We demonstrate that

participation in the training intervention improved knowledge, altered the perceived

barriers and cultural expectations.

42 1. Introduction

Research culture encompasses the values, ideals, norms and thus the resultant

behaviours of a community. This resulting culture in turn has substantial impact on

how research is conducted and communicated. There is a long-established

embedded practice of studying only a single (typically male) sex in preclinical

research and then generalising the results to a wider population (*1-3*).

Since the 1990's, it has been recognised that this approach is highly limited (*4*), with

an emerging body of published evidence highlighting that sex may profoundly

influence biological response (*5, 6*). Consequently, there has been growing

awareness that unless sex is factored into experimental design its influence remains

unknown. Importantly, it is now appreciated that single sex studies result in a

knowledge bias (*7, 8*). The need to drive cultural change has been highlighted, with

the goal of enabling equitable research at the exploratory stage (*9*) which delivers a translational body

of knowledge relevant to the broader population (*10*).

Despite numerous funding bodies establishing initiatives to encourage researchers

to routinely integrate males and females into basic, preclinical, and clinical research,

little progress has been observed (*11, 12*). As a response to the status quo,

numerous funding bodies have recently issued inclusion mandates requiring

justification for single sex studies (*9, 11, 13*). The proportion of published papers

including females and males has improved (*14, 15*). A study comparing inclusion

between 2009 and 2019 across nine biological disciplines, found that six out of the

nine disciplines saw improved representation, and overall inclusion improved from

26% to 48% of published studies (*15*). Whilst a positive step forward, this proportion

still represents a minority of studies, and notably this inclusion was not associated

with an increase in the proportion studies that included data analysed by sex.

Additionally, studies identified methodological issues when males and females were

included. Examples include: a failure to report the proportion of the sexes in the

cohort (*16*), inappropriate data analysis such as disaggregation, pooling or

comparison of *p* values to assess for treatment by sex interaction (*14, 17*). There is a

need to not only improve inclusion but also improve the analysis of the subsequent

data to ensure integrity of the conclusions made.

Notionally scientists support efforts to improve sex representation in research (*15, 18,*

19). However, sociological research suggests that many researchers believe

inclusion is not "doable" (*20, 21*). The perceived barriers cited encompass ethical,

financial, statistical, and practical constraints (*18, 22, 23*). An analysis of 30 single

sex justifications identified six themes: concern around a known sex difference or

sex effect (30%), fear of increased experimental variability (27%), experimental

conditions limiting use of female and male animals (13%), limited sample size (13%),

inability to sex subjects (10%), and issues with animal husbandry (7%) (*15*). Many of

the barriers have been highlighted as cultural ingrained misconceptions (*23*). For

example, a commonly proffered argument for exclusion is that female animals are

inherently more variable, and thus inclusion will lead to the need for larger sample

size (N). This is despite a wealth of studies that reject this thesis of increased

variability across a range of biologically relevant endpoints (*24-26*). Alongside

common perception that inclusion will require a doubling of the sample size as

standard (*20, 27, 28*), researchers have also questioned the value of studying

multiple sexes in preclinical research, suggesting that single sex *in vivo* studies can

be generalisable to the wider population (*20*).

The scale of the misconceptions and statistical misunderstandings that result in

barriers to inclusion have resulted in a call for training (*9, 23*). We developed a

workshop directed at equipping participants with knowledge of the current best

practice for sex inclusion. The workshop included material designed to explicitly

address many of the known misconceptions and barriers to sex inclusive research.

To evaluate the efficacy of this intervention we ran two independent experiments

using a survey to collect quantitative and qualitative data regarding self-efficacy

(confidence to conduct inclusive research), knowledge, and views of scientists on the

issues regarding incorporating males and females into preclinical *in vivo* research.

The first experiment was conducted at an international conference with a parallel

group design testing the general population attending the conference, those who

expressed interest in the topic before attending a related seminar and those who

attended the intervention. The second experiment was conducted at a UK Russell

group university with a pre- and post-intervention design. The survey was based on

the theory of planned behaviour; as an established strategy for evaluating and

understanding human behaviour (*29*). The theory of planned behaviour is a

psychological theory that quantifies behavioural intention as a proximal determinant

of behaviour through three core beliefs (attitude, subjective norms and perceived

behavioural control: Box 1). We then assessed the impact of the intervention, with

three key objectives; (1) to capture and quantify the scale of misconceptions around

sex inclusive studies within the research community, (2) to assess whether the

intervention led to an improvement in knowledge, (3) to explore statistically the

impact of the intervention on the intention to conduct sex inclusive research,

alongside identification of other predictors of intention. Taken together, the studies

demonstrate that the intervention is effective and gives insight into how the research

community can expedite change in research practice.

Box 1: Glossary

Technical terms, used within this manuscript, and their associated definitions are
detailed in the table below. For those terms associated to the theory of planned
behaviour, the definitions provided were originally published by Icek Ajzen (*29*). The
name in brackets represents the term used within the statistical model and
associated output.

Word	Definition
Theory of planned behaviour	An empirically supported framework that predicts an individual's intention to perform a particular behaviour by assessing that participant's attitude, subjective norms, and perceived behavioural control over the behaviour.
Intention (Avg_Intent)	The motivational factors that influence a given behaviour. The stronger the intention, the more likely the behaviour will be performed.
Attitude toward the behaviour (Attitude)	The degree to which a person believes a particular behaviour is good or bad.
Subjective norm (Soc_Norm)	This refers to the perceived social pressure experienced by an individual to engage, or not engage, in the behaviour in question.
Perceived behavioural control (Beh _Control)	This includes both actual and perceived control over the behaviour by the participant. Actual control covers resources and opportunities that are available to the participant. Perceived control can encompass an individual's perception about the ease or difficulty in performing the behaviour,

	which may encompass self-efficacy beliefs or confidence in conducting the behaviour.

2. Results

2.1 Demographic results

2.1.1 Study 1

Across the three test groups, the demographics and distribution for the potential predictor variables were found to be balanced (Supplementary Table 1). Furthermore, no significant correlation ($p > 0.05$) was found between the continuous variables as assessed by a Pearson correlation analysis (Supplementary Analysis 1A). Missingness was observed in two demographic variables: two participants (Baseline:1; Interested:1) did not provide information on the years worked in animal research (missingness = 1.9%) and seven responders (Baseline:4; Interested:3) did not provide their age (missingness = 6.7%). The two that did not report years worked also did not report their age.

The majority of the participants were female (61%), had an average age of 39 years, had a PhD (63%), worked at an academic institution (97%) in Europe (58%) and predominantly studied basic biological mechanisms (68%). A large proportion of the participants (71%) had received some type of formal statistical training but 41% were not, or not very, familiar with factorial designs. Whilst 63% of participants are always, or often, involved, or can influence, the planning of experiments, the majority (62%) have only incorporated females and males in 50% or less of their studies. On average, the participants had been involved in animal research for 13.8 years.

2.1.2 Study 2

Analysis of the demographic data (Supplementary Table 2) found that the majority of the participants identified as female (52%), had an average age of 35 years, had a PhD (68%) and predominantly studied basic biological mechanisms (52%). A large proportion of the participants had received some type of formal statistical training (68%) but 55% were not, or not very, familiar with factorial designs. Whilst 81% of participants are always, or often, involved, or can influence, the planning of experimental designs, the majority (51%) have only incorporated females and males in 50% or less of their studies. On average, the participants had been involved with animal research for 10.6 years. No statistically significant correlation ($p > 0.05$) was found between the continuous variables as assessed by a Pearson correlation analysis (Supplementary Analysis 1B).

2.2 Exploration of advantages and barriers

2.2.1 Study 1

Participants had an opportunity to articulate the barriers and benefits to inclusive research by either selecting preset options or entering free text. For most preset options, there was no statistically significant difference seen in the proportion selected between the treatment groups (Fig 1A-B, Supplementary Analysis 2, Supplementary Analysis 3). We did not expect differences between the baseline and interested group but hypothesised that we would observe a shift in the intervention group following the workshop with a reduction in those identifying misconceptions as barriers and an increase in the advantage statements selected. However, power is low to see a change in this exploratory proportion data due to the low sample size of eligible attendees at the workshop. Two exceptions occurred in the benefit question.

Firstly, for the preset option: 'efficient use of all animals from breeding' we saw a

statistically significant increase for the intervention group selecting this as a benefit

of inclusive research (34.2% and 37.5% selecting this option in the baseline and

interested group respectively which then increasing to 80% in the intervention group).

We also observed a statistically significant increase for the intervention group

selecting the translation option (74% baseline, 68% interested, 100% intervention

group).

To represent the general population, the baseline and interested treatment groups

data was pooled (Fig. 1C-D). A higher proportion of participants selected benefits

that are associated with generalisability (translatability and understanding sex

differences) over robustness (reproducibility) and ethical use of animals. A small

proportion of the sample (1.2%) did not perceive any barriers. The predominate

barrier was associated with cost (48.9%). 35% of the participants raised male animal

aggression. Barriers associated with misconceptions (Female variability and sample

size concerns) were raised by 25% of the participants.

Fig. 1: Exploration of the perceived barriers and benefits of inclusion of males and females in in vivo research from study 1.

Survey data collected in study 1 with 39 participants in the baseline group, 51 in the interested group and 15 in the intervention group. A: Barrier comparison between treatment groups. B: Benefit comparison between treatment groups. C: Proportion time barrier selected in the general population (baseline and interested group combined). D: Proportion time benefit selected in the general population (baseline and interested group combined). A Pearson's chi-squared test was used to compare proportions between the treatment groups. Statistical significance is highlighted with a horizontal bar and if the *p*-value is less than 0.05, it is flagged with one star (*). If the *p*-value is less than 0.01, it is flagged with 2 stars (**). If a *p*-value is less than 0.001 it is flagged with 3 stars (***).

2.2.2 Study 2

The study design allowed a comparison of the pre- versus the post-selection of barriers and benefits. For some barriers, statistically significant differences were

seen in the proportion of time they were selected (Supplementary Analysis 4, Fig. 2.).

The workshop was designed to challenge seven of the most frequently cited barriers

by highlighting that they were based on misconceptions. The intervention produced a

statistically significant reduction in five out of the seven barriers, with an average

reduction of 28% citing the barrier post-workshop (Fig 2A). Participants had the

option of selecting alternative barriers and entering free text options. Of these, there

was no statistically significant change in the proportions selected between the data

collected pre- and post-intervention (Fig 2B). For some of the barriers, the

proportions observed in the general population in study 1 (baseline and interested

community) were similar to that seen in study 2 in pre-intervention data (e.g. 'Cost',

'Female variability')(Fig 2E). Whilst other barriers saw a statistically significant

difference between studies, which could represent a difference in the local culture

and practices (Fig 2C). For example, relative to study 1, there was a 24% increase in

the proportion selecting test material availability in study 2, whilst we observed a

25% reduction in those selecting male animal aggression concerns in study 2.

A comparison of the pre- versus post-data (Fig. 2C), found the intervention had little

impact on the proportion of time benefits were selected except for "reproducibility"

where the intervention led to a statistically significant increase (26%). The proportion

of time benefits were selected were equivalent between study 1 and study 2 (Fig.

2D; Supplementary Analysis 5).

Fig. 2: Exploration of the perceived barriers and benefits on the topic of inclusion of females and males in in vivo research for study 2.

Data collected from study 2 with 29 participants completing the barrier question in the pre-survey and 28 in the post-survey. A: Barriers associated with misconceptions that the intervention was looking to address. B: Other barriers. C: Pre- versus post-comparison of the benefits of inclusion. D: Comparison of the benefits selected in study 1 general population (baseline plus interested group) versus study 2 pre-testing group. E: Comparison of the barriers selected in study 1 general population (baseline plus interested groups) versus study 2 pre-testing group. A McNamar's test of association was used to compare the proportions between the pre and post intervention data in study 2. A chi-squared test was used to compare proportions between the two studies. Statistical significance is highlighted with a horizontal bar and if the *p*-value is less than 0.05, it is flagged with one star (*). If the *p*-value is less than 0.01, it is flagged with 2 stars (**). If a *p*-value is less than 0.001 it is flagged with 3 stars (***).

2.3 Exploration of intention

2.3.1 Study 1

The average intention score, across all treatment groups for study 1 was high (5.67 ± 1.19, possible range 1:7). A statistical analysis assessed which variables could

explain variation in intent to conduct sex inclusive research (Table 2, Fig 3). Two of the three theory of planned behaviour attributes, subjective norm and attitude, positively correlated with intent (Fig 3A-C). The data indicates that as the participant's attitude increases (i.e. feel incorporating males and females is important) their intention to incorporate females and males increases; with a high mean attitude of 6.5 ± 0.87. The subjective norm attribute had a similar positive correlation with intent, indicating that when participants perceived higher societal pressure, they had higher intention. Subjective norm differed from attitude in that at the population level the score was in the mid-range (4.7 ± 1.4). The intent for those who attended the workshop was significantly higher, with approximately 20% increase in the intention score (Fig 3D). However, no significant difference in intent was found between the baseline and interested treatment groups. The only other variable to have some evidence of predictive ability was how often a participant was involved in or could influence the planning of experiments positively correlated with intent to incorporate females and males (Table 2, Fig 3E). Thus, suggesting individuals more involved in planning experiments had a higher intention to run inclusive designs.

Source	Nparm	Df	Sum of Squares	F Ratio	Prob > F	Significance	Eta^2
Attitude	1	1	18.5650	34.9548	**<.0001**	***	0.15
Beh_Control	1	1	0.0814	0.1534	0.6964		0.00
Soc_Norm	1	1	12.2229	23.0137	**<.0001**	***	0.10
Treatment_Group	2	2	3.8643	3.6380	**0.0311**	*	0.03
Age	1	1	0.0995	0.1874	0.6664		0.00
Gender	2	2	2.6196	2.4662	0.0918		0.02

Geography	3	3	0.1955	0.1227	0.9464		0.00
Year_Work	1	1	0.1828	0.3442	0.5592		0.00
Type_Work	3	3	1.1958	0.7505	0.5255		0.01
Education	2	2	2.9339	2.7621	0.0696		0.02
Stats_Training	2	2	0.3810	0.3587	0.6998		0.00
Factorial_Fam	1	1	0.0332	0.0625	0.8033		0.00
Factorial_Incor	1	1	0.2614	0.4921	0.4852		0.00
Ability_Influence	1	1	2.3253	4.3782	**0.0398**	*	0.02

Table 2: Statistical model output for the full model for study 1 data exploring the predictors ability to explain variation in average intent.

Where Beh_control represent the average behavioural control score, Soc_norm the average social normal score, Year_Work represented the number of years the participants have worked in animal research, Type_Work represents the type of research conducted by the participant, Education the highest level of education obtained, Stats_Training represents the level of statistical training received, Factorial_Fam represents how familiar the participants were with factorial experimental design, Factorial_Incor represents how often the participants incorporated males and females into their experiments while studying an intervention, attitude represents the average attitude score, and Ability_Influence represents how often the participants were involved or could influence the planning of experiments involving animals. Nparm stands for the number of parameters, Df represents the degrees of freedom and Prob > F represent the p value associated with the F ratio. Statistical significance shown as * for p-value < 0.05, ** for p< 0.01 and *** for p<0.001.

Fig. 3: Exploration of intent for significant and critical predictor variables for study 1 data

A full model, with all potential predictors and demographics was fitted to explore the variation in intent and assess for evidence of predictive behaviour. The baseline group was set as the reference group. If main effects were significant the variation was explored with Tukey post hoc testing. For graphs A, B and C the grey area indicates the 95% confidence interval for the fitted linear relationship (blue). D: Data are presented as the model estimated mean (Least Square Means) with a standard error bar estimated from the model. Bar represents the planned comparison with * representing statistical significance <0.05. E: Violin plot showing the distribution of intent as a function of the ability to influence the design. The red box indicates the mean.

2.3.2 Study 2

In an exploration of intention for study 2, only subjective norm positively correlated with intent to conduct a sex inclusive study (Table 3, Fig 4A-C). Attitude did not predict intention but was found to have a very high average score (6.7 ± 0.51). Like study 1, behavioural control was not predictive but had a mid-range average score (4.3 ± 1.14). The workshop intervention was not found to explain variation in the intention score (Table 3, Fig 4D). However, the intention score in the pre-test group (5.71 ± 1.08) was only slightly higher than the intention score in the general population in study 1 (baseline and general interest; 5.51 ± 1.2). Age was also found

289 to be negatively correlated indicating older staff were less engaged with sex inclusive
290 research in this dataset (Table 3, Fig 4E).

Source	Nparm	Df	Df Denominator	F Ratio	Prob > F	Significance	Eta^2
Attitude	1	1	38.78	0.0268	0.8708		6.91e-04
Beh_Control	1	1	42.47	1.9080	0.1744		0.04
Soc_Norm	1	1	29.15	19.658	**0.0001**	***	0.40
Intervention	1	1	40.06	0.0758	0.7845		1.89e-03
Age	1	1	28.2	5.3428	**0.0283**	*	0.16
Gender	2	2	33.93	0.2973	0.7447		0.02
Ability_Influence	1	1	27.91	2.2618	0.1438		0.07
Factorial_Fam	1	1	42.98	0.3822	0.5397		8.81e-03
Factorial_Incor	1	1	35.72	0.0725	0.7893		2.03e-03

Table 3: Statistical model output for the full model for study 2 data exploring the predictors
ability to explain variation in average intent.

Where Beh_control represent the average behavioural control score, Soc_norm the average social normal score,
Year_Work represented the number of years the participants have worked in animal research, Type_Work
represents the type of research conducted by the participant, Education the highest level of education obtained,
Stats_Training represents the level of statistical training received, Factorial_Fam represents how familiar the
participants were with factorial experimental design, Factorial_Incor represents how often the participants
incorporated males and females into their experiments while studying an intervention, attitude represents the
average attitude score, and Ability_Influence represents how often the participants were involved or could
influence the planning of experiments involving animals. Nparm stands for the number of parameters, Df
represents the degrees of freedom and Prob > F represent the p value associated with the F ratio. Statistical
significance is flagged with one star (*) if the *p*-value is less than 0.05, with 2 stars (**) if less than 0.01, and with
3 stars (***) if less than 0.001.

$F_{1,40.06} = 0.076; p=0.78$

Fig. 4: Exploration of intent for significant and critical predictor variables for

study 2 data

A full model, with all potential predictors and demographics was fitted to explore the variation in intent and assess

for evidence of predictive behaviour. For graphs A, B, C and E: the grey area indicates the 95% confidence

interval for the fitted linear relationship (blue). Graph D displays density and individual participant pre/post Box-

Cox transformed intent values.

2.4 Exploration of knowledge

2.4.1 Study 1

To complement the evaluation of intention and barriers and advantages, five

questions were included to assess the knowledge of the participants on common

misconceptions and errors around planning and data analysis of sex inclusive

studies (Supplementary Analysis 6, Fig 5). The cumulative knowledge score did not

significantly differ between the baseline and interested treatment groups ($z = - 0.513$;

$p = 0.601$) (Fig 5A). Whilst the workshop intervention group had a significantly higher

321 score relative to the baseline ($z = 3.515$; $p = 0.0044$) with an average increase of

322 1.94 questions correctly answered. This could also be seen in a comparison of

323 proportion of correct answers for each individual question with a statistically

324 significant higher score in the intervention group for 3 out of the 5 questions (Fig. 5B).

2.4.2 Study 2

The knowledge questions allow exploration in dataset 2 on the common

misconceptions around sex inclusive designs and common errors in data analysis of

sex inclusive studies (Supplementary Analysis 7, Fig 5). The workshop intervention

was found to significantly improve the score ($t = 4.328$, $df = 25$, $p = 0.0002$) with an

average increase of 1.769 questions answered correctly (Fig 5C), a similar effect

size to study 1. The impact of the intervention could also be seen at an individual

question level with a statistically significant higher score in the intervention group for

four out of the five questions (Fig. 5D).

Fig. 5: Exploration of intervention impact on the proportion of correctly answered knowledge questions.

A: Study 1: Impact of treatment group on the cumulative knowledge score. Statistical significance assessed with a Poisson regression (Baseline: N=39, interest: N=51 and intervention: N=15). B: Study 1: Proportion of correct answers for each question (Baseline: N=39, interest: N=51 and intervention: N=15). Statistical significance assessed with a Pearson's chi squared test. C: Study 2: Impact of intervention on the cumulative knowledge score. Statistical significance assessed with a paired *t*-test (N=26 with pre and post data available). D: Study 2: Proportion of correct answers for each question (Pre: N=29, Post: N=28). Statistical significance assessed with McNamar's test of association. Statistical significance shown as * for *p*-value < 0.05, ** for p< 0.01 and *** for p<0.001.

3. Discussion

General population conclusions – barriers/benefits/knowledge

The survey provided an opportunity to understand researchers position around sex inclusive research. Across the two studies, the benefits of sex inclusive research

were selected at a similar rate with a higher proportion of participants valuing the

benefits associated with generalisability of the findings (ability to understand sex

differences and translation) over robust ethical research. This recognition of the

benefit of inclusive research aligns with previous suggestions that scientists are

generally supportive of efforts to improve sex representation in biomedical research

(*23, 28*). However, the focus on the ability to understand sex differences aligns with

a common misconception that the goal of inclusion is to identify sex differences. This

misconception causes a dissonance with the recommendation to share the sample

size (N) for a treatment effect between the two sexes as this approach does not

power experiments to identify sex differences. Being explicit on the experimental

design goals and the impact on the design was raised by Rich-Edwards (*9*) who

developed a "4 Cs" decision framework to guide researchers in sex inclusive

research by focusing on four steps: Consider, Collect, Characterise and

Communicate. Within this framework, they identify two pathways depending on

whether the study is an exploratory study (where males and females are included to

improve generalizability) versus confirmatory study (those actively exploring sex-

related differences) and guide researchers in the design and analysis implication of

this. There is a culture shift needed that the primary goal in exploratory sex-inclusion

is to deliver generalisable results where large differences in sex-related variation will

be identified. It is this understanding of the difference between exploratory versus

confirmatory sex inclusive research that in our opinion is deficient.

Research has found that scientists do not believe that inclusion is 'do-able' (*21*) due

to a range of perceived barriers to inclusion (*13, 22, 23, 25, 30*). This study provides

the first quantification of the scale of these perceived barriers, of which many are

misconceptions, and finds them endemic. Seldom did the sampled community

express the absence of barriers. Cost was the most cited barrier, approaching half of

the community raising this as a concern. This was followed by on average a third of

the community selecting barriers related to misconceptions such as experimental

design complexity, sample size concerns and female variability.

For some of the barriers, we observed a statistically significant difference between

studies (test material availability and male animal aggression), which could represent

a difference in the local culture or research interests impacting the perceived barriers.

For example: In study 1, male mouse aggression was raised by 30% of the

respondents but only 10% in study 2. This finding is at odds with the observation that

male animal research predominates (*15*) but could relate to welfare management

practices prioritised within that community. Sharing best practices and working on

housing and husbandry conditions could minimise the impact of this concern (*31-33*).

The responses to the knowledge questions also give an opportunity to directly

assess whether participants are holding a misconception. By focusing on the

baseline and interested group in study 1 or the pre-intervention data in study 2, we

get an estimate of the general research population. This finds that misconceptions

dominate the research community beliefs. For example, the misconception that

inclusion of males and females would double the N (80% of study 1 N=90, 79% of

study 2 N=28) held this view. This is probably arising from the perception that sex

inclusive designs increased variability and therefore require a larger overall sample

size (90% of study 1 N=90, 93% of study 2 N=28). This perception has been

acknowledged in previously published articles (*13, 20, 21, 25, 26*), however, this is

the first known quantitation of the extent of these viewpoints.

Regarding the data analysis, most researchers (75.6% in study 1 general population

N=90, and 72% of the participants in pre-assessment of study 2 N=28) correctly

knew that sex should be included in the statistical model. Questions that are more

specific about analysis strategies finds the proportion of incorrect answers to be very

high. For example, most thought the data could be pooled across the sexes (81.1%

in study 1 general population, 83% study 2 pre-assessment) and that the data should

be disaggregated (80% in study 1 general population, 72% study 2 pre-assessment).

The prevalence of incorrect ideas around data analysis aligns with a published meta-

analysis that analysis mistakes are common when females and males are included

in addition to the intervention of interest in research (*17*). The high proportion of

correct answers for the initial question to include sex but then high proportion of

agreeing with poor analytical strategies suggests that participants have a sense they

need to do something but either don't know what the right analysis strategy is or are

unaware of the problems with the suboptimal analysis strategies.

It is of note that no one selected data analysis concerns in study 1 and infrequently

in study 2 (7% in the pre-test assessment) as a barrier to inclusion which suggests

researchers are unaware of this shortcoming. This aligns with research identifying

that statistical errors in the analysis of sex inclusive datasets are common (*17*). The

difficulty in recognizing one's own incompetence is a known cognitive bias, described

as the Dunning-Kruger effect (*34*). Research has shown that training not only

improves participants skills but also helps the individuals recognize the limitations of

their ability (*34*). This highlights the importance of investing in training researchers in

experimental design and data analysis and that this is a critical step to enable sex

inclusive research.

These findings highlight the scale of the misconceptions hindering engagement with

sex inclusive research and the need for research practice leaders to focus both on

these general misconceptions but also to determine local barriers that need

exploration with their research communities.

Exploration of intention to inclusive research

While training is crucially important, it alone is unlikely to completely solve the

imbedded cultural problem we currently face. Simply knowing that one should

exercise to improve your health, does not mean one will get up every morning to go

to the gym. Thus, knowing exactly what drivers are needed to influence the required

behavioural change are equally as important as making sure people receive training

to enact the change. The theory of planned behaviour provides us with key insights

on how to influence intention and ultimately behavioural change. This is key as the

intention to conduct sex inclusive research in the future was generally positive (5.51

± 1.25 in study 1 general population, 5.71 ± 1.08 study 2 pre-assessment out of a

total score of 7) and attitude was very high (6.41 ± 0.92 in study 1 general population,

6.66 ± 0.67 study 2 pre-assessment out of a score of 7). This indicates that generally,

scientists view sex inclusive research as beneficial, useful, and the right thing to do.

Despite this, the researchers reported only conducting sex inclusive research in ≤

55% of their studies over the past 5 years (55% in study 1 general population and

35% study 2 pre-assessment). Thus, confirming the above posit that simply knowing

sex inclusive research is important is not enough to influence active incorporation,

and since attitude is already high, it leaves little room for further influence on

behavioural intent.

Although attitude was high, subjective norm differed at the population level where the

score was more in the mid-range (4.6 ± 1.4 in study 1 general population, 5.14 ±

1.11 study 2 pre-assessment). This factor was also the only one of the three theory

of planned behaviour's beliefs found to significantly influence intent in both studies

with large effect sizes. Understanding the strong positive correlation between

subjective norm and intent allows for focused interventions to increase societal

pressure on the scientific community which should then improve overall intention

towards sex inclusive research.

While researchers may have the right motivation, performing the behaviour is

dependent on other non-motivational factors such as the time, money, or skills (*29*).

A behaviour can only be conducted if the researcher is the decision maker. Not only

does actual control influence behaviour but so does the perception of behavioral

control, such as self-efficacy. Self-efficacy encompasses both how well a person can

do the behaviour but also how confident they feel about doing it. Self-efficacy

opinions can influence choice of activities, preparation, as well as effort. In this case,

we believe that the lack of significance in our model is due to the false perception of

how to accurately design and analyse factorial designs, evidenced by the low

knowledge score. According to Ajzen, perceived behavioural control may not be

realistic if a participant has little knowledge about the behaviour in question (*29*).

Again, based on the low knowledge scores of how to design and analyse sex

inclusive data may indicate that this measure may not add much accuracy to

behavioural prediction. However, it does indicate the overwhelming need to create a

training that is accessible and practical to all in vivo scientists.

Impact of the intervention on intention

We had an opportunity to incorporate a workshop at an international conference and as part of this we decided to assess how effective that training was. As many have experienced, the practical aspects and decisions that need to be made in an *in vivo* experiment are rarely covered in traditional statistics courses. This often leaves scientists unsure how to apply what they have learned in the real world. Further, 26-29% of the participants in this study had received no formal statistical training, and yet 71-75% can always or often influence the experimental design. Illustrating the need for continuing education and practical instruction for established scientists. In the first study, trained attendees had the highest level of intent to conduct sex inclusive research in the future. Potentially indicating that our workshop was effective. However, it is also plausible that those who signed up inherently had a higher level of intent. Thus, we felt it was important to evaluate attendee's intent before and after the training. So, this naturally led to our second study where we found that intent was not altered by the workshop, because it was already high. Even though intent was not statistically altered (Pre: 5.71 ± 1.08; Post: 6.00 ± 1.03), participant's knowledge increased and indicated fewer barriers after the training. Thus, illustrating that education alone may not be enough to change behaviour but utilizing the insight gained that social pressure may be the best avenue to influence actual behavioural change.

Impact of the intervention on perceived benefits and barriers

The primary goal of the intervention was to challenge misconceptions; however, the workshop material naturally started the learning journey by exploring the drivers for,

and barriers preventing, change. When focused on the benefits, we found that

following the training intervention participants tended to select each of the benefits at

a higher proportion compared to other groups. There was a statistically significant

change in a few of the benefits: a large increase in the selection of efficient use of

animals and translatability in study 1 and reproducibility in study 2. The infrequent

statistical significance of the change at the individual benefit level is likely to arise

from low statistical power due to the high baseline signal and low sample size.

An exploration of the study 2 data found that the intervention did not generally

reduce the selection of barriers but did decrease selection of the barriers associated

with the misconceptions. This demonstrates that the targeted intervention was

effective, with a meaningful effect size (a 28% reduction on average), at improving

the perception that generalisable sex inclusive research is feasible.

Impact of intervention on knowledge

It has been highlighted previously that early career researchers lack awareness of

best practice recommendations across their areas of research (*35*). To address this,

it is important that training initiatives directed at improving knowledge of best practice

are put in place by institutions, with the goal of continuously improving practice, and

supporting cultural change. This means that training must extend beyond that

provided simply as part of graduate and post-graduate studies, and that

implementing training opportunities as part of continuing professional development of

research staff.

In the baseline and pre-intervention data, performance on the panel of knowledge

questions was generally poor, with only the question around whether sex should be

included in analysis answered correctly by more than half the participants. This is

consistent with previous findings that statistical approaches, even studies designed

as sex inclusive, are frequently erroneous (*17*). Encouragingly, post-intervention

participants demonstrated significant improvements in knowledge. For example, far

more post-intervention researchers correctly identified that disaggregation and/or

pooling of the sexes within a statistical analysis was a mistake, particularly in study 2.

Given that the majority of commonly cited barriers stem from misconceptions which

are arising from knowledge gaps, the evidence that targeted interventions can

directly enhance knowledge offers promise for promoting cultural change through

educational initiatives. Despite across-the-board improvements in performance, the

outcomes still leave room for further improvement. For example, nearly 40% of post-

intervention participants still expressed the view that inclusion of both sexes required

a doubling of the sample size. This highlights that a single session is likely

insufficient to equip researchers with all relevant knowledge and would thus support

a future strategy of continuous improvement through multiple training and learning

strategies.

Limitations of the study

Whilst the data indicate clear benefit of the workshop intervention, there are several

caveats and limitations that should be considered and acknowledged. For both

studies, university researchers made up the majority of the population. The

workshop was hosted by a recognised senior leader (Professor Ahluwalia) both at

the World Congress (role: WCP2023 Secretary-General) and within the university

setting (role: Dean for research at the Faculty of Medicine and Dentistry). It is

possible that participant awareness of the views of the senior leader may have

influenced the positive outcome of the intervention. For study 1, the congress was

specifically focussed for pharmacologists, and it is possible that the outcomes reflect

only this sector of biomedical research. Although for study 2, the workshop was

advertised across a multi-faculty university and likely represented a more specialty-

diverse population. Whether the workshop would effect a similar change in

understanding from those that work within the pharmaceutical industry requires

further evaluation. The impact of the workshop could depend on the skills and

experience of the trainers. Furthermore, the results could be specific to a face-to-

face training event and whether a virtual course (which has the benefit of scale)

would have similar benefit would need further exploration. This workshop focused on

the barriers and misconceptions around the design and analysis. The material did

not tackle any of the practical barriers e.g. management of welfare issue with male

mice. There will be a need for future work to explore how to support and address

these barriers. Finally, the questions posed to workshop attendees did not state

whether the queries were being posed for a first exploratory study or not. It is

possible that had this been made explicit then the responses could have differed.

556   4. Conclusions

An endemic and persistent sex bias in early research has been raised as a risk to the validity

of biological knowledge. Sociological research finds that scientists believe sex matters, but

change has been limited. Funders are actively trying to change behaviour by requiring a

justification for exclusion. Changing the status quo requires individuals to buy into a new

direction that aligns with their values and beliefs. Here we demonstrate that a workshop

intervention can rescue the core beliefs that hinder sex inclusive research. We also provide

evidence that institutes, funders, professional bodies can help this journey by raising

awareness of best practice as this will change the perceived cultural norms. All these

resources used in this research both the teaching material and the surveys have been made

freely available as supplementary material to support further research and activities.

5. Materials and Methods

All procedures and informed consent protocols were approved by the Queen Mary

University of London Ethics Research Committee (ID: QMERC23.001). The

protocols obtained informed consent after the nature and possible consequences of

the study were explained. In November 2023, an amendment was approved to run

additional workshops for staff and students at Queen Mary University of London (ID:

QMERC23.002).

5.1 Study 1 Experimental Design

In this study, data from three treatment groups (baseline, interested, intervention)

were collected at the 19th World Congress of Basic & Clinical Pharmacology 2023 in

Glasgow, Scotland with an on-line questionnaire (Supplementary material 1) by

convenience sampling within each treatment group. Participants attending the poster

sessions on 3rd and 4th July were recruited by circulating staff to contribute to the

baseline treatment group, thus sampling the general population that attended the

conference. Participants for the interested group were recruited by approaching

attendees who were in attendance at a related symposium advertised in the

conference proceedings on the 4th of July 2023 (entitled: "The importance of

interrogating sex differences in cardiovascular physiology and disease"). Participants

for the intervention group were recruited by advertising a free educational workshop

in the conference proceedings entitled "Best practice for sex inclusive research"

conducted on the 5th of July 2023 and data were collected post intervention.

Workshop participants were given the option to participate in the survey at the end of

the workshop. To compensate participants, in any of the treatment groups, for their

time, they received an entry into a drawing for one of ten £50 Amazon e-gift cards.

Participants were included if they conducted research on a disease or biological

phenomenon that affects male and females and were able to influence the plan or

conduct of *in vivo* research. At the point of data collection, the organisers were not

masked (blinded) as we knew the hypothesis and the intervention group in the study.

However, data was collected by sharing a QR code with the participants who then completed

the data entry independently and therefore our ability to influence the results were minimal.

A sample size estimate was conducted a priori using the highest standard deviation

(1.4) observed in a similar study (*36*) that also used a survey based on the theory of

planned behaviour. With this standard deviation, to detect a change of one point

between the groups a power calculation based on the student's *t*-Test estimated 31

participants per group (62 total) would be needed to achieve a power of 0.8. We

aimed to collect data from 50 participants in each treatment group, to ensure we had

sufficient data after dropout (failure to attend or meet the inclusion criteria).

605 5.2 Study 2 Experimental Design

In this study, a pre- and post- intervention design was implemented to assess the

impact of the workshop with an online questionnaire (Supplementary material 2 (pre-

questions) and 3 (post-questions)) with convenience sampling of a pre-existing

group. Participants from Queen Mary University of London were recruited by

advertising an educational workshop entitled "Best Practice for sex inclusive

research" conducted on the 14th of February 2024. To compensate participants for

their time, they received an entry into a drawing for a single £50 Amazon e-gift card.

Participants were included if they conducted research on a disease or biological

phenomenon that affects female and males and were able to influence the plan or

conduct of *in vivo* research.

We aimed to recruit 50 participants to the workshop and only collected data during

the workshop. This was the maximum number we felt would be feasible to maintain

the interactive nature of the workshop. The implementation of a pre- and post-design

should have higher power than the previous study (see power calculation section

2.1) to detect changes. At the point of data collection, the organisers were not

masked (blinded) as we knew the hypothesis and the intervention point in the study.

However, data was collected by sharing a QR code with the participants who then

completed the data entry independently and therefore our ability to influence the

results were minimal.

625 5.3 Survey construct

The survey was developed through consulting with experts in survey methodology,

behaviour theory, and sex inclusive research design and analysis. When possible,

validated instruments were used (i.e., questions based on the TBP). When validated

instrumentation did not exist, new items were created, reviewed by statistical experts,

piloted, and revised as necessary. Two cycles of pilots were conducted with *in vivo*

pharmacology researchers (N=9) from both academic and pharmaceutical settings.

Survey 1 consisted of 39 questions and survey 2 a maximum of 34 questions in a

testing phase (Supplementary Material 1). Demographic questions were omitted

from the post-question format in Study 2 and data from the same individual were

aligned between the pre- and post-questions through the 3+ initials inputted by the

researcher (pre- survey: Supplementary Material 2 and post-survey: Supplementary

Material 3).

The initial four questions obtained consent and assessed whether participants met

the inclusion criteria. If the participant did not grant consent or failed to meet the

inclusion criteria on any question the survey was terminated. Thereafter, all

questions were presented to each participant. Participants were asked demographic

questions which included age, gender, geographic region (study 1 only), and highest

level of education. Several questions captured information about their work including

type of institution (e.g., academic, contract research organization; study 1 only),

primary type of research (e.g., applied, basic, regulatory), and number of years

working with laboratory animals. In addition, questions were included to capture data

on potential explanatory variables. These included their ability to influence the

construct of experiments, the amount of statistical education they had received,

familiarity with factorial designs, how often they incorporate males and females into

their experimental designs, their knowledge on how to analyse data when females

and males were collected, and the impact of including males and females on

statistical power. Finally, a list of pre-identified advantages and barriers were

provided to participants to list perceived reasons why incorporating females and

males was considered advantageous or challenging. Participants were provided with

an empty text box to list any other advantages or barriers in addition to the list

provided.

Motivations, outlined by the theory of planned behaviour, regarding sex inclusive

research were assessed using a series of thirteen questions. Participants answered

four close-ended quantitative questions about their behavioural attitudes (relative

perception about the positive or negative valence of sex inclusive research), three

questions on subjective norms (social and professional pressures to conduct sex

inclusive research) and three questions on perceived behavioural control (general

confidence/control over the ability to run sex inclusive studies). The perceived

behavioural control variable is different from self-efficacy as it asks participants about

external control factors, such as whether inclusive designs are under the control of

the participant. One open ended question was included at the end of survey to

assess if there were any other thoughts or comments participants wanted to share

on the topic. Finally, three questions were used to assess participants' future intent

to implement sex inclusive research.

The survey includes a question to determine whether participants are answering the

question as part of the baseline/interested/intervention group (study 1) or pre/post

assessment (study 2). Multiple participation was not anticipated due to the length of

the survey and the time constraints for data collection.

The surveys were constructed to ensure that the only identifiable information was the

optional input of an email address to support the draw of the Amazon vouchers. To

maintain anonymity, once data was downloaded, an ID code was assigned, email

addresses removed, and the email information only used for the draw. The final data

was stored in an encrypted, password-protected file accessible only to the research team.

5.4 Workshop construct

The workshop, or training intervention, was developed by statisticians who have

published, presented internationally, and taught on this topic (Supplementary

Material 4). One of the statisticians has a formal teaching qualification and a

background in education. The training consisted of a 1 hour 45 min workshop with

didactic, interactive activities and introduced a framework to evaluate research

proposals from a sex inclusive perspective (Table 1, Supplementary material 4).

Prior to study 1, it was run as a pilot with a small group of scientists to test the

material. Following the results of study 1, the workshop material was refined by

amending the multiple-choice question one and two to be more explicit in driving a

discussion around whether data should be analysed by pooling, disaggregation or

with a factorial test. Furthermore, multiple-choice question six was added to re-

enforce learning that a baseline sex difference can be separated from an intervention

effect.

Section	Content
Lecture	Terminology Reflection that clinical sex matters. Exploration of the status of sex inclusion in preclinical research. Exploration of what is sex inclusive research. The role of factorial analysis in sex inclusive research.
Interactive Element	Multiple choice questions with voting and discussion on appropriate analysis strategy and understanding factorial output.
Lecture	Exploration of the perceived barriers to sex inclusive research.
Evaluation tool	Introduction of the Sex Inclusive Research Framework (SIRF) which can be used to evaluate research proposals from a sex inclusive perspective.
Interactive element	Multiple choice classification using the SIRF to classify justifications given for one sex designs.

696 **Table 1:** Workshop Intervention construct

5.5 Statistics and Reproducibility

5.5.1 Participant inclusion

In study 1, a total of 194 participants started the survey. Of these, 105 met the

inclusion criteria questions (N=39 baseline, N=51 interested and N=15 intervention).

While all 105 met the inclusion criteria, seven participants left the question about age

blank. Therefore, the final dataset to evaluate intent included 98 participants (N=35

baseline, N=48 interested and N=15 intervention).

In study 2, a total of 63 individuals registered for the workshop, 42 participants

attended all, or part of the workshop, and survey data was collected from 31

individuals. However, due to some not meeting the inclusion criteria, or not

completing one of the surveys, we received data from 28 individuals for the pre-

survey and 28 for the post.

5.5.2 Variable coding

For the theory of planned behaviour questions, each question had a scale from 1 to

7. An average score was calculated for each motivation (attitude, perceived

behavioural control and social norm) and for intent. This strategy required that each

participant answer at least 50% of the questions for each motivation, otherwise their

data was discarded. All participants answered at least 50% of the theory of planned

behaviour questions and therefore no participant's data were discarded based on

these criteria.

To aid analysis, demographic categories with less than ten responses were
combined with similar themed options into larger categories. For example, the
question that asked about previous statistical training, the categories of "No training"
and "Primarily informal/practical" were collapsed into a category called "No courses".
Missing data for categorical variables (gender, geography, etc) were coded as
"unknown".

Within the survey, two (survey 2) to three (survey 1) questions had opportunities for
participants to add a free text answer in response to a question as an additional
option provided to a question. Those questions were about the perceived barriers to
or advantages of using males and females in preclinical research. For these
questions, all free text submitted were reviewed by the research team and
categorized based on perceived theme. Generally, very few free responses (Survey
1: 12 for barrier question only; Survey 2:6 for barrier question only) were provided.

Five knowledge questions were posed to the participants based on misconceptions
(22) concerning using females and males simultaneously in an experiment and
variability in the female sex. With each question having a single correct answer. If a
participant provided the correct answer, they were given a point, and the points were
summed across the five questions. The knowledge question summary metric was
not included in the statistical exploration of intent, as knowledge was anticipated to
correlate with treatment group.

5.5.3 Quantitative analysis

739 Throughout this research, a significance level of $p < 0.05$ was considered statistically
740 significant. Multiple testing correction was not applied as this is an exploratory
741 analysis.

5.5.3.1 Evaluation of intent

All data analyses were conducted in JMP 14.0.0 (SAS Institute Inc., Cary, NC USA). The data for each analysis are provided in Supplementary Analysis 8, including the specific SAS code used to run the statistical model regarding the theory of planned behaviour intent analyses.

The dependent variable for quantitative analysis was the average intent (Avg_Intent) which was Box-Cox transformed to improve the distributional characteristics in study 1 only. As fixed effects of interest, the statistical model included 'Treatment_Group' for study 1 and 'Intervention' for study 2. In addition, three theory of planned behaviour attributes: average attitude (Attitude), average behavioural control score (Beh_Control) and average social norm (Soc_Norm) were included as fixed effects. Furthermore, potential explanatory variables (e.g., demographic variables) were included in the model as fixed effects. A full model containing all possible terms (Eq. [1] study 1 and Eq. [2] study 2) was used as we had no strong prior information that we could use to select predictive variables. For the data from study 2, a mixed model analysis was used as we included a random term for participant to account for the repeat nature of the data. For both studies, the baseline group was set as the reference group. Since a study of this kind has not been conducted before and the social factors (age, gender, previous statistical experience, etc.) are not reported as explaining/not explaining variability in data such as this, we left all potential explanatory variables in the model to provide data on demographic variables for future social science work of this kind. We evaluated the final model with and without these non-significant variables and the inclusion, or exclusion, did not affect the results presented. Model diagnostics (e.g., independence of residuals, homogeneity

of variance, and normality of residuals) were inspected and no concerns on the

model quality were identified.

Eta squared for study 2 was estimated using the F_to_eta2 function in the effect size

library with R4.3.1.

Eq. 1: Avg_Intent = Treatment_Group + Attitude + Beh_Control + Soc_Norm + Age +

Gender + Geography + Year_Work + Education + Type_Work + Stats_Training +

Factorial_Fam + Factorial_Incor + Ability_Influence

Eq.2: Avg_Intent = Intervention + Attitude + Beh_Control + Soc_Norm + Age +

Gender + Ability_Influence + Factorial_Fam + Factorial_Incor + (1|Participant)

Where Year_Work represented the number of year the participants have worked in

animal research, Education the highest level of education obtained ('Doctoral',

'Masters', 'Other'), Type_Work represents the type of research, Training represents

the level of statistical training received ('No courses', '1-2 courses', (>2 courses'),

Factorial_Fam represents how familiar the participants were with factorial

experimental design, Factorial_Incor represents how often the participants

incorporated males and females into their experiments while studying an intervention

and Ability_Influence represents how often the participants were involved or could

influence the planning of experiments involving animals. Participant was included as

a random effect in Eq. 2 and is represented in the model as (1|Participant). For study

2, few demographics could be included in the analysis due to lack of variability and

the fact that several were not asked at the post-intervention time point.

5.5.3.2 Evaluation of knowledge

Using the statistical programming language, R Version 4.3.1, a Poisson regression analysis assessed the role of treatment group in explaining variation in the cumulative knowledge score for study 1 (data: Supplementary Table 3, analysis: Supplementary Analysis 9) and study 2 (data: Supplementary Table 4). A variety of diagnostics (e.g., residual distribution, assessment Cook's distance, Leverage, goodness fit test) were generated to ensure the model was robust and appropriate for the data.

6. References and Notes

1. A. K. Beery, I. Zucker, Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev* **35**, 565-572 (2011).
2. K. J. Berkley, Vive la difference. *Trends Neurosci* **15**, 331-332 (1992).
3. N. C. Woitowich, A. Beery, T. Woodruff, A 10-year follow-up study of sex inclusion in the biological sciences. *Elife* **9**, e56344 (2020).
4. E. A. Nunamaker, P. V. Turner, Unmasking the Adverse Impacts of Sex Bias on Science and Research Animal Welfare. *Animals* **13**, 2792 (2023).
5. N. A. Karp, J. Mason, A. L. Beaudet, Y. Benjamini, L. Bower, R. E. Braun, S. D. M. Brown, E. J. Chesler, M. E. Dickinson, A. M. Flenniken, H. Fuchs, M. H. Angelis, X. Gao, S. Guo, S. Greenaway, R. Heller, Y. Herault, M. J. Justice, N. Kurbatova, C. J. Lelliott, K. C. K. Lloyd, A. M. Mallon, J. E. Mank, H. Masuya, C. McKerlie, T. F. Meehan, R. F. Mott, S. A. Murray, H. Parkinson, R. Ramirez-Solis, L. Santos, J. R. Seavitt, D. Smedley, T. Sorg, A. O. Speak, K. P. Steel, K. L. Svenson, C. International Mouse Phenotyping, S. Wakana, D. West, S. Wells, H. Westerberg, S. Yaacoby, J. K. White, Prevalence of sexual dimorphism in mammalian phenotypic traits. *Nat Commun* **8**, 15475 (2017).
6. *Drug Safety: Most Drugs Withdrawn in Recent Years Had Greater Health Risks for Women* (2001).
7. J. S. Mogil, Qualitative sex differences in pain processing: emerging evidence of a biased literature. *Nature Reviews Neuroscience* **21**, 353-365 (2020).
8. J. A. Clayton, Studying both sexes: a guiding principle for biomedicine. *The FASEB Journal* **30**, 519 (2016).
9. J. W. Rich-Edwards, D. L. Maney, Best practices to promote rigor and reproducibility in the era of sex-inclusive research. *Elife* **12**, e90623 (2023).
10. R. M. Shansky, A. Z. Murphy, Considering sex as a biological variable will require a global shift in science culture. *Nature neuroscience* **24**, 457-464 (2021).
11. S. K. Lee, Sex as an important biological variable in biomedical research. *BMB reports* **51**, 167 (2018).
12. I. Zucker, B. J. Prendergast, A. K. Beery, Pervasive neglect of sex differences in biomedical research. *Cold Spring Harbor perspectives in biology* **14**, a039156 (2022).
13. MRC. (UK Research and Innovation, https://www.ukri.org/councils/mrc/guidance-for-applicants/policies-and-guidance-for-researchers/sex-in-experimental-design/, 2022), vol. 2022.
14. R. K. Rechlin, T. F. Splinter, T. E. Hodges, A. Y. Albert, L. A. Galea, An analysis of neuroscience and psychiatry papers published from 2009 and 2019 outlines opportunities for increasing discovery of sex differences. *Nature communications* **13**, 2137 (2022).
15. N. C. Woitowich, A. Beery, T. Woodruff, A 10-year follow-up study of sex inclusion in the biological sciences. *Elife* **9**, (2020).
16. S. C. Stanford, S. Alexander, G. Cirino, C. H. George, P. A. Insel, D. Kendall, Y. Ji, R. A. Panettieri Jr, H. H. Patel, C. G. Sobey, Considering and reporting sex as an experimental variable II: An update on progress in the British Journal of Pharmacology. *British Journal of Pharmacology*, (2023).
17. Y. Garcia-Sifuentes, D. L. Maney, Reporting and misreporting of sex differences in the biological sciences. *Elife* **10**, (2021).
18. M. W. Group, Working Group on Sex in Experimental Design of Animal Research 2022.
19. N. C. Woitowich, T. K. Woodruff, Implementation of the NIH sex-inclusion policy: attitudes and opinions of study section members. *Journal of Women's Health* **28**, 9-16 (2019).

20. M. Waltz, J. A. Fisher, A. D. Lyerly, R. L. Walker, Evaluating the National Institutes of Health's
sex as a biological variable policy: conflicting accounts from the front lines of animal
research. *Journal of Women's Health* **30**, 348-354 (2021).
21. A. Gompers, "Three years in:"sex as a biological variable" policy in practice-and an invitation
to collaborate," (Genderscilab, 2018).
22. B. Phillips, T. N. Haschler, N. A. Karp, Statistical simulations show that scientists need not
increase overall sample size by default when including both sexes in in vivo studies. *Plos*
Biology **21**, e3002129 (2023).
23. N. A. Karp, N. Reavey, Sex bias in preclinical research and an exploration of how to change
the status quo. *Br J Pharmacol* **176**, 4107-4118 (2019).
24. J. B. Becker, B. J. Prendergast, J. W. Liang, Female rats are not more variable than male rats: a
meta-analysis of neuroscience studies. *Biology of sex differences* **7**, 1-7 (2016).
25. B. J. Prendergast, K. G. Onishi, I. Zucker, Female mice liberated for inclusion in neuroscience
and biomedical research. *Neuroscience & Biobehavioral Reviews* **40**, 1-5 (2014).
26. A. K. Beery, Inclusion of females does not increase variability in rodent research studies.
Current opinion in behavioral sciences **23**, 143-149 (2018).
27. R. D. Fields, NIH policy: mandate goes too far. *Nature* **510**, 340-340 (2014).
28. M. W. Group. (UK Research and Innovation, https://www.ukri.org/publications/sex-in-
experimental-design-summary-report/, 2022), chap. 9th March 2022.
29. I. Ajzen, The theory of planned behavior. *Organizational behavior and human decision*
processes **50**, 179-211 (1991).
30. UKRI, U. R. a. Innovation, Ed. (https://engagementhub.ukri.org/mrc-
regulatorysupportcentre/inclusion-sex-in-research-design/, 2023), vol. 2023.
31. P. Jirkof, N. Bratcher, L. Medina, D. Strasburg, P. Ebert, B. N. Gaskill, The effect of group size,
age and handling frequency on inter-male aggression in CD 1 mice. *Scientific reports* **10**, 2253
(2020).
32. W. B. Blankenberger, E. M. Weber, D. K. Chu, J. T. Geronimo, J. Theil, B. N. Gaskill, K. Pritchett-
Corning, M. A. Albertelli, J. P. Garner, J. Ahloy-Dallaire, Breaking up is hard to do: Does
splitting cages of mice reduce aggression? *Applied animal behaviour science* **206**, 94-101
(2018).
33. P. L. P. Van Loo, L. Van Zutphen, V. Baumans, Male management: coping with aggression
problems in male laboratory mice. *Laboratory animals* **37**, 300-313 (2003).
34. J. Kruger, D. Dunning, Unskilled and unaware of it: how difficulties in recognizing one's own
incompetence lead to inflated self-assessments. *Journal of personality and social psychology*
77, 1121 (1999).
35. H. Morahan, N. Hancock, L. Bero, K. Rooney, Experiences of early career researchers:
Influences on the design and reporting of animal experiments, and the practical and
emotional support needed to enhance best practice methods. *Laboratory Animals*,
00236772241242850 (2024).
36. M. R. LaFollette, S. Cloutier, C. M. Brady, M. E. O'Haire, B. N. Gaskill, Changing human
behavior to improve animal welfare: A longitudinal investigation of training laboratory
animal personnel about Heterospecific play or "rat tickling". *Animals* **10**, 1435 (2020).

7. Acknowledgements

Competing interests statement:

The authors declare the following competing interests: BP and NAK have shareholdings

in AstraZeneca. All other authors declare no competing interests.

Funding

The funders had no role in the design of the study; in the collection, analyses, or

interpretation of data; in the writing of the manuscript, or in the decision to publish

the results. This work was supported by grants for AA from The British Journal of

Pharmacology   (https://bpspubs.onlinelibrary.wiley.com/journal/14765381)   and

AstraZeneca (https://www.astrazeneca.com/).

Contribution

NAK: Conceptualization, Funding acquisition, Investigation, Visualization, Project

administration, Formal analysis, Supervision, Writing – original draft, Writing – review

& editing

BP: Investigation, Visualization, Writing – original draft, Writing – review & editing

BG: Conceptualization, Investigation, Formal analysis, Writing – original draft,

Writing – review & editing

HR: Investigation

OO: Investigation

AR: Investigation

JO: Investigation

AA: Funding acquisition, Investigation, Writing – original draft, Writing – review &

editing

Data availability statement

All data generated during this research has been made available within the
manuscript or via the supplementary information.

8. List of Supplementary Material

• Supplementary Table 1: Summary of demographic and potential predictors between
groups for all survey 1 contributors who met the inclusion criteria.

• Supplementary Table 2: Summary of demographic and potential predictors for all
participants who met the inclusion criteria for Study 2.

• Supplementary Analysis 1: Pearsons's correlation coefficient analysis between
continuous variables.

• Supplementary Analysis 2: For the perceived barriers, the Pearson's chi-square test
of association between treatment groups in study 1

• Supplementary Analysis 3: For the perceived benefits, the Pearson's chi-square test
of association between treatment groups in study 1

• Supplementary Analysis 4: For the perceived barriers, the McNemar's test of
association between treatment groups in study 2.

• Supplementary Analysis 5: For the perceived benefits, the McNemar's test of
association between treatment groups in study 2.

• Supplementary Analysis 6: For the knowledge questions, a Pearson's chi-squared
test to assess association between treatment groups of the proportion of participants
who answered the question correctly in study 1.

• Supplementary Analysis 7: For the knowledge questions, a McNemar's test to assess
association between pre and post answers for each knowledge question in study 2.

• Supplementary Table 3: Cumulative knowledge scores for study 1.

• Supplementary Table 4: Pre and post cumulative knowledge scores for study 2.

• Supplementary Analysis 8: Poisson regression analysis of cumulative knowledge

score for study 1.

• Supplementary Material 1: WCP survey

• Supplementary Material 2: Pre workshop survey

• Supplementary Material 3: Post workshop survey

• Supplementary Material 4: Workshop training material

Supplementary Table 1: Summary of demographic and potential predictors

between groups for all survey 1 contributors who met the inclusion criteria.

The demographic information included is for the full 105 participants that met the inclusion criteria. While all 105

met the inclusion criteria, 7 participants left the question about age blank. As age was included as a predictor in

the analysis of intent the missing values were managed with listwise deletion, assuming missing at random,

reducing the dataset size to 98 participants (N=35 baseline, N=48 interested and N=15 intervention). To test for a

statistically significant difference (association) between the treatment groups, a Pearson's Chi-square test was

used for categorical variables, ordered logistic regression for nominal variables and ANOVA test for continuous

variables. Institute type was collected as a demographic, the resulted population sampled was predominately

academic and this variable was therefore removed from downstream analysis due to the lack of predictive ability

to assess institute type on the outcome of interest. The abbreviation name in bracket, within the demographic

column, indicates the term used within the statistical model and associated output.

Demographic variable	Units or level	Full dataset Mean (range) or Count (% total)	Baseline (N=39)	Interested (N=51)	Intervention (N=15)	Test of association Test statistic & p value
Age	Years	39 (21-77)	41.1	36.4	42.7	$F_{2,95} = 1.9$; p = 0.15
Gender	Man	41 (39)	15	19	7	$X^2 = 3.3$; p = 0.51
	Woman	62 (59)	24	30	8	
	Other	2 (2)	0	2	0	
Geographic	Asia	16 (15)	4	9	3	$X^2 = 4.03$; p =

Region (Geography)	Australia	13 (12)	5	6	2	0.67
	Europe	62 (59)	22	32	8	
	Other	14 (13)	8	4	2	
Years working in animal research (Year_Work)	Years	14.1 (0-53)	16.0	11.7	17.6	$F_{2,100} = 1.9$; p = 0.15
Institution	Academic	101 (96)	38	49	14	$X^2 = 0.46$; p = 0.79
	Other	4 (4)	1	2	1	
Highest Education (Education)	Doctoral	62 (64)	24	28	10	$X^2 = 4.8$; p = 0.31
	Masters	26 (26)	7	14	5	
	Other	10 (10)	4	6	0	
Type of *in vivo* research (Type_Work)	Basic biological investigation	69 (66)	24	36	9	$X^2 = 3.5$; p = 0.75
	Discovery	14 (13)	6	6	2	
	Drug development / Lead optimization	13 (12)	4	7	2	
	Other	9 (9)	5	2	2	
Statistical background – Number of courses (Stats_Training)	>2 courses	18 (17)	7	8	3	Likelihood X^2 ratio = 0.41; p = 0.81
	1-2 courses	57 (54)	22	27	8	
	No formal courses	30 (29)	10	16	4	
Familiarity with Factorials (Factorial_Fam)	Extremely	5 (5)	2	2	1	Likelihood X^2 ratio = 3.73; p = 0.15
	Familiar	25 (24)	8	13	4	
	Somewhat familiar	33 (31)	16	9	8	
	Not very familiar	35 (33)	10	23	2	
	Not at all familiar	7 (7)	3	4	0	
How often previously incorporated sex? (Factorial_Incor)	76-100%	17 (16)	6	7	4	$F_{2,102} = 2.05$; p = 0.13
	51-75%	23 (22)	8	11	4	
	26-50%	22 (20)	9	8	5	
	1-25%	27 (26)	10	16	1	

	0%	16 (15)	6	9	1	
How often can you influence design? (Ability_Influence)	Always	45 (43)	19	19	7	Likelihood X^2 ratio = 0.65; p = 0.72
	Often	34 (32)	9	19	6	
	Sometimes	21 (20)	8	12	1	
	Rarely	5 (5)	3	1	1	
	Never	0 (0)	0	0	0	

Supplementary Table 2: Summary of demographic and potential predictors for all participants who met the inclusion criteria for Study 2.

The demographic information is for the full 31 participants that met the inclusion criteria in study 2. While 31 unique individuals met the criteria, an individual may not have responded to both surveys (N=29 pre-survey and N=28 in the post-survey). For instance, 2 participants did not meet the inclusion criteria for the pre-survey but met the criteria for the post-intervention survey. For the intention analysis, some missing data was observed in the demographic data. To reduce survey burden, the post-survey only included one demographic question (the participant's age) to support alignment of data just in case duplicate initials were used as an identifier. Two responders did not include the age information in the study and were managed with listwise deletion, assuming missing at random. The abbreviation name in bracket, within the demographic column, indicates the term used within the statistical model and associated output. For McNemar's paired analysis we conducted listwise deletion, assuming missing at random, reducing the dataset size to 26.

Demographic variable	Units or level	Participants Mean (range) or Count (% of total); N=31
Age	Years	Avg. 35.4 (range 25-60)
Gender	Man	11 (35)
	Woman	16 (52)
	Other	2 (6)
	Unknown	2 (6)
Type of *in vivo* research	Basic biological investigation	16 (52)
	Other	13 (13)

(Type_Work)	Unknown	2 (6)
Years working in animal research (Year_Work)	Years	Avg. 10.6 (range 0-35)
Highest Education (Education)	Doctoral	21 (68)
	Other	8 (26)
	Unknown	2 (6)
How often can you influence design? (Ability_Influence)	Always	15 (48)
	Often	7 (23)
	Sometimes	3 (10)
	Rarely	4 (4)
	Never	0 (0)
	Unknown	2 (6)
Statistical background – Number of courses (Stats_Training)	>2 courses	4 (13)
	1-2 courses	17 (55)
	No formal courses	8 (26)
	Unknown	2 (6)
Familiarity with Factorials (Factorial_Fam)	Not at all familiar	5 (16)
	Not very familiar	12 (39)
	Somewhat familiar	5 (16)
	Familiar	5 (16)
	Extremely familiar	2 (6)
	Unknown	2 (6)
How often previously incorporated sex? (Factorial_Incor)	Always	9 (29)
	Often	7 (23)
	Sometimes	2 (6)
	Rarely	9 (29)
	Never	2 (6)
	Always	2 (6)

Supplementary Analysis 1: Pearsons's correlation coefficient analysis

between continuous variables.

Where Year_Work represents the number of years the participants have worked in animal research, Education

represents the highest level of education obtained, Type_Work represents the type of research conducted by the

participant, Training represents the level of statistical training received, Factorial_Fam represents how familiar the

participants were with factorial experimental design, Factorial_Incor represents how often the participants

incorporated males and females into their experiments while studying an intervention, attitude represents the

average attitude score, Beh_control represents the average behavioural control score, Soc_norm represents the

average social normal score and Ability_Influence represents how often the participants were involved or could

influence the planning of experiments involving animals. Only variables that were used in the final statistical

model were compared for correlations.

Table A: Study 1

	Ability_Influence	Factorial_Fam	Factorial_Incor	Attitude	Beh_Control	Soc_Norm
Ability_Influence	1.0000	0.3499	0.2581	-0.0138	0.2218	0.1514
Factorial_Fam	0.3499	1.0000	0.3317	0.2172	0.2371	0.2341
Factorial_Incor	0.2581	0.3317	1.0000	0.2539	0.4496	0.4384
Attitude	-0.0138	0.2172	0.2539	1.0000	0.2081	0.1555
Beh_Control	0.2218	0.2371	0.4496	0.2081	1.0000	0.3691
Soc_Norm	0.1514	0.2341	0.4384	0.1555	0.3691	1.0000

987 Table B: Study 2

	Ability_Influence	Factorial_Fam	Factorial_Incor	Attitude	Beh_Control	Soc_Norm	Age
Ability_Influence	1.0000	0.1361	0.1862	0.1305	0.3818	0.2926	1.0000
Factorial_Fam	0.1361	1.0000	-0.2319	0.0714	0.3041	0.2671	0.1361
Factorial_Incor	0.1862	-0.2319	1.0000	-0.0689	0.1139	0.0407	0.1862
Attitude	0.1305	0.0714	-0.0689	1.0000	0.2664	0.0710	0.1305
Beh_Control	0.3818	0.3041	0.1139	0.2664	1.0000	0.1624	0.3818
Soc_Norm	0.2926	0.2671	0.0407	0.0710	0.1624	1.0000	0.2926
Age	0.2047	-0.0511	-0.0616	-0.3604	0.1078	0.1659	0.2047

 Supplementary Analysis 2: For the perceived barriers, the Pearson's chi-

 square test of association between treatment groups in study 1

 This survey question provided several pre-defined options and ability to enter a free-texted option. Participants

 were asked to choose all that applied. Exploration of the free text has grouped the barriers into three additional

 categories: convention, logistic, and none or no barriers. To test for a statistically significant difference

 (association) between the treatment groups, a Pearson's Chi-square test was applied for all options where the

 total N>10.

Barrier offered	Total (count) Possible N=105	Baseline (count) Possible N=39	Interested (count) Possible N=51	Intervention (count) Possible N=15	Test of association Test statistic & p value
Cost	44	14	24	6	$X^2 = 1.157; p = 0.561$
Male animals are more likely to fight and may lead to premature euthanasia	34	8	21	5	$X^2 = 4.317; p = 0.115$
Female animals are more variable	29	6	18	5	$X^2 = 4.668; p = 0.097$
Complexity of experimental design	28	7	18	3	$X^2 = 3.789; p = 0.149$
Model behavior may be different in the other sex	27	10	14	3	$X^2 = 0.337; p = 0.844$
Sample size concerns	24	7	13	4	$X^2 = 0.857; p = 0.651$
Not relevant to the research question	19	9	9	1	$X^2 = 1.98; p = 0.371$
Availability of sample/test material	12	2	7	3	$X^2 = 2.884; p = 0.236$
Welfare issues	5	2	2	1	NA
Data analysis concerns	0	0	0	0	NA
Other - Convention	7	2	3	2	NA
Other – Logistics	1	1	0	0	NA
Other - None	3	1	0	2	NA

 Supplementary Analysis 3: For the perceived benefits, the Pearson's chi-

 square test of association between treatment groups in study 1

1000   This question provided several pre-defined options and the ability to enter a free-texted option. Participants were

1001   asked to choose all that applied. No free text advantages were provided by survey takers for this question. To

1002   test for a statistically significant difference (association) between the treatment groups, a Pearson's chi-square

1003   test of association was applied for all options where the total N >10.

1004

Benefits offered	Total Possible N=105	Baseline (count) Possible N=39	Interested (count) Possible N=51	Intervention (count) Possible N=15	Test of association Test statistic & *p* value
Understanding sex differences	91	33	44	14	$X^2 = 0.726$; $p = 0.695$
Translatability	79	29	35	15	**$X^2 = 6.149$; $p = 0.046$**
Efficient use of all animals from breeding	42	12	18	12	**$X^2 = 11.855$; $p = 0.0026$**
Reproducibility	42	17	16	9	$X^2 = 4.291$; $p = 0.117$
3Rs – Reduction	33	11	14	8	$X^2 = 3.902$; $p = 0.142$
Animal welfare	22	10	7	5	$X^2 = 3.48$; $p = 0.176$
Other	0	0	0	0	NA

1005

Supplementary Analysis 4: For the perceived barriers, the McNemar's test of

association between treatment groups in study 2.

This question provided several pre-defined options and the ability to enter a free-texted option. Participants were

asked to choose all that applied. No free text advantages were provided by survey takers for this question. To

test for a statistically significant difference (association) between the treatment groups, a McNemar's test was

applied for all options where the total N >10.

Barrier offered	Pre-Intervention (count) Possible N=29	Post-Intervention (count) Possible N=28	McNemar's test Test statistic & *p* value
Cost	15	8	$X^2 = 3.57$; $p = 0.060$
Complexity of experimental design	14	5	**$X^2 = 5.33$; $p = 0.021$**
Sample size concerns	13	4	**$X^2 = 4.45$; $p = 0.035$**
Model behaviour may be different in the other sex	12	2	**$X^2 = 5.44$; $p = 0.020$**
Availability of sample/test material	10	9	$X^2 = 0.0$; $p = 1.0$
Female animals are more variable	9	2	**$X^2 = 5.0$; $p = 0.025$**

Experiment would take longer	4	5	NA
Male animals are more likely to fight and may lead to premature euthanasia	3	2	NA
Convention	2	4	NA
Data analysis concerns	2	4	NA
Logistics	3	4	NA
Welfare issues	2	1	NA
Not relevant to the research question	0	4	NA
Free text: Sponsor request	0	1	NA
Free text: Model induction issue	0	1	NA

1012

Supplementary Analysis 5: For the perceived benefits, the McNemar's test of association between treatment groups in study 2.

1013

1014

1015 This survey question provided several pre-defined options and ability to enter a free-texted option. Participants

1016 were asked to choose all that applied. Participants were given the option of a free text response, but none were

1017 submitted. To test for a statistically significant difference (association) between the pre and post measures, a

1018 McNemar's test was applied for all options where the total N >10. This test accounts for the repeat nature of the

1019 data which required the list wise deletion of those individuals with missing data.

Advantages offered	Pre-Intervention (count) Possible N=29	Post-Intervention (count) Possible N=28	McNemar's test Test statistic & p value
Understanding sex differences	26	25	$X^2 = 0.0$; $p = 1.0$
Translatability	21	22	$X^2 = 1.0$; $p = 0.32$
Efficient use of all animals from breeding	13	17	$X^2 = 1.6$; $p = 0.21$
3Rs - Reduction	8	13	$X^2 = 2.78$; $p = 0.096$
Reproducibility	6	13	$\mathbf{X^2 = 6.0}$; $\boldsymbol{p = 0.014}$
Animal welfare	3	5	NA

1020

Supplementary Analysis 6: For the knowledge questions, a Pearson's chi-

squared test to assess association between treatment groups of the

proportion of participants who answered the question correctly in study 1.

Question (Answer)	Total (Baseline & Interest) N=90	Baseline (% correct) N=39	Interest (% correct) N=51	Intervention (% correct) N=15	Test of association [Baseline v Interest] Test statistic & p value	Test of association [across the 3 treatment groups] Test statistic & p value
Do you think inclusion of both sexes requires doubling a study's sample size? (No)	20 (30.3%)	13 (33%)	7 (14%)	10 (67%)	**X^2 = 5.973; p = 0.014**	**X^2 = 16.85; p = 0.00021**
Do you think sex influences data variability, therefore when you include both sexes, more animals are needed? (No)	9 (10%)	5 (13%)	4 (8%)	6 (60%)	X^2 = 0.608; p = 0.4355	**X^2 = 7.994; p = 0.01837**
When analyzing *in vivo* data collected from both sexes, do you think sex should be included in the statistical model? (Yes)	68 (75.6%)	34 (87%)	34 (67%)	12 (80%)	**X^2 =5.035; p = 0.0248**	X^2 =5.266; p = 0.072
When analyzing *in vivo* data, do you think data from the two sexes should be pooled (combined) for an intervention into a single group for the analysis? (No)	17 (18.9%)	6 (15%)	11 (21%)	6 (40%)	X^2 =0.552; p = 0.457	X^2 =3.844; p = 0.1463
When	18	7	11	8	X^2 =0.181; p	**X^2 =7.823; p =**

| analyzing *in vivo* data collected from both sexes, do you think the analysis should be run independently for each sex through separate statistical tests? (No) | (20%) | (18%) | (21%) | (53%) | = 0.671 | **0.020** |

1024

Supplementary Analysis 7: For the knowledge questions, a McNemar's test to assess association between pre and post answers for each knowledge question in study 2.

Question (Answer)	Pre (% correct) N=29	Post (% correct) N=28	McNemar's test of association Test statistic & *p* value
Do you think inclusion of both sexes requires doubling a study's sample size? (No)	6 (21%)	17 (61%)	**X^2 =7.14; *p* = 0.007**
Do you think sex influences data variability, therefore when you include both sexes, more animals are needed? (No)	2 (7%)	13 (46%)	**X^2 =9.31; *p* = 0.0023**
When analyzing *in vivo* data collected from both sexes, do you think sex should be included in the statistical model? (Yes)	21 (72%)	21 (75%)	X^2 =0.143; *p* = 0.71
When analyzing *in vivo* data collected from both sexes, do you think the analysis should be disaggregated (for example run a t-test comparing control and intervention on the male data and then running a separate test on the data from females)? (No)	8 (28%)	22 (79%)	**X^2 =8.0; *p* = 0.0047**
When analyzing *in vivo* data, do you think data from the two sexes should be pooled (combined) for the analysis (for example run a single t-test comparing control and intervention group)? (No)	5 (17%)	18 (64%)	**X^2 =10.29; *p* = 0.0013**

1028

Supplementary Analysis 8: Intention data and SAS analysis code

The following provides the data and SAS code used to analyze study 1 and study 2 intention data. This information can be used for data and analysis transparency. Further, the information below can be cut and pasted directly into SAS.

Study 1 intent data and analysis code

```
DATA Survey_data_no_select_questions2; INPUT  InfluencePlanningExp2 StatsCourses2
&$11. Familiar_with_Factorials2 Prev_Sex_Incorporation2 Exposure_TRT &$17.
AvgAttitude AvgBehavControl AvgSocNorm What_is_your_age_ Gender2 &$ Geography2
&$9. How_many_years_have_you_worked_w Type_of_work2 &$19. Education2 &$
AvgIntent_Box_Cox_3_3_2; Lines;
4  1-2 courses  4  25  Interest in topic  7  5.66666666666667  4.66666666666667
32  Woman  Asia  10  Basic biol invest  Doctoral  3.17416428286284
3  1-2 courses  3  50  Interest in topic  7  4.66666666666667  6  29  Man  Asia  3
Drug devel/Lead opt  MS  3.54700262719911
3  No courses  2  0  Interest in topic  5.5  3.66666666666667  4.66666666666667
29  Woman  Australia  8  Basic biol invest  MS  1.61227392145414
5  1-2 courses  2  0  Interest in topic  7  4.33333333333333  2.33333333333333  26
Woman  Australia  6  Basic biol invest  Other  1.88434514569953
4  1-2 courses  3  25  Interest in topic  7  4.33333333333333  6.33333333333333
28  Woman  Australia  5  Discovery  Doctoral  3.93999439555356
3  1-2 courses  2  0  Interest in topic  5  2.66666666666667  3.66666666666667  27
Man  Australia  3  Basic biol invest  Other  0.916980792827043
5  No courses  2  100  Interest in topic  5.5  5.66666666666667  5.33333333333333
27  Woman  Europe  3  Basic biol invest  MS  4.35313958792618
3  1-2 courses  2  0  Interest in topic  6  4.66666666666667  3  51  Man  Asia  25
Drug devel/Lead opt  Doctoral  2.17656979396309
4  1-2 courses  2  25  Interest in topic  7  3.33333333333333  4  27  Man  Europe
7  Basic biol invest  MS  2.82147936254475
5  No courses  3  75  Interest in topic  7  5.66666666666667  6  53  Woman  Other
20  Basic biol invest  Doctoral  4.35313958792618
3  No courses  2  50  Interest in topic  4  4.66666666666667  5  40  Woman  Asia
15  Discovery  Doctoral  1.61227392145414
2  1-2 courses  2  0  Interest in topic  7  2.66666666666667  2  25  Woman  Europe
4  Other  Doctoral  2.48894786624483
3  1-2 courses  1  50  Interest in topic  7  5.66666666666667  6  56  Woman  Other
10  Drug devel/Lead opt  Doctoral  3.17416428286284
3  >2 courses  1  25  Interest in topic  6  3.33333333333333  2.33333333333333  62
Woman  Europe  35  Basic biol invest  Doctoral  1.1285917450179
5  1-2 courses  4  75  Interest in topic  7  3.33333333333333  5.66666666666667
43  Woman  Europe  13  Basic biol invest  Doctoral  3.54700262719911
5  1-2 courses  2  25  Interest in topic  5.75  4  1.66666666666667  32  Man
Europe  6  Basic biol invest  Doctoral  1.61227392145414
5  1-2 courses  4  75  Interest in topic  7  4  5  36  Woman  Europe  4  Basic
biol invest  Doctoral  3.93999439555356
5  >2 courses  4  50  Interest in topic  6.75  5  4.66666666666667  35  Woman
Europe  15  Basic biol invest  Doctoral  4.35313958792618
4  >2 courses  3  25  Interest in topic  7  4.33333333333333  3  27  Woman  Europe
5  Discovery  MS  3.93999439555356
5  1-2 courses  4  100  Interest in topic  7  4.66666666666667  5.66666666666667
57  Woman  Europe  35  Basic biol invest  Doctoral  4.35313958792618
5  1-2 courses  2  25  Interest in topic  6.75  3  5.33333333333333  30  Man
Europe  7  Basic biol invest  Doctoral  1.61227392145414
5  1-2 courses  4  100  Interest in topic  7  6.66666666666667  5.33333333333333
49  Woman  Europe  20  Discovery  Other  4.35313958792618
5  1-2 courses  5  75  Interest in topic  7  5  6.66666666666667  35  Woman
Europe  12  Basic biol invest  Doctoral  4.35313958792618
3  1-2 courses  3  100  Interest in topic  7  4.66666666666667  5  28  Woman
Europe  3  Basic biol invest  MS  2.82147936254475
4  No courses  2  100  Interest in topic  7  6.33333333333333  6  30  Woman
Europe  5  Basic biol invest  Doctoral  3.17416428286284
4  1-2 courses  2  75  Interest in topic  7  4.33333333333333  2.66666666666667
44  Woman  Europe  5  Drug devel/Lead opt  Doctoral  2.82147936254475
```

3 No courses 2 0 Interest in topic 6.5 2.33333333333333 2 30 Man Europe 7 Discovery MS 1.1285917450179

5 >2 courses 2 25 Interest in topic 4 4.66666666666667 4.33333333333333 35 Woman Asia 12 Basic biol invest Doctoral 1.36035612122693

4 No courses 3 100 Interest in topic 7 6 5.66666666666667 30 Man Europe 8 Basic biol invest Doctoral 3.17416428286284

4 No courses 2 0 Interest in topic 5.25 4.33333333333333 4.33333333333333 29 Man Asia 5 Basic biol invest MS 1.36035612122693

3 1-2 courses 2 0 Interest in topic 7 2 1.66666666666667 24 Woman Europe 2 Basic biol invest MS 2.48894786624483

5 1-2 courses 2 25 Interest in topic 7 5.33333333333333 5.66666666666667 34 Man Europe 13 Basic biol invest Doctoral 3.54700262719911

4 No courses 2 25 Interest in topic 6 4.33333333333333 4 24 Woman Asia 0 Basic biol invest MS 2.17656979396309

5 >2 courses 3 75 Interest in topic 6 5 6 28 Man Asia 5 Basic biol invest MS 3.17416428286284

4 1-2 courses 2 25 Interest in topic 7 3 5.33333333333333 23 Woman Other 4 Basic biol invest Other 4.35313958792618

5 >2 courses 4 50 Interest in topic 5.75 4.66666666666667 5.33333333333333 44 Man Europe 20 Drug devel/Lead opt Doctoral 2.17656979396309

4 1-2 courses 1 25 Interest in topic 5 4 4 26 Man Europe 3 Drug devel/Lead opt Other 2.82147936254475

5 No courses 2 100 Interest in topic 7 6 6 24 Woman Europe 1 Basic biol invest Other 4.35313958792618

3 1-2 courses 3 25 Interest in topic 6 3.66666666666667 5 50 Man Europe 3 Basic biol invest Doctoral 2.48894786624483

3 1-2 courses 1 25 Interest in topic 4.75 4 4 25 Woman Europe 2 Basic biol invest MS 1.88434514569953

4 No courses 4 75 Interest in topic 6 4.33333333333333 5 35 Man Australia 3 Drug devel/Lead opt MS 2.82147936254475

4 No courses 4 25 Interest in topic 6 3.33333333333333 4.66666666666667 65 Man Europe 38 Basic biol invest Doctoral 1.36035612122693

2 1-2 courses 3 0 Gen Pop 6.5 4.66666666666667 2.66666666666667 22 Man Europe 4 Basic biol invest Other 1.88434514569953

2 1-2 courses 3 25 Gen Pop 6.75 4 3.66666666666667 21 Woman Europe 1 Other Other 3.54700262719911

5 >2 courses 3 50 Gen Pop 7 4.66666666666667 7 53 Man Australia 32 Discovery Doctoral 4.35313958792618

5 No courses 3 75 Gen Pop 6.25 3.33333333333333 6 54 Woman Australia 35 Discovery Doctoral 3.93999439555356

5 >2 courses 3 25 Gen Pop 7 7 2.33333333333333 77 Man Other 53 Basic biol invest Doctoral 1.36035612122693

5 1-2 courses 3 50 Gen Pop 7 5 5.66666666666667 54 Woman Australia 25 Basic biol invest Doctoral 3.17416428286284

5 1-2 courses 3 0 Gen Pop 4.25 3.33333333333333 3.33333333333333 27 Man Asia 2 Basic biol invest MS 2.17656979396309

3 1-2 courses 1 50 Gen Pop 7 4 5 21 Man Europe 1 Basic biol invest Other 3.93999439555356

4 >2 courses 4 25 Gen Pop 7 4.33333333333333 4.33333333333333 42 Woman Other 7 Basic biol invest Doctoral 2.17656979396309

5 1-2 courses 2 0 Gen Pop 7 5 1.33333333333333 29 Woman Europe 2 Basic biol invest Doctoral 1.36035612122693

5 1-2 courses 4 50 Gen Pop 6.5 5.66666666666667 4 51 Man Other 30 Basic biol invest Doctoral 3.54700262719911

5 1-2 courses 4 25 Gen Pop 3.75 2.66666666666667 3 56 Woman Europe 25 Discovery Doctoral 0.272071224245386

3 No courses 3 75 Gen Pop 7 3.66666666666667 6 41 Man Other 3 Drug devel/Lead opt Doctoral 3.17416428286284

5 >2 courses 3 50 Gen Pop 7 4.33333333333333 3.66666666666667 37 Man Europe 12 Basic biol invest Doctoral 1.1285917450179
3 1-2 courses 3 100 Gen Pop 7 4 4 24 Woman Other 3 Other MS 2.48894786624483
5 1-2 courses 3 50 Gen Pop 7 6 6.66666666666667 42 Woman Europe 19 Basic biol invest Doctoral 4.35313958792618
5 1-2 courses 2 25 Gen Pop 6.25 3 5.33333333333333 34 Woman Europe 5 Basic biol invest Doctoral 3.17416428286284
3 No courses 2 0 Gen Pop 7 2.33333333333333 5.33333333333333 24 Woman Europe 1 Basic biol invest MS 2.82147936254475
2 No courses 2 100 Gen Pop 7 4.66666666666667 5.66666666666667 26 Woman Europe 1 Drug devel/Lead opt MS 1.88434514569953
4 No courses 4 75 Gen Pop 7 4 5.33333333333333 55 Man Europe 30 Basic biol invest Doctoral 3.17416428286284
4 1-2 courses 2 25 Gen Pop 5.75 4.33333333333333 3.33333333333333 75 Man Europe 52 Discovery Doctoral 1.61227392145414
3 1-2 courses 2 25 Gen Pop 7 3.33333333333333 4.33333333333333 23 Woman Europe 0.5 Basic biol invest Other 3.93999439555356
5 No courses 2 50 Gen Pop 7 5 4.66666666666667 32 Woman Europe 8 Basic biol invest Doctoral 3.17416428286284
5 1-2 courses 5 75 Gen Pop 7 6 5.66666666666667 56 Woman Australia 35 Discovery Doctoral 4.35313958792618
4 1-2 courses 3 0 Gen Pop 2.5 4.33333333333333 6.33333333333333 30 Woman Europe 3 Basic biol invest MS 0.554219160499861
4 No courses 2 25 Gen Pop 6.25 3 4.66666666666667 37 Woman Other 10 Basic biol invest Doctoral 2.17656979396309
4 No courses 2 75 Gen Pop 7 3.66666666666667 5.33333333333333 74 Woman Other 40 Other Doctoral 3.17416428286284
3 1-2 courses 1 0 Gen Pop 7 4.66666666666667 6 38 Woman Other 10 Basic biol invest MS 3.17416428286284
3 No courses 2 75 Gen Pop 6 5 5.33333333333333 51 Woman Australia 30 Basic biol invest Doctoral 3.17416428286284
5 1-2 courses 4 100 Gen Pop 6 5 6.33333333333333 25 Woman Europe 8 Drug devel/Lead opt MS 4.35313958792618
5 1-2 courses 5 75 Gen Pop 7 2.66666666666667 4 45 Woman Europe 20 Basic biol invest Doctoral 4.35313958792618
4 No courses 4 25 Gen Pop 7 6 3.33333333333333 55 Man Asia 35 Basic biol invest Doctoral 3.17416428286284
4 >2 courses 4 100 Gen Pop 6.5 5.33333333333333 4 40 Man Europe 10 Other Doctoral 2.48894786624483
5 1-2 courses 3 75 Gen Pop 7 3.66666666666667 6.33333333333333 39 Man Europe 15 Basic biol invest Doctoral 3.17416428286284
5 1-2 courses 1 100 Gen Pop 5 3.66666666666667 4.33333333333333 27 Woman Asia 4 Other Doctoral 1.36035612122693
5 >2 courses 5 75 Interest in topic 7 4.33333333333333 3.33333333333333 56 Man Europe 30 Basic biol invest Doctoral 2.17656979396309
4 No courses 4 50 Interest in topic 7 4 3 41 Woman Australia 8 Basic biol invest Doctoral 3.17416428286284
4 >2 courses 3 75 Interest in topic 7 4.33333333333333 5 38 Other Europe 15 Basic biol invest Doctoral 2.17656979396309
4 1-2 courses 4 25 Interest in topic 7 4.33333333333333 5.66666666666667 46 Woman Europe 23 Basic biol invest Doctoral 3.17416428286284
4 1-2 courses 4 50 Interest in topic 6 5 4.66666666666667 29 Woman Asia 7 Basic biol invest MS 0.272071224245386
5 No courses 2 0 Interest in topic 6.75 4 5.66666666666667 51 Woman Europe 30 Basic biol invest Doctoral 4.35313958792618
5 No courses 3 100 Workshop 5.5 6 5.66666666666667 27 Woman Europe 3 Basic biol invest MS 4.35313958792618

```
1210   5  1-2 courses  5  50  Workshop  7  4  7  35  Woman  Europe  12  Basic biol invest
1211   Doctoral  3.54700262719911
1212   5  1-2 courses  4  25  Workshop  7  5  6.66666666666667  52  Man  Australia  30
1213   Discovery  Doctoral  4.35313958792618
1214   2  No courses  4  75  Workshop  7  6.33333333333333  6.66666666666667  57  Man
1215   Europe  25  Basic biol invest  Doctoral  3.93999439555356
1216   4  1-2 courses  2  0  Workshop  6  2.33333333333333  1  26  Woman  Other  6  Drug
1217   devel/Lead opt  MS  2.48894786624483
1218   5  1-2 courses  4  75  Workshop  7  5  5.66666666666667  36  Woman  Europe  5
1219   Basic biol invest  Doctoral  4.35313958792618
1220   4  >2 courses  4  50  Workshop  7  5.33333333333333  6.33333333333333  35  Woman
1221   Europe  15  Basic biol invest  Doctoral  3.93999439555356
1222   4  1-2 courses  3  50  Workshop  7  4  3.33333333333333  58  Woman  Europe  35
1223   Other  Doctoral  3.17416428286284
1224   4  No courses  3  75  Workshop  7  4.66666666666667  6  30  Man  Europe  8  Basic
1225   biol invest  Doctoral  3.93999439555356
1226   4  1-2 courses  3  100  Workshop  6.5  4.66666666666667  4.66666666666667  32
1227   Woman  Asia  7  Other  MS  3.17416428286284
1228   5  1-2 courses  3  100  Workshop  7  5.66666666666667  4.33333333333333  57  Woman
1229   Europe  35  Basic biol invest  Doctoral  4.35313958792618
1230   5  No courses  3  100  Workshop  7  6  6  58  Man  Australia  38  Discovery
1231   Doctoral  4.35313958792618
1232   4  >2 courses  3  50  Workshop  7  6  5  33  Man  Other  12  Basic biol invest
1233   Doctoral  3.17416428286284
1234   5  1-2 courses  3  75  Workshop  6  3.66666666666667  3  52  Man  Asia  30  Drug
1235   devel/Lead opt  MS  3.17416428286284
1236   3  >2 courses  2  50  Workshop  7  5  7  52  Man  Asia  3  Basic biol invest  MS
1237   4.35313958792618
1238   ;
1239   RUN;
1240
1241   PROC GLM  DATA=Survey_data_no_select_questions2 ALPHA=0.05;
1242   CLASS StatsCourses2 Exposure_TRT Gender2 Geography2 Type_of_work2 Education2;
1243   MODEL  AvgIntent_Box_Cox_3_3_2 =  AvgAttitude AvgBehavControl AvgSocNorm
1244   Exposure_TRT What_is_your_age_ Gender2 Geography2 How_many_years_have_you_worked_w
1245   Type_of_work2 Education2 StatsCourses2 Familiar_with_Factorials2
1246   Prev_Sex_Incorporation2 InfluencePlanningExp2;
1247   RUN;
1248
1249   Study 2 data and analysis code
1250
1251
1252   DATA TPB_no_select_analysis20240319; INPUT  ParticipantInitials &$ Pre_Post &&$16.
1253   InfluencePlanningExp2 FamilarwithFactorials2 HowOftenIncorpSex2 AvgAttitude
1254   AvgBehavControl AvgSocNorm AvgIntent What_is_your_age_ Gender2 &$; Lines;
1255   MIP  Pre  4  1  25  7  5.33333333333333  4  5.33333333333333  39  Woman
1256   SSS  Pre  4  4  100  5.75  5.66666666666667  6.33333333333333  7  38  Woman
1257   CKTL  Pre  1  2  0  7  2.66666666666667  4  5  27  Man
1258   AAA  Pre  4  4  50  7  3.33333333333333  6.33333333333333  6.33333333333333  36  W
1259   oman
1260   EAS  Pre  4  3  75  7  4  6  5.33333333333333  36  Man
1261   YD  Pre  3  1  100  7  5  5  7  27  Woman
1262   MSU  Pre  4  2  25  7  3  5.66666666666667  6  42  Man
1263   AJH  Pre  4  4  100  7  6.66666666666667  4.66666666666667  6  55  Man
1264   AC  Pre  4  2  75  7  5  6.33333333333333  7  36  Woman
1265   RBN  Pre  3  2  25  5  2  5.66666666666667  4.66666666666667  46  Woman
1266   AJM  Pre  4  3  100  6.5  5  4  6  42  Woman
1267   AMS  Pre  4  4  25  7  3.66666666666667  6.66666666666667  6  50  Other
1268   JJD  Pre  4  5  75  7  4.66666666666667  7  7  25  Woman
```

```
1269    MC   Pre  1  4  100  7  4.33333333333333  5.66666666666667  7  29   Woman
1270    EMJ  Pre  4  2  100  6.75  6  4.66666666666667  6.66666666666667  29   Woman
1271    SFA  Pre  3  5  25  6.5  3  4.33333333333333  4  30  Other
1272    PEE  Pre  3  3  25  7  4.33333333333333  4.66666666666667  5  .  Man
1273    CMG  Pre  4  2  50  7  3.66666666666667  3  4  25  Woman
1274    DDD  Pre  2  2  25  6.25  4  4  6  28  Man
1275    ksk  Pre  3  2  25  5  3.33333333333333  5.66666666666667  5.66666666666667  60  M
1276    an
1277    RYX  Pre  3  1  75  7  2  6  6  38  Woman
1278    DRH  Pre  1  2  75  7  3.66666666666667  5.33333333333333  6  27  Man
1279    SMC  Pre  2  2  75  6.5  3  3  2.66666666666667  33  Woman
1280    NNK  Pre  2  1  75  7  2.66666666666667  4.33333333333333  5.33333333333333  35  W
1281    oman
1282    GJF  Pre  4  2  100  7  4  7  7  28  Man
1283    OMO  Pre  4  3  100  7  4  5.66666666666667  6.33333333333333  35  Woman
1284    SSS  Post  4  4  100  6.25  6.33333333333333  6.33333333333333  6  38  Other
1285    JWH  Pre  1  3  0  4.75  2  4.33333333333333  4.33333333333333  43  Man
1286    SFA  Post  2  5  0  6.75  3.66666666666667  3.66666666666667  3  30  Other
1287    DRH  Post  1  5  0  7  5.33333333333333  6.33333333333333  6.33333333333333  28  O
1288    ther
1289    EAS  Post  4  4  0  7  3.66666666666667  7  7  36  Other
1290    AJH  Post  4  4  0  7  6  5.33333333333333  6  .  Other
1291    MIP  Post  4  4  0  7  6  4  6  39  Other
1292    Tw   Post  3  3  0  7  3.66666666666667  5  7  29  Other
1293    EMJ  Post  4  4  0  6.75  6  4.66666666666667  7  29  Other
1294    AC   Post  4  4  0  7  6.33333333333333  7  7  36  Other
1295    AAA  Post  4  5  0  7  3.66666666666667  7  7  36  Other
1296    PEE  Post  3  3  0  7  6  5.66666666666667  5  56  Other
1297    EJS  Post  4  2  0  7  4.33333333333333  4.33333333333333  6  34  Other
1298    MC   Post  2  3  0  7  4  4.33333333333333  7  29  Other
1299    JWH  Post  1  4  0  7  5  6.66666666666667  4.66666666666667  43  Other
1300    CKTL Post  1  2  0  7  4  4.33333333333333  4.66666666666667  27  Other
1301    AJM  Post  4  4  0  7  5.66666666666667  4.33333333333333  6  42  Other
1302    RYX  Post  3  3  0  7  5  6.33333333333333  7  38  Other
1303    MI   Pre  4  1  25  7  4.33333333333333  4.66666666666667  5.33333333333333  39  Wo
1304    man
1305    ksk  Post  3  3  0  6.5  4.33333333333333  6  6  60  Other
1306    DDD  Post  1  2  0  7  3.66666666666667  5.33333333333333  7  28  Other
1307    JJD  Post  4  5  0  7  5  6.66666666666667  7  26  Other
1308    CMG  Post  3  4  0  7  3.33333333333333  4  5  25  Other
1309    RBN  Post  3  2  0  7  4  6  5.66666666666667  46  Other
1310    NNK  Post  2  3  0  7  4  4.66666666666667  5.66666666666667  35  Other
1311    OMO  Post  4  4  0  7  4  6  6  35  Other
1312    GJF  Post  4  4  0  7  4.33333333333333  7  7  28  Other
1313    SMC  Post  2  4  0  7  3.33333333333333  2.66666666666667  4  33  Other
1314    ABC  Post  3  2  0  7  4  6.33333333333333  6  30  Other
1315    MSU  Post  4  3  0  7  4  6  6  42  Other
1316    ;
1317    RUN;
1318
1319    PROC MIXED ASYCOV NOBOUND  DATA=TPB_no_select_analysis20240319 ALPHA=0.05;
1320    CLASS ParticipantInitials Pre_Post Gender2;
1321    MODEL  AvgIntent =  AvgAttitude AvgBehavControl AvgSocNorm Pre_Post
1322    What_is_your_age_ Gender2 InfluencePlanningExp2 FamilarwithFactorials2
1323    HowOftenIncorpSex2/ SOLUTION DDFM=KENWARDROGER;
1324    RANDOM ParticipantInitials / SOLUTION ;
1325    RUN;
1326
```

1327 **Supplementary Table 3: cumulative knowledge score for study 1.**

Participant ID	Treatment group	Gender	Cumulative Knowledge Score
93	Symposium	Woman	1
96	Symposium	Woman	1
97	Symposium	Man	0
102	Symposium	Man	3
105	Symposium	Man	1
109	Symposium	Woman	2
111	Symposium	Woman	0
113	Symposium	Woman	1
114	Symposium	Woman	0
116	Symposium	Woman	1
117	Symposium	Woman	0
122	Symposium	Woman	1
123	Symposium	Woman	3
124	Symposium	Woman	3
125	Symposium	Woman	2
126	Symposium	Woman	2
127	Symposium	Man	1
129	Symposium	Man	2
130	Symposium	Man	1
132	Symposium	Woman	1
133	Symposium	Man	3
134	Symposium	Woman	1
136	Symposium	Woman	1
138	Symposium	Man	1
140	Symposium	Woman	0
147	Symposium	Man	2
68	Symposium	Man	3
159	Symposium	Woman	3
162	Symposium	Woman	0
201	Symposium	Man	1
72	Conference	Woman	1
73	Conference	Man	1
94	Conference	Woman	1
99	Conference	Woman	2
107	Conference	Man	1
110	Conference	Woman	0
112	Conference	Woman	2

115	Conference	Man	3
118	Conference	Woman	1
120	Conference	Woman	3
121	Conference	Man	2
128	Conference	Woman	1
131	Conference	Man	1
135	Conference	Man	1
139	Conference	Man	1
141	Conference	Man	0
142	Conference	Woman	0
144	Conference	Man	1
10	Conference	Man	1
11	Conference	Woman	2
12	Conference	Man	4
13	Conference	Man	1
14	Conference	Woman	2
15	Conference	Man	2
16	Conference	Woman	1
20	Conference	Man	1
21	Conference	Man	1
22	Conference	Woman	2
23	Conference	Woman	1
24	Conference	Woman	0
27	Conference	Man	3
28	Conference	Man	2
29	Conference	Woman	2
35	Conference	Man	2
36	Conference	Man	2
38	Conference	Woman	1
39	Conference	Woman	2
40	Conference	Woman	1
41	Conference	Woman	1
42	Conference	Woman	0
44	Conference	Man	1
45	Conference	Man	1
46	Conference	Woman	2
47	Conference	Woman	1
49	Conference	Woman	2
50	Conference	Woman	1
52	Conference	Woman	1
53	Conference	Woman	1

54	Conference	Woman	2
55	Conference	Woman	3
56	Conference	Woman	4
58	Conference	Woman	2
62	Conference	Woman	1
63	Conference	Man	3
64	Conference	Man	3
65	Conference	Man	2
67	Conference	Woman	1
69	Conference	Woman	0
153	Conference	Other	2
170	Conference	Woman	2
175	Workshop	Woman	3
176	Workshop	Woman	3
177	Workshop	Man	5
178	Workshop	Man	4
179	Workshop	Woman	1
192	Workshop	Woman	5
193	Workshop	Woman	4
194	Workshop	Woman	3
195	Workshop	Man	2
196	Workshop	Woman	2
197	Workshop	Woman	3
198	Workshop	Man	3
199	Workshop	Man	0
200	Workshop	Man	3

1328

1329 **Supplementary Table 4: Pre- and post-cumulative knowledge scores for study**

1330 **2.**

Participant Initials	Pre-workshop: Cumulative score	Post-workshop: Cumulative score
AAA	4	5
ABC	1	1
AC	3	5
AJH	3	0
AJM	2	2
CKTL	1	3
CMG	2	3

DDD	3	5
DRH	2	4
EAS	2	5
EMJ	0	4
GJF	4	4
JJD	1	2
JWH	1	5
ksk	1	3
MC	1	3
MIP	0	4
MSU	0	5
NNK	1	5
OMO	1	5
PEE	0	3
RBN	1	3
RYX	1	0
SFA	3	0
SMC	1	5
SSS	1	2

1331

Supplementary Analysis 9: Poisson regression analysis of cumulative knowledge score for study 1.

Study 1

Data Preparation

Import:

```
surveydf <- read.csv("Survey1_CumulativeKnowledgeMetric.csv")
```

Wrangling:

```
summary(as.factor(surveydf$Gender2))
```

```
##    Man Other Woman
##     41     1    62
```

```
surveydf_2sexes <- subset(surveydf, surveydf$Gender2 != "Other")
levels(as.factor(surveydf_2sexes$Gender2))
```

```
## [1] "Man"   "Woman"
```

Visualisation study 1

Visual inspection of data suggests an effect of intervention that is not dependent on gender.

```
pcPlot <- ggplot(data = surveydf, aes(y = CumulativeKnowledgeScore,x = Gro
up, col = Group)) + ylab("Cumulative knowledge Score") + xlab("Group")
pcPlot + geom_violin() + theme(axis.text.x = element_text(angle = 45, vjus
t = 1, hjust = 1))
```

```
pcPlot <- ggplot(data = surveydf_2sexes, aes(y = CumulativeKnowledgeScore,
x = Group, col = Group)) + ylab("Cumulative knowledge Score") + xlab("Grou
p")
pcPlot + geom_violin() + theme(axis.text.x = element_text(angle = 45, vjus
t = 1, hjust=1))+ facet_grid(. ~ Gender2)
```

1359

Statistical Analysis Study 1

1360

A Poisson regression analysis assessing the role of treatment group in explaining
variation in the cumulative knowledge score.

Reference group is the conference.

Conclusions:

1.   Symposium group is not significantly different to reference.

2.   Workshop is statistically significantly different with B=0.66342 indicating the
expected count is (exp(B))= 1.94 fold larger than the reference group.

```
1368    #Assumptions of a Poisson distribution
1369    #Poisson Response the response variable is a count per unit of time or spa
1370    ce, described by a Poisson distribution.
1371    #Independence The observations must be independent of one another.
1372    #Mean=Variance By definition, the mean of a Poisson random variable must b
1373    e equal to its variance.
1374    #Linearity The log of the mean rate, log( λ), must be a linear function of
1375     x.
1376
1377    poisson.model <- glm(CumulativeKnowledgeScore ~ Group, surveydf_2sexes, fa
1378    mily = poisson(link = "log"))
1379    summary(poisson.model)
1380
1381    ## Call:
1382    ## glm(formula = CumulativeKnowledgeScore ~ Group, family = poisson(link =
```

```
 "log"),
##     data = surveydf_2sexes)
##
## Coefficients:
##                  Estimate Std. Error z value Pr(>|z|)
## (Intercept)       0.41110    0.10600   3.878 0.000105 ***
## GroupSymposium   -0.09872    0.18875  -0.523 0.600943
## GroupWorkshop     0.66342    0.18875   3.515 0.000440 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##     Null deviance: 91.143  on 102  degrees of freedom
## Residual deviance: 77.614  on 100  degrees of freedom
## AIC: 304.83
##
## Number of Fisher Scoring iterations: 5

#model checks
# 1: Check whether the model has over-dispersion or under-dispersion. If t
he Residual Deviance is greater than the degrees of freedom, then over-dis
persion exists. This means that the estimates are correct, but the standar
d errors (standard deviation) are wrong and unaccounted for by the model.
#the residual deviance < df suggesting over-dispersion is not an issue

#2 goodness fit test testing hypothesis that the model is appropriate - p-
value of Residual deviance goodness-of-fit test
1 - pchisq(deviance(poisson.model), df = poisson.model$df.residual)

## [1] 0.9526611

#3 Diagnostic plots
par(mfrow = c(2,3))
plot(poisson.model, which = 1:6)
```

1416

```
#https://www.statology.org/how-to-identify-influential-data-points-using-c
ooks-distance/#:~:text=A%20general%20rule%20of%20thumb,to%20identify%20inf
luential%20data%20points.
#rule of thumb  - any point with a Cook's Distance over 4/n (where n is th
e total number of data points) is considered to be an outlier.
#4/103
#diagnostics looks acceptable

# Calculating the ES as a meaningful value - If б > 0, then exp(б) > 1, an
d the expected count is exp(б) times larger than when X = 0
exp(0.66342)

## [1] 1.941421

plot_summs(poisson.model, scale = TRUE, exp = TRUE)

## Registered S3 methods overwritten by 'broom':
##   method             from
##   tidy.glht          jtools
##   tidy.summary.glht  jtools
```

1434
```
## Loading required namespace: broom.mixed
```

1435

Session Info

1437
```r
pander(sessionInfo())
```

R version 4.3.1 (2023-06-16 ucrt)

Platform: x86_64-w64-mingw32/x64 (64-bit)

locale: *LC_COLLATE=English_United Kingdom.utf8, LC_CTYPE=English_United*
Kingdom.utf8, LC_MONETARY=English_United Kingdom.utf8, LC_NUMERIC=C
and *LC_TIME=English_United Kingdom.utf8*

attached base packages: *stats, graphics, grDevices, utils, datasets, methods* and
base

other attached packages: *here(v.1.0.1), emmeans(v.1.10.4), pander(v.0.6.5),*
jtools(v.2.2.2) and *ggplot2(v.3.5.1)*

loaded via a namespace (and not attached): *broom.mixed(v.0.2.9.5)*, *utf8(v.1.2.3)*,
future(v.1.34.0), *generics(v.0.1.3)*, *tidyr(v.1.3.0)*, *lattice(v.0.21-8)*, *listenv(v.0.9.0)*,
digest(v.0.6.37), *magrittr(v.2.0.3)*, *evaluate(v.1.0.0)*, *grid(v.4.3.1)*,
estimability(v.1.5.1), *mvtnorm(v.1.2-2)*, *fastmap(v.1.2.0)*, *rprojroot(v.2.0.3)*,
backports(v.1.5.0), *purrr(v.1.0.1)*, *fansi(v.1.0.4)*, *scales(v.1.3.0)*, *codetools(v.0.2-20)*,
cli(v.3.6.1), *rlang(v.1.1.1)*, *crayon(v.1.5.3)*, *parallelly(v.1.38.0)*, *splines(v.4.3.1)*,
munsell(v.0.5.0), *withr(v.2.5.0)*, *yaml(v.2.3.7)*, *parallel(v.4.3.1)*, *tools(v.4.3.1)*,
coda(v.0.19-4.1), *dplyr(v.1.1.2)*, *colorspace(v.2.1-0)*, *forcats(v.1.0.0)*,
globals(v.0.16.2), *broom(v.1.0.7)*, *vctrs(v.0.6.3)*, *R6(v.2.5.1)*, *lifecycle(v.1.0.3)*,
MASS(v.7.3-60), *furrr(v.0.3.1)*, *pkgconfig(v.2.0.3)*, *pillar(v.1.9.0)*, *gtable(v.0.3.3)*,
glue(v.1.6.2), *Rcpp(v.1.0.11)*, *xfun(v.0.39)*, *tibble(v.3.2.1)*, *tidyselect(v.1.2.0)*,
highr(v.0.10), *rstudioapi(v.0.15.0)*, *knitr(v.1.43)*, *farver(v.2.1.2)*, *xtable(v.1.8-4)*,
nlme(v.3.1-162), *htmltools(v.0.5.8.1)*, *rmarkdown(v.2.23)*, *labeling(v.0.4.2)* and
compiler(v.4.3.1)

1461