The carnosinase dipeptidase CNDP1 is a novel metabolic vulnerability in brain

metastasis

Maria A. Gomez-Munoz[1, 2, #], Alcida Karz[1, 2, #], Maya Navarro[1, 2], Victoria Osorio-Vasquez[1, 2, 3],

Olga Katsara[4], Adam Walker[5,8], Pietro Berico[1, 2], A. Paulina Medellin[9], Nicole M. Eskow[1, 2],

Yunjing Wan[5], Milad Ibrahim[6], Somnath Tagore[10], Edridge K. D'Souza[10], Benjamin Izar[10],

Eleazar Vega Saenz de Miera[6], Iman Osman[6], Michael Pacold[3], Xiangpeng Kong[7], Mitchell

Levesque[11], Richard Possemato[1,3], Sheri L. Holmen[9], Robert J.Schneider[4], C. Theresa

Vincent[4,12], Kelly V. Ruggles[5,8], Douglas E. Biancur[1, 3, *], Eva Hernando[1, 2,*].

Departments of [1]Pathology, [3]Radiation Oncology, [4]Microbiology, [5]Medicine, [6]Dermatology,

[7]Biochemistry and Molecular Pharmacology, NYU Grossman School of Medicine, New York, NY

10016, United States

[2] Interdisciplinary Melanoma Cooperative Group, Perlmutter Cancer Center, NYU Langone

Health, New York, NY, 10016, United States

[8] Institute of Systems Genetics, NYU Langone Health, New York, NY 10016, United States

[9] Huntsman Cancer Institute, University of Utah Health Sciences Center, Salt Lake City, UT

84112, USA; Department of Surgery, University of Utah Health Sciences Center, Salt Lake City,

UT 84112, USA

[10] Herbert Irving Comprehensive Cancer Center, Columbia University, New York, NY 10032,

United States

[11] University of Zurich, Switzerland

[12] Department of Laboratory Medicine, Karolinska Institute, 141 86 Stockholm, Sweden

 # these authors contributed equally

*corresponding author and lead contact: Eva Hernando (eva.hernando-monge@nyulangone.org)

*corresponding author: Douglas Biancur (Douglas.Biancur@nyulangone.org)

Abstract

Brain metastatic cells undergo metabolic adaptations, such as increased reliance on

oxidative phosphorylation. Integrating proteomic and transcriptomic profiling of patient samples

revealed a consistent upregulation of the carnosine dipeptidase-1 (CNDP1) in brain versus

extracranial metastases. Carnosine is an abundant metabolite in brain and muscle, known to exert

anti-proliferative effects on cancer cells. Here, we demonstrate that CNDP1 inhibition suppresses

both the establishment and maintenance of melanoma brain metastasis while its ectopic

expression is sufficient to confer brain metastatic potential to poorly metastatic cells. CNDP1

suppression results in activation of the Integrated Stress Response via Heme-Regulated Inhibitor

Kinase and reprogrammed translation towards preferential expression of mitochondrial and

survival transcripts. We further show that CNDP1 upregulation supports mitochondrial activity by

limiting the levels of its substrate carnosine, a copper ionophore, thus protecting metastatic cells

from carnosine-induced copper toxicity. Our studies reveal a novel metabolic adaptation during

brain metastasis, which can be leveraged for therapeutic purposes.

Keywords: melanoma, brain metastasis, metabolic adaptation, mitochondria, metabolism,

carnosinase, carnosine, ionophore, copper.

43
Introduction

While advances in cancer therapy have drastically improved patient care, metastatic

disease is often refractory to current treatment options and remains the most common cause of

cancer-related deaths(1,2). Primary tumor cells may acquire mutations or transcriptional

programs that predispose cells to disseminate(3–5), and once they reach the metastatic niche,

they must undergo further site-specific adaptations to survive in the new microenvironment

(TME)(6–8). Understanding these adaptations is key to uncovering novel and targetable

vulnerabilities of metastatic cells.

Specifically, brain metastases (BMs) are a major cause of cancer-related morbidity and

mortality, affecting nearly 40% of patients with advanced solid tumors, most frequently in patients

with carcinomas of the lung or breast (9–11), and melanoma(12). Despite tremendous advances

in surgical approaches, radiotherapy, and systemic targeted and immune therapies, patients with

symptomatic BMs experience reduced treatment response rates, even when extracranial

metastases (ECMs) are controlled(13–15). Therefore, there is an urgent need of improved

treatment options tailored to the unique BM biology(16–20).

Extensive efforts have been made to identify mechanisms of brain metastasis tropism (21–

25) . For instance, studies have shown that metastatic cells rewire their metabolism to adapt to

the distinctive properties of the brain such as higher oxygen concentration, distinct micronutrient

availability, amino acid deprivation or alternative sources of carbon(22,23,26,27). Genes involved

in oxidative phosphorylation have been found to be upregulated at both mRNA and protein levels

in melanoma brain metastasis (MBM) patient samples compared to extracranial

metastases(22,26), and patients with an increased oxidative phosphorylation signature in their

MBMs exhibit decreased survival(21,26). We have previously shown that melanoma cells derived

from the brain metastatic niche hold a higher oxidative potential than their extracranial

counterparts(25). Moreover, metastatic cells undergo metabolic adaptations that enhance their

capacity to survive under oxidative stress(28,29), such as modulating the expression of NADPH+

generating enzymes. Similarly, cells that metastasize to the leptomeninges are able to compete

for available iron with macrophages in the cerebrospinal fluid (CSF) by modulating lipocalin-2

expression in the hypoxic niche to support their metabolic needs, among other cellular

processes(30). A better understanding of metabolic dependencies in the brain metastatic niche

may reveal novel susceptibilities that can be exploited for therapeutic purposes.

Here, we integrated two datasets: mass spectrometry-based proteomics of melanoma

extracranial (non-brain metastasis; NBM) and MBM Short Term Cultures (STCs)(25), and

transcriptomics data from extracranial and brain metastasis samples(26), reasoning that gene

products consistently upregulated in MBM across different cohorts and two omics modalities

might have functional relevance to the MBM cascade. This integration revealed that expression

of the brain-associated dipeptidase CNDP1 is consistently increased in MBM versus NBM.

CNDP1, or carnosine dipeptidase-1, is a secreted peptidase known to cleave carnosine into its

component amino acids L-histidine and ß-alanine(31–34). Carnosine exerts a protective

physiological function in muscle, brain, liver and kidney, and is reported to have anti-glycation

effects and impact oxidative stress, though the respective underlying mechanisms, as well as

those governing its abundance, remain unclear(35–38). It has been recently suggested that these

properties of carnosine come from its ability to chelate metals such as copper and iron, which can

contribute to the formation of Advanced Glycation End (AGE) products through reactive carbonyl

or reactive oxygen species (ROS)(37–42). Carnosine administration *in vitro* is known to exert an

anti-proliferative effect on cell lines of various cancer types(34,43–45). We therefore hypothesized

that CNDP1 upregulation in MBM may provide an adaptative advantage to cancer cells in the

brain where abundant carnosine exhibits a tissue-protective role. Indeed, our work uncovers a

new role for CNDP1 in overcoming metabolic stress during MBM establishment by cleaving

carnosine excess and protecting cells from intracellular copper overload. Our studies reveal a

novel molecular mechanism of BM adaptation, and CNDP1 as a druggable vulnerability which

can be exploited therapeutically to prevent or treat BM.

Results

Integrative multi-omics approach nominates CNDP1 as a driver of MBM

To uncover mechanisms of adaptation to the brain microenvironment shared across

patients, proteomics data from paired NBM and MBM short-term cultures (STCs) (n=3 unique

patients) were compared to an RNA-sequencing (RNA-seq) dataset of 35 surgically resected

MBM and 41 NBM tissue specimens (n=29 unique patients, Table S1(25,26)). We compared

differentially expressed genes in paired MBM vs NBM samples to those found by proteomics

profiling (Figure 1A). By applying a cutoff of log2FC > 0.4 in both studies, and a p-value threshold

of 0.005 in the differential transcriptomic analysis, we compiled a list of candidate genes

upregulated in both cohorts, with Carnosine dipeptidase-1 (CNDP1) having the highest

upregulation on both datasets (Figure 1A, Table S2). CNDP1 is a secreted zinc metalloprotease

that cleaves the dipeptide carnosine into L-histidine and ß-alanine(46). *CNDP1* is predominantly

expressed in the mammalian brain and liver, while its homologue *CNDP2* and its counterpart in

carnosine metabolism *CARNS1* (carnosine synthase 1) are ubiquitously expressed (Figure S1A).

Data mining of SKCM TCGA (The Cancer Genome Atlas) transcriptomics data revealed that

CNDP1 expression is significantly increased in metastatic (n=367) compared to primary samples

(n=104) (p=0.0054; Figure S1B). *CARNS1* levels were found increased and *CNDP2* levels

decreased during progression from primary to metastasis whereas CNDP1 increased in

metastatic (n=367) compared to primary samples (n=104) (p<0.006, p<0.001 and p<0.06,

respectively; Figure S1B). Neither *CNDP1* nor *CNDP2* transcript levels associate with melanoma

patients' overall survival in that dataset, while lower *CARNS1* correlates with shorter overall

survival in metastatic patients (p<0.002; Figure S1C). In a separate cohort, CNDP1 protein levels

within melanoma-infiltrated lymph nodes associate with shorter patient survival (n=59, p=0.03,

Figure 1B). Additionally, RNA-sequencing analysis of melanoma extended to resected lymph

nodes revealed higher expression of *CNDP1* in melanomas that metastasized to brain relative to

those that recurred to other sites (p=0.03, Figure 1C, Table S4). Single nucleus RNA-sequencing

(snRNA-seq) data from fresh-frozen patient samples(47) showed a higher percentage of MBM

cells expressing *CNDP1* compared to NBM (Figure 1D; p = $6.22e^{-16}$). Extending our findings

beyond melanoma, analysis of lung and breast cancer transcriptomics datasets showed *CNDP1*

upregulation in BM samples relative to primary tissues (p<0.001; Figure 1E-F, S1D).

Immunohistochemistry analysis of a tissue microarray (TMA) of melanoma patient samples

confirmed higher CNDP1 levels in brain metastasis samples (n=70) compared to patient-matched

NBM (n=24), primary tumors (n=43) or healthy skin and cortex (p=0.02, Figure 1G, Table S3). We

confirmed the expression of CNDP1 by qPCR, immunoblotting and carnosinase activity in patient-

matched STCs (Figure S1D-F). Altogether, these data nominated CNDP1 as a potential driver of

brain metastasis.

Next, we investigated potential transcriptional programs driving *CNDP1* upregulation in

melanoma metastasis. *In silico* analysis of TCGA ATAC-seq data revealed open chromatin

regions in the *CNDP1* locus and identified an enhancer, the opening of which strongly correlates

with *CNDP1* expression (Fig. S1H-I). It has been reported that levels of *MITF*, the master regulator

of melanocyte expression (48), are significantly higher in BM versus NBM (47) , suggesting

CNDP1 transcription could be driven by MITF. ChIPseq confirmed direct MITF binding to the

CNDP1 promoter and enhancer (Fig. S1J), and MITF silencing resulted in decreased H3K27ac

at the *CNDP1* promoter and reduced *CNDP1* mRNA levels (Figure S1J-L). Together, our data

suggests that *CNDP1* expression may be at least partially governed by MITF.

CNDP1 depletion and carnosine accumulation decrease *in vitro* proliferation of MBM

patient-derived short-term cultures

To test the potential role of CNDP1 in MBM, we used a loss-of-function approach. CNDP1

silencing with two doxycycline (dox)-inducible shRNAs in two human MBM STCs (Figure 2A)

slowed cell proliferation after 24 h by G1 phase stalling ($p<0.001$, Figure 2B-C), suggesting that

CNDP1 is required for 2D cell growth in MBM cells. Importantly, 24 hours of dox-induced CNDP1

KD resulted in an increase in carnosine extracellular levels measured by ELISA (Figure S2A). We

then confirmed that carnosine supplementation at comparable concentrations to those found

physiologically in brain (49), leads to a proliferation defect in the same cells and increased

intracellular carnosine levels ($p<0.001$, Figure S2B-C). Additionally, CNDP1 dependency is not a

general feature of melanoma cells, as suggested by a proliferation assay in a non-brain tropic

melanoma cell line (Figure S2D). We confirmed the anti-proliferative effect of CNDP1 depletion

using RNA interference with pooled siRNA as an orthogonal approach ($p<0.001$; Figure 2D-E).

Taken together, these data suggest cellular reprogramming to the brain microenvironment

involves CNDP1 induction, exposing a novel and potentially targetable vulnerability of brain

metastases.

CNDP1 inhibition suppresses brain metastasis establishment and maintenance in

xenografts, and Cndp1 ectopic expression confers brain tropism to syngeneic models

To test if CNDP1 is required for BM establishment and/or growth *in vivo*, NSG mice were

injected intracardially with patient-derived 12-273 BM cells stably transduced with either CNDP1-

targeting or non-targeting control shRNA (Figure 3A-H). *In vivo* experiments were carried out

using two approaches: i) constitutive CNDP1 knockdown, in which dox-containing chow was

administered to mice right before tumor cell injection (Figure 3A-D), which addresses the

requirement of CNDP1 for the *establishment* of metastasis, and ii) a more therapeutically relevant

approach in which dox was administered after metastases were seeded, which assesses if

CNDP1 is required for metastasis *maintenance* and growth (Figure 3E-H). Mice were humanely

euthanized when any group started to show symptoms of distress or more than 20% weight loss,

and organs were collected and processed for histological analyses. To quantify metastasis

burden, we used both whole body bioluminescence live imaging (BLI) and MetFinder(50), a

machine learning-powered tool we developed to segment, measure and quantify metastases in

H&E-stained sections of murine tissues at termination. Our data shows strongly reduced brain

and liver metastatic burden upon CNDP1 silencing with both shRNA ($p < 0.03$, n.s. for shCNDP1-

1 in the inducible approach in liver; Fig. 3D,H) A similar phenotype was observed in another

patient-derived model, 10-230 BM, and in a brain-tropic breast cancer model (MDA-231 Brm2),

after either constitutive (Fig S3A-D) or inducible CNDP1 knock-down (Fig S3E-L), supporting a

role for this dipeptidase across cancer types. RNA-seq of GFP positive cells sorted from

metastatic lesions confirmed that *CNDP1* silencing was maintained *in vivo* (Fig. S3M).

Next, we tested whether CNDP1 ectopic expression is sufficient to enhance BM

development. Murine Yumm3.2 melanoma cells (Braf$^{V600E/wt}$ Cdkn2a-/- Pten-/-(51–53)), which do

not metastasize when injected subcutaneously in C57BL/6J mice, were stably transduced with

lentiviral vectors carrying HA-tagged Cndp1 or Akt1-E17K, which was previously shown to

enhance metastasis development in this model(52) (Figure 3I). Cndp1 expression is not detected

in these murine melanoma cells (not shown). Ectopically expressed Cndp1 exhibits carnosinase

activity and confers a proliferation advantage *in vitro* (p<0.01, Figure S2E-G). Control, Akt1-E17K-

and Cndp1-HA-transduced Yumm3.2 cells were subcutaneously injected in C57BL/6 mice. Upon

reaching the maximum tumor volume, mice were euthanized, and organs harvested and

examined by ex-vivo BLI for metastasis burden (Figure 3J-K). We observed that Cndp1-HA

expressing cells gained exclusive brain tropism, with a higher incidence than Akt1-E17K

transduced cells (62% vs 0 % (parental) vs 40% (Akt-E17K), p<0.001), which were also able to

form lung and liver metastasis (Figure 3J-K). All these data points to a specific role for CNDP1 in

enhancing BM development.

Our results support a role for CNDP1 in brain metastasis and expose CNDP1 targeting as

a promising therapeutic intervention against these tumors. Interestingly, the observation that

single nucleotide polymorphisms (SNPs) impairing CNDP1 secretion protect diabetic patients

from developing nephropathy (36,54) has sparked interest in developing CNDP1 inhibitors, some

of which have shown promising activity in preclinical models(55). Our studies provide a rationale

for further developing small molecule compounds or antibodies targeting CNDP1.

CNDP1 knockdown elicits an Integrated Stress Response and translational

reprogramming

To further characterize the role of CNDP1 to tumor progression, we employed an unbiased

approach integrating LC/MS metabolomics and bulk RNA-seq of shCNDP1-transduced cells

induced with doxycycline for 72h (Figure 4A, Tables S5,6). We analyzed the resulting data with

the MetaboAnalyst algorithm(56) (Table S7), which uses hypergeometric tests to identify

enrichment of both transcript-based and mass spectra-based data in metabolic pathways. We

found that "Aminoacyl tRNA-synthesis" was the most deregulated metabolic pathway in tumor

cells upon CNDP1 suppression (Figure 4B, Table S7). To examine if these changes in tRNA

synthesis may have implications for translation, we performed additional MS/MS (proteomics)

analysis of 10-230 BM cells transduced with indicated shRNA (Table S8) and integrated these

data with LC/MS (metabolomics) data (Table S5). This analysis revealed an accumulation of

amino acids such as phenyl-alanine, arginine, tyrosine and leucine together with their respective

aminoacyl tRNA-synthetases (Figure 4C) upon CNDP1 silencing.

Further elucidating the cellular adaptative response triggered by CNDP1 silencing,

transcriptomic and proteomics profiling revealed changes in translation initiation, elongation and

RNA metabolism along with stress response pathways (Figures 4D, S4A, Tables S6,S9-10).

Together, these changes were reminiscent of activation of the Integrated Stress Response (ISR),

a cellular mechanism that halts or rewires protein synthesis in response to various sources of

stress.(57) Indeed, we observed an increase in expression of ATF4 targets (Figure 4E,F) and p-

Ser51-eIF2α levels (Figures 4G, S4B), both hallmarks of ISR activation, upon CNDP1 depletion.

Moreover, a p-eIF2α inhibitor, ISRIB,(58) was able to partially rescue the proliferation defect

induced by CNDP1 knock-down (p<0.02) (Figure 4H).

Although activation of the ISR often leads to a general shutdown of translation, CNDP1

silencing only had a modest impact on overall translation, in comparison to treatment with

cycloheximide (CHX), a *bona fide* translation inhibitor(59) (Figure 5A). CNDP1 KD cells showed

decreased mTOR activity, as revealed by p70 pS6K1, p-4E-BP1 and p-eIF4E levels, key

modulators of translation initiation(60,61) (Figure 5B). mTOR modulation has been previously

observed in response to TCA perturbation, such as a decrease in α-Ketoglutarate or changes in

branched chain amino acid (BCAA) metabolism including isoleucine, leucine and valine, as

observed upon CNDP1 KD (Figure 4B,C, Tables S5(62,63)). Additionally, transcriptomic data

analysis of human melanoma metastases (i.e., TCGA, including various metastatic sites;

GSE185386, including only MBM), revealed that *CNDP1* expression anti-correlates with

translation and stress response gene signatures (Figure 5C). Our data pointed to a selective

reprogramming of translation initiation rather than a complete shut down upon CNDP1 KD. To

better understand specific translation changes elicited by CNDP1 KD, we performed polysome

isolation followed by RNA-seq. Polysome profiling revealed changes in translation of mRNAs

directly involved in mitochondria, ribosomes and cell division genes (Figure 5D-E, Figure S5A-B,

Table S11). Concomitantly, we observed changes in eIF3d expression, suggestive of a shift from

canonical to non-canonical cap-dependent translation possibly mediated by eIF3d/DAP5 (Figure

S5C). It has been shown that eIF3d promotes selective cap-dependent translation of mRNAs

involved in cell survival, such as mitochondrial genes (64,65). Of note, human BM cells expressing

CNDP1^{high} levels are significantly enriched in mitochondrial genes in comparison to CNDP1^{low}

cells (Table S12). Altogether, these data suggest that CNDP1 silencing initiates the ISR which

reprograms the translation machinery to support mitochondrial respiration and cell cycle related

transcripts.

CNDP1 depletion perturbs mitochondrial activity and results in increased intracellular

copper levels

We next sought to understand how CNDP1 KD triggers mitochondrial stress, resulting in

activation of the ISR and translation rewiring. Given our data indicating a shift towards translation

of mitochondrial transcripts, we focused our attention on the mitochondria. Indeed, we observed

profound morphological changes in the mitochondria of CNDP1 KD cells, including a smaller

number of cristae per mitochondria (Figure 6A-E, Figure S6A-E), a phenotype previously reported

as functionally defective(66). *In vivo*, we observed reduced expression of the mitochondrial

transport protein Tom20 in brain metastases upon acute CNDP1 KD (Figure 6F), which is

indicative of dysfunctional mitochondrial structures (67). Taken together, these data raised the

possibility that carnosine accumulation from CNDP1 inhibition causes mitochondrial dysfunction.

Consistent with this hypothesis, both CNDP1 KD and carnosine treatment elicited a significant

decrease in oxygen consumption rate (OCR) measured by Seahorse assays (Figure 6G, Figure

S6F, $p<0.02$).

These findings led us to investigate if CNDP1 modulation also impacts Gln usage in the

TCA cycle of melanoma cells. We performed Gln tracing experiments by using U-^{13}C5 labelled

Gln and traced its incorporation by GC/MS analysis(68) in melanoma brain metastasis derived

cells transduced with shRNA targeting CNDP1 as indicated (Figure 6H). We found that after 24h

of silencing, CNDP1 KD cells display significantly reduced glutamine usage as evidenced by the

lower levels of its glutamine-derived metabolites in the tricarboxylic acid (TCA) cycle: α-

Ketoglutarate (M+5), Fumarate, Malate and Citrate (M+4) (Figure 6I). All these effects were

mimicked by carnosine addition (Figure S6G-H). These data demonstrate that both CNDP1 KD

and carnosine treatment impact TCA cycle activity. In addition to mitochondrial dysfunction,

CNDP1 KD cells accumulated ROS, as indicated by CellROX measurements (Fig. 6J).

Concomitant with ROS accumulation we observed the activation of heme-regulated inhibitor (HRI)

kinase (Figure 6K), likely responsible for the above mentioned ISR activation (Figure 4).

Having observed that CNDP1 silencing impairs mitochondrial respiration and results in

ROS accumulation, translation stalling and reprogramming, we looked for potential mechanisms

directly triggering these effects downstream of CNDP1. Carnosine has been reported to work both

as an ion chelator and ionophore, that is, able to transport iron and copper into the intracellular

space.(69) We hypothesized that carnosine accumulation upon CNDP1 KD could lead to

increased intracellular copper uptake through SLC transporters(70) and result in copper overload.

Intriguingly, we observed CNDP1 KD cells exhibit dysregulation of genes involved in heavy metal

metabolism such as iron (*LCN2, HMOX*) and copper transporters (*SLC46A3, ATP7B*) (Figure S6I-

K). Indeed, by analyzing scRNA-seq data of MBM,(22) we found a significant enrichment in genes

controlling copper homeostasis in CNDP1^{high} vs CNDP1^{low} MBM cells (Figure S6L). By staining

cells with a permeable fluorescent dye (CS1) that binds free intracellular copper, we measured

copper levels in CNDP1 KD and carnosine-treated cells along with cells treated with an

established copper ionophore, Elesclomol(71), as a positive control. Indeed, both CNDP1 KD and

carnosine-treated cells contain significantly higher intracellular copper levels, comparable to

those of cells treated with Elesclomol ($p < 0.05$ or indicated, Figure 6L). Copper mediated toxicity

also affects the lipoylation of TCA enzymes such as Pyruvate dehydrogenase (PDH) and 2-

oxoglutarate dehydrogenase (OGDH)(72), that results in degradation of these enzymes (Figure

S6M), and Fe-S clusters assembly(73), leading to a decrease of both glutamine usage and

OXPHOS dependent metabolism. Additionally, copper overload has been described to cause

ROS accumulation(74). Therefore, intracellular copper overload in CNDP1 KD and carnosine-

treated cells could explain the previously observed phenotypes of mitochondrial dysfunction and

accumulation of ROS (Figure 6G-J, S6F,G).

Altogether, our studies support CNDP1 upregulation as a metabolic adaptation of cancer

cells that increases their fitness during BM, likely by overcoming copper toxicity induced by

carnosine accumulation in the brain metastatic niche (see model in **Figure S7**). CNDP1 inhibition

emerges as a plausible strategy against metastatic progression by inducing metabolic stress.

Discussion

Understanding distinct biological features of site-specific metastatic adaptation offers the

possibility of discovering new targets and therapeutic options. In this study, we integrate two data

sets, RNA-seq and proteomics,(25,26) comparing BM versus extracranial tumor samples to

identify potential genes relevant to BM establishment and development. This analysis nominated

a secreted dipeptidase, carnosine dipeptidase 1 or CNDP1, as a plausible player in melanoma

brain metastasis. We also found *CNDP1* upregulated in lung and breast cancer brain metastases

versus primary tumors across multiple databases. Silencing CNDP1 slows proliferation *in vitro* of

brain metastasis-derived short-term cultures and impairs BM establishment and maintenance. A

defect in liver metastasis upon CNDP1 silencing was also observed in our brain-tropic xenograft

models; not surprisingly, since intracardiac injection favors hematogenous spread which allows

our brain metastasis derived cells to yield not only brain but also liver metastases[25,116]. Moreover,

carnosine is abundant in the liver[3], thus CNDP1 upregulation may also provide a growth

advantage there. However, Cndp1 ectopic expression exclusively conferred brain tropism to non-

metastatic syngeneic models, supporting CNDP1 as a brain metastasis driver. In all, we

demonstrate that high CNDP1 levels confer cancer cells increased ability to seed and colonize

the brain. While CNDP1 target, carnosine, has been studied for its various chemical properties,

the contribution of CNDP1 to human disease and in particular cancer is understudied. A mutant

form of this dipeptidase was found associated with less severe nephropathy in type II diabetes

patients(36,54); however, our study is the first to demonstrate the contribution of CNDP1 to brain

metastasis.

Whether CNDP1 expression is induced once cells arrive to the brain, or if CNDP1[high] cells

have increased ability to seed in the brain parenchyma is unclear. In support of the latter, we

found higher *CNDP1* levels in lymph nodes of melanoma patients that went on to develop brain

metastases than in those that did not have BM. In melanoma, recent studies have shown cells

can adopt different cell states, characterized by distinct transcriptional programs associated with

varying metastatic potential(75,76). We did not find the expression of CNDP1 to be associated

with specific transcriptional states (data not shown), yet we observed a plausible regulation of

CNDP1 by MITF. Indeed, in Biermann et al., we reported an enrichment of the MITF signature in

MBM vs extracranial metastasis.(22) However, our preliminary data suggest *CNDP1* upregulation

in BM across non-melanoma tumors (i.e., breast and lung adenocarcinoma BM), likely by MITF-

independent pathways. Further studies will be necessary to dissect CNDP1 regulation, and

whether CNDP1[high] cells can be detected in primary tumors and perhaps even hold prognostic

value.

In this study, we propose that melanoma BM cells overcome carnosine-mediated copper

overload in the metastatic niche by upregulating CNDP1. Copper uptake has been reported to

enhance MAPK signaling in BRAFV600E mutant cells, and trigger MAPK inhibitor (MAPKi)

resistance in melanoma cells(77). It also known how OXPHOS activation depends on MAPK

signaling(78,79) . We therefore hypothesize that melanoma cells that resist copper overload may

be able to not only escape from MAPKi treatment, but also to survive in the BM niche. By

DepMap(80) data mining we found that MITF[high] melanoma cell lines are more resistant to

Elesclomol treatment (Figure S6N). Moreover, melanocytic-like (MITF[high]) cells may exhibit more

tropism for BM(22) because of their well-known copper dependency, which these cells utilize for

tyrosinase activity and melanin production(81–83) .

Our results reveal a detrimental effect of CNDP1 silencing in the establishment and

maintenance of metastasis *in vivo* along with a mild effect on proliferation *in vitro* in our brain-

derived models*.* The main reported substrate of CNDP1, carnosine, is known to be abundant in

brain (e.g., 3 mg / 100 gr of tissue in rat brain), where it is produced mainly by oligodendrocytes

and displays antioxidant, scavenging, pH buffering and anti-neuroinflammatory properties(38,41–

43) . Carnosine has been studied as a dietary supplement in its native form, as well as in forms

less easily cleaved by CNDP1, such as N-acetyl carnosine and zinc-L-Carnosine(34) . Small

randomized controlled trials have shown that carnosine supplementation can benefit the short-

term memory of healthy elderly patients, as well as those with Mild Cognitive Impairment(84,85).

Our studies support previous evidence of carnosine's ability to function as a copper ionophore(39)

as part of a number of mechanisms that regulate tightly the distribution and transport of this

micronutrient. These findings open a new avenue in the study of carnosine in copper-dependent

mechanisms such as hepatic function or dopamine production and their potential effects in other

diseases(81,82,86). For instance, highly abundant oligodendrocytes in the substantia

nigra(87,88), that express both CARNS1 and CNDP1(49), may contribute to control carnosine-

mediated copper uptake by dopaminergic neurons used for dopamine production. Meanwhile,

carnosine supplementation has been reported to exert anti-proliferative effects on tumor cells(44).

In highly glycolytic tumors, it has been reported that Isoform 2 of CARNS1 (CARNS2) controls

carnosine levels, which have immunomodulatory effects during hepatic cancer progression(89).

We hypothesized that carnosine accumulation could be detrimental to tumor growth, and that

metastatic cells may benefit from cleaving carnosine and mitigating its antitumoral effects. The

regulation of carnosine levels in tissues is an open question; for instance, humans are known to

have higher levels of circulating CNDP1 in serum than other mammals(31,33). In fact, higher apes

and the Syrian hamster have evolved an export signal sequence at the *CNDP1* locus, absent

from other species, for unclear reasons(32,34). Carnostatine is a structural analog of carnosine

that has been described to inhibit CNDP1 dipeptidase activity in preclinical models of diabetic

nephropathy[52]. Future *in vivo* studies could test the therapeutic potential of pharmacological

inhibitors of CNDP1 able to cross the blood brain barrier, alone or in combination with approved

therapies.

CNDP1 is also known to cleave homocarnosine, a dipeptide made of GABA and L-

histidine that exists at higher concentrations in the human than in the mouse brain(37,90,91).

GABA is an inhibitory neuromodulator mainly produced in the cytosol of GABAergic neurons that

regulates the activity of astrocytes and oligodendrocytes(92–94). We and others have reported

that brain metastatic tumors can modulate the activity of neural cells in the metastatic niche by

various mechanisms(25,95). In addition, GABA release from melanocytes has been shown to

promote melanoma initiation by influencing their crosstalk with keratinocytes(96). Future studies

could determine if increased GABA −from CNDP1-mediated homocarnosine cleavage − allows

metastatic cells to hijack the neural microenvironment to their advantage.

To our knowledge, we are the first to report that CNDP1 silencing elicits translation

reprogramming by modulating two interconnected pathways, ISR and mTOR, known to respond

to various sources of cellular stress(61,97–100). ISR activation and mTOR inhibition upon CNDP1

KD result in preferred translation of mitochondrial and survival proteins, most likely controlled by

eIF3d cap-dependent translation mechanisms(64,65), as previously shown under other forms of

cellular stress. Extensive efforts have tried to determine if ISR activation is a barrier or conduit for

tumor progression(29,99,101,102). ISR activation in response to CNDP1 KD leads to changes in

the total AA pool, accumulation of tRNAs, and preferential translation of survival factors without

inducing cell death. Genetic inhibition of CNDP1 significantly reduces brain metastasis burden;

however, ISR activation upon CNDP1 KD could eventually allow cell survival in the metastatic

niche. These adapted cells might be able to persist, acquiring additional changes that may permit

tumor recurrence.

The shift of the translation machinery towards mitochondria-related transcripts paralleling

copper accumulation upon CNDP1 KD particularly captured our attention. Whereas iron

homeostasis is tightly post-transcriptionally regulated by Iron Response Elements (IREs)(103)

and Iron Regulatory Proteins (IRPs)(103,104), similar mechanisms regulating copper metabolism

have not yet been identified in human models(105–107). As discussed above, mitochondrial

metabolism has been shown to play a key role in the establishment of brain metastasis (22,25,26).

Proper oxidative phosphorylation relies on various factors,(21,22,26) including metal transport, a

key limiting process that links mitochondrial function and ROS homeostasis(108). Recent studies

have shown that cancer cells develop different adaptative mechanisms aimed at collecting heavy

metals from the metastatic niche to support their metabolic function(109,110). Their reliance on

heavy metals and particularly copper uptake becomes a vulnerability for brain metastatic cells.

As such, upon CNDP1 silencing, accumulation of the ionophore carnosine (39,40) may induce

copper toxicity(72,74,111) and mitochondrial malfunction. Supporting this model, CNDP1

depletion recapitulates the reported phenotypic effects of copper overload including ROS

accumulation(40,74,112),   reduced   OCR(73)   and   degradation   of   lipoylated   TCA

enzymes(72,111).

Our work exposes the relevance of copper transport and uncovers a mechanism

underlying oxidative phosphorylation reliance as a key metabolic adaptation to the brain

metastatic niche. We found the dipeptidase CNDP1 tipping the balance of complex, interrelated

mechanisms of cell survival emerging under metabolically demanding circumstances. Future

studies will reveal how to further exploit these cell vulnerabilities to maximize the therapeutic

potential of CNDP1 pharmacological inhibitors, alone or in combination with approved therapies.

Methods

Patient samples

Patients' tumor samples from infiltrated lymph nodes were collected from melanoma

patients who were prospectively enrolled in the Interdisciplinary Melanoma Cooperative Group

Database (IMCG) at NYU Langone Health (IRB #10362). Inclusion criteria for this

clinicopathological database included patients who presented to the Laura and Isaac Perlmutter

Cancer Center (2002-2018) for melanoma and were accrued within two months of diagnosis. All

patients signed NYU Langone Health Institutional Review board-approved consent to allow using

their specimens for research and to conduct follow-up.

Tissue microarray samples were acquired by agreement with Dr. Levesque lab from

University of Zurich. All patients included in this study have signed a patient release form, which

has been approved by an ethics committee and assigned the numbers EK647 and EK800.

Data mining

TCGA. Data from The Cancer Genome Atlas Skin Cutaneous Melanoma Firehose Legacy

dataset (n = 472 samples, The Cancer Genome Atlas (RRID:SCR_003193) was analyzed using

cBioPortal (RRID:SCR_014555), (113)), which reports RNA-sequencing data in RSEM. Survival

analysis based on gene expression in the TCGA cohort was carried out using TIMER

(RRID:SCR_018737, (114)), the Tumor Immune Estimation Resource, which draws Kaplan-Meier

curves based on a user-defined cutoff of expression by percentage compared to the median, and

reports p values based on log-rank testing. Primary and metastatic samples were analyzed

separately.

441  *Analysis of scRNA-seq data.* Integration of samples from *Biermann et al. (22)* was carried

442 out using Seurat's CCA function (RRID:SCR_007322) in R studio (R Project for Statistical

443 Computing (RRID:SCR_001905) and Bioconductor (RRID:SCR_006442). Tumor cells were

444 parsed from non-tumor cells: non-tumor cells were excluded based on expression of genes

445 unique to immune and stromal populations (using AddModuleScore and AUCell function

446 (RRID:SCR_021327)). Next, malignant cells were segregated in *CNDP1* high and low using SCT

447 (Single Cell Transform) (expression > 0 as threshold to plot a copper signature ((115)WP3286,

448 WikiPathways (RRID:SCR_002134)) in *CNDP1*-high vs -low cells. CNDP1 expression from Non-

449 Small Cell Lung Cancer primary and brain metastasis was obtained as described in *Tagore,*

450 *Caprio et al.* in press in Nature Medicine.

451  *Other databases.* Raw, normalized or analyzed data were downloaded from Gene

452 Expression Omnibus (GEO) (RRID:SCR_005012) following each study's directions and code

453 indicated in figure legends. Statistical analysis as described in figure legends.

454  *TCGA-ATACseq.* Normalized ATACseq counts from 370 TCGA patients were retrieved

455 from UCSC Xena Browser database (RRID:SCR_018938). 10kb BIN segments around the

456 *CNDP1* and *CNDP2* loci were visualized by heatmap using Prism10 (RRID:SCR_002798) and

457 averaged for the ATACseq counts. Next, ATACseq counts in 10kb BIN segments around *CNDP1*

458 promoters were averaged and plot against matched RSEM expression counts for Pearson's

459 correlation using Prism10 (RRID:SCR_002798).

460  *GTEx, Genotype-Tissue Expression*. We utilized Genotype-Tissue Expression

461 (RRID:SCR_013042) data to assess expression from various tissues to identify tissue-specific

462 and cross-tissue expression patterns of *CNDP1, CNDP2* and *CARNS*1. The data used for the

463 analyses described in this manuscript were obtained from the GTEx Portal on 08/11/2023.

464  *Depmap.* Gene expression data for *MITF* was obtained from the "DepMap Public 22Q2"

465 (Cancer Dependency Map Portal (RRID:SCR_017655)(80)) release, which includes RNA-seq

466 data (TPM, transcripts per million) for melanoma cell lines. *MITF* expression data was filtered to

retain melanoma-derived cell lines only by cross-referencing with metadata on cell line origin. For

each melanoma cell line, the area under the dose-response curve (AUC) was extracted as a

measure of Elesclomol sensitivity, with lower AUC values indicating higher sensitivity to the drug.

Correlation was calculated using Pearson's correlation coefficient.

Plasmid generation

CMV-Luciferase-EF1α-copGFP (GFP-Luc) Lentivector Plasmid was purchased from BD

Biosciences   (BLIV511PA-1).   Tet-pLKO-puro   was   purchased   from   Addgene

(RRID:Addgene_98398). shRNAs were cloned as previously described 115) into Tet- pLKO-puro

using AgeI (NEB, Cat#R3552S) and EcoRI (Thermo, Cat#FD0275) restriction sites. pLKO tet-on

scrambled   non-targeting   control   (shNTC)   was   purchased   from   Addgene

(RRID:Addgene_47541).

The FG12-HA-AKT1^{E17K} Luciferase-IRES-eGFP lentiviral vector has been described(52).

Cndp1-HA cDNA was synthesized by Gene Universal in the pENTR2B plasmid. Gateway cloning

was used to recombine the pENTR2B-Cndp1-HA entry clone and the pDEST FG12-Luciferase-

IRES-eGFP lentiviral destination vector (synthesized in Gene Universal) to generate the lentiviral

vector FG12-Cndp1-HA- Luciferase-IRES-eGFP.

siRNA pools and transfection

NON-targeting control (NTC) plus Human siRNA pools, ON-TARGET plus *CNDP1* and

MITF siRNAs were purchased from Dharmacon (QTE-3158440G, L-005824-00-0005, L-008674-

00-0005). 150,000 to 200,000 melanoma cells were seeded in 6-well plates and transfected with

20μM siRNA pools using Lipofectamine 2000 (Invitrogen,Cat#11668019) following

manufacturer's recommendations.

Lentivirus production and infection

HEK293T (RRID:CVCL_0063) cells at 80% confluency were co-transfected with 12 µg of lentiviral expression constructs, 8 µg of psPAX2 (RRID: Addgene_12260) and 4 µg pMD2.G (RRID: Addgene_ 12259) vectors using Lipofectamine 2000 (Invitrogen, Cat#11668019) following manufacturer's recommendations for Tet-pLKO-puro system. Packaging plasmids psPAX2 and pCMV-VSV-G (RRID: Addgene_8454) were used in combination with FG12-HA-AKT1-Ei7K Luciferase-IRES-eGFP (52,117) to generate virus for stable cell line generation as described below. At 48h and 72h post transfection, supernatants were collected, filtered (0.45 µm) and stored at −80°C. Melanoma cells were infected with lentiviral supernatants supplemented with polybrene (EMD Millipore, Cat#TR-1003-G) at a final concentration of 4µg/mL. Twenty-four hours after infection, cells were selected by cell sorting by GFP fluorescence for GFP-Luc transduced cells or treatment with puromycin (1 ug/mL, Thermo Fisher Scientific, Cat #A1113803), blasticidin (10 ug/mL, Invivogen, Cat#ant-bl-1) or hygromycin (300 ug/mL, Thermo, Cat#10687010) accordingly.

Cell culture

Melanoma short-term cultures (STCs) were derived as described and used at passage (25,51,118). For maintenance cultures, STC cells were grown in DMEM (Corning, Cat #10-013-CM) with 10% fetal bovine serum (regular: Corning, Cat#35-010-CV, Tet-free: Neuromics, Cat#FBS002-T), 1 mM Sodium Pyruvate (Cytiva, Cat#SH30239.01), 4 mM L- Glutamine (Sigma-Aldrich, Cat #59202C), 25 mM D-Glucose (Gibco, Cat#A2494001), 1% Nonessential Amino Acids (Sigma-Aldrich, Cat#M7145-100ML), 100 units/mL penicillin, and 100 ug/mL streptomycin (Gibco, Cat# 15-140-163). For functional assays, cells were seeded in DMEM (Corning, Cat #10-013-CM) with 10% dialyzed Tet-free fetal bovine serum (Neuromics, Cat#FBS002-T), 4 mM L- Glutamine (Sigma-Aldrich, Cat #59202C) and 25 mM D-Glucose (Gibco, Cat#A2494001).

HEK293T cells (RRID:CVCL_0063), MDA-231-Brm2 (RRID:CVCL_VR36) and SKMEL-239 (

RRID:CVCL_6122) melanoma cells(119) were grown in DMEM (Corning, Cat #10-013-CM)with

10% fetal bovine serum (FBS, Corning, Cat#35-010-CV), 1 mM Sodium Pyruvate (Cytiva,

Cat#SH30239.01), 4 mM L-Glutamine (Sigma-Aldrich, Cat #59202C), 25 mM D-Glucose (Gibco,

Cat#A2494001), 100 units/mL penicillin, and 100 ug/mL streptomycin(Gibco, Cat# 15-140-163).

YUMM3.2 cells were grown in F12/DMEM media (Thermo Fisher) supplemented with 10% FBS, 1

µg/mL penicillin-streptomycin (Thermo Fisher), and 1 µg/mL NEAA (Thermo Fisher). Cell lines

were maintained in a 5% CO2 incubator at 37°C and were routinely tested for Mycoplasma

contamination by using ATCC Kit (30-1012K) following manufacturer instructions.

In vitro proliferation assay

3,500 to 4,000 cells per well were seeded in 96 wells and monitored for 72 to 96 hours.

Cells transduced with Tet-pLKO-puro shRNAs were treated with doxycycline 1 ug/mL (Goldbio,

Cat #24390-14-5) for at least 48 hours before seeding or with indicated treatment.

Incucyte. After allowing cells to adhere overnight, Incucyte Software V2022.1, 2022.2, and

2023.1 (Sartorius IncuCyte SX5 Live Cell Analysis System (RRID:SCR_026298)), were used to

capture brightfield whole-well images. Masks for delineating confluence were manually adjusted

to ensure only live cells were captured, such as a minimum of 100 μM size per object. Images

were taken and analyzed every 6-12 hrs. Percent confluency was normalized to time=0 reading

to obtain a fold change value.

Crystal violet staining. After allowing cells to adhere overnight, a baseline plate was

obtained by removal of media and fixation in 0.1% glutaraldehyde for 15 minutes, then stored with

PBS(Corning, Cat #21-031-CM) in each well at 4C. Remaining plates were fixed every 24 hours

thereafter and/or at the experimental endpoint. Wells were stained with 0.5% crystal violet in PBS

for one hour and washed extensively with water. Crystal violet retained by cells was then eluted

by incubation in 250 μL 15% acetic acid for 1 hour with shaking, and absorbance at 590 nm was

measured. Absorbances were normalized to the absorbances from the baseline plate to obtain a

fold change value.

Cell cycle analysis

Cell cycle analysis was carried out using the Click iT EdU kit (Thermo Fischer, Cat

#C10636) to approximate DNA replication, and FxCycle Far Red (Thermo Fischer, Cat #F10348)

to measure total DNA content. Briefly, STC cells expressing control or CNDP1-targeting Tet-On

shRNA's were plated at 250,000 cells per well in 6-well plates with 1 ug/mL doxycycline. An

additional dose of doxycycline was added at 48 h post plating. At 72h post-plating, cells were

incubated with 10 μM EdU for 4 h. Then, cells were collected by trypsinization, washed in 3 mL

3% BSA in PBS, spun at 1200 rpm for 4 mins, fixed in 250μL Fixation buffer (Biolegend, Cat#

420801) for 20 mins in the dark at room temperature, washed in 3 mL 3% BSA, spun again.

Permeabilization and Click-iT reaction were executed according to kit instructions, followed by

FxCycle Far Red staining and analysis on BD LSR II Flow Cytometer.

In vitro treatments

Crystalline >99% L-carnosine (Sigma-Aldrich, Cat #C9625-25G) was dissolved into Media

or PBS (see Cell Culture) at a concentration of 200 mM by rocking in a light-protected 50 mL

conical tube. Solutions were pH adjusted to match base Media by adding HCl dropwise (~pH

7.45), then filtered using a 0.2 μm filter. Stocks were light-protected and maintained at -20C until

use, when they were thawed at room temperature and diluted into media. Cells were treated in a

10 μM to 50 mM range. ISRIB (phospho-eIF2a inhibitor, Selleck Chemicals, Cat#S7400) was

diluted in DMSO at 1mM stock solution and added *in vitro* at 1 μM final concentration for 72 hours.

Elesclomol (Selleck Chemicals, Cat#S1052) was diluted in DMSO 10 mM and added *in vitro* in a

range of 10 nM to 50 μM for 24 hours.

L-Histidine assay

Carnosinase activity was analyzed by measuring liberated L-histidine using a protocol

adapted from Teufel et al 2003(33). Briefly, conditioned media was collected, centrifuged to

remove dead cells and debris, and supernatant isolated. Adherent cells were rinsed in PBS and

scraped, and pellets were lysed in 50 mM Tris pH 8 with protease inhibitor (Roche,

Cat#11836145001) using a Bioruptor sonicator for 5 rounds of 1 min with 1 min off in between

each cycle. Lysates were centrifuged at 20,000 rpm for 15 mins at 4C and supernatants collected.

Lysates or conditioned media were diluted 1:1 (150 uL: 150 uL) in 50 mM Tris. 50μL of 20mM

MnCl2 was added to each tube, and samples were incubated for 30 mins at 37C with gentle

shaking. L-Histidine diluted in culture media was used as a positive control. 100μL of 50mM L-

Carnosine in Tris was then added to each sample, followed by incubation at 37C for 120 min with

gentle shaking. Finally, 500 μL of 10% trichloroacetic acid in 50 mM Tris was added to each

sample. Samples were centrifuged at 3000 rpm for 10 mins to remove precipitated proteins.

100μL of each supernatant was transferred to fresh 5 mL conical tubes. In a dark chemical hood,

2mL 0.3N NaOH was added to each sample, followed by 400 μL of freshly made 5 mg/mL o-

phthaldialdehyde solution (OPA) in 2 N NaOH. Solutions were allowed to incubate at room

temperature for exactly 3 mins in the dark before addition of 400 μL 6M HCl, at which point

samples were capped and shaken vigorously for 5s each. 100 μL of each sample was transferred

into a well of a black bottom 96 well plate. L-Histidine diluted in culture media was used as a

positive control. L-Histidine fluorescence was detected using excitation wavelength of 344 λ and

emission wavelength of 420 λ.

AHA integration translation assay

591   10-230 and 12-273 BM cells expressing control or CNDP1-targeting Tet-On shRNA's. 72h

after doxycycline treatment initiation, cells were labeled according to recommendations from the

Click-iT AHA labeling kit (Invitrogen Cat, #C10102). Briefly, cells were pre-incubated in

methionine-free, serum-free media (Fisher Scientific with repleted glutamine, Cat#21-012-026),

with or without doxycycline/carnostatine/carnosine depending on group condition, with or without

cycloheximide (Sigma-Aldrich, Cat #C7698-5G) for 30 mins. Then, cells were labeled with AHA

at a 50 μM final concentration for 4 hrs. Cells were trypsinized and permeabilized in 100 μL

saponin-based permeabilization buffer, washed in 1 mL 1% BSA in PBS, and incubated on 500

μL alkyne detection reaction cocktail with 5 μM Alexa Fluor 647 Alkyne for 30 mins at room

temperature. After washing with 1 mL 1% BSA in PBS, cells were resuspended in 100 μL 1%

BSA and stained with DAPI immediately prior to FACS analysis on a BD LSR II Flow Cytometer

using the APC and DAPI channels to gate for singlets. AHA incorporation in place of methionine

was quantified as the mean fluorescence intensity in the APC channel.

Quantitation of Reactive Oxygen Species

606   STC cells expressing control or CNDP1-targeting Tet-On shRNA's were plated at 250,000

cells per well in 6-well plates with 1 ug/mL doxycycline. An additional dose of doxycycline was

added at 48h post plating. At 72h post-plating, cells were incubated by replacing media with 1 mL

fresh serum-free media containing indicated doses of Tert-butyl hydrogen peroxide (TBHP,

positive control, ROS-stimulating agent, Thermo Scientific, Cat#180340050) or N-Acetyl Cysteine

(NAC, antioxidant, Sigma-Aldrich, Cat #180340050) for 2.5 hrs. Fresh resuspensions of THBP

and NAC were made on the day of each experiment in deionized sterile water. Then, CellRox

DeepRed reagent was added to each well (excluding negative controls) at a final concentration

of 500nM and allowed to incubate with cells for 30 mins at 37C protected from light. Media was

then collected, then adherent cells were rinsed with PBS, trypsinized, and added to their

respective conditioned media. After centrifugation at 1200 rpm for 4 minutes, cells were

resuspended in 100 μL 3% BSA and kept on ice until analysis on the BS LSRII Flow Cytometer.

DAPI was added immediately prior to analysis to detect dead cells. Cells were analyzed on the

cytometer in order of treatment group, not by shRNA expression group. Only live singlet cells

were compared, using the Mean Fluorescence Intensity of APC channel signal to approximate

ROS burden.

Copper detection

20K cells were seeded in black wall clear bottom 96 well plates treated for cell culture. All

conditions were treated with doxycycline and/or indicated Elesclomol and carnosine doses. After

24 treatment, Coppersensor-1 (CS1, MedChem Express, Cat#HY-141511) was utilized at 5 µM

following manufacturer indications.

RT-qPCR

RNA was isolated from cells at 80% confluence in 6 well plates using the RNeasy Mini Kit

(Qiagen, Cat #74104) with DNase I treatment. 1 ug of RNA was subjected to reverse transcription

following TaqMan™ Reverse Transcription Reagents manufacturer indications (Applied

Biosystems, Cat #4304134). Real time quantitative PCR was performed in 384 wells by mixing

Applied Biosystems™ Power SYBR™ Green PCR Master Mix (Applied Biosystems, Cat

#4367659), forward and reverse target gene primers at 0.2 nM final concentration (See Materials

Table) and 100 to 150 ng of cDNA up to 15 μl final volume.

Immunohistochemistry

Immunohistochemistry was performed using mouse anti-human CNDP1 Monoclonal

(OTI1A5, LSBio) and Tom20 (D8T4N) Rabbit mAb (Cell Signaling Technology) on formalin fixed,

paraffin embedded tissues. In brief, after deparaffinization and rehydration, heat-induced epitope

retrieval was performed in 0.01 M citrate buffer (pH 6.0) in a 1200-W microwave oven at 100%

power for 20 min. Sections were then cooled in tap water for 5 min, quenched in hydrogen

peroxide for 30 min, washed with PBS, and incubated with blocking serum for 30 min followed by

each primary antibody (Ab) diluted in buffer at room temperature for 1 h and at 4 °C overnight.

Abs used were Tom20 (D8T4N) Rabbit mAb (Cell Signaling Technology, Cat #42406,

RRID:AB_2687663) and CNDP1 Mouse anti-Human Monoclonal (OTI1A5) ab (LSBio, Cat#LS-

C172361, RRID: AB_3676497). Slides were washed in buffer and incubated with diluted

biotinylated for Tom20 (Sigma-Aldrich, Cat#A0545, RRID:AB_257896) or alkaline phosphatase

for CNDP1 secondary antibodies for 1 h (Alkaline Phosphatase Ab mouse, LSBio, LS-C44264,

RRID:AB_1050644). Avidin-biotinylated horseradish peroxidase complexes diluted at 1:500 were

added, and complexes were visualized with diaminobenzidine for Tom20 staining or with Fast

Red (Abcam, Cat# ab64254) for CNDP1 following manufacturer indications. Slides were then

washed in distilled water, counterstained with hematoxylin, dehydrated, and mounted with

permanent media. Appropriate positive and negative controls were included with the study

sections.

Western Blot Analysis

Cells were harvested in cold RIPA lysis buffer (Pierce, Cat # 89900) supplemented with

protease inhibitor and protein was quantified by DC assay. Cell lysates (30 ug of protein) were

resolved in 4%- 12% Bis-Tris gels (Thermo Fisher Scientific, Cat#NP0321BOX) and transferred

to PVDF membranes using wet transfer.

Membranes were blocked in either 5% non-fat milk or 5% BSA in Tris-buffered saline/1%

Tween 20 (TBST) for 1 hour. Membranes were incubated overnight with primary antibodies (Ab)

diluted in 5% milk or 5% BSA TBST. Primary ab used were anti-Human CNDP1 (352216) Ab

(Novus Biological, Cat#MAB2489-SP, RRID: AB_3676498), PhosphoeIF2α (Ser51) Ab (Cell

Signaling Technology Cat#9721S, RRID: AB_330951), EIF2S1 Monoclonal Ab (EIF2-alpha)

(Invitrogen, Cat#AHO0802, RRID: AB_2536316), Monoclonal Anti-Vinculin Ab produced in

mouse (Sigma-Aldrich, Cat#V4505-100UL. RRID: AB_477617), Atf 4 (D4B8) Rabbit mAb (Cell

Signaling Technology, Cat#11815S, RRID: AB_2616025), Phospho-p70 S6 Kinase (Thr389)

(108D2) Rabbit mAb (Cell Signaling Technology, Cat#9234T, RRID:AB_2269803), p70 S6

Kinase Ab (Cell Signaling Technology, Cat#9202, RRID: AB_AB_331676) 4E-BP1 (53H11)

Rabbit mAb (Cell Signaling Technology, Cat#9644T, RRID:AB_2799054), Phospho-4E-BP1

(Ser65) Ab (Cell Signaling Technology, Cat#9451T, RRID: AB_330947), Hsp90 Ab (Cell Signaling

Technology, Cat#4874S, RRID: AB_2121214), Rabbit HRI Polyclonal Ab (MyBioSource.com,

Cat#MBS2538144, RRID:AB_2885168), Monoclonal Anti-beta-Actin-Peroxidase Ab produced in

mouse, clone AC-15, purified from hybridoma cell culture (Sigma-Aldrich, Cat#A3854-200UL,

RRID:AB_262011, Rabbit anti-eIF3D/EIF3S7 Ab, Affinity Purified (Bethyl Laboratories,

Cat#A301-758A, RRID:AB_1210970, Eif4E Mouse, Unlabeled, Clone: 87, (BDCat#610270,

RRID:AB_397665), eIF4G Ab (Cell Signaling Technology, Cat#2498, RRID: AB_2096025), DAP-

5 Ab (F-2) (Santa Cruz Technology, Cat#sc-137011, RRID:AB_2095908 Pyruvate

Dehydrogenase (PDH) Ab (Cell Signaling Technology, Cat#2784, RRID: AB_2162928), 2-

oxoglutarate dehydrogenase (OGDH) Ab (Cell Signaling Technology, Cat#26865,

RRID:AB_2737585), Anti-HA-Tag Rabbit Monoclonal Ab (Cell Signaling Technology, Cat

#C29F4, RRID:AB_10693385). Membranes were washed 3 times for 10 min in TBST and

incubated in secondary antibody for 1 hour in 5% milk or BSA TBST. Membranes were washed

3 times for 10 minutes in TBST and incubated with ECL substrate for 1 min and exposed to film

or BioRad chemiluminescence dock station for imaging. Antibodies for phosphorylated proteins

and their respective antibodies for total protein were blotted in different membranes or

phosphorylated antibodies were blotted first, membrane was stripped and then the total was

blotted. Representative housekeeping shown.

The following secondary antibodies were used: Goat anti-Rabbit IgG-Peroxidase (Sigma-

Aldrich, Cat#A0545, RRID:AB_257896), Goat anti-mouse IgG kappa-light chain (Santa Cruz

Technology, Cat#sc-516102, RRID:AB_2687626) 1:10,000.

Quantification was performed by inverting the membrane raw image, measuring the

integrated density of each band using the same quantification area for all bands for the

housekeeping bands and the protein of interest. Quantification is represented as a ratio of the

signal of the protein of interest and the housekeeping protein signal and finally normalized to each

control condition.

Carnosine ELISA

Media (see Cell Culture above) was conditioned with melanoma cells for 24-72 hours (until

reaching ~80% confluency) and spun at 1200 rpm for 4 minutes to remove cells and debris.

Carnosine ELISA was performed following manufacturer's indications (Novus Biologicals,

Cat#NBP2-75013).

Metabolomics

LC-MS/MS with the hybrid metabolomics method. Samples were subjected to an LCMS

analysis to detect and quantify known peaks. A metabolite extraction was carried out on each

sample based on a previously described method(120). The LC column was a Millipore™ ZIC-

pHILIC (2.1 x 150 mm, 5 µm) coupled to a Dionex Ultimate 3000™ system and the column oven

temperature was set to 25 °C for the gradient elution. A flow rate of 100 µL/min was used with the

following buffers; A) 10 mM ammonium carbonate in water, pH 9.0, and B) neat acetonitrile. The

gradient profile was as follows; 80-20% B (0-30 min), 20-80% B (30-31 min), 80-80% B (31-42

min). Injection volume was set to 2 µL for all analyses (42 min total run time per injection).

MS analyses were carried out by coupling the LC system to a Thermo Q Exactive HF™

mass spectrometer operating in heated electrospray ionization mode (HESI). Method duration

was 30 min with a polarity switching data-dependent Top 5 method for both positive and negative

modes. Spray voltage for both positive and negative modes was 3.5 kV and capillary temperature

was set to 320 °C with a sheath gas rate of 35, aux gas of 10, and max spray current of 100 µA.

The full MS scan for both polarities utilized 120,000 resolution with an AGC target of 3e6 and a

maximum IT of 100 ms, and the scan range was from 67-1000 m/z. Tandem MS spectra for both

positive and negative mode used a resolution of 15,000, AGC target of 1e5, maximum IT of 50

ms, isolation window of 0.4 m/z, isolation offset of 0.1 m/z, fixed first mass of 50 m/z, and 3-way

multiplexed normalized collision energies (nCE) of 10, 35, 80. The minimum AGC target was 1e4

with an intensity threshold of $2e^5$. All data were acquired in profile mode.

The resulting ThermoTM RAW files were converted to SQLite format using an in-house

python script to enable downstream peak detection and quantification. The available MS/MS

spectra were first searched against the NIST17 MS/MS (121), METLIN (RRID:SCR_010500)

(122)and respective Decoy spectral library databases using an in-house data analysis python

script adapted from our previously described approach for metabolite identification false discovery

rate control (FDR) (123,124). Metabolite peaks were extracted based on the theoretical m/z of

the expected ion type, e.g., [M+H]+, with a 15 part-per-million (ppm) tolerance and a ± 0.2 min

peak apex retention time tolerance within an initial retention time search window of ±0.5 min. For

all the group-wise comparisons, t-tests were performed using the Python SciPy SciPy

(RRID:SCR_008058), (1.5.4) (125) library to test for differences and generate statistics for the

downstream analyses. For the pairwise t-tests, any metabolite with a p-value < 0.01 was

considered significantly regulated (up- or down-) for prioritization in the subsequent analyses.

Heatmaps were generated by hierarchical clustering, performed based on the imputed matrix

values utilizing R studio (R Project for Statistical Computing (RRID:SCR_001905) and

Bioconductor (RRID:SCR_006442), (1.0.12). GraphPad Prism 9 (9.4.1, GraphPad Software

GraphPad Prism (RRID:SCR_002798).

Data analysis. Metabolite peak intensities were extracted according to a library of m/z

values and retention times developed with authentic standards. Intensities were extracted with an

in-house script with a 10 ppm tolerance for the theoretical m/z of each metabolite, and a maximum

30 sec retention time window.

RNA-sequencing analysis

After 72 h induction of shRNA expression with doxycycline, cells were scraped and RNA

was harvested using the RNeasy Mini Kit (Qiagen, Cat #74104). For RNA-seq in cells from *in vivo*

experiments, tissues were harvested after intracardiac injection and 7 days of doxycycline

treatment. Tissues were homogenized by using a mechanic/enzymatic protocol (Collagenase

Sigma, Cat#C2674-100MG and DNAse I, EMD Millipore, 260913-10MU) and sorted for GFP

positive expression with adequate controls. RNA was extracted using the abovementioned kit.

RNA-Seq library preps were made using the Illumina TruSeq Stranded total RNA with RiboZero

Gold kit on a Beckman Biomek FX instrument, using 100 ng of total RNA as input, amplified by

12 cycles of PCR, and run on an Illumina NovaSeq 6000. Reads were aligned and gene

expression compared using the Seq-n-Slide pipeline (Seq-N-Slide (RRID:SCR_021752)). Briefly,

FastQC v0.11.7 was used to check fastq files for poor sequencing quality; all samples had high

quality. Illumina adapter sequences and poor-quality bases were then trimmed using trimmomatic

v0.36. Trimmed sequences were mapped to mm10 using STAR v2.6.0a (RRID:SCR_004463),

indexed using samtools v1.9, then quantified for UCSC genes using htseq-count

(RRID:SCR_011867) v0.11.1. Comparative analysis between conditions was performed using

DESeq2 (RRID:SCR_015687) v1.24.0 with default parameters.

Polysome Sequencing

Polysome isolation was performed by separation of ribosome-bound mRNAs via sucrose

gradient as previously described(126). Briefly, 14X89mm Beckman Ultra-Clear centrifuge tubes

were loaded with 5.5 ml of 50% and 5.5 ml 15% sucrose in Polysome Lysis Buffer (PLB) [20 mM

Tris (pH 7.4) in Nuclease Free H2O, 10 mM NaCl, and 3 mM MgCl2 supplemented with RiboLock

RNase inhibitor (ThermoFisher, Cat#EO0384) and 100µg/ml cycloheximide (Sigma-Aldrich, Cat

#C7698-5G) and incubated at 4 °C horizontally overnight. 10-230 BM and 12-273 BM expressing

shNTC and shCNDP1-1,2 were pre-treated with 100 µg/ml cycloheximide (Sigma-Aldrich, Cat

#C7698-5G) for 15 min at 37oC, washed twice in phosphate-buffered saline (PBS) containing

100 µg/ml CHX, pelleted, and flash frozen (10% was separated for total RNA and protein

analysis). Pellets were allowed to thaw on ice and resuspended in 750 µl of PLB. After 5 min of

incubation on ice, 250 µl of detergent buffer (1.2% Triton, and 0.2 M sucrose in polysome isolation

buffer) was added. Cells were further lysed for 10 min and lysate was clarified. Equal amount of

clarified lysate was layered onto 15–50% sucrose gradients and sedimented at 36 000 rpm for 2 h

in a SW41 rotor (Beckman Coulter) at 4 °C. Gradients were collected in 12 × 750 µl fractions by

pumping 60% sucrose into the bottom of the gradient and collecting from the top using an ISCO

fraction collector while simultaneously monitoring absorbance at 254 nm. RNA was isolated by

extraction using TRIZOL (ThermoFisher, Cat# 15596026) as per manufacturer's instructions.

Light (poorly translated) polysome fractions (2–3 ribosomes) and well-translated heavy polysome

fractions (≥4 ribosomes) were pooled. In parallel total RNA was extracted to normalize for

transcriptional changes. All experiments were performed in triplicate. RNA quality and amount

were determined by the Agilent Technologies kit and Nanodrop. RNAseq was carried out by the

NYU School of Medicine Genome Technology Core using the Illumina Hi-Seq 2500 Single Read.

Multiple triplicate samples were pooled for 10-230 BM. PCA analysis was performed using the

"stats" R package (Bioconductor (RRID:SCR_006442)) and multiple individual triplicates (12-273

BM shCNDP11-1 and shNTC-2 bulk) were identified as outliers and removed from further

processing.

Polysome data was analyzed using RIVET (RRID:SCR_026488, a R-Shiny-based

tool(126). Utilizing the matched bulk and light + heavy polysome data, RIVET uses the "limma" R

statistical package (RRID:SCR_010943) to determine transcripts that were over or under

expressed in both transcription and translation. Within the RIVET app, the transcripts for both

shRNAs targeting CNDP1 were combined to create a "combined" sample. The differentially

expressed genes identified by RIVET were exported for further analysis. Using the R package

"enrichplot," GSEA (Gene Set Enrichment Analysis (RRID:SCR_003199)) analyses were

performed on the differentially expressed gene lists to identify associated pathways possibly

affected by the changed genes.

Proteomics workflow

Protein isolation. 10-230BM cells expressing shNTC, shCNDP1-1, or shCNDP1-2 were

seeded at 1e6 cells per 15cm plate in triplicate and incubated in DMEM + 10% dialyzed serum +

1 ug/mL doxycycline for 72 hrs. STCs at ~80% confluence were scraped from 10 cm plates on

ice, washed once with cold PBS, and scraped in 1mL additional cold PBS into an eppendorf tube.

After centrifugation at 1200 rpm for 4 minutes, pellets were flash-frozen in liquid nitrogen. Upon

thawing, pellets were lysed in cold RIPA buffer supplemented with protease inhibitor, for 15

minutes with vortexing every 5 minutes. Protein concentration was determined using Micro BCA

Protein Assay Kit (Thermo Scientific, Cat #23235).

812   *Sample preparation for mass spectrometry analysis.* 150 μg of each protein lysate were

813 proteolytically digested and subjected to quantitative mass spectrometry on an Orbitrap Fusion

814 Lumos mass spectrometer using isobar tandem mass tags (TMT). Samples were prepared using

815 the filter-aided sample preparation (FASP) method. Briefly, each sample was reduced with DTT

816 (final concentration of 20 mM) at 57°C for 1 hour and loaded onto a MicroCon 30-kDa centrifugal

817 filter unit (Millipore) pre- equilibrated with 200 μl of FASP buffer [8 M urea and 0.1 M tris-HCl (pH

818 7.8)]. Following three washes with FASP buffer, lysates were alkylated on a filter with 50 mM

819 iodoacetamide for 45 min in the dark. Filter-bound lysates were washed three times each with

820 FASP buffer followed by 100 mM ammonium bicarbonate (pH 7.8). The samples were digested

821 overnight at room temperature with trypsin (Promega) at a 1:100 ratio of enzyme to protein.

822 Peptides were eluted twice with 100 μl of 0.5 M NaCl. The tryptic peptides were subsequently

823 desalted using an UltraMicro Spin Column, C18 (Harvard Apparatus) and concentrated in a

824 SpeedVac concentrator.

825   *TMT labeling.* The dried peptide mixture was re-suspended in 100 μl of 100 mM

826 Triethylammonium bicarbonate (TEAB) (pH 8.5). Each sample was labeled with TMT reagent

827 according to the manufacturer's protocol. In brief, each TMT reagent vial (0.8 mg) was dissolved

828 in 41 μL of anhydrous ethanol and was added to each sample. The reaction was allowed to

829 proceed for 60 min at room temperature and then quenched using 8 μL of 5% (w/v)

830 hydroxylamine. The samples were combined at a 1:1 ratio and the pooled sample was

831 subsequently desalted using strong-cation exchange and strong-anion exchange solid- phase

832 extraction columns (Strata, Phenomenex) as described.

833   *Global Proteome Analysis.* A 500 μg aliquot of the pooled sample was fractionated using

834 basic pH reverse-phase HPLC (as described) 250. Briefly, the sample was loaded onto a 4.6 mm

835 × 250 mm Xbridge C18 column (Waters, 3.5 μm bead size) using an Agilent 1260 Infinity Bio-inert

836 HPLC and separated over a 90 min linear gradient from 10 to 50% solvent B at a flow rate of 0.5

837 ml/min (Buffer A = 10 mM ammonium formate, pH 10.0; Buffer B = 90% ACN, 10 mM ammonium

formate, pH 10.0). A total of 120 fractions were collected and non-concatenated fractions

combined into 40 final fractions. The final fractions were concentrated in the SpeedVac and stored

at -80°C until further analysis.

LC-MS/MS analysis. An aliquot of each final fraction was loaded onto a trap column

(Acclaim® PepMap 100 pre-column, 75 µm × 2 cm, C18, 3, 100 Å, Thermo Scientific) connected

to an analytical column (EASY-Spray column, 50 m × 75 µm ID, PepMap RSLC C18, 2 µm, 100

Å, Thermo Scientific) using Easy nLC 1000 (Thermo Scientific) with solvent A consisting of 2% in

0.5% acetic acid and solvent B consisting of 80% acetonitrile in 0.5% acetic acid. The peptide

mixture was gradient eluted into the Orbitrap Lumos Fusion mass spectrometer (Thermo

Scientific) using the following gradient: 5%-23% solvent B in 100 min, 23% - 34% solvent B in 20

min, 34% - 56% solvent B in 10 min, followed by 56%- 100% solvent B for 20 min. Full scans

were acquired with a resolution of 60,000 (@ m/z 200), a target value of 4e5 and a maximum ion

time of 50 ms. After each full scan the most intense ions above 5E4 were selected for

fragmentation with HCD using the "Top Speed" algorithm. The MS/MS were acquired in the

Orbitrap with a resolution of 60,000 (@ m/z 200), isolation window of 1.5 m/z, target value of 1e5,

maximum ion time of 60 ms, normalized collision energy (NCE) of 35, and dynamic exclusion of

30 s.

Data analysis. The MS/MS spectra were searched against the UniProt human reference

proteome with the Andromeda 252 search engine integrated into the MaxQuant 253 environment

(version 1.5.2.8) using the following settings: oxidized methionine (M), TMT-labeled N- term and

lysine, acetylation (protein N-term) and deamidation (asparagine and glutamine) were selected

as variable modifications, and carbamidomethyl (C) as fixed modifications; precursor mass

tolerance was set to 10 ppm; fragment mass tolerance was set to 0.01 Th. The identifications

were filtered using a false-discovery rate (FDR) of 0.01 using a target-decoy approach at the

protein and peptide level. Only unique peptides were used for quantification and only proteins

with at least two unique peptides were reported. Data analysis was performed using Perseus.

Protein levels were median centered and log2 normalized. To identify proteins differentially

expressed between groups, a Welch's t-test was performed.

ChIP-seq

Around 1×10^7 501mel cells per IP were double crosslinked in 25mM EGS (Thermo Fisher

Scientific, 21565) and 1% formaldehyde (Sigma Aldrich, Cat#47608). Cells were lysed in ChIP

lysis buffer (CLB = 10mM Tris HCl pH8, 100mM NaCl, 1mMEDTA pH8, 0.5mM EGTA pH8, 0.1%

Sodium Deoxycholate, 0.5% N-lauroylsarcosine, 1% Triton X-100) and sonicated with Diagenode

Bioruptor for 15-30 cycles (30" ON +30" OFF max intensity). H3K27ac IP was conducted using

5ug of Ab/IP H3K27Ac Ab (Active Motif, Cat#91193, RRID:AB_2793797) against an IgG control

(Cell Signaling Technology, Cat#3900, RRID:AB_1550038) prebound to ProteinA/G magnetic

beads (Pierce Thermo Fisher Scientific, Cat#88803). After overnight IP, beads were washed

twice in Wash Buffer I (WBI, 20mM Tris HCl pH7.5, 150mM NaCl, 2mM EDTA pH8, 0.1% SDS,

1% Triton X100), Wash Buffer II (WBII, 10mM Tris HCl pH8, 250mM LiCl, 1mM EDTA, 1% Triton

X100, 0.7% Sodium Deoxycholate) and TET Buffer (10mM Trist HCl pH8, 1mM EDTA, 0.2%

Tween20) and eluted in Elution buffer (EB, 10mM Tris HCl pH8, 300mM NaCl, 5mM EDTA pH8,

0.5% SDS, 10ug RNAse A, 100ug proteinase K). ChIP elutions were purified using MinElute

Reaction Cleanup kit (Qiagen, Cat#28204) followed by library preparation using NEBNext Ultra II

DNA Library Prep Kit for Illumina (NEB, Cat# E7600S) following manufacture instructions.

Analysis. H3K27ac and MITF fastqs (MITF fastq obtained from GSE61967) were trimmed,

filtered and aligned to hg38 human genome using using Trim Galore (RRID:SCR_011847),

FastQC (RRID:SCR_014583) and Bowtie 2 (RRID:SCR_016368) (2.4.1). Sam output files were

converted into bam using SAMTOOLS (RRID:SCR_002105) (1.16124) and bigwig files were

generated using bamcoverage (deeptools 3.5.6125). H3K27ac and MITF peaks were called using

MACS (RRID:SCR_013291) (2.2.9.1126) using paired-end mode. IgG served as controls. During

all steps, genomic regions were filtered for blacklisted regions.

Seahorse Metabolic Analysis

30 to 50 K cells were plated on a XF96 Cell Culture Microplate. The Seahorse MitoStress test protocol was followed using a Seahorse XF96 instrument (Agilent). Concentrations of inhibitors injected were as follows: 2uM oligomycin, 500 nM FCCP and 1 uM antimycin A and rotenone (Cayman Chemical Company, Cat#11342, Cat#15218, Cat #19433, Cat#13995, respectively). After the MitoStress test, cells were fixed in the plated and stained using crystal violet and dissolved with Acetic Acid 15%. Absorbance was measured at 565l. Oxygen consumption rate (OCR) values for every condition were normalized to final confluence.

Glutamine tracing

For ^{13}C5-Gln tracing experiments cells were plated in a six-well plate at 2.0 × 105 cells per well and allowed to attach overnight in DMEM. Cells were treated as indicated in figure legends. Next, cells were washed with PBS twice and cultured for 30 mins in DMEM without phenol (Gibco, Cat#A1443001) supplemented with 4 mM 13C5-Gln and 10% dialyzed serum was added. The tracing medium was removed and washed with PBS. Polar metabolites were collected and derivatized using MOX-TBDMS (Thermo Scientific, Cat#TS-45950, Regis Technologies, Cat#77377-52-7; Cat#18162-48-6, respectively) and analyzed using a labeled amino acid standard as previously described 119–122.

Animal Studies

NOD/SCID/IL2yR- /- male mice (RRID:IMSR_JAX:005557) at 8-12 weeks were used for *in vivo* studies involving injection of 12-273BM STC. Male athymic nude mice

(RRID:IMSR_JAX:002019) were used for in vivo studies involving injection of 10-230BM STC and

MDA-231-Brm2. Male mice were used exclusively because injection of these short-term cultures

into female mice results in overwhelming ovarian tumor burden that causes mice decline before

brain metastasis develops.

All experiments were conducted following protocols approved by the NYU Institutional

Animal Care Use Committee (IACUC) (protocol number s16-00051). Animals experiments with

YUMM3.2 were conducted in AAALAC approved facilities at the University of Utah. All animal

protocols were reviewed and approved prior to experimentation by the Institutional Animal Care

and Use Committee (IACUC) at the University of Utah.

Metastasis in the YUMM3.2 syngeneic model. C57BL/6J (RRID:MGI:3028467) male and

female mice were housed in cages containing up to five animals of the same sex and maintained

at room temperature. Mice were fed a combination of Teklad Global 2920X and Teklad 3980X

(Inotiv), supplemented with DietGel76A and HydroGel (Clear H2O) post-weaning. One-week

post-weaning, mice were transitioned to Teklad Global 2920X.

In vivo metastasis assays. 100,000 cells suspended in 100 μl of PBS were injected with

ultrasound guidance (Visualsonics Vevo 770 Ultrasound Imaging System) into the left ventricle of

mice anesthetized with isoflurane. Mice were monitored weekly for metastatic progression by in

vivo Bioluminescence imaging (BLI). Upon weight loss > 20% of their maximum weight over the

course of the experiment, and/or signs of distress (neurological signs, abnormal locomotion) in

any experimental mice, experimental endpoint was established. At experimental endpoint, all

mice in all experimental groups were euthanized with ketamine/xylazine (see "Perfusion").

Generally, 12 mice per group underwent intracardiac injection with cancer cells. Figure

legends indicate where group size may vary due to unattended death or failed intracardiac

injection.

In vivo syngeneic allografts. Seven to nine weeks-old glowing head mice were

subcutaneously injected into the dorsal area near the scapula with 5×10^4 Yumm3.2 BrafV600E

Cdkn2a-/- Pten-/- cell expressing Cndp1-HA and Akt1E17K GFP-Luciferase in Matrigel and

observed for tumor growth. Tumors were visualized and measured weekly using bioluminescence

imaging (BLI), and tumor burden was quantified using bioluminescent photon output values and

digital caliper measurements up to 2000 mm^3. The following formulas was used to calculate tumor

volume: (Length X Width^2)/2.

In-vivo BLI. 10 minutes prior to imaging, luciferin substrate (150 mg/kg) was administered

to mice by intraperitoneal injection. Mice were anesthetized with isoflurane (Covetrus, Cat#11695-

6777-2) and imaged by IVIS Illumina instrument (PerkinElmer, (RRID:SCR_025239)) for an

automatically-determined duration (1-120 sec). Signal was quantified by measurement of total

luminescent flux (p/sec/cm2/sr) in drawn brain and body regions of interest using Living Image

software. For the Yumm3.2 syngeneic model organ visualization, at euthanasia, the primary tumor

as well as lungs, livers and brains were imaged ex vivo using the IVIS Spectrum to confirm

luciferase expression and to detect metastases. Living image software (version 4.5.2) was used

to compile images.

Perfusion. Mice were anesthetized with a ketamine (100 mg/kg, Covetrus Cat#071069)

and xylazine (10 mg/kg, Anased, Cat#NDC59399-110-20) cocktail by intraperitoneal injection.

The heart was exposed by gross dissection and an incision was made in the right atrium.

Subsequently, 10 mL of PBS followed by 10 mL of 4% PFA was injected into the left ventricle.

Continuous shRNA-mediated gene silencing. Starting two days prior to intracardiac

injection and continuing through the experimental duration, doxycycline hyclate (200 mg/kg/day)

was administered to mice in food pellets.

Induction of gene silencing in growing metastases. Protocol for *in vivo* metastasis assay

(as described above) was performed. Mice were administered standard chow at the beginning of

the experiment. 14-17 days (indicated in figure legend) after intracardiac injection of cancer cells,

gene silencing was induced by administration of doxycycline hyclate (200 mg/kg/day) in food

pellets.

Mouse tissue processing. Organs harvested from PFA-perfused mice were fixed in 10%

formalin for 72 hours, followed by several washes PBS and resuspension in 70% ethanol for 24

hours. Prior to embedding, brains were sectioned grossly into thirds coronally and livers were

sectioned by lobe. Organs were embedded in paraffin and cut into 5 μM thick sections. The

embedded brain thirds were sectioned coronally through the entire length of tissue at an interval

of 50 μM. For livers, sections were obtained from two levels 100 μM apart. For brain, sections

were obtained from 2 evenly spaced levels of the embedded coronal thirds. In total, this resulted

in 6 serial coronal brain section levels for metastatic quantification. All sections were stained with

H&E.

Statistical Analysis

Statistical analyses were performed with Prism 9 (RRID:SCR_002798), specific tests are

indicated in figure legends. Unless otherwise stated, values are averages and error bars are +/-

standard deviation.

979
Resource Availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Eva Hernando (eva.hernando-monge@nyulangone.org).

Material availability

Requests for resources and reagents should be directed to and will be fulfilled by the lead contact,

Eva Hernando (eva.hernando-monge@nyulangone.org)

Data and code availability

RNA-sequencing/Proteomics/Metabolomics data have been deposited at [TBD] as

[Database:TBD] and are publicly available as of the date of publication. Additional dataset utilized

are detailed indicated in references, figure legends and Methods.

De-Identified human data is available in Supplementary Tables. They are publicly available as of

the date of publication until [TBD].

Original western blots images are available as supplementary information in Document S1.

All code and packages are available in this paper's methods

Any additional information required to reanalyze the data reported in this paper is available from

the lead contact upon request. Eva Hernando (eva.hernando-monge@nyulangone.org).

Acknowledgements. We thank the NYU Grossman School of Medicine core facilities, funded by

Cancer Center Support Grant (NCI/NIH; 5P30CA016087), including the Preclinical Imaging

Laboratory (Youssef Zaim-Wadghiri, Orlando Aristizabal) , the Experimental Pathology Research

Laboratory (Director, Dr. Cindy Loomis) for tissue processing and histology, the Proteomics Core

(Director, Dr. Beatrix Ueberheide) for mass spectrometry data, Metabolomics Core (Director, Dr.

Drew Jones) for metabolic profiling, Genomics Technology Center (Director, Adriana Heguy) for

transcriptomics. This work was funded by NIH/NCI U54 CA263001, R21CA286244,

5R01CA277425, the V Foundation, and the Melanoma Research Alliance. I.O was supported by

NIH Melanoma SPORE grant NCI P50 CA225450. M.L. was funded by the University Research

Priority Program (URPP). A.K was supported by T32CA009161 and T32GM136573. V.O-V was

supported by 5F31CA250364. P.B. has been funded by a National Cancer Center fellowship and

a Melanoma Research Alliance fellowship. S.L.H. is funded by NIH R01CA121118 and the

Melanoma Research Alliance. D.E.B is funded by the NCI Predoctoral to Postdoctoral Fellow

Transition Award (F99/K00) K99 CA245822-07. Uta Mackensen designed and performed the

illustrations that can be found in Figure S7.

Authors contributions

A.K., M.A.G-M and E.H conceived and designed the study. M.A.G-M, A.K., M.N., V.O-V.,

O.K., P.B., A. P.M and E.V.S-dM performed experiments. M.A.G-M., A.K., M.N., A.W., P.B.,

Y.W., M.I., X.K., S.T. and D.E.B analyzed experimental or genomic data. D.E.B., M.P., R.P.,

R.J.S., S.L.H., C.T.V., K.V.R. contributed to data interpretation. I.O., M.L, E.dS and B.I.

provided patients tissues and/or data. A.K., M.A.G-M., D.E.B. interpreted the data. M.A.G-M,

A.K., and E.H. wrote the manuscript. All authors read and approved the manuscript.

Declaration of interest

The authors declare no competing interests.

1022

References

1. Steeg PS. Targeting metastasis. Nat Rev Cancer. 2016;16:201–18.
2. Zemek RM, Anagnostou V, Pires da Silva I, Long G V., Lesterhuis WJ. Exploiting
temporal aspects of cancer immunotherapy. Nat Rev Cancer. 2024;24:480–97.
3. Soave A, Kluwe L, Yu H, Rink M, Gild P, Vetterlein MW, et al. Copy number variations in
primary tumor, serum and lymph node metastasis of bladder cancer patients treated with
radical cystectomy. Sci Rep. 2020;10:21562.
4. Gui P, Bivona TG. Evolution of metastasis: new tools and insights. Trends Cancer
[Internet]. 2022;8:98–109. Available from: http://www.ncbi.nlm.nih.gov/pubmed/34872888
5. Karras P, Bordeu I, Pozniak J, Nowosad A, Pazzi C, Van Raemdonck N, et al. A cellular
hierarchy in melanoma uncouples growth and metastasis. Nature. 2022;610:190–8.
6. Gerstberger S, Jiang Q, Ganesh K. Metastasis. Cell. 2023;186:1564–79.
7. Bergers G, Fendt S-M. The metabolism of cancer cells during metastasis. Nat Rev
Cancer. 2021;21:162–80.
8. Di Martino JS, Nobre AR, Mondal C, Taha I, Farias EF, Fertig EJ, et al. A tumor-derived
type III collagen-rich ECM niche regulates tumor cell dormancy. Nat Cancer. 2021;3:90–
107.
9. Achrol AS, Rennert RC, Anders C, Soffietti R, Ahluwalia MS, Nayak L, et al. Brain
metastases. Nat Rev Dis Primers [Internet]. 2019;5:5. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/30655533
10. Berghoff AS, Schur S, Füreder LM, Gatterbauer B, Dieckmann K, Widhalm G, et al.
Descriptive statistical analysis of a real life cohort of 2419 patients with brain metastases
of solid cancers. ESMO Open. 2016;1:e000024.
11. Schouten LJ, Rutten J, Huveneers HAM, Twijnstra A. Incidence of brain metastases in a
cohort of patients with carcinoma of the breast, colon, kidney, and lung and melanoma.
Cancer. 2002;94:2698–705.

12.   Lamba N, Wen PY, Aizer AA. Epidemiology of brain metastases and leptomeningeal
1050   disease. Neuro Oncol. 2021;23:1447–56.
1051   13.   Long G V., Atkinson V, Lo S, Guminski AD, Sandhu SK, Brown MP, et al. Five-year
1052   overall survival from the anti-PD1 brain collaboration (ABC Study): Randomized phase 2
1053   study of nivolumab (nivo) or nivo+ipilimumab (ipi) in patients (pts) with melanoma brain
1054   metastases (mets). Journal of Clinical Oncology. 2021;39:9508–9508.
1055   14.   Tawbi HA, Forsyth PA, Hodi FS, Algazi AP, Hamid O, Lao CD, et al. Long-term outcomes
1056   of patients with active melanoma brain metastases treated with combination nivolumab
1057   plus ipilimumab (CheckMate 204): final results of an open-label, multicentre, phase 2
1058   study. Lancet Oncol. 2021;22:1692–704.
1059   15.   Long G V, Atkinson V, Lo SN, Guminski AD, Sandhu SK, Brown MP, et al. Ipilimumab
1060   plus nivolumab versus nivolumab alone in patients with melanoma brain metastases
1061   (ABC): 7-year follow-up of a multicentre, open-label, randomised, phase 2 study. Lancet
1062   Oncol. 2025;
1063   16.   Rauschenberg R, Bruns J, Brütting J, Daubner D, Lohaus F, Zimmer L, et al. Impact of
1064   radiation, systemic therapy and treatment sequencing on survival of patients with
1065   melanoma brain metastases. Eur J Cancer. 2019;110:11–20.
1066   17.   Vosoughi E, Lee JM, Miller JR, Nosrati M, Minor DR, Abendroth R, et al. Survival and
1067   clinical outcomes of patients with melanoma brain metastasis in the era of checkpoint
1068   inhibitors and targeted therapies. BMC Cancer. 2018;18:490.
1069   18.   Tio M, Wang X, Carlino MS, Shivalingam B, Fogarty GB, Guminski AD, et al. Survival and
1070   prognostic factors for patients with melanoma brain metastases in the era of modern
1071   systemic therapy. Pigment Cell Melanoma Res [Internet]. 2018;31:509–15. Available
1072   from: http://www.ncbi.nlm.nih.gov/pubmed/29277979
1073   19.   Bander ED, Yuan M, Carnevale JA, Reiner AS, Panageas KS, Postow MA, et al.
1074   Melanoma brain metastasis presentation, treatment, and outcomes in the age of targeted
1075   and immunotherapies. Cancer. 2021;127:2062–73.
1076   20.   Knisely JPS, Yu JB, Flanigan J, Sznol M, Kluger HM, Chiang VLS. Radiosurgery for
1077   melanoma brain metastases in the ipilimumab era and the possibility of longer survival. J
1078   Neurosurg. 2012;117:227–33.
1079   21.   Fukumura K, Malgulwar PB, Fischer GM, Hu X, Mao X, Song X, et al. Multi-omic
1080   molecular profiling reveals potentially targetable abnormalities shared across multiple
1081   histologies of brain metastasis. Acta Neuropathol. 2021;141:303–21.
1082   22.   Biermann J, Melms JC, Amin AD, Wang Y, Caprio LA, Karz A, et al. Dissecting the
1083   treatment-naive ecosystem of human melanoma brain metastasis. Cell [Internet].
1084   2022;185:2591-2608.e30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/35803246
1085   23.   Ngo B, Kim E, Osorio-Vasquez V, Doll S, Bustraan S, Liang RJ, et al. Limited
1086   Environmental Serine and Glycine Confer Brain Metastasis Sensitivity to PHGDH
1087   Inhibition. Cancer Discov. 2020;10:1352–73.
1088   24.   Karz A, Dimitrova M, Kleffman K, Alvarez-Breckenridge C, Atkins MB, Boire A, et al.
1089   Melanoma central nervous system metastases: An update to approaches, challenges,
1090   and opportunities. Pigment Cell Melanoma Res. 2022;35:554–72.

1091   25.   Kleffman K, Levinson G, Rose IVL, Blumenberg LM, Shadaloey SAA, Dhabaria A, et al. Melanoma-Secreted Amyloid Beta Suppresses Neuroinflammation and Promotes Brain Metastasis. Cancer Discov. 2022;12:1314–35.

26.   Fischer GM, Jalali A, Kircher DA, Lee W-C, McQuade JL, Haydu LE, et al. Molecular Profiling Reveals Unique Immune and Metabolic Features of Melanoma Brain Metastases. Cancer Discov [Internet]. 2019;9:628–45. Available from: http://www.ncbi.nlm.nih.gov/pubmed/30787016

27.   Burdette SC, Lippard SJ. Meeting of the minds: Metalloneurochemistry. Proceedings of the National Academy of Sciences. 2003;100:3605–10.

28.   Piskounova E, Agathocleous M, Murphy MM, Hu Z, Huddlestun SE, Zhao Z, et al. Oxidative stress inhibits distant metastasis by human melanoma cells. Nature. 2015;527:186–91.

29.   Nease LA, Church KP, Delclaux I, Murakami S, Astorkia M, Zerhouni M, et al. Selenocysteine tRNA methylation promotes oxidative stress resistance in melanoma metastasis. Nat Cancer. 2024;

30.   Chi Y, Remsik J, Kiseliovas V, Derderian C, Sener U, Alghader M, et al. Cancer cells deploy lipocalin-2 to collect limiting iron in leptomeningeal metastasis. Science (1979) [Internet]. 2020;369:276–82. Available from: https://www.science.org/doi/10.1126/science.aaz2193

31.   Pegova A, Abe H, Boldyrev A. Hydrolysis of carnosine and related compounds by mammalian carnosinases. Comp Biochem Physiol B Biochem Mol Biol. 2000;127:443–6.

32.   Jackson MC, Kucera CM, Lenney JF. Purification and properties of human serum carnosinase. Clinica Chimica Acta. 1991;196:193–205.

33.   Teufel M, Saudek V, Ledig J-P, Bernhardt A, Boularand S, Carreau A, et al. Sequence Identification and Characterization of Human Carnosinase and a Closely Related Non-specific Dipeptidase. Journal of Biological Chemistry. 2003;278:6521–31.

34.   Regazzoni L. State of the Art in the Development of Human Serum Carnosinase Inhibitors. Molecules. 2024;29:2488.

35.   Reddy VP, Garrett MR, Perry G, Smith MA. Carnosine: A Versatile Antioxidant and Antiglycating Agent. Science of Aging Knowledge Environment. 2005;2005.

36.   Peters V, Yard B, Schmitt CP. Carnosine and Diabetic Nephropathy. Curr Med Chem. 2020;27:1801–12.

37.   Bauer K. Carnosine and Homocarnosine, the Forgotten, Enigmatic Peptides of the Brain. Neurochem Res. 2005;30:1339–45.

38.   Berezhnoy DS, Stvolinsky SL, Lopachev A V., Devyatov AA, Lopacheva OM, Kulikova OI, et al. Carnosine as an effective neuroprotector in brain pathology and potential neuromodulator in normal conditions. Amino Acids. 2019;51:139–50.

39.   Naletova I, Greco V, Sciuto S, Attanasio F, Rizzarelli E. Ionophore Ability of Carnosine and Its Trehalose Conjugate Assists Copper Signal in Triggering Brain-Derived Neurotrophic Factor and Vascular Endothelial Growth Factor Activation In Vitro. Int J Mol Sci [Internet]. 2021;22:13504. Available from: https://www.mdpi.com/1422-0067/22/24/13504

40.   Bellia F, Lanza V, Naletova I, Tomasello B, Ciaffaglione V, Greco V, et al. Copper(II) Complexes with Carnosine Conjugates of Hyaluronic Acids at Different Dipeptide Loading

Percentages Behave as Multiple SOD Mimics and Stimulate Nrf2 Translocation and Antioxidant Response in In Vitro Inflammatory Model. Antioxidants. 2023;12:1632.

41. Anderson EJ, Vistoli G, Katunga LA, Funai K, Regazzoni L, Monroe TB, et al. A carnosine analog mitigates metabolic disorders of obesity by reducing carbonyl stress. Journal of Clinical Investigation. 2018;128:5280–93.

42. Prokopieva VD, Yarygina EG, Bokhan NA, Ivanova SA. Use of Carnosine for Oxidative Stress Reduction in Different Pathologies. Oxid Med Cell Longev. 2016;2016:1–8.

43. Caruso G, Di Pietro L, Cardaci V, Maugeri S, Caraci F. The therapeutic potential of carnosine: Focus on cellular and molecular mechanisms. Current Research in Pharmacology and Drug Discovery. 2023;4:100153.

44. Gaunitz F, Hipkiss AR. Carnosine and cancer: a perspective. Amino Acids. 2012;43:135–42.

45. Renner C, Zemitzsch N, Fuchs B, Geiger KD, Hermes M, Hengstler J, et al. Carnosine retards tumor growth in vivo in an NIH3T3-HER2/neu mouse model. Mol Cancer. 2010;9:2.

46. Teufel M, Saudek V, Ledig J-P, Bernhardt A, Boularand S, Carreau A, et al. Sequence Identification and Characterization of Human Carnosinase and a Closely Related Non-specific Dipeptidase. Journal of Biological Chemistry. 2003;278:6521–31.

47. Biermann J, Melms JC, Amin AD, Wang Y, Caprio LA, Karz A, et al. Dissecting the treatment-naive ecosystem of human melanoma brain metastasis. Cell [Internet]. 2022;185:2591-2608.e30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/35803246

48. Goding CR. Mitf from neural crest to melanoma: signal transduction and transcription in the melanocyte lineage. Genes Dev. 2000;14:1712–28.

49. De Marchis S, Cosimo Melcangi R, Modena C, Ilaria Cavaretta, Peretto P, Agresti C, et al. Identification of the glial cell types containing carnosine-related peptides in the rat brain. Neurosci Lett. 1997;237:37–40.

50. Karz A, Coudray N, Bayraktar E, Galbraith K, Jour G, Shadaloey AAS, et al. MetFinder: a neural network-based tool for automated quantitation of metastatic burden in histological sections from animal models. bioRxiv [Internet]. 2023;2023.09.07.555360. Available from: http://biorxiv.org/content/early/2023/09/08/2023.09.07.555360.abstract

51. Meeth K, Wang JX, Micevic G, Damsky W, Bosenberg MW. The YUMM lines: a series of congenic mouse melanoma cell lines with defined genetic alterations. Pigment Cell Melanoma Res. 2016;29:590–7.

52. Almazan J, Turapov T, Kircher DA, Stanley KA, Culver K, Medellin AP, et al. Combined inhibition of focal adhesion kinase and RAF/MEK elicits synergistic inhibition of melanoma growth and reduces metastases. Cell Rep Med. 2025;6:101943.

53. Parkman GL, Turapov T, Kircher DA, Burnett WJ, Stehn CM, O'Toole K, et al. Genetic Silencing of AKT Induces Melanoma Cell Death via mTOR Suppression. Mol Cancer Ther. 2024;23:301–15.

54. Mooyaart AL, Zutinic A, Bakker SJL, Grootendorst DC, Kleefstra N, van Valkengoed IGM, et al. Association Between *CNDP1* Genotype and Diabetic Nephropathy Is Sex Specific. Diabetes. 2010;59:1555–9.

55. Iacobini C, Menini S, Blasetti Fantauzzi C, Pesce CM, Giaccari A, Salomone E, et al. FL-926-16, a novel bioavailable carnosinase-resistant carnosine derivative, prevents onset

and stops progression of diabetic nephropathy in *db* / *db* mice. Br J Pharmacol.
2018;175:53–66.

56. Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, et al. MetaboAnalyst
5.0: narrowing the gap between raw spectra and functional insights. Nucleic Acids Res.
2021;49:W388–96.

57. Falletta P, Sanchez-del-Campo L, Chauhan J, Effern M, Kenyon A, Kershaw CJ, et al.
Translation reprogramming is an evolutionarily conserved driver of phenotypic plasticity
and therapeutic resistance in melanoma. Genes Dev. 2017;31:18–33.

58. Rabouw HH, Langereis MA, Anand AA, Visser LJ, de Groot RJ, Walter P, et al. Small
molecule ISRIB suppresses the integrated stress response within a defined window of
activation. Proceedings of the National Academy of Sciences. 2019;116:2097–102.

59. Schneider-Poetsch T, Ju J, Eyler DE, Dang Y, Bhat S, Merrick WC, et al. Inhibition of
eukaryotic translation elongation by cycloheximide and lactimidomycin. Nat Chem Biol.
2010;6:209–17.

60. Musa J, Orth MF, Dallmayer M, Baldauf M, Pardo C, Rotblat B, et al. Eukaryotic initiation
factor 4E-binding protein 1 (4E-BP1): a master regulator of mRNA translation involved in
tumorigenesis. Oncogene. 2016;35:4675–88.

61. Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. Cell.
2017;168:960–76.

62. Ericksen RE, Han W. Malignant manipulaTORs of metabolism: suppressing BCAA
catabolism to enhance mTORC1 activity. Mol Cell Oncol. 2019;6:1585171.

63. Vanweert F, Schrauwen P, Phielix E. Role of branched-chain amino acid metabolism in
the pathogenesis of obesity and type 2 diabetes-related metabolic disturbances BCAA
metabolism in type 2 diabetes. Nutr Diabetes. 2022;12:35.

64. Shah M, Su D, Scheliga JS, Pluskal T, Boronat S, Motamedchaboki K, et al. A Transcript-
Specific eIF3 Complex Mediates Global Translational Control of Energy Metabolism. Cell
Rep. 2016;16:1891–902.

65. Ma S, Liu J-Y, Zhang J-T. eIF3d: A driver of noncanonical cap–dependent translation of
specific mRNAs and a trigger of biological/pathological processes. Journal of Biological
Chemistry. 2023;299:104658.

66. Mukhopadhyay S, Encarnación-Rosado J, Lin EY, Sohn ASW, Zhang H, Mancias JD, et
al. Autophagy supports mitochondrial metabolism through the regulation of iron
homeostasis in pancreatic cancer. Sci Adv. 2023;9.

67. Schleiff E, Turnbull JL. Functional and Structural Properties of the Mitochondrial Outer
Membrane Receptor Tom20. Biochemistry. 1998;37:13043–51.

68. Zhang J, Ahn WS, Gameiro PA, Keibler MA, Zhang Z, Stephanopoulos G. 13C Isotope-
Assisted Methods for Quantifying Glutamine Metabolism in Cancer Cells. 2014. page
369–89.

69. Naletova I, Greco V, Sciuto S, Attanasio F, Rizzarelli E. Ionophore Ability of Carnosine
and Its Trehalose Conjugate Assists Copper Signal in Triggering Brain-Derived
Neurotrophic Factor and Vascular Endothelial Growth Factor Activation In Vitro. Int J Mol
Sci. 2021;22:13504.

70. Barca A, Ippati S, Urso E, Vetrugno C, Storelli C, Maffia M, et al. Carnosine modulates
the Sp1-Slc31a1/Ctr1 copper-sensing system and influences copper homeostasis in

murine CNS-derived cells. American Journal of Physiology-Cell Physiology.
2019;316:C235–45.

71. Zheng P, Zhou C, Lu L, Liu B, Ding Y. Elesclomol: a copper ionophore targeting
mitochondrial metabolism for cancer therapy. Journal of Experimental & Clinical Cancer
Research. 2022;41:271.

72. Li S-R, Bu L-L, Cai L. Cuproptosis: lipoylated TCA cycle proteins-mediated novel cell
death pathway. Signal Transduct Target Ther [Internet]. 2022;7:158. Available from:
https://www.nature.com/articles/s41392-022-01014-x

73. Du J, Huang Z, Li Y, Ren X, Zhou C, Liu R, et al. Copper exerts cytotoxicity through
inhibition of iron-sulfur cluster biogenesis on ISCA1/ISCA2/ISCU assembly proteins. Free
Radic Biol Med. 2023;204:359–73.

74. Vo TTT, Peng T-Y, Nguyen TH, Bui TNH, Wang C-S, Lee W-J, et al. The crosstalk
between copper-induced oxidative stress and cuproptosis: a novel potential anticancer
paradigm. Cell Communication and Signaling. 2024;22:353.

75. Karras P, Bordeu I, Pozniak J, Nowosad A, Pazzi C, Van Raemdonck N, et al. A cellular
hierarchy in melanoma uncouples growth and metastasis. Nature. 2022;610:190–8.

76. Pozniak J, Pedri D, Landeloos E, Van Herck Y, Antoranz A, Vanwynsberghe L, et al. A
TCF4-dependent gene regulatory network confers resistance to immunotherapy in
melanoma. Cell. 2024;187:166-183.e25.

77. Brady DC, Crowe MS, Greenberg DN, Counter CM. Copper Chelation Inhibits
BRAFV600E-Driven Melanomagenesis and Counters Resistance to BRAFV600E and
MEK1/2 Inhibitors. Cancer Res. 2017;77:6240–52.

78. Sundstrøm T, Prestegarden L, Azuaje F, Aasen SN, Røsland GV, Varughese JK, et al.
Inhibition of mitochondrial respiration prevents BRAF-mutant melanoma brain metastasis.
Acta Neuropathol Commun. 2019;7:55.

79. Chrzan N, Hartman ML. Copper in melanoma: At the crossroad of protumorigenic and
anticancer roles. Redox Biol. 2025;81:103552.

80. Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, et al.
Defining a Cancer Dependency Map. Cell. 2017;170:564-576.e16.

81. Rihel J. Copper on the brain. Nat Chem Biol. 2018;14:638–9.

82. Zeid CA, Yi L, Kaler SG. Copper Metabolism and the Liver. The Liver. Wiley; 2020. page
207–14.

83. Chen L, Min J, Wang F. Copper homeostasis and cuproptosis in health and disease.
Signal Transduct Target Ther. 2022;7:378.

84. Rokicki J, Li L, Imabayashi E, Kaneko J, Hisatsune T, Matsuda H. Daily Carnosine and
Anserine Supplementation Alters Verbal Episodic Memory and Resting State Network
Connectivity in Healthy Elderly Adults. Front Aging Neurosci. 2015;7.

85. Masuoka N, Yoshimine C, Hori M, Tanaka M, Asada T, Abe K, et al. Effects of
Anserine/Carnosine Supplementation on Mild Cognitive Impairment with APOE4.
Nutrients. 2019;11:1626.

86. An Y, Li S, Huang X, Chen X, Shan H, Zhang M. The Role of Copper Homeostasis in
Brain Disease. Int J Mol Sci. 2022;23:13850.

87.   Han S, Gim Y, Jang E-H, Hur E-M. Functions and dysfunctions of oligodendrocytes in
1266   neurodegenerative diseases. Front Cell Neurosci [Internet]. 2022;16:1083159. Available
1267   from: http://www.ncbi.nlm.nih.gov/pubmed/36605616
1268   88.   Agarwal D, Sandor C, Volpato V, Caffrey TM, Monzón-Sandoval J, Bowden R, et al. A
1269   single-cell atlas of the human substantia nigra reveals cell-specific pathways associated
1270   with neurological disorders. Nat Commun. 2020;11:4183.
1271   89.   Yan R, Zhang P, Shen S, Zeng Y, Wang T, Chen Z, et al. Carnosine regulation of
1272   intracellular pH homeostasis promotes lysosome-dependent tumor immunoevasion. Nat
1273   Immunol. 2024;25:483–95.
1274   90.   Kish SJ, Perry TL, Hansen S. Regional distribution of homocarnosine, homocarnosine-
1275   carnosine synthetase and homocarnosinase in human brain. J Neurochem.
1276   1979;32:1629–36.
1277   91.   Perry TL, Kish SJ, Sjaastad O, Gjessing LR, Nesbakken R, Schrader H, et al.
1278   Homocarnosinosis: Increased content of homocarnosine and deficiency of
1279   homocarnosinase in brain. J Neurochem. 1979;32:1637–40.
1280   92.   Ordaz RP, Garay E, Limon A, Pérez-Samartín A, Sánchez-Gómez MV, Robles-Martínez
1281   L, et al. GABAA Receptors Expressed in Oligodendrocytes Cultured from the Neonatal
1282   Rat Contain α3 and γ1 Subunits and Present Differential Functional and Pharmacological
1283   Properties. Mol Pharmacol. 2021;99:133–46.
1284   93.   Murata Y, Colonnese MT. GABAergic interneurons excite neonatal hippocampus in vivo.
1285   Sci Adv [Internet]. 2020;6:eaba1430. Available from:
1286   http://www.ncbi.nlm.nih.gov/pubmed/32582852
1287   94.   Tzilivaki A, Tukker JJ, Maier N, Poirazi P, Sammons RP, Schmitz D. Hippocampal
1288   GABAergic interneurons and memory. Neuron [Internet]. 2023;111:3154–75. Available
1289   from: https://linkinghub.elsevier.com/retrieve/pii/S0896627323004750
1290   95.   Barron T, Yalçın B, Su M, Byun YG, Gavish A, Shamardani K, et al. GABAergic neuron-
1291   to-glioma synapses in diffuse midline gliomas. Nature. 2025;
1292   96.   Tagore M, Hergenreder E, Perlee SC, Cruz NM, Menocal L, Suresh S, et al. GABA
1293   Regulates Electrical Activity and Tumor Initiation in Melanoma. Cancer Discov.
1294   2023;13:2270–91.
1295   97.   Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM. The integrated
1296   stress response. EMBO Rep [Internet]. 2016;17:1374–95. Available from:
1297   https://www.embopress.org/doi/10.15252/embr.201642195
1298   98.   Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, et al. An Integrated Stress
1299   Response Regulates Amino Acid Metabolism and Resistance to Oxidative Stress. Mol
1300   Cell. 2003;11:619–33.
1301   99.   Lines CL, McGrath MJ, Dorwart T, Conn CS. The integrated stress response in cancer
1302   progression: a force for plasticity and resistance. Front Oncol. 2023;13.
1303   100.   Ma XM, Blenis J. Molecular mechanisms of mTOR-mediated translational control. Nat
1304   Rev Mol Cell Biol. 2009;10:307–18.
1305   101.   Gorrini C, Harris IS, Mak TW. Modulation of oxidative stress as an anticancer strategy.
1306   Nat Rev Drug Discov. 2013;12:931–47.
1307   102.   Tian X, Zhang S, Zhou L, Seyhan AA, Hernandez Borrero L, Zhang Y, et al. Targeting the
1308   Integrated Stress Response in Cancer Therapy. Front Pharmacol. 2021;12.

1309   103.   Anderson CP, Shen M, Eisenstein RS, Leibold EA. Mammalian iron metabolism and its
1310   control by iron regulatory proteins. Biochimica et Biophysica Acta (BBA) - Molecular Cell
1311   Research. 2012;1823:1468–83.
1312   104.   Pantopoulos K. Iron Metabolism and the IRE/IRP Regulatory System: An Update. Ann N
1313   Y Acad Sci. 2004;1012:1–13.
1314   105.   González M, Reyes-Jara A, Suazo M, Jo WJ, Vulpe C. Expression of copper-related
1315   genes in response to copper load. Am J Clin Nutr. 2008;88:830S-834S.
1316   106.   Oc S, Eraslan S, Kirdar B. Dynamic transcriptional response of Saccharomyces
1317   cerevisiae cells to copper. Sci Rep. 2020;10:18487.
1318   107.   Roy G, Antoine R, Schwartz A, Slupek S, Rivera-Millot A, Boudvillain M, et al.
1319   Posttranscriptional Regulation by Copper with a New Upstream Open Reading Frame.
1320   mBio. 2022;13.
1321   108.   Seiwert N, Wecklein S, Demuth P, Hasselwander S, Kemper TA, Schwerdtle T, et al.
1322   Heme oxygenase 1 protects human colonocytes against ROS formation, oxidative DNA
1323   damage and cytotoxicity induced by heme iron, but not inorganic iron. Cell Death Dis
1324   [Internet]. 2020;11:787. Available from: https://www.nature.com/articles/s41419-020-
1325   02950-8
1326   109.   Hanai J, Mammoto T, Seth P, Mori K, Karumanchi SA, Barasch J, et al. Lipocalin 2
1327   Diminishes Invasiveness and Metastasis of Ras-transformed Cells. Journal of Biological
1328   Chemistry. 2005;280:13641–7.
1329   110.   Xu W-X, Zhang J, Hua Y-T, Yang S-J, Wang D-D, Tang J-H. An Integrative Pan-Cancer
1330   Analysis Revealing LCN2 as an Oncogenic Immune Protein in Tumor Microenvironment.
1331   Front Oncol. 2020;10.
1332   111.   Tsvetkov P, Coy S, Petrova B, Dreishpoon M, Verma A, Abdusamad M, et al. Copper
1333   induces cell death by targeting lipoylated TCA cycle proteins. Science (1979).
1334   2022;375:1254–61.
1335   112.   Kong R, Sun G. Targeting copper metabolism: a promising strategy for cancer treatment.
1336   Front Pharmacol. 2023;14.
1337   113.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio Cancer
1338   Genomics Portal: An Open Platform for Exploring Multidimensional Cancer Genomics
1339   Data. Cancer Discov. 2012;2:401–4.
1340   114.   Li T, Fan J, Wang B, Traugh N, Chen Q, Liu JS, et al. TIMER: A Web Server for
1341   Comprehensive Analysis of Tumor-Infiltrating Immune Cells. Cancer Res. 2017;77:e108–
1342   10.
1343   115.   Agrawal A, Balcı H, Hanspers K, Coort SL, Martens M, Slenter DN, et al. WikiPathways
1344   2024: next generation pathway database. Nucleic Acids Res. 2024;52:D679–89.
1345   116.   Robert Jan Lebbink MM. Cloning small hairpains into lentiviral vectors .
1346   https://mcmanuslab.ucsf.edu/protocol/cloning-small-hairpins-lentiviral-vectors.
1347   117.   Kircher DA, Trombetti KA, Silvis MR, Parkman GL, Fischer GM, Angel SN, et al.
1348   AKT1E17K Activates Focal Adhesion Kinase and Promotes Melanoma Brain Metastasis.
1349   Mol Cancer Res [Internet]. 2019;17:1787–800. Available from:
1350   http://www.ncbi.nlm.nih.gov/pubmed/31138602

1351   118.   de Miera EV, Friedman EB, Greenwald HS, Perle MA, Osman I. Development of five new
1352   melanoma low passage cell lines representing the clinical and genetic profile of their
1353   tumors of origin. Pigment Cell Melanoma Res. 2012;25:395–7.
1354   119.   Fontanals-Cirera B, Hasson D, Vardabasso C, Di Micco R, Agrawal P, Chowdhury A, et
1355   al. Harnessing BET Inhibitor Sensitivity Reveals AMIGO2 as a Melanoma Survival Gene.
1356   Mol Cell. 2017;68:731-744.e9.
1357   120.   Jones DR, Wu Z, Chauhan D, Anderson KC, Peng J. A nano ultra-performance liquid
1358   chromatography-high resolution mass spectrometry approach for global metabolomic
1359   profiling and case study on drug-resistant multiple myeloma. Anal Chem. 2014;86:3667–
1360   75.
1361   121.   Simón-Manso Y, Lowenthal MS, Kilpatrick LE, Sampson ML, Telu KH, Rudnick PA, et al.
1362   Metabolite profiling of a NIST Standard Reference Material for human plasma (SRM
1363   1950): GC-MS, LC-MS, NMR, and clinical laboratory analyses, libraries, and web-based
1364   resources. Anal Chem. 2013;85:11725–31.
1365   122.   Smith CA, Maille GO, Want EJ, Qin C, Trauger SA, Brandon TR, et al. METLIN. Ther
1366   Drug Monit. 2005;27:747–51.
1367   123.   Wang X, Jones DR, Shaw TI, Cho J-H, Wang Y, Tan H, et al. Target-Decoy-Based False
1368   Discovery Rate Estimation for Large-Scale Metabolite Identification. J Proteome Res.
1369   2018;17:2328–34.
1370   124.   Wang X, Cho J-H, Poudel S, Li Y, Jones DR, Shaw TI, et al. JUMPm: A Tool for Large-
1371   Scale Identification of Metabolites in Untargeted Metabolomics. Metabolites. 2020;10.
1372   125.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al.
1373   SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods.
1374   2020;17:261–72.
1375   126.   Ernlund AW, Schneider RJ, Ruggles K V. RIVET: comprehensive graphic user interface
1376   for analysis and exploration of genome-wide translatomics data. BMC Genomics.
1377   2018;19:809.
1378   127.   Fernandez CA, Rosiers C Des, Previs SF, David F, Brunengraber H. Correction of13C
1379   Mass Isotopomer Distributions for Natural Stable Isotope Abundance. Journal of Mass
1380   Spectrometry. 1996;31:255–62.
1381   128.   Parker SJ, Encarnación-Rosado J, Hollinshead KER, Hollinshead DM, Ash LJ, Rossi
1382   JAK, et al. Spontaneous hydrolysis and spurious metabolic properties of α-ketoglutarate
1383   esters. Nat Commun. 2021;12:4905.
1384   129.   Biancur DE, Paulo JA, Małachowska B, Quiles Del Rey M, Sousa CM, Wang X, et al.
1385   Compensatory metabolic networks in pancreatic cancers upon perturbation of glutamine
1386   metabolism. Nat Commun. 2017;8:15965.
1387   130.   Betancourt LH, Gil J, Sanchez A, Doma V, Kuras M, Murillo JR, et al. The Human
1388   Melanoma Proteome Atlas—Complementing the melanoma transcriptome. Clin Transl
1389   Med. 2021;11.
1390   131.   Hayashi A, Ruppo S, Heilbrun EE, Mazzoni C, Adar S, Yassour M, et al. GENI: A web
1391   server to identify gene set enrichments in tumor samples. Comput Struct Biotechnol J.
1392   2023;21:5531–7.

1393   132.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative
1394   Analysis of Complex Cancer Genomics and Clinical Profiles Using the cBioPortal. Sci
1395   Signal. 2013;6.
1396   133.   de Bruijn I, Kundra R, Mastrogiacomo B, Tran TN, Sikina L, Mazor T, et al. Analysis and
1397   Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE
1398   Biopharma Collaborative in cBioPortal. Cancer Res. 2023;83:3861–7.
1399   134.   Li B, Severson E, Pignon J-C, Zhao H, Li T, Novak J, et al. Comprehensive analyses of
1400   tumor immunity: implications for cancer immunotherapy. Genome Biol. 2016;17:174.
1401   135.   Varešlija D, Priedigkeit N, Fagan A, Purcell S, Cosgrove N, O'Halloran PJ, et al.
1402   Transcriptome Characterization of Matched Primary Breast and Brain Metastatic Tumors
1403   to Detect Novel Actionable Targets. JNCI: Journal of the National Cancer Institute.
1404   2019;111:388–98.
1405   136.   Xiao L, Zhou J, Liu H, Zhou Y, Chen W, Cui W, et al. RNA Sequence Profiling Reveals
1406   Unique Immune and Metabolic Features of Breast Cancer Brain Metastases. Front Oncol.
1407   2021;11.
1408

1409

Figure legends

Figure 1

Figure 1. Multi-omics approach nominates *CNDP1* as driver of melanoma brain metastasis.

(A) Data integration strategy showing log2Fold expression changes between patient-matched

Brain Metastasis (BM) and Non-Brain Metastasis (NBM) tissue samples (RNA-sequencing n = 29

patients with the following samples: 35 brain, 7 intestine, 18 lymph node, 5 lung, 12 skin.

Proteomics, n = 3 patients with the following samples: 3 brain, 1 subcutaneous, 1 bowel). NBM

sources in RNA-sequencing data included P value (edgeR) < 0.005 for differential expression in

RNA-sequencing (EGAS000010036720, Fischer et al.) No p-value threshold was applied for

proteomics data. Genes labeled are log2FC > 0.4 in both studies. Additional data can be found in

Table S1 and S2.

(B) Kaplan-Meier curve showing patient Overall Survival stratified by high and low CNDP1 protein

levels in lymph nodes metastasis on MM500 study(130). Cut off established between the lower

and upper quartiles of protein intensity values. T test by log-rank test. Corrected for 50% minimum

tumor content.

(C) *CNDP1* differential mRNA expression represented as normalized counts in melanoma

samples spread to the lymph node that later on metastasize to the brain or to other organs.

Statistical analysis by Well's t-test. Additional data can be found in Table S4.

(D) Single Nucleus RNA-sequencing (snRNA-seq) data of human patient samples mined from

Biermann et al (Series GSE185386). UMAP representing *CNDP1* expression in extracranial

metastasis (NBM) vs intracranial metastasis (MBM). Statistics derived from Seurat's FindMarkers

function which applies a Wilcoxon Rank Sum test.

(E) *CNDP1* normalized counts of single cell RNA-seq data of malignant cells (n=333050 cells)

from lung adenocarcinoma primary tumors (LPT) and lung adenocarcinoma brain metastasis

(LBM). Data from Tagore, Caprio et al. *in press* in Nature Medicine. Statistical analysis by Welch's

test.

(F) *CNDP1* differential mRNA expression (2log TMMCPM) in breast cancer (BC) versus breast

cancer brain metastasis (BCBM) paired patients' samples from indicated data set. Statistical

analysis by paired t-test.

(G) Immunohistochemistry (IHC) showing differential expression of CNDP1 in patient-matched

primary, extracranial metastasis and brain metastatic samples (TMA, n samples indicated). **Left.**

Representative primary, extracranial and brain metastasis IHC DAB images (BM003, BM0026)

Scale=10uM**. Right.** CNDP1 expression quantification in healthy skin, healthy cortex, primary

tumor, NBM (no brain metastasis) and BM (brain metastasis) samples represented as % of

positive cells. P value by Paired One way ANOVA. Additional data included in Table S3.

See Figure S1

1446

1447

1448

1449

1450

1451

1452

1453

1454

Figure 2

1455

Figure 2. CNDP1 modulation impacts 2D proliferation of melanoma cells.

(A) qRT-PCR quantification of *CNDP1* mRNA, relative to GAPDH and normalized to shNTC, in

10-230 BM (top) and 12-273 (bottom) cells expressing indicated shRNA.

(B) Top. Proliferation curve performed by analyzing % confluency extracted from Incucyte image

analysis normalized to day 0 of 10-230 BM cells expressing indicated tet-On shRNA (n=3, 96 h).

Bottom. Proliferation curve performed by serial fixing and crystal violet staining of 12-273BM cells

expressing indicated tet-On shRNA. Representative experiment shown of n=3 biological

replicates. Statistics derived from one-way ANOVA testing between groups on the final time point.

(C) Bar plots representing % cells distributed along cell cycle phases assessed by Edu differential

staining (n=2) in 10-230 and 12-273 BM cells expressing indicated tet-On shRNA. Statistical

analysis by one-way ANOVA with Dunnett multiple hypothesis testing correction.

(D) Left. qRT-PCR quantification of *CNDP1* mRNA, relative to hPPIA and then normalized to

siNTC, in 10-230 BM. **Right.** Proliferation ratio of 10-230 BM cells transfected with indicated

siRNAs after 72 h of culture normalized to 0h (n=3), indicated as % confluency extracted from

Incucyte image analysis. Multiple biological replicates represented. Statistical analysis by one-

way ANOVA.

(E). Left. qRT-PCR quantification of *CNDP1* mRNA, relative to hPPIA and then normalized to

sINTC, in 12-273 BM. **Right.** Proliferation ratio of 12-273 BM cells transfected with indicated

siRNAs after 96 h culture normalized to 0h (n=4), indicated as % confluency extracted from

Incucyte image analysis technology. Multiple biological replicates represented. Statistical analysis

by one-way ANOVA.

See Figure S2

1478

Figure 3. CNDP1 silencing impairs establishment and maintenance of *in Vivo* melanoma

metastasis, and Cndp1 ectopic expression drives melanoma brain metastasis.

(A) Schematic representation of the *in vivo* xenograft constitutive approach experiment (USG ICI:

Ultra Sound Guided Intra-Cardiac Injection).

(B) Representative images of whole-body *in vivo* BLI of NSG mice injected with 12-273 BM cells

transduced with indicated tetON-shRNA 28 days post-injection (n = 12 mice per group, BLI

represented by radiance: p/sec/cm^2/sr).

(C) Representative H&E images and corresponding MetFinder-generated heatmaps of brain and

liver sections of NSG mice injected with indicated cells (n = 12 mice per group, two sections at 3

levels analyzed per organ/mice. Scale bars represent 1000 μm and 2000 μm for brain and liver,

respectively).

(D) Tumor burden in brain parenchyma, leptomeninges and liver quantified as the percent of area

occupied by tumor and average of 2 levels of sectioning per organ (100 μM separation), obtained

by MetFinder analysis. Statistical analysis by one-way ANOVA.

(E) Schematic representation of the *in vivo* xenograft therapeutic approach experiment (USG ICI).

(F) Representative images of whole-body *in vivo* BLI of NSG mice injected with 12-273 BM cells

transduced with indicated tetON-shRNA at day 36 post-injection (n = 12 mice per group, BLI

expressed as radiance: p/sec/cm^2/sr).

(G) Representative H&E and MetFinder brain and liver metastasis sections of NSG mice injected

with indicated vectors (n = 12 mice per group; duplicated sections and levels analyzed per

organ/mice. Scale bar representative of 1000 μm and 2000 μm for brain and liver, respectively)

(H) Tumor burden in brain parenchyma, leptomeninges and liver quantified as the percent of

tumor-occupied area vs total organ area and averaged of 2 levels of sectioning per organ.

Obtained by performing MetFinder analysis. Statistical analysis by one-way ANOVA.

(I) Schematic representation of *in vivo* allograft experiment with indicated conditions. S.I:

Subcutaneous injection.

(J) Ex vivo BLI of organs in the corresponding groups of mice injected with YUMM3.2 cells control

or transduced with Akt1-E17K or Cndp1-HA in indicated organs (Pt=Primary tumor). (n = 9,10

&11 mice per group, respectively. BLI expressed as radiance: p/sec/cm^2/sr).

(K) Bar graph representing % of tumor incidence for primary (Pt), brain, lung and liver metastasis

in mice injected with indicated conditions. Statistical analysis results from one-way ANOVA

comparing metastasis incidence at end point (1=yes, 0=no).

See Figure S3

1512

1513

1514

1515

1516

1517

1518

1519

1520

1521

1522

Figure 4. CNDP1 knockdown elicits an integrative stress response in melanoma cells.

(A) Workflow of RNA-sequencing and LC/MS-based nonpolar metabolomics integrative studio in

10-230BM cells transduced with indicated tetON-shRNA.

(B) Differential gene expression/metabolite abundance between shNTC vs shCNDP1-1 and

shCNDP1-2 by indicated adjusted p values. MetaboAnalyst executed hypergeometric testing by

combining p-values for metabolites and transcripts according to their relative proportion in each

pathway. (RNA-seq: DESeq2, Metabolomics: T-test, pathways enrichment FDR < 0.05). See

Tables S5-7

(C) Heatmap of t-Aminoacyl synthetases expression across different conditions paired with a

heatmap of expression of their corresponding amino acids. MS/MS-based proteomics and

LC/MS-based metabolomics were conducted on lysates of 10-230 BM cells expressing the

indicated shRNA treated with doxycycline for 72 hours in 10% dialyzed serum DMEM.

Differentially expressed proteins were identified by unpaired, two- tailed t-test comparing 2 groups

normalized abundance: shCNDP1-1 and shCNDP1-2 vs shNTC (Data represented by z-score as

indicated).

(D) Integrated Gene Set Enrichment analysis (GSEA) of 12-273 and 10-230 BM shNTC vs both

shCNDP1 tetON-shRNA RNA sequencing represented by Normalized Enrichment Score using

Reactome Gene Ontology Analysis (RNA-seq: DESeq2, adjust p value < 0.001 and FC > 1.5).

See Table S6.

(E) Differential expression testing by DESeq2 nominated ATF4 targets in both shCNDP1 vs

shNTC conditions in 10-230 BM RNA sequencing. (Data represented by z-score as indicated).

(F) Pathway analysis via the Seq-n-Slide pipeline tested for pathway enrichment in differentially

expressed genes between shCNDP1 and shNTC 10-230 BM RNA sequencing (Adjust p value

indicated).

(G) Representative immunoblots of phosphoSerine51 eIF2a, total-eIF2a, ATF4 and

housekeeping (HK) protein Vinculin in lysates of 10-230 BM transduced with indicated tetON-

shRNA. See Document S1 for details.

(H) Proliferation ratio of 12-273 BM cells transfected with indicated siRNAs for CNDP1 silencing

after 96 h of culture normalized to 0h (n=4), either treated with vehicle (DMSO) or ISRIB (phospho-

eIF2a inhibitor, 1 uM) performed by analyzing % confluency extracted from Incucyte image

analysis. Statistical analysis by one-way ANOVA. Proliferation ratio for 12-230 BM

siNTC/siCNDP1 also shown in Figure 2 E.

See also Figure S4

1556

1557

Figure 5

1558

Figure 5. CNDP1 knockdown rewires translation program towards mitochondrial transcripts.

(A) 10-230 and 12-273 BM transduced with indicated tetON-shRNAs and treated with doxycycline during 72h, labeled with AHA (APC) for 4 hours and analyzed by flow cytometry. Cycloheximide (CHX, 50 ug/ml) was used as a positive control of translation shutdown. Statistics rendered by one-way ANOVA with Dunnet's multiple hypothesis testing correction.

(B) Left. Representative immunoblots of phospho-S6K1 p70 (Threonine 389), S6K1 p70, phosphoSer65 4E-BP, total 4EBP and housekeeping (HK) protein Hsp90 in lysates of 10-230 BM transduced with indicated tetON-shRNA. **Right.** Representative immunoblots of phospho-S6K1 p70 (Threonine 389), S6K1 p70, phosphoSer65 4E-BP, 4EBP and housekeeping (HK) protein Hsp90 in lysates of 10-230 BM cells transfected with siRNA NTC or CNDP1 as indicated. Band

correspondent to p70 S6K1 is indicated. Results for HRI, phosphoSer51 and total eiF2α

expression from same samples shown in Figure 6K and Figure S4H, respectively. Ratio of protein

phosphorylation shown normalized to total protein and HK protein levels. See Document S1 for

details.

(C) GENI (gene set enrichment identifier124) was applied to TCGA melanoma samples (n = 472)

using the Reactome 2022 pathway database. Five anticorrelating gene sets with FDR < 0.05 were

represented.

(D) Genes differentially expressed in shCNDP1 (combined) vs shNTC polysome heavy chain (log

fc Translation vs log fc Transcription; p<0.05, lfc > 1) at translation, transcription and both

transcription and translation levels in 10-230 BM cells. See Table 11.

(E) Integrated Gene Enrichment analysis of Heavy Chain RNA sequencing of 10-230 BM

shCNDP1 (combined) vs shNTC represented by Normalized Enrichment Score using Reactome

Gene Ontology Analysis (RNA-seq: adjust p value < 0.05).

See also Figure S5

Figure 6

1584

Figure 6. CNDP1 knockdown melanoma cells display profound mitochondrial defects, copper overload and ROS accumulation.

(A) Mitochondria imaging using electron microscopy of 10-230 BM transduced with indicated tetON-shRNA. Scale =1 μM.

(B-E) Quantification of number of mitochondria per cell **(B)**, mitochondria area **(C)**, cristae number

per mitochondria **(D)** and cristae area per mitochondria **(E)** area in 10-230 BM. Statistical analysis

by one-way ANOVA (p<0.05 or indicated).

(F) Tom20 expression by IHC in BM from 12-273 BM cells transduced with indicated shRNAs

induced with doxycycline during 5 days. Scale = 200 μM. Quantification of Tom20 cytosolic

staining DAB intensity per cell and per mice. Statistical analysis by two-way ANOVA (p<0.001).

(G) Top. Seahorse MitoStress analysis of OCR in 12-273 BM transduced with indicated tetON-

shRNA. Statistical analysis by one-way ANOVA. Representative replicate shown (n=3, p<0.05 or

indicated). **Bottom** Seahorse MitoStress analysis of OCR in 12-273 BM treated with Carnosine

40 mM overnight. Statistical analysis by one-way ANOVA. Representative replicate shown (n=3,

p<0.05 or indicated).

(H) Schematic model of glutamine metabolism in cancer cells. Green (M+5) and purple circles

(M+4) represent carbons derived from [U-13C5] glutamine. The black arrows indicate oxidative

and reductive carboxylation flux from glutamine. Cells were cultured with [U-^{13}C5] glutamine for 6

h before metabolite extraction and gas chromatography-mass spectrometry (GC-MS) analyses.

Normalized distribution of glutamine in mentioned conditions in 12-273 BM cells transduced with

indicated tetON-shRNAs treated with doxycycline for 24h are shown. (n = 3); data are shown as

the mean ± SD.

(I) Normalized distributions of glutamine, glutamate, and TCA cycle metabolites, including α-

ketoglutarate (αKG), succinate, fumarate, malate and citrate, in 12-273 BM cells transduced with

indicated tetON-shRNAs treated with doxycycline for 24h are shown. (n = 3); data are shown as

the mean ± SD. Statistical analysis by one-way ANOVA. P value indicated.

(J) Mean fluorescence intensity of CellROX in live cells as determined by DAPI- status in 10-230

(left) and 12-273 BM (right) cells. Statistical analysis by one-way ANOVA. Representative

replicate shown (n=3).

(K) Left. Representative immunoblots of II & I HRI fractions and housekeeping (HK) protein b-

actin in lysates of 12-273 BM transduced with indicated tetON-shRNA. **Right.** Representative

immunoblots of II & I HRI fractions and housekeeping (HK) protein b-actin in lysates of 10-230

BM cells transfected with siRNA NTC or CNDP1 as indicated. II fraction considered as the

autophosphorylated. Results for translation related proteins, phosphoSer51 and total eiF2α

expression from same samples shown in Figure 5B and Figure S4H, respectively. Representative

HK shown. See Document S1 for details.

(L) Copper abundance represented as ratio of emission signal at 563 nm in indicated conditions

in 10-230 BM (left) and 12-273 BM (right) normalized to total number of cells quantified by crystal

violet (595 nm absorbance values). Statistical analysis by one-way ANOVA. Representative

replicate shown (n=3).

See also Figure S6

1626

1627

1628

Figure S1. *CNDP1 In silico* analysis in melanoma databases. Related to Figure 1.

(A) Comparative analysis of log 10 transcription *CNDP1, CNDP2* and *CARNS1* expression levels

(TPM+1) in human normal tissues mined from GTEx, the Genotype-Tissue Expression project.

(B) *CNDP1, CNDP2, CARNS1* (RSEM analysis in normalized counts) expression in primary vs

metastatic melanoma human samples from Melanoma dataset Cancer Genome Atlas RNA

sequencing (n = 472 total samples from 469 patients. n = 369 metastases and n = 103 primary).

P value by unpaired, two-tailed t-test with Welch's correction.

(C) Kaplan-Meier curves showing patient Overall Survival stratified by high and low *CNDP1,*

CNDP2 and *CARNS1* gene expression (RNA-seq) in primary and metastatic tumor on

TCGA(113,132,133) Melanoma (SKCM Firehouse Legacy, n = 472) and log-rank test by

TIMER2.0(114,131,134).

(D) 2logFC *CNDP1* expression in breast cancer data bases described in the indicated references

(135,136) (adj p value p<0.0001).

(E) *CNDP1* expression (qRT-PCR) in matched extracranial (NBM) and brain metastasis (BM)

derived short term cultures 12-273, WM4527, 10-230. Expression represented as Fold Change

relative to housekeeping gene (hPPIA) and to extracranial derived matched short-term culture

(n=3). Additional data can be found in Table S1

(F) Immunoblot showing CNDP1 expression in patient-matched extracranial (NBM) and brain

metastasis (BM) derived short term cultures 12-273BM, WM-4527, 10-230BM. Additional data

can be found in Table S1. See also Document S1 for details.

(G). L-Histidine liberation assay as carnosinase activity read-out in conditioned media of 10-230

and 12-273 paired matched STC. Additional data can be found in Table S1. Two technical

replicates shown.

(H) *CNDP1* expression significantly correlates with the average epigenetic accessibility around

CNDP1 promoters. In red is shown the linear regression line. R represents the Spearman

coefficient.

(I) Combined heatmap showing the epigenetic accessibility surrounding *CNDP1* and *CNDP2* loci

in a subset of TCGA patients (left) and *CNDP1* expression in the corresponding samples (right),

(n=404)

(J) IGV snapshot of MITF and the active chromatin marker H3K27ac genomic occupancy around

CNDP1 and *CNDP2* loci generated in 501mel. MITF binds to *CNDP1* promoter and H3K27ac loss

upon MITF silencing at *CNDP1* promoter strongly suggests a direct MITF-dependent CNDP1

regulation.

(K) Bar plot of the mRNA expression of *CNDP1, CNDP2* and *MITF* upon MITF silencing in 501mel

cell. *CNDP1* expression is strongly reduced following MITF depletion compared to *CNDP2* (data

extrapolated from Laurette et al., 2015, GSE61967).

(L) *CNDP1* expression (qRT-PCR) in 501Mel melanoma cells after 24, 48 and 72 h MITF silencing

(siMITF). Expression represented as Fold Change relative to housekeeping gene (hPPIA) and

non-targeting control silencing (siNTC).

1669

1670

Figure S2. CNDP1 modulation effect in melanoma cells proliferation. Related to Figure 2.

(A) Carnosine accumulation measured by ELISA represented as normalized values to Carnosine

concentration in shNTC conditions (30 mins under culture) in 12-273 transduced with indicated

tetON-shRNAs treated with doxycycline for 24h. Mean of two technical replicates shown.

(B) Proliferation ratio of 10-230 BM cells cultured with 10 μm, 100 μm and 50 mM Carnosine

concentrations during 0.5, 2, 6 and 24 h of culture normalized to not treated (NT) of their

respective time points (n=3, technical replicates). Confluency obtained by treating the cells with

Resazurin and measuring absorbance after 3 hours of treatment at 570 nm. Statistical analysis

by one-way ANOVA.

(C) Carnosine accumulation shown as normalized units to STC media in conditioned media and

pellet after 30 minutes of treatment.

(D) Proliferation ratio of SKMEL-239 cells transfected with indicated shRNAs after 96 h of culture

normalized to 0h, indicated as % confluency extracted from Incucyte image analysis.

(E) Proliferation ratio performed by serial fixing and crystal violet staining of Yumm3.2

BrafV600E/wt Cdkn2a-/- Pten-/- control or ectopically expressing Cndp1-HA after 96h as

indicated.

(F) Immunoblot showing Cndp1-HA expression in conditioned media and pellets of Yumm3.2

BrafV600E/wt Cdkn2a-/- Pten-/- control or ectopically expressing Cndp1-HA. See also Document

S1 for details.

(G) L-Histidine liberation assay as carnosinase activity read-out in pellets of Yumm3.2

BrafV600E/wt Cdkn2a-/- Pten-/- control or ectopically expressing Cndp1-HA. Three technical

replicates shown.

One representative technical replicate shown for all the above assays.

Figure S3

1694

Figure S3. CNDP1 Silencing Decreases *In Vivo* Melanoma Metastasis. Related to Figure 3.

(A) Schematic representation of the *in vivo* xenograft constitutive approach experiment (USG ICI).

(B) Representative images of whole-body *in vivo* BLI of athymic/nude mice injected with 10-230

BM cells transduced with indicated tetON-shRNA 52 days post-injection (n = 12 mice per

condition, BLI represented by radiance: $p/sec/cm^2/sr$).

(C) Representative H&E and MetFinder brain metastasis sections of nude mice injected with

indicated (n = 12 mice per condition, duplicated sections and levels analyzed per organ/mice,

Scale bar 1000 μm and 2000 μm for brain and liver, respectively).

(D) Tumor burden in brain, leptomeningeal and liver quantified as the percent of area occupied

by the tumor of the total organ area and averaged of 2 levels of sectioning per organ obtained by

performing MetFinder analysis. Statistical analysis by one-way ANOVA.

(E) Schematic representation of the *in vivo* xenograft therapeutic approach experiment (USG ICI:

Ultra Sound Guided Intra-Cardiac Injection).

(F) Representative images of whole-body *in vivo* BLI of nude mice injected with 10-230 BM cells

transduced with indicated tetON-shRNA day 59 post-injection (n = 12 mice per condition, BLI

expressed as radiance: $p/sec/cm^2/sr$).

(G) Representative H&E and MetFinder brain and liver metastasis sections of NSG mice injected

with indicated (n = 12 mice per condition, duplicated sections and levels analyzed per organ/mice,

Scale bar 1000 μm and 2000 μm for brain and liver, respectively).

(H) Tumor burden in brain, leptomeningeal and liver quantified as the percent of area occupied

by the tumor in the total organ area and averaged of 2 levels of sectioning per organ. Obtained

by performing MetFinder analysis. Statistical analysis by one-way ANOVA.

(I) Percent change in IVIS signal in the brain (top) and whole body (bottom) from days 12 to 36 of

nude mice injected with 10-230 BM cells transduced with indicated tetON-shRNA. One bar = one

mouse, n = 10 mice/group.

(J) Schematic representation of the *in vivo* xenograft therapeutic approach experiment (USG ICI:

Ultra Sound Guided Intra-Cardiac Injection).

(K) Representative H&E images and corresponding quantitation of metastasis burden in brain sections of NSG mice injected with MDA-231 BrM2 cells transduced with indicated tetON-shRNA obtained by manual analysis. Not burden in lung, liver or kidneys detected.

(L) Tumor burden in brain quantified as the percent of area occupied by tumor and average of two levels of sectioning per organ (100 µM separation) obtained by manual analysis (n=10, 8, and 12 mice per group, two sections and two levels analyzed per organ/mice). Statistical test: one-way ANOVA. Scale bars represent 1000 µm. Not burden in lung, liver or kidneys detected.

(M) *CNDP1* expression (counts) in RNA sequencing of 12-273 BM GFP positive cells harvested from 4 mice livers and 3 mice brains of the indicated conditions after USG ICI and 7 days of doxycycline treatment.

1742

Figure S4. Proteomics changes upon CNDP1 silencing in melanoma cells. Related to

Figure 4.

(A) Representation of DAVID functional annotation algorithm of upregulated (right) and

downregulated (left) differentially expressed proteins identified by unpaired, two- tailed t-test

comparing 2 groups: shCNDP1-1 and shCNDP1-2 versus shNTC. Represented by 1/FDR as

indicated. See Table S10.

(B) Representative immunoblots of phophoSerine51 eIF2α and total eIF2α, housekeeping (HK)

protein β-actin in lysates of 10-230 BM cells transfected with siRNA NTC or CNDP1 as indicated.

Results for HRI and translation related proteins expression from samples shown in Figure 6K and

Figure 5B, respectively. Total and phospho-proteins were blotted in different membranes.

Representative HK shown. See Document S1 for details.

1754

Figure S5. Extended characterization of translation phenotype of CNDP1 KD melanoma

cells. Related to Figure 5.

(A) Integrated Gene Enrichment analysis of 12-273 BM heavy chain both shCNDP1 vs shNTC

RNA sequencing represented by Normalized Enrichment Score using Reactome Gene Ontology

Analysis (RNA-seq: adjust p value < 0.05).

(B) Venn diagram including common transcripts changing as indicated at translation level (top)

and transcription and translation level (bottom) in both 10-230 and 12-273 BM shCNDP1

(combined) vs shNTC polysome heavy chain (adjusted p value <0.05, n.i - non-identified). See

Table S11.

(C) Representative immunoblot of eIFs proteins and phophoSerine51 4E-BP1, 4E-BP1 and

housekeeping (HK) protein Vinculin in lysates of 12-273 BM cells transfected with siRNA NTC or

CNDP1 as indicated. See Document S1 for details.

1767

Figure S6. Extended characterization of mitochondrial phenotype of CNDP1 KD melanoma

cells. Related to Figure 6.

(A-E) Quantification of total mitochondria per cell **(A)**, mitochondria area **(B)**, circularity **(C)**, cristae

number per mitochondria **(D)** and cristae area per mitochondria area **(E)** in 12-273 BM. Statistical

analysis by one-way ANOVA.

(F) Seahorse MitoStress analysis of OCR in 10-230 BM transduced with indicated tetON-shRNA

and MDA-231 Brm2 treated as indicated. Statistical analysis by one-way ANOVA. Representative

replicate shown (n=3, p<0.05 or indicated).

(G) Normalized distributions of glutamine, glutamate, and TCA cycle metabolites, including α-

ketoglutarate (αKG), succinate, fumarate, malate and citrate, in 10-230 BM cells treated with 10

10 μm, 100 μm and 50 mM Carnosine for 30 minutes are shown. (n = 3); data are shown as the

mean ± SD. Statistical analysis by one-way ANOVA. P value indicated. Cells were cultured with

[U-^{13}C5] glutamine for 6 h before metabolite extraction and gas chromatography-mass

spectrometry (GC-MS) analyses.

(H) Carnosine accumulation measured by ELISA represented as normalized values to carnosine

concentration in dialyzed media supplemented with indicated concentrations in 10-230 BM cells

cultured with 10 μm, 100 μm and 50 mM Carnosine concentrations during 0.5, 2 h. Mean of two

technical replicates shown. Results for 0.5 h also shown in Figure S2C.

(I) 2logFold Change of HMOX protein abundance measured by proteomics (MS/MS) in 10-230

BM cells transduced with indicated tetON-shRNAs. Differentially expressed proteins identified by

unpaired, two- tailed t-test comparing 2 groups: shCNDP1-1 and shCNDP1-2 versus shNTC.

(J) 2logFoldChange of *LCN2* transcript expression of both shCNDP1 tetON-shRNA vs shNTC in

10-230 BM and 12-230 BM cells. RNA sequencing data. p value < 0.05.

(K) 2logFold Change of *ATP7B* and *SLC46A3* transcript of Heavy Chain RNA sequencing of 10-

230 BM shCNDP1 vs shNTC. p value < 0.05.

(L) AUCell (Area Under the Curve) measurement of copper homeostasis gene signature

(WikiPaths, WP3286) in *CNDP1* high expression versus *CNDP1* low expression in MBM cells

from Biermann et al., 2022 data mining. Statistical analysis by one-way ANOVA**,** p<0.001.

(M) Representative immunoblots of OGDH, PDH and housekeeping (HK) protein β-actin in lysates

of 10-230 BM cells transfected with indicated sh-RNAs. See document S1 for details.

(N) R Pearson correlation between melanoma cell lines classified by their MITF expression and

resistance to Elesclomol treatment expressed as AUC of cell viability after treatment. Data mining

from Depmap.

1801

1802   **Figure S7. Graphical representation of CNDP1 role in brain metastasis.**

1803

1804

1805

1806

1807

1808

1809

1810

1811

1812

1813

1814

Supplementary information.

Document S1. Raw western blot images

Supplementary tables.

Table S1. Extracranial (NBM) and Brain metastasis (BM) samples included in analysis. Related

to Figure 1

Table S2. Extracranial (NBM) and Brain metastasis (BM) samples integration results. Significantly

changing candidates shown. Related to Figure 1

Table S3. Tissue microarray codified tumor sites. Related to Figure 1

Table S4. CNDP1 normalized counts from RNA-sequencing of melanoma extended to the lymph

node. Metastasis follow up information. Related to Figure S1

Table S5. Significantly differentially produced metabolites in CNDP1 KD (shCNDP1-1 and

shCNDP1-2) vs control cells (shNTC) in 10-230 BM cells after 72 hours of doxycycline treatment

(LC-MS, Hybrid Metabolomics). Related to Figure 4

Table S6. Differentially expressed genes (DEGs) in CNDP1 KD (shCNDP1-1 and shCNDP1-2)

vs control cells (shNTC) in 10-230 and 12-273 BM cells after 72 hours of doxycycline treatment

(RNA-sequencing). Related to Figure 4

Table S7. MetaboAnalyst analysis 10230 BM CNDP1 KD. Related to Figure 4

Table S8. Differentially expressed proteins in CNDP1 KD (shCNDP1-1 and shCNDP1-2) vs

control cells (shNTC) in 10-230 after 72 hours of doxycycline treatment (LC-MS/MS). Related to

Figure 4

Table S9. Gene Ontology analysis (GO) from differentially expressed genes (DEGs) in CNDP1

KD (shCNDP1-1 and shCNDP1-2) vs control cells (shNTC) in 10-230 and 12-273 BM cells after

72 hours of doxycycline treatment (RNA-sequencing)

Table S10. DAVID analysis 10230 BM CNDP1 KD LC-MS/M. Related to Figure 4S

Table S11. Differentially expressed transcripts in the heavy chain polysomes of CNDP1 KD

(shCNDP1-1 and shCNDP1-2) vs control cells (shNTC) in 10-230 and 12-273 BM after 72 hours

of doxycycline treatment (RNA sequencing). Related to Figure 5 and S5

Table S12. Differential gene expression and gene ontology of CNDP1 high versus CNDP1 low

expression cells from Biermann 2022. Related to Figure 5.

Materials. All materials used in this manuscript are referenced by catalog number in this file.

1845

1846