Silicon regulation of pectin methyl-esterification and cellulose microfibril organization modulates FERONIA-mediated cell wall integrity signaling for rice adaptation to salt stress

Yuyan Lei[a, b, 1], Yuan Liu[a, b, 1], Jing Wei[a, b, 1], Wenbing Li[b], Shaoshan Zhang[c], Zhengming Yang[a], Jingqiu Feng[c], Ying Li[c], Huachun Sheng [a, b, *]

[a]Institute of Qinghai-Tibetan Plateau, Southwest Minzu University, Chengdu, Sichuan, 610225, PR China

[b]Sichuan Provincial Qiang-Yi Medicinal Resources Protection and Utilization Technology and Engineering Laboratory, Southwest Minzu University, Chengdu, Sichuan 610225, PR China

[c]Tibetan Plateau Ethnic Medicinal Resources Protection and Utilization Key Laboratory of National Ethnic Affairs Commission of the People's Republic of China, Southwest Minzu University, Chengdu, Sichuan, 610225, PR China

[1]**Co-first Authors**:

These authors should be considered joint first author.

*Corresponding Authors:

Huachun Sheng, Email: huachunsheng99@126.com, Orcid-ID: 0000-0002-4684-5055

Abstract

The beneficial element silicon (Si) plays a crucial role in mitigating salt stress in

plants and can activate intracellular signal transduction pathways that reprogram the

transcriptome of salt-stressed plants. However, the mechanisms by which plants

perceive Si nutrition, induce the expression of Si-responsive genes, and regulate the

activity of Si-responsive proteins remain inadequately understood. In this study, we

found that Si enhances cell wall integrity by increasing the degree of pectin

methyl-esterification and changing the cellulose microfibril arrangement and

functions as a regulatory switch for the activities of cell wall integrity sensors (FER

homologs OsFLR1 and OsFLR2) during rice plant adaptation to high salinity.

Furthermore, the inhibition of OsFLR1 and OsFLR2 through FER-specific inhibitors

could negate the beneficial effects of Si and affect the uptake and accumulation of Si

in rice seedlings. In conclusion, these findings suggest that Si signaling initiation

involves FER-mediated cell wall integrity signaling pathways during salt stress

adaptation in rice.

Keywords: Si signaling initiation, Cell wall integrity, FER, pectin

methyl-esterification, Salt stress

1. Introduction

The plant cell walls, composed of cellulose, hemicellulose, pectin, lignin, and structural proteins, provide a physical barrier that separates plant cells from their environment. Consequently, they protect plant cells against environmental stresses such as high salinity, and damage to cell wall integrity results in decreased salt stress tolerance [1, 2]. For instance, arabinose serves as a structural unit in several cell wall components and glycoproteins, and mutations in the UDP-arabinose synthesis-related gene *MUR4* render plants hypersensitive to salt stress due to an intrinsic mechanism whereby polysaccharides containing arabinose residues are essential for maintaining cell wall integrity under high-salinity conditions [3]. Additionally, defects in cellulose synthesis and excessive accumulation of β-1,4-galactan within the cell wall disrupt cell wall integrity and further diminish plant salt tolerance [4, 5].

Furthermore, cell wall integrity-sensing mechanisms have been proposed and several families of receptor kinases are identified as cell wall integrity sensors in plants, such as the *Catharanthus roseus* receptor-like kinase 1-like (CrRLK1L) family, wall-associated kinases (WAKs), and leucine-rich repeat receptor-like kinases (LRR-RLKs) [6, 7]. FERONIA (FER), a member of CrRLK1L family in Arabidopsis, is a well-known cell wall integrity sensor involved in plant tolerance to high salinity, which can interact directly with the stress-induced de-methyl-esterified pectic polysaccharides and then transduce the environmental cues to intracellular signals [8]. Very recently, it was proposed that FER mediated-phosphorylation of companion of cellulose synthase1 (CC1) controls microtubule array behavior in response to salt stress, by modulating CC1 trafficking and affecting the ability of CC1 to engage with microtubules [9]. There are two FER homologs with the highest homology in rice plants, OsFLR1 and OsFLR2, involved in root immunity and plant architecture [10, 11]. To date, no studies have shown that they are related to the rice response to salt stress.

All terrestrial plants contain a certain amount of silicon (Si), with most Si deposited in the epidermal cell walls and silica cells as hydrated SiO□ [12] while trace amounts of Si covalently crosslink with cell wall components as organosilicon [13, 14]. Si

association and interaction with plant cell wall can alter the mechanical properties,

surface potential and composition of plant cell wall [14-16]. Rice is a typical Si

accumulator, with Si content reaching up to 10% of its dry weight, and it is also one

of the most salt-sensitive cereals [17]. In salt-stressed rice plants, Si deposition into

the root cell wall and Si-enhanced formation of Casparian strips can effectively block

the apoplastic bypass flow of Na^+, thereby reducing shoot Na^+ concentration [18-20].

This suggests that Si alleviates ionic toxicity through an apoplastic obstruction

mechanism. Additionally, Si nutrition stimulates the Salt-Overly Sensitive (SOS)

pathway in the SOS1-dependent manner, further mitigating ionic toxicity in rice [21].

Beyond Na^+ toxicity, salinity-induced osmotic stress can also be mitigated by Si

regulation of root morphological traits and root osmotic potential, as well as Si

promotion of transpiration force in the shoot [22]. Furthermore, Si increases the

activity of anti-oxidative enzymes, which scavenge reactive oxygen species (ROS),

leading to reduced oxidative damage and enhanced photosynthetic rates [22]. Kim et

al. (2014) reported that prolonged exposure to increased Si concentrations can

mitigate salinity-induced stress through the modulation of phytohormonal responses

[23].

To sum up, Si plays a significant role in alleviating salt stress in plants by

regulating ion and ROS homeostasis, enhancing root hydraulic conductance, and

improving the metabolism of osmoregulatory substances and phytohormones, as well

as increasing photosynthetic rates and the physicochemical properties of cell walls [24,

25]. Literature reviews indicate that the integration of Si with ROS metabolism is part

of a signaling cascade that regulates phytohormone biosynthesis and morphological,

biochemical, and molecular responses [25, 26]. However, how plants sense Si

nutrition, induce the expression of Si-responsive genes, and regulate the activity of

Si-responsive proteins remains poorly understood. Given the role of Si in modifying

plant cell walls and the importance of cell wall integrity in salt tolerance, we

hypothesize that Si signaling may initiate cell wall integrity signaling pathways

during salt stress adaptation in rice. In this study, we provide evidence that Si

regulates pectin methyl-esterification and cellulose microfibril organization to

enhance cell wall integrity and influences FER-mediated cell wall integrity signaling

to mitigate salt stress in rice plants.

2. Materials and Methods

2.1 Plant material and growth conditions

The japonica rice (*Oryza sativa* L.) variety Zhonghua 11 was utilized in this study.

Rice seeds were initially treated with 75% ethanol for 2 minutes, followed by

disinfection with 1% NaClO for 30 min, and finally immersed in deionized water for

an additional 30 min. Subsequently, the seeds were soaked in deionized water and

incubated in a constant temperature chamber at 30°C for two days to promote

germination. Following seed germination, rice seedlings exhibiting consistent growth

were selected and cultivated in Yoshida solution for 5 days before being transplanted

into fresh nutrient solution with various treatments. The concentrations of Si

($Na_2SiO_3·9H_2O$, Sigma-Aldrich, St. Louis, MO, USA), NaCl, and FER inhibitors

(reversine and staurosporine; AbMole, Houston, TX, USA) were set at 1.5 mM, 150

mM, and 5 μM respectively. The rice seedlings were grown in an artificial climate

chamber under the following conditions: a light/dark cycle of 16 hours/8 hours;

relative humidity of 65%; day/night temperatures of 30°C/22°C; and light intensity of

450 μmol m□² s□¹.

2.2 Root cell wall chemistry analysis

For cell wall separation, 5-day-old rice seedlings were transplanted into fresh

Yoshida solution containing Si, NaCl, or a combination of both. Four days later, the

roots of the rice plants were harvested and ground to a powder in liquid nitrogen. The

samples were vacuum-dried overnight and subsequently subjected to successive

washes with 70% ethanol, a 1:1 (v/v) chloroform/methanol mixture, and acetone, each

followed by centrifugation at 10,000 rpm for 10 minutes. After another round of

vacuum drying, the cell wall extracts were prepared for subsequent analysis.

Fourier-transform infrared microscopy. The FT-IR spectra were obtained using a

Thermo Nicolet iN10 FTIR spectrometer, utilizing cell wall extracts (100 mg)

prepared by compressing a blend of fine dust with potassium bromide (KBr; 100 mg)

powder. The wave number data derived from FTIR spectroscopy of cell walls, along

with their corresponding assignments to the primary components of the cell wall,

were presented in Table S1 [27].

Determination of the Content of Crystalline Cellulose. 2 mg of cell wall powder

were resuspended in 400 µL of a 2 M trifluoroacetic acid (TFA) solution and

subjected to incubation at 121°C for a duration of 90 minutes. The material that

remained insoluble in TFA was subsequently isolated, dried, and then redispersed in 1

mL of Updegraff reagent. After being incubated at 100°C for 30 minutes in boiling

water, the samples underwent centrifugation at a force of 2500 g for 5 minutes. The

residual crystalline cellulose was then dissolved in a solution containing 72% (v/v)

H_2SO_4. Finally, the obtained crystalline cellulose was quantified colorimetrically

using anthrone reagent at a wavelength of 625 nm with the aid of a

spectrophotometer.

Pectin Content Measurements. To extract pectin, 2 mg of the cell wall extract was

boiled with 1 mL of ultrapure water three times for 1 hour each. After each boiling,

the supernatant was centrifuged at 16,800 g for 3 minutes and collected in a single

tube. Galacturonic acid served as the standard to quantify the pectin content. A

volume of 50 µL of the aforementioned pectin extract was added to 250 µL of 98%

H_2SO_4 containing 0.0125 M $Na_2B_4O_7$ and incubated at 100°C for 5 min in boiling

water. Once cooled to room temperature, 5 µL of a solution containing 0.15% (w/v)

meta-hydroxybiphenyl dissolved in 0.5% NaOH was added to the samples and

incubated at room temperature for an additional 20 minutes. The absorbance of these

samples at a wavelength of 520 nm was measured using a microplate reader.

Determination of the degree of pectin methyl-esterification. 1 mg of cell wall

extracts were initially washed with ultrapure water and centrifuged at 16,800 g for 10

minutes at room temperature to remove the supernatant. The samples were then

incubated with 200 µL of NaOH at room temperature for 1 hour for pectin

de-methyl-esterification, followed by adding 200 µL of 1 M HCl to stop the process.

Subsequently, 125 µL of the reaction mixture was extracted and combined with 125

µL of a 20 mM HEPES solution (pH 7.5). Then, 250 µL of a HEPES solution

containing 0.03 units of ethanol oxidase (Sigma-Aldrich) was added and incubated at

room temperature for 15 minutes. After the reaction was done, an additional 250 µL of

assay reagent (comprising 20 mM acetoacetone, 50 mM acetic acid, and 2 M

ammonium acetate) was introduced and incubated at 60 °C for 15 minutes. After

cooling to room temperature, absorbance at 412 nm was measured using a microplate

reader. Methanol was used to construct a calibration curve to quantify

methyl-esterification levels.

Hemicellulose Content Measurements. Hemicellulose extraction was performed

on the residue obtained from pectin extraction. An aqueous solution containing 4%

KOH and 0.02% KBH_4 (1 mL) was added to the residue, and the resulting suspension

was incubated at room temperature for 12 hours. Subsequently, the supernatant was

collected by centrifugation at 16,800 g for 10 minutes at room temperature. This

procedure was repeated once more, and the extracts from both rounds were combined

to yield the hemicellulose extract. The hemicellulose content was quantified using

glucose as a standard via the phenol-sulfuric acid method. To this end, 1 mL of 98%

H_2SO_4 and 10 µL of an 80% phenol solution were added to 200 µL of the

hemicellulose extract. The samples were initially incubated at room temperature for

15 minutes before being boiled for another 15 minutes. After cooling to room

temperature, absorbance measurements of the samples were taken at a wavelength of

490 nm using a microplate reader.

2.3 Cell wall integrity assay

Propidium iodide (PI) staining. 5-day-old rice seedlings cultivated on Yoshida

medium with or without 1.5 mM Si were subsequently transferred to a medium

enriched with 150 mM NaCl. Following a growth period of 16 hours, the roots were

stained with PI prior to cell integrity evaluation using confocal microscopy (Zeiss,

Oberkochen, Germany). Cell death was characterized by complete staining with PI.

The number of burst cells in the elongation zone of the roots was quantified.

Atomic force microscopy. The protocol for atomic force microscopy (AFM) was

adapted from the methodologies established by He et al. (2015) [28] and Pu et al.
(2021) [29]. The cell wall powder was re-suspended in ultrapure water and
subsequently deposited onto a poly-l-lysine-coated mica substrate. AFM imaging was
conducted using the NanoScope 8 atomic force microscope (Bruker, Santa Barbara,
CA) in ScanAsyst-Air mode. To assess mechanical properties, representative
Force-Distance (F-D) curves were acquired and analyzed with Nanoscope analysis
software version 1.9 (Bruker, Santa Barbara, CA). Young's modulus was determined
utilizing the Sneddon model as described by Sneddon (1965) [30].

2.4 RNA Isolation and Quantitative Real-Time PCR Analysis

5-day-old rice seedlings were transferred to new nutrient solutions containing
various treatments. For RNA extraction, samples were collected at 12-hour and
24-hour intervals following the treatment. Total RNA was extracted using the RNeasy
Plant Mini Kit (QIAGEN, Hilden, Germany). Complementary DNA (cDNA) was
synthesized (Invitrogen, Carlsbad, CA, USA) and subsequently utilized as a template
for quantitative real-time PCR (qRT-PCR), conducted on a CFX96 Real-Time System
(Bio-Rad Laboratories, Hercules, CA, USA) in conjunction with the SYBR Premix
Ex Taq II Kit (Takara Bio, Otsu, Japan). The rice *Histone H3* gene served as an
internal control for normalizing the expression levels of target genes. The sequences
of primers used for amplification in qRT-PCR analyses are provided in Table S2.

2.5 Si content measurements

100 mg of dried rice seedlings were ground into a fine powder, placed in a Teflon
crucible, and combined with 6 mL of NaClO and 2 mL of 2 M NaOH. The crucible
was subsequently heated at 150°C for 30 minutes to facilitate digestion of the sample.
After cooling, the resulting solution was transferred to a 250 mL volumetric flask
containing 9 mL of 1 M H_2SO_4 and 100 mL of distilled water, followed by dilution
to the mark. A volume of 1.5 mL from this solution was then transferred to a separate
50 mL volumetric flask and mixed with an additional 2 mL of 98% H_2SO_4 until
homogeneous. Five minutes later, another addition of distilled water (20 mL) was

made, followed by thorough mixing with 5 mL of 5.0% ammonium molybdate

solution and 2 mL of 0.5% ascorbic acid. The mixture stood for 20 minutes before

being diluted to the mark, after which absorbance measurements were conducted at

810 nm.

2.6 Analysis of ABA content

Rice seedling was homogenized using a glass rod. Subsequently, the homogenates

were each subjected to two rounds of methanol extraction, and the extracts were

pooled separately. The pooled extracts were then centrifuged. The supernatants were

passed through an HLB column, followed by passage through an MCX column. The

collected samples were evaporated to dryness and re-dissolved in 500 μL of

chromatographic-grade methanol for subsequent UPLC-MS/MS analysis (Shimadzu

UPLC-MS/MS 8050, Kyoto, Japan). Moreover, a concentration gradient of ABA

standard (5, 2, 1, 0.5, 0.2, 0.1, 0.05, 0.02, and 0.01 μg/mL) was prepared in

chromatographic-grade methanol. MS detection was performed in multiple reaction

monitoring (MRM) mode, and the compound-specific MRM parameters were set as

previously reported [31].

2.7 Statistical analysis

For the measurements of plant height and fresh weight, all experiments were

conducted at least in triplicate. The sample size, denoted as n, for each experiment is

provided in the respective figure legends. For root cell wall chemistry and integrity

analysis, qRT-PCR, and content measurements of Si and endogenous ABA, all

experiments were performed at least three times using independent biological

replicates, with consistent results observed across trials. All reported experimental

data represent means ± standard deviation (SD) and were statistically analyzed using

Welch's one-way ANOVA.

3.　Results

3.1 Si effects on pectin methyl-esterification and cellulose microfibril organization of root cell wall under salt stress condition

The effects of Si, NaCl, or their combination on cell wall chemistry were assessed using Fourier transform infrared (FTIR) spectroscopy and chemical methods. The highest peaks for cellulose (approximately 1050 cm^{-1}) were observed in rice plants across all treatment conditions, indicating elevated cellulose levels (Fig. 1A). No significant differences were detected at the wavenumbers corresponding to hemicelluloses (between 1161 and 1442 cm^{-1}), pectin (between 950 and 990 cm^{-1}), and esterified uronic acid (1712 and 1735 cm^{-1}) under the three treatment conditions (Fig. 1A). Specific measurements of hemicellulose, and pectin largely exhibited a similar trend as the FTIR results (Fig. 1C-D), and the cellulose content rose slightly but without statistical significance (Fig. 1B). Notably, Si appeared to mitigate the NaCl-induced de-methyl-esterification of pectin (Fig. 1E).

To further investigate the Si effects on cell wall integrity under salt stress conditions, NaCl-induced cell bursting and cell wall softening were detected by PI staining and AFM, respectively. It was observed that the addition of Si could reduce the NaCl-induced cell bursting (Fig. 2A, B). Moreover, the AFM results indicated that Si-modified cell walls possess denser cellulose microfibrils (Fig. 2C) and enhanced mechanical properties under high salinity conditions (Fig. 2D). According to the above, Si could maintain the cell wall integrity under salt stress conditions by increasing the methyl-esterification degree of pectin and changing the cellulose microfibril arrangement.

3.2 Si and NaCl influence the expression of FER homolog genes

In rice plants, OsFLR1 and OsFLR2 are the homologs of FER (Fig. 3A). Salt stress reduces the activity of FER kinase [32], while the expression of the FER gene is upregulated to sustain its activity in plant response to high salinity [33]. This also applies to its homologous genes *OsFLR1* and *OsFLR2* in rice (Fig. 3B, C), suggesting that the FER homologs OsFLR1 and OsFLR2 also play a role in salt stress responses. Notably, Si enhances the NaCl-induced expression of *OsFLR1* and *OsFLR2* genes

during the early stages when rice plants are exposed to high salinity within 12 hours

(defense phase), whereas after 24 hours of treatment (re-growth phase), Si suppresses

the NaCl-induced expressions of these FER homolog genes (Fig. 3B, C). Considering

that Si downregulates *OsFLR1* and *OsFLR2* gene expressions under normal

conditions, it can be concluded that Si plays a dual role in regulating the expression of

FER homolog genes, thereby mediating the trade-off between plant growth and

defense.

3.3 Si mitigation of salt stress depends on the activation of FER kinase

Under salt stress conditions, Si-mediated cell wall integrity maintenance may be

perceived by FER homologs. To test this hypothesis, reversine and staurosporine were

employed in this study as FER-specific inhibitors that can inactivate FER kinase and

its homologs by targeting the conserved ATP binding pocket within the kinase

structure [34]. The addition of reversine had no significant effects on plant growth

(Fig. 4A-C), whereas staurosporine exhibited inhibitory effects on both plant height

and fresh weight (Fig. 4G). Notably, Si was able to mitigate salt stress in rice

seedlings; however, its beneficial effects were negated by the application of both FER

inhibitors (Fig. 4D-J), suggesting that Si mitigation of salt stress is contingent upon

the activation of FER kinase.

3.4 FER kinases could regulate the uptake of Si

To investigate the interactions between Si and FER, we assessed the expression of

Si transporter genes (*OsLsi1* and *OsLsi2*) and Si uptake in rice seedlings treated with

NaCl, a FER inhibitor (reversine), or both. The expression levels of *OsLsi1* and

OsLsi2 genes were downregulated by treatment with NaCl or the FER inhibitor, with

a more pronounced effect observed when both treatments were applied

simultaneously (Fig. 5A, B). We also measured Si content in rice seedlings subjected

to NaCl, the FER inhibitor, or both and found that Si uptake was decreased under

these conditions (Fig. 5C). Collectively, these findings suggest that FER homologs

positively regulate the expression of Si transporter genes and the uptake of Si in rice

plants.

3.5 ABA biosynthesis is mediated by Si and FER kinases

Previous studies have reported that the expression of *OsLsi1* and *OsLsi2* genes

negatively correlates with endogenous abscisic acid (ABA) contents due to the

presence of ABA-responsive motifs in their promoter regions [35]. Therefore, we

detected the ABA contents under the normal and salt stress (Fig. 6) conditions. It has

been reported that salt stress and FER inactivation lead to increased ABA production

[33], and our results align with those previously reported. Interestingly, Si could

markedly enhance ABA biosynthesis under the normal condition (Fig. 6A) but inhibit

it under the long-term salt stress condition. Moreover, when FER is rendered inactive

by its specific inhibitor, the role of Si is conspicuously attenuated under the long-term

salt stress condition (Fig. 6B), suggesting Si regulation of ABA production in a FER

activation-dependent manner. And, these findings are in accordance with the results

shown in Fig. 4.

4.   Discussion

4.1   Si regulates pectin methyl-esterification and cellulose microfibril

organization for cell wall integrity maintenance under salt stress conditions

It is well established that the cell wall plays a critical role in plant defense against

environmental stresses including salt stress, and remodeling of the cell wall through

alterations in its composition and organization constitutes a vital adaptive strategy [1,

2, 36]. Prior to the recovery of growth rates in salt-stressed plants, the cell wall

particularly that of root epidermal cells, undergoes a transition from softening to

stiffening [8]. Initially, excess sodium ions (Na$^+$) can disrupt the Ca$^{2\square}$-crosslinked

pectin network by competing with Ca$^{2\square}$ for negative charges on galacturonic acid

residues [37], leading to a softened cell wall. To maintain cell wall integrity, pectic

polysaccharides are subsequently de-methyl-esterified through the activation of pectin

methylesterases (PMEs) in plants, which polysaccharides can then be cross-linked by

divalent cations to restore cell wall stiffness [38, 39].

The beneficial element Si plays a positive role in alleviating salt stress in plants [24,
25], and its association and interaction with cell wall components have been proposed
[13, 14]. Previous studies have regarded deposited silica as supplementary material
that strengthens plant cell walls [40]. Si also collaborates with lignin to enhance cell
wall stiffness: it can induce the biosynthesis of lignin, a component of the secondary
cell wall; conversely, lignin interacts with mono-silicic acid to regulate silica
deposition within the cell walls [41, 42]. Increased deposition of both lignin and silica
forms apoplastic obstruction that prevent Na$\square$ from entering cells [40]. In addition to
inorganic silica, our research group focuses more on cell wall-bound Si. Recently, Pu
et al. (2021) reported that xyloglucan serves as the ligand for bonding Si in rice cell
walls, participating in establishing and remodeling the cell wall architecture through
cross-linking with cellulose microfibrils, thereby contributing to increased stiffness
[29]. It has been demonstrated that Si-xyloglucan complexes not only enhance
resistance to enzymatic degradation and salt-alkali digestion but also improve the
nanomechanical properties of rice plant cell walls [13, 29].

In addition to lignin and the hemicellulose xyloglucan, pectic polysaccharides can
serve as silicification templates for SiO$\square$ deposition [43] and crosslink with
mono-silicic acid to form Si-pectin complexes [44]. Here, we aim to investigate the
relationship between salt stress adaptation and Si-pectin interactions in rice plants.
Our data indicate that Si application can counteract NaCl-induced
de-methyl-esterification of pectin (Fig. 1), thereby rearranging cellulose microfibril
organization and increasing root cell wall stiffness (Fig. 2) and suggesting that Si
maintains cell wall integrity in the cellulose- and pectin-dependent manner. Similar
results were observed in aluminum-stressed rice and lanthanum-stressed
Dicranopteris linearis: the addition of Si increases the degree of pectin
methyl-esterification and preserves cell wall integrity under stress conditions, thus
facilitating recovery of plant growth [45, 46].

4.2 The initiation of Si signaling involves FER-mediated cell wall integrity
signaling pathways in rice plants response to salt stress

Changes in cell wall composition and organization trigger the cell wall integrity

signaling pathway [6, 7]. FER is a plasma membrane-localized receptor kinase with

an extracellular ligand-binding domain that senses cell wall integrity, perceiving salt

stress-mediated cell wall damage through binding to de-methylesterified

homogalacturonan [8, 47]. In this study, we found that the FER homolog genes of

OsFLR1 and *OsFLR2* are induced by NaCl treatments (Fig. 3) and OsFLR1 and

OsFLR2 positively regulate the salt stress tolerance in rice seedlings (Fig. 4).

Moreover, recent reports indicate that the pectin-FER complex plays a role in plant

mechanical force perception and activates ROP6 GTPase signaling during

Arabidopsis pavement cell morphogenesis, suggesting that FER kinase also functions

as a mechanoreceptor [48].

Considering the effects of Si on pectin methyl-esterification and the mechanical

properties of rice plant cell walls, it appears that Si may influence FER-mediated cell

wall integrity signaling under stress conditions [49]. Indeed, when the activity of FER

kinase is inhibited by specific inhibitors, the beneficial effects of Si are abolished (Fig.

4), implying a direct link between Si and FER. Interestingly, Si acts as a regulatory

switch for the activities of FER homologs OsFLR1 and OsFLR2 (Fig. 3): Si can

enhance their activities during stress response stages and reduce them during growth

recovery periods. Herein we propose a hypothesis that Si signaling initiation involves

FER-mediated cell wall integrity signaling pathways during salt stress adaptation in

rice plants.

Very recently, a homolog of the flowering hormone "florigen" FT, specifically

FLOWERING LOCUS T-LIKE 12 (OsFTL12), has been identified as the

Shoot-Silicon-Signal protein [50]. This protein is predominantly expressed in the

shoot; however, under Si-deficient conditions, it also reaches the phloem and roots,

thereby inducing increased expression of transporters (OsLsi1 and OsLsi2)

responsible for Si uptake in roots [50]. It can be concluded that OsFTL12 is a

potential component of the Si signaling process but not the receptor of Si signaling,

because how low Si induces the expression of *OsFTL12* gene is still unknown. It is

worth noting that FER modulates the flowering time by regulating the expression of

FT in Arabidopsis and the lower peak in *FT* expression was observed in the *fer-4*

mutant [51]. By coincidence, the inactivation of FER homologs OsFLR1 and OsFLR2

down-regulates the expression levels of Si transporter genes *OsLsi1* and *OsLsi2*,

leading to decreased uptake and accumulation of Si in rice seedlings (Fig. 5). Similar

results were also observed in *OsFTL12* rice mutants [50], suggesting an underlying

association between OsFLR1/2 and OsFTL12.

Furthermore, the ABA content in *fer-4* mutants is significantly higher than that in

wild-type Arabidopsis plants under both control and salt stress conditions [33]. For Si

nutrition, its addition may inhibit ABA biosynthesis by down-regulating key ABA

biosynthesis-related genes, including *zeaxanthin epoxidase* (*ZEP*) and

9-cis-epoxycarotenoid dioxygenases (*NCED1* and *NCED4*), when rice plants are

exposed to salt stress for a long time (> 24 h) [23]. Given that NaCl-induced ABA

negatively regulates the expression of Si transporters [35], ABA levels were measured

under various conditions in this study (Fig. 6). Our results demonstrate that both Si

and FER are requirement for ABA homeostasis, thereby promoting the growth

recovery of salt-stressed rice plants. Moreover, the mitigation of NaCl-induced ABA

activity by Si depends on the activation of FER kinases (Fig. 6B), which supports the

involvement of OsFLR1/2 in initiating Si signaling, regulating the expression of Si

transporters, and facilitating the uptake and accumulation of Si.

As early as 2020, we were among the first to propose that Si alternation of the cell

wall — encompassing changes in its composition and mechanical properties —

coordinates with cell wall integrity signaling to dynamically regulate plant growth and

development in response to environmental changes [14]. This proposition highlighted

the potential role of cell wall integrity receptors in Si sensing. Subsequently, several

colleagues concurred with our hypothesis and further suggested that

mechanoreception molecules on the cell membrane may function as Si sensing

receptors [52]. Prior to this study, there was no direct evidence supporting these

hypotheses. Our current findings represent a significant advancement in our

fundamental understanding of Si signaling initiation. Future research will likely focus

on whether other cell wall integrity receptors, such as WAKs, are involved in Si signaling transduction. Additionally, while we have conducted preliminary studies on the relationship between Si nutrition and OsFLR1/2, further investigation is required to elucidate the underlying mechanisms by which OsFLR1/2 functions in Si sensing.

5. Conclusions

Our study demonstrates that Si functions as an enhancer of cell wall integrity by regulating pectin methyl-esterification and cellulose microfibril arrangement under salt stress conditions. This regulation modulates the FER homologs OsFLR1/2-mediated cell wall integrity signaling pathway (Fig. 7). Furthermore, Si plays a dual role in modulating the activity of OsFLR1/2 under different stage during salt stress response. In turn, OsFLR1/2 regulates the expression of Si transporter genes, thereby influencing the subsequent uptake and accumulation of Si in salt-stressed rice plants. These findings suggest the existence of a feedback loop between Si nutrition and FER kinase activity, implying the involvement of FLR1/2 in initiating Si signaling.

Acknowledgements

This work was supported by the Key Research and Development Program of Sichuan Province (Grant No. 2024YFFK0190), the first batch of Scientific and Technological Innovation Team for Qinghai-Tibetan Plateau Research in Southwest Minzu University (Grant No. 2024CXTD04) and the Southwest Minzu University Research Startup Funds (Grant No. RQD2023019).

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Table S1 Some FT-IR wavenumbers from the main plant cell wall components.

Table S2 Primers used in this study.

References

[1] L. Colin, F. Ruhnow, J. K. Zhu, C. Zhao, Y. Zhao, S. Persson, The cell biology of primary cell walls during salt stress, Plant Cell 35 (2023) 201-217. https://doi.org/10.1093/plcell/koac292

[2] W. Li, J. Wei, Y. Lei, Z. Yang, S. Zhang, J. Feng, Y. Li, Y. Liu, H. Sheng, Phosphorylation of cellulose synthases in plant responses to environmental changes, Int. J. Biol. Macromol. 292 (2025) 139313. https://doi.org/10.1016/j.ijbiomac.2024.139313

[3] C. Zhao, O. Zayed, F. Zeng, C. Liu, L. Zhang, P. Zhu, C. C. Hsu, Y. E. Tuncil, W. A. Tao, N. C. Carpita, J. K. Zhu, Arabinose biosynthesis is critical for salt stress tolerance in Arabidopsis, New Phytol. 224 (2019) 274-290. https://doi.org/10.1111/nph.15867

[4] A. Endler, C. Kesten, R. Schneider, Y. Zhang, A. Ivakov, A. Froehlich, N. Funke, S. Persson, A mechanism for sustained cellulose synthesis during salt stress, Cell 162 (2015) 1353-1364. https://doi.org/10.1016/j.cell.2015.08.028

[5] J. Yan, Y. Liu, L. Yang, H. He, Y. Huang, L. Fang, H. V. Scheller, M. Jiang, A. Zhang, Cell wall β-1, 4-galactan regulated by the BPC1/BPC2-GALS1 module aggravates salt sensitivity in *Arabidopsis thaliana*, Mol. Plant 14 (2021) 411-425. https://doi.org/10.1016/j.molp.2020.11.023

[6] Y. Rui, J. R. Dinneny, A wall with integrity: surveillance and maintenance of the plant cell wall under stress, New Phytol. 225 (2020) 1428-1439. https://doi.org/10.1111/nph.16166

[7] S. Wolf, Cell wall signaling in plant development and defense, Ann. Rev. Plant Biol. 73 (2022) 323-353. https://doi.org/10.1146/annurev-arplant-102820-095312

[8] W. Feng, D. Kita, A. Peaucelle, H. N. Cartwright, V. Doan, Q. Duan, M. C. Liu, J. Maman, L. Steinhorst, I. Schmitz-Thom, R. Yvon, J. Kudla, H. M. Wu, A. Y. Cheung, J. R. Dinneny, The FERONIA receptor kinase maintains cell-wall integrity during salt stress through Ca^{2+} signaling, Curr. Biol. 28 (2018) 666-675. https://doi.org/10.1016/j.cub.2018.01.023

[9] X. Liu, L. Wang, L. Liu, Y. Li, M. Ogden, M. Somssich, Y. Liu, Y. Zhang, M. Ran,

S. Persson, C. Zhao, FERONIA adjusts CC1 phosphorylation to control microtubule

array behavior in response to salt stress, Sci. Adv. 10 (2024) eadq8717.

https://doi.org/10.1126/sciadv.adq8717

[10] Z. Yang, J. Jiang, L. Wang, Y. Liu, J. Qu, Y. Tan, X. Fu, Q. Lin, H. Deng, F. Yu,

Mutations of two FERONIA-like receptor genes enhance rice blast resistance without

growth penalty, J. Exp. Bot. 71 (2020) 2112-2126. https://doi.org/10.1093/jxb/erz541

[11] C. Li, L. Wang, Y. Cui, L. He, Y. Qi, J. Zhang, J. Lin, H. Liao, Q. Lin, T. Yang, F.

Yu, X. Liu, Two FERONIA-like receptor (FLR) genes are required to maintain

architecture, fertility, and seed yield in rice, Mol. Breeding 36 (2016) 1-11.

https://doi.org/10.1007/s11032-016-0580-x

[12] J. F. Ma, N. Yamaji, Silicon uptake and accumulation in higher plants, Trends

Plant Sci. 11 (2006) 392-397. https://doi.org/10.1016/j.tplants.2006.06.007

[13] H. Sheng, J. Ma, J. Pu, L. Wang, Cell wall-bound silicon optimizes ammonium

uptake and metabolism in rice cells, Ann. Bot. 122 (2018) 303-313.

https://doi.org/10.1093/aob/mcy068

[14] H. Sheng, S. Chen, Plant silicon-cell wall complexes: Identification, model of

covalent bond formation and biofunction, Plant Physiol. Biochem. 155 (2020) 13-19.

https://doi.org/10.1016/j.plaphy.2020.07.020

[15] J. Cui, Y. Li, Q. Jin, F. Li, Silica nanoparticles inhibit arsenic uptake into rice

suspension cells via improving pectin synthesis and the mechanical force of the cell

wall, Environ Sci: Nano 7 (2020) 162-171. https://doi.org/10.1039/C9EN01035A

[16] J. Ma, H. Cai, C. He, W. Zhang, L. Wang, A hemicellulose-bound form of silicon

inhibits cadmium ion uptake in rice (*Oryza sativa*) cells. New Phytol. 206 (2015)

1063-1074. https://doi.org/10.1111/nph.13276

[17] S. J. Thorne, P. M. Stirnberg, S. E. Hartley, F. J. Maathuis, The ability of silicon

fertilisation to alleviate salinity stress in rice is critically dependent on cultivar, Rice

15 (2022) 8. https://doi.org/10.1186/s12284-022-00555-7

[18] H. J. Gong, D. P. Randall, T. J. Flowers, Silicon deposition in the root reduces

sodium uptake in rice (*Oryza sativa* L.) seedlings by reducing bypass flow, Plant Cell

Environ. 29 (2006) 1970-1979. https://doi.org/10.1111/j.1365-3040

[19] R. Flam-Shepherd, W. Q. Huynh, D. Coskun, A. M. Hamam, D. T. Britto, H. J.

Kronzucker, Membrane fluxes, bypass flows, and sodium stress in rice: the influence

of silicon, J. Exp. Bot. 69 (2018) 1679-1692. https://doi.org/10.1093/jxb/erx460

[20] G. Yan, X. Fan, L. Tan, C. Yin, T. Li, Y. Liang, Root silicon deposition and its

resultant reduction of sodium bypass flow is modulated by OsLsi1 and OsLsi2 in rice,

Plant   Physiol.   Biochem.   158   (2021)   219-227.

https://doi.org/10.1016/j.plaphy.2020.11.015

[21] B.K. Gupta, K.K. Sahoo, K. Anwar, R.C. Nongpiur, R. Deshmukh, A. Pareek,

S.L. Singla-Pareek, Silicon nutrition stimulates Salt-Overly Sensitive (SOS) pathway

to enhance salinity stress tolerance and yield in rice, Plant Physiol. Biochem. 166

(2021) 593-604. https://doi.org/10.1016/j.plaphy.2021.06.010

[22] G. Yan, X. Fan, M. Peng, C. Yin, Z. Xiao, Y. Liang, Silicon improves rice salinity

resistance by alleviating ionic toxicity and osmotic constraint in an organ-specific

pattern, Front. Plant Sci. 11 (2020) 260. https://doi.org/10.3389/fpls.2020.00260

[23] Y. H. Kim, A. L. Khan, M. Waqas, J. K. Shim, D. H. Kim, K. Y. Lee, I. J. Lee,

Silicon application to rice root zone influenced the phytohormonal and antioxidant

responses under salinity stress, J. Plant Growth Regul. 33 (2014) 137-149.

https://doi.org/10.1007/s00344-013-9356-2

[24] Y. Zhu, H. J. Gong, J. L. Yin, Role of silicon in mediating salt tolerance in plants:

a review, Plants 8 (2019) 147. https://doi.org/10.3390/plants8060147

[25] P. Dhiman, N. Rajora, S. Bhardwaj, S. S. Sudhakaran, A. Kumar, G. Raturi, K.

Chakraborty, O. P. Gupta, B. N. Devanna, D. K. Tripathi, R. Deshmukh, Fascinating

role of silicon to combat salinity stress in plants: An updated overview, Plant Physiol.

Biochem. 162 (2021) 110-123. https://doi.org/10.1016/j.plaphy.2021.02.023

[26] D.K. Tripathi, K. Vishwakarma, V.P. Singh, V. Prakash, S. Sharma, S. Muneer, M. Miroslav Nikolic, R. Deshmukh, M. Vaculík, F.J. Corpas, Silicon crosstalk with reactive oxygen species, phytohormones and other signaling molecules, J. Hazard Mater. 408 (2021) 124820. https://doi.org/10.1016/j.jhazmat.2020.124820

[27] X. Liu, H. Cui, B. Zhang, M. Song, S. Chen, C. Xiao, Y. Tang, J. Liesche, Reduced pectin content of cell walls prevents stress-induced root cell elongation in Arabidopsis, J. Exp. Bot. 72 (2021) 1073-1084. https://doi.org/10.1093/jxb/eraa533

[28] C. He, J. Ma, L. Wang, A hemicellulose-bound form of silicon with potential to improve the mechanical properties and regeneration of the cell wall of rice, New Phytol. 206 (2015) 1051-1062. https://doi.org/10.1111/nph.13282

[29] J. Pu, L. Wang, W. Zhang, J. Ma, X. Zhang, C. V. Putnis, Organically-bound silicon enhances resistance to enzymatic degradation and nanomechanical properties of rice plant cell walls, Carbohyd. Polym. 266 (2021)118057. https://doi.org/10.1016/j.carbpol.2021.118057

[30] I. N. Sneddon, The relation between load and penetration in the axisymmetric Boussinesq problem for a punch of arbitrary profile, Int. J. Eng. Sci. 3 (1965) 47-57. https://doi.org/10.1016/0020-7225(65)90019-4

[31] J. Huang, J. Hua, L. Peng, L. Bai, S. Luo, The Diterpene Isopimaric Acid Modulates the Phytohormone Pathway to Promote Oryza sativa L. Rice Seedling Growth, Curr. Issues Mol. Biol. 46 (2024) 9772-9784. https://doi.org/10.3390/cimb46090580

[32] X. Liu, W. Jiang, Y. Li, H. Nie, L. Cui, R. Li, L. Tan, L. Peng, C. Li, J. Luo, M. Li, H, Wang, J, Yang, B. Zhou, P. Wang, H. Liu, J. K. Zhu, C. Zhao, FERONIA coordinates plant growth and salt tolerance via the phosphorylation of phyB, Nat. Plants 9 (2023) 645-660. https://doi.org/10.1038/s41477-023-01390-4

[33] C. Zhao, W. Jiang, O. Zayed, X. Liu, K. Tang, W. Nie, Y. Li, S. Xie, Y. Li, T. Long, L. Liu, Y. Zhu, Y. Zhao, J. K. Zhu, The LRXs-RALFs-FER module controls plant growth and salt stress responses by modulating multiple plant hormones, Natl. Sci. Rev. 8 (2021) nwaa149. https://doi.org/10.1093/nsr/nwaa149

[34] H. B. Liu, X. Li, J. Cai, L. L. Jiang, X. Zhang, D. Wu, L. Wang, A. Yang, C. Guo, J. Chen, W. Pu, F. Yu, A screening of inhibitors targeting the receptor kinase FERONIA reveals small molecules that enhance plant root immunity, Plant Biotech. J. 21 (2023) 63-77. https://doi.org/10.1111/pbi.13925

[35] M. E. F. Abdel-Haliem, H. S. Hegazy, N. S. Hassan, D. M. Naguib, Effect of silica ions and nano silica on rice plants under salinity stress, Ecol. Eng. 99 (2017) 282-289. https://doi.org/10.1016/j.ecoleng.2016.11.060

[36] D. M. Oliveira, T. R. Mota, F. V. Salatta, et al., Cell wall remodeling under salt stress: Insights into changes in polysaccharides, feruloylation, lignification, and phenolic metabolism in maize, Plant Cell Environ. 43 (2020) 2172-2191. https://doi.org/10.1111/pce.13805

[37] C. S. Byrt, R. Munns, R. A. Burton, M. Gilliham, S. Wege, Root cell wall solutions for crop plants in saline soils, Plant Sci. 269 (2018) 47-55. https://doi.org/10.1016/j.plantsci.2017.12.012

[38] L. Hocq, J. Pelloux, V. Lefebvre, Connecting homogalacturonan-type pectin remodeling to acid growth, Trends Plant Sci. 22 (2017) 20-29. https://doi.org/10.1016/j.tplants.2016.10.009

[39] T, Zhou, P. J. Wu, J. F. Chen, X. Q. Du, Y. N. Feng, Y. P. Hua, Pectin demethylation-mediated cell wall Na^+ retention positively regulates salt stress tolerance in oilseed rape, Theor. Appl. Genet. 137 (2024) 54. https://doi.org/10.1007/s00122-024-04560-w

[40] D. Coskun, R. Deshmukh, H. Sonah, J. G. Menzies, O. Reynolds, J. F. Ma, H. J. Kronzucker, R. R. Bélanger, The controversies of silicon's role in plant biology, New Phytol. 221 (2019) 67-85. https://doi.org/10.1111/nph.15343

[41] N. Zexer, R. Elbaum, Unique lignin modifications pattern the nucleation of silica in sorghum endodermis, J. Exp. Bot. 71 (2020) 6818-6829. https://doi.org/10.1093/jxb/eraa127

[42] H. Sheng, Y. Li, J. Feng, Y. Liu, Regulation of thermodynamics and kinetics of silica nucleation during the silicification process in higher plants, Plant Physiol. Biochem. 198 (2023) 107674. https://doi.org/10.1016/j.plaphy.2023.107674

[43] O. Leroux, F. Leroux, A. A. Mastroberti, F. Santos-Silva, D. Van Loo, A. Bagniewska-Zadworna, L. V. Hoorebeke, S. Bals, Z. A. Popper, J. E. de Araujo Mariath, Heterogeneity of silica and glycan-epitope distribution in epidermal idioblast cell walls in *Adiantum raddianum laminae*, Planta 237 (2013) 1453-1464. https://doi.org/10.1007/s00425-013-1856-6

[44] H. Sheng, Y. Lei, J. Wei, Z. Yang, L. Peng, W. Li, Y. Liu, Analogy of silicon and boron in plant nutrition, Front. Plant Sci. 15 (2024) 1353706. https://doi.org/10.3389/fpls.2024.1353706

[45] Z. Xiao, G. Yan, M. Ye, Y. Liang, Silicon relieves aluminum-induced inhibition of cell elongation in rice root apex by reducing the deposition of aluminum in the cell wall, Plant Soil 462 (2021) 189-205. https://doi.org/10.1007/s11104-021-04850-y

[46] H. X. Zheng, Y. L. Yang, W. S. Liu, Y. Zhong, Y. Cao, R. L. Qiu, C. Liu, A. Van Der Ent, M. J. Hodson, Y. T. Tang, Rare earth elements detoxification mechanism in the hyperaccumulator *Dicranopteris linearis*: [silicon-pectin] matrix fixation, J. Hazard. Mater. 452 (2023) 131254. https://doi.org/10.1016/j.jhazmat.2023.131254

[47] N. Gigli-Bisceglia, E. van Zelm, W. Huo, J. Lamers, C. Testerink, Arabidopsis root responses to salinity depend on pectin modification and cell wall sensing, Development 149 (2022) dev200363. https://doi.org/10.1242/dev.200363

[48] W. Tang, W. Lin, X. Zhou, J. Guo, X. Dang, B. Li, D. Lin, Z. Yang, Mechano-transduction via the pectin-FERONIA complex activates ROP6 GTPase signaling in Arabidopsis pavement cell morphogenesis, Curr. Biol. 32 (2022) 508-517. https://doi.org/10.1016/j.cub.2021.11.031

[49] Y. Hailai, Y. Liu, Z. Yang, Y. Li, J. Feng, W. Li, H. Sheng, Silicon regulation of manganese homeostasis in plants: mechanisms and future prospective, Front. Plant Sci. 15 (2024) 1465513. https://doi.org/10.3389/fpls.2024.1465513

[50] N. Yamaji, N. Mitani-Ueno, T. Fujii, T. Shinya, J. F. Shao, S. Watanuki, Y. Saitoh, J. F. Ma, Shoot-Silicon-Signal protein to regulate root silicon uptake in rice, Nat. Commun. 15 (2024) 10712. https://doi.org/10.1038/s41467-024-55322-7

[51] L. Wang, T. Yang, Q. Lin, B. Wang, X. Li, S. Luan, F. Yu, Receptor kinase

FERONIA regulates flowering time in Arabidopsis, BMC Plant Biol. 20 (2020) 1-16.

https://doi.org/10.1186/s12870-019-2223-y

[52] Z. Pang, H. Peng, S. Lin, Y. Liang, Theory and application of a Si-based defense

barrier for plants: Implications for soil-plant-atmosphere system health, Crit. Rev. Env.

Sci. Tec. 54 (2024) 722-746. https://doi.org/10.1080/10643389.2023.2267939

Figures 1-7

Fig. 1 Si alters cell wall chemistry in a pectin-dependent manner in salt-stressed rice

plants. (A) FTIR spectroscopy of root cell wall. (B-D) Cellulose, hemicellulose and

uronic acid content measured using the colorimetric method. (E) Fractions of

methylesterified pectic polysaccharides. Values are means $\pm$ SD ($n = 3$). Data were

statistically evaluated using Welch's one-way ANOVA, and the significance of

differences (*, $p < 0.05$; ns, not significant) is indicated.

Fig. 2 Si improvement of cell wall integrity. (A) Images of cells in the root elongation

zone stained with PI. (B) The number of burst cells in the root elongation zone. (C)

Images of root cell wall by atomic force microscopy. (D) Average Young's modulus of

the root cell wall. Values are means ± SD (n =10). Data were statistically evaluated

using Welch's one-way ANOVA, and the significance of differences (***, p < 0.001)

is indicated. Scale bar = 50 μm (A) and 1 μm (C).

Fig.3 Relationship between Si and FER kinase activity. (A) Comparative phylogenetic analysis of rice and Arabidopsis CrRLK1L proteins. (B, C) The expression patterns of FER homolog gene *OsFLR1* and *OsFLR2* under the NaCl, Si, and Si+NaCl conditions. Values are means $\pm$ SD ($n = 3$). Data were statistically evaluated using Welch's one-way ANOVA, and the significance of differences (**, $p < 0.01$; *, $p < 0.05$; ns, not significant) is indicated.

Fig. 4 Si roles depend on the activation of FER kinase. (A) Images of hydroponic seedlings treated with reversine. 5-day-old rice seedings were transplanted into the medium containing reversine for 4 days. (B, C) Plant height and fresh weight of rice seedlings after being transferred to the medium with reversine for 4 days. (D) Images of hydroponic seedlings treated with NaCl-Si, NaCl+Si, NaCl-Si+reversine, and NaCl+Si+reversine. 5-day-old rice seedings were transplanted into the medium containing the aforementioned four treatments for 4 days. (E, F) Plant height and fresh weight of rice seedlings measured in those shown in (D). (G) Images of hydroponic seedlings treated with staurosporine, NaCl-Si+staurosporine, and NaCl+Si+staurosporine. 5-day-old rice seedings were transplanted into the medium with above three treatments for 4 days. (H, J) Plant height and fresh weight of rice seedlings measured in those shown in (G). Values are means $\pm$ SD ($n > 12$). Data were statistically evaluated using Welch's one-way ANOVA, and the significance of differences (**, $p < 0.01$; ns, not significant) is indicated. Scale bar = 3 cm.

Fig.5 FER kinase regulates the uptake and accumulation of Si. (A, B) The expression patterns of Si transporter gene *OsLsi1* and *OsLsi2* under the NaCl, reversine, and Si+reversine conditions. (C) Si contents in rice seedlings treated with Si, Si+NaCl, Si+reversine, Si+NaCl+reversine. 5-day-old rice seedings were transplanted into the medium with above treatments for 4 days and then the Si contents were measured using the colorimetric method. Values are means ± SD (*n* = 3). Data were statistically evaluated using Welch's one-way ANOVA, and the significance of differences (***, p < 0.001; **, p < 0.01; *, p < 0.05) is indicated.

Fig.6 ABA biosynthesis is affected by Si and FER kinase under normal (A) and salt stress (B) conditions. 5-day-old rice seedlings were transplanted into fresh Yoshida solution containing Si, Reversine, NaCl, NaCl+Si, NaCl+Si+Reversine. Four days later, rice plants were harvested to detect ABA content. Values are means ± SD (n = 3). Data were statistically evaluated using Welch's one-way ANOVA, and the significance of differences (***, p < 0.001; **, p < 0.01) is indicated.

Fig.7 Si mitigates salt stress by modulating pectin methyl-esterification and FLR1/2-mediated cell wall integrity signaling. (A) Rice adaption to salt stress without Si addition. Salt stress induces pectin de-methyl-esterification, leading to cell wall softening and activation of OsFLR1/2-mediated cell wall integrity signaling. Additionally, high concentrations of NaCl increase ABA production, which subsequently inhibits the expression of Si transporter genes *OsLsi1* and *OsLsi2*. (B) The switch roles of Si in regulating the activity of FER homologs OsFLR1/2 in response to salt stress. In the early stage of salt stress, Si application increases pectin methyl-esterification and stiffens the cell wall. Concurrently, Si enhances the activity of OsFLR1/2, although the underlying mechanism remains to be elucidated. In turn, OsFLR1/2 positively regulate the expression of Si transporter genes by inhibiting the synthesis of ABA, thereby promoting Si accumulation. However, in the late stages, Si can suppress the expression of *OsFLR1/2* genes, facilitating growth recovery. The presence of a feedback loop between Si nutrition and FER kinase activity suggests the involvement of OsFLR1/2 in initiating Si signaling.