Learning millisecond protein dynamics from what is missing in NMR spectra

Hannah K. Wayment-Steele[1]*, Gina El Nesr[2]*, Ramith Hettiarachchi[3,4], Hasindu Kariyawasam[5,6], Sergey Ovchinnikov[3], Dorothee Kern[1#]

[#]To whom correspondence should be addressed: dkern@scripps.edu
*These authors contributed equally.

[1]Department of Integrated Structural and Computational Biology, Scripps Research & Howard Hughes Medical Institute, La Jolla, CA
[2]Biophysics Program, Stanford University, Stanford, CA
[3]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA
[4]Current address: Ray and Stephanie Lane Computational Biology Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA
[5]Center for Advanced Imaging, Faculty of Arts and Sciences, Harvard University, Cambridge, MA
[6]Current address: Cornell Bowers College of Computing and Information Science, Gates Hall, Ithaca, NY

Abstract. Many proteins' biological functions rely on interconversions between multiple conformations occurring at micro- to millisecond (µs-ms) timescales. A lack of standardized, large-scale experimental data has hindered obtaining a more predictive understanding of these motions. After curating >100 Nuclear Magnetic Resonance (NMR) relaxation datasets, we realized an observable for µs-ms motion was hiding in plain sight. Millisecond motions can cause NMR signals to broaden beyond detection, leaving some residues not assigned in the chemical shift datasets of ~10,000 proteins deposited in the Biological Magnetic Resonance Data Bank (BMRB)[1]. We made the bold assumption that residues missing assignments are exchange-broadened due to µs-ms motions, and trained various deep learning models to predict missing assignments as markers for such dynamics. Strikingly, these models also predict µs-ms motion directly measured in NMR relaxation experiments. The best of these models, which we named Dyna-1, leverages an intermediate layer of the multimodal language model ESM-3[2]. Notably, dynamics directly linked to biological function — including enzyme catalysis and ligand binding — are particularly well predicted by Dyna-1, which parallels our findings that residues with µs-ms motions are highly

conserved. We anticipate the datasets and models presented here will be transformative in unlocking the common language of dynamics and function.

Introduction. The functions of proteins often depend on their ability to interconvert between multiple conformations. Our ability to understand fundamental biological mechanisms would be significantly improved if we had greater ability to predict multiple conformations and the timescales at which they interconvert. Computationally modelling the motions of proteins has been a longstanding goal since the first molecular dynamics (MD) simulations of proteins[3,4]. However, our predictive power for the motions of proteins pales in comparison to our predictive power for single structures. AlphaFold-2 (AF2)[5] demonstrated that the experimental data available in the Protein Data Bank (PDB)[6], coupled with large-scale sequencing data and modern deep-learning architectures, could achieve unprecedented accuracy with notable impact in real-world tasks[7,8]. Glimpses that AF2 might have learned information about multiple conformations abound [9-11]. However, the task of predicting *protein dynamics* lacks what the task of predicting single structures had: large-scale, standardized benchmarks of experimental observables to compare to. Any computational method, be it deep-learning-based or simulation-based, currently faces a paucity of standardized experimental data available on dynamics to evaluate on, let alone train from (**Fig. 1a**).

We focus here on µs-ms dynamics, since it is well established that motions on this timescale are directly linked to biological function[12] and are often concerted in nature[13]. Where can we find experimental data on µs-ms dynamics? The main source of structural data used in deep learning to date, structure information in the PDB, lacks time information. Time-resolved X-ray diffraction and cryo-electron microscopy methods are gaining increasing traction, yet processing these datasets requires nontrivially deconvolving signal to identify multiple conformations. In contrast, the last 50 years of methods development in NMR has led to experiments in which the

presence of μs-ms dynamics *creates* signal in the form the observable R_{ex}. R_{ex} is an increase in R_2 relaxation times that unambiguously indicates that the monitored atom is experiencing changes in its chemical environment at the μs-ms timescale[14,15].

We wished to address the lack of experimental dynamics benchmarks by curating as many datasets as we could of standardized R_{ex} measurements. In analyzing our "RelaxDB" dataset of 133 proteins, we found that residues with μs-ms motion were more conserved than residues with no motion, which in turn were more conserved than residues with pico-nanosecond motion. This suggested to us that if we could find more experimental data on these μs-ms dynamics, it might be feasible to use an evolutionary-guided deep learning approach, such as a protein language model, to predict μs-ms dynamics. We then realized that such an observable was available in the ~10,000 proteins with deposited NMR chemical shift information in the BMRB. To train a new predictive model for dynamics, we made the key assumption that missing ^{15}N backbone assignments are caused by exchange-broadening due to μs-ms motions! We used this "missing" label as motivation to delineate a classification task: can a deep learning model predict which residues are missing? Our results demonstrate that hallmarks of millisecond-timescale dynamics are learnable by modern deep learning approaches. The developed model, Dyna-1, is able to predict biologically relevant μs-ms motions as assessed on proteins that have been characterized with multiple types of experiments to measure μs-ms motion collected by many independent labs. Notably, our model led to reinterpretation of published NMR relaxation data in several ways. While investigating "false negatives" in the RelaxDB dataset, we noticed that some proteins interact with phosphate ions as part of their biological function, yet the NMR relaxation experiments were conducted in phosphate buffer. Dyna-1 did not predict exchange from phosphate binding. Conversely, Dyna-1 did predict millisecond motions that typical analysis of Carr-Purcell-Meiboom-Gill (CPMG) experiments missed, but more careful consideration of the existing NMR data validated. We anticipate Dyna-1 as

well as the datasets used for training and evaluation will be transformative in better understanding how biomolecular dynamics leads to function.

Results

Residues with micro-millisecond dynamics are more conserved than other dynamics categories. In setting out to curate as many datasets as we could of standardized R_{ex} measurements, we started with the experiment where R_{ex} is perhaps most directly measured: Carr-Purcell-Meiboom-Gill (CPMG) relaxation dispersion[14-17]. However, we could only find fewer than 10 sets of complete CPMG dispersion data either publicly available or from private correspondences. We next turned to a more abundant type of experiments that have been collected for the last 30 years, which we will refer to as R_1/R_2/NOE experiments, that provides information on both pico-nanosecond (ps-ns) and μs-ms motions. We compiled data on 163 distinct protein domains (**Fig. 1b**), the earliest of which were published in 1990[18]. Roughly half of these were deposited in the BMRB, though many required further curation such as converting from incorrectly identified units. The other half came from private correspondences, including a benchmark of 20 proteins compiled by Goodman et al. in 2000[19], supplemental tables of published papers, and plot digitization from publications. Because R_{ex} reports on any process causing a change in chemical environment, which can include monomer-dimer equilibrium or the binding-unbinding of a ligand, we only included datasets of apo proteins with no cofactors present, and where the authors had verified the protein is monomeric under the experimental conditions used. We excluded intrinsically disordered proteins.

Data collected at one magnetic field strength results in three values per residue: a R_1, R_2, and NOE value. R_{ex} is the increase in R_2, the backbone amide's transverse relaxation (see Methods), caused by exchange between states with different chemical shifts on the μs-ms timescale. A common practice to estimate the presence of R_{ex} is to look for high outliers in R_2/R_1 per residue. R_1

does not contain R_{ex} and decreases if motion is present at slower timescales. However, anisotropic tumbling can also alter the R_2/R_1 ratio. The model-free formalism[20-22] was developed to estimate R_{ex} from these data; however, we were unable to robustly apply it at this scale of datasets since relaxation data from multiple field strengths—important for accurate estimations—were often not available. Therefore, we developed the following processing method to label residues with R_{ex}. We generated AF2 models for the 163 proteins and calculated their R_2 and R_1 values accounting for rigid-tumbling only with HYDRONMR[23,24], fitting one constant per protein to best match the rigid-overall tumbling time with the experimental time. We then designated a residue as having R_{ex} present if the difference between experimental R_2 and rigid-tumbling R_2 (dR_2) was greater than one standard deviation for that protein. We also labeled residues as having ps-ns motion if their hetNOE was less than 0.65.

Fig. 1c depicts the excellent degree to which HYDRONMR accounts for anisotropic effects for four example proteins from RelaxDB. Experimental data is in black, R_2/R_1 calculated from rigid tumbling is in grey, and labels for ps-ns motion and µs-ms motion are in blue and orange, respectively. The protein GB3 is small yet has pronounced anisotropy in its R_2/R_1 arising from its helix; it was a system first used to demonstrate that anisotropy in tumbling could lead to heightened R_2/R_1[25]. This "zigzag" pattern due to differing directions of backbone N-H amide bond vectors in a helix is also evident in the highly-anisotropic helical bundle of alpha spectrin[26]. Nevertheless, our fitting process discerns between anisotropic data features and increases in R_2/R_1 due to exchange, exemplified in AcrIIA4 Cas9 inhibitor[27] and beta-lactamase[28]. We filtered the original 163 proteins to 133 for which this process was successful by visually assessing each dataset. All 133 datasets are in **Extended Data Fig. 1**. **Fig. 1d** depicts the distribution of all residues from these proteins along the two experimental observables defining the presence of different timescales of motion: namely, change in R_2 (see Methods) for µs-ms motion, and hetNOE for ps-ns

motion. As expected, many residues with fast motion have a decrease in dR_2. A few residues have both fast motions detected via NOE and elevated dR_2. If anything, we anticipate that this method of looking at elevated R_2/R_1 is under-estimating exchange for residues where motions are present at more than one timescale.

Despite comprising of only 133 proteins, the RelaxDB benchmark represents an unprecedented amount of experimental information on dynamics in one place. With this curated dataset in hand, we were interested in seeing if there were any eminent relationships between sequence and dynamics. When calculating sequence conservation, we noticed a striking trend: the residues with μs-ms motion were more conserved than residues with no designated motion, which in turn were more conserved than residues with ps-ns motion. Overall, solvent-exposed residues are less conserved than non-solvent-exposed residues (**Fig. 1e**). Residues involved in μs-ms processes are often involved in function, whether it be an allosteric signaling network, catalysis, or a ligand binding site undergoing motion in the absence of its binding partner. Nevertheless, we were surprised that this could be quantitatively detected in as few as 133 proteins. Furthermore, the predicted local distance difference test (pLDDT) measure from AF2 is lowest for residues with ps-ns motion, and shows little discrepancy between residues with no motion and μs-ms motion (**Extended Data Fig. 2**), supporting a previous finding that pLDDT is a reasonable predictor of backbone order parameters in 10 proteins[29].

Though we hoped RelaxDB would be useful for evaluating models, any model trained with only 133 proteins is unlikely to generalize. In looking at the data, we realized that our initial curation of the data lacked a key observable that could also give information on μs-ms dynamics. Some residues in the proteins of RelaxDB could not be assigned a dynamics label because they did not have

measured $R_1/R_2/NOE$ values. These residues lacking relaxation data fall into two fundamentally different categories. In the first category, a residue is assigned, but the authors did not report $R_1/R_2/NOE$ values for a variety of reasons: possibly the peaks were too overlapped to fit an exponential to each reliably, or an exponential did not fit well. In the second category, a residue is missing $R_1/R_2/NOE$ values because the residue was unable to be assigned in the first place. Residues can be missing from assignment if they are "exchange-broadened", i.e., they are experiencing exchange between two states where the chemical shift difference of the two states is comparable to the frequency at which the residue is interconverting. We therefore identified published assignments corresponding to each protein from the same authors, and created labels to distinguish if residues without $R_1/R_2/NOE$ data were assigned or missing from assignments. We hypothesized that if the residues missing assignments were also reporting on residues with μs-ms motion, they would show similar trends in conservation to residues with R_{ex}. Indeed, sequence conservation is statistically indistinguishable between residues with R_{ex} and residues with missing peaks (**Fig. 1e**). Fundamentally, whether a residue has a peak with R_{ex} or a peak that is missing and exchange-broadened depends on the chemical shift difference of its multiple states relative to their interconversion rate.

Curating labels for μs-ms exchange across thousands of proteins.

After the painstaking work of finding a set of assignments to match each dataset in RelaxDB (and rejecting several candidate datasets where we could not reliably identify corresponding published assignments), we realized that these missing assignments were present at orders-of-magnitude higher counts than in just the relaxation measurements (**Fig. 2a**); the Biological Magnetic Resonance Bank (BMRB) contains ~12,000 proteins with deposited chemical shifts. Could we train a model on all the proteins in the BMRB, where the label is, "is this residue assigned or not", that

could predict µs-ms exchange? The key bold assumption we make is that backbone amide assignments are primarily missing due to exchange, as opposed to incomplete assignments or other reasons. We next curated a dataset where this might be reasonable: we started with all protein entries in the BMRB as of May 2023 and removed entries with selective isotope labeling and methyl labelling. We also removed entries with deuterated labelling identified in the metadata, as incomplete back-exchange in H_2O results in erroneously missing amide peaks in deuterated proteins. After further filtering to remove entries with outlier sets of missing peaks for other reasons (see Methods, **Extended Data Fig 3**), our dataset contained 9,381 proteins, two orders of magnitude more proteins than RelaxDB (**Fig. 2b**). Some residues are labeled as missing but are not used in evaluation: because prolines lack an amide proton, the proline backbone amide assignments are by default missing from $^1H^{15}N$ HSQCs. We also did not consider unassigned termini.

After applying these filters, we observed that a substantial fraction of proteins across the BMRB had significant numbers of non-proline and non-terminal missing assignments (**Fig. 2c**). Disaggregating residues by residue and secondary structure type (calculated from structures predicted by ESMFold[30]) reveals that residues are missing across both loops and structured elements for all amino acid types, though most frequently in loops (**Fig. 2d**).

We wished to compare the information in these "missing peak" labels to sources of heterogeneity and uncertainty present in other forms of publicly-available structure data to inform on what models trained on the PDB such as AF2 may or may not have already been exposed to. We used MMseqs[31] to align sequences in the mBMRB dataset (filtered to 80% sequence identity) against all proteins in the PDB. 95% of the mBMRB had one or more similar PDB structures (alignment bitscore

> 50, sequence identity > 90%). 45% of these proteins (2680/5906) had no X-ray or EM structures, but had one or more NMR structures (**Fig. 2e**). NMR structures were removed from the AF2 training set, signifying that much of the mBMRB comprises proteins that AF2 and other algorithms adopting similar training splits have not been trained on.

For proteins characterized by X-ray and/or EM and also assigned by NMR, we found that residues with missing peaks are distinct from residues unresolved in X-ray or EM structure models (**Fig. 2f,g**). Of the residues that are resolved, the distributions of B-factors are similar for residues assigned or unassigned (**Fig. 2h**). In summary, the mBMRB dataset contains unique information on thousands of proteins in comparison to information present in X-ray and EM structures in the PDB.

Deep learning models can predict missing assignments.

With this data in hand, we tested if current deep learning models have the capacity to predict which peaks are missing, which we framed as a binary classification task. We created an initial train/validation/test data split, holding out any sequences with >80% sequence identity to the RelaxDB or RelaxDB-CPMG datasets to use later as higher quality evaluations with actual labels of exchange (**Fig. 3a**). To contextualize model performance on this new task, we began by calculating a set of naïve baselines. Given that residues on each of the secondary structures (loop, sheet, helix) had different propensities of being missing in the training data (cf. **Fig. 2d**), we created a naïve classifier that classifies a residue as "missing" based on training set frequencies. For example, 14% of histidines in loops are missing in the training set, so this naïve classifier randomly assigns a histidine in loop to be missing 14% of the time. This classifier achieved an AUROC of 0.501 on the validation set, equivalent to a random predictor.

We next designed architectures to test the predictive capacity of three pretrained flagship models in deep learning for proteins: AF2[5], a supervised structure prediction algorithm; ESM-2[30], a protein language model trained on sequence alone; and ESM-3[2], a multimodal language model trained on sequence, structure, and function information. We tested ESM-2 and ESM-3 by using the fixed embeddings from each layer as input to a series of Transformer layers (**Fig. 3b, Extended Data Fig. 4**). We created an architecture based on AF2 which we viewed to be significantly different from what ESM-2 or ESM-3 may have learned. This "AF2-pair" model, which uses structure input as templates but no MSA information, learns an attention matrix to select features from AF2's pair representation, which are input to a similar series of Transformer layers (see Methods, **Extended Data Fig. 4**). As a control, we also trained a classifier with the same Transformer architecture, but with one-hot sequence encoding input instead of language model embeddings, "OHE AA" in **Fig. 3c**. We first compared the architectures described above by training with data at 80% sequence identity cutoff to the validation data, as this is a novel prediction task, and we wanted to first assess these models' predictive capability given a majority of the data.

To our surprise, many modelling approaches showed predictive power: the AF2-pair model achieved an AUROC of 0.71 and the embeddings of the last layer of ESM-2 achieved an AUROC of 0.75. Remarkably, even the "OHE AA" baseline, simply a transformer accepting sequence inputs, achieved an AUROC of 0.62. However, our highest performance was achieved using the intermediate layer embeddings of ESM-3 with both sequence and structure input. We wanted to select one layer for further model development and selected it based on the highest AUPRC achieved on the validation set (**Fig. 3c**, right), since this metric had slightly more discrimination than AUROC. This selection led to using the embeddings from layer 22 in ESM-3, which achieved an AUROC of 0.77 on our validation set. In following analyses, our conclusions from AUROC and

AUPRC are quite comparable, and AUPRC visualizations for all subsequent per-protein comparisons discussed are included in **Extended Data Fig. 5**. We re-trained a final model with slightly updated train/validation/test splits and hyperparameters. We named this final model – which is based on the embeddings of layer 22 of ESM-3 and uses sequence and structure model inputs – "Dyna-1".

We probed the effect of sequence and structure similarity in the training data on Dyna-1 performance by training with more stringent training splits compared to our original split of 80% sequence identity. We tested 50% and 30% sequence identity, as well as structure-based similarity cutoffs of TM-score at 0.5 and 0.7. These significantly reduced the size of the training set, with the smallest including only ~2700 proteins. However, the most stringent of these training splits only decreased the AUROC of Dyna-1 on the validation set from 0.755 to 0.74 (**Fig. 3d**), and comparably on the remaining evaluation datasets in this work (**Extended Data Fig. 4**). We did not observe any individual proteins with significant jumps in AUROC between our most stringent training set (30% sequence identity, 0.5 TM-score cutoff) and our most lenient (80% sequence identity, 1.0 TM-score cutoff), which would have indicated model memorization (**Extended Data Fig. 4d**). We proceeded with analyzing the model trained with the most stringent training set. However, we note that model performance does not improve significantly when given almost three-fold more data (cf. **Extended Data Figs. 4-6**).

Fig. 3e depicts AUROC per protein for those with two or more missing assignments in the mBMRB-Test dataset. **Fig. 3f** depicts the structures of some of the best-performing proteins in the test set with tube sized by the output of Dyna-1, i.e., the predicted probability each residue is missing, "p(missing)". The missing assignments in example (iv) are related to the function of this protein,

ribonuclease toxin MqsR: loop β2-β3 is known to have exchange, and its flexibility is reported to be related to its ability to bind mRNA[32]. The antitoxin MqsA acts on this protein not by directly inhibiting its active site, but rather by sequestering the β2-β3 loop and prohibiting its motion, which allows the ribonuclease to show broad non-specificity towards a variety of mRNA substrates[32].

Though these results appeared promising, missing assignments are an incomplete indicator of μs-ms motion, and "false positives" in this data cannot be held as ground truth. This is exemplified in (v), the telomerase protein Pof8[33]. Dyna-1 predicts high p(missing) for the α2-β4 loop, yet most of these residues are assigned. Using the missing peak label as our sole source of ground truth data on μs-ms motion would create many false positives. However, the authors described that assigning the α2-β4 loop and analyzing subsequent relaxation data for the loop was difficult "due to severe line broadening"[33], a clear indicator of μs-ms exchange. We therefore needed to evaluate Dyna-1's predictions using the more-informative relaxation data in RelaxDB to gain a better sense of the model's predictive capability.

Dyna-1 can predict assigned residues with μs-ms exchange.

We found that when evaluating only missing assignments (**Fig. 4ai**), representative models achieved comparable performance to the mBMRB validation and test sets. We then devised a more stringent test: we held out missing residues and asked how well it could predict which assigned residues had exchange (**Fig. 4aii**). Indeed, AF2-pair, ESM-2, and ESM-3-based models achieved a normalized AUPRC well above the controls, indicating that they are able to generalize learning from predicting missing assignments (e.g. p(missing)) to predicting exchange within assigned residues (e.g. p(exchange)). Normalized AUPRC is calculated by subtracting the baseline AUPRC from the AUPRC for each protein. As a summary evaluation, we consider both missing peaks and residues

with measured μs-ms exchange in RelaxDB (**Fig. 4aiii**). Out of the representative model options, Dyna-1 fared the best in normalized AUPRC in this evaluation. Equivalent AUROC values are in **Extended Data Fig. 5**. **Fig. 4a** excludes evaluation on 14 proteins which we realized contained exchange from phosphate buffer binding, described next. The equivalent plots containing those proteins are in **Extended Data Fig. 5**.

Fig. 4b depicts AUROC for all proteins in RelaxDB vs. sequence length. We first noticed there were numerous proteins in RelaxDB with an AUROC of < 0.5, i.e. worse than a random predictor. By carefully reading the corresponding NMR papers, we noticed that some of these were DNA- or RNA-binding proteins measured in phosphate buffer, and that the unpredicted exchange often occurred in positively charged surface residues (**Extended Data Fig. 7**). Several works directly state this about either their own or others' datasets being a limitation to interpreting measured exchange in the system they investigate.[34,35] We annotated these low-performing datasets further, and found that out of the 29 proteins with AUROC ≤ 0.5, 14 of these (48%) are proteins that during their biological function would be expected to interact with phosphate groups (i.e. DNA- or RNA-binding proteins or signaling domains), yet were experimentally characterized in phosphate buffer. Removing these 14 proteins from RelaxDB increased Dyna-1's overall performance from an AUROC of 0.62 to 0.65.

Figs. 4c,d,f depict representative proteins where structures are sized by Dyna-1 predicted p(exchange) and colored by experimental data. In Bovine Pancreatic Trypsin Inhibitor (BPTI), Cys14, Lys15, Cys38, and Arg39 are known to undergo millisecond transitions from disulfide bond isomerization.[36] Promisingly for our automated labeling scheme, these four residues are also labeled as having μs-ms motion from relaxation data in ref. [37] (depicted in **Fig. 4c**, top). Shaw et al.

identified 5 kinetically-distinct states on the micro-millisecond timescales via molecular dynamics simulations[38] (depicted in **Fig. 4c**, left). Strikingly, Dyna-1 predicts high p(exchange) for the two loops containing these residues (**Fig. 4c,** right), to achieve an AUROC of 0.92 for predicting millisecond exchange in BPTI.

While inspecting predictions for other small proteins, we noticed that Dyna-1 predicted high p(exchange) for some residues with assignments but no relaxation data. This prompted us to revisit our initial curation. For instance, in the Tudor domain of Survival of Motor Neuron protein (SMNTD)[39], some of the highest p(exchange) from Dyna-1 were in two residues that were assigned yet had no relaxation data reported: Glu104 and Tyr130 (depicted in black, **Fig. 4d**). The amide hydrogen of Glu104 was assigned at 4.9 ppm, which is a highly unlikely value for an amide hydrogen. We posit that for this small domain and well-resolved spectra that the authors report, this residue is more likely exchange-broadened with an incorrect assignment reported. Also, no R_1 value was reported for Tyr130, yet the authors categorized it as fitting to "model 4" from the model-free formalism, which is a model containing a term for R_{ex} and requires an R_1 value. If we consider both residues as having evidence of exchange, Dyna-1's AUROC increases from 0.74 to 0.79.

In the NtrC receiver domain, Dyna-1 achieved an AUROC of 0.60. We revisited the 10 residues that were assigned yet had no relaxation data. We found that 2/10 had been erroneously assigned in chemical shift depositions corresponding to ref. [40] despite being exchange-broadened. 5/10 had very small peaks, indicative of exchange-broadening, and the remaining 3/10 were unanalyzable due to peak overlap. Taking these into account increases Dyna-1's AUROC to 0.62. Dyna-1 missed predicting exchange in Tyr101 posited to be from ring-flipping. This is exemplary of what we found

more broadly across all of RelaxDB, that Dyna-1 is weakest at predicting exchange in beta-sheets (**Fig. 4e**).

In assessing what motifs Dyna-1 was able to predict in larger proteins (**Fig. 4f**), we were intrigued to see that many of the dynamic motifs it correctly predicts have biological relevance. In flavodoxin YcqA, Dyna-1 predicts high exchange in the loop regions that correspond to the FMN ligand-binding site. In Olfactory Marker Protein from humans, Dyna-1 predicts high p(exchange) in the omega-loop, which is described in ref. [41] to have substantial R_{ex} and is highly conserved. More recent studies indicate that the omega loop serves as a nuclear export signal[42]. In APOBEC-2, Dyna-1 predicts high p(exchange) in a loop that in structure homologues of APOBEC-2 is known to control the specificity of DNA substrates[43].

Another example is Tyrosine-phosphatase A from *M. tuberculosis* (MptpA)[44] (**Fig. 4f**). Dyna-1 has high p(exchange) in loops containing residues with differing orientations between the determined apo and holo structures (apo: 2LUO, holo: 1U2P) and which are postulated to be involved in regulating enzyme specificity[44]. Some of Dyna-1's high p(exchange) is in the P-loop of MptpA, which our curation indicated had been assigned (BMRB:18533) yet had no relaxation data (BMRB: 26513). However, the authors noted that "[d]ue to exchange broadening the P-loop, residues Cys^{11}–Gly^{13} and Ile^{15}–Ser^{18} were not detectable"[44]. If we consider these labels, the AUROC for MptpA increases from 0.65 to 0.73.

To test our hypothesis that Dyna-1 was performing better in predicting conserved motions, we calculated AUROC across different sets of residues categorized by conservation score and core and surface designation. Indeed, we found that Dyna-1 had the most predictive power for highly

conserved residues in both the core and surface, supporting our observations on individual proteins (**Fig. 4g**).

Dyna-1 predictions in the MAP kinase p38 gamma (MK12) are indicative of room for future improvement. For MK12, we used labels derived from the authors' model-free fitting to multiple field strengths[45]. MK12 is a highly dynamic protein with ps-ns and μs-ms processes occurring throughout its structure, and intriguingly, the p(exchange) predicted by Dyna-1 was high across the whole protein. Improving discrimination between ps-ns and μs-ms motions is the subject of future work. We identified numerous proteins where Dyna-1's highest p(exchange) is in regions with ps-ns motion (blue in **Fig. 4b**, structures in **Extended Data Fig. 6c**). One challenge in interpreting these is that residues can experience both ps-ns and μs-ms motion, and a single field strength is insufficient to quantify motions at two timescales simultaneously.

Dyna-1 predicts exchange not typically considered.

Due to the challenges in discerning different types of motion in single-field-strength R_1/R_2/NOE experiments, we lastly wished to test Dyna-1 on proteins with μs-ms motions characterized by CPMG relaxation dispersion. Unlike R_1/R_2/NOE experiments, CPMG experiments allow for directly measuring contribution from conformational exchange to R_2[14,16,17]. In brief, data are collected with increasingly short gaps between spin-echo pulses, which increasingly suppress contributions from exchange between states with different chemical shifts (R_{ex}) to R_2. In typical data processing (orange trace, **Fig. 5a**), $R_{2,eff}$ plateaus at a value that can be understood to be R_2 in the absence of chemical exchange ($R_{2,inf}$). The difference between $R_{2,eff}$ at the lowest B_1 field strength and $R_{2,inf}$ is the contribution to R_2 that was suppressed, i.e., R_{ex}.

We first evaluated Dyna-1 by creating labels for residues with exchange following this data processing. **Fig. 5b** depicts Cyclophilin A (CypA) in a tube representation sized by Dyna-1 prediction probabilities, where residues with statistically significant R_{ex} from ref. [46] are colored orange. We were thrilled to see that Dyna-1 predicted higher p(exchange) in the beta-sheet with a concerted process involving the sidechains of residues Ser99, Phe113, Met61, and Arg55 (shown in grey in **Fig. 5b**).[47] In **Fig. 5b**, we also show the beta-sheet on the back of CypA, which contains no motion, for comparison. We were curious about residue Arg148: Dyna-1 predicted high probability of it being exchange-broadened, but Arg148 did not register as having significant R_{ex} as evaluated by the above method. When looking at the raw data, we found that Arg148 had high $R_{2,inf}$ compared to other residues. We propose that Arg148 has exchange in a process that is too fast to be suppressed by the B_1 fields used in the CPMG experiment for the given temperature. However, to confirm this requires a reasonable estimate for what $R_{2,inf}$ should be if $R_{2,eff}$ is never sufficiently suppressed to estimate it. We sought a data-processing method that could be simply applied across many proteins, and again used HYDRONMR to calculate R_2 from rigid tumbling and fit these values to best match the overall tumbling time. **Fig. 5c** depicts CPMG data for CypA collected at 25˚C and 600 MHz from a B_1 field strength of 33 (red) to 1000 Hz (blue), overlaid with predicted R_2 from rigid tumbling from HYDRONMR in black. CypA's tumbling is highly isotropic and deviations from $R_{2,inf}$ can be reasonably inferred. Indeed, several residues have unsuppressed R_2 (in green) which conventional data processing would not have identified as having exchange.

We applied this new processing pipeline to 6 datasets where we had CPMG dispersion data for all non-overlapped residues. This included CypA[48], beta-lactamase[28], Adenylate Kinase from *Aquifex Aeolicus*[49], Biliverdin reductase B[50], K-Ras[51], and Dual-specificity phosphatase 3[52]. For 5/6 of these, Dyna-1's performance improved when considering residues with unsuppressed R_2, some quite

dramatically: for instance, Dyna-1's AUROC on beta-lactamase from *M. tuberculosis* increases from 0.65 to 0.72 (**Fig. 5d**, structures in **5e**). Dyna-1 also predicts substantial unsuppressed R_2 dynamics in Biliverdin reductase B, a cellular redox regulator, where millisecond motions are implicated in efficient coenzyme engagement and catalysis[50]. Intriguingly, Dyna-1 also predicts concerted millisecond dynamics in the switch I and II regions of K-Ras that is crucial for its signaling activity[51]. All datasets are depicted in **Extended Data Fig. 8.** We additionally analyzed 4 proteins in the literature where CPMG had been performed but the raw data was unavailable and found that Dyna-1 also had predictive power for all of these when assigning labels based on residues that had been reported to contain exchange. Structures sized by Dyna-1 predictions with experimental labels are shown in **Extended Data Fig. 9**.

Discussion

AlphaFold-2 (AF2) fundamentally altered the course of molecular biology and was made possible thanks to the clear task of protein structure prediction defined in the Critical Assessment for Structure Prediction (CASP) challenges and the standardized data deposited in the Protein Data Bank (PDB). Though widely acknowledged that understanding protein dynamics is essential for understanding protein function, the experimental measurements of dynamics that have uncovered foundational principles of allostery, specificity, catalysis, and more have resisted being collected into standardized datasets. Given the predictive power displayed in Dyna-1, it is essential that the field shares and deposits experimental data on dynamics in macromolecules. One barrier to this in NMR has been the perceived diversity and complexity of experiments that measure dynamics. We have laid out in this work a simple task – classify residues with µs-ms exchange – and showed that

we could train a model with predictive power across multiple experimental data types: missing

assignments, R_{ex} from R_1/R_2/NOE, and multiple exchange indicators from CPMG.

Further improvements to Dyna-1 doubtlessly exist for the task of classifying residue dynamics: for

instance, ESM-3, the base of Dyna-1, was trained with C-alpha structure data, and not sidechain

information. Incorporating sidechain-aware training may improve performance. Additionally, Dyna-

1 is trained on proteins under 400 residues in length, possibly limiting the scope in which Dyna-1

can predict the dynamics involved in larger proteins. Dyna-1's sensitivity to within-family variations

in dynamics and its ability to predict dynamics of *de novo* proteins is the subject of ongoing

research. Our comparison of the Dyna-1 model architecture with a training split of 30% sequence

identity, 0.5 TM-score cutoff to a more lenient training split of 80% sequence identity, 1.0 TM-score

cutoff showed only a marginal increase in performance. This indicates that future model

improvements are likely to require innovations in model architectures, rather than simply

increasing the dataset size alone.

We showed that using one of the simplest possible NMR observables for dynamics – if a residue ^{15}N

backbone assignment is missing – resulted in a model with predictive power for exchange because

that observable was available on a scale of ~10,000 proteins. We also showed that this observable

is distinct from other measures of heterogeneity available in X-ray and EM structures. Our

demonstrations were in spite of many issues with the current publicly-available data. The data that

we used to obtain our labels – lists of atom assignments – are the end point of NMR data

processing. The spectra themselves are not required to be deposited, and the lack of standardized

requirements to deposit them is a major loss for science. In each spectra, the peak width is related

to R_2[53]. Had the raw data been required to be deposited, we could have had roughly the information

density of RelaxDB (i.e., R_{ex} labels) for all 10,000 proteins. Careful work[54] has curated a set of 100

raw spectra, two orders of magnitude lower. There is much room for improvement in metadata

labeling in the BMRB as well. We filtered out entries that were indicated to be based on a

deuterated sample, but while inspecting our test set, we found at least one entry (25080) where

assignments were transferred from a deuterated sample, though it itself was not deuterated. Such

curation is a useful direction for future machine learning methods development. Looking forward, if

one could prepare samples in higher throughput and diversity, in combination with emerging

automated assignment algorithms[55], one could imagine collecting data with intentionally lower

resolution to quickly collect information on exchange-broadened peaks and peak widths. More

broadly beyond *in vitro* structural biology, there are increasingly many high-throughput readouts of

fitness that reflect underlying mechanisms involving exchange between multiple states. Combining

this data in conjunction with *in vitro* structure data will undoubtedly be a fruitful direction to

improve understanding of dynamics in both contexts.

Competing Interests. D.K. is a co-founder of Relay Therapeutics and MOMA Therapeutics. H.K.W.-

S. and D.K. are listed as inventors on a provisional patent application related to the methods

described in this manuscript. The remaining authors declare no competing interests.

Data availability. The RelaxDB dataset is publicly available for noncommercial use at

https://github.com/WaymentSteeleLab/Dyna-1. The mBMRB datasets will be made available upon

publication.

Code availability. Code to run Dyna-1 is available for noncommercial use at

https://github.com/WaymentSteeleLab/Dyna-1.

Author contributions. H.K.W.-S., G.E.N., S.O., and D.K. designed the research. H.K.W.-S. curated the datasets. G.E.N., H.K.W.-S., R.H., H.K. contributed to model training and development. H. K.W.-S. and D.K. interpreted all existing NMR data during model validation. H.K.W.-S., G.E.N., D.K. wrote the manuscript. All authors reviewed the manuscript.

Acknowledgments. We thank Ricardo Padua, Hannes Ludewig, Lucy Colwell, Jeffrey Hoch, Art Palmer, Anthony Gitter, Sameer D'Costa, and Kern lab and Ovchinnikov lab members for useful discussions and advice. We thank Martin Stone for sharing the Indiana Dynamics Database data his group curated in 2000. We thank Katie Henzler-Wildman, Magnus Wolf-Watz, Elan Eisenmesser, J. Patrick Loria, Marcellus Ubbelink, George Lisi, Sam Butcher, and Nicolas Doucet for sharing data. We thank Zeming Lin for assistance with ESM-3. We thank Possu Huang for financial and cluster support. Model training was performed on the Stanford Sherlock cluster. H.K.W-S. acknowledges funding from the Jane Coffin Childs fellowship. G. E. N. acknowledges funding from the NSF GRFP. This work was supported by the Howard Hughes Medical Institute (HHMI) to D.K.

Fig. 1. A benchmark of 133 protein NMR backbone relaxation measurements demonstrates residues with ps-ns motion are less conserved, and residues with µs-ms motion are more conserved. (a) Many biological processes occur at a µs-ms time regime, yet experimental data in this regime have not existed at a sufficient scale for machine learning. NMR R_1, R_2 and heteronuclear NOE (hetNOE) measure ps to low ns motion, while millisecond timescale exchange causes increased R_2 (R_{ex}). (b) The "RelaxDB" benchmark we curated consists of 133 single-domain protein backbone datasets with associated labels for the nature of motion present at each residue. Pie chart contains the origin of the data in RelaxDB. (c) Example raw R_2/R_1 data (black) and comparison to calculation from rigid tumbling using HYDRONMR (grey line)[23,24] to obtain labels. Labels are shown in the different colors in legend and plotted onto the AF2 predicted structures. Error bars on R_2/R_1 data are reported by authors. (d) Distribution of hetNOE vs. calculated change in

R_2 between experiment and rigid tumbling calculation. (e) Calculating sequence conservation for these residues demonstrates that residues with µs-ms motion are more conserved than residues with no assigned motion, which in turn are more conserved than residues with ps-ns motion. Residues with missing backbone amide assignments are no different in conservation than residues with µs-ms motion. Box plots depict median and 25/75% interquartile range, whiskers = 1.5 *interquartile range.

Fig. 2. Missing NMR amide backbone assignments in BMRB used as label for µs-ms motion. (a) In seeking more experimental data for µs-ms motions, we realized that data on roughly 10,000 proteins exists in the form of missing backbone amide assignments in the BMRB. Theoretical NMR line-shapes for 2-site exchange on different timescales shown. (b) Scheme for curating the "missing assignment BMRB" (mBMRB) dataset. (c) Fraction backbone amides missing vs. sequence length, not considering prolines or unassigned termini. (d) Fraction of backbone amides missing,

disaggregated by amino acid and secondary structure type, in structures predicted by ESMFold[30].

(e) The majority of proteins in the BMRB have only one NMR structure and no X-ray or EM structure.

(f) Residues with missing NMR peaks are distinct from residues missing in X-ray and EM structures, as exemplified by the MAP kinase binding domain of DUSP 16 (BMRB:19330, PDB:3TG3). Residues with missing assignments are in red, residues with missing X-ray density are in blue. (g) Fraction of residues assigned/unassigned in the mBMRB compared with resolved/unresolved in X-ray or EM structures shows little overlap. (h) B-factors of residues with present or missing NMR peaks have similar distributions. Solid lines: median, dotted lines: 25/75% percentile.

Fig. 3. Deep learning has predictive power for the "missing assignment BMRB" (mBMRB). (a) Train/validation/test split scheme to train from the mBMRB while holding out 100 proteins from mBMRB, and all RelaxDB and RelaxDB-CPMG data as test sets. (b) Schematic of deep learning architectures tested. (c) AUROC and AUPRC on validation set testing layers from ESM-2[30] and ESM-3[2] models, AlphaFold2[5] pair representation, and baselines. AA: amino acid. DSSP: secondary structure assignment. OHE AA: one-hot encoded amino acid. Layer 22 was chosen as the best-performing model (Dyna-1). (d) Performance is not significantly impacted by removing sequence or structure homologues more stringently from training. (e) AUROC from Dyna-1, trained using the 30% sequence identity and 0.5 TM-score dataset split, on the mBMRB-Test set. Proteins in blue are shown in (f). (f) Examples of Dyna-1 predictions. Structures are generated with ESMFold from mBMRB sequence, the thickness of the tube is the probability missing as predicted by Dyna-1. Residues are colored as follows: red: peak missing, grey: peak present, purple: proline (no data).

a RelaxDB

b

Phosphate buffer
High p(missing) for ps-ns motion

Dyna-1
p(exchange)
1
0.5
0

Experiment
Peak missing
μs-ms motion
ps-ns motion
No motion label
Proline
No R₂/R₁ data
↘ other evidence for exchange

c Bovine Pancreatic Trypsin Inhibitor

millisecond-separated states (Shaw, 2010)

d SMN tudor domain

NtrC receiver domain

e

f Mercuric transport protein

YcqA flavodoxin

Olfactory marker protein

g

DNA-binding loops in homologues

APOBEC-2

Protein Tyrosine-phosphatase A *M. tuberculosis*

Map-12 kinase

Fig 4. Dyna-1 predicts biologically-relevant micro-millisecond dynamics in NMR relaxation data. (a) Representative models evaluated on (i) missing peaks in RelaxDB, (ii) residues with assigned R_{ex}, and (iii) both missing and R_{ex}. Error bars represent 95% confidence interval for mean evaluated over proteins in RelaxDB. Normalized AUPRC is calculated by subtracting the baseline AUPRC from the AUPRC for each protein, and the metric by which the final model was chosen. Error bars represent 95% confidence interval for mean normalized AUPRC evaluated over proteins in RelaxDB. (b) AUROC per protein vs. sequence length for Dyna-1 in RelaxDB for both missing peaks and residues with R_{ex}. Only shown are proteins with 2 or more residues with missing assignment or exchange. Brown: proteins with exchange from phosphate buffer. Blue: proteins with high p(exchange) primarily in ps-ns regions (cf. Extended Data Fig. 7). Orange arrows: proteins where follow-up on residues with no data resulted in increased AUROC (see d-f). (c) Dyna-1 prediction for BPTI agrees with experimental R_2/R_1 data[37] (top) and heterogeneity among kinetically-distinct substates in molecular dynamics[38] (left). (d) The SMN Tudor Domain and NtrC offer prospective tests of residues with high p(exchange) for which no data was available (black), but other evidence for R_{ex} (orange arrow). R_2/R_1 uncertainties in (c,d) are from author datasets. (e) AUROC for residues binned by core and surface designation and secondary structure type indicates that Dyna-1 has the most predictive power in core helices and surface loops. (f) Representative Dyna-1 predictions for larger proteins. Biologically-relevant features are annotated in grey. The Dyna-1 prediction for MptpA, a phosphatase from *M. tuberculosis*, recapitulates measurements from NMR as well as structural heterogeneity in sidechains regulating specificity (W48, Y128, Y129) (apo: 2LUO, holo: 1UPA). (g) AUROC binned by core/surface location and sequence conservation indicate that Dyna-1 has the most predictive power for more conserved residues. In (e,g), error bars represent standard error evaluated via bootstrapping.

Fig. 5. Dyna-1 even predicts exchange in residues that conventional CPMG data analysis misses. (a) Example ^{15}N CPMG data for Cyclophilin A (CypA)[48]. Error bars represent background noise in spectrum. R_{ex} is typically estimated via difference in $R_{2,eff}$ at lowest field strength and R_2^0 value to which $R_{2,eff}$ plateaus. However, plotting $R_{2,eff}$ for Arg148 (green) demonstrates different behavior: $R_{2,eff}$ remains high over all B_1 field strengths used, indicative of exchange on the microsecond regime that is too fast to be suppressed with the CPMG field strengths used. (b) AF2 structure of CypA with tube thickness by Dyna-1 p(exchange). Legend coloring according to the

experimental data is shown to the right. Dyna-1 predicts high exchange for Arg148. (c) We devised a

data-processing method using R_2 predicted by HYDRONMR to systematically assign residues with

exchange with $R_{2,eff}$ that get suppressed (conventional R_{ex}, in orange) or residues with elevated,

unsuppressed R_2 (green). Spectra depicted with representative data for CypA was collected at

25˚C. (d) Incorporating labels for unsuppressed R_2 increases reported AUROC for 5/6 proteins

where complete dispersion data was available for reprocessing (shown by green arrows). Also

shown is the AUROC for proteins where we could not find dispersion data for all residues, and

instead curated residue labels based on descriptions in literature. (e) Representative experimental

data and Dyna-1 predictions for a selection of proteins with dynamics characterized with CPMG.

For Adenylate Kinase, structure predictions of the open and closed states from AF2 were used as

input into Dyna-1.

References

Hoch, J. C. *et al*. Biological Magnetic Resonance Data Bank. *Nucleic Acids Res* **51**, D368-D376 (2023). https://doi.org/10.1093/nar/gkac1050
Hayes, T. *et al*. Simulating 500 million years of evolution with a language model. *Science*, eads0018 (2025). https://doi.org/10.1126/science.ads0018
McCammon, J. A., Gelin, B. R. & Karplus, M. Dynamics of folded proteins. *Nature* **267**, 585-590 (1977). https://doi.org/10.1038/267585a0
McCammon, J. A. & Karplus, M. Internal motions of antibody molecules. *Nature* **268**, 765-766 (1977). https://doi.org/10.1038/268765a0
Jumper, J. *et al*. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021). https://doi.org/10.1038/s41586-021-03819-2
Berman, H. M. *et al*. The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000). https://doi.org/10.1093/nar/28.1.235
Marks, D. S. *et al*. Protein 3D structure computed from evolutionary sequence variation. *PLoS One* **6**, e28766 (2011). https://doi.org/10.1371/journal.pone.0028766
Morcos, F. *et al*. Direct-coupling analysis of residue coevolution captures native contacts across many protein families. *Proc Natl Acad Sci U S A* **108**, E1293-1301 (2011). https://doi.org/10.1073/pnas.1111471108
Del Alamo, D., Sala, D., McHaourab, H. S. & Meiler, J. Sampling alternative conformational states of transporters and receptors with AlphaFold2. *Elife* **11** (2022). https://doi.org/10.7554/eLife.75751
Wayment-Steele, H. K. *et al*. Predicting multiple conformations via sequence clustering and AlphaFold2. *Nature* **625**, 832-839 (2024). https://doi.org/10.1038/s41586-023-06832-9

Zheng, S. *et al.* Predicting equilibrium distributions for molecular systems with deep learning. *Nature Machine Intelligence* **6**, 558-567 (2024). https://doi.org:10.1038/s42256-024-00837-3

Henzler-Wildman, K. & Kern, D. Dynamic personalities of proteins. *Nature* **450**, 964-972 (2007). https://doi.org:10.1038/nature06522

van den Bedem, H. & Fraser, J. S. Integrative, dynamic structural biology at atomic resolution--it's about time. *Nat Methods* **12**, 307-318 (2015). https://doi.org:10.1038/nmeth.3324

Hansen, D. F., Vallurupalli, P. & Kay, L. E. Using relaxation dispersion NMR spectroscopy to determine structures of excited, invisible protein states. *J Biomol NMR* **41**, 113-120 (2008). https://doi.org:10.1007/s10858-008-9251-5

Tollinger, M., Skrynnikov, N. R., Mulder, F. A., Forman-Kay, J. D. & Kay, L. E. Slow dynamics in folded and unfolded states of an SH3 domain. *J Am Chem Soc* **123**, 11341-11352 (2001). https://doi.org:10.1021/ja011300z

Loria, J. P., Rance, M. & Palmer, A. G., 3rd. A TROSY CPMG sequence for characterizing chemical exchange in large proteins. *J Biomol NMR* **15**, 151-155 (1999). https://doi.org:10.1023/a:1008355631073

Vallurupalli, P., Bouvignies, G. & Kay, L. E. Increasing the exchange time-scale that can be probed by CPMG relaxation dispersion NMR. *J Phys Chem B* **115**, 14891-14900 (2011). https://doi.org:10.1021/jp209610v

Clore, G. M., Driscoll, P. C., Wingfield, P. T. & Gronenborn, A. M. Analysis of the backbone dynamics of interleukin-1 beta using two-dimensional inverse detected heteronuclear 15N-1H NMR spectroscopy. *Biochemistry* **29**, 7387-7401 (1990). https://doi.org:10.1021/bi00484a006

Goodman, J. L., Pagel, M. D. & Stone, M. J. Relationships between protein structure and dynamics from a database of NMR-derived backbone order parameters. *J Mol Biol* **295**, 963-978 (2000). https://doi.org:10.1006/jmbi.1999.3419

Mandel, A. M., Akke, M. & Palmer, A. G., 3rd. Backbone dynamics of Escherichia coli ribonuclease HI: correlations with structure and function in an active enzyme. *J Mol Biol* **246**, 144-163 (1995). https://doi.org:10.1006/jmbi.1994.0073

Lipari, G. & Szabo, A. Model-free approach to the interpretation of nuclear magnetic resonance relaxation in macromolecules. 1. Theory and range of validity. *Journal of the American Chemical Society* **104**, 4546-4559 (1982). https://doi.org:10.1021/ja00381a009

Cole, R. & Loria, J. P. FAST-Modelfree: a program for rapid automated analysis of solution NMR spin-relaxation data. *J Biomol NMR* **26**, 203-213 (2003). https://doi.org:10.1023/a:1023808801134

Bernado, P., Garcia de la Torre, J. & Pons, M. Interpretation of 15N NMR relaxation data of globular proteins using hydrodynamic calculations with HYDRONMR. *J Biomol NMR* **23**, 139-150 (2002). https://doi.org:10.1023/a:1016359412284

Garcia de la Torre, J., Huertas, M. L. & Carrasco, B. HYDRONMR: prediction of NMR relaxation of globular proteins from atomic-level structures and hydrodynamic calculations. *J Magn Reson* **147**, 138-146 (2000). https://doi.org:10.1006/jmre.2000.2170

Hall, J. B. & Fushman, D. Characterization of the overall and local dynamics of a protein with intermediate rotational anisotropy: Differentiating between conformational exchange and anisotropic diffusion in the B3 domain of protein G. *J Biomol NMR* **27**, 261-275 (2003). https://doi.org:10.1023/a:1025467918856

Brenner, A. K., Kieffer, B., Trave, G., Froystein, N. A. & Raae, A. J. Thermal stability of chicken brain alpha-spectrin repeat 17: a spectroscopic study. *J Biomol NMR* **53**, 71-83 (2012). https://doi.org:10.1007/s10858-012-9620-y

Kim, I. *et al*. Solution structure and dynamics of anti-CRISPR AcrIIA4, the Cas9 inhibitor. *Sci Rep* **8**, 3883 (2018). https://doi.org:10.1038/s41598-018-22177-0

Elings, W. *et al*. Two beta-Lactamase Variants with Reduced Clavulanic Acid Inhibition Display Different Millisecond Dynamics. *Antimicrob Agents Chemother* **65**, e0262820 (2021). https://doi.org:10.1128/AAC.02628-20

Ma, P., Li, D. W. & Bruschweiler, R. Predicting protein flexibility with AlphaFold. *Proteins* **91**, 847-855 (2023). https://doi.org:10.1002/prot.26471

Lin, Z. *et al*. Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science* **379**, 1123-1130 (2023). https://doi.org:10.1126/science.ade2574

Hauser, M., Steinegger, M. & Soding, J. MMseqs software suite for fast and deep clustering and searching of large protein sequence sets. *Bioinformatics* **32**, 1323-1330 (2016). https://doi.org:10.1093/bioinformatics/btw006

Yu, V., Ronzone, E., Lord, D., Peti, W. & Page, R. MqsR is a noncanonical microbial RNase toxin that is inhibited by antitoxin MqsA via steric blockage of substrate binding. *J Biol Chem* **298**, 102535 (2022). https://doi.org:10.1016/j.jbc.2022.102535

Basu, R., Eichhorn, C. D., Cheng, R., Peterson, R. D. & Feigon, J. Structure of S. pombe telomerase protein Pof8 C-terminal domain is an xRRM conserved among LARP7 proteins. *RNA Biol* **18**, 1181-1192 (2021). https://doi.org:10.1080/15476286.2020.1836891

Sakhrani, V. V. *et al*. Backbone assignments and conformational dynamics in the S. typhimurium tryptophan synthase alpha-subunit from solution-state NMR. *J Biomol NMR* **74**, 341-354 (2020). https://doi.org:10.1007/s10858-020-00320-2

Morin, S. & Gagne, S. M. NMR dynamics of PSE-4 beta-lactamase: an interplay of ps-ns order and mus-ms motions in the active site. *Biophys J* **96**, 4681-4691 (2009). https://doi.org:10.1016/j.bpj.2009.02.068

Grey, M. J., Wang, C. & Palmer, A. G., 3rd. Disulfide bond isomerization in basic pancreatic trypsin inhibitor: multisite chemical exchange quantified by CPMG relaxation dispersion and chemical shift modeling. *J Am Chem Soc* **125**, 14324-14335 (2003). https://doi.org:10.1021/ja0367389

Beeser, S. A., Goldenberg, D. P. & Oas, T. G. Enhanced protein flexibility caused by a destabilizing amino acid replacement in BPTI. *J Mol Biol* **269**, 154-164 (1997). https://doi.org:10.1006/jmbi.1997.1031

Shaw, D. E. *et al*. Atomic-level characterization of the structural dynamics of proteins. *Science* **330**, 341-346 (2010). https://doi.org:10.1126/science.1187409

Kawale, A. A. & Burmann, B. M. Inherent backbone dynamics fine-tune the functional plasticity of Tudor domains. *Structure* **29**, 1253-1265 e1254 (2021). https://doi.org:10.1016/j.str.2021.06.007

Volkman, B. F., Lipson, D., Wemmer, D. E. & Kern, D. Two-state allosteric behavior in a single-domain signaling protein. *Science* **291**, 2429-2433 (2001). https://doi.org:10.1126/science.291.5512.2429

Baldisseri, D. M., Margolis, J. W., Weber, D. J., Koo, J. H. & Margolis, F. L. Olfactory marker protein (OMP) exhibits a beta-clam fold in solution: implications for target peptide interaction and olfactory signal transduction. *J Mol Biol* **319**, 823-837 (2002). https://doi.org:10.1016/S0022-2836(02)00282-6

Nakashima, N., Nakashima, A., Nakashima, K. & Takano, M. Olfactory marker protein contains a leucine-rich domain in the Omega-loop important for nuclear export. *Mol Brain* **15**, 89 (2022). https://doi.org:10.1186/s13041-022-00973-0

Salter, J. D. & Smith, H. C. Modeling the Embrace of a Mutator: APOBEC Selection of Nucleic Acid Ligands. *Trends in Biochemical Sciences* **43**, 606-622 (2018). https://doi.org:10.1016/j.tibs.2018.04.013

Stehle, T. *et al.* The apo-structure of the low molecular weight protein-tyrosine phosphatase A (MptpA) from Mycobacterium tuberculosis allows for better target-specific drug development. *J Biol Chem* **287**, 34569-34582 (2012). https://doi.org:10.1074/jbc.M112.399261

Aoto, P. C., Stanfield, R. L., Wilson, I. A., Dyson, H. J. & Wright, P. E. A dynamic switch in inactive p38γ leads to an excited state on the pathway to an active kinase. *Biochemistry* **58**, 5160-5172 (2019).

Eisenmesser, E. Z. *et al.* Intrinsic dynamics of an enzyme underlies catalysis. *Nature* **438**, 117-121 (2005). https://doi.org:10.1038/nature04105

Fraser, J. S. *et al.* Hidden alternative structures of proline isomerase essential for catalysis. *Nature* **462**, 669-673 (2009). https://doi.org:10.1038/nature08615

Eisenmesser, E. Z., Bosco, D. A., Akke, M. & Kern, D. Enzyme dynamics during catalysis. *Science* **295**, 1520-1523 (2002). https://doi.org:10.1126/science.1066176

Henzler-Wildman, K. A. *et al.* Intrinsic motions along an enzymatic reaction trajectory. *Nature* **450**, 838-844 (2007). https://doi.org:10.1038/nature06410

Paukovich, N. *et al.* Biliverdin Reductase B Dynamics Are Coupled to Coenzyme Binding. *J Mol Biol* **430**, 3234-3250 (2018). https://doi.org:10.1016/j.jmb.2018.06.015

Hansen, A. L., Xiang, X., Yuan, C., Bruschweiler-Li, L. & Bruschweiler, R. Excited-state observation of active K-Ras reveals differential structural dynamics of wild-type versus oncogenic G12D and G12C mutants. *Nat Struct Mol Biol* **30**, 1446-1455 (2023). https://doi.org:10.1038/s41594-023-01070-z

Beaumont, V. A. *et al.* Allosteric Impact of the Variable Insert Loop in Vaccinia H1-Related (VHR) Phosphatase. *Biochemistry* **59**, 1896-1908 (2020). https://doi.org:10.1021/acs.biochem.0c00245

Dudley, J. A., Park, S., MacDonald, M. E., Fetene, E. & Smith, C. A. Resolving overlapped signals with automated FitNMR analytical peak modeling. *J Magn Reson* **318**, 106773 (2020). https://doi.org:10.1016/j.jmr.2020.106773

Klukowski, P. *et al.* The 100-protein NMR spectra dataset: A resource for biomolecular NMR data analysis. *Sci Data* **11**, 30 (2024). https://doi.org:10.1038/s41597-023-02879-5

Klukowski, P., Riek, R. & Guntert, P. Rapid protein assignments and structures from raw NMR spectra with the deep learning technique ARTINA. *Nat Commun* **13**, 6151 (2022). https://doi.org:10.1038/s41467-022-33879-5

Kneller, J. M., Lu, M. & Bracken, C. An effective method for the discrimination of motional anisotropy and chemical exchange. *J Am Chem Soc* **124**, 1852-1853 (2002). https://doi.org:10.1021/ja017461k

Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**, 261-272 (2020). https://doi.org:10.1038/s41592-019-0686-2

Pei, J. & Grishin, N. V. AL2CO: calculation of positional conservation in a protein sequence alignment. *Bioinformatics* **17**, 700-712 (2001). https://doi.org:10.1093/bioinformatics/17.8.700

Shrake, A. & Rupley, J. A. Environment and exposure to solvent of protein atoms. Lysozyme and insulin. *J Mol Biol* **79**, 351-371 (1973). https://doi.org:10.1016/0022-2836(73)90011-9

McGibbon, R. T. *et al.* MDTraj: A Modern Open Library for the Analysis of Molecular Dynamics Trajectories. *Biophys J* **109**, 1528-1532 (2015). https://doi.org:10.1016/j.bpj.2015.08.015

Chen, H. & Zhou, H. X. Prediction of solvent accessibility and sites of deleterious mutations from protein sequence. *Nucleic Acids Res* **33**, 3193-3199 (2005). https://doi.org:10.1093/nar/gki633

Beaumont, V. A. *Functional Implications of Conformational Dynamics of Protein Tyrosine Phosphatases*, Yale University, (2019).

McDonald, L. R., Boyer, J. A. & Lee, A. L. Segmental motions, not a two-state concerted switch, underlie allostery in CheY. *Structure* **20**, 1363-1373 (2012). https://doi.org:10.1016/j.str.2012.05.008

Aspholm, E. E., Lidman, J. & Burmann, B. M. Structural basis of substrate recognition and allosteric activation of the proapoptotic mitochondrial HtrA2 protease. *Nat Commun* **15**, 4592 (2024). https://doi.org:10.1038/s41467-024-48997-5

Beach, H., Cole, R., Gill, M. L. & Loria, J. P. Conservation of mus-ms enzyme motions in the apo- and substrate-mimicked state. *J Am Chem Soc* **127**, 9167-9176 (2005). https://doi.org:10.1021/ja0514949

Davulcu, O., Flynn, P. F., Chapman, M. S. & Skalicky, J. J. Intrinsic domain and loop dynamics commensurate with catalytic turnover in an induced-fit enzyme. *Structure* **17**, 1356-1367 (2009). https://doi.org:10.1016/j.str.2009.08.014

Zhang, Y. & Skolnick, J. TM-align: a protein structure alignment algorithm based on the TM-score. *Nucleic Acids Res* **33**, 2302-2309 (2005). https://doi.org:10.1093/nar/gki524

Loshchilov, I. Decoupled weight decay regularization. *arXiv preprint arXiv:1711.05101* (2017).

Menon, A. K. *et al.* Long-tail learning via logit adjustment. *arXiv preprint arXiv:2007.07314* (2020).

Waskom, M. L. Seaborn: statistical data visualization. *Journal of Open Source Software* **6**, 3021 (2021).

Methods

Curating the "RelaxDB" dataset.

Theoretical basis for relaxation measurements. The following is condensed from ref. [56]. The longitudinal (R_1) and transverse (R_2) relaxation rates of a backbone nitrogen atom attached to a single proton may be written in a simplified form as

$$R_1 = \frac{A}{10}J(\omega_N),$$

$$R_2 = \frac{A}{15}[J(0) + \frac{3}{4}J(\omega_N)] + R_{ex},$$

where $J(\omega) = S^2\tau_C/(1 + \omega^2\tau_C{}^2)$. $J(\omega)$ is the spectral density function describing the intensity of motions occurring at frequency ω. These expressions do not include terms on the order of $J(\omega_H)$, which are small on the timescale of protein tumbling ($\omega_N\tau_C \gg 1$). In the above, $A = 3d^2 + 4c^2$, where d=$\frac{\mu_0 h \gamma_H \gamma_N}{r_{NH}8\pi^2}$ and $c = \frac{\gamma_N B_0 \Delta\sigma}{\sqrt{3}}$. μ_0 is the permeability of free space, h is Planck's constant, γ_X is the gyromagnetic ratio of atom X (nitrogen N or hydrogen H), r_{NH} is the bond length of the N-H bond, B_0 is the static magnetic field strength, and $\Delta\sigma$ is the chemical shift anisotropy of nitrogen.

Note that the ratio R_2/R_1 for residues that do not demonstrate dynamics can be used to estimate the global tumbling time τ_C via the following relationship:

$$\frac{R_2}{R_1} = \frac{2}{3}\frac{J(0)}{J(\omega_N)} + \frac{1}{2} = \frac{2}{3}\omega_N{}^2\tau_C{}^2 + \frac{7}{6}.$$

BMRB entries.

To get entries from the BMRB that contained R_1, R_2 and NOE data, we used

https://bmrb.io/software/query/ to download a list of all entries with one entity of polymer type polypeptide(L) that contained R_2 relaxation data (referred to as T_2 in the user interface). This resulted in 344 entries when queried in January 2023. We manually went through the list and

identified entries that were monomers, had no cofactors present, were not in micelles/bicelles/nanodisks, were not in denaturing conditions, and had at least one set of $R_1/R_2/NOE$ data all at the same field strength. This resulted in 91 entries. We included data collected at multiple field strengths when available. Several of the BMRB entries had incorrect units, which we fixed by visually inspecting data and comparing to figures of plotted data in the corresponding publications. For three entries, we realized that they contained two or more domains, but that these two or more domains could be modelled as tumbling independently. We therefore separated these into two separate entries.

Data from literature.

To find data from literature, we searched in PubMed and Google Scholar for keywords "NMR", "backbone", and "relaxation". Upon finding a dataset of an apo, monomeric, stably folded protein and if the data was only available in figure form, we used PlotDigitizer (https://plotdigitizer.com/) to compile values for $R_1/R_2/NOE$. We did not include datasets if the figures that we found were too low-resolution to allow for accurately using PlotDigitizer. We only used datasets where we could identify a corresponding set of assignments either cited in the same paper or clearly referencing another set of assignments.

Fitting data / assigning labels.

Generating MSAs.

We generated MSAs using the `run_mmseqs2` command in ColabFold with `filter=True`. We used these MSAs to predict AF2 models. Prior to calculating conservation metrics (below), we removed sequences with more than 10% gaps and used MMseqs easy-cluster to further filter with options `-c 0.9 --cov-mode 1 --min-seq-id 0.9`.

Generating structure models.

We used AF2 as implemented in ColabDesign

(https://github.com/sokrypton/ColabDesign.git@gamma) to generate the models used for

HYDRONMR calculations. We ran AF2 for 3 recycles, seed=0. We used model 1 for each protein. For

the Adenylate Kinase datasets (both from *E. coli* and *A. Aquifex*), AF2 predicts the closed state

though it is known to occupy the open state predominantly in solution. Therefore, we templated AF2

using structure model PDB ID 4AKE of the open state.

We used these structure models as inputs into HYDRONMR[23,24] with default settings to estimate

R2/R1 from rigid tumbling, denoted as $\widehat{R_2^0/R_1^0}$. We trimmed unstructured termini of the 161 proteins

by removing residues from the N or C termini until the pLDDT of the next residue was over 70, or

over 80 for CBFB and PCF11. Residues corresponding to unstructured termini are not considered

during evaluation.

We needed to account for a constant offset in the overall tumbling time between each experimental

dataset and the corresponding prediction. To do so, we fit $\widehat{R_2^0/R_1^0}$ to the experimental data R_2/R_1 as

follows: for each dataset, we minimized the squared difference between R_2/R_1 and $C\,\widehat{R_2^0/R_1^0}$ for

residues where R_2/R_1 was within one standard deviation of the mode, where C is a scalar that is fit

to minimize:

$$\min_C \frac{1}{n} \sum_{i=1}^{n} \left[\frac{R_2}{R_1}(i) - C\frac{\widehat{R_2^0}}{R_1^0}(i)\right],$$

$$i \in \frac{R_2}{R_1}(i) - \ mode\ \left(\frac{R_2}{R_1}\right) < \ \sigma\left(\frac{R_2}{R_1}\right).$$

C was initialized as the ratio of the mean of the experimental R_2/R_1 values to the mean of the predicted R_2/R_1 values. The mode of the experimental R_2/R_1 values was estimated using a kernel density estimation approach. The optimization was performed using the `minimize` function in SciPy[57].

R_2/R_1 is unitless, so we calculate the estimated difference in R_2 as

$$dR_2(i) = [\frac{R_2}{R_1}(i) - C\frac{\widehat{R_2^0}}{R_1^0}(i)]R_1(i).$$

For each protein, we calculated the mode value of dR_2 plus one standard deviation and labeled any residues with dR_2 higher than this as having µs-ms motion. We did this in terms of dR_2, as opposed to dR_2/R_1 to be able to consider this value in terms of s^{-1} units. The majority of the RelaxDB proteins had a cutoff of 3 Hz or lower, indicating our fitting was reasonable.

We assigned a label for residues with ps-ns motion as residues with hetNOE < 0.65.

For the RelaxDB proteins and the analysis shown in **Fig. 1e**, we calculated conservation scores following the method described in ref. [58]. To determine buried/exposed residues, we calculated solvent accessible surface area (SASA) using the Shrake-Rupley method[59] implemented in MDtraj[60] and normalized values by the maximum value per secondary structure type (loop, helix, sheet) over the whole dataset, following the practice in ref. [61]. We assigned residues with a normalized SASA ≤ 0.2 as core and > 0.2 as surface residues, following ref. [61].

The "mBMRB" dataset

Curating data.

We downloaded a list of all entries consisting of polymer type polypeptide(L) from the BMRB[1] using

https://bmrb.io/software/query/. We processed this to retrieve all entities within each entry which

were type polypeptide(L). On May 31, 2023, this resulted in 12102 entities.

We removed entities with isotope labels corresponding to deuterated samples, as a deuterated

sample that has not completely back-exchanged with solvent would lead to missing peaks. We also

removed methyl-labeled samples or samples otherwise partially labeled based on keywords in the

isotope metadata. Additionally, we removed any sequences shorter than 30 residues. After this

filtering, we had 10123 entities.

We removed any entities with "X" in the sequence (242 removed). Some sequences deposited

included multiple copies if the entity consisted of a multimer. We detected and curated these. We

realized that 1834 sequences contained His-tags, many of which were unassigned. We therefore

trimmed the N- or C-termini of sequences if 6 or more contiguous histidines were present,

including everything before the His tag if it was an N-terminal His tag, or everything after if it was a

C-terminal His tag. We further filtered the remaining entities to include only entities with 10 or more

assignments, to remove entries that included only a few chemical shifts (removed 273).

 For each residue, we assigned labels corresponding to whether the ^{15}N backbone chemical shift is

missing or not based on if that residue had a chemical shift entry corresponding to an "Atom_ID" of

N.

We further removed entities that had 15 or more contiguous missing assignment labels between any assigned residues. This was to filter entities where we noticed that entire domains were unassigned. We selected the cutoff of 15 contiguous residues because the most residues missing in a row in the RelaxDB dataset was 14. This removed 227 entities.

Of the remaining 9,381 entries, 5% did not have isotope information present. However, we elected to keep these, as we saw that they had on average fewer missing peaks and shorter sequences than the 95% of other remaining sequences which had 1H labeling (**Extended Data Figure 3**). Roughly 60% of these are older entries (entry_ID < 10000).

Structure models for each protein in the mBMRB dataset were generated using ESMFold v1[30].

Comparing to structure models in PDB

We used MMseqs[31] to align all the mBMRB sequences (9,381) against all protein chains in the PDB (curated based on PDB data in NMRbox, October 2024). We kept all hits with a bitscore greater than 50 and sequence identity greater than 90%. We kept one chain per protein sequence per model.

RelaxDB-CPMG curation

Complete dispersion data available

AdK: $R_{2,eff}$ vs v_{CPMG} for apo AdK from *Aquifex aeolicus* at 10˚C collected at 800 MHz, presented in ref. [49], as well as a corresponding peak list, was found in the Kern lab data archive, with thanks to K. Henzler-Wildman.

BlaC: $R_{2,eff}$ vs v_{CPMG} of apo BlaC collected at 600 MHz and 850 MHz, presented in ref. [28], was kindly provided by M. Ubbink. A peak list to identify missing assignments was taken from BMRB entry 27888 (same as in RelaxDB).

BLVRB: $R_{2,eff}$ vs v_{CPMG} of apo BLVRB, presented in ref. [50], was kindly provided by E. Eisenmesser. A peak list to identify missing assignments was taken from BMRB entry 27462.

CypA: $R_{2,eff}$ vs v_{CPMG} of apo CypA, collected at 10˚C and 25˚C at 600 MHz in ref. [48], as well as peak lists to identify missing assignments at both temperatures, were found in the Kern lab data archive. Because different processes are more prominent at 10˚C or 25˚C, the final labels created represent whether a residue experienced exchange at either temperature.

VHR: $R_{2,eff}$ vs v_{CPMG} was taken from 800 MHz data provided in Table 7.10.3 of ref. [62]. Missing peaks were taken from assignments in Table 7.10. Note that these differ from assignments by the authors in BMRB 27950; the assignments in the dissertation table are more complete.

K-Ras: $R_{2,eff}$ vs v_{CPMG} of K-Ras with GTP bound, presented in ref. [51], had been publicly deposited at https://doi.org/10.5061/dryad.j6q573nm0.

Complete dispersion data unavailable

CheY: Residues with missing peaks were taken as the black residues in Fig. 2 from ref. [63]. Residues were labeled as having exchange when CPMG could individually be fit (all residues in Table 2 in ref. [63]). This is CheY in the presence of 1 mM EDTA to remove any Mg^{2+}, which alters the dynamics.

HtrA2 PDZ domain: Values for Rex were taken from source data corresponding to Fig. 2a in ref. [64]. Assignments were taken from BMRB:51320.

RNase: Residues were labeled as having exchange when CPMG data was >3 Hz at 20˚C in either 500MHz or 600 MHz data. This recreates Fig. 4B in ref. [65]. The $R_{2,eff}$ vs v_{CPMG} data for these residues was kindly provided by P. Loria, but the dispersion data for all residues was not available.

Arginine kinase: Labels for residues with exchange were taken from Table 1 in ref. [66], which summarized all residues for which CPMG data could be fit individually. Table S1 in ref. [66] indicated if residues were missing due to exchange-broadening or overlap, and we used those labels.

Identifying unsuppressed $R_{2,eff}$

We first compared model predictions to labels for exchange where we assigned labels using a conventional assignment from the literature: any residue where R_{ex} is significantly apparent, calculated as R_2 (first field strength) - R_2 (last field strength) is greater than background noise.

However, we noticed that in some datasets there were residues with noticeably high R_2's across all field strengths but $R_{2,eff}$(first) - $R_{2,eff}$(last) was not substantial enough to be greater than 2-3 Hz. These correspond to residues that have motion at the microsecond timescale: at the fast end of the micro-millisecond regime that CPMG can detect.

However, assigning these residues with confidence requires determining what residues have elevated $R_{2,eff}$ when there is no suppression, the equivalent of measuring R_{ex} as is conventionally

done. Given our success using HYDRONMR to quantitatively fit R_2/R_1 ratios in RelaxDB, we used

HYDRONMR to calculate R_2 values and fit a scalar to find the closest fit between $R_{2,inf}$ and $R_{2,rigid}$.

The $R_{2,eff}$ data for all residues, corresponding $R_{2,rigid}$ values, and labels are in **Extended Data Fig. 8**.

Data splitting

We clustered sequences using MMseqs[31] `easy-cluster` with arguments `--cluster-mode 1 --min-seq-id 0.8 -k 5`. We used the resulting clusters to remove all sequences similar to those in RelaxDB or RelaxDB-CPMG. From the remaining sequences, we randomly separated out 500 clusters for the validation set and 100 clusters for the test set.

For training with a 50% or 30% sequence identity cutoff, we used –min-seq-id 0.5 [or 0.3] to cluster but kept the same proteins in the validation and test set. To create training splits with structure homologues removed at TM-score = 0.5 or 0.7, we calculated all pairwise TM-scores using TMalign[67] between the mBMRB-Train set and mBMRB-Val, mBMRB-Test, RelaxDB, and RelaxDB-CPMG sets. We used TM-score normalized by the average length of the query and target protein, as recommended in ref. [67].

The model sweep depicted in **Fig. 3c** was performed with a different dataset split that considered all 163 sequences initially curated for RelaxDB and used a split of 80% sequence identity, 1.0 TM-score cutoff. The splits used for the model sweep in **Fig. 3c** can be found in the `oct2024` data repository. Subsequent model training was reperformed using a data split considering only the later filtered 133 sequences in RelaxDB and used the most stringent training data split (30% sequence identity, 0.5 TM-score cutoff). These subsequent models are depicted in Figs. 3d and onwards.

Training

We trained Dyna-1 on the mBMRB-Train dataset, sampling using a sequence-cluster-based approach. One protein per sequence cluster is chosen randomly during each epoch. We used the validation set to compute the validation loss and validation metrics. The final model weights for all models are chosen using max AUROC on the validation set. We excluded unassigned termini and prolines (which have no backbone amide and therefore are always missing) from model evaluations.

We used the ESM2-t30-150M model for ESM-2 embeddings, which has 31 layers and 150M parameters. Initial attempts using ESM2-t6-8M and ESM2-t12-35M indicated that smaller models did not perform as well, and ESM2-t33-650M performed marginally similar. We used the outputs of AF2 model 1 to construct the AF-pair representation. The weights from ESM and AF2 were fixed (frozen) during training.

Models were trained with the AdamW optimizer[68] with a learning rate of 1e-6. The AF2-pair model was trained with batch size of 4 with an accumulation step every 4 epochs, while the ESM-2 and ESM-3 based models were trained with a batch size of 16 and accumulation each epoch. Training is completed within 500 epochs, and we use early stopping when validation loss begins to overfit. We choose our best performing model by comparing normalized AUPRC on the validation set, as we saw this had slightly more discrimination than AUROC. Normalized AUPRC is calculated by subtracting the baseline AUPRC from the AUPRC for each protein.

The transformer architecture used for the ESM-2 embeddings has 8 heads and 12 layers. The transformer for the ESM-3 embeddings has 6 heads and 12 layers, and the transformer for AF2-pair representation has 4 heads and 4 layers. We note that increase in number of heads or number of layers had negligible improvements for ESM-based models, and we were limited by GPU memory for AF2-pair. The hidden dimension for the ESM-based models is equivalent to the hidden size of the embedding, while for the AF2-pair model is 128. We apply a 0.1 dropout rate. All models were trained on a single NVIDIA A40 GPU, which takes between 12 hours and 2 days to complete training.

Adjusting logits to account for class imbalance.

When converting from logits to probabilities, we adjusted the raw logit outputs predicted by the model to account for the significant class imbalance in the problem following the practice in ref. [69] (cf. **Extended Data Fig. 10**). This practice seeks probabilities that minimizes the "balanced error", the prediction error averaged over all classes. In brief, for data *x* labelled with classes *y*, the native posterior probability *P(y | x)* is proportional to the prior probability of class y, $\pi(y)$:

$$P(y|x) \propto \pi(y)P(x|y).$$

In seeking to minimize the balanced error, the balanced class-probability function instead implicitly becomes

$$P^{bal}(y|x) \propto P(x|y),$$

where *L* is the number of classes (L=2 in our binary classification context).

Menon et al. describe how $P^{bal}(y|x)$ is a Bayes-optimal scorer for minimizing the balanced error. For a Bayes-optimal scorer, the accuracy of any given class is not proportional to its degree of representation in the training data, which is desirable in our context.

Consider our model to be a scoring function $\hat{s}$, with learned class probabilities $\hat{P}(y|x) \propto e^{\hat{s}(x)}$. Via the definition of P^{bal}, we can approximate the Bayes-optimal scorer using our given scorer by using an adjusted scorer s':

$$P^{bal}(y|x) \approx e^{s'(x)} = \frac{\hat{P}(y|x)}{\pi(y)} = \frac{e^{\hat{s}(x)}}{\pi(y)}.$$

Accordingly, the adjusted logits may be written as $s'(x) = s'(x) - \log \pi(y)$.

In the context of our binary classification problem, let π^m be the fraction missing in the training data set and let $l(x)$ be the raw logit output given input x. Dyna-1 is structured to just predict one logit, the logit of the "missing" class, hence the logit predicted for the "not missing" class is always zero. We calculate our probability P(missing) as

$$P(missing|x) = \frac{p(missing|x)}{p(present|x) + p(missing|x)} = \frac{e^{l(x) - \log \pi^m}}{e^{-\log(1 - \pi^m)} + e^{l(x) - \log \pi^m}} = \frac{e^{l(x)}}{\frac{\pi^m}{1 - \pi^m} + e^{l(x)}}.$$

This can be rewritten in terms of the sigmoid function $\sigma(x) = (1 + e^{-x})^{-1}$, which is typically used to convert logits to probabilities:

$$P(missing|x) = \sigma(x - \log \frac{\pi_m}{1 - \pi_m}).$$

Uncertainty estimation.

We estimated uncertainty for comparing models in the RelaxDB dataset (**Fig. 4a, Extended Data Fig. 5d,e**) using a 95% confidence interval calculated in Seaborn v0.11.2 [70].

We estimated uncertainty in binned residue analyses (**Fig. 4e, 4g**) using bootstrapping. We resampled residue subsets with replacement to obtain 100 bootstrap iterations that contained at least one of each class (required for calculating AUROC).

All depicted p-values (**Fig. 1e, Fig. 2h**) are for a 2-sided hypothesis test whose null hypothesis is that two sets of data are uncorrelated, performed using statannotations v0.2.3 (https://pypi.org/project/statannotations/).

We curated uncertainties for the RelaxDB datasets in the following way: if authors reported R_1/R_2/NOE uncertainties, we used those. In the BMRB, we found a few with a deposited uncertainty of zero. We set these to be the smallest nonzero uncertainty value given for that data type in the dataset. If no uncertainties were given, we set the uncertainty for R_1 as 10% the measured value, and 5% for R_2 and hetNOEs. If the data was plot digitized and had error bars, we digitized the error bars as uncertainties.

Extended Data

Extended Data Figure 1: (page 1/3) ^{15}N R_2/R_1 data and labels for all 133 proteins in RelaxDB. Calculated R_2/R_1 from HYDRONMR[23,24] for rigid tumbling is depicted by grey line. Colors used are the same as in Fig. 1. RelaxDB entry ID is in title for each.

Extended Data Figure 1: (page 2/3)

Extended Data Figure 1: (page 3/3)

Extended Data Figure 2: pLDDT (predicted local distance difference test) from AF2 by secondary structure and motion type. Low AF2 pLDDT is most indicative of ps-ns motion in RelaxDB.

Extended Data Figure 3: Statistics from curating the "missing assignment BMRB" (mBMRB). We removed proteins with deuteration or methyl or specific labelling, based on metadata. Some proteins had no isotope information, yet we elected to keep them, as they had fewer missing assignments compared to proteins with [1]H isotope labels and were overall shorter than other isotope categories. (a) Fraction missing split by isotope category. (b) Sequence length by isotope category. In (a,b): Box plots depict median and 25/75% interquartile range, whiskers = 1.5 times the interquartile range. (c) We removed entries with 15 or more consecutive missing residues (orange).

Extended Data Figure 4: Details of training Dyna-1. (a) Architectures of tested ESM and AF2 inputs, outlining the model architecture. At training time, either a sequence-structure pair is passed through AF2 or ESM-3, or a sequence is passed through ESM-2 or ESM-3. The corresponding embedding from the k[th] layer from ESM or AF2-pair output is fixed ("frozen") and used as input into a one-dimensional Transformer architecture. Output is one value per residue, the logit for that residue being missing. (b) Comparing Dyna-1 performance given both sequence and structure

(blue) vs. sequence only (orange) or structure only (green). Performance is best when given both sequence and structure. (c) Dyna-1 performance given different training split cutoffs based on sequence identity and structure similarity (cf. Fig. 3d) across all datasets. (d) AUROC per protein for either training with the 80% sequence identity and 1.0 TM-score split or 30% sequence identity and 0.5 TM-score split. Performance on individual proteins in the validation, test, RelaxDB, and RelaxDB-CPMG does not significantly change between more stringent sequence similarity cutoffs.

Extended Data Figure 5. Dyna-1 trained with 80% sequence identity cutoff is comparable to results presented in main text for 30% sequence identity and 0.5 TM-score cutoff. (a) AUROC for mBMRB-Test (cf. Fig. 3e), RelaxDB (cf. Fig. 4b), RelaxDB-CPMG (cf. Fig. 5d). Coloring for each follows respective main-text figures. (b) Normalized AUPRC per protein for Dyna-1, trained with 30% sequence identity and 0.5 TM-score cutoff. (c) Normalized AUPRC per protein for Dyna-1, trained with 80% sequence identity cutoff.

a Norm. AUPRC, Phosphate buffer data included

b AUROC, Phosphate buffer data removed

c

Extended Data Figure 6. (a) RelaxDB evaluated by normalized AUPRC across different models tested, using all 133 proteins including those with exchange from phosphate binding (cf. **Fig. 4a**). (b) RelaxDB evaluated across different models evaluated by AUROC, with phosphate binding proteins removed. In (a,b), error bars represent 95% confidence interval evaluated over proteins in RelaxDB. (c) ^{1}H-^{15}N HSQC of NtrC at 25˚C, annotated with residues for which assignments exist in BMRB entry 4527 (non-phosphorylated NtrC), but relaxation data is missing from BMRB entry 4762. The Arg56 and Gly97 assignments in entry 4762 (shown in red here) are presumed to be erroneously copied from entry 4528, which are assignments corresponding to phosphorylated NtrC. All residues labeled orange are very weak peaks indicative of R_{ex}. For these, we concluded exchange broadening was the reason for no reported relaxation data. Black residues are overlapped peaks.

Extended Data Figure 7: Some proteins with poorly predicted exchange in RelaxDB reflect phosphate-binding exchange or high probability for ps-ns motion. 14/29 of proteins in RelaxDB predicted with an AUROC < 0.5 are these proteins that bind phosphate ion moieties during their biological function, yet the NMR experiments were measured in phosphate buffer. (a) R_2/R_1 data for 6 of these proteins which bind RNA or DNA. (b) All positively-charged sidechains (arginine, lysine) are depicted in stick form. Many of the residues with experimental exchange (orange) that are not predicted by Dyna-1 are on the surface of these proteins and can be explained by phosphate binding. (c) Proteins with low AUROC and high p(exchange) predicted in disordered N- or C- termini, or areas with characterized ps-ns motion, labeled in blue in Fig. 4b and here. In (b,c): structures are sized by Dyna-1 p(exchange) and colored by experimental labels as in Fig. 4.

Extended Data Figure 8: RelaxDB-CPMG datasets where dispersion data was available for all residues. Legend: Missing assignments (red star), prolines (purple "P"), exchange via R_{ex} calculated conventionally (light orange), residues with unsuppressed R_2 (light green), residues with no CPMG data due to peak overlap (gray). The color gradient represents B_1 field strength from lowest (red,

e.g., 50 Hz) to highest (purple, e.g., 1000 Hz). The AqAdk and CypA datasets are annotated with residue groups reported in refs. [49] and [46], respectively.

Arg Kin   RNase A   HT PDZ domain

Extended Data Figure 9: Additional [15]N CPMG datasets with no dispersion data published for all residues (cf. Figure 5e).

Extended Data Figure 10: Adjusting native logits from Dyna-1 to account for class imbalance. (a) We follow the practice described in ref. [69] to adjust probabilities calculated from logits to maximize the "balanced accuracy", i.e. accuracy averaged across both classes, rather than accuracy averaged over all datapoints. We set our prior for the missing class based on the training data, π, as 5%. Comparing the adjusted curve (black) to the probability densities in (b) demonstrates how this intuitively matches where a discriminator between the two classes should be (at x ~ -3). (b) Histograms and probability densities of logits predicted for residues labeled as "present" and "missing" over train and validation sets. (c) Densities of resulting p(missing) if calculated using native logits or adjusted logits.