Is Britain still Great for Pine marten? A Habitat Suitability Assessment

Ella Lewis[1], Amy Fitzmaurice[1], Rachel Gardner[3], Suzanne Kynaston[2], Neal Armour-Chelu[4], Dave J.I. Seaman[5], Kirsty Swinnerton[1], Lawrence Ball[1]

[1]Kent Wildlife Trust Group, Tyland Barn, Chatham Rd, Sandling, Maidstone ME14 3BD

[2]Kent Wildwood Trust, Wealden Forest Park, Canterbury Rd, Herne Common, Herne Bay CT6 7LQ

[3]Forestry England, National Office, 620 Bristol Business Park, Coldharbour Lane, BS16 1EJ

[4]Forestry England, East Forest District, Santon Downham, Brandon, Suffolk IP27 0TJ

[5]Durrell Institute of Conservation and Ecology, School of Natural Sciences, University of Kent, Canterbury, UK

Abstract

Context

Species distribution models are used to predict habitat suitability for a species, by quantifying the environmental characteristics that allow a species to occupy a geographical area. The abundance and range of pine marten *(Martes martes)* has declined substantially in Great Britain, with remaining populations restricted to Scotland.

Objectives

Here, we perform species distribution modelling using BIOMOD2 platform to determine habitat

suitability, and inform the identification of potential reintroduction sites for pine marten in Great

Britain.

Methods

Using a global range dataset of 4,189 occurrences and seven environmental variables,

ensemble species distribution models were used to predict habitat suitability across Europe at 1

km resolution and Great Britain at 100 m resolution.

Results

Across the extent of both Europe and Britain, results indicate high suitability in areas with woody

vegetation cover in low topographic positions, and notably low in urban areas and extensive

areas of arable land. In Britain, high habitat suitability is identified across substantial areas in the

South East of England, parts of South West of England, East Yorkshire and Gloucestershire,

with pockets of suitable habitat along the West Coast of Britain. The results indicate that

elevation and land cover are important drivers of suitability.

Conclusion

Habitat suitability modelling at a high resolution of 100m proves effective for informing potential

reintroduction sites for pine marten in Britain. We also demonstrate the importance of using

occurrence data from pine martens' global range to predict optimal habitat suitability.

Introduction

The Pine marten *(Martes martes)* is a mustelid species that typically occupies woodland habitats, distributed across mainland Europe extending to Western Siberia (Buskirk, 1992; Delibes,1983). In Britain, pine marten were once abundant, however their population suffered severe decline during the late 1800s due to habitat loss and hunting practices, particularly predator control for game birds (Langley and Yalden, 1977). The species is classified as critically endangered in England and Wales according to the Red List for British Mammals (Mathews and Harrower, 2020). It is a priority species in the UK Biodiversity Action Plan and has been fully protected since 1988 under Schedule 5 of the Wildlife and Countryside Act 1981. The fragmented populations that have persisted are restricted to north-west Scotland and the north of England and Wales. As an opportunistic omnivore, their diet is comprised of small mammals, particularly voles, small birds, berries and insects (Twining et al., 2019; Jedrzejewski et al., 1993). They play an important role within a woodland ecosystem by providing several functions including seed dispersal and balancing prey populations as a mesopredator (Paine, 1969; Hastings et al., 2007). For example, in Scotland and Ireland, they have been shown to suppress non-native invasive grey squirrel (*Sciurus carolinensis*) populations (Sheehy et al., 2018; Vincent Wildlife Trust, 2020, Reilly and Lawton, 2025). This in turn, allows indirect benefits for vegetation recovery due to reductions in tree damage, and red squirrel (*Sciurus vulgaris*) population recovery by reducing interspecific competition with grey squirrel (Sheehy et al., 2018; Schaumann and Heinken, 2002; Crowley et al., 2018).

Pine martens are considered an indicator species for woodland ecosystems where their presence can denote a biologically diverse ecosystem in good condition (Buskirk, 1992; DEFRA, 2021). As a flagship species, pine marten can act as an effective conservation tool, not only for reintroduction approaches, but also to promote conservation of woodland habitats.

Woodland, defined as land with at least 20% tree canopy cover (Forestry Commission, 2024),

accounts for only 13% of land in Britain (Forest Research, 2022), compared to the European

average of 37% (Forestry Research 2010), therefore woodland creation and restoration is a

national priority (Defra, 2022).

Reintroduction projects can be a key tool for species recovery, and pine marten reintroductions

have already occurred across Britain in Gloucestershire, Wales, Devon and Cumbria

(MacPherson et al., 2014; Stringer et al., 2015). Reintroductions require extensive research,

primarily around social and ecological feasibility pre-release, population genetics, disease risk,

dispersal potential, and post-release monitoring (Kirkwood, 2013; Klein and Arts, 2021;

Armstrong and Seddon, 2008, Seddon and Armstrong, 2019; IUCN, 2013).

Species distribution models (SDM), sometimes referred to as environmental niche models

(ENM) or habitat suitability models (HSM), are a useful tool to identify suitable habitats or

environments for a species. SDMs estimate the relationship between observed species

occurrences and the environmental conditions at these locations, to characterise the

environmental conditions that are suitable for that species. These estimated relationships are

then used to identify where suitable environments are distributed in space, in the form of a

habitat suitability map (Anderson et al., 2003; Elith and Leathwick, 2009).

A key application for SDMs is to identify sites with high suitability for a species, as an

informative tool for conservation planning (Guisan et al., 2006). For example, they can be used

to assist in identifying priority areas for conservation management that will have the most

impact, based on species richness and suitability. SDMs are also used to predict species'

distributional changes over time and space (Mammola et al., 2021), for instance, to understand

how species will respond to future climatic conditions (Raes and Aguirre-Gutiérrez, 2018).

SDMs generally use a single statistical technique to predict habitat suitability for a species.

However, the field of habitat suitability mapping has evolved rapidly, giving access to more

complex and potentially reliable approaches (Meller et al., 2014). One approach is ensemble

species distribution models (eSDMs), which use an assemblage of statistical models to produce

suitability predictions. This method has been shown to improve the accuracy of model

predictions compared to individual models (Kwon, 2014; Araújo and New, 2007; Hao et al.,

2020). Given that there is often significant variation in the predictions across model techniques

and there is no single outstanding model algorithm, it is sensible to combine several, well-

performing techniques in order to generate accurate predictions (Meller et al., 2014; Li and

Wang, 2012). In addition, ensemble forecasting has become more widely used for its ability to

combat the limitations of individual algorithms, whilst allowing for comparisons between the

performance of different algorithms (Araújo & New, 2007).

Here, we aim to use eSDMs to identify areas in Britain with sufficient suitable habitat to support

viable breeding populations of pine marten and to guide the selection of potential reintroduction

sites. Recently founded populations in southern England are small, fragmented and highly

vulnerable to environmental, genetic and demographic factors (Birks et al., 2025, MacPherson

et al., 2014; Gilpin and Soulé, 1986). Interventions, including translocation and reintroduction

projects, are therefore necessary for the long-term viability of populations in Britain

(MacPherson and Wright, 2021). Previous feasibility assessments for the reintroduction of pine

marten to Britain have used occurrence records from populations in Scotland, Ireland and the

Netherlands to predict habitat suitability in Britain at 1 km resolution (MacPherson et al., 2014;

MacPherson et al., 2024; Scopes et al., 2025). Our methodology uses occurrence data from

across the global range of pine marten, to predict suitability at 100 m resolution. Pine marten

populations in the UK and Ireland have likely been driven to suboptimum, or 'refuge', areas due

to anthropogenic disturbance, with loss of forest cover as a main contributor (European

Environment Agency, 2018a). Across mainland Europe, we assume that pine marten can more

easily disperse and occupy their preferred habitat due to greater forest cover and

connectedness. Consequently, we expect our model predictions, using global occurrences and

a high resolution reflecting the habitat scale used by pine martens, to accurately identify areas

of optimal habitat suitability. resolution capturing the habitat scale pine martens operate at, will

reflect the optimal habitat suitability requirements of pine marten.

Methods

We assessed pine marten habitat suitability at two distinct spatial scales and extents. First, we

fit eSDMs across Europe at 1 km resolution to represent the species' global distribution, before

creating fine scale maps for Great Britian at 100 m resolution to better inform potential release

sites for future reintroductions.

<u>Occurrence data</u>

The occurrence data was collated from the Global Biodiversity Information Facility (GBIF, 2023),

which holds a vast array of georeferenced species occurrence records, and the NBN Atlas

(NBN, 2023), the UK's main biodiversity data centre. GBIF provided 26,277 presence records

across Europe (https://doi.org/10.15468/dd.32w6qw), and NBN Atlas provided 11,048 records

across Britain and Ireland (Supplementary material 1). Although these databases hold large

datasets, many records may be subject to bias from uneven sampling or inaccurate

georeferencing (Glon et al., 2017; Beck et al., 2014). For example, certain areas may be the

focus of targeted pine marten surveys (Austin, 2007).

Data cleaning is essential so that models are built on robust and spatially-uniform data. To clean

the GBIF data we followed the methodology of Zizka et al (2019) and used the

'CleanCoordinates' package v2.0-11 (Zizka et al., 2019). This included removing records made

prior to 1945 (which have low accuracy), removing duplicate records, removing coordinates

assigned to country and province centroids, removing records with 0,0 coordinates, coordinates

at sea, and coordinates assigned to biodiversity institutions. We incorporated additional cleaning

steps to the remaining data including removing records with coordinate uncertainty above 1 km,

removing records of unknown source, records derived from fossil specimens, and records with

noted geospatial issues. After data cleaning, 32,000 out of 37,325 records were retained (Fig.

1).

Fig 1. PRISMA diagram illustrating the necessary cleaning steps with the number (n) of occurrences

remaining after each cleaning process.

Sampling bias can skew model accuracy, as it may not reflect the complete species' range of

environmental conditions and may over-represent environmental conditions associated with

regions of higher sampling (Kadmon et al., 2004; Anderson and Gonzalez, 2011; Duque-Lazo et

al., 2016). To reduce the influence of spatial bias, occurrence data should undergo spatial

thinning. To do so, the 'spThin' function in R was applied to the cleaned occurrences, which

used a randomised approach to remove occurrence points within 10 km of one another, making

sure the maximum number of occurrence points were returned (Aiello-Lammens et al., 2015).

We identified 10 km as an appropriate thinning distance by visually inspecting a variogram of

the environmental variables (Fig. 2). The variogram shows the variation in environmental

conditions with increasing Euclidian distance between points. Therefore, the distance at which

the variance ceases to increase and begins to plateau can inform the distance at which

occurrences are geospatially independent from one another, and therefore the distance to use

for thinning (Anderson and Gonzalez, 2011; Bio et al., 2002). After data cleaning and spatial

thinning, occurrences were clipped to the model area of Europe, retaining 4,189 occurrences for

use in the modelling (Fig. 3).

Fig 2. The variogram shows that the semivariance reached a plateau after 10 km from points. This 'practical range' was used to thin the occurrence data in order to remove spatial autocorrelation, but retain the most occurrence data.

Fig 3. The locations of occurrence records before (left) and after (right) cleaning and spatial thinning within the modelled area.

Environmental predictors

SDMs are based on the ecological niche concept, which posits that species occupy spatial

ranges with suitable environmental conditions for their existence. Thus, the environmental

variables used to predict suitability herein were chosen based on the ecological requirements of

pine marten. It is hypothesised that the suitability of habitat for pine marten is determined by

risks of mortality, food availability, and availability of denning sites (Stringer et al., 2018;

MacPherson et al., 2014). A total of fourteen environmental variables were chosen based on

these factors. Processing of environmental predictors was conducted in QGIS v3.22 (QGIS.org,

2021). All environmental variables were reprojected to ETRS89-extended /LAEA Europe,

resampled to a resolution of 1 km (nearest neighbour for categorical variables and bilinear for

continuous variables using WARP tool in QGIS), cropped to the same extent of Europe and

aligned. Four of these environmental variables were used for the British projection, whilst the

remainder were sourced from different datasets (Table 1).

Multi-collinearity was tested between environmental variables using Pearson's r-correlation test

with r >0.7 considered as unacceptably collinear. Collinear variables cannot be included in

statistical models because it would be impossible to determine the unique effect of each

variable (Dormann et al., 2013; Brun et al., 2019; Glon et al., 2017). This process led to the

removal of five environmental variables: average monthly maximum and minimum temperature

(°C) derived from WorldClim database v2.1 (Fick and Hijmans, 2017), average monthly

precipitation (mm) derived from WorldClim database v2.1 (Fick and Hijmans, 2017), and

northness and eastness derived from SRTM WorldClim database v2.1 (Fick and Hijmans,

2017). Removal of these variables was not a concern as average elevation, collinear to climatic

variables, was retained, whilst northness and eastness are unlikely to influence pine marten

habitat suitability.

Two environmental variables were removed from the eSDM due to spatial autocorrelation of

sampling efforts (sampling bias); distance to roads and urban areas. Distance to roads was

derived from a shapefile subset to major roads (A and B roads) using UN Spatial Data

Infrastructure Transport (UNSDI-T) v2 (CIESIN, 2013). Distance to urban areas was derived

from a shapefile from North American Cartographic Information Society (Patterson et al., 2012).

Both urban areas and major roads vector layers were rasterized, and distance was calculated

from the proximity (raster distance) tool, both derived from GDAL in QGIS.

This process resulted in a total of seven environmental variables used to inform suitability in the

model (Table 1): elevation, slope, topographic position index, normalised difference vegetation

index (NDVI), forest density, land cover, canopy height.

Environmental predictor	Source	Original resolution (m)	Resolution for European modelling (m)	Resolution for UK modelling (m)	Processing	Justification for inclusion
Elevation	SRTM DEM WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000			Elevation is strongly correlated with climatic conditions. Pine marten avoid thermal stresses and lack of prey abundance at high altitudes (Vergara et al., 2015).
Slope	WorldClim database v2.1	1000	1000		GDAL geoprocessing tool in QGIS.	Slope is closely linked to soil nutrients, water-holding capacity and erosion, which in

					turn influence vegetation communities and pine marten prey abundance (particularly small-rodents) and availability of denning sites (Krebs, 2009; Lee et al., 2020).
Topographic Position Index (TPI)	WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000	GDAL geoprocessing tool in QGIS.	TPI is used to classify slope position and landform types. It is a powerful proxy for local geology, hydrology, nutrient availability, and topoclimatic conditions which in turn influence ecological communities. High values represent hilltops and valley shoulders, whilst lower values represent valley beds and depressions (Ball and Tzanopoulos, 2020). TPI also influences the mobility of animals with implications for species interactions (Sultaire et al., 2022)..
Normalised Difference Vegetation Index (NDVI)	Landsat satellite imagery (USGS, 2022)	30	1000	Calculated in Google Earth Engine as the mean annual maximum greenest pixel from 2010 to 2022 (USGS, 2022).	NDVI indicates vegetation greenness (relative plant biomass). Pine marten prefer habitat with high vegetation cover fraction (Storch et al.,1990).
Forest cover density (%)	Copernicus land monitoring (European Environment Agency, 2018b)	100	1000	Secondary mapping product. Percentage tree density per pixel.	Pine marten are predominantly found in diversely-structured deciduous and coniferous woodland, which provide a combination of abundant food resources, well-insulated denning sites and protection from predators (Buskirk, 1992; Delibes,1983; Balestr ieri et al., 2016; Virgos et al., 2012).
Land cover	CORINE 2018 (European Environment Agency, 2018a)	100	1000	Consisting of 44 land cover type categories amalgamated into 17 broad categories that align as far as possible with UK recognised habitat nomenclature (Appendix 3).	Pine marten have been shown to avoid open habitat and urban areas (Brainerd and Rolstad, 2002; Storch et al.,1990; Vergara et al., 2015; O'Mahony, 2017). Certain land cover types support the dispersal of pine marten, for example riparian corridors and hedgerows allow range expansion in landscapes dominated by urbanisation and agriculture (Balestrieri et al., 2016). Land cover distinguishesconiferous, deciduous and mixed woodland categories.
Canopy height (mm)	GLAS and ICESat satellite imagery (Simard et al., 2011)	1000	1000	Secondary mapping product	Pine marten prefer mature woodland for protection from predators and for den and resting sites within the forest canopy (Brainerd and Rolstad, 2002).

Elevation	Japan Aerospace Exploration Agency, using ALOS satellite (JAXA, 2021)	30	100	
Topographic Position Index (TPI)	Japan Aerospace Exploration Agency, using ALOS satellite (Japan Aerospace Exploration Agency, 2021)	30	100	GDAL geoprocessing tool in QGIS.
Slope	Japan Aerospace Exploration Agency, using ALOS satellite (JAXA, 2021)	30	100	GDAL geoprocessing tool in QGIS.
Land cover	UKCEH 2020 (Morton et al., 2021)	25	100	Consisting of 21 land type categories, amalgamated into the same 17 broad land categories as the Europe raster (Appendix 4).

225   Table 1: Environmental predictors used in the modelling or removed after multicollinearity assessment,

226   with their justification for inclusion based on ecological relevance.

Modelling

Model algorithms:

To build the ensemble model, we used 9 algorithms within the BIOMOD2 v 4.2 package in R (R

Core Development Team, 2014; Thuiller et al., 2016).These were generalised linear model

(GLM), generalised boosted regression model (GBM), generalised additive model (GAM),

classification tree analysis (CTA), artificial neural networks (ANN), flexible discriminant analysis

(FDA), multivariate adaptive regression splines (MARS), random forest (RF), and maximum

entropy model (MaxEnt), using the default parameters specific to each algorithm.

Pseudo absences:

Most SDM algorithms require both presence and absence data. For many species, true

absence data is rarely available or unreliable and so presence-only models are used, wherein

artificial absence data, known as pseudo-absences, are generated (Barbet-Massin et al., 2012).

Pseudo-absences aid the model's ability to differentiate the environmental conditions under

which the species occurs (Brown and Yoder, 2015). We used a large number of pseudo

absences, equally weighted to presences (4,189), as has been shown to produce the most

accurate predictions (Barbet-Massin et al., 2012). Pseudo-absences were randomly generated

within the entire study area. We used BIOMOD's disk method to prohibit pseudo-absence

generation within 5 km of occurrence points, respective of a typical pine marten home range for

females (Stringer et al., 2018). This random assignment of geographically stratified pseudo-

absences is recommended for machine-learning and classification methods (Barbet-Massin et

al., 2012). To reduce bias towards a single distribution of pseudo-absences, five sets were

generated (Guisan et al., 2017). Prevalence was assigned as 0.5, where presences and

pseudo-absences have the same weighting in the model calibration process (Liu et al., 2005;

McPherson et al., 2004).

Model evaluation:

To test model accuracy when no independent data is available, cross-validation is performed,

where the data is randomly partitioned into a set of training data and testing data (Raes and

Aguirre-Gutiérrez, 2018). For our analysis, data was randomly partitioned to 70% for training

and 30% for testing using repeated split samples. This cross-validation process was repeated

10 times for each modelling run (Yu et al., 2024). We chose the widely used repeated random

cross-validation method. Model quality was evaluated based on kappa statistic (Cohen, 1960),

the area under the receiver operating characteristic curve (AUC), and the true skill statistic

(TSS) (Allouche, et al., 2006). Only single models with a TSS score greater than the upper

quartile (0.324) of TSS scores (0%: 0.218, 25%: 0.286, 50%: 0.307, 75%: 0.324, 100%: 0.368)

were retained for the final ensemble and ensemble projections (Tanaka et al., 2020; Hill et al.,

2017). TSS corresponds to the sum of sensitivity and specificity minus one (Barbet-Massin et

al., 2012). TSS scores can be classified according to Ben Rais Lasram et al., (2010) as

excellent (TSS > 0.8), good (TSS = 0.6–0.8), fair (TSS = 0.4–0.6), poor (TSS = 0.2–0.4), and no

predictive ability (TSS < 0.2). We chose TSS to evaluate model accuracy for inclusion in the

ensemble models as it is a threshold-dependent measure of model accuracy and independent

of prevalence (the ratio of presence to pseudo-absence data in the presence-absence

predictions) (Allouche et al., 2006).

We combined the single models into ensemble models based on mean, median and weighted

average. Mean and median produce the ensemble results by averaging or taking the median of

the single model probability of suitability values, respectively, whereas weighted mean weights

the individual models based on their cross-validation performance, in our case the TSS values

(Marmion et al., 2009; Thuiller et al., 2009). For the ensemble model evaluation, the same

evaluation statistics from the single models were used to assess model performance. The

method used to combine the models that performed best, should be used to present the

projection results. The ensemble model included 111 single models out of 450 individual

models.

To evaluate the relative importance of each environmental variable, a simple Pearson's

correlation *r* was calculated between predictions of the full model and a model with each

variable removed (Thuiller et al., 2009). We also generated species response curves, which

show how suitability for pine marten changes as a given environmental variable changes.

Habitat suitability projections:

We generated single and weighted mean ensemble model projections in Britain using the 100 m

resolution environmental rasters. Suitability index has been re-scaled to a 0 to 100 range, where

larger values indicate higher suitability.

Results

Our study used eSDM with 4,189 pine marten occurrences from across the species' range and

7 environmental variables to predict pine marten habitat suitability in Europe and Great Britain.

The results indicate that elevation, land cover and vegetation greenness (NDVI) are the

strongest predictors of habitat suitability. Across the extent of Europe at 1km and Britain at

100m, the eSDM results illustrate high suitability for pine marten in areas with woody vegetation

cover in low topographic positions, and notably low suitability in urban areas and extensive

areas of arable land.

The mapped relative probability of habitat suitability in Europe on a 1km scale shows markedly

low habitat suitability for pine marten at high altitudes in the Alps and the Scandinavian

Mountains, in heavily modified and farmed landscapes, in dry and arid zones including much of

Spain, and over open water bodies. Higher habitat suitability occurs in areas with woody

vegetation cover, in lowlands, low topographic positions, and notably along Europe's West coast

(Fig. 4).

The mapped relative probability of habitat suitability in Great Britain on a 100m scale shows

high suitability across substantial areas in the South East of England, South West of England,

East Yorkshire, and Gloucestershire, with pockets of suitable habitat along the West Coast of

Britain. Suitability is greatest in wooded landscapes and lower in urban and intensively farmed

areas. Substantial areas of the Scottish Highlands, North Pennines, Central Midlands and South

Lincolnshire show relatively low suitability (Fig. 5).

Fig 4. Habitat suitability map indicating pine marten suitability across Europe at a resolution of 1km x 1km. Suitability index ranged from 0 to 100, where larger values indicate higher suitability.

Fig 5. Habitat suitability map indicating pine marten suitability across Great Britain at a resolution of 100 x 100 m. Suitability index ranged from 0 to 100, where larger values indicate higher suitability.

TSS and ROC validation scores were used to assess the performance of the models. The mean

TSS value of the single models was 0.3046 (Min: 0.218, Q1: 0.286, Median: 0.307, Q3: 0.324,

Max: 0.368). The mean ROC value of the single models was 0.7087 (Min: 0.628, Q1: 0.698,

Median: 0.713, Q3: 0.723, Max: 0.756). The calibration scores for the mean ensemble model

were TSS = 0.425 and ROC = 0.786. The calibration scores for the median ensemble model

were TSS = 0.361 and ROC = 0.751. The calibration scores for the weighted-mean ensemble

model were TSS = 0.426 and ROC = 0.786.

The ensemble model variable importance results, averaged across the ensemble types, showed

elevation was the strongest predictor of suitability, as were land cover, vegetation greenness

(NDVI) and slope but to a lesser degree. TPI and canopy height were of relatively low

importance, with forest density being of least importance (Fig. 6).

Fig 6. Box plot showing the ensemble variable importance, across the mean, median and weighted

mean.

The response curves, averaged across the ensemble results (Fig. 7), indicate that pine marten

favour lower elevations, with a sharp decline in suitability at elevations above 500 m above sea

level (a.s.l.).

Woodland was favoured over other land cover types, with mixed woodland as the most suitable,

followed by broadleaf woodland and coniferous woodland (Fig. 7). Preference for arable and

pasture land cover was lower, with suitability values below 0.5. Urban greenspace,

sclerophyllous vegetation and peat bogs were the least favoured land cover type.

There is a fairly consistent medium probability of occurrence of pine marten across all values of

tree density, although at low densities (<25% tree cover) probability starts to decrease, and at

very low densities (<6% tree cover), the probability of occurrence drops sharply from about 0.42

to 0.36 (Fig. 7). NDVI showed a peak in suitability at 0.75 NDVI, where the probability of

occurrence exceeds 0.4 (Fig. 7) and drops suddenly to 0.25 suitability probability. The

probability of occurrence of pine marten increases with canopy height, until canopy height

reaches above 20 m and probability of occurrence drops from 0.4 at 25 m, to 0.3 at 35 m

(Fig.7).

The species response curve for TPI, indicates a preference of pine marten for lower topographic

positions within valleys and depressions (Fig. 7). The probability of occurrence of pine marten is

fairly consistent across all values of slope, apart from a slight decrease in very flat areas (Fig.7).

Fig 7. The response curves, averaged from the final ensemble models. Probability of suitability is on a 0-1 scale.

Discussion

In this study, eSDMs were used to map environmental suitability for pine marten in Great

Britain, to inform conservation and restoration of the species, with a focus on reintroduction site

suitability. Across the extent of Europe, the 1km eSDM results indicate high suitability in the

British isles, northern parts of continental Europe including France, Denmark and Germany and

lower altitudes of Scandinavia. Low suitability regions were often found at lower latitudes

including the Mediterranean regions such as large parts of the Iberian Peninsula and South East

Italy, as well as areas of high altitudes including the Alps, the North Western Highlands across

Scandinavia and extents of the Scottish Highlands (Fig. 4). Across Britain, the 100m eSDM

results indicated high suitability in most regions of the South East of England, parts of the South

West of England, East Yorkshire, Gloucestershire and eastern borders of East Anglia, with

pockets of suitable habitat on the West coast of England (Fig. 5). High suitability identified in the

South East of England could help to inform potential release areas of the South East Pine

Marten Restoration project, who are currently scoping a reintroduction to the High Weald

National Landscape. Here, we found areas of high suitability to be extensively wooded

environments in protected areas (National Landscapes and NPs), with lowest suitability in urban

and especially low-lying coastal urban areas (Fig. 6). Notable hotspots of suitability were in the

South Downs National Park, the High Weald National Landscape, and the western end of the

Kent Downs National Landscape which may serve as suitable reintroduction locations.

Model results show low suitability at high altitudes, particularly in the Scottish Highland regions.

Significant distributions of pine marten across Scotland does not necessarily correspond to

high-quality habitat, rather, it may represent remnant or refuge areas where populations persist

in sub-optimal habitats in Britain. By incorporating European occurrence data to build our

eSDMs, the model provides a more ecologically representative assessment of habitat suitability

for the species, and one which is relevant to expected poleward range shifts under future

climatic conditions.

The variable importance results showed that elevation is the strongest predictor of habitat

suitability for pine marten across its global range. Our results show substantially higher

probability of pine marten habitat suitability at lower elevations (< 500 m a.s.l). This preference

for lowland wooded habitats in Europe is likely to be linked to the availability of denning sites

which is higher in mature broadleaved woodland which typically occurs at lower elevations

(Harmer et al., 2010; Portoghesi, 2006). Coniferous woodlands and alpine tundra and meadows

generally occur at higher altitudes but lack the abundance of prey and denning sites required by

pine marten (Fonda et al., 2021). Based on our results we see that pine marten are unlikely to

occur in substantial populations at altitudes above 2000 m a.s.l.

Although its relative importance was low, the response curve indicates that pine marten prefer

lower topographic positions, representing areas within valleys and depressions. We expect this

preference is linked to the greater presence of woody vegetation in these locations compared to

sparser tree cover that tends to occur in higher topographic positions, on ridges and valley

shoulders. This can be seen in the South East of England where areas with highest suitability

for pine marten are in topographically complex, often forested, areas (Appendix 1).

Land cover is the second strongest predictor of suitability, with mixed, broadleaf and coniferous

woodland being highly favoured (Fig. 7). As forest-dwellers, pine marten are found primarily in

well-structured deciduous and coniferous woodland, with well-insulated denning sites and

abundant food resources, and with a complex understorey providing a low risk of predation

(Buskirk, 1992, Virgo et al., 2012; Delibes,1983; Balestrieri et al., 2015). The results show a

lower preference for coniferous woodland compared to mixed and broadleaf woodland, which

could reflect their preference towards more structurally diverse woodland systems and lower

canopy heights associated with avoidance to coniferous-dominant plantation habitat. Planted

coniferous woodland is often thinned and structurally uniform, which serves as a poor habitat for

pine marten due to the lack of diverse ground flora, foraging opportunities and denning sites.

Interestingly, arable, pasture and grassland land cover types were found to be moderately

favoured, contradicting previous models which have shown that they avoid open habitats

(Brainerd and Rolstad, 2002; Storch et al.,1990; Vergara et al., 2015; Wereszczuk and

Zalewski, 2015; O'Mahony, 2017). This may be explained by the species' ecological flexibility

and ability to occupy heterogenous landscapes. Evidence suggests that diverse habitats can

promote higher diversity of food resources than a continuous forest habitat (Rosalino and

Santos-Reis, 2009; Vergara et al., 2015). For instance, we know that pine marten travel up to

150 m from woodland edge to forage for prey species such as field voles in their favoured

habitats, provided there is some low-level vegetation cover (Caryl, 2008).

Linear or small habitat features will be underrepresented in our model due to the analytical

resolution of the modelling across both projections (1km and 100m). For example, shrub or

hedge features within a grassland or arable pixel will not be represented. These features can

serve as important facilitators of connectivity for individuals to move through the landscape to

access other patches of woodland, therefore our model likely underpredicts habitat suitability.

Notably, urban greenspace, sclerophyllous vegetation and peat bog land cover types were least

suitable. These land types are also rare. If repeating this study, one may consider further

reducing the total number of land cover types by conducting pairwise correlations between the

broad land cover types to combine correlated categories to simplify the model (Buckley and

Lundy, 2012).

Vegetation greenness (NDVI) was the third strongest predictor of suitability because pine

marten inhabit woodland and adjacent edge habitat. As well as providing denning sites and

shelter, vegetation greenness is also strongly correlated with prey abundance. For example,

rodent density is strongly correlated with net productivity of ground vegetation (Zalewski et al.,

2004). The response curve showed a peak in suitability around 0.55-0.8 NDVI which indicates

high vegetation cover fraction like that of tall grassland and woodland canopy.

Our results also suggested that pine marten will occupy a broad range of forest densities, with a

consistent medium probability of occurrence of pine marten across all values of tree density

(Fig. 7). We expect this links strongly with their ability to adapt to possibly sub-optimum

environments (MacPherson et al., 2014). For example, Southern European populations can be

found across ancient woodland, coppiced woodland and cultivated land with forest fragments,

where they are not restricted to dense woodland cover (Pittiglio, 1996). Even within countries,

the adaptation to varying vegetation structures is evident. For example, French populations

have genetically diverged to occupy varying woodland cover. In the Bressie region, individuals

occupy human-dominated landscapes with fragmented forest whilst in the Ariège region

individuals occupy densely forested habitat (Larroque et al., 2015).

The results show canopy height was a relatively unimportant variable, however suitability

increased with increasing canopy height, up to 20 m, after which it decreased with increasing

canopy height. Pine marten favour mature forest for protection from predators and access to

denning and resting sites within the forest canopy (Brainerd and Rolstad, 2002).. However, the

decrease in suitability in very tall forests (>20 m) may be due to fewer denning sites in less

structurally diverse, coniferous-dominant plantation habitats (Croose et al., 2016).

Mean monthly temperature (°C) and mean monthly precipitation (mm) were excluded from the

eSDM. Although these variables have strong predictive power at the European scale, the

climate of Great Britain is already highly suitable for pine marten. Consequently, including

broad-scale climatic predictors led to their undue dominance in the model and reduced its ability

to map fine-scale habitat suitability within Great Britain. In a previous model iteration with these

variables included, mean monthly temperature accounted for 37% of model influence and mean

monthly precipitation for 17%, making them the first and third most important predictors,

respectively. Model evaluation metrics were also slightly higher when these variables were

included (mean single-model validation TSS ≈ 0.43). However, mean monthly precipitation

showed collinearity with vegetation greenness (NDVI) and elevation, further justifying its

removal.Sampling bias is a major challenge for species distribution modelling, and we identified

disproportionate sampling effort in our dataset. Specifically, pine marten occurrence records

were more common near urbanised areas and roads, possibly linked to accessibility for

recorders (Araújo and Guisan, 2006). This could lead to spatial bias in the model,

misrepresenting the environmental conditions which reflect suitability. We therefore decided to

remove the distance to roads and urban areas variables to prevent the influence of spatial

autocorrelation. One may also argue that roads and urban areas should not be included as they

serve as barriers rather than predictors of suitable habitat. In other words, pine marten do not

deliberately choose to be close to or far from roads, rather, territory is chosen based on

predominantly the abundance of woodland for denning potential and availability of prey. The

impacts of roads and urban areas on habitat suitability and connectivity are better examined

using connectivity modelling to quantify the effects of movement barriers, which includes

analysing the risks of mortality and how it effects overall population viability. Whilst

comprehensive efforts have been made to reduce bias in the occurrence data, other causes of

bias may still be present. For instance, we identified low environmental suitability at high

altitudes above 2000 m, which may be due to a lower number of observations in these difficult

to reach locations.

Our models did not perform to a good or excellent standard according to Kappa, TSS and AUC

evaluation scores, which may be due to the large spatial extent of our analysis (Europe), which

can lead to the selection of a higher proportion of less informative background points (Barbet-

Massin et al., 2012). This can lead to difficulties detecting commission and omission errors

(false presences and false absences), leading to low-moderate model performance. Moreover,

climatic predictor variables were removed due to multi-collinearity, which had the strongest

predictive power. Due to these limitations, the suitability maps should be used to guide further

ecological feasibility research to assess suitability in the local context.

Raster data with higher resolutions than is typical for this type of study were used for a number

of reasons. Firstly, higher resolution environmental variables provide more accurate information

about the environment, whereas down-sampled environmental variables average across

neighbouring cells, often leading to a poor measure of local conditions. This is particularly

pertinent in highly fragmented landscapes, where environmental predictors that change over

small spatial distances may become meaningless if a coarse resolution averages a broad range

of values. Secondly, high-resolution projections provide suitability estimates for smaller land

areas, which is particularly important in highly fragmented landscapes such as in England.

Producing eSDM maps at 100m resolution, improves our ability to inform potential reintroduction

areas at a local scale.

To inform potential reintroduction locations, these results must be coupled with further research

into barriers to movement, such as infrastructure, to inform the viability of a reintroduction.

MacPherson and Wright's (2021) 'Long-term strategic recovery plan for pine martens in Britain'

report recommends further research into the effects of roads on risk of mortality in more

densely-populated parts of the Great Britain. Since male pine martens hold large territories,

ranging from 2 km^2 to up to 25 km^2 (Zalewski and Jędrzejewski, 2006; Balestrieri et al., 2010),

the requirement for connectivity between identified highly suitable habitat areas is essential for

their long-term establishment (McNicol et al., 2020) and should therefore also be investigated.

Connectivity modelling can also assess proximity to existing populations which is important for

ensuring genetic diversity and the long-term viability of any reintroduced populations.

Appendices

<u>Appendix 1:</u>

Habitat suitability map indicating pine marten suitability across regions of Scotland, England and Wales at

a resolution of 100 x 100 m. Suitability index ranged from 0 to 100, where larger values indicate higher

suitability.

<u>North Scotland</u>

546 South Scotland

557 Northern England

568   <u>Yorkshire</u>

579   <u>Central England</u>

590   <u>Wales</u>

594

601   <u>South West England</u>

605

612   <u>South East England</u>

623   <u>East Anglia</u>

634   <u>Appendix 2:</u>

635   Reclassification of CORINE land classes and UKCEH land classes to form 17 land classifications. NA

636   indicates the categories which were not available in the UKCEH layer.

UKCEH land class number	Alternative layer for UKCEH land class	CORINE land class number	Merged land class categories (combined CORINE and UKCEH)	Classification code of combined categories
20		1 to 9, 11	Urban	1
NA	OS Open Greenspace (Ordnance Survey, 2022)	10	Urban green	2
3		12, 19 to 22	Arable	3
NA		18	Pastures	4
1		23	Broadleaf	5
2		24	Conifer	6
NA		25	Mixed woodland	7
4 to 7		26	Natural grassland	8
9		27	Moor and heath	9
NA		28	Sclerophyllous vegetation	10
NA		29	Transitional woodland-scrub	11
12		30 to 34	Bare (bare rock/sparse veg/burnt area/glacial/perpetual snow)	12
8		35	Inland marsh	13
11		36	Peat bogs	14
19		37	Salt marsh	15

13, 15, 16		38 to 39, 42 to 44	Coastal	16
14		40 to 41	Freshwater	17

638   Appendix 3:

639   Ensemble model evaluation results from the ensemble model outputs, indicating model performance

640   across TSS, ROC and Kappa evaluations.

	filtered.by	algo	metric.eval	cutoff	sensitivity	specificity	calibration	validation	evaluation
1	TSS	EMmean	ROC	496.5	76.940	65.643	0.786	NA	NA
2	TSS	EMmean	TSS	491.0	78.133	64.327	0.425	NA	NA
3	TSS	EMmean	KAPPA	597.0	51.444	84.548	0.323	NA	NA
4	TSS	EMmedian	ROC	493.5	70.972	65.294	0.751	NA	NA
5	TSS	EMmedian	TSS	491.0	71.545	64.595	0.361	NA	NA
6	TSS	EMmedian	KAPPA	607.0	44.235	85.094	0.273	NA	NA
7	TSS	EMwmean	ROC	494.5	77.894	64.734	0.786	NA	NA
8	TSS	EMwmean	TSS	493.0	78.205	64.241	0.426	NA	NA
9	TSS	EMwmean	KAPPA	600.0	51.516	84.558	0.323	NA	NA

641   (row labels above correspond to lines 642–650)

653   Appendix 4:

654   Ensemble variable importance results from ensemble model outputs across mean, median and weighted

655   average evaluations.

	filtered.by	algo	expl.var	rand	var.imp
1	TSS	EMmean	elevation	1	0.424409
2	TSS	EMmean	canopy_height	1	0.036039
3	TSS	EMmean	forest_density	1	0.022745
4	TSS	EMmean	ndvi	1	0.184354
5	TSS	EMmean	slope	1	0.183473
6	TSS	EMmean	tpi	1	0.040790
7	TSS	EMmean	land_cover	1	0.210616
8	TSS	EMmean	elevation	2	0.419510
9	TSS	EMmean	canopy_height	2	0.036504
10	TSS	EMmean	forest_density	2	0.022720
11	TSS	EMmean	ndvi	2	0.184362
12	TSS	EMmean	slope	2	0.184106
13	TSS	EMmean	tpi	2	0.040971
14	TSS	EMmean	land_cover	2	0.209221
15	TSS	EMmean	elevation	3	0.414192
16	TSS	EMmean	canopy_height	3	0.036326
17	TSS	EMmean	forest_density	3	0.023042
18	TSS	EMmean	ndvi	3	0.185699
19	TSS	EMmean	slope	3	0.182867
20	TSS	EMmean	tpi	3	0.040130
21	TSS	EMmean	land_cover	3	0.209279
22	TSS	EMmedian	elevation	1	0.419994
23	TSS	EMmedian	canopy_height	1	0.035677

```
680   24        TSS EMmedian forest_density    1 0.022718
681   25        TSS EMmedian          ndvi     1 0.189380
682   26        TSS EMmedian         slope     1 0.176616
683   27        TSS EMmedian           tpi     1 0.039820
684   28        TSS EMmedian    land_cover     1 0.208859
685   29        TSS EMmedian     elevation     2 0.419096
686   30        TSS EMmedian canopy_height     2 0.035526
687   31        TSS EMmedian forest_density    2 0.022522
688   32        TSS EMmedian          ndvi     2 0.186160
689   33        TSS EMmedian         slope     2 0.177070
690   34        TSS EMmedian           tpi     2 0.039465
691   35        TSS EMmedian    land_cover     2 0.209999
692   36        TSS EMmedian     elevation     3 0.418853
693   37        TSS EMmedian canopy_height     3 0.035591
694   38        TSS EMmedian forest_density    3 0.022522
695   39        TSS EMmedian          ndvi     3 0.187979
696   40        TSS EMmedian         slope     3 0.175561
697   41        TSS EMmedian           tpi     3 0.038905
698   42        TSS EMmedian    land_cover     3 0.212062
699   43        TSS  EMwmean     elevation     1 0.422913
700   44        TSS  EMwmean canopy_height     1 0.036650
701   45        TSS  EMwmean forest_density    1 0.022307
702   46        TSS  EMwmean          ndvi     1 0.184124
703   47        TSS  EMwmean         slope     1 0.181986
704   48        TSS  EMwmean           tpi     1 0.040894
705   49        TSS  EMwmean    land_cover     1 0.208688
706   50        TSS  EMwmean     elevation     2 0.424065
707   51        TSS  EMwmean canopy_height     2 0.035704
708   52        TSS  EMwmean forest_density    2 0.022511
709   53        TSS  EMwmean          ndvi     2 0.185314
710   54        TSS  EMwmean         slope     2 0.183928
711   55        TSS  EMwmean           tpi     2 0.040741
712   56        TSS  EMwmean    land_cover     2 0.207191
713   57        TSS  EMwmean     elevation     3 0.420573
714   58        TSS  EMwmean canopy_height     3 0.036186
715   59        TSS  EMwmean forest_density    3 0.022882
716   60        TSS  EMwmean          ndvi     3 0.186369
717   61        TSS  EMwmean         slope     3 0.182078
718   62        TSS  EMwmean           tpi     3 0.040748
719   63        TSS  EMwmean    land_cover     3 0.209171
```

References

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B. and Anderson, R.P. (2015). spThin: an R package for spatial
thinning of species occurrence records for use in ecological niche models. *Ecography*, 38, pp.541–545.

Allouche, O., Tsoar, A. and Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa and the
true skill statistic (TSS). *Journal of Applied Ecology*, 43(6), pp.1223–1232.

Anderson, R.P. and Gonzalez, I.J. (2011). Species-specific tuning increases robustness to sampling bias in models of species
distributions: an implementation with Maxent. *Ecological Modelling*, 222, pp.2796–2811.

Anderson, R.P., Lew, D., and Peterson, A.T. (2003). Evaluating predictive models of species' distributions: criteria for selecting
optimal models. *Ecological Modelling*, 162, pp.211–232.

Araújo, M.B. and Guisan, A. (2006). Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, 33(10),
pp.1677–1688.

Araújo, M.B. and New, M. (2007). Ensemble forecasting of species distributions. *Trends in Ecology & Evolution*, 22(1), pp.42–47.

Armstrong, D.P. and Seddon, P.J. (2008). Directions in reintroduction biology. *Trends in Ecology and Evolution*, 23, pp.20–25.

Austin, M. (2007). Species distribution models and ecological theory: a critical assessment and some possible new approaches.
Ecological Modelling, 200(1–2), pp.1–19.

Balestrieri, A., Bogliani, G., Boano, G., Ruiz-González, A., Saino, N., Costa, S. and Milanesi, P. (2016). Modelling the distribution of
forest-dependent species in human-dominated landscapes: patterns for the pine marten in intensively cultivated lowlands. *PLOS*
ONE, 11(7), p.e0158203.

Balestrieri, A., Remonti, L., Ruiz-González, A., Gómez-Moliner, B.J., Vergara, M. and Prigioni, C. (2010). Range expansion of the
pine marten (*Martes martes*) in an agricultural landscape matrix (NW Italy). *Mammal Biology*, 75, pp.412–419.

Balestrieri A, Remonti L, Ruiz-Gonza´lez A, Zenato M, Gazzola A, Vergara M, Detorri E.E, Saino N, Capelli E, Go´mezMoliner BJ,
Guidali F, Prigioni C (2015) Distribution and habitat use by pine marten Martes martes in a riparian corridor crossing intensively
cultivated lowlands. Ecological Research, 29(6), pp.1-10

Ball, S. and Tzanopoulos, J. (2020). Interplay between topography, fog and vegetation in the central South Arabian mountains
revealed using a novel Landsat fog detection technique. *Remote Sensing in Ecology and Conservation*, 6(4), pp.498–513.

Barbet-Massin, M., Jiguet, F., Albert, C.H. and Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: how,
where and how many? *Methods in Ecology and Evolution*, 3(2), pp.327–338.

Beck, J., Böller, M., Erhardt, A. and Schwanghart, W. (2014). Spatial bias in the GBIF database and its effect on modelling species'
geographic distributions. *Ecological Informatics*, 19, pp.10–15.

Ben Rais Lasram, F., Guilhaumon, F., Albouy, C., Somot, S., Thuiller, W. and Mouillot, D. (2010). The Mediterranean Sea as a "cul-
de-sac" for endemic fishes facing climate change. *Global Change Biology*, 16, pp.3233–3245.

Birks, J., Johnstone, S., Snell, E. and MacPherson, J. (2025). What is the future for the *Martes* complex (*Guloninae*) in the face of
climate change and ecological breakdown? *Stacks Journal*, 25001. https://doi.org/10.60102/stacks-25006

Bio, A.M.F., Becker, P.T., Bie, E.D. and Wassen, M.J. (2002). Prediction of plant species distribution in lowland river valleys in
Belgium: modelling species response. *Biodiversity and Conservation*, 11(12), pp.2189–2216.

Brainerd, S.M. and Rolstad, J. (2002). Habitat selection by Eurasian pine martens *Martes martes* in managed forests of southern
boreal Scandinavia. *Wildlife Biology*, 8(4), pp.289–297.

Brown, J.L. and Yoder, A.D. (2015). Shifting ranges and conservation challenges for lemurs in the face of climate change. *Ecology*
and Evolution, 5(6), pp.1131–1142.

Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R.O., Wang, Z. and Zimmermann, N.E. (2019). Model complexity affects
species distribution projections under climate change. *Journal of Biogeography*, 47(1), pp.130–142.

Buckley, D.J. and Lundy, M. (2012). The current distribution and potential for future range expansion of feral ferret *Mustela putorius*
furo in Ireland. *European Journal of Wildlife Research*, 59(3), pp.323–330.

Buskirk, S.W. (1992). Conserving circumboreal forests for martens and fishers. *Conservation Biology*, 6, pp.318–320.

Caryl, F.M. (2008). *Pine marten diet and habitat use within a managed coniferous forest*. PhD thesis, University of Stirling. Available
at:

https://dspace.stir.ac.uk/bitstream/1893/2261/1/Fiona%20Mae%20Caryl.%20pine%20marten%20Diet%20%26%20Habitat%20Use
%20Within%20A%20Managed%20Coniferous%20Forest.pdf (Accessed 7 July 2023).

Center for International Earth Science Information Network (CIESIN) – Columbia University, and Information Technology Outreach
Services (ITOS) – University of Georgia. (2013). *Global Roads Open Access Data Set, Version 1 (gROADSv1).* NASA SEDAC.
Available at: https://doi.org/10.7927/H4VD6WCT [Accessed 23 Mar 2023].

Cohen, J. (1960). A coefficient of agreement for nominal scales. *Educational and Psychological Measurement*, 20, pp.37–46.

Croose, E. et al. (2016). Den boxes as a tool for pine marten *Martes martes* conservation and population monitoring in a commercial
forest in Scotland. *Conservation Evidence*, 13, pp.57–61.

Crowley, S.L., Hinchliffe, S. and McDonald, R.A. (2018). Killing squirrels: exploring motivations and practices of lethal wildlife
management. *Environment and Planning E: Nature and Space*, 1, pp.120–143.

DEFRA (2021) *UK Biodiversity Indicators 2021*. Available at: https://data.jncc.gov.uk/data/31925413-3fa4-4382-962f-
6ac2c168d542/ukbi2021-summary-booklet.pdf

DEFRA. (2022). *Keepers of Time: Ancient and Native Woodland and Trees Policy in England.* Available at:
https://www.gov.uk/government/publications/keepers-of-time-ancient-and-native-woodland-and-trees-policy-in-england [Accessed
28 Mar 2023].

Delibes, M (1983) Interspecific competition and the habitat of the stone marten Martes foina (Erxleben 1777). Environmental
Science. 174, pp. 229–231

Dormann, C.F., Elith, J., Bacher, S. and Lautenbach, S. (2013). Collinearity: a review of methods to deal with it and a simulation
study evaluating their performance. *Ecography*, 36(1), pp.27–46.

Duque-Lazo, J., van Gils, H., Groen, T.A. and Navarro-Cerrillo, R.M. (2016). Transferability of species distribution models: the case
of *Phytophthora cinnamomi* in Southwest Spain and Southwest Australia. *Ecological Modelling*, 320, pp.62–70.

Elith, J. and Leathwick, J.R. (2009). Species distribution models: ecological explanation and prediction across space and time.
Annual Review of Ecology, Evolution, and Systematics, 40(1), pp.677–697.

European Environment Agency. (2018a). *CORINE Land Cover 2018 – Copernicus Land Monitoring Services.* Available at:
https://land.copernicus.eu/pan-european/corine-land-cover/clc2018

European Environment Agency. (2018b). *Tree Cover Density 2018 – Copernicus Land Monitoring Services.* Available at:
https://land.copernicus.eu/pan-european/high-resolution-layers/forests/tree-cover-density/status-maps/tree-cover-density-2018

Fick, S.E. and Hijmans, R.J. (2017). WorldClim 2: new 1 km spatial resolution climate surfaces for global land areas. *International*
Journal of Climatology, 37(12), pp.4302–4315.

Fonda, F., Chiatante, G., Meriggi, A.. (2021) Spatial distribution of the pine marten (*Martes martes*) and stone marten (*Martes foina*)
in the Italian Alps. Mammal Biology 101, pp.345–356

Forest Research. (2022). *Woodland Statistics.* Available at: https://www.forestresearch.gov.uk/tools-and-
resources/statistics/statistics-by-topic/woodlandstatistics/ [Accessed 31 Jan 2023].

Forest Research (2010) Forestry Facts & Figures. Available at: https://cdn.forestresearch.gov.uk/2010/08/fcfs210.pdf [Accessed 31
Jan 2023].

Forestry Commission. (2024). *Definition of Trees and Woodland.* Available at:
https://www.gov.uk/government/publications/definition-of-trees-and-woodland [Accessed 27 Nov 2024].

GBIF Secretariat. GBIF Backbone Taxonomy. Checklist dataset. Available at: https://www.gbif.org/dataset/d7dddbf4-2cf0-4f39-
9b2a-bb099caae36c2023-09-01. [Accessed 28 Mar 2023]

Gilpin, M.E. and Soulé, M.E. (1986) Minimum viable populations: processes of species extinction. Conservation Biology: the science
of scarcity & diversity (eds M.E. Soulé & M.A. Sunderland), pp. 19-34. Sinauer Associates.

Glon, H.E., Heumann, B.W., Carter, J.R., Bartek, J.M. and Monfils, A.K. (2017). The contribution of small collections to species
distribution modelling: A case study from Fuireneae (Cyperaceae). Ecological Informatics, 42, pp.67–78.

Guisan A, Lehmann A, Ferrier S, Austin M, Overton JMC, Aspinall R, Hastie T. (2006) Making better bio-geographical predictions of
species' distributions. Journal of Applied Ecology. 43, 386–392.

Guisan, A., Thuiller, W. and Zimmermann, N. (2017). *Habitat Suitability and Distribution Models: With Applications in R.* Cambridge:
Cambridge University Press.

Hao, T., Elith, J, Lahoz-Monfort, J.J.,Guillera-Arroita, G. (2020) Testing whether ensemble modelling is advantageous for
maximising predictive performance of species distribution models. Ecography 2020, 43, pp.549–558.

Harmer, R., Kerr, G. and Thompson, R. (2010) Managing Native Broadleaved Woodland. The Stationary Office, Edinburgh

Hastings, A. et al. (2007). Ecosystem engineering in space and time. *Ecology Letters*, 10, pp.153–164.

Hill, L., Hector, A., Hemery, G., Smart, S., Tanadini, M., & Brown, N. (2017). *Ecology and Evolution*, *7*(4), pp.1043–1056.
doi:10.1002/ece3.2661

IUCN. (2013). *Guidelines for Reintroductions and Other Conservation Translocations.* Version 1.0. Gland, Switzerland: IUCN
Species Survival Commission.

Jedrzejewski, W., A. Zalewski and B. Jedrzejewska, (1993) Foraging by pine marten Martes martesin relation to food
resources in Bialowieza National Park, Poland. Acta Theriologica, 38, pp.405-426

Kadmon, R, Farber, O, and Danin, A (2004). Effect of roadside bias on the accuracy of predictive maps produced by bioclimatic
models. – Ecological Applications. 14, pp. 401–413.

Kirkwood, J. (2013). Guidelines for reintroductions and other conservation translocations. Animal Welfare, 22(4), pp.489–490

Klein, L. and Arts, K. (2021). Public participation in decision-making on conservation translocations: the importance and limitations
of a legislative framework. *Restoration Ecology*, 30(1).

Krebs, C.J. (2009). *Ecology: The Experimental Analysis of Distribution and Abundance.* 6th edn. Pearson Benjamin Cummings, San
Francisco, CA.

Kwon, H. S. (2014) 47. Species Distribution Modeling of Endangered Mammals for Ecosystem Services Valuation - Focused on
National Ecosystem Survey Data. Journal of the Korean Society of Environmental Restoration Technology 17(1) pp. 111-122

Lariviere, S. and Jennings, A.P. (2009). European pine marten (*Martes martes*). In: Wilson, D.E. and Mittermeier, R.A. (eds)
Handbook of the Mammals of the World: Volume 1, Carnivores. Lynx Edicions, Barcelona, Spain.

Langley, P.J.W. and Yalden, D.W. (1977). The decline of the rarer carnivores in Great Britain during the nineteenth century.
Mammal Review, 7, pp.95–116.

Lee, J.-K., Hwang, H.-S., Eum, T.-K., Bae, H.-K. and Rhim, S.-J. (2020). Cascade effects of slope gradient on ground vegetation
and small-rodent populations in a forest ecosystem. Animal Biology, 70(2), pp.203–213.

Li, X. and Wang, Y. (2012). Applying various algorithms for species distribution modelling. *Integral Zoology*, 8, pp.124–135.

Liu, C., Berry, P.M., Dawson, T.P. and Pearson, R.G. (2005). Selecting thresholds of occurrence in the prediction of species
distributions. Ecography, 28(3), pp.385–393.

MacPherson, J. and Wright, P. (2021). *Long-term Strategic Recovery Plan for Pine Martens in Britain.* The Vincent Wildlife Trust,
Natural England, NatureScot.

MacPherson, J. (2014) Feasibility assessment for reinforcing pine marten numbers in England and Wales. Vincent Wildlife Trust.
Available at: https://www.vwt.org.uk/wp-content/uploads/2015/04/Feasibility-Assessment-for-Reinforcing-pine-martens-in-England-
and-Wales.pdf [Accessed 28 Mar 2023]

MacPherson, J., Jetz, W. and Rogers, D.J. (2004). The effects of species' range sizes on the accuracy of distribution models.
Journal of Applied Ecology, 41, pp.811–823.

MacPherson, J., Wright, P., Schumaker, N., & Watkins, S. (2024). Use of multi-modelling methods to inform conservation and
reintroductions of pine marten *Martes martes* in Britain. *Stacks Journal*: 24004. https://doi.org/10.60102/stacks-24004

Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R.K., Thuiller, W., 2009. Evaluation of consensus methods in predictive species
distribution modelling. Divers. Distrib. 15 (1), pp.59–69

Mathews, F. and Harrower, C. (2020). *IUCN–compliant Red List for Britain's Terrestrial Mammals.* Natural England, Peterborough.

McNicol, C.M. et al. (2020). Postrelease movement and habitat selection of translocated pine martens *Martes martes*. *Ecology and*
Evolution, 10(11), pp.5106–5118.

McPherson J, Jetz W, Rogers D.J (2004) The effects of species' range sizes on the accuracy of distribution models: ecological
phenomenon or statistical artefact? Journal of Applied Ecology 41, pp.811–823

Meller L, Cabeza M, Pironon S, Barbet-Massin M, Maiorano L, Georges D, Thuiller W. (2014) Ensemble distribution models in
conservation prioritization: from consensus predictions to consensus reserve networks. Diversity Distribution. 20(3):309-321.

National Biodiversity Network (NBN) Trust (2023). The National Biodiversity Network (NBN) Atlas. Available at:
https://ror.org/00mcxye41 [Accessed 28 Mar 2023].

O'Mahony, D.T. (2017). A species distribution model for pine marten (*Martes martes*) in the least forested region of Europe.
Mammal Research, 62(2), pp.195–200.

Paine, R.T. (1969) A note on trophic complexity and community stability. The American Naturalist, 103. pp.91–93.

Patterson, Tom. Kelso, Nathaniel Vaughn. (2012). World Urban Areas, 1:10 million [Shapefile]. North American Cartographic
Information Society. Available at: https://maps.princeton.edu/catalog/stanford-xg070wh7159. [Accessed 23 Mar 2023]

Pittiglio, C. (1996). Differential use of food and habitat by sympatric carnivores in the eastern Italian Alps. *Journal of Zoology*, 75(2),
pp.173–184.

Portoghesi L, (2006). European Forest Types. Categories and types for sustainable forest management reporting and policy. Forest
3 (4): 462-463

QGIS Development Team. (2021). *QGIS Geographic Information System.* Open Source Geospatial Foundation Project.

R Development Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical
Computing.

Raes, N. and Aguirre-Gutiérrez, J.A. (2018). Modelling framework to estimate and project species distributions in space and time. In:
Hoorn, C., Perrigo, A. and Antonelli, A. (eds) *Mountains, Climate and Biodiversity.* John Wiley & Sons, Chichester, UK.

Reilly, E. and Lawton, C. (2025). *Royal Society Open Science*, 12(6). doi:10.1098/rsos.250661.

Rosalino, L. and Santos-Reis, M. (2009). Fruit consumption by carnivores in Mediterranean Europe. *Mammal Review*, 39, pp.67–78.

Schaumann, F. and Heinken, T. (2002). Endozoochorous seed dispersal by martens (Martes foina, M. martes) in two woodland
habitats. Flora - Morphology, Distribution, Functional Ecology of Plants, 197(5), pp.370–378.

Scopes, E. R., MacPherson, J., Wright, P. G., McNicol, C. M., Kingscott-Edmunds, J., Ham, C., … Bellamy, C. C. (2025). *Journal of
Applied Ecology*. doi:10.1111/1365-2664.70160

Seddon, P.J, and Armstrong, D.P (2019) The role of translocation in rewilding. In: du Toit, J.T., Pettorelli, N., Durant S.M. (eds)
Rewilding (pp. 303-324). Cambridge University Press, Cambridge

Sheehy, E., Sutherland, C., O'Reilly, C. and Lambin, X. (2018). The enemy of my enemy is my friend: native pine marten recovery
reverses the decline of the red squirrel by suppressing grey squirrel populations. *Proceedings of the Royal Society B: Biological
Sciences*, 285, 20172603.

Simard, M., Pinto, N., Fisher, J.B. and Baccini, A. (2011). Mapping forest canopy height globally with spaceborne lidar. *Journal of
Geophysical Research*, 116(G4).

Stringer, A. P., MacPherson, J., Carter, S., Gill, R., Ambrose-Oji, B. and Wilson, R. (2015). The Feasibility of Reintroducing pine
martens (Martes martes) to the Forest of Dean and Lower Wye Valley. Herefordshire: Forestry Commission
England, Gloucestershire Wildlife Trust, Vincent Wildlife Trust.

Stringer, A.P., MacPherson J., Carter, S., Gill, R., Ambrose-Oji., Wilson. R., Kelsall. P.,Feirn. W.G., Galbraith. L.C., Hilário, C.M.,

Parry, G., and Taylor, A. (2018). *The Feasibility of Reintroducing Pine Martens (Martes martes) to the Forest of Dean and Lower*

Wye Valley. Herefordshire: Forestry Commission England, Gloucestershire Wildlife Trust, Vincent Wildlife Trust

Storch I, Lindström E.R, Jounge,J (1990) Diet and habitat selection of the pine marten in relation to competition with the red fox.

Acta Theriologica 35, pp.311–320

Sultaire, S.M., Millspaugh, J.J., Jackson, P.J. and Montgomery, R.A. (2022). The influence of fine-scale topography on detection of

a mammal assemblage at camera traps in a mountainous landscape. *Wildlife Biology*, 2023(2).

Tanaka, K. R., Torre, M. P., Saba, V. S., Stock, C. A., & Chen, Y. (2020). *Diversity and Distributions*, *26*(8), 987–1001.

doi:10.1111/ddi.13069

Thuiller, W. et al. (2009). BIOMOD: a platform for ensemble forecasting of species distributions. *Ecography*, 32(3), pp.369–373.

Thuiller, D. Georges, R. Engler, F. Breiner biomod2: ensemble platform for species distribution modeling R package version, 3

(2016), pp. 3-7

Twining, J. P., Montgomery, I., Fitzpatrick, V., Marks, N., Scantlebury, D. M., & Tosh, D. G. (2019). *European Journal of Wildlife*

Research, *65*(3). doi:10.1007/s10344-019-1289-z

Vergara, M., Cushman, S.A., Urra, F. and Ruiz-González, A. (2015). Shaken but not stirred: multiscale habitat suitability modeling of

sympatric *Martes martes* and *Martes foina* across Europe. *Ecography*, 39, pp.545–556.

Vincent Wildlife Trust (2020) Red squirrels making comeback as return of native pine marten spells bad news for invasive grey

squirrel. Available at: https://www.vincentwildlife.ie/news/red-squirrels-making-comeback-as-return-of-native-pine-marten-spells-

bad-news-for-invasive-grey-squirrel

Virgos E, Zalewski A, Rosalino L, Mergey M (2012) Habitat ecology of genus Martes in Europe: a review of the evidences. In: Aubry

KB, Zielinski WJ, Proulx G, Buskirk SW (eds) Biology and conservation of martens, sables, and fishers: a new synthesis. Cornell

University Press, New York, pp. 255–266

Wereszczuk, and A, Zalewski, A. (2015) Spatial Niche Segregation of Sympatric Stone marten and pine marten– Avoidance of

Competition or Selection of Optimal Habitat? PLOS ONE. 10(10)

Yu B, Dai W, Li S, Wu Z, Wang J. (2024) A new threshold selection method for species distribution models with presence-only data:

Extracting the mutation point of the P/E curve by threshold regression. Ecological Evolution. 14(4):e11208.

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn, R.,

Svantesson, S., Wengström, N., Zizka, V. and Antonelli, A. (2019). CoordinateCleaner : Standardized cleaning of occurrence

records from biological collection databases. Methods in Ecology and Evolution, 10(5), pp.744–751.

Zalewski, A. and Jędrzejewski, W. (2006) Spatial organisation and dynamics of the pine marten Martes martes population in

Białowieza Forest (E Poland) compared with other European woodlands. Ecography, 29,pp. 31-43.

Zalewski, A., Jedrzejewski, W. and Jedrzejewski, B. (2004). Mobility and home range use by pine martens (Martes martes) in a

Polish primeval forest. Écoscience, 11(1), pp.113–122.