Is Britain still Great for Pine Marten? A Habitat Suitability Assessment

Ella Lewis[1] Amy Fitzmaurice[1] Kirsty Swinnerton[1], Rachel Gardner[2], Lawrence Ball[1], Neal

Armour-Chelu[3]

[1]Kent Wildlife Trust Group, Tyland Barn, Chatham Rd, Sandling, Maidstone ME14 3BD

[2]Forestry England, National Office, 620 Bristol Business Park, Coldharbour Lane, BS16 1EJ

[3]Forestry England, East Forest District, Santon Downham, Brandon, Suffolk IP27 0TJ

Abstract

Species distribution models are used to predict habitat suitability for a species, by quantifying

the environmental characteristics which allow a species to occupy a geographical area. The

abundance and range of Pine Marten *(Martes martes)* has declined substantially in Great

Britain, with the remaining populations restricted to Scotland. Here, we perform species

distribution modelling to determine habitat suitability, and identify potential reintroduction sites,

for Pine Marten in Great Britain. Using a global range dataset of 4,189 occurrences, and seven

environmental variables, ensemble species distribution models were used to predict habitat

suitability across Europe at 1 km resolution and Great Britain at 100 m resolution. High habitat

suitability was identified across substantial areas in the South East, East of England and South

West, with pockets of suitable habitat along the West Coast of Britain. The results indicate that

elevation and land cover are important drivers of suitability.

Introduction

Pine Marten *(Martes martes)* are a mustelid species that typically occupy woodland habitats,
distributed across mainland Europe extending to Western Siberia (Buskirk, 1992; Delibes,1983).
In Britain, Pine Marten were once abundant, however their populations suffered severe decline
during the late 1800's due to habitat loss and hunting practices, particularly predator control for
game birds (Langley and Yalden, 1977). The species is classified as critically endangered in
England and Wales according to the Red List for British Mammals (Mathews and Harrower,
2020). It is a priority species in the UK Biodiversity Action Plan and has been fully protected
since 1988 under Schedule 5 of the Wildlife and Countryside Act 1981. The remaining
populations are restricted to north-west Scotland and the north of England and Wales. As a
generalist apex predator, their diet is comprised of small mammals, particularly voles, small
birds, berries and insects (Twining et al., 2019). They play an important role within a woodland
ecosystem as a mesopredator, by providing several functions including seed dispersal and
balancing prey populations (Paine 1969; Hastings et al. 2007). In Scotland for example, they
have been shown to suppress non-native invasive Grey Squirrel (*Sciurus carolinensis*)
populations (Sheehy et al., 2018; Vincent Wildlife Trust, 2020). This in turn, allows indirect
benefits for vegetation recovery due to reductions in tree damage, and Red Squirrel (*Sciurus
vulgaris*) population recovery by reducing interspecific competition with Grey Squirrel (Sheehy et
al. 2018; Schaumann and Heinken, 2002; Crowley et al., 2018).

Pine Martens are considered a keystone species for woodland ecosystems, where their
presence can indicate a biologically diverse ecosystem in good condition (Buskirk, 1992; JNCC,
2021). As a flagship species, Pine Marten can act as an effective conservation tool, not only for
reintroduction approaches, but also to promote conservation of woodland habitats. Woodland

covers only 13% of land in Britain (Forest Research, 2022), compared to the European average

of 34% (Forestry Commission 2010), therefore woodland recovery is a national priority (Defra,

2022). Reintroduction programmes are a key strategy for population recovery and Pine Marten

reintroductions have already occurred across Britain in the Forest of Dean, Wales, Devon and

Cumbria (MacPherson et al., 2014; Stringer et al., 2015). However, they require extensive

research, primarily around social and ecological feasibility pre-release, population genetics,

disease risk, dispersal potential, and post-release monitoring (Kirkwood, 2013; Klein and Arts,

2021; Armstrong and Seddon, 2008; 2019).

This research aims to identify areas in Britain with sufficient suitable habitat to support viable

breeding populations of Pine Marten. Current populations in southern England are small and

fragmented and thus are considered highly vulnerable to environmental, genetic and

demographic factors (MacPherson et al., 2014) which could potentially lead to an extinction

vortex (MacPherson et al., 2014; Gilpin and Soulé, 1986). Interventions, including translocation

and reintroduction projects, are therefore necessary for the long-term viability of these

populations (MacPherson and Wright, 2021). Previous feasibility assessments for the

reintroduction of Pine Marten to Britain have used occurrence records from the populations in

Scotland, Ireland and the Netherlands only (Stringer et al., 2015; MacPherson et al., 2014). Our

methodology uses occurrence data indiscriminately, from across their global range, to predict

suitability at 100 m resolution in Britain. Pine Marten populations in the UK and Ireland have

likely been driven to suboptimum, or 'refuge', areas due to anthropogenic disturbance, with loss

of forest cover as a main contributor (European Environment Agency, 2018a). Across mainland

Europe, we assume that Pine Marten can more easily disperse and occupy the most suitable

areas. Therefore, we anticipate our model predictions, which are based on their global range,

will more closely reflect the optimal habitat suitability requirements of Pine Marten.

Commented [RG1]: Add disease risk to list?

Commented [RG2]: Inherently novel by what you're saying this research does.

Commented [RG3]: The UK doesn't include the Republic of Ireland. British Isles would be all encompassing, but isn't used often.

I've had to look this up previously to get it right for various things - I find the map on this page handy! https://www.britannica.com/story/whats-the-difference-between-great-britain-and-the-united-kingdom

Species distribution models (SDM), sometimes referred to as environmental niche models

(ENM) or habitat suitability models (HSM), are a useful tool to identify suitable habitats or

environments for a species. SDMs estimate the relationship between observed species

occurrences and the environmental conditions at these locations, to characterise the

environmental conditions that are suitable for that species. These estimated relationships are

then used to identify where suitable environments are distributed in space, in the form of a

habitat suitability map (Anderson et al., 2003; Elith and Leathwick 2009).

A key application for SDMs is to identify sites with high suitability for a species, as an

informative tool for conservation planning (Guisan et al., 2006). For example, they can be used

to assist in identifying priority areas for conservation management that will have the most

impact, based on species richness and suitability. SDMs are also used to predict species'

distributional changes over time and space (Mammola et al., 2021), for instance, to understand

how species will respond to future climatic conditions (Raes and Aguirre-Gutiérrez, 2018).

SDMs use a single statistical technique to predict habitat suitability for a species. However, the

field of habitat suitability mapping has evolved rapidly, giving access to more complex and

potentially reliable approaches (Meller et al., 2014). One approach is ensemble species

distribution models (eSDMs), which use an assemblage of statistical models to produce

suitability predictions. This method has been shown to improve the accuracy of model

predictions compared to individual models (Kwon, 2014; Araújo and New 2007; Hao et al.,

2020). Given that there is often significant variation in the predictions across model techniques

and there is no single outstanding model algorithm, it is sensible to combine several, well-

performing techniques in order to generate accurate predictions (Meller et al., 2014). In addition,

ensemble forecasting has become more widely used for its ability to combat the limitations of

individual algorithms, whilst allowing for comparisons between the performance of different

algorithms (Araújo & New, 2007).

This research aimed to use eSDMs to map environmental suitability for Pine Marten, to inform

the selection of optimum release site locations for reintroductions in Britain.

Methods

BIOMOD2 v 4.2, a widely used 'ensemble platform for species distribution modelling' available

in the R statistical programming language (R Development Core Team, 2020; Thuiller et al.,

2016), was used to build eSDMs with 9 algorithms, 4,189 occurrences and 7 environmental

variables, at a 1 km resolution across Europe. They were ensembled based on a threshold

model evaluation score, and weighted according to these scores, then ensembled predictions of

habitat suitability were projected across Europe at 1 km resolution and Great Britain at 100 m

resolution. The fine scale (100 m) predictions for Britain provide a more detailed view of habitat

suitability to facilitate understanding of habitat fragmentation and connectivity at local scales to

support conservation practice.

Environmental predictors

SDMs are based on the ecological niche concept, which posits that species occupy spatial

ranges with suitable environmental conditions for their existence. Thus, the environmental

variables used to predict suitability herein are chosen based on the ecological requirements of

Pine Marten. It is hypothesised that the suitability of a habitat for Pine Marten is determined by

risks of mortality, food availability, and availability of denning sites (Stringer et al., 2018;

MacPherson et al., 2014). Seven environmental variables were chosen based on these factors,

derived from open data sources (Table 1). Processing of environmental predictors was

conducted in QGIS v3.22 (QGIS.org, 2021). All environmental variables were reprojected to

ETRS89-extended /LAEA Europe, resampled to a resolution of 1 km (nearest neighbour for

categorical variables and bilinear for continuous variables using WARP tool in QGIS), cropped

to the same extent of Europe and aligned. Four of these environmental variables were used for

the British projection, whilst the remainder were sourced from different datasets (Table 1). Pre-

processing of these environmental variables followed the same aforementioned workflow.

Multicollinearity was tested between environmental variables using Pearson's r-correlation test

with r >0.7 considered as unacceptably collinear. Collinear variables cannot be included in

statistical models because it would be impossible to determine the unique effect of each

variable (Dormann et al., 2013; Brun et al., 2019; Glon et al., 2017). This process led to the

removal of four environmental variables: average monthly maximum and minimum temperature

(°C), and northness and eastness. Removal of these variables was not a concern, as average

mean annual temperature was retained, whilst northness and eastness were not assumed to

influence Pine Marten habitat preference.

During the iterative process, two environmental variables were removed from the eSDM due to

spatial autocorrelation of sampling efforts (sampling bias); distance to roads and urban areas.

Distance to roads was derived from a shapefile subset to major roads (A and B roads) using UN

Spatial Data Infrastructure Transport (UNSDI-T) v2 (CIESIN, 2013). Distance to urban areas

was derived from a shapefile from North American Cartographic Information Society (Patterson

et al., 2012). Both urban areas and major roads vector layers were rasterized, and distance was

calculated from the proximity (raster distance) tool, both derived from GDAL in QGIS.

Commented [ACN4]: Have these been removed from the analysis?
These are two of the most critical factors for risk of mortality.

Environmental predictor	Source	Original resolution (m)	Resolution for European modelling (m)	Resolution for UK modelling (m)	Processing	Justification for inclusion
Elevation	SRTM DEM WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000			Elevation is strongly correlated with climatic conditions. Pine Martens avoid thermal stresses and lack of prey abundance at high altitudes (Vergara et al., 2015).
Slope	WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000		GDAL geoprocessing tool in QGIS.	Slope is closely linked to soil nutrients, water-holding capacity and erosion, which in turn influence vegetation communities and Pine Marten prey abundance (particularly small-rodents) and availability of denning sites (Krebs, 2009; Lee et al., 2020).
Topographic Position Index (TPI)	WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000		GDAL geoprocessing tool in QGIS.	TPI is used to classify slope position and landform types. It is a powerful proxy for local geology, hydrology, nutrient availability, and topoclimatic conditions which in turn influence ecological communities. High values represent hilltops and valley shoulders, whilst lower values represent valley beds and depressions (Ball and Tzanopoulos, 2020). TPI also influences the mobility of animals with implications for species interactions (Sultaire et al., 2022)..
Normalised Difference Vegetation Index (NDVI)	Landsat satellite imagery (USGS, 2022)	30	1000		Calculated in Google Earth Engine as the mean annual maximum greenest pixel from 2010 to 2022 (USGS, 2022).	NDVI indicates vegetation greenness (relative plant biomass). Pine Martens prefer habitat with high vegetation cover fraction (Storch et al.,1990).
Forest cover density (%)	Copernicus land monitoring (European Environment Agency, 2018b)	100	1000		Secondary mapping product. Percentage tree density per pixel.	Pine Martens are predominantly found in diversely-structured deciduous and coniferous woodland, which provide a combination of abundant food resources, well-insulated denning sites and protection from predators (Buskirk, 1992;

Commented [RG5]: Include full term in table, not just abbreviation? Same for TPI.

Delibes,1983; Balestrieri et al., 2015; Virgos et al., 2012).

Variable	Source				Description	Ecological justification
Land cover	CORINE 2018 (European Environment Agency, 2018a)	100	1000		Consisting of 44 land cover type categories amalgamated into 17 broad categories that align as far as possible with UK recognised habitat nomenclature (Appendix 3).	Pine Marten have been shown to avoid open habitat and urban areas (Brainerd and Rolstad, 2002; Storch et al.,1990; Vergara et al., 2015). Certain land cover types support the dispersal of Pine Marten, for example riparian corridors and hedgerows allow range expansion in landscapes dominated by urbanisation and agriculture (Balestrieri et al., 2016). Land cover distinguishes coniferous, deciduous and mixed woodland categories.
Canopy height (mm)	GLAS and ICESat satellite imagery (Simard et al., 2011)	1000	1000		Secondary mapping product	Pine Marten prefer taller spruce-dominated forest for protection from predators and for den and resting sites within the forest canopy (Brainerd and Rolstad, 2002).
Average monthly temperature (°C)	WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000		Averaged across 1970 to 2000.	Pine Martens deal poorly with thermal stress, with small fat reserves, poor fur insulation and high low critical temperatures, so avoid areas of extreme temperature ranges (Harlow, 1994). Snow also reduces prey catching efficiency (Jedrzejewski et al., 1993).
Average monthly precipitation (mm)	WorldClim database v2.1 (Fick and Hijmans, 2017)	1000	1000		Averaged across 1970 to 2000.	Rainfall strongly influences vegetation abundance, which in turn influences prey abundance.
Elevation (DEM)	Japan Aerospace Exploration Agency, using ALOS satellite (JAXA, 2021)	30		100		
Topographic Position Index (TPI)	Japan Aerospace Exploration Agency, using ALOS satellite (Japan Aerospace Exploration Agency, 2021)	30		100	GDAL geoprocessing tool in QGIS.	
Slope	Japan Aerospace Exploration Agency, using ALOS satellite (JAXA, 2021)	30		100	GDAL geoprocessing tool in QGIS.	
Land cover	UKCEH 2020 (Morton et al., 2021)	25		100	Consisting of 21 land type categories, amalgamated into the same 17 broad	

Commented [LB6]: Is this correct?

Commented [RG7]: How did you resample the cell size, from 30/25 to 100 m?

land categories as
the Europe raster
(Appendix 4).

Table 1: Environmental predictors used in the modelling or removed after multicollinearity assessment,

with their justification for inclusion based on ecological relevance.

<u>Occurrence data</u>

The occurrence data was collated from the Global Biodiversity Information Facility (GBIF, 2023),

which holds a vast array of georeferenced species occurrence records, and the NBN Atlas

(NBN, 2023), the UK's main biodiversity data centre. GBIF provided 26,277 presence records

across Europe and NBN Atlas provided 11,048 records across Britain and Ireland. Although

these databases hold large datasets, many records may be subject to bias from uneven

sampling or inaccurate georeferencing (Glon et al., 2017; Beck et al., 2014). For example,

certain areas may be the focus of targeted Pine Marten surveys (Austin, 2007).

Data cleaning is essential so that models are built on robust and spatially-uniform data. To clean

the GBIF data we followed the methodology of Zizka et al (2019) and used the

'CleanCoordinates' package v2.0-11 (Zizka et al., 2019). This included removing records made

prior to 1945 (which have low accuracy), removing duplicate records, removing coordinates

assigned to country and province centroids, removing records with 0,0 coordinates, coordinates

at sea, and coordinates assigned to biodiversity institutions. We incorporated additional cleaning

steps to the remaining data including removing records with coordinate uncertainty of above 1

km, , removing records of unknown source, records derived from fossil specimens, and records

with noted geospatial issues. After data cleaning, 32,000 out of 37,325 records were retained

(Figure 1).

Figure 1: PRISMA diagram illustrating the necessary cleaning steps with the number (n) of occurrences

remaining after each cleaning process.

Sampling bias can skew model accuracy, as it may not reflect the complete species' range of

environmental conditions and may over-represent environmental conditions associated with

regions of higher sampling (Kadmon et al. 2004; Anderson and Gonzalez, 2011; Duque-Lazo et

al., 2016). To reduce the influence of spatial bias, occurrence data should undergo spatial

thinning. To do so, the 'spThin' function in R was applied to the cleaned occurrences, which

used a randomised approach to remove occurrence points within 10 km of one another, making

sure the maximum number of occurrence points were returned (Aiello-Lammens et al., 2015).

We identified 10 km as an appropriate thinning distance by visually inspecting a variogram of

the environmental variables (Figure 2). The variogram shows the variation in environmental

conditions with increasing Euclidian distance between points. Therefore, the distance at which

the variance ceases to increase and begins to plateau can inform the distance at which

occurrences are geospatially independent from one another, and therefore the distance to use

for thinning (Anderson and Gonzalez, 2011; Bio et al., 2002). After data cleaning and spatial

thinning, 4,189 occurrences were retained for use in the modelling (Figure 3).

193 Figure 2: The variogram shows that the semivariance reached a plateau after 10 km from points. This

194 'practical range' was used to thin the occurrence data in order to remove spatial autocorrelation, but

195 retain the most occurrence data.

Figure 3: The locations of occurrence records before (left) and after (right) cleaning and spatial thinning.

Modelling

Model algorithms:

For the ensemble model, we used 9 algorithms within the BIOMOD2 package in R (R Core

Development Team, 2014). These were generalised linear model (GLM), generalised boosted

regression model (GBM), generalised additive model (GAM), classification tree analysis (CTA),

artificial neural networks (ANN), flexible discriminant analysis (FDA), multivariate adaptive

regression splines (MARS), random forest (RF), and maximum entropy model (MaxEnt), using

the default parameters specific to each algorithm.

Pseudo absences:

Most SDM algorithms require both presence and absence data. For many species, true

absence data is rarely available or unreliable and so presence-only models are used, wherein

artificial absence data, known as pseudo-absences, are generated (Barbet-Massin et al., 2012).

They aid the model's ability to differentiate the environmental conditions under which the

species occurs (Brown and Yoder, 2015). We used a large number of pseudo absences, equally

weighted to presences (4,189), as has been shown to produce the most accurate predictions

(Barbet-Massin et al., 2012). Pseudo-absences are randomly generated within the entire study

area. We used BIOMOD's disk method to prohibit pseudo-absence generation within 5 km of

occurrence points, respective of a typical Pine Marten home range for females (Stringer et al.,

2018). This random assignment of geographically stratified pseudo-absences is recommended

for machine-learning and classification methods (Barbet-Massin et al., 2012). To reduce bias

towards a single distribution of pseudo-absences, five sets were generated (Guisan et al.,

2017). Prevalence was assigned as 0.5, where presences and pseudo-absences have the

same weighting in the model calibration process (Liu et al., 2005).

Model evaluation:

In order to test model accuracy when no independent data is available, cross-validation is

performed, where the data is randomly partitioned into a set of training data and testing data

(Raes and Aguirre-Gutiérrez, 2018). For our analysis, data was randomly partitioned to 70% for

training and 30% for testing using repeated split samples. This cross-validation process was

repeated 10 times for each modelling run (Yu et al., 2024). We chose the widely used repeated

random cross-validation method. Model quality was evaluated based on kappa statistic (Cohen,

1960), the area under the receiver operating characteristic curve (AUC), and the true skill

statistic (TSS) (Allouche, et al., 2006). Only single models with a TSS score greater than the

upper quartile (0.324) of TSS scores (0%: 0.218, 25%: 0.286, 50%: 0.307, 75%: 0.324, 100%:

0.368) were retained for the final ensemble and ensemble projections (Tanaka et al., 2020; Hill

Commented [EL12]: Lawrence is this the upper quantile? metric.select.thresh = unname(quantile(subset(get_evaluations(myBiomodModelOut), metric.eval == "TSS")$validation)[4])

et al., 2017). TSS corresponds to the sum of sensitivity and specificity minus one (Barbet-

Massin et al., 2012). TSS scores can be classified according to Ben Rais Lasram et al., (2010)

as excellent (TSS > 0.8), good (TSS = 0.6–0.8), fair (TSS = 0.4–0.6), poor (TSS = 0.2–0.4), and

no predictive ability (TSS < 0.2). We chose TSS to evaluate model accuracy for inclusion in the

ensemble models as it is a threshold-dependent measure of model accuracy and independent

of prevalence (the ratio of presence to pseudo-absence data in the presence-absence

predictions) (Allouche et al., 2006).

We combined the single models into ensemble models based on mean, median and weighted

average. Mean and median produce the ensemble results by averaging or taking the median of

the single model probability of suitability values, respectively, whereas weighted mean weights

the individual models based on their cross-validation performance, in our case the TSS values

(Marmion et al., 2009; Thuiller et al., 2009). For the ensemble model evaluation, TSS, ROC and

Kappa were used to assess model performance. The method used to combine the models that

performed best, should be used to present the projection results. The ensemble model included

111 single models out of 450 individual models.

To evaluate the relative importance of each environmental variable (the variable importance), a

simple Pearson's correlation *r* was calculated between predictions of the full model and a model

with each variable removed (Thuiller *et al.*, 2009). We also generated species response curves,

which show how suitability for Pine Marten changes as a given environmental variable changes.

Commented [EL13]: I think its 111 models as it says in the em.models.kept in the Ensemble model output in R

Habitat suitability projections:

We generated single and weighted mean ensemble model projections in Britain using the 100 m

resolution environmental predictors. Suitability index has been re-scaled 0 to 100, where larger

values indicate higher suitability.

Results

The mapped relative probability of habitat suitability in Europe show markedly low habitat

suitability for Pine Marten at high altitudes in the Alps and the Scandinavian Mountains, in

heavily-modified and farmed landscapes, in dry and arid zones including much of Spain, and

over open water bodies. Higher habitat suitability occurs in areas with woody vegetation cover,

in lowlands, low topographic positions, and notably along Europe's West coast (Figure 4).

The cartographic outputs of relative probability of habitat suitability in Great Britain show high

suitability across substantial areas in the South East, East of England and South West, with

pockets of suitable habitat along the West Coast of Britain. Suitability is greatest in wooded

landscapes and lower in urban and intensively farmed areas. Substantial areas of the Scottish

Highlands and central Wales show relatively low suitability (Figure 5).

In the South East, where the feasibility of a reintroduction is currently being assessed, highest

suitability was in woodland habitats and other areas with tree cover (Figure 6). Notable hotspots

of suitability were in the South Downs National Park, the High Weald AONB, and the western

end of the Kent Downs AONB. Low-lying coastal agricultural areas of low suitability include

Thanet, Romney, Dungeness, and along the North Kent coast, as well as agricultural areas

around Andover. Greater London and other urban centres are unsuitable, however, those along

Commented [RG14]: The results currently don't highlight land cover or any vegetation measure as a key variable - is this a general observation looking at satellite imagery?

285 the south coast such as Southampton, Portsmouth and Eastbourne are closely neighboured by

286 areas of high suitability. Across the region, highest suitability was in woodland habitats and

287 other areas with tree cover.

297 Figure 4: Habitat suitability map indicating Pine Marten suitability across Europe at a resolution of 1km x
298 1km. Suitability index ranged from 0 to 100, where larger values indicate higher suitability.

307 Figure 5: Habitat suitability map indicating Pine Marten suitability across Great Britain at a resolution of
308 100 x 100 m. Suitability index ranged from 0 to 100, where larger values indicate higher suitability .

Figure 6: Habitat suitability map indicating Pine Marten suitability across the South East of England, at a

resolution of 100 x 100 m. Suitability index ranged from 0 to 100, where larger values indicate higher

suitability

TSS and ROC validation scores were used to assess the performance of the models. The mean

TSS value of the single models was 0.3046 (Min: 0.218, Q1: 0.286, Median: 0.307, Q3: 0.324,

Max: 0.368). The mean ROC value of the single models was 0.7087 (Min: 0.628, Q1: 0.698,

Median: 0.713, Q3: 0.723, Max: 0.756). The calibration scores for the mean ensemble model

were TSS = 0.425 and ROC = 0.786. The calibration scores for the median ensemble model

were TSS = 0.361 and ROC = 0.751. The calibration scores for the weighted-mean ensemble

model were TSS = 0.426 and ROC = 0.786.

The ensemble model variable importance results, averaged across the ensemble types, showed

elevation was the strongest predictor of suitability, followed by land cover, vegetation greenness

(NDVI) and slope, but to a lesser degree. TPI and canopy height were of relatively low

importance, with forest density being of least importance (Figure 7).

Commented [LB16]: Figure 4 should be the Europe map. Figure 5 should be the UK map.

Figure 7: Box plot showing the ensemble variable importance, across the mean, median and weighted

mean.

The response curves, averaged across the ensemble results (Figure 8), indicate that Pine

Marten favour lower elevations, with a sharp decline in suitability at elevations above 500 m

above sea level (a.s.l.).

Woodland was favoured over other land cover typed, with mixed woodland as the most suitable,

followed by broadleaf woodland and coniferous woodland (Figure 8). Preference for arable and

pasture land cover was lower, with suitability values below 0.5. Urban greenspace,

sclerophyllous vegetation and peat bogs were the least favoured land cover type.

There is a fairly consistent medium probability of occurrence of Pine Marten across all values of

tree density, although at low densities (<25% tree cover) probability starts to decrease, and at

very low densities (<6% tree cover), the probability of occurrence drops sharply from about 0.42

to 0.36 (Figure 8). NDVI showed a weak positive correlation with suitability, such that the

probability of occurrence of Pine Marten increased slightly with increasing vegetation

greenness. Between NDVI values of 0.55 and 0.8 the probability of occurrence exceeds 0.4

(Figure 8). The probability of occurrence of Pine Marten is fairly consistent across values of

canopy height, although above a canopy height of 20 m, there is a slight decrease in the

probability of occurrence, from 0.4 at 25 m, to 0.3 at 35 m (Figure 8).

The species response curve for TPI, indicates a preference of Pine Marten for lower

topographic positions within valleys and depressions (Figure 8). The probability of occurrence of

Pine Marten is fairly consistent across all values of slope, apart from a slight decrease in very

flat areas (Figure 8).

Figure 8: The response curves, averaged from the final ensemble models. Probability of suitability is on a

0-1 scale.

Discussion

In this study, eSDMs were used to map environmental suitability for Pine Marten in Great

Britain, to inform conservation and restoration of the species, with a focus on reintroduction site

suitability. Across the extent of Europe, the results indicate high suitability in the British isles,

northern parts of continental Europe including France, Denmark and Germany and lower

altitudes of Scandinavia. Low suitability regions were often found at lower latitudes including the

Mediterranean regions such as large parts of the Iberian Peninsula and South East Italy, as well

as areas of high altitudes including the Alps, the North Western Highlands across Norway and

Sweden and extents of the Scottish Highlands (Figure 4). Across Britain, the results indicated

high suitability in most regions of Northern and Republic of Ireland and lower altitudes of

Scotland as well as parts of South West of England, North Yorkshire, South Cumbria and North

and South Wales (Figure 5). In the South East, we found areas of high suitability to be

extensively wooded environments in protected areas (National Landscapes and NPs), with

lowest suitability in urban especially low-lying coastal areas (Figure 6). Hotspots of suitability

were in the South Downs National Park, the High Weald AONB, and the western end of the

Kent Downs AONB, which may serve as suitable reintroduction locations. The South East

region holds the second highest density of woodland in the UK with 22.1% forest cover (Forest

Research, 2023), which suggests why a large proportion of the landscape indicates medium to

high suitability. Evaluation scores indicated an overall moderate performance of model

accuracy, likely due to the limitation of mapping at a fine scale with a large spatial extent.

Commented [EL17]: Lawrence - should it say something along the lines of interpretations should be taken with caution?

Commented [LB18]: This needs reshuffling to discuss Europe, GB, then South East.

The variable importance results showed that elevation is the strongest predictors of habitat

suitability for Pine Marten across its global range. Our results show substantially higher

probability of Pine Marten habitat suitability at lower elevations (< 500 m a.s.l). We expect that

elevation acts as a proxy for the complex nuances around the characteristics of woodland

habitats across Europe, with the preference for lower elevations likely being associated with =

their preference for broadleaved woodland habitats which typically occur at lower altitudes

(Harmer et al., 2010; Portoghesi, 2006). Coniferous woodlands and alpine tundra and meadows

generally occur at higher altitudes, but lack the abundance of prey and denning sites required

by Pine Marten (Fonda et al., 2021). Based on our results, which we assume to be free of

sampling bias with altitude, we see that Pine Marten are unlikely to occur in substantial

populations at altitudes above 2000 m a.s.l. In the South East of England, the maximum

elevation is 276m, and thus the altitude of the region is highly suitable for Pine Marten.

Although its relative importance was low, the response curve indicates that Pine Marten prefer

lower topographic positions, representing environments within valleys and depressions. We

expect this preference is linked to the greater presence of woody vegetation in these locations

compared to sparser tree cover that tends to occur in higher topographic positions, on ridges

and valley shoulders. Interestingly, the areas with highest suitability for Pine Marten in the South

East of England were in topographically complex areas (Appendix 1).

Land cover is the second strongest predictor of suitability, with mixed, broadleaf and coniferous

woodland being highly favoured (Figure 8). As forest-dwellers, Pine Marten are found primarily

in well-structured deciduous and coniferous woodland, with well-insulated denning sites and

abundant food resources, and with a complex understorey providing a low-risk of predation

(Buskirk, 1992, Virgo et al., 2012; Delibes,1983; Balestrieri et al., 2015). The results show a

lower preference for coniferous woodland compared to mixed and broadleaf woodland, which

Commented [RG19]: You're saying elevation is accounting for significant variation because it acts as a proxy for a more complex nuance around wooded habitats across pine marten global range. I think spell that out very clearly.

could reflect their preference towards more structurally diverse woodland systems. Planted

coniferous woodland is often thinned and structurally uniform, which serves as a poor habitat for

Pine Marten due to the lack of diverse ground flora and foraging opportunities. Interestingly,

arable, pasture and grassland land cover types were found to be moderately favoured,

contradicting previous models which have shown that they avoid open habitats (Brainerd and

Rolstad, 2002; Storch et al.,1990; Vergara et al., 2015; Wereszczuk and Zalewski, 2015). This

may be explained by the species' ecological flexibility and ability to occupy heterogenous

landscapes. Evidence suggests that diverse habitats can promote higher diversity of food

resources than a continuous forest habitat (Rosalino and Santos-Reis, 2009; Vergara et al.,

2015). For instance, we know that pine marten travel up to150 m from woodland edge to forage

for prey species such as voles, provided there is some low-level vegetation cover (Caryl, 2008).

Linear or small habitat features will be underrepresented in our model due to the analytical

resolution of the modelling (1x1 km). For example, shrub or hedge features within a grassland

or arable pixel will not be represented. These features can serve as important facilitators of

connectivity for pine martens to move through the landscape to access other patches of

woodland, therefore our model likely underpredicts habitat suitability. Notably, urban

greenspace, sclerophyllous vegetation and peat bogs land cover types indicated the lowest

suitability. These land types are rare which may explain why they indicate significantly lower

suitability compared to other land types. Amalgamating the land categories into 17 classes may

have led to increased background noise within the environmental data creating biased habitat

predictions. An improvement to the methodology would be to further reduce the total number of

categories in the model by conducting pairwise correlations between the broad land cover types

to identify highly correlated categories (Buckley and Lundy, 2012).

Commented [LB20]: I don't think I agree, unless im misunderstanding. Reducing the number of categories should reduce noise in the data I think?
How do you test correlation between land cover types?
Do you mean if there are types that commonly neighbour each other then merge them? Like urban and suburban?

NDVI (vegetation greenness) was the third strongest predictor of suitability because Pine

Marten inhabit woodland and adjacent edge habitat. As well as providing denning sites and

shelter, vegetation greenness is also strongly correlated with prey abundance. For example,

rodent density is strongly correlated with net productivity of ground vegetation (Zalewski et al.,

2004). The response curve showed a peak in suitability around 0.55-0.8 NDVI which indicates

high vegetation cover fraction like that of closed-canopy woodland.

Our results also suggested that Pine Marten will occupy a broad range of forest densities, with a

consistent medium probability of occurrence of Pine Marten across all values of tree density

(Figure 8). We expect this links strongly with their ability to adapt to possibly sub-optimum

environments (MacPherson et al., 2014). For example, South Europe populations can be found

across ancient woodland, coppiced woodland and cultivated land with forest fragments, where

they are not restricted to dense woodland cover (Pittiglio, 1996). Even within countries, the

adaptation to varying vegetation structures is evident. For example, French populations have

genetically diverged to occupy varying woodland cover. In the Bressie region, individuals

occupy human-dominated landscapes with fragmented forest whilst in the Ariège region

individuals occupy densely forested habitat (Larroque et al., 2015).

The results show canopy height was a relatively unimportant variable. Suitability increased with

increasing canopy height, up to 20 m, where it then decreased with increasing canopy height

Pine Marten favour taller spruce-dominated forest for protection from predators and access to

den and resting sites within the forest canopy (Brainerd and Rolstad, 2002).

Mean average monthly temperature (°C) and mean average monthly precipitation (mm) were

removed from the eSDM. While these variables have high predictive power at the European

scale, we know that Great Britain has a highly favourable climate for Pine Martens, and thus the

predictive dominance of the climatic variables in the model were unfavourable for mapping

suitability in Great Britain. In previous model versions, mean average monthly temperature (°C)

was the strongest (37%), and mean average monthly precipitation (mm) (17%), the third

strongest predictor of suitability. Model performance evaluation metrics were also higher (mean

single model validation TSS of ~0.43) when these variables were included. Mean average

monthly precipitation (mm) was also collinear with vegetation greenness (NDVI) and elevation.

Sampling bias is a major challenge for species distribution modelling, and we identified

disproportionate sampling effort in our dataset. Specifically, Pine Marten occurrence records

were more common near urbanised areas and roads, possibly linked to accessibility for

recorders (Araújo and Guisan, 2006). This could lead to spatial bias in the model,

misrepresenting the environmental conditions which reflect suitability. We therefore decided to

remove the distance to roads and urban areas variables to prevent the influence of spatial

autocorrelation. One may also argue that roads and urban areas should not be included as they

serve as barriers rather than predictors of suitable habitat. In other words, Pine marten do not

deliberately choose to be close to or far from roads, rather, territory is chosen based on

predominantly the abundance of woodland for denning potential and availability of prey. The

impacts of roads and urban areas on habitat suitability and connectivity are better examined

using connectivity modelling to quantify the intensity of movement barriers, which includes

analysing the risks of mortality and how it effects overall population viability. Whilst

comprehensive efforts have been made to reduce bias in the occurrence data, other causes of

bias may still be present. For instance, we identified low environmental suitability at high

altitudes, which may be due to a lower number of observations in these difficult to reach

locations, as well as low importance of canopy height which may be linked to fewer

observations of Pine Marten high in the canopy.

Commented [ACN21]: Were RTA records stripped out?

Commented [RG22R21]: This is likely to have affected every variable, particularly land cover/woodland variables.

Our models did not perform to a good or excellent standard, which may be due to the large

spatial extent of our analysis (Europe), which can lead to the selection of a higher proportion of

less informative background points (Barbet-Massin et al. 2012). This can lead to difficulties

detecting commission and omission errors (false presences and false absences), leading to low-

moderate model performance. Moreover, climatic predictor variables were removed which had

the strongest predictive power. Due to these imitations, the suitability maps should be used to

guide further ecological feasibility research to assess suitability in the local context.

Raster data with higher resolutions than is typical for this type of study were used for a number

of reasons. Firstly, higher resolution environmental variables provide more accurate information

about the environment, whereas down-sampled environmental variables average across

neighbouring cells, often leading to a poor measure of local conditions. This is particular

pertinent in highly fragmented landscapes, where environmental predictors which change over

small spatial distances may become meaningless if a coarse resolution averages a broad range

of values. Secondly, high-resolution projections provide suitability estimates for smaller land

areas, which is particularly important in highly fragmented landscapes such as in England. At

the 100m scale, the results show substantial connected suitable habitat in the South East,

particularly in the High Weald National Landscape. Moreover, it is expected that Nature

Recovery Networks will increasingly connect fragmented habitat patches in future years.

Our results reveal contiguous areas of suitable habitat in the South East and East of England.

These results must be coupled with further research into connectivity and road mortality to

inform the viability of a reintroduction. Since male Pine Martens hold large territories, ranging

from 2 km^2 to up to 25 km^2 (Zalewski and Jędrzejewski, 2006; Balestrieri et al. 2010), the

requirement for connectivity between highly suitable habitat areas is essential for their long-term

establishment (McNicol et al., 2020) and should therefore be investigated. Through the

connectivity modelling, we can also assess proximity to existing populations, for example in the

New Forest, which is important for ensuring genetic diversity and the long-term viability of the

reintroduced populations. Additionally, MacPherson and Wright's (2021) 'Long-term strategic

recovery plan for Pine Martens in Britain' report recommends further research into the effects of

roads on risk of mortality in more densely-populated parts of the Great Britain, like the South

East region. The South East has the second highest density of woodland (Forest Research,

2023), but also the highest traffic levels in the UK (Department for Transport, 2023). Our

suitability models do not account for roads, and therefore further research into road mortality is

essential prior to restoration activity.

Appendices

<u>Appendix 1:</u>

Habitat suitability map indicating Pine Marten suitability across the Southeast of England overlayed with
DEM Terrain model 30m (Jaxa, 2021), at a resolution of 100 x 100 m. Suitability index ranged from 0 to
100, where larger values indicate higher suitability

<u>Appendix 2:</u>

Habitat suitability map indicating Pine Marten suitability across regions of Scotland, England and Wales at

a resolution of 100 x 100 m. Suitability index ranged from 0 to 100, where larger values indicate higher

suitability.

534   <u>Scotland</u>

544   <u>Wales</u>

556 <u>Southwest England</u>

567 <u>East Anglia</u>

578 <u>South Cumbria</u>

589 <u>North Yorkshire</u>

600   Appendix 3:

601   Reclassification of CORINE land classes and UKCEH land classes to form 17 land classifications. NA

602   indicates the categories which were not available in the UKCEH layer.

UKCEH land class number	Alternative layer for UKCEH land class	CORINE land class number	Merged land class categories (combined CORINE and UKCEH)	Classification code of combined categories
20		1 to 9, 11	Urban	1
NA	OS Open Greenspace (Ordnance Survey, 2022)	10	Urban green	2
3		12, 19 to 22	Arable	3
NA		18	Pastures	4
1		23	Broadleaf	5
2		24	Conifer	6
NA		25	Mixed woodland	7
4 to 7		26	Natural grassland	8
9		27	Moor and heath	9
NA		28	Sclerophyllous vegetation	10
NA		29	Transitional woodland-scrub	11
12		30 to 34	Bare (bare rock/sparse veg/burnt area/glacial/perpetual snow)	12
8		35	Inland marsh	13
11		36	Peat bogs	14

19		37	Salt marsh	15
13, 15, 16		38 to 39, 42 to 44	Coastal	16
14		40 to 41	Freshwater	17

607 Appendix 6:

608 Ensemble model evaluation results from the ensemble model outputs, indicating model performance
609 across TSS, ROC and Kappa evaluations.

```
610     filtered.by     algo metric.eval cutoff sensitivity specificity calibration validation evaluation
611  1       TSS    EMmean        ROC  496.5      76.940      65.643       0.786          NA         NA
612  2       TSS    EMmean        TSS  491.0      78.133      64.327       0.425          NA         NA
613  3       TSS    EMmean      KAPPA  597.0      51.444      84.548       0.323          NA         NA
614  4       TSS  EMmedian        ROC  493.5      70.972      65.294       0.751          NA         NA
615  5       TSS  EMmedian        TSS  491.0      71.545      64.595       0.361          NA         NA
616  6       TSS  EMmedian      KAPPA  607.0      44.235      85.094       0.273          NA         NA
617  7       TSS   EMwmean        ROC  494.5      77.894      64.734       0.786          NA         NA
618  8       TSS   EMwmean        TSS  493.0      78.205      64.241       0.426          NA         NA
619  9       TSS   EMwmean      KAPPA  600.0      51.516      84.558       0.323          NA         NA
```

622 Appendix 7:

623 Ensemble variable importance results from ensemble model outputs across mean, median and weighted
624 average evaluations.

```
625     filtered.by     algo       expl.var rand  var.imp
626  1       TSS   EMmean      elevation    1 0.424409
627  2       TSS   EMmean  canopy_height    1 0.036039
628  3       TSS   EMmean forest_density    1 0.022745
629  4       TSS   EMmean           ndvi    1 0.184354
630  5       TSS   EMmean          slope    1 0.183473
631  6       TSS   EMmean            tpi    1 0.040790
632  7       TSS   EMmean     land_cover    1 0.210616
633  8       TSS   EMmean      elevation    2 0.419510
634  9       TSS   EMmean  canopy_height    2 0.036504
```

```
635  10      TSS   EMmean  forest_density  2 0.022720
636  11      TSS   EMmean           ndvi  2 0.184362
637  12      TSS   EMmean          slope  2 0.184106
638  13      TSS   EMmean            tpi  2 0.040971
639  14      TSS   EMmean     land_cover  2 0.209221
640  15      TSS   EMmean      elevation  3 0.414192
641  16      TSS   EMmean  canopy_height  3 0.036326
642  17      TSS   EMmean  forest_density  3 0.023042
643  18      TSS   EMmean           ndvi  3 0.185699
644  19      TSS   EMmean          slope  3 0.182867
645  20      TSS   EMmean            tpi  3 0.040130
646  21      TSS   EMmean     land_cover  3 0.209279
647  22      TSS EMmedian      elevation  1 0.419994
648  23      TSS EMmedian  canopy_height  1 0.035677
649  24      TSS EMmedian  forest_density  1 0.022718
650  25      TSS EMmedian           ndvi  1 0.189380
651  26      TSS EMmedian          slope  1 0.176616
652  27      TSS EMmedian            tpi  1 0.039820
653  28      TSS EMmedian     land_cover  1 0.208859
654  29      TSS EMmedian      elevation  2 0.419096
655  30      TSS EMmedian  canopy_height  2 0.035526
656  31      TSS EMmedian  forest_density  2 0.022522
657  32      TSS EMmedian           ndvi  2 0.186160
658  33      TSS EMmedian          slope  2 0.177070
659  34      TSS EMmedian            tpi  2 0.039465
660  35      TSS EMmedian     land_cover  2 0.209999
661  36      TSS EMmedian      elevation  3 0.418853
662  37      TSS EMmedian  canopy_height  3 0.035591
663  38      TSS EMmedian  forest_density  3 0.022522
664  39      TSS EMmedian           ndvi  3 0.187979
665  40      TSS EMmedian          slope  3 0.175561
666  41      TSS EMmedian            tpi  3 0.038905
667  42      TSS EMmedian     land_cover  3 0.212062
668  43      TSS   EMwmean      elevation  1 0.422913
669  44      TSS   EMwmean  canopy_height  1 0.036650
670  45      TSS   EMwmean  forest_density  1 0.022307
671  46      TSS   EMwmean           ndvi  1 0.184124
672  47      TSS   EMwmean          slope  1 0.181986
673  48      TSS   EMwmean            tpi  1 0.040894
674  49      TSS   EMwmean     land_cover  1 0.208688
675  50      TSS   EMwmean      elevation  2 0.424065
676  51      TSS   EMwmean  canopy_height  2 0.035704
677  52      TSS   EMwmean  forest_density  2 0.022511
678  53      TSS   EMwmean           ndvi  2 0.185314
679  54      TSS   EMwmean          slope  2 0.183928
680  55      TSS   EMwmean            tpi  2 0.040741
681  56      TSS   EMwmean     land_cover  2 0.207191
682  57      TSS   EMwmean      elevation  3 0.420573
683  58      TSS   EMwmean  canopy_height  3 0.036186
684  59      TSS   EMwmean  forest_density  3 0.022882
685  60      TSS   EMwmean           ndvi  3 0.186369
686  61      TSS   EMwmean          slope  3 0.182078
687  62      TSS   EMwmean            tpi  3 0.040748
688  63      TSS   EMwmean     land_cover  3 0.209171
```

References

1.Langley, P.J.W. & Yalden, D.W. (1977) The decline of the rarer carnivores in Great Britain during the nineteenth century.

Mammal Review, 7, 95-116

2.Twining, J.P., Montgomery, I., Fitzpatrick, V., Marks, N., Scantlebury, D.M. and Tosh, D.G. (2019). Seasonal, geographical,

and habitat effects on the diet of a recovering predator population: the European Pine Marten (Martes martes) in Ireland.

European Journal of Wildlife Research, 65(3).

3.Thuiller, W. (2014). Editorial commentary on 'BIOMOD - optimizing predictions of species distributions and projecting

potential future shifts under global change'. Global Change Biology, 20(12), pp.3591–3592.

4.Raes, N.; Aguirre-Gutiérrez, J. A (2018) Modeling framework to estimate and project species distributions in space and time.

In Mountains, Climate and Biodiversity; Hoorn, C., Perrigo, A., Antonelli, A., Eds.; John Wiley & Sons: Chichester, UK

5.Dormann, C.F., Elith, J., Bacher, S. and Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a

simulation study evaluating their performance. *Ecography,* 36(1), pp.27-46

6.Allouche, O., Tsoar, A. and Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa

and the true skill statistic (TSS). Journal of Applied Ecology, 43(6), pp.1223–1232.

7.Anderson R. P. et al. 2003. Evaluating predictive models of species' distributions: criteria for selecting optimal models. *Ecol.*

Model. 162, pp.211– 232.

8.Brown, J.L. and Yoder, A.D. (2015). Shifting ranges and conservation challenges for lemurs in the face of climate change.

Ecology and Evolution, 5(6), pp.1131–1142.

9.Forest Research. (2022). Woodland Statistics - Forest Research. Available at: https://www.forestresearch.gov.uk/tools-and-

resources/statistics/statistics-by-

topic/woodlandstatistics/#:~:text=The%20area%20of%20woodland%20in,and%209%25%20in%20Northern%20Ireland.

[Accessed 31 Jan. 2023].

10.Araújo, M. & New, M. (2007) Ensemble forecasting of species distributions. Trends in Ecology & Evolution. 22 (1), 42–47.

11.Elith, J and Leathwick, J.R (2009) Species Distribution Models Ecological Explanation and Prediction Across Space and

Time. Annual Review of Ecology, Evolution, and Systematics 40(1),pp. 677-697

12.Kwon, H. S. (2014) 47. Species Distribution Modeling of Endangered Mammals for Ecosystem Services Valuation -

Focused on National Ecosystem Survey Data**.** Journal of the Korean Society of Environmental Restoration Technology 17(1)

pp. 111-122

13.Araújo, M. B., and M. New. (2007). Ensemble forecasting of species distributions. Trends in Ecology & Evolution 22:42–47.

14. Yu B, Dai W, Li S, Wu Z, Wang J. (2024) A new threshold selection method for species distribution models with presence-

only data: Extracting the mutation point of the P/E curve by threshold regression. Ecological Evolution. 14(4):e11208.

15.Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J.J. and Elith, J. (2021). Predictive performance of presence-only species

distribution models: a benchmark study with reproducible code. Ecological Monographs, 92(1).

16.Hughes-Oliver, J. M. (2018). Population and empirical PR curves for assessment of ranking algorithms.

ArXiv abs/1810.08635

17.Barbet-Massin, M., Jiguet, F., Albert, C.H. and Thuiller, W. (2012). Selecting pseudo-absences for species distribution
models: how, where and how many? Methods in Ecology and Evolution, 3(2), pp.327–338.

18. Rosalino, Luís and Santos-Reis, Margarida. (2009). Fruit consumption by carnivores in Mediterranean Europe. Mammal
Review. 39. 67 - 78.

19. Glon, H.E., Heumann, B.W., Carter, J.R., Bartek, J.M. and Monfils, A.K. (2017). The contribution of small collections to
species distribution modelling: A case study from Fuireneae (Cyperaceae). Ecological Informatics, 42, pp.67–78.

20.Austin, M. (2007). Species distribution models and ecological theory: A critical assessment and some possible new
approaches. Ecological Modelling, 200(1-2), pp.1–19.

21.Thuiller, W. et al. (2009) 'BIOMOD - A platform for ensemble forecasting of species distributions', Ecography, 32(3), pp.
369–373.

22. Liu, C., Berry, P.M., Dawson, T.P. and Pearson, R.G. (2005). Selecting thresholds of occurrence in the prediction of
species distributions. Ecography, 28(3), pp.385–393.

23.Cohen, J. (1960) A coefficient of agreement of nominal scales. Education and Psychological Measurement, 20, pp.37-46.

24.Thuiller, W., Lafourcade, B., Engler, R. & Araújo, M.B. (2009) BIOMOD – a platform for ensemble forecasting of species
distributions. Ecography, 32, 369–373.

25. Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., and Anderson, R. P. (2015). spThin: an R package for
spatial thinning of species occurrence records for use in ecological niche models. Ecography 38, 541–545.

26. Kadmon, R, Farber, O, and Danin, A (2004). Effect of roadside bias on the accuracy of predictive maps produced by
bioclimatic models. – Ecological Applications. 14, pp. 401–413.

27.Anderson, R. P. and Gonzalez, I. J. (2011). Species-specific tuning increases robustness to sampling bias in models of
species distributions: an implementation with Maxent. – Ecological Modelling. 222, pp-2796–2811.

28. Duque-Lazo, J., van Gils, H., Groen, T.A. and Navarro-Cerrillo, R.M. (2016). Transferability of species distribution models:
The case of Phytophthora cinnamomi in Southwest Spain and Southwest Australia. Ecological Modelling, 320, pp.62–70.

29. R Core Development Team. R: A Language and Environment for Statistical Computing R Foundation for Statistical
Computing, Vienna, Austria (2014)

30. Bio, A.M.F. Becker, P.T, Bie, E.D., and Wassen, M.J. (2002). Prediction of plant species distribution in lowland river valleys
in Belgium: Modelling species response. Biodiversity and Conservation, 11(12), pp.2189-2216

31.Sheehy, E., Sutherland, C., O'Reilly, C. and Lambin, X. (2018) The enemy of my enemy is my friend: native Pine Marten

recovery reverses the decline of the red squirrel by suppressing grey squirrel populations. Proceedings of the Royal Society B:

Biological Sciences, 285, 20172603.

32.Schaumann, F. and Heinken, T. (2002). Endozoochorous seed dispersal by martens (Martes foina, M. martes) in two

woodland habitats. Flora - Morphology, Distribution, Functional Ecology of Plants, 197(5), pp.370–378.

33. Fick, S.E. and R.J. Hijmans, 2017. WorldClim 2: new 1km spatial resolution climate surfaces for global land areas.

International Journal of Climatology 37 (12), 4302-4315.

34. Guisan A, Lehmann A, Ferrier S, Austin M, Overton JMC, Aspinall R, Hastie T. (2006) Making better bio-geographical

predictions of species' distributions. Journal of Applied Ecology. 43, 386–392.

35.Meller L, Cabeza M, Pironon S, Barbet-Massin M, Maiorano L, Georges D, Thuiller W. (2014) Ensemble distribution models

in conservation prioritization: from consensus predictions to consensus reserve networks. Diversity Distribution. 20(3):309-321.

36. Beck, J., Böller, M., Erhardt, A., & Schwanghart, W. (2014). Spatial bias in the GBIF database and its effect on

modeling species' geo-graphic distributions. Ecological Informatics, 19, 10–15

37.Aarts, G., Fieberg, J., Brasseur, S. and Matthiopoulos, J. (2013). Quantifying the effect of habitat availability on species

distributions. Journal of Animal Ecology, 82(6), pp.1135–1145.

38. Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R.K., Thuiller, W., 2009. Evaluation of consensus methods in predictive

species distribution modelling. Divers. Distrib. 15 (1), 59–69

39. S.J. Phillips, R.P. Anderson, R.E. (2006) Schapire Maximum entropy modelling of species geographic distributions

Ecological Modelling, 190 (3–4), pp. 231-259,

40. Guisan, A., Thuiller, W., and Zimmermann, N. (2017). The Biomod2 Modeling Package Examples. In Habitat Suitability and

Distribution Models: With Applications in R (Ecology, Biodiversity and Conservation, pp. 357-400). Cambridge: Cambridge

University Press.

41.Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn,

R., Svantesson, S., Wengström, N., Zizka, V. and Antonelli, A. (2019). CoordinateCleaner : Standardized cleaning of

occurrence records from biological collection databases. Methods in Ecology and Evolution, 10(5), pp.744–751.

42. Ball and Tzanopoulos (2020) Interplay between topography, fog and vegetation in the central South Arabian mountains

revealed using a novel Landsat fog detection technique, Remote Sensing in Ecology and Conservation 6(4): 498-513.

43.Paine, R.T. (1969) A note on trophic complexity and community stability. The American Naturalist, 103:91–93.

44.Hastings, A., Byers J.E, Crooks J.A, Cuddington K, Jones C.G, Lambrinos J.G, Talley T.S, Wilson W.G. (2007) Ecosystem
engineering in space and time. Ecology Letters 10, pp.153–164.

45. Crowley, S.L, Hinchliffe S, McDonald R.A. (2018) Killing squirrels: Exploring motivations and practices of lethal wildlife
management. Environment and Planning E: Nature and Space 1, pp.120–143.

46. Sandom C, Donlan C.J, Svenning J.C, Hansen D (2013) Rewilding. Key topics in conservation biology. Wiley-Blackwell,
Chichester, UK. pp.430–451

47.Davies H.T, Du Toit J.T. (2004) Anthropogenic factors affecting wild dog Lycaon Pictus reintroductions: A case study in
Zimbabwe. Oryx. 38, pp. 32–39

48. Simard, M., Pinto, N., Fisher, J.B. and Baccini, A. (2011). Mapping forest canopy height globally with spaceborne lidar.
Journal of Geophysical Research, 116(G4).

49. Brus, D.J., Hengeveld, G.M., Walvoort, D.J.J., Goedhart, P.W., Heidema, A.H., Nabuurs, G.J. and Gunia, K. (2011).
Statistical mapping of tree species over Europe. European Journal of Forest Research, 131(1), pp.145–157.

50.Morton, R.D.; Marston, C.G.; O'Neil, A.W.; Rowland, C.S. (2021). Land Cover Map 2020 (25m rasterised land parcels, GB).
NERC EDS Environmental Information Data Centre

51. F. Ben Rais Lasram, F. Guilhaumon, C. Albouy, S. Somot, W. Thuiller, D. Mouillot (2010), The Mediterranean Sea as a
"cul-de-sac" for endemic fishes facing climate change Global Change Biol., 16. pp. 3233-3245,

52.W. Thuiller, D. Georges, R. Engler, F. Breiner biomod2: ensemble platform for species distribution modeling R package
version, 3 (2016), pp. 3-7

53. Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R.O., Wang, Z. and Zimmermann, N.E. (2019). Model complexity
affects species distribution projections under climate change. Journal of Biogeography, 47(1), pp.130–142.

54. Duan, R.Y., Kong, X.Q., Huang, M.Y., Fan, W.Y. and Wang, Z.G. (2014). The Predictive Performance and Stability of Six
Species Distribution Models. PLoS ONE, 9(11), p.e112764.

55.A.T. Peterson, J. Soberón, R.P. Anderson, R.G. Pearson, E. Martínez-Meyer, M. Nakamura, M.B. Araújo Ecological Niches
and Geographic Distributions: A Modeling Perspective Princeton University Press, Princeton (2012) 328 pp.

56.Hao, T., Elith, J, Lahoz-Monfort, J.J.,Guillera-Arroita, G. (2020) Testing whether ensemble modelling is advantageous for
maximising predictive performance of species distribution models. Ecography 2020, 43, pp.549–558.

57. Buskirk, S.W (1992) Conserving circumboreal forests for martens and fishers. Conservation Biology 6, pp.318–320

58.Delibes, M (1983) Interspecific competition and the habitat of the stone marten Martes foina (Erxleben 1777). Environmental
Science. 174, pp. 229–231

59. Pittiglio, C (1996). Differential use of food and habitat by sympatric carnivores in the eastern Italian Alps. Journal of
Zoology, 75(2), pp. 173–184

60. Storch I, Lindström E.R, Jounge,J (1990) Diet and habitat selection of the Pine Marten in relation to competition with the
red fox. Acta Theriologica 35, pp.311–320

61. Balestrieri A, Remonti L, Ruiz-González A, Gómez-Moliner B.J, Vergara M, Prigioni, C (2010) Range. expansion of the
Pine Marten (Martes martes) in an agricultural landscape matrix (NW Italy). Mammal Biology 75, pp. 412–419

62. Zalewski, A., Jedrzejewski, W. and Jedrzejewski, B. (2004). Mobility and home range use by Pine Martens (Martes martes)
in a Polish primeval forest. Écoscience, 11(1), pp.113–122.

63. Jedrzejewski, W., A. Zalewski and B. Jedrzejewska, (1993) Foraging by pine marten Martes martesin relation to
foodresources in Bialowieza National Park, Poland. ActaTheriologica, 38, pp.405-426

64. Harlow, H.J. (1994). Trade-offs associated with the size and shape of American martens. Pages 391-403 inS.
W.Buskirk, A. S. Harestad, M. G. Raphael & R. A. Powell(eds.). Martens, Sables, and Fishers: Biology and
Conservation. Cornell University Press, Ithaca, New York.

65. Vergara M, Cushman S.A, Urra F, Ruiz-González A. (2015) Shaken but not stirred: multiscale habitat suitability modeling of
sympatric marten species (Martes martes and Martes foina) in the northern Iberian Peninsula. Landscape Ecology. 31,
pp.1241–1260

66. Wereszczuk, A, Zalewski, A. (2015) Spatial Niche Segregation of Sympatric Stone Marten and Pine Marten– Avoidance of
Competition or Selection of Optimal Habitat? PLOS ONE. 10(10)

67. Virgos E, Zalewski A, Rosalino L, Mergey M (2012) Habitat ecology of genus Martes in Europe: a review of the evidences.
In: Aubry KB, Zielinski WJ, Proulx G, Buskirk SW (eds) Biology and conservation of martens, sables, and fishers: a new
synthesis. Cornell University Press, New York, pp. 255–266

68. Balestrieri A, Remonti L, Ruiz-Gonza´lez A, Zenato M, Gazzola A, Vergara M, Detorri E.E, Saino N, Capelli E,
Go´mezMoliner BJ, Guidali F, Prigioni C (2015) Distribution and habitat use by Pine Marten Martes martes in a riparian corridor
crossing intensively cultivated lowlands. Ecological Research, 29(6), pp.1-10

69. JNCC (2021) UK Biodiversity Indicators 2021. JNCC.

70. Mammola, S., Pétillon, J., Hacala, A., Monsimet, J., Marti, S., Cardoso, P. and Lafage, D. (2021). Challenges and
opportunities of species distribution modelling of terrestrial arthropod predators. Diversity and Distributions, 27(12), pp.2596–
2614.

71. European Environment Agency (2018a) CORINE land cover 2018--copernicus land monitoring services
https://land.copernicus.eu/pan-european/corine-land-cover/clc2018

72. European Environment Agency (2018b) Tree Cover Density 2018--copernicus land monitoring services
https://land.copernicus.eu/pan-european/high-resolution-layers/forests/tree-cover-density/status-maps/tree-cover-density-2018

73. QGIS Development Team, 2021. QGIS Geographic Information System. Open Source Geospatial Foundation Project.

74.Li X, Wang Y. (2012) Applying various algorithms for species distribution modelling. Integral Zoology. 8, pp.124–135

75. Zalewski, A. and Jędrzejewski, W. (2006) Spatial organisation and dynamics of the Pine Marten Martes martes population
in Białowieza Forest (E Poland) compared with other European woodlands. Ecography, 29, pp. 31-43.

76. McNicol, C.M., Bavin, D., Bearhop, S., Bridges, J., Croose, E., Gill, R., Goodwin, C.E.D., Lewis, J., MacPherson, J.,
Padfield, D., Schofield, H., Silk, M.J., Tomlinson, A.J. and McDonald, R.A. (2020). Postrelease movement and habitat selection
of translocated Pine Martens Martes martes. Ecology and Evolution, 10(11), pp.5106–5118.

77.Caryl, F.M. (2008). Pine Marten diet and habitat use within a managed coniferous forest. Ph. D. Thesis. University of Stirling
Available at:
https://dspace.stir.ac.uk/bitstream/1893/2261/1/Fiona%20Mae%20Caryl.%20Pine%20Marten%20Diet%20%26%20Habitat%20
Use%20Within%20A%20Managed%20Coniferous%20Forest.pdf (Accessed 7th July 2023)

78. Forest Research (2023) The canopy cover Webmap of the United Kingdom's towns and cities. Available at:
https://www.forestresearch.gov.uk/publications/the-canopy-cover-webmap-of-the-united-kingdoms-towns-and-cities/ Accessed
23 Mar 2024]

79. Department for Transport (2023) Road Traffic Statistics. Available at: https://roadtraffic.dft.gov.uk/regions Accessed 23 Mar
2024]

80. Buckley, D.J. and Lundy, M. (2012). The current distribution and potential for future range expansion of feral ferret Mustela
putorius furo in Ireland. European Journal of Wildlife Research, 59(3), pp.323–330

81. Kirk T.A and Zielinski W.J (2009) Developing and testing a landscape habitat suitability model for the American marten
(Martes americana) in the Cascades mountains of California. Landscape Ecology 24, pp.759–773

82.Carroll C, Zielinski W.J, Noss R.F (1999) Using presence-absence data to build and test spatial habitat models for the fisher
in the Klamath region, USA. Conservation Biology, 13, pp.1344–1359

83. O'Mahony, D.T. (2017). A species distribution model for Pine Marten (Martes martes) in the least forested region of Europe.
Mammal Research, 62(2), pp.195–200.

84. McPherson J, Jetz W, Rogers D.J (2004) The effects of species' range sizes on the accuracy of distribution models:
ecological phenomenon or statistical artefact? Journal of Applied Ecology 41, pp.811–823

85. GDAL/OGR contributors (2022). GDAL/OGR Geospatial Data Abstraction software Library. Open Source Geospatial
Foundation. URL: https://gdal.org

86. USGS (2022) *Landsat 8 Collection 1 Tier 1 Landsat 8 Collection 1 Tier 1 Annual Greenest-Pixel TOA Reflectance.*
Available at: https://developers.google.com/earth-
engine/datasets/catalog/LANDSAT_LC08_C01_T1_ANNUAL_GREENEST_TOA [Accessed 28 Mar. 2023]

87. GBIF Secretariat. GBIF Backbone Taxonomy. Checklist dataset. Available at: https://www.gbif.org/dataset/d7dddbf4-2cf0-
4f39-9b2a-bb099caae36c2023-09-01. [Accessed 28 Mar 2023]

88. National Biodiversity Network (NBN) Trust (2023). The National Biodiversity Network (NBN) Atlas. Available at:
https://ror.org/00mcxye41 [Accessed 28 Mar 2023]

89. 75. Seddon, P.J, Armstrong, D.P (2019) The role of translocation in rewilding. In: du Toit JT, Pettorelli N, Durant SM (eds)
Rewilding (pp. 303-324). Cambridge University Press, Cambridge

90. Klein, L. and Arts, K. (2021). Public participation in decision-making on conservation translocations: the importance and
limitations of a legislative framework. Restoration Ecology, 30(1).

91. Kirkwood, J. (2013). Guidelines for reintroductions and other conservation translocations. Animal Welfare, 22(4), pp.489–
490

92. D.P. Armstrong, P.J. and Seddon (2008) Directions in reintroduction biology. Trends in Ecology and Evolution, 23, pp. 20-
25

93. Mathews F, and Harrower C. (2020). IUCN – compliant Red List for Britain's Terrestrial Mammals. Assessment by the
Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage. Natural England,
Peterborough ISBN 978-1-78354-485-1

94. MacPherson, J. (2014) Feasibility assessment for reinforcing Pine Marten numbers in England and Wales. Vincent Wildlife
Trust. Available at: https://www.vwt.org.uk/wp-content/uploads/2015/04/Feasibility-Assessment-for-Reinforcing-Pine-Martens-
in-England-and-Wales.pdf [Accessed 28 Mar 2023]

95. Stringer, A. P., MacPherson, J., Carter, S., Gill, R., Ambrose-Oji, B. and Wilson, R.(2015). The Feasibility of Reintroducing
Pine Martens (Martes martes) to the Forest of Dean and Lower Wye Valley. Herefordshire: Forestry Commission
England, Gloucestershire Wildlife Trust, Vincent Wildlife Trust.

96. MacPherson, J., Croose, E., Bavin, D., O'Mahony, D., Somper, J. and Buttriss, N. (2014). Feasibility assessment for
reinforcing Pine Marten numbers in England and Wales. Ledbury, UK: Vincent Wildlife Trust.

97. MacPherson, J. and Wright, P. (2021) Long-term strategic recovery plan for Pine Martens in Britain. The Vincent Wildlife
Trust, Natural England, NatureScot. pp.1-72.

98. Gilpin, M.E. and Soulé, M.E. (1986) Minimum viable populations: processes of species extinction. Conservation Biology:
the science of scarcity & diversity (eds M.E. Soulé & M.A. Sunderland), pp. 19-34. Sinauer Associates.

99. Birks, J.D. (2020) Pine Martens. Whittet Books Ltd, Essex, UK. Reprint (with updates)

100. Lariviere, S. & Jennings, A.P. (2009). European Pine Marten (Martes martes). Family Mustelidae (Weasels and
Relatives). In: D.E. Wilson & R. A. Mittermeier (Eds), Handbook of the Mammals of the World. Volume 1: Carnivores. Lynx
Edicions, Barcelona, Spain.

101. Brainerd, S.M. and Rolstad, J. (2002). Habitat selection by Eurasian Pine Martens *Martes martes* in managed forests of
southern boreal Scandinavia. Wildlife Biology, 8(4), pp.289–297.

102. Lee, J.-K., Hwang, H.-S., Eum, T.-K., Bae, H.-K. and Rhim, S.-J. (2020). Cascade effects of slope gradient on ground
vegetation and small-rodent populations in a forest ecosystem. Animal Biology, 70(2), pp.203–213.

103. Krebs, C.J. (2009) *Ecology: the Experimental Analysis of Distribution and Abundance*. 6th Edition. Pearson Benjamin
Cummings, San Francisco, CA, USA.

104. Báez, S., Fadrique, B., Feeley, K. and Homeier, J. (2022). Changes in tree functional composition across topographic
gradients and through time in a tropical montane forest. PLOS ONE, 17(4), p.e0263508.

105. Sultaire, S.M., Millspaugh, J.J., Jackson, P.J. and Montgomery, R.A. (2022). The influence of fine-scale topography on
detection of a mammal assemblage at camera traps in a mountainous landscape. *Wildlife Biology*, 2023(2).

106. Balestrieri, A., Bogliani, G., Boano, G., Ruiz-González, A., Saino, N., Costa, S. and Milanesi, P. (2016). Modelling the
Distribution of Forest-Dependent Species in Human-Dominated Landscapes: Patterns for the Pine Marten in Intensively
Cultivated Lowlands. *PLOS ONE*, 11(7), p.e0158203.

107. DEFRA (2022). Keepers of time: ancient and native woodland and trees policy in England Government's statement on
England's ancient and native woodland and ancient and veteran trees. Available at:
https://www.gov.uk/government/publications/keepers-of-time-ancient-and-native-woodland-and-trees-policy-in-
england/keepers-of-time-ancient-and-native-woodland-and-trees-policy-in-england [Accessed 28 Mar 2023]

108. Ferrier, S. and Guisan, A. (2006) 'Spatial modelling of biodiversity at the community level', *Journal of Applied Ecology*,
43(3), pp. 393–404.

109. Japan Aerospace Exploration Agency (JAXA) (2021). ALOS World 3D 30 meter DEM. V3.2, Jan 2021. Distributed by
OpenTopography. https://doi.org/10.5069/G94M92HB. [Accessed 23 Mar 2023]

110. R Development Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical
Computing.

111. Center for International Earth Science Information Network - CIESIN (2013)- Columbia University, and Information
Technology Outreach Services - ITOS - University of Georgia. 2013. Global Roads Open Access Data Set, Version 1
(gROADSv1). Palisades, New York: NASA Socioeconomic Data and Applications Center (SEDAC). Available at:
https://doi.org/10.7927/H4VD6WCT. [Accessed 23 Mar 2023]

112. Patterson, Tom. Kelso, Nathaniel Vaughn. (2012). World Urban Areas, 1:10 million [Shapefile]. North American
Cartographic Information Society. Available at: https://maps.princeton.edu/catalog/stanford-xg070wh7159. [Accessed 23 Mar
2023]

113.Araújo, M.B. and Guisan, A. (2006). Five (or so) challenges for species distribution modelling. Journal of Biogeography,
33(10), pp.1677–1688.

114. Stringer A.P, MacPherson, J , Carter, S , Gill, R , Ambrose-Oji, B , Wilson, R , Kelsall, P, Feirn, W.G, Galbraith, L.C,
Hilário CM, Parry, G and Taylor, A (2018) The feasibility of reintroducing pine martens (Martes martes) to the Forest of Dean
and lower Wye Valley. Gloustershire Wildlife Trust Report

115. Fonda, F., Chiatante, G., Meriggi, A.. (2021) Spatial distribution of the pine marten (*Martes martes*) and stone marten
(*Martes foina*) in the Italian Alps. Mammal Biology 101, pp.345–356

116. Croitor, R and Brugal J.P, (2010) Ecological and evolutionary dynamics of the carnivore community in Europe during the
last 3 million years. Quaternary International, 212, pp.98–108.

117. Harmer, R., Kerr, G. and Thompson, R. (2010) Managing Native Broadleaved Woodland. The Stationary Office,
Edinburgh

118. Portoghesi L, (2006). European Forest Types. Categories and types for sustainable forest management reporting and
policy. Forest 3 (4): 462-463

119. Allaby,, M.(2006) Biomes of the Earth: Temperate Forests. New York, NY: Chelsea House

120. Ordnance Survey (2022) *OS Green Open Greenspace.* Available at: https://www.ordnancesurvey.co.uk/products/os-
open-greenspace

121.Vincent Wildlife Trust (2020) Red squirrels making comeback as return of native pine marten spells bad news for invasive
grey squirrel. Available at: https://www.vincentwildlife.ie/news/red-squirrels-making-comeback-as-return-of-native-pine-marten-
spells-bad-news-for-invasive-grey-squirrel

122. Larroque, J., Ruette, S., Vandel, J.-M. and Devillard, S. (2015) *Landscape Ecology*, 31(3), pp. 517–531. doi:
10.1007/s10980-015-0281-6.