Cucumber extracellular vesicles-localized tetraspanin and phloem protein 2 mediates the

cross-kingdom trafficking of a pathogenic non-coding RNA

Neha Devi[1], Vasudha Sharma[1], Nisha Devi[1], Ravi Gupta[2]*, Sunny Dhir[1]*, Yashika Walia[1]*

[1]Molecular Plant Virology Group, R&D Lab, Department of Bio-Sciences & Technology, Maharishi

Markandeshwar (Deemed to be University), Mullana, Ambala-133207 HR India

[2]College of General Education, Kookmin University, Seoul, South Korea

Neha Devi: nehadevi611999@gmail.com

Vasudha Sharma: vasu.3096sharma@gmail.com

Nisha Devi: nishapahal99@gmail.com

*Corresponding Authors

Dr. Yashika Walia; yashikawalia@gmail.com, yashika.dhir@mmumullana.org

ORCID ID: 0000-0002-0796-0050

Dr. Sunny Dhir; sunny72155@gmail.com, sunny.dhir@mmumullana.org

ORCID ID: 0000-0002-3790-8333

Dr. Ravi Gupta; ravigupta@kookmin.ac.kr

ORCID ID: 0000-0001-5242-9528

Abstract

The plant apoplast and Extracellular Vesicles (EVs) harbor various RNA types, including small RNAs, long non-coding RNAs, and circular RNAs that traverse kingdom boundaries via encapsulation within EVs. Viroid RNAs, which are pathogenic circular non-coding RNAs, move across kingdoms, from plants to insects and fungi. However, the molecular mechanism underlying this cross-kingdom trafficking remains largely elusive. Using apple scar skin viroid (ASSVd) as a model for circular non-coding RNAs and cucumber as its host, we show that viroid RNAs, in their circular, linear, and viroid-derived small RNA forms, are present in cucumber EVs. Additionally, we revealed that ASSVd RNA exhibits extracellular RNA signatures, such as m6A methylation and five EXOmotifs, aiding their packaging into EVs. Our characterization of the cucumber EV proteome validated the role of tetraspanin 8 (TET8) as a conserved marker protein in cucumber EVs. Additionally, we report PP2-a phloem lectin with RNA binding domain as a novel protein associated with cucumber EVs. Through RNA-Co-immunoprecipitation and immunoenrichment assays, we show that PP2 and ASSVd are present in TET8-specific exosomes, with direct interactions occurring between PP2-TET8 and PP2-ASSVd. We propose that PP2 may act as an RNA sorting protein, aiding in the RNA loading into EVs. The potential for cross-kingdom viroid RNA transmission via EVs was also investigated and confirmed. In conclusion, our findings suggest that viroid RNAs could serve as a model RNA system for exploring the complexities of cross-kingdom RNA trafficking in plants.

Keywords: Extracellular vesicles, Viroid RNA, Phloem Protein 2, Tetraspanin 8, RNA

Introduction

The role of the apoplast and EVs in mediating cross-kingdom RNA trafficking has gained attention recently (Zhang et al., 2016, Wang et al., 2024). EVs are small, membrane-bound vesicles secreted by cells that can encapsulate a variety of biomolecules, including RNA (Zhang et al., 2016; He et al., 2021, Karimi et al., 2022; Wang et al., 2024). Majorly three classes of EVs have been identified in plants (Ambrosonethe et al., 2023). The first class is represented by exosomes with sizes ranging from 30-150nm which are specifically enriched in TET8, an ortholog of mammalian CD63, CD81, and CD151 (Chen et al., 2022, Jankovičová et al., 2020). The second class is represented by micro-vesicles with sizes ranging from 150-300nm and are associated with syntaxins proteins, of which Penetration 1 (PEN1) is well characterized (Rutter et al., 2017). The third class is represented by EXPO vesicles which are double membrane bodies fused with the plasma membrane (PM), releasing a single membrane vesicle into the cell wall (Wang et al., 2010). Exo70E2 is essential for exocyst subunit recruitment and EXPO formation in both plants and animals (Ding et al., 2014).

The process of cross-kingdom RNA transfer via EVs was first observed during the interaction between Arabidopsis and *Botrytis cinerea*, where pathogen-derived small RNAs (sRNAs) were shown to cross kingdoms and interact with plant argonaute (AGO) to target mitogen-activated protein kinase transcripts, promoting pathogen colonization (Weiberg et al., 2013). Plant apoplast and EVs are enriched in various RNA species, such as long non-coding RNAs (lncRNA), circular RNAs, (circRNA) micro RNAs (miRNAs), and tasiRNAs (Karimi et al., 2022). Subsequent studies revealed that plants can send their small RNAs in TET8-specific exosomes to suppress fungal virulence genes (Cai et al., 2018). In addition, they carry mRNAs that were translated into fungi to reduce their pathogenicity (Wang et al., 2024). Recent studies have identified various mechanisms involved in the loading of RNA into EVs. Specifically, RNA-binding proteins such as AGO1, annexins, and RNA helicases play significant roles in the incorporation of plant small RNAs into TET8-specific exosomes (He et al., 2021; Karimi et al., 2022). Furthermore, glycine-rich binding protein (GRP7) and AGO2 have been linked to the secretion of N6-methyladenosine (m6A) modified circular and long non-coding RNAs, as well as small RNAs in the apoplast (Karimi et al., 2022). Similar to findings in animals, circular RNAs in the apoplast are notably enriched with m6A modifications (Sabrina Garbo et al., 2024;

Karimi et al., 2022). Certain factors have been identified in animals and plants that could assist the sorting of RNAs inside the vesicles. These factors encompass characteristics such as a high GC content, the presence of an EXOmotif (a motif known to facilitate preferential sorting of RNA into exosomes or EVs), and their ability to modify RNAs and interact with RNA binding proteins (O'Grady et al., 2022, Ruben Garcia-Martin et al., 2022, Sabrina Garbo et al., 2024, Xiao-Man Liu 2025).

A class of non-coding circular RNAs is represented by viroids which are RNA-only pathogens infecting plants with distinct structural and functional properties (Flores et al., 2016; Navarro et al., 2021; Hammond, 2017). Despite their small size (246-401nts) and non-coding properties, they autonomously replicate within infected plant cells and cause a variety of plant diseases. The ability of viroids to infect plants is enabled by their secondary structure (Branch et al., 1984; Daros et al., 1994; Warrilow et al., 1999; Navarro et al., 2000; Rodio et al., 2007), which is crucial for their replication and movement and allows hijacking of plant cellular machinery for their propagation (Schnölzer et al., 1985; Schmitz and Riesner, 1998; Sano et al., 1992; Reanwarakorn & Semancik, 1998). Recent studies have shown that viroids are not limited to infecting plants but can also affect other biological kingdoms, including fungi, insects, and bacteria (Tian et al., 2020; Walia et al., 2015; Leichtfried et al., 2022; Afanasenko et al., 2022). This cross-kingdom infection potential suggests that viroids, much like other RNA entities, can traverse biological barriers and may play a role in complex ecological interactions (Cai et al., 2019, Wang et al., 2024). For example, research on apple scar skin viroid (ASSVd) demonstrated that viroids get transmitted from infected apple trees to fungi, such as *Alternaria alternata*, a common plant pathogen (Tian et al., 2022). We have previously reported that the glass-house whitefly, *Trialeurodes vaporariorum*, mediates the transmission of ASSVd and cucumber PP2 assists in its acquisition and transmission (Walia et al., 2015). Moreover, certain aphid species and codling moth larvae, which feed directly on infected apple trees, were found to transmit apple chlorotic fruit spot viroid (ACFSVd), highlighting the role of insect vectors in cross-kingdom viroid transmission (Leichtfried et al., 2022). Potato spindle tuber viroid (PSTVd) transmission from infected plants to *Phytophthora infestans* also emphasizes the inter-kingdom dynamics of viroid spread (Afanasenko et al., 2022).

All these findings together suggest that viroids, as circular non-coding RNAs, may take
advantage of RNA sorting and encapsulation mechanisms within EVs to facilitate their
intercellular movement and potential cross-kingdom dissemination. By associating with EVs,
viroids could bypass the hostile intracellular environment of the host, enabling long-distance
transport or transmission across different organisms. This study uses ASSVd as a model RNA
pathogen to explore its presence in the apoplast and EVs of its host. We provide evidence that
PP2 acts as an RNA sorting protein, facilitating viroid RNA loading into TET8 exosomes, and
demonstrate that whiteflies can transmit viroids via EVs.

Results

ASSVd RNA localizes in the apoplast and EVs of cucumber plants

To investigate the presence of viroid RNA in apoplast and EVs, we used cucumber plants
infected with dimeric infectious transcripts of ASSVd (Walia et al., 2014). The apoplastic wash
fluid (AWF) was isolated from viroid-infected plants, 3-week post ASSVd inoculation, using an
infiltration centrifugation method (Fig. S1; O'Leary et al., 2014). SDS-PAGE analysis of the
isolated apoplastic proteins showed the absence of the large subunit of RuBisCO (approximately
55 kDa), indicating that the AWF was free from cytosolic contamination (Fig. S2). Trypan blue
staining of the infiltrated leaves further revealed absence of any cell damage and potential
cytosolic leakage (Fig.S3). Using size exclusion chromatography (SEC), the isolated AWF was
fractionated into 23 fractions (Fig. S4). Of these, fractions 1-7 contained the void volume,
fractions 8-11 contained EVs, and fractions 12-23 contained larger protein complexes. Fractions
8-11, corresponding to EVs, were pooled together and examined using Transmission Electron
Microscopy (TEM) which confirmed the presence of vesicle-like structures at 50nm, 200nm, and
500nm size corresponding to the size of exosomes and micro-vesicular bodies in plants (Fig. 1a;
Rutter et al., 2017). The pooled EV fractions were also subjected to Dynamic Light Scattering
(DLS) for particle size measurement, which revealed their size ranging 50 -1000 nm (Fig. 1b).
Three peaks corresponding to the size 15nm, 100nm, and 500nm were obtained. The percentage
distribution for the 500nm size EVs was maximum compared to 100nm size EVs.

To examine the presence of ASSVd RNA in the apoplast and EVs, total RNA was isolated from
both AWF and pooled EV fractions. Given that RNA can be present either inside EVs or bound
to their surface as protein-associated RNA complexes (Karimi et al., 2022), the EVs were further

split in two fractions to pinpoint the localization of ASSVd RNA within these vesicles (Fig. 1c).

First, the EV fraction was treated with trypsin to release the protein associated RNA and the

RNA was isolated from the outer surface of the EVs without breaking the vesicles which

represented the RNA associated with the EV surface or EV outside RNA (EV^O). The remaining

EV fraction was treated with trypsin to remove proteins, followed by RNase treatment to degrade

protein-bound RNA from the EV surface. The EVs were then ruptured by treatment with

detergent and the inside RNA was isolated which represented the RNA inside EVs (EV^I). The

RT-PCR results indicated that ASSVd RNA is present in the apoplast, adhered to the surface of

EVs (as EV surface RNA), and is also found inside the EVs (Fig. 1d and 1e). However, the

intensity of the ASSVd band was weaker in the inside EV fraction compared to AWF and

outside EV fractions, suggesting that the mature form of ASSVd RNA is present at lower

concentrations inside the vesicles. To further validate these findings, a denaturing Northern blot

analysis of total RNA from AWF and inside and outside EV fractions was carried out. The

analysis revealed the presence of different forms of ASSVd RNA (including circular, linear, and

small RNAs) in both the apoplast and EV fractions. Ethidium bromide staining of the denaturing

PAGE gel with total RNA from AWF, EV^O, and EV^I fractions showed an enrichment of small

RNAs inside EVs compared to the other fractions, consistent with earlier findings in Arabidopsis

(Fig. 1f) (Karimi et al., 2022). Furthermore, the Northern blot results confirmed the presence of

both circular and linear forms of ASSVd RNA in the apoplast and in both the outer and inner EV

fractions, implicating that ASSVd RNA is indeed associated with EVs in these plants (Fig. 1g

and 1h). Small RNA species corresponding to ASSVd were detected only within the EVs,

suggesting that ASSVd-derived small RNAs (sRNAs) are packaged into these vesicles. The lack

of probe hybridization with the RNA from the mock leaf confirmed that the bands observed in

the Northern blot were specific to ASSVd and not from any other plant-derived RNAs.

Figure 1: Isolation of EVs from cucumber apoplast and detection of ASSVd.

(a)TEM images illustrating the morphology of EVs isolated from cucumber plants infected with ASSVd
using the SEC method. EVs are observed to range in size from 50 nm to 500 nm. (b) DLS analysis of the
isolated EVs showing particle sizes between 10 nm and 1000 nm, consistent with the sizes observed in
TEM. (c) Graphical method illustrating the protocol followed for isolation of RNA from outside surface
of EVs and from EVs inside. (d and e) RT-PCR detection of ASSVd in the AWF, total EV fraction
obtained via SEC (EV), EV^O fraction after trypsin treatment and RNA isolation without EV disruption,
and EV^I fraction after trypsin, RNase and detergent treatment followed by RNA isolation from inside the
EVs. (f and g) Northern blot analysis of the RNA samples described above: left panel – EtBr-stained gel
showing RNA from infected leaves and their corresponding AWF and EV fractions, with mock leaf RNA
as a control; right panel – Northern blot of the same gel using a 3'-labeled biotin-negative sense ASSVd
probe, revealing both circular and linear forms of ASSVd in infected leaves, AWF, and inside and
outside EVs. Small RNAs of ASSVd are distinctly visible within the EV fraction. (h) is the
representation of the northern blot in tabulated form.

ASSVd RNA exhibits characteristics of EV loading including m6A modification and EXOmotifs

An immunoprecipitation (IP) experiment using antibodies specific to m6A with equal RNA from

viroid-infected AWF and EVs (Fig. 2a) revealed distinct patterns of m6A modification. In the

apoplastic RNA, m6A-modified RNA was not detected in the pull-down samples and all of it

remained in the flow-through that suggested that ASSVd RNA in the apoplast is mostly

unmodified or is likely modified by modifications other than m6A. In contrast, northern blot

analysis of the EV fractions showed that ASSVd RNA was present in both the pull-down

fractions (containing m6A-modified RNA) and the flow-through (containing unmodified RNA).

However, the intensity of m6A-modified ASSVd in EVs was weaker compared to its unmodified

form. Despite this, since m6A-modified ASSVd was detected in EVs but not in AWF, we

hypothesized that this m6A-modified ASSVd RNA is cell originated. To confirm, we performed

m6A immunoprecipitation from total cell lysates and probed for ASSVd. Our results showed that

m6A modification was present at very low levels within the cell, indicating that some portion of

ASSVd RNA is modified with m6A inside the cell (Fig. 2b). Additionally a sequence-specific

screening revealed an enrichment of five EXOmotifs in the ASSVd genome viz. UCCC, CUCC,

CCCU, UGUG, and GGUG (Fig. 2c). The UCCC and UGUG motif lies in the terminal left

domain, CUCC and CCCU lies in the central conserved region whereas GGUG lies in the

pathogenicity domain of ASSVd (Fig. 2d).

Figure 2: **Enrichment of m6A modification and EXOmotifs in the ASSVd genome.**

(a) Immunoprecipitation of m6A-enriched RNAs from the ASSVd-infected AWF and EVs fractions using an m6A antibody (upper panel), followed by detection of ASSVd in the immunoprecipitated samples using a 3'-labeled biotin-negative sense ASSVd probe (middle panel). The middle panel shows that ASSVd RNA in EVs is modified with m6A, while the lower panel demonstrates that unmodified ASSVd RNA, which did not bind to the m6A antibody, remains in the flow-through fraction. Most of the RNA in AWF is unmodified and remains in flow through fraction, and a subset of ASSVd RNA in the EVs fraction is modified with m6A. (b) Immunoprecipitation of m6A-enriched RNAs from the ASSVd-infected leaf cell extract showing very insignificant modification of ASSVd RNA in cell extract. (c) Overview of various EXOmotifs identified in animal studies from extracellular RNA and the motifs found within the ASSVd genome. d. The location of the identified EXOmotifs in the secondary structure of ASSVd is highlighted by colored boxes.

Proteome analysis of the cucumber EVs infected with viroid

To characterize the proteome of EVs upon viroid stress, AWF and EVs were isolated from ASSVd-infected cucumber plants and mock plants three weeks post viroid infection (Fig. S5, S6). A total of 2454 proteins were identified from healthy cucumber EVs and 2238 proteins from viroid-infected EVs. Of these, 816 proteins were common to both groups (Fig. 3a) suggesting ~33% of the proteome was conserved among viroid-infected and healthy EVs. The identified proteins included exosome marker proteins such as tetraspanins (Tet16/PLAT/LH2 domain-containing lipoxygenase family protein) in healthy EVs, and lipoxygenase 3 (orthologue of TET8 in cucumber) in infected EVs (Table 1). Additionally, markers for microvesicles, including syntaxins (SYP21/71/31, SNARE-like superfamily protein) and target SNARE coiled-coil domain protein in healthy EVs, and SYP124/21/42/71 and SNARE-like super-family proteins were also found in infected EVs. Proteins associated with EXPO vesicles, such as exocyst complex components (SEC84B, SEC15B, SEC5, EXO70 family protein in healthy EVs, and SEC10, SEC15B in infected EVs) were also identified. These findings align with the TEM data, which confirmed that EVs isolated through SEC had a size of 50-500nm consistent with the size of exosomes, micro-vesicles, and EXPO vesicles in plants. Notably, enrichment of RNA-binding proteins including Argonaute family proteins (AGO2, AGO3, and AGO7), DEAD-box RNA helicases, sorting nexins, and glycine-rich RNA-binding proteins were also observed in both healthy and infected plant EVs. Apart from these RNA-binding proteins, chaperonins such as heat shock proteins (HSP60/70/81/89.1/70, DnaJ) were also identified in the EVs (Table 1). Interestingly, a phloem protein -PP2 was also identified in the cucumber EVs which is known to be involved in viroid transmission to whiteflies (Walia et al., 2015) and contains a dsRNA binding motif (ref). The absence of markers for the trans-Golgi network (syntaxin 61, SYP61), late endosomes (Ara6), Golgi bodies (GmMan149), and plasma membrane (LTI6) confirmed the integrity of the EV proteome (Table 1, File: S1). Gene ontology (GO) enrichment analysis based on cellular components revealed that the proteins of both healthy and infected EVs were associated with the endomembrane system, cell-cell junctions, the apoplast, the plasma membrane, and extracellular space, among others (Fig. 3b and 3c). In the biological process category, the enriched proteins in healthy EVs were primarily linked to the regulation of Ras protein signal transduction and GTPase-mediated signal transduction, while infected EVs proteins were mainly involved in cytoskeleton-dependent intracellular transport and microtubule-

based movement (Fig. 3d and 3e). A deeper analysis of biological processes uncovered a particularly interesting finding: infected EVs were notably enriched in proteins related to non-coding RNA processing/modification and RNA processing/metabolic processes, a category not observed in healthy EVs (Fig. 3d and 3e).

Figure 3: Proteome analysis of ASSVd-infected and healthy EVs using bioinformatic tools. (a) Venn diagram illustrating the shared and unique peptide between ASSVd-infected and healthy EVs. (b and c) Pie chart

showing classification of viroid infected and healthy EV proteome based on sub-cellular component. (d and e)
Bar graph showing the classification of viroid infected and healthy EV proteome based on biological processes
with notable enrichment of proteins related to non-coding RNA processes in viroid infected EVs compared to
healthy EVs.

	Healthy		Infected	
Accession	Protein name	Accession	Protein name	
	Syntaxins related			
AT5G16830	syntaxin of plants 21 (PEP12 VACULAOR TRANSPORT)	AT1G61290	syntaxin of plants 124	
AT3G09740	syntaxin of plants 71	AT5G16830	syntaxin of plants 21	
AT5G05760	syntaxin of plants 31	AT4G02195	syntaxin of plants 42	
AT5G02280	SNARE-like superfamily protein	AT5G02280	SNARE-like superfamily protein	
AT1G15370	SNARE-like superfamily protein	AT2G35190	novel plant snare 11	
AT1G16230	Target SNARE coiled-coil domain	AT1G16230	Target SNARE coiled-coil domain protein	
		AT3G09740	syntaxin of plants 71	
	Tetraspanin related			
AT3G22400	lox5/PLAT/LH2 domain	AT1G17420	lipoxygenase 3 (PLAT/LH2 domain)	
AT1G18510	tetraspanin 16 (TET8 RELATED)	AT4G02380	senescence-associated gene 21	
AT4G02380	senescence-associated gene 21	AT2G29350	senescence-associated gene 13	
AT2G29350	senescence-associated gene 13			
	EXPO Related			
AT5G49830	exocyst complex component 84B	AT5G12370	exocyst complex component sec10	
AT4G02350	exocyst complex component sec15B	AT2G32750	Exostosin family protein	
AT2G28110	Exostosin family protein	AT4G02350	exocyst complex component sec15B	
AT1G21170	Exocyst complex component SEC5			
AT5G58430	exocyst subunit exo70 family protein B1			
	RNA binding proteins			
AT1G69440	Argonaute family protein	AT1G31280	Argonaute family protein	
AT1G31290	ARGONAUTE 3	AT1G31290	ARGONAUTE 3	
		AT1G69440	Argonaute family protein	
AT3G01540	DEAD box RNA helicase 1	AT5G26742	DEAD box RNA helicase (RH3)	
AT5G07120	sorting nexin 2B	AT5G07120	sorting nexin 2B	
AT1G15240	Sorting nexin, C-terminal	AT5G58440	sorting nexin 2A	
AT3G23830	glycine-rich RNA-binding protein 4	AT3G23830	glycine-rich RNA-binding protein 4	
	Heat Shock proteins			
AT3G07770	HEAT SHOCK PROTEIN 89.1	AT3G13860	heat shock protein 60-3A	
AT1G11660	heat shock protein 70 (Hsp 70)	AT1G11660	heat shock protein 70 (Hsp 70)	
AT4G02100	Heat shock protein DnaJ	AT3G07770	HEAT SHOCK PROTEIN 89.1	
AT3G13860	heat shock protein 60-3A	AT4G24280	chloroplast heat shock protein 70-1	

Table 1: displays the list of proteins identified in mass spectrometry data from both viroid-infected and
healthy EVs. It includes plant EV-related proteins such as Syntaxins, Tetraspanins, and Expo proteins.
Additionally, RNA-binding proteins and Heat shock proteins, previously reported to be associated with plant
EVs, are also included. The table further highlights PP2 proteins identified specifically in the EVs from viroid-
infected plants.

TET8 is a conserved EV marker protein in cucumber while PP2 is a newly identified RNA-binding protein localized within EVs

As tetraspanins were detected in the proteome analysis, the association of cucumber TET8 (CsTET8) with cucumber EVs was examined through immunoblotting. First, it was confirmed through immunoblot of transiently expressed CsTET8 that AtTET8 can effectively recognize CsTET8 in cucumber plants (Supplementary data S7-S11). Subsequent western blotting with AtTET8 antibodies on cucumber samples revealed a band around 30 kDa in cucumber leaf and AWF, consistent with the size of AtTET8 in Arabidopsis (Fig. 3a and 3b). To explore the association of TET8 with cucumber EVs, different fractions from SEC columns were analyzed by western blotting. In the western blot, EV fractions F9 and F11 were positive for TET8 protein (Fig. 3c-upper panel), suggesting that TET8 is localized within the EVs. To investigate whether TET8 is exposed on the surface of cucumber EVs, the EVs were treated with trypsin to cleave surface proteins. However, protein remained detectable in the EV fractions after trypsin treatment, suggesting that these proteins were not easily accessible to the enzyme (Fig. 3c). This indicates that TET8 is embedded within the EVs and protected by the vesicle membrane. To further confirm this localization, the EVs were disrupted using detergent followed by trypsin treatment, which resulted in the degradation of TET8 protein bands. This confirmed that TET8 is located inside the EVs and is protected by the vesicle membrane under normal conditions, but becomes exposed when the vesicles are disrupted. This experiment provides strong evidence that TET8 is an integral component of cucumber EVs, with its function conserved across plant species.

PP2 is another interesting protein identified in the proteome analysis. In order to investigate the association of PP2 with cucumber EVs, a western blot analysis using cucumber cell lysate and AWF was performed, which confirmed the presence of CsPP2 in both the cells and the apoplast (Fig. 3b). Further, analysis of CsPP2 in EV fractions revealed that PP2 is co-eluted with TET8 in SEC fractions 9 and 11 (Fig. 3c). Moreover, a protease protection assay revealed that CsPP2 is located inside the vesicles, as an intact PP2 band remained detectable after treatment with trypsin followed by detergent. However, the protein was degraded when the EVs were ruptured with detergent and then treated with trypsin (Fig. 3d). Overall, these assays suggest that PP2 is present both in the apoplast and inside the vesicles of cucumber plants.

Figure 4: Characterization of TET8 and PP2 from EVs infected with ASSVd.

(a) Western blot detection of TET8 in arabidopsis cell lysate (At-CL) cucumber cell lysates (Cs-CL) using Arabidopsis anti-AtTET8 antibody, revealing a band around 30 kDa, which corresponds to the size of TET8 in both Arabidopsis and cucumber CL (lower panel) western blot of the same extracts with anti-Actin antibody. (b) Western blot of cucumber CL and AWF with anti-AtTET8 and anti-CsPP2 antibody. Western blot of the same extract with anti-actin antibody are used as loading controls. (c)Western blot analysis of EV fractions (8-11) isolated by SEC, using AtTET8 antibodies to detect TET8 and PP2 in fractions 9 and 11 (upper lane in upper and lower blot). The fractions were treated with trypsin to degrade the surface proteins of the EVs, followed by detergent-induced rupture before loading onto the membrane. The lower lane shows the degradation of TET8 and PP2 from inside the EVs when ruptured with detergent before trypsin digestion, suggesting both proteins are localized inside the EVs. (d) Western blot analysis using anti-AtTET8 and anti-CsPP2 antibodies of pooled EV fractions (9 and 11) with different treatments—no treatment, trypsin followed by detergent, and detergent followed by trypsin—indicating that TET8 and PP2 are localized both inside and outside the EVs. Western blot of the same extract with anti-actin antibody are used as loading controls.

PP2 may aid in the sorting of viroid and other plant RNAs into TET8-specific exosomes and TET8 expression is triggered during viroid infection

The co-localization of TET8 and PP2 in the same EV fractions indicates a potential interaction between these two. To validate this we carried out immunoprecipitation of PP2 using CsPP2 antibody from the total cell lysate and AWF (treated with detergent to release the EV proteome) of viroid infected plants and detected the presence of TET8 in the pull down sample. Our immunoblot analysis revealed the presence of TET8 protein in the PP2 immunoprecipitated from the cell lysate as well as AWF (Figs. 4a). We further corroborated this interaction using the AlphaFold3 tool that predicted a pTM score of 0.47 (close to a good structure score of 0.5) for the PP2-TET8 interaction (Fig. 4b). The CsTET8 ortholog in Arabidopsis, TET8_ARATH, contains an extracellular domain (EC2) spanning amino acids 97-235 and the CsPP2 likely interact with CsTET8 at its EC2 domain (Fig. 4b). These results validate the interaction between TET8 and PP2 in the cucumber cell and apoplast.

Considering that PP2 is an RNA-binding protein, the interaction between TET8 and PP2 suggests that PP2 may play a role in sorting RNAs into TET8 exosomes. To explore this interaction, RNA was isolated from PP2-immunoprecipitated samples and the presence of ASSVd RNA was detected by RT-PCR. The results confirmed the co-immunoprecipitation of ASSVd RNA along with PP2 from the total cell lysate and apoplast (Fig. 4a). Overall these results suggested that PP2, ASSVd, and TET8 might exist as a trimeric complex in cucumber plants. To get a deeper insight into this interaction we specifically carried out immuno-enrichment of TET8 labeled exosomes from the total EV pool as reported in the arabidopsis (He et al., 2021). Our immunoblot analysis of the enriched TET8 exosomes revealed co-presence of PP2 and RT-PCR analysis of the same extracts showed presence of ASSVd in the TET8 specific exosomes (Fig. 4c).

We further validated this model through interactions using Alphafold3, which revealed the tripartite interaction, in which PP2 interacts with the viroid RNA on one side and TET8 on the other (Fig. 4d). However, TET8 did not show any direct interaction with the viroid RNA. The model predicted the central conserved region (CCR) of the viroid RNA, spanning nucleotides 81-96, was in closest proximity to PP2 at amino acids 1-8, 41-47, and 91-101, with CsPP2 motifs spanning amino acids 41-47 and 91-101 containing known double-stranded RNA-binding motifs

(dsRBM). Additionally, TET8 protein motifs 186-193 and 208-211 were in close proximity to

PP2 regions 78-86, 107-111, and 202-206, which lie within the EC2 of CsTET8 (Fig. 4d). Thus,

using a combination of biochemical and bioinformatic assays, we demonstrate that TET8

specific exosomes harbor ASSVd and PP2 where PP2-TET8 directly interact with each other at

EC2 domain of TET8 but there is no direct interaction between ASSVd-TET8 further supporting

the notion that PP2 might facilitate loading of RNAs in TET8 specific exosomes. To further

demonstrate that CsPP2 could be a potential protein facilitating the sorting of RNAs inside EVs

the PP2-immunoprecipitated RNAs were separated using denaturing PAGE and stained with

Diamond nucleic acid stain, a binding between PP2 and RNAs of various sizes was observed in

both whole cell lysate and the apoplast/EV fractions (Fig. 4e) further strengthening the role of

PP2 as a RNA sorting protein inside vesicles.

The expression of TET8 induces upon fungal infection in arabidopsis (He et al., 2021). We

monitored the dynamics of TET8 protein expression in plants infected with ASSVd transcripts.

Western blot analysis revealed that TET8 expression gradually increased from 7 to 28 dpi in

ASSVd-infected plants, whereas minimal expression was observed in mock-inoculated plants

throughout the 21-day period (Fig. 4f). These results suggest that TET8 expression is induced

relatively late in the plant's development, around 28 dpi, in cucumber. However, TET8

expression was elevated as early as 14 dpi in viroid-infected plants, indicating that ASSVd

infection triggers the up-regulation of TET8 protein expression at an earlier stage compared to

mock-infected plants. However, PP2 expression did not change in the same plants following

viroid infection (Fig. S12).

Figure 5: Interaction between PP2-TET8 and ASSVd

(a) Western blot analysis showing the detection of PP2 in immunoprecipitates of CsPP2 from the cell lysate (upper panel) along with the detection of TET8 via western blot and ASSVd RNA by RT-PCR in the same sample. The lower panel shows Western blot analysis of PP2 detection in immunoprecipitates from the apoplast+EVs fraction (upper panel), with TET8 detection via western blot and ASSVd RNA via RT-PCR in the same sample. (b) Interaction between CsTET8 and CsPP2 predicted using the Alphafold 3 tool, with a pTM score of 0.47, indicating a structure close to the true model. (c) Western blot showing the immuno-enrichment of TET8 exosomes from the total EV pool, with the detection of PP2 in the same extract. ASSVd

detection in the immunoenriched TET8 exosomes by RT-PCR is also presented. (d) Interaction between CsTET8, CsPP2, and ASSVd predicted using Alphafold 3, with a pTM score of 0.33, indicating a structure close to a good model. (e) Total RNA was extracted from the CsPP2-immunoprecipitated sample and analyzed on a denaturing gel. The left panel shows RNA immunoprecipitated with PP2 from leaf tissue, while the right panel shows RNA immunoprecipitated with PP2 from the apoplast + EVs fraction. (f) Western blot analysis showing the simultaneous detection of TET8 and anti-actin as a control in ASSVd-infected samples. The results indicate an upregulation of TET8 expression upon viroid infection compared to mock-infected plants.

EV-encapsulated ASSVd RNA is efficiently delivered via mechanical inoculation and is readily acquired and transmitted by whiteflies

To confirm the role of extracellular vesicles (EVs) in viroid transmission, we conducted an experiment where a group of ten whiteflies (*Trialeurodes vaporariorum*) were fed for 48 hours with three different treatments: intact EVs, AWF (apoplast with EV fractions depleted), and water, in addition to an artificial diet (Walia et al., 2015). After the feeding period, five whiteflies were analyzed for ASSVd presence using RT-PCR, while the remaining five were allowed to feed on healthy cucumber plants. The RT-PCR results revealed that whiteflies were able to acquire ASSVd from the intact EVs (Fig. 6a). Interestingly, a form of infectious ASSVd was also present in the apoplast that had been depleted of EVs, and whiteflies were able to acquire it from there as well (Fig. 6a). However, the PCR signal was significantly stronger for RNA acquired from intact EVs compared to that from the apoplast, indicating that the viroid is more stable and protected from RNases when encapsulated within EVs. This suggests that EVs might facilitate more efficient acquisition of the viroid by whiteflies compared to naked RNA or as RNP complex. No ASSVd was detected in whiteflies fed with water.

Furthermore, plants that were fed with viroid-carrying whiteflies were tested for viroid transmission at 14dpi using RT-PCR, which confirmed that the viroid could be actively acquired and transmitted from EVs through the whiteflies (Fig. 6b). Additionally, we demonstrated that mechanical inoculation of healthy cucumber plants with EVs could also deliver the viroid. For this, we inoculated healthy cucumber plants with intact EVs, AWF, or water. At 14 dpi, RT-PCR on the plants confirmed the presence of ASSVd infection, showing that both the apoplast and EVs of infected plants contained infectious forms of the viroid (Fig. 6c). In contrast, plants inoculated with water showed no viroid-specific amplification. Overall our results suggest that EV encapsulated viroid RNA can be readily acquired and transmitted by whiteflies to healthy

391 plants.

a

Healthy whiteflies fed with

Ladder EV^I water EV^I AWF-EV

500bp

300bp   330bp

b

Infected whiteflies fed   Mock whiteflies fed
on Healthy Plants   on Healthy Plants

P1   P2   Ladder   P3   P4

                                              330bp

c

Healthy plant infected with sap of

Ladder Leaf AWF EV Mock

500bp

300bp   330bp

d

ASSVd export in apoplast

Figure 6: Active transmission of ASSVd to healthy plants and whiteflies via infected EVs.

(a) RT-PCR showing detection of the viroid within EVs, whiteflies fed with intact EVs, apoplast lacking EVs, and water as the mock control. (b) RT-PCR showing detection of ASSVd in plants fed with whiteflies that were pre-fed on viroid-infected EVs, with water serving as the control. (c) RT-PCR analysis of cucumber plant sap inoculated with leaf tissue, AWF, and EVs from viroid-infected plants, revealing a 330 bp amplification corresponding to the ASSVd genome using viroid-specific primers. Plants inoculated with water, used as a mock control, did not show the presence of the viroid, serving as a negative control in the assay. (d)

A graphical summary of the key findings from the study. In brief, ASSVd RNA (both circular and linear
forms) and VdsRNAs can be packaged into EVs. PP2 interacts with ASSVd RNA and the EC2 domain of
TET8 within the cell, potentially facilitating the packaging of ASSVd RNAs into EVs. Additionally, ASSVd
RNA is modified with m6A inside the cell and contains exomotifs that may aid in its loading into EVs. Once
inside the EVs, the viroid RNA is transported to the extracellular space via these vesicles. Viroid RNAs also
exist in the apoplast independently of EVs, either as naked RNA species or as RNP complexes. These viroid
RNAs can be acquired by whiteflies, and likely by fungi as well, in the form of EV-encapsulated RNA or
naked RNAs as RNP complexes, and then transmitted to healthy plants.

Discussion

Viroids, unlike bacteria and fungi, are intracellular pathogens that replicate within the nucleus or

chloroplast of the cell. They traffic cell to cell through cytoplasmic connections, plasmodesmata,

and through phloem-mediated routes for their long-distance movement. Our previous studies

have highlighted the potential role of viroids in the cross-kingdom trafficking, such as the

acquisition and transmission of ASSVd RNA by whiteflies to neighboring plants (Walia et al.,

2015). Later, this phenomenon was also reported in natural interactions between apple plants and

their fungal pathogens (Tian et al., 2022). These observations, combined with research on the

role of EVs in RNA trafficking during plant-fungal interactions (Cai et al., 2018; Wang et al.,

2024), raised important questions about whether the viroid RNAs, typically considered

intracellular pathogens, could also be transported in EVs. Our study aimed to investigate the

presence of ASSVd RNA in the extracellular space including EVs. Interestingly, our results

revealed that viroid RNA also exists in the apoplast, and different forms of ASSVd RNA—

circular, linear, and small RNAs—are packaged into the EVs of cucumber plants. This indicated

that viroids may employ EVs for *in planta* and cross-kingdom trafficking, as EVs could protect

the RNA from host RNases and help viroids cross biological barriers. In animals, certain factors

have been identified that assist in the selective loading of RNAs into EVs. For example, certain

sequences have been identified in animals that code for the loading of microRNAs into EVs

(Martin et al., 2022). These motifs are referred to as EXOmotifs. In search for the identification

of such factors that could preferentially facilitate the transport and packaging of viroid RNAs

inside EVs, we observed an enrichment of five EXOmotifs in the ASSVd genome. Interestingly,

sequence-specific screening of most of the viroids revealed a pattern of exomotif enrichment in

their genomes (Fig. S13). For example, the UGUG motif is present in 24 out of 30 viroids,

CGGGAG in 2 out of 30, CCGGGG in 16 out of 30, and CCGGUG in 20 out of 30. Notably, the

majority of viroids, on average, possess five enriched exomotifs in their genomes. Furthermore,

the 25-nt long ACCCUGCCGCCUGGACUCCGCCUGU motif is specifically present in TASVd,

which is known to be transmitted by bumblebees through encapsulation in potato leaf roll virus. Apart from EXOmotifs in animals as well as in plants, the m6A modification of RNA has been linked to the sorting of RNAs into EVs. For example, it has been reported that m6A modification of miRNAs alters their binding affinity to AGO1 protein and increases their association with RNA-binding proteins subsequently facilitating their extracellular export (Sabrina Garbo et al., 2024). In plants, the apoplastic localized circular RNAs and long noncoding RNAs are enriched in m6A which likely facilitates their export into EVs (Karimi et al., 2022). In this study, we found that some proportion of EV-localized viroid RNA is modified with m6A, however, which of the three forms, whether circular, linear, or small RNA form is modified with m6A still needs to be investigated. Although the biological relevance of this modification is unclear, it is possible that it might have implications on viroid encapsulation within the EVs. Some viroid RNAs are known to undergo reversible modifications, such as m6A methylation (Pavel Vopalensky et al., 2024), thus it is plausible that ASSVd RNA is modified with m6A to enhance its export.

An intriguing finding of our study is the identification of PP2, a phloem-specific protein, in the apoplast and EVs of cucumber. A recent study examining the enrichment of the 26S proteasome from melon EVs reported the presence of PP2 in EVs from aphid-infested plants (Sánchez López, 2024). This suggests that the localization of PP2 in the EVs of the *Cucurbitaceae* species could represent a conserved phenomenon. PP2 is a phloem lectin with dsRNA binding motifs, capable of increasing the size exclusion limit of plasmodesmata, and has been implicated in the movement of various endogenous and pathogenic RNAs within the phloem (Gomez and Pallas, 2004). In our previous work, we demonstrated that the viroid-PP2 complex is more efficiently acquired and transmitted by whiteflies compared to the naked viroid RNAs (Walia et al., 2015). In the current study, through a feeding assay of EVs to whiteflies, we observed that viroid RNA is acquired and transmitted more efficiently by the whiteflies when packaged into EVs rather than naked RNA or as an RNP complex. This suggests that although viroid RNA can be acquired by whiteflies in various forms—either as a free RNA molecule, an RNP complex or encapsulated within EVs—the transmission through EVs provides a dual advantage: protection from host RNases and a safer route into new host kingdoms. Interestingly, it has been proposed that cross-kingdom trafficking of RNAs can occur via both EVs and RNP complexes. One research group reported that RNA-binding proteins such as AGO1, RNA helicases, and annexins can mediate RNA loading into vesicles and facilitate subsequent cross-kingdom trafficking (He et al., 2021;

Cai et al., 2019; Wang et al., 2024). Additionally, another group demonstrated that apoplast-
localized proteins like GRP7 and AGO2 can bind to the RNAs in the apoplast, potentially aiding
in the cross-kingdom movement of RNAs as RNP complexes (Karimi et al., 2022). Combining
these findings, we show using viroid RNA as a model, that cross-kingdom trafficking of RNAs
can occur both as RNP complexes and via EVs, although potentially at different intensities.
However, the relevance and function of viroid RNAs and PP2 within EVs, as well as the
potential for viroids to be transmitted through EV encapsulation, requires further investigation.
The presence of the PP2-viroid complex in the apoplast and its association with TET8 suggests
that PP2 may be a novel protein that assists in sorting RNA into EVs in *Cucurbitaceae* plants. It
is quite possible that viroids can channelize the EV-mediated route for their systemic movement
inside plants, as hypothesized for the movement of endogenous RNAs and viral RNAs in plants
(Kehr et al., 2018).

The EVs proteome from viroid-infected plants implicated changes in nuclear proteins
specifically related to non-coding RNA-related biological processes. Viroids are nuclear-
replicating pathogens and their effect on nuclear protein turnover and their subsequent export via
autophagy or extracellular vesicles warrant further investigation. These findings correlate with
cancer disease progression, where genomic DNA and nuclear components are exported by the
formation of amphisome-like structures and their fusion to vesicles (Akira et al., 2019). It is quite
possible that viroids may influence similar processes during infection, and hence,
characterization of DNA and RNA from EVs during viroid infection might shed light on these
unexplored processes related to viroid pathogenicity in plants. The induced expression of TET8
protein suggest that it acts as a positive regulator of viroid stress and could be used as a marker
protein for viroid stress, while CsPP2, despite its binding with viroid RNA, may not play a direct
role in the defense against viroid infection.

Overall, through our study using ASSVd as the model RNA system, cucumber as its host, and
whitefly as its vector, we propose that viroid RNAs are exported into the plant apoplast and
packaged into EVs from where they can be acquired by the whiteflies (Fig. 6d). They may use
EVs as a means of cross-kingdom trafficking and their interactions with RNA-binding proteins
such as PP2 could assist their loading into exosomes which improves their transmission
efficiency. Moreover, the presence of factors like EXOmotifs or m6A modification may assist

viroid RNA in their packaging inside EVs. The precise functional roles of small RNAs in EVs,
particularly those derived from viroids and their characterization, remain unclear. These small
RNAs may influence the plant's immune response or aid in the viroid's transmission to other
organisms, such as insects or fungi. Understanding the mechanisms by which viroids and other
RNA molecules are sorted into vesicles and transferred across biological boundaries will offer
valuable insights into cross-kingdom RNA trafficking, with implications for viroid biology,
RNA-based inter-organism communication, and the ecological impact of RNA transfer.

Material and Methods

Plant Growth

504 *Cucumis sativus* (Var. Summer Green) and *Nicotiana benthamiana* seeds were cultivated in pots
505 containing a mixture of cocopeat, compost, and vermiculite/Perlite in a 1:1:1 ratio. The plants
506 were kept in an insect-proof plant growth chamber equipped with fluorescent tubes under
507 controlled conditions with a temperature of $24\pm2°C$, relative humidity of 70-80%, and a 12-hour
508 light/12-hour dark cycle.

In vitro synthesis of dimeric ASSVd (+) RNA and transcript inoculation

To generate positive-sense RNA transcripts of ASSVd, recombinant plasmids containing the
dimeric ASSVd insert in the correct positive orientation within the pGEMT-easy vector were
linearized using *Spe*I. In vitro transcription was carried out using T7 Highscribe Kit (NEB, USA)
according to the manufacturer's protocol. Approximately 100 ng of the RNA transcripts were
subsequently used to infect the first true leaf of cucumber plants through mechanical inoculation
using carborundum-dust. The water-inoculated plants were used as controls in all experiments.

RNA extraction and RT-PCR for ASSVd detection

Total RNA was extracted from 100 mg of cucumber leaves using the RNAiso plus (TakaraBio,
USA; Walia et al., 2020). Briefly, 100mg of the cucumber leaves (viroid infected and mock)
were frozen in liquid nitrogen, ground in Trizol buffer, and incubated with chloroform. After
centrifugation, the supernatant was mixed with isopropanol for RNA precipitation and stored at -
20°C. The RNA pellet was then washed with 75% ethanol and eluted in 20 µL of nuclease-free
water. The RNA was quantified using Quantus™ Fluorometer (Promega) and 1µg of RNA was

used for RT-PCR with viroid-specific primers ICc (GTCGACGACGACAGGTGAGTT) and ICh

(GTCGACGAAGGCCGGTGAGAA) (Walia et al., 2014). Reverse transcription was performed

using verso cDNA synthesis kit (Thermo Scientific, USA) at 42°C for 60 minutes, followed by

PCR using Phusion DNA polymerase (Thermo Scientific, USA) under specified cycling

conditions for ASSVd detection (Walia et al., 2014). The PCR products were analyzed on 1.5%

agarose gel electrophoresis.

Isolation of AWF

Apoplastic washing fluid (AWF) from cucumber leaves was isolated as described previously

(O'Leary et al., 2014). The leaf petioles were excised, followed by washing and drying the leaves

(Fig. S1). The leaves were placed in a 60 mL syringe filled with an infiltration buffer (20mM

MES, 2mM $CaCl_2$, 0.1M NaCl, pH 6.0). Carefully without damaging the leaves, a negative

pressure was created by pulling the plunger gently until the leaves turned dark green. The leaves

were removed and dried and 3-4 leaves rolled were placed in a 12 mL syringe inside a 50 mL

falcon tube. The tube was wrapped with parafilm and centrifuged at 4,000 rpm for 30 minutes to

collect the AWF.

Trypan blue staining of leaves after AWF isolation for cell death analysis

The integrity of the isolated AWF was confirmed by cell death assay by staining the leaves in

trypan blue solution using the protocol as described (http://resources.rothamsted.ac.uk).

RNA isolation and RT-PCR detection of ASSVd from AWF

One ml of isolated AWF was concentrated to 100µl using an Amicon® Ultra Centrifugal Filter,

10 kDa MWCO (Millipore, USA). RNA was isolated using RNAiso plus (Takara Bio, USA).

Briefly, 1ml of RNAiso plus was added to 100µl of AWF followed by the addition of the

chloroform. The RNA was precipitated using isopropanol and suspended in 30µl of RNase-free

water. One microgram RNA was used for RT-PCR as mentioned earlier.

Isolation of EVs

Pure-EVs SEC columns were used to isolate EVs from cucumber plants (mock/viroid infected).

For EV isolation, 20 mL of isolated AWF was concentrated to 2 mL using an Amicon filter unit

(10kDa cutoff) (Millipore, USA) and was subjected to Pure-EVs SEC columns (HansaBiomed

Life Sciences, Estonia) according to the manufacturer's instructions. Briefly, the column was
washed with 3 volumes of 1x PBS buffer to remove preservative buffer residues. The sample
containing EVs (up to 2 mL) was then loaded into the column and eluted with $1\times$ PBS buffer as
the mobile phase. Fractions collected were labeled 1-7 as void volume, 8-13 as the EV fraction,
and 14-23 as the protein sample (Fig. S3).

Dynamic Light Scattering (DLS) analyses of EVs

The DLS analyses were performed in a Litesizer 500 (Anton Paar, Germany). 10 µL of isolated
EVs were suspended in 990 µL of 1x sterile PBS buffer, loaded into a disposable cuvette, and
placed in the litesizer 500. For reproducibility, and standardization, the parameter values were
fixed as follows; Temp-25°C, equilibration time-30s, processed runs-5, and Time for each run
10s. The size or homogeneity of exosomes was recognized through the radius or the % Intensity,
respectively.

Transmission electron microscopy of EVs

564 Purified EVs (10 µl) were diluted with an equal volume of 4% paraformaldehyde at 4°C for 30
565 min, and then carefully placed on a carbon-coated 300-mesh copper grid for 20 min, followed by
566 fixed by 1% glutaraldehyde for 5 min. After two washings, the grids were contrasted with 2%
567 uranyl acetate and then washed again two times. The morphology of isolated exosomes was
568 visualized with TEM (MAKE: JEOL MODEL: JEM 2100 plus).

Trypsin, RNase treatment, RNA isolation, and RT-PCR from EVs

To determine whether the viroid RNA was present inside or outside EVs, RNase protection
assays were performed (Fig. 1c). 100µl of Purified EVs were treated with 25 µg/µL trypsin
(Sigma Life Sciences, USA). Samples were incubated at 37°C for 16-18 hours, followed by the
addition of 1U of Proteinase K (Invitrogen, USA)/1mM of PMSF (Sigma, USA) to inactivate
trypsin. For isolating the RNA from the EV surface the above EV samples were precipitated with
3 volume ethanol and 1/10 sodium acetate overnight followed by isolation of RNA with RNAiso
plus next day. For isolating the RNA inside EVs the purified EVs (100µL) were treated with
trypsin + trypsin inhibitor + RNaseA + RNase inhibitor to degrade the outside RNA followed by
treatment with 1% (v/v) Triton X-100 (Himedia) to rupture the EVs. RNA isolation using
RNAiso plus was carried out to isolate the RNA inside EVs. The isolated RNA was quantified

and 200ng of the total RNA was used for RT-PCR from the detection of ASSVd from EV^O and

EV^I as described above. The concentration of trypsin and RNase used in the assays was

standardized by taking different dilutions of trypsin and RNase for protein degradation and RNA

degradation respectively (Fig. S14 and S15).

Urea PAGE and Northern Blot for ASSVd detection

Viroid-infected samples from total leaf, AWF, and EVs (EV^O and EV^I) along with mock RNA

from leaf were analyzed using Denaturing Polyacrylamide gel electrophoresis (15%

polyacrylamide and 7M Urea). The RNA samples (3μg for AWF, EV^O, EV^I 10ug, and 3ug for

CL from infected and 3μg from Mock) were denatured at 65°C in 2x RNA dye (Thermo

Scientific, USA) for 10 minutes and were separated in 0.5× TBE buffer for 2h at 90V. The total

RNAs were visualized by staining with Diamond Nucleic acid stain (Promega) and then

transferred onto a nylon membrane using 1× TBE transfer buffer. The RNAs were cross-linked

on a nylon membrane in a UV crosslinker followed by pre-hybridization at 65°C for 3hrs. The

membrane was incubated overnight at 65°C in hybridization buffer with gentle shaking along

with a 3' labeled biotin probe. The membrane was rinsed with washing buffer (1× SSC. 0.1%

SDS) thrice for 10 minutes at room temperature and blocked in a blocking buffer (0.1% Malic

acid buffer (Roche). The membrane was incubated with HRP-conjugated anti-streptavidin

antibody (Sigma, USA) at 1:5000 dilutions for 1h. After a final wash in maleic acid and 0.3%

Tween-20 containing buffer, the blot was visualized using ChemiDoc (Protein Simple, Biotechne,

USA).

Generation of negative sense 3'-biotin labeled ASSVd probe

A 3'-biotin-labeled RNA probe was prepared and used for viroid detection using northern

hybridization. For probe synthesis, a PCR was carried out using ASSVd plasmid as a template

and   primers   FP1   and   RP4   (FP1:

AGTAATACGACTCACTATAGGGGGGAAACACCTATTGTGTT   and   RP4:

GGTAAACACCGTGCGGTTC) to generate negative sense viroid amplicon along with T7

promoter sequence. The desired amplicon of 330bp was cut and eluted. Two micrograms of gel

eluted product was directly used to generate RNA transcript using a T7 in vitro RNA

transcription kit (Thermo Scientific, USA). The transcript at the 3' end was ligated to pcp-biotin

(Jena Biosciences) using T4 RNA Ligase (Promega, USA). The reaction was kept at 16ºC

overnight. The labeled transcript was purified using a Monarch RNA cleanup column (New

England Biolabs, USA) and was used for hybridization at a concentration of 100ng/10ml of

hybridization buffer (Fig. S15).

Immunoprecipitation with m6A antibody

Total RNA was isolated from leaves, apoplastic fluid, and EVs using RNAisol plus as described

earlier and quantified using a Quantus fluorimeter (Promega, USA). For leaf samples, 200µg

RNA and for AWF and EVs, 5µg RNA was incubated with an m6A antibody (Epigenetek-A-

1801-020) in IP buffer (50mM Tris HCl pH 7.5, 150mM NaCl, 0.5% NP-40) at 4°C for 4-5

hours, followed by the addition of Protein A/G agarose beads (Merck, USA) and incubated

overnight. The samples were washed with IP buffer and eluted from beads using RNAiso plus.

The isolated RNA was suspended in 100µl of RNase-free water. For spotting on the membrane,

10µl of the RNA was denatured at 65°C for 10 minutes and applied to a nylon membrane (for

detecting with ASSVd probe) and cross-linked. The membrane was pre-hybridized, incubated

overnight with a negative sense ASSVd-specific biotin probe, washed, and blocked. After

incubation with HRP-labelled streptavidin (Sigma, USA), m6A-modified RNA was visualized

using a ChemiDoc system (Protein Simple, Biotechne, USA). For detecting with anti m6A

antibody the RNA was spotted on the nitrocellulose membrane and the membrane was blocked

in 5% skim milk solution with overnight incubation with 1:5000 dilution of anti m6A antibody

(Epigenetek-A-1801-020). The next day membranes were washed and incubated with goat anti-

rabbit HRP-conjugated secondary antibodies, washed and visualized using ECL substrate

(Biorad, USA), and imaged with a ChemiDoc system (Protein Simple, Biotechne, USA) or X-ray

film.

Mass spectrometry

EVs fraction (100µl) was treated with trypsin followed by incubation with trypsin inhibitor and

were then treated with 0.1% Triton X-100. The proteins were TCA precipitated and partially

resolved by polyacrylamide gel electrophoresis till they entered the resolving gel. The region was

excised and sequenced using HR-LCMS, SAIF facility at IIT-BOMBAY. Briefly, the gel slices

were subjected to in gel-tryptic digestion followed by desalting and purification and subjected to

LC/MS/MS analysis in Q-Exactive Plus Orbitrap high resolution MS (Thermo Scientific). The

MS spectra were acquired at a resolution of 70,000 in a mass range of 350-2500 m/z. Eluted

samples were used for MS/MS events (resolution of 17,500), measured in a data-dependent

mode for the 15 most abundant peaks (Top15 method).

Proteome analysis and GO enrichment tools

The proteome analysis was carried out using the software Thermo Proteome Discoverer 2.2

against the Arabidopsis (TAIR) and cucumber database (Cucurbit genomics-

http://cucurbitgenomics.org/). The ShinyGO v0.82 (ShinyGO 0.82 (sdstate.edu)) and string tools

were used (STRING: functional protein association networks (string-db.org) for GO analyses.

Venn diagram was constructed using the software Bioinformatics & Evolutionary Genomics

(Draw Venn Diagram (ugent.be).

Immunoblots

For immunoblots, cell lyaste was prepared by grinding 50mg of leaf tissue in 500ɥl of protein

extraction buffer (50mM Tris HCl pH 7.5, 150mM NaCl, 10mM $MgCl_2$, 2.5mM EDTA, 1mM

DTT, 10% Glycerol, 1x protease inhibitor, 1mM PMSF). 20ɥl of cell lysate, concentrated AWF,

and EVs (EVO and EVI) were mixed with 3 µL of 6× SDS loading buffer, heated at 99°C for 10

minutes, and loaded onto a 15% PAGE, and after run proteins were transferred to a nitrocellulose

membrane (for dot blots, the dots were applied on the nitrocellulose membrane directly without

the SDS dye). Membranes were blocked with 5% skim milk in TBST and incubated overnight at

4°C with primary antibodies (1:1000 for TET8 (PhytoAb-PHY2750A), 1:10000 for CsPP2

(Walia et al., 2015, 1:10000 anti-GFP (Bio Bharati Life Science, India) and 1:3000 anti-HA (Bio

Bharati, Life Science, India). Membranes were washed and incubated with goat anti-rabbit HRP-

conjugated secondary antibodies (Bio Bharati, Life Science, India). After a final wash, proteins

were visualized using the ECL substrate (Biorad, USA) and imaged with a ChemiDoc system

(Protein Simple, Biotechne, USA) or X-ray film.

RNA immunoprecipitation assay (RIP assay)

For RNA immunoprecipitation from the leaf, 1g/ml of viroid-infected leaf tissue was ground in

protein extraction buffer supplemented with RNase inhibitor and 500µl of lysate was used for

RNA immunoprecipitation assay. For RNA immunoprecipitation from infected AWF, 5ml of

AWF isolated from viroid-infected leaves was concentrated to 500µl solution using an Amicon

filter. Both the lysate and AWF were incubated with 5mg of CsPP2 antibody for 4h at 4°C with

rotation followed by the addition of protein A/G agarose beads overnight. The next day the beads

were pelleted with centrifugation at 3000 rpm for 5 minutes and washed three times with protein

extraction buffer supplemented with RNase inhibitor. The beads were eluted in 100ul of RNAse-

free water. Half of the beads were heated at 100°C for 10 minutes to release bead-bound proteins

for subsequent immunoblot. The rest of the beads were proceeded for RNA isolation using

RNAiso plus. The RNA was quantified and used for the detection of ASSVd by RT-PCR or

loaded on 15% denaturing PAGE to visualize the total immunoprecipitated RNAs using

Diamond Nucleic acid stain (Promega) as described earlier.

Immunoenrichment of TET8 exosomes

The isolated intact EVs were treated with trypsin overnight, followed by the addition of a trypsin

inhibitor. Next, RNase was added for one hour, and then RNase inhibitor was used to inactivate

the RNase. The intact EVs were incubated with anti-AtTET8 antibody in IP buffer without NP-

40 for 4 hours at 4°C with rotation, followed by the addition of protein A/G agarose beads

overnight. The next day, the beads were pelleted and washed with IP buffer containing 1%Triton

X-100 and RNase inhibitor. The beads were then eluted in RNase-free water, with part of the

eluate used for RNA isolation as previously described. The isolated RNA was tested for the

presence of ASSVd by RT-PCR, as described earlier. The remaining beads were heated and

loaded onto an SDS-PAGE gel for immunoblotting with anti-AtTET8 and anti-CsPP2 antibodies.

Dynamics of CsTET8 and CsPP2 upon viroid infection

Three cucumber plants were infected with ASSVd transcripts along with mock as control. The

samples were harvested at 0dpi, 7dpi, 14dpi, 21dpi and 28dpi. The lysates were prepared in 8M

urea, mixed with SDS dye, heated, and loaded on 15% PAGE. The proteins were transferred to

the membrane and immunoblotted with CsPP2 and AtTET8 antibodies as described earlier.

Bioinformatic analysis of the ASSVd-PP2-TE8 interactions

The amino acid sequences of the proteins were obtained from the following accession numbers

in GenBank: Tetraspanin-8 (TET8) *Cucumis sativus* Accession XP_004151158.1, Lectin

(phloem protein 2) *Cucumis sativus* Accession NP_001292700.1, and ASSVd RNA sequence

was of an isolate from Solan with Accession FM178284. The models were generated from the

above mentioned sequences by the Alphafold2 algorithm (Jumper et al., 2021). The quality was

assessed using pLDDT scores. The models were visualized using ChimeraX software v 1.7.1 (Meng et al., 2023). The interactions of PP2 with TET8 and trimeric interaction of PP2, TET8 with viroid RNA was studied using AlphaFold3 (Abramson et al., 2024; https://alphafoldserver.com/fold/20a59528fa980c38) that allows structural modeling and interactions of protein and RNA. The quality was assessed using pTM scores. The models were visualized using ChimeraX software v. 1.7.1 (Meng et al., 2023).

EV acquisition and viroid transmission assay

For ASSVd acquisition by whiteflies, ten whiteflies were fed each with AWF (depleted of EVs) and EV (depleted of outside RNAs) and water as control. The whiteflies were trapped in 50ml falcon tubes with their top covered with stretched parafilm. The AWF, EV, and water were placed on the parafilm along with food color (turmeric) and covered with a coverslip. The tubes were kept under the light for 48h and whiteflies were allowed to feed on the solution. After 48h the five whiteflies were killed using the cold shock method and RNA isolation and RT-PCR for ASSVd detection was done as described earlier. The remaining five whiteflies were made to feed on healthy cucumber plants in an insect proof racks. The whitefly fed plants were checked for the viroid presence using RT-PCR method using ASSVd specific primers as described above.

For transmission assay using viroid-infected EVs and AWF, three cucumber plants each with AWF (depleted of EVs) and EV (depleted of outside RNAs) were sap inoculated along with leaf sap as positive control and water as negative control. The plants were checked for the ASSVd presence at 14dpi by RT-PCR method as described earlier.

Author contributions

Neha Devi: Formal analysis, methodology, and validation. **Vasudha Sharma**: Formal analysis and methodology. **Nisha Devi:** Formal analysis and methodology. **Ravi Gupta:** writing-review and editing (supporting). **Sunny Dhir:** conceptualization, data analysis, investigation, methodology, supervision (equal), writing- review and editing. **Yashika Walia:** conceptualization (lead), data curation and analysis (lead), investigation (lead), methodology, supervision (equal), validation, writing- original draft, writing- review and editing, funding acquisition.

Acknowledgments

We would like to thank the Management MMDU, for providing necessary research facilities. We would like to acknowledge SAIF at IIT Bombay for mass-spectrometry analysis and CIL facility at Panjab University Chandigarh for TEM imaging. We would also like to thank I-STEM facility, Jamia Hamdard, New Delhi for confocal microscopy. We acknowledge special thanks to Dr. Vipin Hallan, CSIR-IHBT Palampur for providing CsPP2 antibodies.

Funding

The work was supported by State University Research Excellence Award to Dr. Yashika Walia by Anusandhan National Research Foundation (ANRF), Government of India thorugh project number SUR/2022/001747, Department of Biotechnology, and Government of India through EMR grant BT/PR40936/AGIII/103/1255/2020 to Dr. Sunny Dhir, and to Bayer for providing MEDHA fellowship to Ms. Nisha Devi.

Disclosure of Interest: The authors report no conflict of interest

Ethics Statement: not applicable

Data availability statement: The data that support the findings of this study are available from the corresponding author (YW) upon reasonable request.

References

1. Wang, S., et al., *Plant mRNAs move into a fungal pathogen via extracellular vesicles to reduce infection.* Cell host & microbe, 2024. **32**(1): p. 93-105. e6.
2. Zhang, M., et al., *Plant derived edible nanoparticles as a new therapeutic approach against diseases.* Tissue barriers, 2016. **4**(2): p. e1134415.
3. Zand Karimi, H., et al., *Arabidopsis apoplastic fluid contains sRNA-and circular RNA–protein complexes that are located outside extracellular vesicles.* The Plant Cell, 2022. **34**(5): p. 1863-1881.
4. He, B., et al., *RNA-binding proteins contribute to small RNA loading in plant extracellular vesicles.* Nature plants, 2021. **7**(3): p. 342-352.
5. Ambrosone, A., et al., *Plant extracellular vesicles: Current landscape and future directions.* Plants, 2023. **12**(24): p. 4141.
6. Chen, A., B. He, and H. Jin, *Isolation of extracellular vesicles from arabidopsis.* Current protocols, 2022. **2**(1): p. e352.
7. Jankovičová, J., et al., *Tetraspanins, more than markers of extracellular vesicles in reproduction.* International Journal of Molecular Sciences, 2020. **21**(20): p. 7568.

8. Rutter, B.D. and R.W. Innes, *Extracellular vesicles isolated from the leaf apoplast carry stress-response proteins.* Plant physiology, 2017. **173**(1): p. 728-741.

9. Wang, J., et al., *EXPO, an exocyst-positive organelle distinct from multivesicular endosomes and autophagosomes, mediates cytosol to cell wall exocytosis in Arabidopsis and tobacco cells.* The Plant Cell, 2010. **22**(12): p. 4009-4030.

10. Ding, Y., et al., *Exo70E2 is essential for exocyst subunit recruitment and EXPO formation in both plants and animals.* Molecular biology of the cell, 2014. **25**(3): p. 412-426.

11. Weiberg, A., et al., *Fungal small RNAs suppress plant immunity by hijacking host RNA interference pathways.* Science, 2013. **342**(6154): p. 118-123.

12. Cai, Q., et al., *Plants send small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes.* Science, 2018. **360**(6393): p. 1126-1129.

13. Garbo, S., et al., *m6A modification inhibits miRNAs' intracellular function, favoring their extracellular export for intercellular communication.* Cell Reports, 2024. **43**(6).

14. O'Grady, T., et al., *Sorting and packaging of RNA into extracellular vesicles shape intracellular transcript levels.* BMC biology, 2022. **20**(1): p. 72.

15. Garcia-Martin, R., et al., *MicroRNA sequence codes for small extracellular vesicle release and cellular retention.* Nature, 2022. **601**(7893): p. 446-451.

16. Liu, Y.-f., et al., *CYP2A6 suppresses hepatocellular carcinoma via inhibiting SRC/Wnt/β-Catenin pathway.* Acta Pharmacologica Sinica, 2025: p. 1-12.

17. Flores, R., R.A. Owens, and J. Taylor, *Pathogenesis by subviral agents: viroids and hepatitis delta virus.* Current opinion in virology, 2016. **17**: p. 87-94.

18. Navarro, B., R. Flores, and F. Di Serio, *Advances in viroid-host interactions.* Annual Review of Virology, 2021. **8**(1): p. 305-325.

19. Hammond, R.W., *Economic significance of viroids in vegetable and field crops*, in *Viroids and satellites*. 2017, Elsevier. p. 5-13.

20. Branch, A.D. and H.D. Robertson, *A replication cycle for viroids and other small infectious RNA's.* Science, 1984. **223**(4635): p. 450-455.

21. Warrilow, D. and R. Symons, *Citrus exocortis viroid RNA is associated with the largest subunit of RNA polymerase II in tomato in vivo.* Archives of virology, 1999. **144**: p. 2367-2375.

22. Navarro, J.A. and R. Flores, *Characterization of the initiation sites of both polarity strands of a viroid RNA reveals a motif conserved in sequence and structure.* The EMBO Journal, 2000.

23. Rodio, M.-E., et al., *A viroid RNA with a specific structural motif inhibits chloroplast development.* The Plant Cell, 2007. **19**(11): p. 3610-3626.

24. Daròs, J.-A., et al., *Replication of avocado sunblotch viroid: evidence for a symmetric pathway with two rolling circles and hammerhead ribozyme processing.* Proceedings of the National Academy of Sciences, 1994. **91**(26): p. 12813-12817.

25. Sano, T., et al., *Identification of multiple structural domains regulating viroid pathogenicity.* Proceedings of the National Academy of Sciences, 1992. **89**(21): p. 10104-10108.

26. Reanwarakorn, K. and J. Semancik, *Regulation of pathogenicity in hop stunt viroid-related group II citrus viroids.* Journal of General Virology, 1998. **79**(12): p. 3163-3171.

27. SCHMITZ, A. and D. RIESNER, *Correlation between bending of the VM region and pathogenicity of different potato spindle tuber viroid strains.* RNA, 1998. **4**(10): p. 1295-1303.

28. Schnölzer, M., et al., *Correlation between structure and pathogenicity of potato spindle tuber viroid (PSTV).* The EMBO Journal, 1985. **4**(9): p. 2181-2190.

29. Afanasenko, O., et al., *Transmission of potato spindle tuber viroid between Phytophthora infestans and host plants.* Vavilov Journal of Genetics and Breeding, 2022. **26**(3): p. 272.

30. Leichtfried, T., et al., *Apple chlorotic fruit spot viroid: a putative new pathogenic viroid on apple characterized by next-generation sequencing.* Archives of Virology, 2019. **164**: p. 3137-3140.

31.   Walia, Y., et al., *Apple scar skin viroid naked RNA is actively transmitted by the whitefly*
Trialeurodes vaporariorum. Rna Biology, 2015. **12**(10): p. 1131-1138.

32.   Tian, M., et al., *Natural cross-kingdom spread of apple scar skin viroid from apple trees to fungi.*
Cells, 2022. **11**(22): p. 3686.

33.   Walia, Y., et al., *Identification of the herbaceous host range of Apple scar skin viroid and analysis*
of its progeny variants. Plant pathology, 2014. **63**(3): p. 684-690.

34.   O'Leary, B.M., et al., *The infiltration-centrifugation technique for extraction of apoplastic fluid*
from plant leaves using Phaseolus vulgaris as an example. Journal of visualized experiments:
JoVE, 2014(94): p. 52113.

35.   Vopalensky, P., et al., *Exploring RNA modifications in infectious non-coding circular RNAs.* RNA
biology, 2025(just-accepted).

36.   Sánchez-López, C.M., et al., *Phloem sap from melon plants contains extracellular vesicles that*
carry active proteasomes which increase in response to aphid infestation. Journal of Extracellular
Vesicles, 2024. **13**(10): p. e12517.

37.   Gomez, G., H. Torres, and V. Pallas, *Identification of translocatable RNA-binding phloem proteins*
from melon, potential components of the long-distance RNA transport system. The Plant Journal,
2005. **41**(1): p. 107-116.

38.   Kehr, J. and F. Kragler, *Long distance RNA movement.* New Phytologist, 2018. **218**(1): p. 29-40.

39.   Yokoi, A., et al., *Mechanisms of nuclear content loading to exosomes.* Science advances, 2019.
5(11): p. eaax8849.

Supporting data

Cucumber extracellular vesicles-localized tetraspanin and phloem protein 2 mediates the cross-kingdom trafficking of a pathogenic non-coding RNA

Neha Devi, Vasudha Sharma, Nisha Pahal, Ravi Gupta*, Sunny Dhir*, Yashika Walia*

Fig. S1 showing the procedure followed for isolation of AWF from cucumber leaves

Fig. S2: Coomassie staining of SDS PAGE with total protein extract isolated from AWF and whole cell lysate (CL).
Absence of Rubisco protein at 55kDa in AWF fraction compared to CL indicate purity of AWF and absence of
cytosolic contamination

Fig. S3: Trypan blue staining of leaves used for AWF isolation. Absence of any cell damage as indicated by clear
leaves suggest purity of isolated AWF free from cellular content.

856 **Fig. S4** showing the procedure followed for isolation of EVs from isolated AWF of cucumber

Fig. S5: TEM images of Healthy EVs and Viroid infected EVs. The size corresponding to exosomes app. 50nM and micro vesicles app. 200nM was observed in both isolated fractions

Fig. S6: Size measurement of Healthy EVs and Viroid infected EVs using DLS method. An average size of app. 200nM was observed in both healthy and viroid infected EVs.

We tested whether Arabidopsis TET8 (AtTET8) antibodies would cross-react with cucumber

Tet8 (CsTet8) protein and found that the Arabidopsis Tet8 (AT2G23810) ortholog in cucumber

is Tet8/Lipoxygenase domain 3 (Csa1g287020) that shared 65.67% homology with AtTet8 (Fig.

S7). Additionally, the 18 amino acids used to generate the peptide from the EC2 domain

(Phytoab) were over 90% identical between the Arabidopsis and cucumber Tet8, suggesting that

AtTet8 antibodies would cross-react with cucumber Tet8 (Fig. S8).

BLAST of Csa1G287020 vs. TAIR10
Match: AT2G23810.1 (AT2G23810.1 tetraspanin8)

876   **HSP 1** Score: 397.5 bits (1020), Expect = 6.5e-111
Identity = 176/268 (65.67%), Postives = 216/268 (80.60%), Query Frame = 1

```
Query: 1    MVKLSNNLVGILNFITFLLSIPIIAGGIWLSRQGTSDCEKFLDTPVIVIGVFLLLVSLAG 60
            M + SNNLVGILNF+ FLLSIPI+AGGIWLS++G+++CE+FLD PVI +GVFL++V++AG
Sbjct: 1    MARCSNNLVGILNFLVFLLSIPILAGGIWLSQKGSTECERFLDKPVIALGVFLMVVAIAG 60
```

```
Query: 61   FIGACCRVRWLLWIYLFVMFLLILLLFVFTIFAFAVTNKGAGKVLSNRGYKEYRLGDYSN 120
             IG+CCRV WLLW+YLFVMFLLILL+F  T+FAF VTNKGAG+ +  +GYKEY+LGDYS
Sbjct: 61   LIGSCCRVTWLLWVYLFVMFLLILLVFCITVFAFVVTNKGAGEAIEGKGYKEYKLGDYST 120
```

```
Query: 121  WLQNRVRNNKDWNRIRSCLVDGKVCNEVSQKFVGETVEQFYLEHLSSIQSGCCKPADECK 180
            WLQ RV N K+WN+IRSCLV+ KVC+++  KFV   V  FY EHL+++QSGCCKP+DEC
Sbjct: 121  WLQKRVENGKNWNKIRSCLVESKVCSKLEAKFVNVPVNSFYKEHLTALQSGCCKPSDECG 180
```

```
Query: 181  FEYKGPIQWDKPANVSSFANPDCGLWDNDPQKLCFDCESCKGGVLDNLKRNWKKVAIINI 240
            FEY  P  W K   +  NPDC  WDN +KLCFDC+SCK G+LDN+K  WKKVAI+NI
Sbjct: 181  FEYVNPTTWTK-NTTGTHTNPDCQTWDNAKEKLCFDCQSCKAGLLDNVKSAWKKVAIVNI 240
```

```
Query: 241  VILVFLIIVYSIGCCAFRNSKEENHYPR 269
            V LVFLIIVYS+GCCAFRN+K ++ Y R
Sbjct: 241  VFLVFLIIVYSVGCCAFRNNKRDDSYSR 267
```

884   **Fig. S7** Sequence Homology between CsTET8 and AtTET8 showing 65.67% identity between the two

885   proteins. The EC2 domain froM which the antibody of Phytoab AtTET8 is made is marked in red boxes

EC2 domain-97aa-235aa

Fig. S8: Multiple alignment between EC2 domain of CsTET8 and AtTET8 showing consensus sequences between

the two proteins

To further confirm this cross-reactivity, we cloned CsTET8 (Csa_1g287020), the cucumber

ortholog of AtTET8, into the pSITE2CA and HApBA plant binary vector with a GFP tag for

visualization. These recombinant plasmid and were then transiently expressed in Nicotiana plants,

and western blot analysis using anti-GFP, anti-TET8 and anti-AtTET8 antibodies confirmed that

the AtTET8 antibodies effectively recognized CsTET8 (Fig. S7) at size of approximately 55kDa

and 25kDa, respectively. Based on these findings, we proceeded to use the available AtTET8

antibodies in further experiments.

Generation of constructs

The CsTET8 gene (Csa1g287020) was amplified from cucumber plants using the RT-PCR

method. In brief, RNA was extracted from healthy cucumber plants with RNAiso Plus. Reverse

transcription was performed with the Verso cDNA synthesis kit (Thermo). PCR amplification

was carried out using Phusion DNA polymerase (Thermo) with specific primers (CsTET8F/R)

(Table S1). The PCR products were resolved on a 1.5% agarose gel, and the desired bands were

excised and purified using the GeneJET Thermo kit. The purified fragments were then cloned

into the pJET1.2 vector (Thermo) and transformed into TOP10 competent cells. The recombinant

plasmid was sequenced, confirming 100% identity to CsTET8.

The CsTET8 gene was further amplified using adaptor primers (CsTET8att F and CsTET8att R)

and universal primers (Adaptor F/R) (Table S1). The amplified fragment was ligated into the

PDONR207 vector using BP clonase (Invitrogen) and transformed into TOP10 competent cells.

The transformed cells were plated on LA plates containing 100 mg/ml kanamycin. Recombinant

plasmids were verified through colony PCR using the vector-specific primer (Nru R) and the

gene-specific primer (CsTET8 F).

Subsequently, the pDONR-CsTET8 clone was sub cloned into the pSITE2CA vector and the

HA-pBA vector (Chakrabarty et al., 2007) for expression under the CaMV35S promoter, with

N-terminal GFP and HA tags, respectively, using the LR clonase enzyme (Thermo Scientific,

USA). The ligated products were transformed into LA plates containing 50 mg/ml spectinomycin.

Recombinant clones were screened by colony PCR with the CaMV35S forward primer and

CsTET8R primer. The GFP-TET8 and HA-TET8 constructs were then transformed into the

Agrobacteriun strain GV3101 using chemical transformation and used for transient expression

analysis in *Nicotiana benthamiana* plants, as described by Walia et al. (2020).

Detection of transiently expressed CsTET8-pSITE2CA in *N.bentahamiana* with anti-GFP and anti AtTET8 antibody

Fig. S9: a and b: western blot of *N.benthamiana* cell lysate (CL) and AWF showing transient expression of CsTET8-pSITE2CA in and empty vector (pSITE2CA) with 25kDa band corresponding to the size of only GFP and app. 55kDa size of CsTET8+GFP detected with (a) anti-GFP antibody. (b)AtTE8 antibody. The band corresponding to the size of CsTET8+GFP is detected in AWF and marked in box. The arabidopsis TET8 band detected with AtTET8 antibody at 30kDa as positive control is marked in green box (Fig.S9b)

Fig. S10: western blot of *N. benthamiana* cell lysate (CL) showing transient expression of HA-CsTET8 and arabidopsis and cucumber cell lysate as control showing 30kDa band corresponding to the size of TET8 with anti-TET8 antibody and anti-HA antibody.

The sub cellular localization of CsTET8 was monitored by transiently expressing the CsTET8-
pSITE2CA in *N. benthamiana* which revealed the cytoplasmic and plasma membranous localization of
CsTET8.

Fig. S11: Localization of GFP-CsTET8 in *N. benthamiana* showing membranous localization. Speckles observed on plasma
membrane points towards its localization inside vesicles.

Fig. S12: western blot of cucumber cell lysate with CsPP2 antibody showing expression of PP2 protein in mock and ASSVd
infected cucumber plants at 0, 7, 14, 21 and 28 dpi.

Fig. S13: EXOmotifs observed in various viroids

Standaridization of trypsin treatment

Standaridization of RNAse treatment

Fig. S14 and S15: Standardization of trypsin and RNase concentration for treatment of EVs with trypsin and RNase for protease protection assay.

952
953

Table S1: Primers used during the study

Primer name	Sequence
ASSVdICC	GTCGACGAAGGCCGGTGAGAA
ASSVd ICH	GTCGACGACGACAGGTGAGTT
CsTET8 f	ATGGTCAAACTCAGCAATAATCTC
CsTET8 r	TCATTTCCATCTAGGATAATG
CsTET8 att F	AAAAAGCAGGCTCAATGGTCAAACTCAGCAATAATC
CsTET8 att R	AGAAAGCTGGGTATCATTTCCATCTAGGATAATG
Adaptor F	GGG GAC AAG TTT GTA CAA AAA AGC AGG CT
Adaptor R	GGG GAC CAC TTT GTA CAA GAA AGC TGG GT
CaMV35S	GACTCACTATAGGGAGAG
Nru R	ACGGAAACAGCTATGAC

961