UFMylation of 14-3-3ε coordinates MAVS signaling complex assembly to promote

antiviral innate immune induction

Isaura Vanessa Gutierrez[1], Moonhee Park[1], Lauren Sar[1], Ryan E. Rodriguez[2], Hannah M.

Schmidt[2], Daltry L. Snider[2], Gabriella Torres[1], K. Matthew Scaglione[2,3], Stacy M. Horner[1,2,4]

[1] Department of Integrative Immunobiology, Duke University School of Medicine, Durham, NC,

USA

[2] Department of Molecular Genetics and Microbiology, Duke University School of Medicine,

Durham, NC, USA

[3] Duke Center for Neurodegeneration and Neurotherapeutics, Duke University School of

Medicine, Durham, NC, USA

[4] Department of Medicine, Duke University School of Medicine, Durham, NC, USA

Correspondence: stacy.horner@duke.edu

Stacy M. Horner, Ph.D.

Duke University School of Medicine

213 Research Dr., Box 3053 DUMC

Durham, NC USA 27710

Phone: 919-684-1921

Abstract

Post-translational modifications regulate RIG-I signaling in diverse ways. We previously showed

that UFMylation, the covalent attachment of the ubiquitin-fold modifier UFM1 to proteins,

enhances RIG-I signaling by promoting its interaction with its membrane-targeting adaptor 14-3-

3ε. Here, we map UFM1 conjugation to lysines K50 and K215 on 14-3-3ε and demonstrate how

these UFMylation events control RIG-I signaling. Using *in vitro* and cellular UFMylation assays,

we reveal that K50R/K215R mutations abolish UFMylation and reduce type I and III interferon

induction following RIG-I activation. Unexpectedly, these mutations do not disrupt 14-3-3ε-RIG-I

interaction. Instead, they paradoxically enhance RIG-I interaction with MAVS while

simultaneously reducing 14-3-3ε-MAVS interaction. These findings establish UFMylation of 14-3-

3ε as an important control that shapes MAVS complex architecture to ensure optimal RIG-I

signaling and highlights the broader regulatory role of UFMylation in antiviral innate immunity.

Importance

Post-translational modifications provide regulatory control of antiviral innate immune responses.

Our study reveals that UFMylation of 14-3-3ε is required for RIG-I-mediated innate immune

signaling. We demonstrate that conjugation of UFM1 to specific lysine residues on 14-3-3ε

enhances downstream signaling events that facilitate interferon induction. It does this by

stabilizing 14-3-3ε association with the MAVS signaling complex and coordinating productive

complex architecture. By identifying the precise sites of UFMylation on 14-3-3ε and their

functional consequences, we provide insights into the regulatory layers governing antiviral innate

immunity. These findings complement emerging evidence that UFMylation serves as a versatile

modulator across diverse immune pathways. Furthermore, our work highlights how protein

chaperones like 14-3-3ε can be dynamically modified to orchestrate complex signaling cascades,

suggesting potential therapeutic approaches for targeting dysregulated innate immunity.

Introduction

The antiviral innate immune response is tightly regulated to ensure effective defense against viral infections while preventing excessive inflammation. This antiviral response to RNA viruses is activated by cytosolic RIG-I-like receptors (RLRs) RIG-I and MDA5, which detect pathogen-associated molecular patterns from RNA viruses and initiate the formation of a MAVS-signaling complex (1). Upon activation, MAVS undergoes oligomerization and recruits downstream signaling proteins to form a large multiprotein assembly known as the MAVS signalosome, which serves as a platform for signal amplification and propagation. This signaling cascade ultimately leads to the transcriptional activation of type I and type III interferons (IFN) and the production of IFN-stimulated genes (ISGs) that limit viral infection. Robust innate immune signaling is essential for effective adaptive immunity, but dysregulation of these pathways can result in autoimmune diseases and interferonopathies (2-4). Therefore, the induction of IFN is carefully regulated by mechanisms that temporarily activate, dampen, and ultimately shut down the signaling pathway.

Both 14-3-3ε and 14-3-3η are RLR regulatory proteins that control IFN induction. These proteins facilitate the translocation of RIG-I and MDA5, respectively, from the cytosol to intracellular membrane signaling sites to interact with the MAVS signalosome (5, 6). 14-3-3 proteins are highly conserved scaffolding proteins involved in diverse cellular processes, including signal transduction, apoptosis, and stress responses. Typically, they coordinate diverse cellular processes by binding to phosphorylated client proteins, although they are also known to bind to non-phosphorylated client proteins (7-13). Recent work has also suggested that 14-3-3 proteins can serve as protein chaperones, preventing aggregation and phase separation of their client proteins (14). The essential role of 14-3-3ε and 14-3-3η in antiviral innate immunity is underscored by the observation that multiple viruses have evolved specific mechanisms to antagonize these proteins. For example, orthoflavivirus NS3 proteins specifically target and inactivate both 14-3-3ε and 14-3-3η to prevent RLR signaling, the enterovirus 3C protease cleaves 14-3-3ε, and the influenza A virus NS1 protein binds to 14-3-3ε (15-18). These diverse viral targeting strategies highlight how critical 14-3-3 proteins are for mounting an antiviral innate immune response.

Post-translational modifications play a crucial role in regulating RIG-I signaling (19, 20). Recent studies by our group have shown that ubiquitin-fold modifier 1 (UFM1) and the proteins involved in UFM1 conjugation promote RIG-I signaling (21). UFMylation involves the covalent conjugation of ubiquitin-fold modifier 1 (UFM1), an 85-amino acid peptide, to lysine residues on proteins, utilizing an E1 activating enzyme (UBA5), an E2 conjugating enzyme (UFC1), and an E3 scaffold-like ligase complex (UFL1 and the membrane-anchored protein UFBP1) (22-30).

UFM1 is a non-degradative mark, and its addition to proteins can promote protein stability by

antagonizing ubiquitination or stabilizing interactions with other proteins (31-33). UFMylation

modulates various cellular processes, including the DNA damage response and genomic

instability, various aspects of ER homeostasis and translation, ER-Golgi transport, viral infection,

cancer progression, and both adaptive and innate immune responses (24, 25, 27, 28, 33-46).

In the context of innate immunity, UFMylation plays diverse regulatory roles. For example,

UFMylation promotes NLRP3 inflammasome activation (47), while in other contexts it can limit

LPS-mediated inflammation via its roles in regulating ER homeostasis and the unfolded protein

response (48-51). Additionally, UFL1 can bind STING during herpes simplex virus 1 infection to

maintain IFN induction (52). However, the overall relationship between UFMylation and viral

infection is complex. In some cases, UFMylation can promote viral infection. For example,

UFMylation of the ribosomal protein RPL26 enhances hepatitis A virus translation (53), and in

orthoflaviviruses, UFMylation promotes the production of infectious virions (54). Reflecting the

importance of UFMylation in host defense, viruses have evolved mechanisms to manipulate host

UFMylation pathways by targeting UFL1 to specific substrates. For example, orthoflaviviruses co-

opt UFL1 to aid in viral assembly (54), while Epstein-Barr virus targets UFL1 to MAVS for

UFMylation to dampen inflammasome activation (55). These diverse roles highlight UFMylation

as a key regulatory system in host-pathogen interactions.

Our previous work on UFMylation in the context of RIG-I signaling revealed that UFL1

translocates to intracellular membranes upon RIG-I activation and that UFL1 and UFM1 broadly

facilitate RIG-I interaction with 14-3-3ε, with 14-3-3ε exhibiting increased UFMylation in response

to RIG-I activation (21, 56). In this study, we established the molecular mechanism by which

UFMylation of 14-3-3ε promotes RIG-I signaling. Using cellular and *in vitro* UFMylation assays,

we identify lysine residues K50 and K215 as the mono-UFMylation sites of 14-3-3ε. Through

mutagenesis studies, we demonstrate that preventing UFMylation at these residues by

introducing K50R/K215R mutations impairs both type I and type III interferon induction in

response to RIG-I activation. Unexpectedly, these UFMylation-deficient mutations do not disrupt

the 14-3-3ε-RIG-I interaction. Instead, UFMylation of 14-3-3ε promotes signaling downstream of

RIG-I recruitment by stabilizing an interaction between 14-3-3ε and MAVS. In the absence of 14-

3-3ε UFMylation, the 14-3-3ε-MAVS interaction is weakened while RIG-I-MAVS binding is

paradoxically enhanced, suggesting that UFMylated 14-3-3ε coordinates optimal MAVS

signalosome assembly to enable robust interferon production. These findings reveal that

UFMylation of 14-3-3ε itself promotes signaling from the RIG-I-MAVS complex, highlighting the

broader role of UFMylation in antiviral innate immunity.

Results

UFM1 conjugation is required to promote 14-3-3ε-mediated enhancement of RIG-I
signaling to IFN-β. To define the molecular mechanism by which UFMylation to 14-3-3ε
specifically regulates RIG-I signaling, we first established the functional requirement for UFM1
and its conjugation activity in signaling to the IFN-β promoter. To test if UFM1 enhances RIG-I
signaling through 14-3-3ε, we performed IFN-β promoter luciferase assays in 293T cells following
ectopic expression of UFM1, 14-3-3ε, and RIG-I. We found that UFM1 ectopic expression
significantly enhanced RIG-I driven signaling to the IFN-β promoter. This signaling was further
amplified when UFM1 was co-expressed with both RIG-I and 14-3-3ε (Figure 1A). Next, to
determine whether UFM1-conjugation is required for this enhancement of RIG-I signaling, we
generated a conjugation-deficient UFM1 mutant by deleting three amino acids from its C-terminus
to prevent covalent attachment of UFM1 to target proteins (ΔC3) (22). Importantly, both WT and
conjugation-deficient UFM1 are expressed at similar levels in cells (Figure 1B). Using IFN-β
promoter luciferase assays in 293T cells, we found that unlike wild-type UFM1, the conjugation-
deficient UFM1 mutant failed to enhance 14-3-3ε-driven RIG-I signaling (Figure 1B). These
results demonstrate that UFM1 must be conjugation-competent to facilitate 14-3-3ε-mediated
enhancement of RIG-I signaling to the IFN-β promoter.

UFM1 conjugation to 14-3-3ε requires residues K50 and K215. Having established that active
UFM1 conjugation is required for 14-3-3ε to enhance RIG-I signaling, we next sought to verify our
previous published results that 14-3-3ε is directly UFMylated (21) and determine which specific
lysine residues serve as UFM1-conjugation sites. First, we verified 14-3-3ε UFMylation in cells.
For this, we developed a modified UFMylation assay based on previous studies (57). Because
UFMylation can be low stoichiometry and transient, overexpression of pathway components and
the use of a deconjugation-resistant UFM1 (G83A) are typically used to reveal substrate
modification. Thus, we co-expressed HA-tagged 14-3-3ε, Flag-UFM1 (WT, G83A, or ΔC3), and
the UFMylation ligase complex proteins Myc-UFL1 and Myc-UFBP1 in 293T-UFM1 KO cells (21).
Critically, lysates were boiled prior to immunoprecipitation to denature protein complexes,
ensuring that only proteins covalently linked to the immunoprecipitated protein would be detected
in subsequent analyses. Following anti-HA immunoprecipitation under denaturing conditions, we
identified a single apparent band with a shift of approximately 15 kDa, consistent with
predominantly a mono-UFMylation event on 14-3-3ε (Figure 2A). Importantly, the conjugation-
deficient Flag-UFM1-ΔC3 showed no attachment of 14-3-3ε, while a deconjugation-resistant

mutant, Flag-UFM1-G83A (22), demonstrated significantly increased levels of 14-3-3ε

UFMylation.

We next sought to identify the specific lysine residues on 14-3-3ε that undergo UFMylation

by using mass spectrometry analysis. We used Huh7 cells stably expressing Flag-UFM1 with a

two amino acid deletion (ΔC2). This mutant enhances detection of UFMylated proteins by

eliminating the prerequisite UFSP1/UFSP2-mediated cleavage of the last two amino acids in

UFM1 required to initiate UFM1 conjugation, (22). After immunoprecipitation of HA-14-3-3ε, mass

spectrometry analysis suggested that two lysine residues, K50 and K215, were covalently

modified by UFM1. Both K50 and K215 reside within clusters of basic residues within a positively

charged pocket of the conserved amphipathic ligand binding groove of 14-3-3 proteins (Figure

2B) (58). This groove mediates interactions with phosphorylated and non-phosphorylated client

proteins, suggesting that UFMylation at these sites could directly modulate 14-3-3ε protein-

protein interactions.

To confirm these specific sites as bona fide UFMylation targets, we employed both *in vitro*

and cellular UFMylation assays. Using an *in vitro* UFMylation assay with purified UFMylation

proteins (UBA5, UFC1, UFBP1/UFL1, and UFM1), we found that wild-type 14-3-3ε is mono-

UFMylated, whereas lysine-to-arginine (K to R) mutations at K50 and K215 (K50R/K215R)

abolished this mono-UFMylation (Figure 2C). Similarly, in the cellular UFMylation assay, the

K50R-K215R mutations significantly abrogated 14-3-3ε UFMylation (Figure 2D). These

complementary approaches confirm that K50 and K215 are necessary for detectable mono-

UFMylation of 14-3-3ε both *in vitro* and in cell-based assays. However, as we only detect mono-

UFMylation of 14-3-3ε, these results indicate that each individual 14-3-3ε molecule is UFMylated

at either K50 or K215, but not both simultaneously. This is consistent with the structural

positioning of these residues on opposite faces of the ligand-binding groove (58). We note that

the relative occupancy and functional contribution of each site remain to be determined and may

vary under different cellular conditions. Therefore, we used the K50R/K215R double point mutant

for further exploration of the biological impacts of 14-3-3ε UFMylation.

K50 and K215 are required for 14-3-3ε-mediated RIG-I activation of IFN. Having identified

K50 and K215 as the UFMylation sites on 14-3-3ε, we next investigated whether these residues

are functionally required for 14-3-3ε to enhance RIG-I-mediated signaling. We used two

complementary approaches to assess the impact of UFMylation-deficient 14-3-3ε on interferon

induction. First, we evaluated IFN-β promoter activation using a luciferase reporter assay in 293T-

RIG-I KO cells (59). We used these cells to establish a reconstitution system where RIG-I

signaling is specifically derived from co-expressed constructs, eliminating confounding effects

from endogenous RIG-I. Importantly, RIG-I ectopic expression can be sufficient to drive signaling.

We ectopically expressed 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$, along with RIG-I, and measured luciferase

activity (Figure 3A). While 14-3-3ε^{WT} significantly increased RIG-I activation of the IFN-β promoter,

as expected, 14-3-3ε$^{K50/215R}$ displayed a significantly impaired ability to enhance this signaling

(Figure 3A). This result suggests that UFMylation at K50 or K215 is critical for 14-3-3ε to promote

RIG-I-mediated transcriptional activation of the IFN-β promoter.

193   To validate these findings using a complementary approach, we measured endogenous

interferon transcript levels by RT-qPCR. We co-expressed RIG-I with 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$

in 293T-RIG-I KO cells and infected them with Sendai virus (SenV). SenV is an RNA virus that is

a potent activator of RIG-I signaling (60, 61). Importantly, SenV is sensed specifically by RIG-I,

ensuring our findings reflect RIG-I-driven signaling. Neither 14-3-3ε induced basal *IFNB1* and

IFNL1 expression without RIG-I. However, in the presence of RIG-I, 14-3-3ε^{WT} enhanced

interferon transcript levels while 14-3-3ε$^{K50/215R}$ did not (Figure 3B), confirming the luciferase

results. To validate these findings in another cell type, we measured the RNA levels of *IFNB1* in

A549-RIG-I KO cells, generated by CRISPR/Cas9, in response to SenV infection. Here, we co-

expressed RIG-I with 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$. Consistent with our earlier results, we found

that 14-3-3ε^{WT} enhances RIG-I-mediated induction of *IFNB1* in response to SenV, while 14-3-

3ε$^{K50/215R}$ does not (Figure 3C). Collectively, these findings demonstrate K50 and K215 are critical

for 14-3-3ε to promote transcriptional activation of type I and type III IFNs in response to RIG-I

activation, suggesting that UFMylation enhances RIG-I-mediated induction of IFN.

K50 and K215 are not required for 14-3-3ε interaction with RIG-I. Having established that K50

and K215 are required for 14-3-3ε to promote RIG-I signaling, we next investigated the

mechanism underlying this functional requirement. Since our previous findings revealed that both

UFM1 and UFL1 promote the interaction of RIG-I with 14-3-3ε (21), we hypothesized that

UFMylation of 14-3-3ε at K50/K215 would be required for its interaction with RIG-I during SenV-

mediated RIG-I activation. To test this hypothesis, we performed co-immunoprecipitation

experiments comparing the interaction between RIG-I and either 14-3-3ε^{WT} or the UFMylation-

deficient 14-3-3ε$^{K50/215R}$ mutant. Unexpectedly, we found that 14-3-3ε$^{K50/215R}$ maintained

interaction with RIG-I at levels comparable to the wild-type protein (Figure 4A-B), indicating that

UFMylation at these residues is not essential for the association of 14-3-3ε and RIG-I. To validate

our experimental approach, we included a control mutant of 14-3-3ε where key residues at the C-

terminal end (EDE to AAA) were altered, as these previously have been reported to mediate this

interaction (62). As expected, this control mutant showed significantly reduced RIG-I association,

confirming the work of others, as well as the specificity of the assay. These results revealed that

UFMylation at K50 or K215 are not critical for the interaction between 14-3-3ε and RIG-I. Further,

since 14-3-3ε$^{K50/215R}$ retains RIG-I binding comparable to wild-type, the signaling defects observed

in Figure 3 are unlikely to result from global disruptions to the 14-3-3ε ligand-binding groove.

UFMylation of 14-3-3ε differentially regulates MAVS-complex interactions. Since 14-3-3ε

facilitates the interaction between RIG-I and MAVS (6), we hypothesized that the UFMylation-

deficient 14-3-3ε$^{K50/215R}$ mutant may impair RIG-I-MAVS complex formation. To test this, we

performed co-immunoprecipitation assays during SenV infection using a MAVS construct

designed to capture CARD-dependent interactions, such as those mediated by the MAVS and

RIG-I N-terminal CARDs (63). As expected, overexpression of 14-3-3ε^{WT} enhanced the RIG-I and

MAVS interaction compared to the control (Figure 5A-5B). Surprisingly, 14-3-3ε$^{K50/215R}$ did not

reduce the association of RIG-I and MAVS, and instead it significantly enhanced this interaction

beyond 14-3-3ε^{WT} levels (Figure 5A-5B). This unexpected finding suggests that when 14-3-3ε

cannot be UFMylated, RIG-I associates more stably with MAVS but in a potentially non-productive

configuration. In contrast to its effect on RIG-I, the K50R/K215R mutation consistently reduced

14-3-3ε interaction with MAVS compared to wild-type (Figure 5A-5B). Together, these results

reveal that UFMylation of 14-3-3ε at K50/K215 is associated with MAVS signalosome assembly

in an unexpected way: it stabilizes 14-3-3ε-MAVS association while paradoxically dampening

RIG-I-MAVS interaction.

Discussion

Our study identifies UFMylation of 14-3-3ε as a novel regulatory mechanism that promotes RIG-

I-mediated innate immune signaling. Upon viral RNA recognition, activated RIG-I must translocate

from the cytosol to intracellular membranes to interact with MAVS within a specialized signaling

complex termed the "RIG-I translocon." This complex includes the K63 ubiquitin ligases TRIM25

and Riplet, as well as 14-3-3ε (6, 64). Through biochemical and functional analyses, we

demonstrated that UFM1 is covalently conjugated to either lysine residue K50 or K215 of 14-3-3ε

as a mono-modification. We further found that mutation of these lysine residues inactivates 14-3-

3ε UFMylation and limits the ability of 14-3-3ε to facilitate RIG-I-dependent IFN induction.

Surprisingly, we found these UFMylation-inactivating mutations did not affect the interaction

between 14-3-3ε and RIG-I. Furthermore, since the 14-3-3ε K50R/K215R mutant retains RIG-I

binding comparable to wild-type, the signaling defects we observed cannot be attributed to global

disruption of the 14-3-3ε ligand-binding groove. Instead, the K50R/K215R mutations reduced the
interaction of 14-3-3ε with MAVS. Our findings reveal that, in addition to facilitating RIG-I
translocation to MAVS (6), 14-3-3ε may also associate with MAVS itself, either directly or more
likely as part of the RIG-I-MAVS complex. UFMylation of 14-3-3ε stabilizes this 14-3-3ε-MAVS
association, as the UFMylation-deficient K50R/K215R mutant shows reduced MAVS complex
interaction. Intriguingly, loss of 14-3-3ε UFMylation paradoxically enhances RIG-I-MAVS
interaction while simultaneously reducing 14-3-3ε-MAVS binding and impairing downstream
signaling. This suggests that UFMylated 14-3-3ε coordinates signaling complex architecture
rather than solely promoting RIG-I-MAVS interaction. These data support a model by which
UFMylated 14-3-3ε functions beyond simply bringing RIG-I to MAVS; it likely coordinates the
proper architecture of the MAVS signalosome ensuring optimal configuration for downstream
signal propagation.

Our findings reveal an important distinction between global loss of UFMylation machinery
and site-specific inactivation of 14-3-3ε UFMylation. In our previous work, depletion of UFL1 or
UFM1 reduced both RIG-I-14-3-3ε and RIG-I-MAVS interactions (21), whereas the current study
shows that preventing 14-3-3ε UFMylation through K50R/K215R mutation maintains normal RIG-
I-14-3-3ε binding while paradoxically enhancing RIG-I-MAVS interaction. This apparent
discrepancy likely reflects the pleiotropic effects of global UFMylation loss versus targeted
disruption of a single substrate. The UFMylation machinery likely modifies multiple proteins within
the RIG-I signaling pathway, such as other MAVS-associated proteins. Thus, complete loss of
the UFMylation machinery would disrupt multiple protein-protein interactions simultaneously,
perhaps affecting earlier steps in RIG-I activation and translocation. In contrast, selective loss of
14-3-3ε UFMylation allows other UFMylation-dependent interactions to proceed normally,
permitting RIG-I-14-3-3ε complex formation and translocation to MAVS. However, without
UFMylated 14-3-3ε to properly coordinate the MAVS signalosome architecture, RIG-I interacts
more stably with MAVS. However, our results suggest that this enhanced interaction presumably
facilitates a non-productive configuration of the MAVS signalosome as it fails to support robust
downstream signaling. This model positions 14-3-3ε UFMylation as a critical quality control
checkpoint that ensures productive, rather than merely physical, RIG-I-MAVS complex assembly.

Previously, we demonstrated that UFL1 is recruited to ER-mitochondrial contact sites
following RIG-I activation (21, 56). Our current work extends this model by identifying 14-3-3ε as
one of the UFL1 target proteins that regulates RIG-I signaling. Interestingly, the UFMylated lysine
residues on 14-3-3ε are strategically located within the conserved amphipathic ligand-binding
groove, suggesting that UFMylation may introduce negative charges that could alter local

electrostatic interactions or induce conformational changes in this region (14). These
modifications likely influence how 14-3-3ε coordinates protein partners at MAVS-localized
membrane signaling sites. The observation that only mono-UFMylation occurs on 14-3-3ε aligns
with structural constraints within the protein: K50 and K215 reside on opposite faces of the ligand-
binding groove, which likely makes it structurally feasible for only one site to be UFMylated per
molecule (10). While the K50R/K215R mutations abolish UFMylation at these sites, we cannot
definitively exclude that these mutations might also disrupt other PTMs. Indeed, K50 in 14-3-3ε
and other 14-3-3 isoforms can undergo acetylation, SUMOylation, and ubiquitination (8, 65, 66),
suggesting that competing PTMs at the same residue could mediate context-dependent
functional switching between cellular pathways. The presence of two potential UFMylation sites
may provide redundancy, ensuring that 14-3-3ε can be UFMylated even if one lysine is occupied
by another PTM. Nevertheless, the strong correlation between loss of UFMylation and impaired
MAVS interaction, combined with our previous findings that UFL1/UFM1 depletion reduces RIG-
I-MAVS interaction (21), supports UFMylation as a critical modification that occurs on these
residues during RIG-I signaling. Our findings align with emerging evidence that 14-3-3 proteins
function as molecular chaperones that prevent client protein aggregation (14). UFMylation may
enhance these chaperone activities during RIG-I signaling by regulating MAVS signalosome
architecture and optimizing the molecular environment for assembly of a productive MAVS
signalosome.

Several important questions remain for future investigation. First, while the K50R/K215R
mutations abolish UFMylation, we cannot entirely exclude that these mutations might also prevent
other PTMs (65, 66). Definitive proof that UFMylation is sufficient to enhance MAVS engagement
would require complementary approaches, such as *in vitro* reconstitution interaction studies with
UFMylated 14-3-3ε. Second, the exact molecular function by which UFM1 conjugation on 14-3-
3ε promotes functional MAVS signalosome assembly remains to be fully elucidated. Our findings
suggest it may modulate interactions with MAVS-associated regulatory proteins or stabilize the
architecture of this multi-protein complex at membrane signaling sites (67). Future studies are
needed to identify the specific protein partners involved. Third, the molecular events and timing
that trigger site-specific UFMylation of 14-3-3ε following RIG-I activation warrant investigation, as
does determining whether other 14-3-3 isoforms undergo similar UFMylation-dependent
regulation. Finally, while overexpression studies were necessary for dissecting mechanism,
validation with endogenous proteins will be important to determine the physiological magnitude,
stoichiometry, and occupancy of 14-3-3ε UFMylation following RIG-I activation during viral
infection. Such studies await development of improved reagents, as currently available 14-3-3ε

antibodies have limited utility for endogenous co-immunoprecipitation studies. Despite these
open questions, our findings establish UFMylation of 14-3-3ε as a critical regulatory mechanism
in RIG-I signaling and position this modification system as an important node in innate immune
regulation.

The regulatory role of UFMylation in RIG-I signaling parallels its function in other immune
pathways. Similar to how UFMylation of PD-1 affects T cell activation (42, 44), modification of 14-
3-3ε demonstrates how this ubiquitin-like system provides context-dependent tuning of immune
responses. By identifying K50 and K215 on 14-3-3ε as critical UFMylation sites that promote RIG-
I signaling, our study reveals how this post-translational modification system tunes diverse innate
immune pathways. These findings have potential therapeutic implications, as the UFMylation
pathway represents a target for modulating antiviral innate immune responses in conditions
characterized by either insufficient immunity or excessive inflammation. Moreover, given that
multiple viruses have evolved mechanisms to manipulate UFMylation (54, 55), targeting this
pathway could potentially disrupt various aspects of viral infection. Understanding the molecular
details of UFMylation-dependent immune regulation, as demonstrated here for 14-3-3ε, may
ultimately guide development of selective immunomodulatory approaches.

Data Availability Statement
Plasmids, cell lines, and other reagents described in this study are available from the
corresponding author upon reasonable request.

Acknowledgements
We thank those colleagues who generously provided reagents, as indicated in the Methods, the
Duke Functional Genomics Core, the Duke Proteomics and Metabolomics Core Facility, and
members of the Horner Lab for valuable feedback and discussion.

Use of Generative AI: During manuscript preparation, the authors used Perplexity AI (Claude-
based model, October 2025 version) to assist with editing, improving clarity, and optimizing
sentence structure. All AI-generated suggestions were reviewed, revised, and approved by the
authors. The authors take full responsibility for the content of this publication.

Conflict of Interest: The authors declare no conflicts of interest relevant to this study.

This work was supported by National Institutes of Health grants R01AI155512 (S.M.H),
R35GM119544 (K.M.S), and T32CA00911 (HMS; IVG; DLS).

Figures

Figure 1. UFM1 conjugation is required to promote 14-3-3ε-mediated enhancement of RIG-

I signaling to IFN-β. (A) IFN-β promoter reporter luciferase expression (relative to *Renilla)* from

293T cells expressing Vector, Flag-UFM1^{WT}, Flag-RIG-I, or HA-14-3-3ε. (B) IFN-β promoter

reporter luciferase expression (relative to *Renilla*) from 293T cells expressing Vector, Flag-

UFM1^{WT}, Flag-UFM1$^{\Delta C3}$, Flag-RIG-I, or HA-14-3-3ε. Bars represent the mean ± SD, *n* = 3

independent replicates. *p ≤ 0.05, ** ≤ 0.01, ***p ≤ 0.001 determined by one way ANOVA followed

by Šidák's multiple comparisons test (A) or Tukey's multiple comparisons test (B).

Figure 2. UFM1 conjugation to 14-3-3ε requires residues K50 and K215 on 14-3-3ε. (A) UFMylation assay of anti-HA immunoprecipitated lysates from 293T UFM1 KO cells expressing Vector (V) or HA-14-3-3ε along with Myc-UFL1, Myc-UFBP1, and Flag-UFM1-WT, -G83A, or -ΔC3. (B) Ribbon representation of 14-3-3ε (PDB entry 3UAL), with amino acids K50 and K215 indicated in orange. (C) Purified UFMylation proteins (UFBP1, UFL1, UFC1, UBA5, and UFM1) as well as 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$ were incubated in UFMylation assay buffer in the presence or absence of ATP, followed by immunoblot analysis for 14-3-3ε (top) or using an antibody cocktail against all UFMylation proteins (bottom). (D) UFMylation assay of anti-HA immunoprecipitated lysates from 293T UFM1 KO cells expressing Vector (-), Flag-UFM1-WT, -G83A, or -ΔC3, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$. Representative blots from $n = 3$ independent replicates are shown (A, C, and D).

Figure 3. K50 and K215 are required for 14-3-3ε-mediated RIG-I activation of IFN. (A) IFN-β promoter reporter *Gaussia* luciferase expression from 293T-RIG-I KO cells expressing Vector (V), Flag-RIG-I, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$, with samples expressing RIG-I + V normalized to 100. (B) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from 293T-RIG-I KO cells transfected with Flag-RIG-I, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$ followed by SenV infection (6 h), with RIG-I + V samples normalized to 100. (C) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from A549-RIG-I KO cells transfected with low amounts of Flag-RIG-I, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$, followed by SenV infection (6 h), with RIG-I +V samples normalized to 100. *p ≤ 0.05, ** ≤ 0.01, ***p ≤ 0.005 determined by one way ANOVA followed by (A) Bonferroni or (B-C) Tukey's multiple comparisons test. Mean ± SD, *n* = 3 independent replicates are shown.

Figure 4. K50 and K215 are not required for 14-3-3ε interaction with RIG-I. (A) Immunoblot analysis of anti-HA immunoprecipitated extracts and inputs from 293T RIG-I KO cells expressing Flag-RIG-I, and HA-14-3-3ε WT, K50R/K215R, or AAA that were SenV-infected (4 h). A representative blot from $n = 3$ independent replicates is shown. (B) Relative quantification of the ratio of RIG-I (IP/relative input) to immunoprecipitated 14-3-3ε (from A) (Mean ± SD, $n = 3$). *$p \leq$ 0.05 or non-significant (ns) determined by one way ANOVA followed by Tukey's multiple comparisons test.

Figure 5. UFMylation of 14-3-3ε differentially regulates MAVS complex interactions. (A)
Immunoblot analysis of anti-Myc immunoprecipitated extracts and inputs from 293T RIG-I KO
cells expressing MAVS aa1-266-Myc, Flag-RIG-I, and HA-14-3-3ε that were SenV-infected (4 h).
A representative blot from $n = 3$ independent replicates is shown. (B) Relative quantification of
the ratio of RIG-I (IP/relative input) to immunoprecipitated MAVS. The dashed line indicates the
baseline level of RIG-I interaction with MAVS. (C) Relative quantification of the ratio of 14-3-3ε
(IP/relative input) to immunoprecipitated MAVS. For (B) and (C), the bars display the data spread,
with mean and individual data points shown from independent replicates; *p ≤ 0.05 by ratio paired
t test.

Methods

Cell lines, viruses, and transfection. 293T, 293T-UFM1 KO, 293T-RIG-I KO, and A549-RIG-I KO were grown in Dulbecco's modification of Eagle's medium (DMEM; Mediatech) supplemented with 10% fetal bovine serum, 1X minimum essential medium nonessential amino acids, and 25 mM of HEPES buffer. The 293T cells (CRL-3216) were obtained from American Type Culture Collection (ATCC). 293T-UFM1 KO and 293T-RIG-I KO were previously described (21, 59). A549-RIG-I KO were generated by using a purified ribonucleoprotein complex consisting of Cas9 protein and single-guide RNAs (sgRNAs) targeting RIG-I (Synthego). Cas9 protein and sgRNAs were mixed at a ratio of 1:6 and then added to 1×10^6 A549 cells in Neon Resuspension Buffer R, followed by electroporation using the Neon Transfection System (Invitrogen). Following recovery, single cell clones were isolated and validated by immunoblot. All cell lines were verified as mycoplasma-free using the Look-Out Mycoplasma PCR detection kit (Sigma-Aldrich). Sendai virus Cantell strain was obtained from Charles River Laboratories and used at 200 HAU/ml. Sendai virus infections were performed in serum-free media (30 to 60 min), after which complete medium was added. DNA transfections were performed using FuGene 6 (Promega). Cells were incubated for 24-48 hours post transfection prior to experimental treatment.

Plasmids. The following plasmids have been previously described: pIFN-β-luc (68), pCMV-Renilla (Promega), pcTak_Myc-5_UFL1 (21), pETDuet-His$_6$-TEV-UFL1-StrepII-3C-UFBP1$^{(21\text{-end})}$ (gift of Dr. Yogesh Kulathu) (29), pTRIPZ IFNB1^{2x}-Gluc (gift of Dr. Nandan Gokhale) (59), and pET-NT*-HRV3CP (gift of Dr. Gottfried Otting; Addgene #162795) (62). The following plasmids were generated by insertion of PCR-amplified fragments into BamHI-to-NotI digested pGEX-GST using InFusion cloning (Takara): pGEX-GST-14-3-3ε^{WT}, pGEX-GST-14-3-3ε$^{K50/215R}$, pGEX-GST-UBA5, pGEX-GST-UFC1, and pGEX-GST-UFM1. The following plasmids were generated by site-directed mutagenesis: pEF-TAK-HA-14-3-3ε$^{K50/215R}$, pEF-TAK-HA-14-3-3ε^{AAA} (E251-253A), and pEF-TAK-Flag-UFM1$^{\Delta C2}$. The following were cloned by InFusion: pEF-MAVS-1-261-Myc; pEF-TAK-Flag-UFM1^{WT}, pEF-TAK-Flag-UFM1$^{\Delta C3}$, and pEF-TAK-Flag-UFM1 G83A from pEF-TAK-HA-UFM1 (21); pEF-TAK-Flag-RIG-I from pEF-Bos-Flag-RIG-I (69), pEF-TAK-HA-14-3-3ε^{WT} from pEF-TAK-Flag-14–3-3ε (21), and pEF-TAK-Myc-UFBP1 by PCR from BC115617. Plasmid sequences were verified by DNA sequencing.

IFN-β promoter luciferase assay. IFN-β promoter luciferase assays were performed at 18 to 24 hours after treatment and transfection of pIFN-β-luc and pCMV-Renilla by using the Dual-Luciferase Reporter System (Promega). *Firefly* luciferase was normalized to *Renilla* luciferase,

whose expression is driven by the CMV promoter. For the *Gaussia* luciferase assays, 48 hours

post transfection of pTRIPZ IFNB1^{2x}-Gluc, supernatant was assayed for *Gaussia* luciferase using

the *Renilla* Luciferase Assay System (Promega). All samples were read using a BioTek Synergy2

microplate reader.

Immunoblotting. Cells were lysed with either RIPA (10 mM Tris HCl pH [7.5], 150 mM NaCl,

0.5% sodium deoxycholate, and 1% Triton X-100) or NP-40 Lysis Buffer (20 mM Tris HCl [pH

7.4], 100 mM NaCl, 0.5% NP-40) supplemented with protease inhibitor mixture (Sigma-Aldrich),

and Halt inhibitor phosphatase (Thermo Fisher Scientific), and lysates were isolated by

centrifugation. Quantified protein was resolved by SDS/PAGE and transferred to nitrocellulose

membranes. Membranes were stained with Revert 700 total protein stain (Li-COR Biosciences)

and then blocked with 3% bovine serum albumin in phosphate-buffered saline containing 0.01%

Tween-20 (PBS-T). Following incubation with primary antibodies, membranes were incubated

with species-specific horseradish-peroxidase (HRP)-conjugated antibodies (Jackson Immuno

Research 1:5000), followed by membrane treatment with Clarity Western ECL substrate (Bio-

Rad) and imaging on a LICOR Odyssey FC. The following antibodies were used for

immunoblotting: R-anti-SenV (MBL, 1:1000), HA-HRP (Sigma-Aldrich 1:1000-1:3000), Flag-HRP

(1:1000-1:5000), and M-anti-Myc (Cell Signaling Technology 1:1000).

Protein immunoprecipitation. Cells were lysed with either RIPA (10 mM Tris HCl [pH 7.5], 150

mM NaCl, 0.5% sodium deoxycholate, and 1% Triton X-100) or NP-40 Lysis Buffer (20 mM Tris

HCl [pH 7.4], 100 mM NaCl, 0.5% NP-40) supplemented with protease inhibitor mixture (Sigma-

Aldrich), and Halt phosphatase inhibitor, and 10% glycerol. Quantified protein (between 100-200

µg) was incubated anti-HA magnetic beads (Invitrogen) in lysis buffer either at room temperature

for 1-2 hours or 4°C overnight with head-over-tail rotation. Beads were washed 3X in lysis buffer

and eluted with 2X Laemmli buffer (Bio-Rad) with 5% 2-mercaptoethanol at 95°C for 5 min.

Proteins were resolved by SDS/PAGE and immunoblotting, as above. For UFMylation

precipitations, cells were lysed UFMylation lysis buffer (50 mM Tris [pH 8], 150 mM NaCl, 0.5%

sodium deoxycholate, and 1% NP-40) supplemented with 10% glycerol, 1:100 protease and

phosphatase inhibitors, as above, and 1:100 *N*-ethylmaleimide (10 mM final) (Sigma-Aldrich).

Post-nuclear lysates were boiled at 95°C for 10 min and incubated with Pierce anti-HA magnetic

beads with UFMylation lysis buffer. Beads were washed 3X in high salt buffer (10 mM Tris [pH 8],

1 mM EDTA, 500 mM NaCl) and eluted with 2X Laemmli buffer with 5% 2-mercaptoethanol at

95°C for 5 min.

Quantification of immunoblots. Immunoblots were imaged using the LICOR Odyssey FC system and quantified using ImageStudio software, and raw values were normalized to relevant controls for each antibody. RIG-I and 14-3-3ε interaction was quantified as ratio of the RIG-I levels in the IP relative to the input normalized to the 14-3-3ε pull down. RIG-I or 14-3-3ε interaction with MAVS was quantified as ratio the RIG-I or 14-3-3ε levels relative to the input normalized to the MAVS pull down. The RIG-I interaction with MAVS was normalized to the vector control (shown as a dashed line) to display the relative enhanced interaction in the presence of 14-3-3ε, while the 14-3-3ε interaction with MAVS shows raw values normalized to display the interaction differences between conditions.

Purification of GST/His-tagged proteins. pET28A-His$_6$-UBA5, pET28A-His$_6$-UFC1, pET28A-His$_6$-UFM1^{dSC}, pGEX-GST-14-3-3ε^{WT}, pGEX-GST-14-3-3ε$^{K50/215R}$, and pETDuet-His$_6$-TEV-UFL1-StrepII-3C-UFBP1$^{(21-end)}$ were transformed in *E. coli* Rosetta cells. Cells were cultured at 37°C until OD600 reached 0.6 to 0.8. Then, cells were induced with a final concentration of 1 mM IPTG. Cells were further cultured at 30°C for 3.5-4 hours. Cells were collected by centrifugation for 30 min at 7,000 rpm. The pellets were lysed in NET-N buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 0.5% IGEPAL, supplemented with Aprotinin, Leupeptin, PMSF, and dithiothreitol (DTT) at 1:100. Cells were sonicated 3 times for 30 sec at output 6-7 and a duty cycle of 80%. Then, lysates were centrifuged for 30 min at 12,000 rpm at 4°C. Supernatant was incubated with GST beads that were washed with NET-N buffer with rotation at 4°C for 1-2 hours. The beads were washed 3 times with NET-N buffer and 3 times with PBS supplemented with 1:100 DTT. After washes, GST tag was cleaved with purified human rhinovirus 3C protease (purified as described (62)) and protein was eluted with PBS by incubation with rotation at 4°C overnight.

***In vitro* UFMylation.** 2 µM of substrate, 0.25 µM of UBA5, 5 µM UFC1, 20 ng UFL1/UFBP1, 30 µM UFM1 were incubated in a reaction buffer containing 50 mM HEPES [pH 7.5], 10 mM MgCl$_2$, with or without 5 mM ATP for 1 hour at 37°C. The reaction was stopped by the addition of 2X SDS loading with 5% 2-mercaptoethanol and incubated at 95°C for 5 min. The reaction products were then separated by SDS/PAGE and analyzed by immunoblotting, as above.

RNA Analysis. Total cellular RNA was extracted using RNeasy Plus mini kit (Qiagen). RNA was then reverse transcribed using iScript complementary DNA (cDNA) synthesis kit (Bio-Rad), following manufacturer's instructions. The resulting DNA was diluted 1:3 in double distilled water.

RT-qPCR was performed in triplicate using the Power SYBR Green PCR master mix (Thermo

Fisher Scientific) and QuantStudio 6 Flex RT-PCR system. Oligonucleotide sequences are as

follows: IFNB1 F-5'-CTTTGCTATTTTCAGACAAGATTCA; R-5'GCCAGGAGGTTCTCAACAAT,

IFNL1 F-5'-CTTCCAAGCCCACCACAACT; R-5'-GGCCTCCAGGACCTTCAGC, and

GAPDH F-5'-AAGGTGAAGGTCGGAGTCAAC; R-5'-GGGGTCATTGATGGCAACAATA.

Sample Preparation and LC-MS/MS Analysis. Samples were spiked with undigested bovine

casein at a total of either 120 or 240 pmol as an internal quality control standard. Next, samples

were supplemented with 10 µL of 20% SDS, reduced with 10 mM DTT for 30 min at 80°C,

alkylated with 20 mM iodoacetamide for 30 min at room temperature, then supplemented with a

final concentration of 1.2% phosphoric acid and 546 µL of S-Trap (Protifi) binding buffer (90%

MeOH/100 mM TEAB). Proteins were trapped on the S-Trap micro cartridge, digested using 20

ng/µL sequencing grade trypsin (Promega) for 1 hour at 47°C, and eluted using 50 mM TEAB,

followed by 0.2% FA, and lastly using 50% ACN/0.2% FA. All samples were then lyophilized to

dryness. Samples were resolubilized using 48 µL of 1% TFA/2% ACN with 25 fmol/µL yeast ADH.

Quantitative LC/MS/MS was performed on 1.5 µL (3.13% of total sample) using an MClass UPLC

system (Waters Corp) coupled to an Orbitrap Fusion Lumos high resolution accurate mass

tandem mass spectrometer (Thermo Fisher Scientific) via a nanoelectrospray ionization source.

Briefly, the sample was first trapped on a Symmetry C18 20 mm × 180 µm trapping column (5

µl/min at 99.9/0.1 v/v water/acetonitrile), after which, the analytical separation was performed

using a 1.8 µm Acquity HSS T3 C18 75 µm × 250 mm column (Waters Corp.) with a 90-min linear

gradient of 5 to 30% acetonitrile with 0.1% formic acid at a flow rate of 400 nanoliters/min with a

column temperature of 55°C. Data collection on the Fusion Lumos mass spectrometer was

performed in a data-dependent acquisition (DDA) mode of acquisition with a r=120,000 (@ m/z

200) full MS scan from m/z 375 – 1500 with a target AGC value of 4e5 ions was performed.

MS/MS scans were acquired in the linear ion trap in "rapid" mode with a target AGC value of 1e5

and max fill time of 100 ms. The total cycle time 2s, with total cycle times of 2 sec between like

full MS scans. A 20s dynamic exclusion was employed to increase depth of coverage. The total

analysis cycle time for each sample injection was approximately 2 hours.

Proteomics Data Analysis. Following UPLC-MS/MS analyses, data were imported into

Proteome Discoverer 2.5 (Thermo Fisher Scientific). In addition to quantitative signal extraction,

the MS/MS data was searched against the SwissProt *H. sapiens* database (downloaded in Nov

2019) and a common contaminant/spiked protein database (bovine albumin, bovine casein, yeast

ADH, etc.), and an equal number of reversed-sequence "decoys" for false discovery rate
determination. Sequest (v 2.5, Thermo PD) was utilized to produce fragment ion spectra and to
perform the database searches. Database search parameters included fixed modification on Cys
(carbamidomethyl) and variable modification on Met (oxidation) and Lys (UFMylation). Search
tolerances were 2ppm precursor and 0.8Da product ion with semi-trypsin enzyme rules.
Discoverer were used to annotate the data at a maximum 1% protein false discovery rate based
on q-value calculations. Note that peptide homology was addressed using razor rules in which a
peptide matched to multiple different proteins was exclusively assigned to the protein has more
identified peptides. Protein homology was addressed by grouping proteins that had the same set
of peptides to account for their identification. A master protein within a group was assigned based
on % coverage.

Statistical analysis. We used ratio paired t test or one-way ANOVA followed by appropriate
multiple comparisons test using GraphPad Prism software. Graphed values are presented as
mean ± SD (n = 3) or as individual data points with spread; *$P \leq 0.05$, **$P \leq 0.01$, and ***$P \leq 0.001$.

References

1.   Rehwinkel J, Gack MU. 2020. RIG-I-like receptors: their regulation and roles in RNA
sensing. Nat Rev Immunol 20:537-551.

2.   Crow YJ, Stetson DB. 2022. The type I interferonopathies: 10 years on. Nat Rev
Immunol 22:471-483.

3.   Iwasaki A, Medzhitov R. 2015. Control of adaptive immunity by the innate immune
system. Nat Immunol 16:343-53.

4.   Taft J, Bogunovic D. 2018. The Goldilocks Zone of Type I IFNs: Lessons from Human
Genetics. J Immunol 201:3479-3485.

5.   Lin JP, Fan YK, Liu HM. 2019. The 14-3-3eta chaperone protein promotes antiviral
innate immunity via facilitating MDA5 oligomerization and intracellular redistribution.
PLoS Pathog 15:e1007582.

6.   Liu HM, Loo YM, Horner SM, Zornetzer GA, Katze MG, Gale M, Jr. 2012. The
mitochondrial targeting chaperone 14-3-3epsilon regulates a RIG-I translocon that
mediates membrane association and innate antiviral immunity. Cell Host Microbe
11:528-37.

7.   Aitken A. 2011. Post-translational modification of 14-3-3 isoforms and regulation of
cellular function. Semin Cell Dev Biol 22:673-80.

8.   Pennington KL, Chan TY, Torres MP, Andersen JL. 2018. The dynamic and stress-
adaptive signaling hub of 14-3-3: emerging mechanisms of regulation and context-
dependent protein-protein interactions. Oncogene 37:5587-5604.

9.   Henriksson ML, Francis MS, Peden A, Aili M, Stefansson K, Palmer R, Aitken A,
Hallberg B. 2002. A nonphosphorylated 14-3-3 binding motif on exoenzyme S that is
functional in vivo. Eur J Biochem 269:4921-9.

10.   Ottmann C, Yasmin L, Weyand M, Veesenmeyer JL, Diaz MH, Palmer RH, Francis MS,
Hauser AR, Wittinghofer A, Hallberg B. 2007. Phosphorylation-independent interaction
between 14-3-3 and exoenzyme S: from structure to pathogenesis. EMBO J 26:902-13.

11.   Liu J, Cao S, Ding G, Wang B, Li Y, Zhao Y, Shao Q, Feng J, Liu S, Qin L, Xiao Y. 2021.
The role of 14-3-3 proteins in cell signalling pathways and virus infection. J Cell Mol Med
25:4173-4182.

12.   Obsilova V, Obsil T. 2022. Structural insights into the functional roles of 14-3-3 proteins.
Front Mol Biosci 9:1016071.

13.   Toleman CA, Schumacher MA, Yu SH, Zeng W, Cox NJ, Smith TJ, Soderblom EJ,
Wands AM, Kohler JJ, Boyce M. 2018. Structural basis of O-GlcNAc recognition by
mammalian 14-3-3 proteins. Proc Natl Acad Sci U S A 115:5956-5961.

14.   Segal D, Maier S, Mastromarco GJ, Qian WW, Nabeel-Shah S, Lee H, Moore G,
Lacoste J, Larsen B, Lin ZY, Selvabaskaran A, Liu K, Smibert C, Zhang Z, Greenblatt J,

Peng J, Lee HO, Gingras AC, Taipale M. 2023. A central chaperone-like role for 14-3-3
proteins in human cells. Mol Cell 83:974-993 e15.

15.   Chan YK, Gack MU. 2016. A phosphomimetic-based mechanism of dengue virus to
antagonize innate immunity. Nat Immunol 17:523-30.

16.   Riedl W, Acharya D, Lee JH, Liu G, Serman T, Chiang C, Chan YK, Diamond MS, Gack
MU. 2019. Zika Virus NS3 Mimics a Cellular 14-3-3-Binding Motif to Antagonize RIG-I-
and MDA5-Mediated Innate Immunity. Cell Host Microbe 26:493-503 e6.

17.   Andrews DDT, Vlok M, Akbari Bani D, Hay BN, Mohamud Y, Foster LJ, Luo H, Overall
CM, Jan E. 2023. Cleavage of 14-3-3epsilon by the enteroviral 3C protease dampens
RIG-I-mediated antiviral signaling. J Virol 97:e0060423.

18.   Tam EH, Liu YC, Woung CH, Liu HM, Wu GH, Wu CC, Kuo RL. 2021. Role of the
Chaperone Protein 14-3-3epsilon in the Regulation of Influenza A Virus-Activated Beta
Interferon. J Virol 95:e0023121.

19.   Liu J, Qian C, Cao X. 2016. Post-Translational Modification Control of Innate Immunity.
Immunity 45:15-30.

20.   Zhou Y, He C, Wang L, Ge B. 2017. Post-translational regulation of antiviral innate
signaling. Eur J Immunol 47:1414-1426.

21.   Snider DL, Park M, Murphy KA, Beachboard DC, Horner SM. 2022. Signaling from the
RNA sensor RIG-I is regulated by ufmylation. Proc Natl Acad Sci U S A
119:e2119531119.

22.   Komatsu M, Chiba T, Tatsumi K, Iemura S, Tanida I, Okazaki N, Ueno T, Kominami E,
Natsume T, Tanaka K. 2004. A novel protein-conjugating system for Ufm1, a ubiquitin-
fold modifier. EMBO J 23:1977-86.

23.   Tatsumi K, Sou YS, Tada N, Nakamura E, Iemura S, Natsume T, Kang SH, Chung CH,
Kasahara M, Kominami E, Yamamoto M, Tanaka K, Komatsu M. 2010. A novel type of
E3 ligase for the Ufm1 conjugation system. J Biol Chem 285:5417-27.

24.   Komatsu M, Inada T, Noda NN. 2024. The UFM1 system: Working principles, cellular
functions, and pathophysiology. Mol Cell 84:156-169.

25.   Wei Y, Xu X. 2016. UFMylation: A Unique & Fashionable Modification for Life. Genomics
Proteomics Bioinformatics 14:140-146.

26.   Banerjee S, Kumar M, Wiener R. 2020. Decrypting UFMylation: How Proteins Are
Modified with UFM1. Biomolecules 10.

27.   Millrine D, Peter JJ, Kulathu Y. 2023. A guide to UFMylation, an emerging
posttranslational modification. FEBS J 290:5040-5056.

28.   Zhou X, Mahdizadeh SJ, Le Gallo M, Eriksson LA, Chevet E, Lafont E. 2024.
UFMylation: a ubiquitin-like modification. Trends Biochem Sci 49:52-67.

29. Peter JJ, Magnussen HM, DaRosa PA, Millrine D, Matthews SP, Lamoliatte F,
Sundaramoorthy R, Kopito RR, Kulathu Y. 2022. A non-canonical scaffold-type E3
ligase complex mediates protein UFMylation. EMBO J 41:e111015.

30. Zhang Y, Yan M, Yu Y, Wang J, Jiao Y, Zheng M, Zhang S. 2024. 14-3-3epsilon: a
protein with complex physiology function but promising therapeutic potential in cancer.
Cell Commun Signal 22:72.

31. Jiang Q, Wang Y, Xiang M, Hua J, Zhou T, Chen F, Lv X, Huang J, Cai Y. 2023. UFL1,
a UFMylation E3 ligase, plays a crucial role in multiple cellular stress responses. Front
Endocrinol (Lausanne) 14:1123124.

32. Liu J, Guan D, Dong M, Yang J, Wei H, Liang Q, Song L, Xu L, Bai J, Liu C, Mao J,
Zhang Q, Zhou J, Wu X, Wang M, Cong YS. 2020. UFMylation maintains tumour
suppressor p53 stability by antagonizing its ubiquitination. Nat Cell Biol 22:1056-1063.

33. Qin B, Yu J, Nowsheen S, Wang M, Tu X, Liu T, Li H, Wang L, Lou Z. 2019. UFL1
promotes histone H4 ufmylation and ATM activation. Nat Commun 10:1242.

34. Liang Z, Ning R, Wang Z, Kong X, Yan Y, Cai Y, He Z, Liu XG, Zou Y, Zhou J. 2024.
The emerging roles of UFMylation in the modulation of immune responses. Clin Transl
Med 14:e70019.

35. Qin B, Yu J, Nowsheen S, Zhao F, Wang L, Lou Z. 2020. STK38 promotes ATM
activation by acting as a reader of histone H4 ufmylation. Sci Adv 6:eaax8214.

36. Ishimura R, El-Gowily AH, Noshiro D, Komatsu-Hirota S, Ono Y, Shindo M, Hatta T, Abe
M, Uemura T, Lee-Okada HC, Mohamed TM, Yokomizo T, Ueno T, Sakimura K,
Natsume T, Sorimachi H, Inada T, Waguri S, Noda NN, Komatsu M. 2022. The UFM1
system regulates ER-phagy through the ufmylation of CYB5R3. Nat Commun 13:7857.

37. Liang JR, Lingeman E, Luong T, Ahmed S, Muhar M, Nguyen T, Olzmann JA, Corn JE.
2020. A Genome-wide ER-phagy Screen Highlights Key Roles of Mitochondrial
Metabolism and ER-Resident UFMylation. Cell 180:1160-1177 e20.

38. Scavone F, Gumbin SC, Da Rosa PA, Kopito RR. 2023. RPL26/uL24 UFMylation is
essential for ribosome-associated quality control at the endoplasmic reticulum. Proc Natl
Acad Sci U S A 120:e2220340120.

39. Walczak CP, Leto DE, Zhang L, Riepe C, Muller RY, DaRosa PA, Ingolia NT, Elias JE,
Kopito RR. 2019. Ribosomal protein RPL26 is the principal target of UFMylation. Proc
Natl Acad Sci U S A 116:1299-1308.

40. Wang L, Xu Y, Rogers H, Saidi L, Noguchi CT, Li H, Yewdell JW, Guydosh NR, Ye Y.
2020. UFMylation of RPL26 links translocation-associated quality control to endoplasmic
reticulum protein homeostasis. Cell Res 30:5-20.

41. Xu X, Huang W, Bryant CN, Dong Z, Li H, Wu G. 2024. The ufmylation cascade controls
COPII recruitment, anterograde transport, and sorting of nascent GPCRs at ER. Sci Adv
10:eadm9216.

42.   He C, Xing X, Chen HY, Gao M, Shi J, Xiang B, Xiao X, Sun Y, Yu H, Xu G, Yao Y, Xie
Z, Xing Y, Budiarto BR, Chen SY, Gao Y, Lee YR, Zhang J. 2024. UFL1 ablation in T
cells suppresses PD-1 UFMylation to enhance anti-tumor immunity. Mol Cell 84:1120-
1138 e8.

43.   Yoo HM, Kang SH, Kim JY, Lee JE, Seong MW, Lee SW, Ka SH, Sou YS, Komatsu M,
Tanaka K, Lee ST, Noh DY, Baek SH, Jeon YJ, Chung CH. 2014. Modification of ASC1
by UFM1 is crucial for ERalpha transactivation and breast cancer development. Mol Cell
56:261-274.

44.   Zhou J, Ma X, He X, Chen B, Yuan J, Jin Z, Li L, Wang Z, Xiao Q, Cai Y, Zou Y, Cong
YS. 2023. Dysregulation of PD-L1 by UFMylation imparts tumor immune evasion and
identified as a potential therapeutic target. Proc Natl Acad Sci U S A 120:e2215732120.

45.   Lee L, Perez Oliva AB, Martinez-Balsalobre E, Churikov D, Peter J, Rahmouni D, Audoly
G, Azzoni V, Audebert S, Camoin L, Mulero V, Cayuela ML, Kulathu Y, Geli V, Lachaud
C. 2021. UFMylation of MRE11 is essential for telomere length maintenance and
hematopoietic stem cell survival. Sci Adv 7:eabc7371.

46.   Wang Z, Gong Y, Peng B, Shi R, Fan D, Zhao H, Zhu M, Zhang H, Lou Z, Zhou J, Zhu
WG, Cong YS, Xu X. 2019. MRE11 UFMylation promotes ATM activation. Nucleic Acids
Res 47:4124-4135.

47.   Jing J, Yang F, Wang K, Cui M, Kong N, Wang S, Qiao X, Kong F, Zhao D, Ji J, Tang L,
Gao J, Cong YS, Ding D, Chen K. 2025. UFMylation of NLRP3 Prevents Its Autophagic
Degradation and Facilitates Inflammasome Activation. Adv Sci (Weinh)
doi:10.1002/advs.202406786:e2406786.

48.   Wang X, Li C, Wang Y, Li L, Han Z, Wang G. 2020. UFL1 Alleviates LPS-Induced
Apoptosis by Regulating the NF-kappaB Signaling Pathway in Bovine Ovarian
Granulosa Cells. Biomolecules 10.

49.   Cai Y, Zhu G, Liu S, Pan Z, Quintero M, Poole CJ, Lu C, Zhu H, Islam B, Riggelen JV,
Browning D, Liu K, Blumberg R, Singh N, Li H. 2019. Indispensable role of the Ubiquitin-
fold modifier 1-specific E3 ligase in maintaining intestinal homeostasis and controlling
gut inflammation. Cell Discov 5:7.

50.   Li C, Li L, Chen K, Wang Y, Yang F, Wang G. 2019. UFL1 Alleviates
Lipopolysaccharide-Induced Cell Damage and Inflammation via Regulation of the
TLR4/NF-kappaB Pathway in Bovine Mammary Epithelial Cells. Oxid Med Cell Longev
2019:6505373.

51.   Balce DR, Wang YT, McAllaster MR, Dunlap BF, Orvedahl A, Hykes BL, Jr., Droit L,
Handley SA, Wilen CB, Doench JG, Orchard RC, Stallings CL, Virgin HW. 2021.
UFMylation inhibits the proinflammatory capacity of interferon-gamma-activated
macrophages. Proc Natl Acad Sci U S A 118.

52.   Tao Y, Yin S, Liu Y, Li C, Chen Y, Han D, Huang J, Xu S, Zou Z, Yu Y. 2023. UFL1
promotes antiviral immune response by maintaining STING stability independent of
UFMylation. Cell Death Differ 30:16-26.

53. Kulsuptrakul J, Wang R, Meyers NL, Ott M, Puschnik AS. 2021. A genome-wide
CRISPR screen identifies UFMylation and TRAMP-like complexes as host factors
required for hepatitis A virus infection. Cell Rep 34:108859.

54. Schmidt HM, Sorensen GC, Lanahan MR, Bland KM, Aufgebauer CJ, Park M,
Grabowski J, Horner SM. 2025. UFMylation promotes orthoflavivirus infectious particle
production. J Virol 99:e0065425.

55. Yiu SPT, Zerbe C, Vanderwall D, Huttlin EL, Weekes MP, Gewurz BE. 2023. An
Epstein-Barr virus protein interaction map reveals NLRP3 inflammasome evasion via
MAVS UFMylation. Mol Cell 83:2367-2386 e15.

56. Horner SM, Wilkins C, Badil S, Iskarpatyoti J, Gale M, Jr. 2015. Proteomic analysis of
mitochondrial-associated ER membranes (MAM) during RNA virus infection reveals
dynamic changes in protein and organelle trafficking. PLoS One 10:e0117963.

57. Zhou J, Liang Q, Dong M, Ma X, Jin Y, Guan D, Liu J, Wang M, Cong YS. 2022.
Optimized protocol to detect protein UFMylation in cells and in vitro via immunoblotting.
STAR Protoc 3:101074.

58. Yang X, Lee WH, Sobott F, Papagrigoriou E, Robinson CV, Grossmann JG, Sundstrom
M, Doyle DA, Elkins JM. 2006. Structural basis for protein-protein interactions in the 14-
3-3 protein family. Proc Natl Acad Sci U S A 103:17237-42.

59. Gokhale NS, Sam RK, Somfleth K, Thompson MG, Marciniak DM, Smith JR, Genoyer E,
Eggenberger J, Chu LH, Park M, Dvorkin S, Oberst A, Horner SM, Ong SE, Gale M, Jr.,
Savan R. 2024. Cellular RNA interacts with MAVS to promote antiviral signaling.
Science 386:eadl0429.

60. Cantell K, Hirvonen S, Kauppinen HL, Myllyla G. 1981. Production of interferon in
human leukocytes from normal donors with the use of Sendai virus. Methods Enzymol
78:29-38.

61. Sumpter R, Jr., Loo YM, Foy E, Li K, Yoneyama M, Fujita T, Lemon SM, Gale M, Jr.
2005. Regulating intracellular antiviral defense and permissiveness to hepatitis C virus
RNA replication through a cellular RNA helicase, RIG-I. J Virol 79:2689-99.

62. Abdelkader EH, Otting G. 2021. NT*-HRV3CP: An optimized construct of human
rhinovirus 14 3C protease for high-yield expression and fast affinity-tag cleavage. J
Biotechnol 325:145-151.

63. Gack MU, Shin YC, Joo CH, Urano T, Liang C, Sun L, Takeuchi O, Akira S, Chen Z,
Inoue S, Jung JU. 2007. TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-
mediated antiviral activity. Nature 446:916-920.

64. Oshiumi H, Matsumoto M, Hatakeyama S, Seya T. 2009. Riplet/RNF135, a RING finger
protein, ubiquitinates RIG-I to promote interferon-beta induction during the early phase
of viral infection. J Biol Chem 284:807-17.

65.   Fan K, Zhu K, Wang J, Ni X, Shen S, Gong Z, Cheng X, Zhang C, Liu H, Suo T, Ni X,
794   Liu H. 2022. Inhibition of 14-3-3epsilon by K50 acetylation activates YAP1 to promote
795   cholangiocarcinoma growth. Exp Cell Res 421:113404.

796   66.   Hendriks IA, Lyon D, Young C, Jensen LJ, Vertegaal AC, Nielsen ML. 2017. Site-
797   specific mapping of the human SUMO proteome reveals co-modification with
798   phosphorylation. Nat Struct Mol Biol 24:325-336.

799   67.   Vazquez C, Horner SM. 2015. MAVS Coordination of Antiviral Innate Immunity. J Virol
800   89:6974-7.

801   68.   Fredericksen B, Akkaraju GR, Foy E, Wang C, Pflugheber J, Chen ZJ, Gale M, Jr. 2002.
802   Activation of the interferon-beta promoter during hepatitis C virus RNA replication. Viral
803   Immunol 15:29-40.

804   69.   Saito T, Hirai R, Loo YM, Owen D, Johnson CL, Sinha SC, Akira S, Fujita T, Gale M, Jr.
805   2007. Regulation of innate antiviral defenses through a shared repressor domain in RIG-
806   I and LGP2. Proc Natl Acad Sci U S A 104:582-7.
807