14-3-3ε UFMylation promotes RIG-I-mediated signaling

Isaura Vanessa Gutierrez[1], Moonhee Park[1], Lauren Sar[1], Ryan Rodriguez[2], Daltry L. Snider[2], Gabriella Torres[1], K. Matthew Scaglione[2,3], Stacy M. Horner[1,2,4]

[1] Department of Integrative Immunobiology, Duke University School of Medicine, Durham, NC, USA

[2] Department of Molecular Genetics and Microbiology, Duke University School of Medicine, Durham, NC, USA

[3] Duke Center for Neurodegeneration and Neurotherapeutics, Duke University School of Medicine, Durham, NC, USA

[4] Department of Medicine, Duke University School of Medicine, Durham, NC, USA

Correspondence: stacy.horner@duke.edu

Stacy M. Horner, Ph.D.

Duke University School of Medicine

213 Research Dr., Box 3053 DUMC

Durham, NC USA 27710

Phone: 919-684-1921

Abstract

Post-translational modifications are critical for regulating the RIG-I signaling pathway. Previously, we identified a role for the post-translation modification UFM1 (UFMylation) in promoting RIG-I signaling by stimulating the interaction between RIG-I and its membrane-targeting protein 14-3-3ε. Here, we identify UFMylation of 14-3-3ε as a novel regulatory mechanism promoting RIG-I signaling. We demonstrate that UFM1 conjugation to lysine residue K50 or K215 results in mono-UFMylation on 14-3-3ε and enhances its ability to promote RIG-I signaling. Importantly, we show that mutation of these residues (K50R/K215R) abolishes UFMylation and impairs induction of type I and III interferons without disrupting the interaction between 14-3-3ε and RIG-I. This suggests that UFMylation of 14-3-3ε likely stabilizes signaling events downstream of RIG-I activation to promote induction of interferon. Collectively, our work suggests that UFMylation-driven activation of 14-3-3ε facilitates innate immune signaling and highlights the broader role of UFMylation for antiviral defense and immune regulation.

Importance

Post-translational modifications provide regulatory control of antiviral innate immune responses. Our study reveals that UFMylation of 14-3-3ε is a control point for RIG-I-mediated antiviral signaling. We demonstrate that conjugation of UFM1 to specific lysine residues on 14-3-3ε enhances downstream signaling events that facilitate interferon induction, but surprisingly it does not affect 14-3-3ε binding to RIG-I. By identifying the precise sites of UFMylation on 14-3-3ε and their functional consequences, we provide insights into the regulatory layers governing antiviral innate immunity. These findings complement emerging evidence that UFMylation serves as a versatile modulator across diverse immune pathways. Furthermore, our work highlights how protein chaperones like 14-3-3ε can be dynamically modified to orchestrate complex signaling cascades, suggesting potential therapeutic approaches for targeting dysregulated innate immunity.

Introduction

The antiviral innate immune response is tightly regulated to ensure effective defense against viral infections while preventing excessive inflammation. This antiviral response to RNA viruses is activated by cytosolic RIG-I-like receptors (RLRs) RIG-I and MDA5, which detect pathogen-associated molecular patterns from RNA viruses and initiate the formation of a MAVS-signaling complex (1). This complex activates a downstream signaling cascade that leads to the transcriptional activation of type I and type III interferons (IFN) and the production of IFN-stimulated genes (ISGs) that limit viral replication. Robust innate immune signaling is essential for effective adaptive immunity, but dysregulation of these pathways can result in autoimmune diseases and interferonopathies (2-4). Therefore, the induction of IFN is carefully regulated by mechanisms that temporarily activate, dampen, and ultimately shut down the signaling pathway.

Both 14-3-3ε and 14-3-3η are examples of RLR regulatory proteins that regulate IFN induction. These proteins facilitate the translocation of RIG-I and MDA5, respectively, from the cytosol to intracellular membrane signaling sites to interact with MAVS (5, 6). 14-3-3 proteins are highly conserved regulatory molecules involved in diverse cellular processes, including signal transduction, apoptosis, and stress responses. Typically, they regulate cellular processes by binding to phosphorylated client proteins, although they are also known to bind to non-phosphorylated client proteins, including those containing other post-translational modifications (PTMs) beyond phosphorylation (7-13). Recent work has also suggested that 14-3-3 proteins can serve as protein chaperones, preventing aggregation and phase separation of their client proteins (14). Orthoflavivirus NS3 proteins specifically target and inactivate both 14-3-3ε and 14-3-3η to prevent RLR signaling, and the enterovirus 3C protease cleaves 14-3-3ε, underscoring the essential role of these membrane-targeting proteins in antiviral innate immunity (15-17).

Post-translational modifications play a crucial role in regulating RIG-I signaling (18, 19). Recent studies by our group have shown that ubiquitin-fold modifier 1 (UFM1) and the proteins involved in UFM1 conjugation promote RIG-I signaling (20). UFMylation involves the covalent conjugation of ubiquitin-fold modifier 1 (UFM1), an 85-amino acid peptide, to lysine residues on proteins, utilizing an E1 activating enzyme (UBA5), an E2 conjugating enzyme (UFC1), and an E3 scaffold-like ligase complex (UFL1 and the membrane-anchored protein UFBP1) (21-29). UFM1 is a non-degradative mark, and its addition to proteins can alter their interactions. UFMylation modulates various cellular processes, including the DNA damage response and genomic instability, various aspects of ER homeostasis and translation, ER-Golgi transport, viral infection, cancer progression, and both adaptive and innate immune responses (23, 24, 26, 27, 30-43). For example, UFMylation of the ribosomal protein RPL26 enhances hepatitis A virus

translation (44). In orthoflaviviruses, UFMylation promotes the production of infectious virions
(45). Interestingly, viruses have evolved mechanisms to manipulate host UFMylation pathways
by targeting UFL1 to specific substrates. For example, orthoflaviviruses co-opt UFL1 to aid in viral
assembly (45), while Epstein-Barr virus (EBV) targets UFL1 to MAVS for UFMylation to dampen
inflammasome activation (46). Others have shown that UFMylation promotes NLRP3 activation
or that it can limit LPS-mediated inflammation via its roles in regulating ER homeostasis and the
unfolded protein response (47-51). UFL1 can also bind STING during HSV-1 infection to maintain
IFN induction (52). Thus, UFL1 or UFMylation can influence viral infection and innate immune
responses in diverse ways.

Our previous work on UFMylation in the context of RIG-I signaling revealed that UFL1
translocates to intracellular membranes upon RIG-I activation (20, 53). We identified roles for
UFL1 and UFM1 in facilitating RIG-I interaction with 14-3-3ε and found that 14-3-3ε exhibits
increased UFMylation in response to RIG-I activation. However, several critical questions
remained unanswered. First, does UFMylation directly modulate 14-3-3ε function during RIG-I
signaling? Second, which specific lysine residues on 14-3-3ε serve as UFMylation sites? Third,
how does UFMylation functionally impact 14-3-3ε's role in RIG-I-mediated signaling? We
hypothesized that direct UFMylation of 14-3-3ε at specific lysine residues enhances its ability to
promote RIG-I signaling. Here, we demonstrate that UFM1 conjugation enhances 14-3-3ε-
mediated RIG-I signaling. We then identify lysine residues on 14-3-3ε that are UFMylated. Then,
we show how UFMylation of these lysine residues regulates RIG-I-dependent IFN induction

Results
UFM1 conjugation is required to promote 14-3-3ε-mediated enhancement of RIG-I
signaling to IFN-β. To determine whether UFM1 plays a functional role in enhancing RIG-I
signaling through 14-3-3ε, we first assessed its effect on IFN-β promoter activation**.** Using IFN-β
promoter luciferase assays in 293T cells, we found that ectopic expression of UFM1 significantly
enhanced RIG-I driven signaling to IFN-β. This signaling was further amplified when UFM1 was
co-expressed with both RIG-I and 14-3-3ε, suggesting these three proteins cooperate for
signaling regulation (Figure 1A). Next, to determine whether the conjugation of UFM1 is
specifically required for this enhancement, we generated a conjugation-deficient UFM1 mutant
(ΔC3) by deleting three amino acids from its C-terminus to prevent covalent attachment of UFM1
to target proteins (21). Using IFN-β promoter luciferase assays in 293T cells, we found that unlike
wild-type UFM1, the conjugation-deficient UFM1 mutant failed to enhance 14-3-3ε-driven RIG-I

signaling (Figure 1B). These results demonstrate that UFM1 must be covalently conjugated to

proteins to facilitate 14-3-3ε-mediated enhancement of RIG-I signaling to the IFN-β promoter.

UFM1 conjugation to 14-3-3ε requires residues K50 and K215 on 14-3-3ε. Having established

that UFM1 conjugation is required for 14-3-3ε-mediated enhancement of RIG-I signaling, we next

sought to verify our previous results that 14-3-3ε is directly UFMylated (20) and determine which

specific lysine residues serve as UFM1-conjugation sites. First, we wanted to verify 14-3-3ε

UFMylation in cells. For this, we developed a modified UFMylation assay based on previous

studies (54). In this assay, we co-expressed HA-tagged 14-3-3ε, Flag-UFM1, and the UFMylation

ligase complex proteins UFL1 and UFBP1 in 293T-UFM1 KO cells generated by CRISPR/Cas9

targeting. Critically, lysates were boiled prior to immunoprecipitation to denature protein

complexes, ensuring that only proteins covalently linked to the immunoprecipitated protein would

be detected in subsequent analyses. Following HA-immunoprecipitation, we identified a single

Flag-UFM1 conjugation to 14-3-3ε, consistent with a mono-UFMylation event (Figure 2A).

Importantly, the conjugation-deficient Flag-UFM1-ΔC3 showed no attachment of 14-3-3ε, while a

deconjugation-resistant Flag-UFM1-G83A mutant (21) demonstrated significantly increased

levels of 14-3-3ε UFMylation.

133   Having established that UFM1 conjugation is required for RIG-I signaling, we next sought

to identify the specific lysine residues on 14-3-3ε that undergo UFMylation by using mass

spectrometry analysis. We used Huh7 cells stably expressing Flag-UFM1 with a two amino acid

deletion (ΔC2) that enhances detection of UFMylated proteins by eliminating the prerequisite of

G-SC cleavage in UFM1 by the UFSP1/UFSP2 proteins that needs to occur to initiate the

UFMylation pathway (21). After HA-immunoprecipitation of HA-tagged 14-3-3ε, mass

spectrometry analysis identified three peptides containing two lysine residues, K50 and K215,

that were covalently modified by UFM1. Both K50 and K215 reside within clusters of basic

residues within a positively charged pocket of the conserved amphipathic ligand binding groove

of 14-3-3 proteins.

143   To confirm these specific sites as bona fide UFMylation targets, we employed both *in vitro*

and cellular UFMylation assays. Using an *in vitro* UFMylation assay with purified UFMylation

proteins, we demonstrated that wild-type 14-3-3ε has mono-UFMylation, whereas lysine-to-

arginine mutations at K50 and K215 (K50R/K215R) abolished this mono-UFMylation (Figure 2B).

Similarly, in our cellular UFMylation assay, the K50R-K215R mutations significantly abrogated

14-3-3ε mono-UFMylation (Figure 2C). These complementary approaches confirm that K50 and

K215 are the principal sites for UFM1 conjugation to 14-3-3ε. However, as we only detect mono-

UFMylation of 14-3-3ε, these results also suggest that either K50 or K215 undergo UFM1-
conjugation in a molecule of 14-3-3ε.

K50 and K215 are not required for 14-3-3ε interaction with RIG-I. Based on our previous
findings that both UFM1 and UFL1 are important for the interaction of RIG-I with 14-3-3ε (20), we
hypothesized that UFMylation of 14-3-3ε at either K50 or K215 would be required for its interaction
with RIG-I. To test this hypothesis, we performed co-immunoprecipitation experiments comparing
RIG-I binding between 14-3-3ε^{WT} or the UFMylation-deficient 14-3-3ε$^{K50/215R}$ mutant.
Unexpectedly, we found that 14-3-3ε$^{K50/215R}$ maintained interaction with RIG-I at levels
comparable to the wild-type protein (Figure 3A-B), indicating that UFMylation at these residues is
not essential for the association of 14-3-3ε and RIG-I. To validate our experimental approach, we
included a control mutant of 14-3-3ε where key residues in the C-terminal end (EDE to AAA) were
altered, as these previously have been reported to mediate this interaction (55). As expected, this
control mutant showed significantly reduced RIG-I binding, confirming the work of others and the
specificity of our assay. These results revealed that UFMylation at K50 or K215 are not critical for
the interaction between 14-3-3ε and RIG-I.

K50 and K215 are required for 14-3-3ε-mediated RIG-I activation of IFN. Since UFMylation
did not affect the binding of for 14-3-3ε to RIG-I, we next investigated whether the K50/K215
residues nevertheless impact the ability of 14-3-3ε to enhance RIG-I-mediated signaling using
two complementary approaches. First, we evaluated IFN-β promoter activation using a luciferase
reporter assay in 293T-RIG-I KO cells. We ectopically expressed 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$,
along with RIG-I, and measured luciferase activity. While 14-3-3ε^{WT} significantly increased RIG-I
signaling to the IFN-β promoter as expected, 14-3-3ε$^{K50/215R}$ showed significantly impaired ability
to enhance this signaling (Figure 4A). This result suggests that UFMylation at K50 or K215 is
critical for 14-3-3ε to promote RIG-I-mediated transcriptional activation. To directly confirm this
finding at the transcriptional level, we measured the induction of endogenous IFN genes by RT-
qPCR in response to infection by Sendai virus, a potent activator of RIG-I signaling (56, 57). We
ectopically expressed 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$, along with RIG-I in 293T-RIG-I KO cells and
analyzed the levels of *IFNB1* and *IFNL1* transcripts. Consistent with our luciferase assay results,
14-3-3ε$^{K50/215R}$ significantly limited RIG-I-mediated induction of both *IFNB1* and *IFNL1* compared
to 14-3-3ε^{WT} (Figure 4B-C). Collectively, these findings demonstrate that while lysine residues
required for 14-3-3ε UFMylation do not affect RIG-I binding, they are nevertheless critical for
promoting RIG-I-mediated transcriptional activation of type I and type III IFNs, suggesting that

UFMylation of 14-3-3ε at K50 and K215 enhances downstream signaling events following initial
RIG-I binding.
186
Discussion.
Our study identifies UFMylation of 14-3-3ε as a novel regulatory mechanism that promotes
RIG-I-mediated antiviral signaling. Through a series of biochemical and functional analyses, we
have demonstrated that UFM1 is covalently conjugated to lysine residues K50 or K215 on 14-3-
3ε, and that this modification is critical for 14-3-3ε to enhance RIG-I-dependent IFN induction.
Upon viral RNA recognition, activated RIG-I must translocate from the cytosol to intracellular
membranes to interact with MAVS within a specialized signaling complex termed the "RIG-I
translocon," which includes the K63 ubiquitin ligases TRIM25 and Riplet. One of our most
intriguing findings was that mutation of the UFMylation sites (K50R/K215R) abolished the ability
of 14-3-3ε to enhance RIG-I signaling without disrupting their physical interaction. This
unexpected result suggests that UFMylation modulates 14-3-3ε function through mechanisms
beyond simple binding affinity regulation.
Previously, we demonstrated that the UFMylation E3 ligase complex member UFL1 is
recruited to ER-mitochondrial contact sites following RIG-I activation (20, 53). Our current work
extends this model by identifying UFMylation sites on 14-3-3ε (K50 and K215) that are critical for
its function in this pathway. The mechanism by which UFMylation enhances 14-3-3ε function may
involve electrostatic or conformational changes within its ligand-binding groove as the negative
charge of UFM1 in the presence of these positively charged residues is likely to influence how
14-3-3ε interacts with and coordinates protein partners at membrane signaling sites (14).
Interestingly, our results suggest that only mono-UFMylation occurs on 14-3-3ε. As K50 and K215
can reside on opposite faces within the conserved amphipathic ligand-binding groove of 14-3-3ε
when 14-3-3ε binds non-phosphorylated proteins, it seems possible that only one site can be
UFMylated on any given molecule (10). While the exact function of UFM1-conjugation on 14-3-
3ε is still unclear, it may play a role in coordinating multiple protein partners at membrane signaling
sites, potentially including known interactors such as Riplet, TRIM25, or MAVS, whose binding to
14-3-3ε might be modulated by its UFMylation status. This model aligns with observations in other
systems where UFMylation has been shown to serve as a scaffold for optimizing signaling
complex organization (40), suggesting a similar role in RIG-I signaling.
Our results highlight the sophisticated regulation of 14-3-3 proteins through various PTMs.
Interestingly, K50 on 14-3-3ε can be acetylated and sumoylated (58, 59), and this residue in other
14-3-3 proteins has been shown to be ubiquitinated (8). This suggests that competing PTMs at

the same residue could mediate context-dependent functional switching between distinct cellular

pathways. Recent studies have also revealed intrinsic chaperone-like activities for 14-3-3 proteins

that broadly prevent aggregation or aberrant phase separation of client proteins (14). We

speculate that UFMylation could also enhance these intrinsic chaperone functions of 14-3-3

proteins during antiviral responses by stabilizing protein complexes critical to innate immune

signaling. This could explain how UFMylation promotes RIG-I signaling without affecting direct

binding interactions, by optimizing the molecular environment around the signaling complex

rather than altering binding interfaces directly. Future studies will be needed to investigate the

molecular events triggering site-specific UFMylation of 14-3-3 following RIG-I activation and to

determine whether other 14-3-3 isoforms undergo similar regulation.

In summary, by identifying K50 and K215 on 14-3-3ε as a critical target residues of

UFMylation that promote RIG-I signaling, our study reveals another mechanism for how this post-

translational modification system tunes diverse innate immune pathways, advancing our

fundamental understanding of innate immune regulation. The UFMylation pathway represents a

potential target for modulating antiviral responses in conditions characterized by either insufficient

immunity or excessive inflammation. Furthermore, as viruses have evolved mechanisms to

manipulate UFMylation, targeting this pathway could potentially disrupt various aspects of viral

infection. Future investigations of UFMylation in both infectious and inflammatory diseases may

yield important insights for developing novel immunomodulatory approaches.

Acknowledgements

We thank those colleagues who generously provided reagents, as indicated in the Methods, the

Duke Functional Genomics Core, the Duke Proteomics and Metabolomics Core Facility, and

members of the Horner Lab for valuable feedback and discussion. During the preparation of this

work, we used Perplexity for editing purposes. After using this tool/service, we reviewed and

sometimes made edits to our text. We take full responsibility for the content of the publication.

This work was supported by National Institutes of Health grants R01AI155512 (S.M.H),

R35GM119544 (K.M.S), and T32CA00911 (IVG; DLS).

Figures

Figure 1. UFM1 conjugation is required to promote 14-3-3ε-mediated enhancement of RIG-

I signaling to IFN-β. (A) IFN-β-promoter reporter luciferase expression (relative to *Renilla)* from

293T cells expressing (Vec) Vector, Flag-UFM1^{WT}, Flag-RIG-I, or HA-14-3-3ε. (B) IFN-β–

promoter reporter luciferase expression (relative to *Renilla*) from 293T cells expressing (Vec)

Vector, Flag-UFM1^{WT}, Flag-UFM1$^{\Delta C3}$, Flag-RIG-I, or HA-14-3-3ε. Bars represent the mean ± SD,

n = 3 independent replicates. *p ≤ 0.05, ** ≤ 0.01, ***p ≤ 0.001 determined by one way ANOVA

followed by Šidák's multiple comparisons test (A) or Tukey's multiple comparisons test (B).

Figure 2. UFM1 conjugation to 14-3-3ε requires residues K50 and K215 on 14-3-3ε. (A) UFMylation assay of anti-HA immunoprecipitated lysates from 293T UFM1 KO cells expressing Vector (Vec) or HA-14-3-3ε along with Myc-UFL1, Myc-UFBP1, and Flag-UFM1-WT, -G83A, or -ΔC3. (B) Purified UFMylation proteins (UFBP1, UFL1, UFC1, UBA5, and UFM1) as well as 14-3-3ε^{WT} or 14-3-3ε$^{K50/215R}$ were incubated in UFMylation assay buffer in the presence or absence of ATP. Following reaction termination, the samples were subjected to immunoblot analysis for 14-3-3ε (top) or using an antibody cocktail to all UFMylation proteins (bottom). (C) UFMylation assay of anti-HA immunoprecipitated lysates from 293T UFM1 KO cells expressing Vector (-), Flag-UFM1-WT, -G83A, or -ΔC3, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$. Representative blots from *n* = 3 independent replicates are shown.

Figure 3. K50 and K215 are not required for 14-3-3ε interaction with RIG-I. (A) Immunoblot

analysis of anti-HA immunoprecipitated extracts and inputs from 293T RIG-I KO cells expressing

Flag-RIG-I, and HA-14-3-3ε WT, K50/215R, or AAA. Representative blot from $n = 3$ independent

replicates is shown. (B) Relative quantification of the ratio of RIG-I (IP/relative input) to

immunoprecipitated 14-3-3ε is shown on the right (Mean ± SD, $n = 3$). *p ≤ 0.05 or non-significant

(ns) determined by one way ANOVA followed by Tukey's multiple comparisons test.

Figure 4. K50 and K215 are required for 14-3-3ε-mediated RIG-I activation of IFN. (A) IFN-β-promoter reporter luciferase expression from 293T RIG-I KO cells expressing Vector (-), Flag-RIG-I, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$, with RIG-I samples normalized to 100. (B-C) RT-qPCR analysis (rel. to *GAPDH*) of RNA extracted from 293T RIG-I KO cells transfected with Flag-RIG-I, and HA-14-3-3ε^{WT} or HA-14-3-3ε$^{K50/215R}$ followed by Sendia virus infection (6 h), with RIG-I samples normalized to 100. *p ≤ 0.05, ** ≤ 0.01, ***p ≤ 0.005 determined by one way ANOVA followed by (A) Bonferroni or (B-C) Tukey's multiple comparisons test. Mean ± SD, *n* = 3 independent replicates are shown.

Methods

Cell lines, viruses, and transfection. 293T, 293T-UFM1 KO, and 293T-RIG-I KO were grown in Dulbecco's modification of Eagle's medium (DMEM; Mediatech) supplemented with 10% fetal bovine serum, 1X minimum essential medium nonessential amino acids, and 25 mM of HEPES buffer. The 293T cells (CRL-3216) were obtained from American Type Culture Collection (ATCC). 293T-UFM1 KO and 293T-RIG-I KO were previously described (20, 60). All cell lines were verified as mycoplasma-free using the Look-Out Mycoplasma PCR detection kit (Sigma-Aldrich). Sendai virus Cantell strain was obtained from Charles River Laboratories and used at 200 HAU/ml. Sendai virus infections were performed in serum-free media (30 to 60 mins), after which complete medium was added. DNA transfections were performed using FuGene 6 (Promega). Cells were incubated for 24-48 hours post transfection prior to experimental treatment.

Plasmids. The following plasmids have been previously described: pIFN-β-luc (61), pCMV-Renilla (Promega), pcTak_Myc-5_UFL1 (20), pETDuet-His$_6$-TEV-UFL1-StrepII-3C-UFBP1$^{(21\text{-end})}$ (gift of Dr. Yogesh Kulathu) (28), pTRIPZ IFNB1^{2x}-Gluc (gift of Dr. Nandan Gokhale) (60), and pET-NT*-HRV3CP (gift of Dr. Gottfried Otting; Addgene #162795) (55). The following plasmids were generated by insertion of PCR-amplified fragments into NotI-to-PmeI digested pEF-TAK-Flag or into BamHI-to-NotI digested pGEX-GST using InFusion cloning (Takara): pGEX-GST-14-3-3ε^{WT}, pGEX-GST-14-3-3ε$^{K50/215R}$, pGEX-GST-UBA5, pGEX-GST-UFC1, and pGEX-GST-UFM1. The following plasmids were generated by site-directed mutagenesis: pEF-TAK-HA-14-3-3ε$^{K50/215R}$, pEF-TAK-HA-14-3-3ε^{AAA} (E251-253A), and pEF-TAK-Flag-UFM1$^{\Delta C2}$. The following were cloned by InFusion: pEF-TAK-Flag-UFM1^{WT}, pEF-TAK-Flag-UFM1$^{\Delta C3}$, and pEF-TAK-Flag-UFM1 G83A from pEF-TAK-HA-UFM1 (20); pEF-TAK-Flag-RIG-I from pEF-Bos-Flag-RIG-I (62), pEF-TAK-HA-14-3-3ε^{WT} from pEF-TAK-Flag-14–3-3ε (20), and pEF-TAK-Myc-UFBP1 by PCR from BC115617. Plasmid sequences were verified by DNA sequencing.

IFN-β promoter luciferase assay. IFN-β promoter luciferase assays were performed at 18 to 24 hours after treatment and transfection of pIFN-β-luc and pCMV-Renilla by using the Dual-Luciferase Reporter System (Promega). *Firefly* luciferase was normalized to *Renilla* luciferase, whose expression is driven by the CMV promoter. For the *Gaussia* luciferase assays, 48 hours post transfection of pTRIPZ IFNB1^{2x}-Gluc, supernatant was assayed for *Gaussia* luciferase using the *Renilla* Luciferase Assay System (Promega). All samples were read using a BioTek Synergy2 microplate reader.

Immunoblotting. Cells were lysed with either RIPA (10 mM tris HCl pH [7.5], 150 mM NaCl, 0.5% sodium deoxycholate, and 1% Triton X-100 or NP-40 Lysis Buffer (20 mM Tris HCl [pH 7.4], 100 mM NaCl, 0.5% NP-40) supplemented with protease inhibitor mixture (Sigma-Aldrich), and Halt inhibitor phosphatase (Thermo Fisher Scientific), and lysates were isolated by centrifugation. Quantified protein was resolved by SDS/PAGE and transferred to nitrocellulose membranes. Membranes were stained with Revert 700 total protein stain (Li-COR Biosciences) and then blocked with 3% bovine serum albumin in phosphate-buffered saline containing 0.01% Tween-20 (PBS-T). Following incubation with primary antibodies, membranes were incubated with species-specific horseradish-peroxidase (HRP)-conjugated antibodies (Jackson Immuno Research 1:5000), followed by membrane treatment with Clarity Western ECL substrate (Bio-Rad) and imaging on a LICOR Odyssey FC. The following antibodies were used for immunoblotting: R-anti-SenV (MBL, 1:1000), HA-HRP (Sigma-Aldrich 1:1000-1:3000), Flag-HRP (1:1000-1:5000), and M-anti-Myc (Cell Signaling Technology 1:1000).

Protein immunoprecipitation. Cells were lysed with either RIPA (10 mM tris HCl [pH 7.5], 150 mM NaCl, 0.5% sodium deoxycholate, and 1% triton X-100 or NP-40 Lysis Buffer (20 mM Tris HCl [pH 7.4], 100 mM NaCl, 0.5% NP-40) supplemented with protease inhibitor mixture (Sigma-Aldrich), and Halt phosphatase inhibitor, and 10% glycerol. Quantified protein (between 100-200 µg) was incubated anti-HA magnetic beads (Invitrogen) in lysis buffer either at room temperature for 1-2 hrs or 4°C overnight with head-over-tail rotation. Beads were washed 3X in lysis buffer and eluted with 2X Laemmli buffer (Bio-Rad) with 5% 2-mercaptoenthanol at 95°C for 5 minutes. Proteins were resolved by SDS/PAGE and immunoblotting, as above. For UFMylation precipitations, cells were lysed UFMylation lysis buffer (50 mM Tris [pH 8], 150 mM NaCl, 0.5% sodium deoxycholate, and 1% NP-40) supplemented with 10% glycerol, 1:100 protease and phosphatase inhibitors, as above, and 1:100 *N*-ethylmaleimide (10 mM final) (Sigma-Aldrich). Post-nuclear lysates were boiled at 95°C for 10 mins and incubated with Pierce anti-HA magnetic beads with UFMylation lysis buffer. Beads were washed 3X in high salt buffer (10 mM Tris [pH 8], 1 mM EDTA, 500 mM NaCl) and eluted with 2X Laemmli buffer with 5% 2-mercaptoenthanol at 95°C for 5 minutes.

Quantification of immunoblots. Immunoblots were imaged using the LICOR Odyssey FC system and quantified using ImageStudio software, and raw values were normalized to relevant controls for each antibody. RIG-I and 14-3-3ε interaction were quantified as ratio of inputs of RIG-

I and 14-3-3ε normalized to total protein and displayed as the relative ratio of RIG-I over 14-3-3ε

pull down.

Purification of GST/His-tagged proteins. pGEX-GST-UBA5, pGEX-GST-UFC1, pGEX-GST-

UFM1, pGEX-GST-14-3-3ε, pGEX-GST-14-3-3ε K50/215R, and pETDuet-His$_6$-TEV-UFL1-

StrepII-3C-UFBP1$^{(21-end)}$ were transformed in *E. coli* Rosetta cells. Cells were cultured at 37°C

until OD600 reached 0.6 to 0.8. Then, cells were induced with a final concentration of 1 mM IPTG.

Cells were further cultured at 30°C for 3.5-4 hours. Cells were collected by centrifugation for 30

mins at 7,000 rpm. The pellets were lysed in NET--N buffer (50 mM Tris-HCl [pH 7.5], 150 mM

NaCl, 0.5% IGEPAL, supplemented with Aprotinin, Leupeptin, PMSF, and dithiothreitol (DTT) at

1:100. Cells were sonicated 3 times for 30 sec at output 6-7 and a duty cycle of 80%. Then,

lysates were centrifuged for 30 mins at 12,000 rpm at 4°C. Supernatant was incubated with GST

beads that were washed with NETN buffer with rotation at 4C for 1-2 hours. The beads were

washed 3 times with NETN buffer and 3 times with PBS supplemented with 1:100 DTT. After

washes, GST tag was cleaved with purified human rhinovirus 3C protease (purified as described

(55)) and protein was eluted with PBS by incubation with rotation at 4°C overnight.

***In vitro* UFMylation.** 2 µM of substrate, 0.25 µM of UBA5, 5 µM UFC1, 20 ng UFL1/UFBP1, 30

µM UFM1 were incubated in a reaction buffer containing 50 mM HEPES [pH 7.5], 10 mM MgCl$_2$,

with or without 5 mM ATP for 1 hr at 37°C. The reaction was stopped by the addition of 2X SDS

loading with 5% 2-mercaptoethanol and incubated at 95°C for 5 minutes The reaction products

were then separated by SDS/PAGE and analyzed by immunoblotting, as above.

RNA Analysis. Total cellular RNA was extracted using RNeasy Plus mini kit (Qiagen). RNA was

then reverse transcribed using iScript complementary DNA (cDNA) synthesis kit (Bio-Rad),

following manufacturer's instructions. The resulting DNA was diluted 1:3 in double distilled water.

RT-qPCR was performed in triplicate using the Power SYBR Green PCR master mix (Thermo

Fisher Scientific) and QuantStudio 6 Flex RT-PCR system. Oligonucleotide sequences are as

follows: IFNB1 F-5'-CTTTGCTATTTTCAGACAAGATTCA; R-5'GCCAGGAGGTTCTCAACAAT,

IFNL1 F-5'-CTTCCAAGCCCACCACAACT; R-5'-GGCCTCCAGGACCTTCAGC, and

GAPDH F-5'-AAGGTGAAGGTCGGAGTCAAC; R-5'-GGGGTCATTGATGGCAACAATA.

Sample Preparation and LC-MS/MS Analysis. Samples were spiked with undigested bovine

casein at a total of either 120 or 240 pmol as an internal quality control standard. Next, samples

were supplemented with 10 µL of 20% SDS, reduced with 10 mM DTT for 30 min at 80°C,

alkylated with 20 mM iodoacetamide for 30 min at room temperature, then supplemented with a

final concentration of 1.2% phosphoric acid and 546 µL of S-Trap (Protifi) binding buffer (90%

MeOH/100 mM TEAB). Proteins were trapped on the S-Trap micro cartridge, digested using 20

ng/µL sequencing grade trypsin (Promega) for 1 hr at 47°C, and eluted using 50 mM TEAB,

followed by 0.2% FA, and lastly using 50% ACN/0.2% FA. All samples were then lyophilized to

dryness. Samples were resolubilized using 48 µL of 1% TFA/2% ACN with 25 fmol/µL yeast ADH.

Quantitative LC/MS/MS was performed on 1.5 µL (3.13% of total sample) using an MClass UPLC

system (Waters Corp) coupled to an Orbitrap Fusion Lumos high resolution accurate mass

tandem mass spectrometer (Thermo Fisher Scientific) via a nanoelectrospray ionization source.

Briefly, the sample was first trapped on a Symmetry C18 20 mm × 180 µm trapping column (5

µl/min at 99.9/0.1 v/v water/acetonitrile), after which, the analytical separation was performed

using a 1.8 µm Acquity HSS T3 C18 75 µm × 250 mm column (Waters Corp.) with a 90-min linear

gradient of 5 to 30% acetonitrile with 0.1% formic acid at a flow rate of 400 nanoliters/minute

(nL/min) with a column temperature of 55°C. Data collection on the Fusion Lumos mass

spectrometer was performed in a data-dependent acquisition (DDA) mode of acquisition with a

r=120,000 (@ m/z 200) full MS scan from m/z 375 – 1500 with a target AGC value of 4e5 ions

was performed. MS/MS scans were acquired in the linear ion trap in "rapid" mode with a target

AGC value of 1e5 and max fill time of 100 ms. The total cycle time 2s, with total cycle times of 2

sec between like full MS scans. A 20s dynamic exclusion was employed to increase depth of

coverage. The total analysis cycle time for each sample injection was approximately 2 hours.

Proteomics Data Analysis. Following UPLC-MS/MS analyses, data were imported into

Proteome Discoverer 2.5 (Thermo Fisher Scientific). In addition to quantitative signal extraction,

the MS/MS data was searched against the SwissProt *H. sapiens* database (downloaded in Nov

2019) and a common contaminant/spiked protein database (bovine albumin, bovine casein, yeast

ADH, etc.), and an equal number of reversed-sequence "decoys" for false discovery rate

determination. Sequest (v 2.5, Thermo PD) was utilized to produce fragment ion spectra and to

perform the database searches. Database search parameters included fixed modification on Cys

(carbamidomethyl) and variable modification on Met (oxidation) and Lys (UFMylation). Search

tolerances were 2ppm precursor and 0.8Da product ion with semi-trypsin enzyme rules.

Discoverer were used to annotate the data at a maximum 1% protein false discovery rate based

on q-value calculations. Note that peptide homology was addressed using razor rules in which a

peptide matched to multiple different proteins was exclusively assigned to the protein has more

identified peptides. Protein homology was addressed by grouping proteins that had the same set

of peptides to account for their identification. A master protein within a group was assigned based

on % coverage.

Statistical analysis. We used one-way ANOVA followed by appropriate multiple comparisons

test using GraphPad Prism software. Graphed values are presented as mean ± SD (n = 3); *$P \leq$

0.05, **$P \leq 0.01$, and ***$P \leq 0.001$.

References

1. Rehwinkel J, Gack MU. 2020. RIG-I-like receptors: their regulation and roles in RNA
sensing. Nat Rev Immunol 20:537-551.

2. Crow YJ, Stetson DB. 2022. The type I interferonopathies: 10 years on. Nat Rev
Immunol 22:471-483.

3. Iwasaki A, Medzhitov R. 2015. Control of adaptive immunity by the innate immune
system. Nat Immunol 16:343-53.

4. Taft J, Bogunovic D. 2018. The Goldilocks Zone of Type I IFNs: Lessons from Human
Genetics. J Immunol 201:3479-3485.

5. Lin JP, Fan YK, Liu HM. 2019. The 14-3-3eta chaperone protein promotes antiviral
innate immunity via facilitating MDA5 oligomerization and intracellular redistribution.
PLoS Pathog 15:e1007582.

6. Liu HM, Loo YM, Horner SM, Zornetzer GA, Katze MG, Gale M, Jr. 2012. The
mitochondrial targeting chaperone 14-3-3epsilon regulates a RIG-I translocon that
mediates membrane association and innate antiviral immunity. Cell Host Microbe
11:528-37.

7. Aitken A. 2011. Post-translational modification of 14-3-3 isoforms and regulation of
cellular function. Semin Cell Dev Biol 22:673-80.

8. Pennington KL, Chan TY, Torres MP, Andersen JL. 2018. The dynamic and stress-
adaptive signaling hub of 14-3-3: emerging mechanisms of regulation and context-
dependent protein-protein interactions. Oncogene 37:5587-5604.

9. Henriksson ML, Francis MS, Peden A, Aili M, Stefansson K, Palmer R, Aitken A,
Hallberg B. 2002. A nonphosphorylated 14-3-3 binding motif on exoenzyme S that is
functional in vivo. Eur J Biochem 269:4921-9.

10. Ottmann C, Yasmin L, Weyand M, Veesenmeyer JL, Diaz MH, Palmer RH, Francis MS,
Hauser AR, Wittinghofer A, Hallberg B. 2007. Phosphorylation-independent interaction
between 14-3-3 and exoenzyme S: from structure to pathogenesis. EMBO J 26:902-13.

11. Liu J, Cao S, Ding G, Wang B, Li Y, Zhao Y, Shao Q, Feng J, Liu S, Qin L, Xiao Y. 2021.
The role of 14-3-3 proteins in cell signalling pathways and virus infection. J Cell Mol Med
25:4173-4182.

12. Obsilova V, Obsil T. 2022. Structural insights into the functional roles of 14-3-3 proteins.
Front Mol Biosci 9:1016071.

13. Toleman CA, Schumacher MA, Yu SH, Zeng W, Cox NJ, Smith TJ, Soderblom EJ,
Wands AM, Kohler JJ, Boyce M. 2018. Structural basis of O-GlcNAc recognition by
mammalian 14-3-3 proteins. Proc Natl Acad Sci U S A 115:5956-5961.

14. Segal D, Maier S, Mastromarco GJ, Qian WW, Nabeel-Shah S, Lee H, Moore G,
Lacoste J, Larsen B, Lin ZY, Selvabaskaran A, Liu K, Smibert C, Zhang Z, Greenblatt J,

Peng J, Lee HO, Gingras AC, Taipale M. 2023. A central chaperone-like role for 14-3-3
proteins in human cells. Mol Cell 83:974-993 e15.

15.   Chan YK, Gack MU. 2016. A phosphomimetic-based mechanism of dengue virus to
antagonize innate immunity. Nat Immunol 17:523-30.

16.   Riedl W, Acharya D, Lee JH, Liu G, Serman T, Chiang C, Chan YK, Diamond MS, Gack
MU. 2019. Zika Virus NS3 Mimics a Cellular 14-3-3-Binding Motif to Antagonize RIG-I-
and MDA5-Mediated Innate Immunity. Cell Host Microbe 26:493-503 e6.

17.   Andrews DDT, Vlok M, Akbari Bani D, Hay BN, Mohamud Y, Foster LJ, Luo H, Overall
CM, Jan E. 2023. Cleavage of 14-3-3epsilon by the enteroviral 3C protease dampens
RIG-I-mediated antiviral signaling. J Virol 97:e0060423.

18.   Liu J, Qian C, Cao X. 2016. Post-Translational Modification Control of Innate Immunity.
Immunity 45:15-30.

19.   Zhou Y, He C, Wang L, Ge B. 2017. Post-translational regulation of antiviral innate
signaling. Eur J Immunol 47:1414-1426.

20.   Snider DL, Park M, Murphy KA, Beachboard DC, Horner SM. 2022. Signaling from the
RNA sensor RIG-I is regulated by ufmylation. Proc Natl Acad Sci U S A
119:e2119531119.

21.   Komatsu M, Chiba T, Tatsumi K, Iemura S, Tanida I, Okazaki N, Ueno T, Kominami E,
Natsume T, Tanaka K. 2004. A novel protein-conjugating system for Ufm1, a ubiquitin-
fold modifier. EMBO J 23:1977-86.

22.   Tatsumi K, Sou YS, Tada N, Nakamura E, Iemura S, Natsume T, Kang SH, Chung CH,
Kasahara M, Kominami E, Yamamoto M, Tanaka K, Komatsu M. 2010. A novel type of
E3 ligase for the Ufm1 conjugation system. J Biol Chem 285:5417-27.

23.   Komatsu M, Inada T, Noda NN. 2024. The UFM1 system: Working principles, cellular
functions, and pathophysiology. Mol Cell 84:156-169.

24.   Wei Y, Xu X. 2016. UFMylation: A Unique & Fashionable Modification for Life. Genomics
Proteomics Bioinformatics 14:140-146.

25.   Banerjee S, Kumar M, Wiener R. 2020. Decrypting UFMylation: How Proteins Are
Modified with UFM1. Biomolecules 10.

26.   Millrine D, Peter JJ, Kulathu Y. 2023. A guide to UFMylation, an emerging
posttranslational modification. FEBS J 290:5040-5056.

27.   Zhou X, Mahdizadeh SJ, Le Gallo M, Eriksson LA, Chevet E, Lafont E. 2024.
UFMylation: a ubiquitin-like modification. Trends Biochem Sci 49:52-67.

28.   Peter JJ, Magnussen HM, DaRosa PA, Millrine D, Matthews SP, Lamoliatte F,
Sundaramoorthy R, Kopito RR, Kulathu Y. 2022. A non-canonical scaffold-type E3
ligase complex mediates protein UFMylation. EMBO J 41:e111015.

29.   Zhang Y, Yan M, Yu Y, Wang J, Jiao Y, Zheng M, Zhang S. 2024. 14-3-3epsilon: a
protein with complex physiology function but promising therapeutic potential in cancer.
Cell Commun Signal 22:72.

30.   Liang Z, Ning R, Wang Z, Kong X, Yan Y, Cai Y, He Z, Liu XG, Zou Y, Zhou J. 2024.
The emerging roles of UFMylation in the modulation of immune responses. Clin Transl
Med 14:e70019.

31.   Qin B, Yu J, Nowsheen S, Wang M, Tu X, Liu T, Li H, Wang L, Lou Z. 2019. UFL1
promotes histone H4 ufmylation and ATM activation. Nat Commun 10:1242.

32.   Qin B, Yu J, Nowsheen S, Zhao F, Wang L, Lou Z. 2020. STK38 promotes ATM
activation by acting as a reader of histone H4 ufmylation. Sci Adv 6:eaax8214.

33.   Ishimura R, El-Gowily AH, Noshiro D, Komatsu-Hirota S, Ono Y, Shindo M, Hatta T, Abe
M, Uemura T, Lee-Okada HC, Mohamed TM, Yokomizo T, Ueno T, Sakimura K,
Natsume T, Sorimachi H, Inada T, Waguri S, Noda NN, Komatsu M. 2022. The UFM1
system regulates ER-phagy through the ufmylation of CYB5R3. Nat Commun 13:7857.

34.   Liang JR, Lingeman E, Luong T, Ahmed S, Muhar M, Nguyen T, Olzmann JA, Corn JE.
2020. A Genome-wide ER-phagy Screen Highlights Key Roles of Mitochondrial
Metabolism and ER-Resident UFMylation. Cell 180:1160-1177 e20.

35.   Scavone F, Gumbin SC, Da Rosa PA, Kopito RR. 2023. RPL26/uL24 UFMylation is
essential for ribosome-associated quality control at the endoplasmic reticulum. Proc Natl
Acad Sci U S A 120:e2220340120.

36.   Walczak CP, Leto DE, Zhang L, Riepe C, Muller RY, DaRosa PA, Ingolia NT, Elias JE,
Kopito RR. 2019. Ribosomal protein RPL26 is the principal target of UFMylation. Proc
Natl Acad Sci U S A 116:1299-1308.

37.   Wang L, Xu Y, Rogers H, Saidi L, Noguchi CT, Li H, Yewdell JW, Guydosh NR, Ye Y.
2020. UFMylation of RPL26 links translocation-associated quality control to endoplasmic
reticulum protein homeostasis. Cell Res 30:5-20.

38.   Xu X, Huang W, Bryant CN, Dong Z, Li H, Wu G. 2024. The ufmylation cascade controls
COPII recruitment, anterograde transport, and sorting of nascent GPCRs at ER. Sci Adv
10:eadm9216.

39.   He C, Xing X, Chen HY, Gao M, Shi J, Xiang B, Xiao X, Sun Y, Yu H, Xu G, Yao Y, Xie
Z, Xing Y, Budiarto BR, Chen SY, Gao Y, Lee YR, Zhang J. 2024. UFL1 ablation in T
cells suppresses PD-1 UFMylation to enhance anti-tumor immunity. Mol Cell 84:1120-
1138 e8.

40.   Yoo HM, Kang SH, Kim JY, Lee JE, Seong MW, Lee SW, Ka SH, Sou YS, Komatsu M,
Tanaka K, Lee ST, Noh DY, Baek SH, Jeon YJ, Chung CH. 2014. Modification of ASC1
by UFM1 is crucial for ERalpha transactivation and breast cancer development. Mol Cell
56:261-274.

41. Zhou J, Ma X, He X, Chen B, Yuan J, Jin Z, Li L, Wang Z, Xiao Q, Cai Y, Zou Y, Cong
YS. 2023. Dysregulation of PD-L1 by UFMylation imparts tumor immune evasion and
identified as a potential therapeutic target. Proc Natl Acad Sci U S A 120:e2215732120.

42. Lee L, Perez Oliva AB, Martinez-Balsalobre E, Churikov D, Peter J, Rahmouni D, Audoly
G, Azzoni V, Audebert S, Camoin L, Mulero V, Cayuela ML, Kulathu Y, Geli V, Lachaud
C. 2021. UFMylation of MRE11 is essential for telomere length maintenance and
hematopoietic stem cell survival. Sci Adv 7:eabc7371.

43. Wang Z, Gong Y, Peng B, Shi R, Fan D, Zhao H, Zhu M, Zhang H, Lou Z, Zhou J, Zhu
WG, Cong YS, Xu X. 2019. MRE11 UFMylation promotes ATM activation. Nucleic Acids
Res 47:4124-4135.

44. Kulsuptrakul J, Wang R, Meyers NL, Ott M, Puschnik AS. 2021. A genome-wide
CRISPR screen identifies UFMylation and TRAMP-like complexes as host factors
required for hepatitis A virus infection. Cell Rep 34:108859.

45. Schmidt HM, Sorensen GC, Lanahan MR, Grabowski J, Park M, Horner SM. 2025.
UFMylation promotes orthoflavivirus infectious particle production. bioRxiv
doi:10.1101/2025.01.09.632082:2025.01.09.632082.

46. Yiu SPT, Zerbe C, Vanderwall D, Huttlin EL, Weekes MP, Gewurz BE. 2023. An
Epstein-Barr virus protein interaction map reveals NLRP3 inflammasome evasion via
MAVS UFMylation. Mol Cell 83:2367-2386 e15.

47. Jing J, Yang F, Wang K, Cui M, Kong N, Wang S, Qiao X, Kong F, Zhao D, Ji J, Tang L,
Gao J, Cong YS, Ding D, Chen K. 2025. UFMylation of NLRP3 Prevents Its Autophagic
Degradation and Facilitates Inflammasome Activation. Adv Sci (Weinh)
doi:10.1002/advs.202406786:e2406786.

48. Wang X, Li C, Wang Y, Li L, Han Z, Wang G. 2020. UFL1 Alleviates LPS-Induced
Apoptosis by Regulating the NF-kappaB Signaling Pathway in Bovine Ovarian
Granulosa Cells. Biomolecules 10.

49. Cai Y, Zhu G, Liu S, Pan Z, Quintero M, Poole CJ, Lu C, Zhu H, Islam B, Riggelen JV,
Browning D, Liu K, Blumberg R, Singh N, Li H. 2019. Indispensable role of the Ubiquitin-
fold modifier 1-specific E3 ligase in maintaining intestinal homeostasis and controlling
gut inflammation. Cell Discov 5:7.

50. Li C, Li L, Chen K, Wang Y, Yang F, Wang G. 2019. UFL1 Alleviates
Lipopolysaccharide-Induced Cell Damage and Inflammation via Regulation of the
TLR4/NF-kappaB Pathway in Bovine Mammary Epithelial Cells. Oxid Med Cell Longev
2019:6505373.

51. Balce DR, Wang YT, McAllaster MR, Dunlap BF, Orvedahl A, Hykes BL, Jr., Droit L,
Handley SA, Wilen CB, Doench JG, Orchard RC, Stallings CL, Virgin HW. 2021.
UFMylation inhibits the proinflammatory capacity of interferon-gamma-activated
macrophages. Proc Natl Acad Sci U S A 118.

52. Tao Y, Yin S, Liu Y, Li C, Chen Y, Han D, Huang J, Xu S, Zou Z, Yu Y. 2023. UFL1
promotes antiviral immune response by maintaining STING stability independent of
UFMylation. Cell Death Differ 30:16-26.

53. Horner SM, Wilkins C, Badil S, Iskarpatyoti J, Gale M, Jr. 2015. Proteomic analysis of
mitochondrial-associated ER membranes (MAM) during RNA virus infection reveals
dynamic changes in protein and organelle trafficking. PLoS One 10:e0117963.

54. Zhou J, Liang Q, Dong M, Ma X, Jin Y, Guan D, Liu J, Wang M, Cong YS. 2022.
Optimized protocol to detect protein UFMylation in cells and in vitro via immunoblotting.
STAR Protoc 3:101074.

55. Abdelkader EH, Otting G. 2021. NT*-HRV3CP: An optimized construct of human
rhinovirus 14 3C protease for high-yield expression and fast affinity-tag cleavage. J
Biotechnol 325:145-151.

56. Cantell K, Hirvonen S, Kauppinen HL, Myllyla G. 1981. Production of interferon in
human leukocytes from normal donors with the use of Sendai virus. Methods Enzymol
78:29-38.

57. Sumpter R, Jr., Loo YM, Foy E, Li K, Yoneyama M, Fujita T, Lemon SM, Gale M, Jr.
2005. Regulating intracellular antiviral defense and permissiveness to hepatitis C virus
RNA replication through a cellular RNA helicase, RIG-I. J Virol 79:2689-99.

58. Fan K, Zhu K, Wang J, Ni X, Shen S, Gong Z, Cheng X, Zhang C, Liu H, Suo T, Ni X,
Liu H. 2022. Inhibition of 14-3-3epsilon by K50 acetylation activates YAP1 to promote
cholangiocarcinoma growth. Exp Cell Res 421:113404.

59. Hendriks IA, Lyon D, Young C, Jensen LJ, Vertegaal AC, Nielsen ML. 2017. Site-
specific mapping of the human SUMO proteome reveals co-modification with
phosphorylation. Nat Struct Mol Biol 24:325-336.

60. Gokhale NS, Sam RK, Somfleth K, Thompson MG, Marciniak DM, Smith JR, Genoyer E,
Eggenberger J, Chu LH, Park M, Dvorkin S, Oberst A, Horner SM, Ong SE, Gale M, Jr.,
Savan R. 2024. Cellular RNA interacts with MAVS to promote antiviral signaling.
Science 386:eadl0429.

61. Fredericksen B, Akkaraju GR, Foy E, Wang C, Pflugheber J, Chen ZJ, Gale M, Jr. 2002.
Activation of the interferon-beta promoter during hepatitis C virus RNA replication. Viral
Immunol 15:29-40.

62. Saito T, Hirai R, Loo YM, Owen D, Johnson CL, Sinha SC, Akira S, Fujita T, Gale M, Jr.
2007. Regulation of innate antiviral defenses through a shared repressor domain in RIG-
I and LGP2. Proc Natl Acad Sci U S A 104:582-7.
622