Projecting spatiotemporal bioclimatic niche dynamics of endemic Pyrenean plant species under

climate change: how much will we lose?

Noèmie Collette[1], Sébastien Pinel[2], Valérie Delorme-Hinoux[1], Joris A.M. Bertrand[1]

[1]Laboratoire Génome et Développement des Plantes (LGDP), UMR 5096, Université de Perpignan Via Domitia

(UPVD) - Centre National de la Recherche Scientifique (CNRS), Perpignan, France

[2]Centre de Formation et de Recherche sur les Environnements Méditerranéens (CEFREM), UMR 5110,

Université de Perpignan Via Domitia (UPVD) - Centre National de la Recherche Scientifique (CNRS),

Perpignan, France

Noèmie Collette[1], noemie.collette@univ-perp.fr , ORCID iD: https://orcid.org/0009-0000-4226-3878
Sébastien Pinel[2], sebastien.pinel@univ-perp.fr , ORCID iD: https://orcid.org/0000-0001-9608-754X
Valérie Delorme-Hinoux[1], valerie.hinoux@univ-perp.fr , ORCID iD: https://orcid.org/0009-0005-9994-6852
Joris A.M. Bertrand[1], joris.bertrand@univ-perp.fr , ORCID iD: https://orcid.org/0000-0002-3379-1019
14

Abstract

Species distributions are shifting under global change, with mountain ecosystems among the most vulnerable.

In such landscapes, ability to track changing conditions is limited, threatening narrowly distributed species. As

a mountain biodiversity hotspot in southwestern Europe, the Pyrenees harbors many such species, making it

a key case study for climate vulnerability assessments.

This study implements a bioclimatic niche modeling pipeline to evaluate climate change impact on endemic

Pyrenean plant species by 2100. Objectives are to (i) map current bioclimatic niche suitability, (ii) forecast its

future spatial dynamics, and (iii) identify potential climate refugia for conservation. Species occurrences were

combined with 19 bioclimatic variables (1x1 km resolution) to characterize bioclimatic niche suitability, using

an ensemble modeling approach integrating five algorithms (MaxEnt, Generalized Linear Model, Generalized

Additive Model, Gradient Boosting Machine, and Random Forest). Their future spatiotemporal dynamics were

projected under four climate scenarios (Shared Socioeconomic pathways 126, 245, 370, 585) for four

successive periods spanning 2021 to 2100.

By 2100, 69% of endemic species are projected to lose over 75% of their bioclimatic niche, and half to face

complete losses under high-emission scenarios. Only two species may gain suitable areas, highlighting the

need for species-specific conservation strategies. Bioclimatic niches are projected to shift by ~180 m upslope

and ~3 km in latitude on average, with areas of highest multi-species suitability, referred to as bioclimatic hotspots, becoming restricted to elevation above 2000 m. These trends intensify after 2041-2060 period, reflecting escalating climate pressures as the century progresses.

Our findings highlight the profound threat climate change may pose to endemic Pyrenean flora, with widespread bioclimatic niche losses projected by the century's end and high elevation refugia emerging as key conservation priorities. Anticipating these shifts and integrating them into conservation planning will be crucial to mitigating high-elevation biodiversity loss in a rapidly changing world.

Keywords: bioclimatic niche, climate change, endemic species, mountain, Pyrenees, species distribution models

1 | Introduction

Climate change represents one of the greatest environmental challenges of our time, with a significant and far-reaching impact on world's ecosystems (INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE, 2022 ; LI *et al.*, 2018). Among these ecosystems, mountain regions are of special concern as they are experiencing some of the highest and fastest rates of warming, a phenomenon widely recognized as elevation-dependent warming (PEPIN *et al.*, 2015). This refers to the tendency for warming rates at higher altitudes to surpass those observed in lowland regions, a pattern identified across multiple mountainous regions worldwide (PEPIN *et al.*, 2022 ; QIXIANG *et al.*, 2018). Such warming raises significant concerns for mountain ecosystems, which are particularly sensitive to these rapid changes (INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE (IPCC), 2023 ; KNIGHT, 2022). A cascade of ecological consequences follows (DAINESE *et al.*, 2024 ; ROGORA *et al.*, 2018), including shifts in species distribution, a well-documented response to climate change (LAWLOR *et al.*, 2024 ; PEARSE & DAVIES, 2019 ; RUBENSTEIN *et al.*, 2023).

Climate velocity describes the speed and direction in which a species needs to move to stay within its climatic niche as climate conditions change, which, in mountainous contexts, forces species to occupy progressively higher elevations (CHAN *et al.*, 2024), resulting in shifts upward in their core distribution (FREEMAN *et al.*, 2018). However, these elevational shifts are unlikely to keep pace with the rapidly changing climate (CARROLL *et al.*, 2015 ; GOODWIN *et al.*, 2025 ; KELLNER *et al.*, 2023). Mountain plants, in particular, face even greater challenges than other organisms due to their limited dispersal capacity (DI MUSCIANO *et al.*, 2020 ; MORGAN & VENN, 2017). Their migration is further constrained by the complex topography of mountainous landscapes and habitat fragmentation (BUENO DE MESQUITA *et al.*, 2018 ; ELSEN *et al.*, 2020), which create substantial barriers to

movement by breaking the spatial continuity of suitable areas. Even when they successfully reach climatically suitable habitat, or are already located there, establishment and persistence may be constrained by several factors. Biotic mismatches with pollinators (DAINESE *et al.*, 2024 ; MARSHALL *et al.*, 2020 ; TRUNSCHKE *et al.*, 2024 ; VITASSE *et al.*, 2021), competition with resident or incoming species (ALEXANDER *et al.*, 2015 ; SCHUCHARDT *et al.*, 2023) or the absence of key mutualists can hinder recruitment. Abiotic constraints, including unsuitable edaphic conditions (EICHEL *et al.*, 2023 ; NI & VELLEND, 2024) or microtopography may also reduce habitat suitability. Finally, intrinsic species traits such as germination requirements (ALEXANDER *et al.*, 2018) can further restrict successful colonization. Moreover, if its establishment occurs within an isolated microclimatic refugia, prolonged isolation of these persistent populations may compromise their long-term survival by limiting genetic mixing, thereby reducing genetic diversity (PINTO *et al.*, 2024 ; TURNOCK *et al.*, 2024). This, in turn, limits their adaptability and resilience to environmental changes (BIJLSMA & LOESCHCKE, 2012 ; FRANKHAM, 2005 ; MATHUR *et al.*, 2023). Such prolonged isolation mirrors patterns observed on a broader scale among mountaintop species, which persist in isolated high-elevation habitats often referred to as "sky island". Habitat fragmentation and progressive loss of physical space with elevation amplify extinction risk for these species (ELSEN & TINGLEY, 2015), as they lack alternative areas to persist when facing unfavorable conditions or competition with other species. This scenario, referred to as "mountaintop extinction", primarily threatens species with narrow distribution, residing in the upper limits of latitude gradients, or restricted to highest elevation.

The Pyrenees mountain chain, Europe's second-largest biodiversity hotspot with 3,652 indigenous vascular plants, is particularly concerning due to the number of mountaintop species, including a panel of endemics and specialized flora (GÓMEZ *et al.*, 2017). The region's complex climate and geodiversity have historically favored this exceptional diversity (KÖRNER & HILTBRUNNER, 2021 ; RAHBEK *et al.*, 2019a) and served as both refuge and corridor for organisms during past climatic changes (GARCÍA *et al.*, 2022 ; SCHMITT, 2009). However, intensifying global pressures such as climate, land use, and socio-economic changes, including tourism, urban development and agricultural abandonment (OPCC, 2018 ; ZANGO-PALAU *et al.*, 2024), may compromise its capacity to buffer future environmental shifts (RAHBEK *et al.*, 2019b). The upslope shift of plants in the Pyrenees (AMEZTEGUI *et al.*, 2016 ; LENOIR *et al.*, 2008 ; PAULI *et al.*, 2012) appears to contribute to increasing species richness on summits (CAMARERO *et al.*, 2006 ; STEINBAUER *et al.*, 2018). Yet, this dynamic often comes at the cost of endemic and range-restricted species.

Endemic species represent about 5 % of the Pyrenean flora and species at their upper distribution limits represent another 20 % (GÓMEZ *et al.*, 2017). Although not all of them are strictly confined to high elevations,

94	their limited distribution makes them particularly vulnerable to range contraction. Altogether, up to a quarter of
95	the regional flora may face mountain top extinction scenario in the coming decades. Observations in the Alps,
96	another European mountain range, support the potential mountaintop extinction risk for endemic and range-
97	restricted plant species in the Pyrenees. Red-listed plants have experienced rapid contraction of their elevation
98	margins, resulting in significant habitat loss in lowland areas (GEPPERT *et al.*, 2023), and endemic ones
99	undergoing substantial range reductions, with projections suggesting that 40 % of their current distribution
100	could become climatically unsuitable by the end of the 21st century (DULLINGER *et al.*, 2012). Similar dynamics
101	in the Pyrenees could drive extinctions, altering summit biodiversity and ecosystems. Although alpine plants
102	show traits promoting resilience, such as adaptation to harsh conditions, clonal growth, and access to diverse
103	thermal microhabitats (KÖRNER & HILTBRUNNER, 2021), these ongoing dynamics raise concerns about their
104	ability to track or adapt to climatic changes, with potential consequences for mountain biodiversity and
105	ecological communities. In this context, identifying potential responses of species to these changes is essential
106	for anticipating and guiding conservation efforts.
107	To assess the impact of climate change on plant distributions, the use of modeling tools is a widely adopted
108	approach. Species distribution models (SDMs), most often implemented as correlative approaches, relate
109	species occurrences to environmental variables to predict and describe habitat suitability or probability of
110	presence of a species across spatial and temporal scales (VASCONCELOS *et al.*, 2024). The robust
111	methodological foundations and predictive power of these models have led to their application across diverse
112	fields, including agriculture, public health, and, notably, conservation biology (FRANKLIN, 2023). In the
113	Pyrenees, SDMs have primarily been used for conservation purposes. Research has explored species
114	distribution drivers (DALIBARD *et al.*, 2021 ; KOUBA *et al.*, 2011 ; WILLIAMS-TRIPP *et al.*, 2012), projected climate-
115	driven shifts (DE POUS *et al.*, 2015 ; GLAD & MALLARD, 2022 ; MARTÍNEZ *et al.*, 2012 ; PÉREZ-GARCÍA *et al.*, 2013
116	; RUSSO *et al.*, 2023 ; SALVADO *et al.*, 2025, 2022), and reconstructed distribution over geological time
117	(BIDEGARAY-BATISTA *et al.*, 2016 ; CHARBONNEL *et al.*, 2016 ; GUTIERREZ *et al.*, 2022). Together, these studies
118	underscore the value of SDMs in informing management and conservation strategies for many species. By
119	providing insights into species-environment relationships and distribution patterns for past, present and future,
120	SDMs become relevant tools for anticipating and addressing climate change impacts, and, more generally,
121	the impact of global change on biodiversity.
122	Open-access environmental and occurrence databases have expanded possibilities for studying species-
123	environment relationships. Although various tools exist for each SDMs stage (KASS *et al.*, 2024), fully
124	integrated pipeline (i.e. framework), especially those suited for rare or range-restricted species, such as

endemics, remain scarce. Default settings in existing frameworks are often not fully suited for these taxa and

could require manual adjustments that become time-consuming when applied to many species. To address

this, we proposed a fully integrated and reproducible SDMs pipeline specifically tailored to species with few

occurrences. It automates algorithm-specific tuning, provides auxiliary scripts to guide key decisions (e.g. block

size for spatial cross-validation) and used evaluation metrics suited for low-prevalence data (e.g. Boyce Index,

AUC-PR). It also incorporates spatial visualization of uncertainties related to both algorithms and climate

projections, addressing the frequent lack of uncertainty reporting in SDMs for rare and endemic species (QAZI

et al., 2022). In addition to streamlining the modeling process, this pipeline produces harmonized outputs

across species, enhancing understanding of spatial and temporal biodiversity dynamics and enabling

consistent comparisons of climate-driven range shifts, making it well suited for multi-species assessments.

Ultimately, it provides natural area managers practical tools to preserve floristic diversity and sustain the

ecological resilience of mountain ecosystems in the face of accelerating climate change.

Endemic species, typically restricted to narrow geographical and climatic niches, serve as indicators of regional

biodiversity (LAMOREUX *et al.*, 2006). Their concentration in mountainous areas highlights the key role these

regions play in enhancing floristic richness and uniqueness. As irreplaceable components of the region's

natural heritage, their conservation is therefore crucial. In this context, the proposed framework, which enables

parallel analysis of SDMs outputs across species, is particularly valuable for identifying areas where several

endemic species may overlap and could be most at risk, thus supporting more effective and targeted

management efforts. Consequently, using SDMs, this study aims to: (i) map current distribution of bioclimatic

suitable areas for 59 endemic Pyrenean plant species, (ii) forecast future extent and dynamics of these areas

by the end of the 21st century under multiple climate scenarios, and (iii) identify where these species could find

refuge by 2081-2100 period.

2 | Material and methods

Study area and species occurrences data

This study follows the mountain delimitation of Körner *et al.* (2011), updated by Snethlage *et al.* (2022). Within

this framework, the Pyrenees span ~40,000 km² along the France-Spain-Andorra border (Figure 1), with over

150 peaks above 3000 m (GÓMEZ *et al.*, 2017). Their steep northern slopes contrast with broader southern

valleys, featuring diverse landscapes. At the Atlantic-Mediterranean-continental interface, this mountain range

displays diverse climatic conditions. Mean annual temperatures range from >10 °C in lowlands to <4 °C on

peaks, with precipitation varying from >2000 mm (Atlantic slopes) to ~1000 mm (sheltered areas) (CUADRAT

et al., 2024).

Among the 2,682 species listed in the Atlas of Pyrenean Flora (Pironon *et al.*, 2022), 84 were classified as endemic. Excluding endemic subspecies reduced the set to 78 species. Occurrence data were obtained from the Global Biodiversity Information Facility (GBIF.ORG, 2025), supplemented with research-grade data from iNaturalist (https://www.inaturalist.org). Occurrences since 1970 within the study area and ≤ 1 km coordinate uncertainty were retained, fitting the resolution of environmental variables. Spatial thinning retained one occurrence per grid cell. Species with <4 occurrences and/or highly restricted distributions were also removed, as selected SDMs assessment method requires presence of species in each spatial block during cross-validation, and therefore requires sufficient spatial representation. Ultimately, 59 species were included.

Bioclimatic data

WorldClim 2.1 (FICK & HIJMANS, 2017) provided 19 rasterized bioclimatic variables (see Supplemental Information, Appendix A) for the current (1970-2000) and future periods (2021-2040, 2041-2060, 2061-2080, 2081-2100 designated as 2030, 2050, 2070, 2090 in this study). Future datasets based on CMIP6 General Circulation Models (GCMs) at 0.5 arcs/min resolution (~1x1 km at the equator) were combined into median raster to capture model variability (CARROLL & MAHONY, 2025 ; SONG *et al.*, 2024). Projections follow four Shared Socio-economic Pathways (SSP, 126, 245, 370, 585, best to worst-case, respectively), representing socio-economic trajectory and radiative forcing (W.m^{-2}) expected in 2100 (O'NEILL *et al.*, 2017 ; WELCH, 2024). Under these scenarios, mean annual temperature in the Pyrenees (9.4 °C) is projected to rise by +18.69 % (11.56°C) to +39.51 % (15.54 °C) in 2081-2100 period, while current annual precipitation (975.41 mm) may decrease by -2.18 % (971.85 mm) to -13.57 % (843 mm) (Figure 2).

To reduce redundancy, highly positively correlated variables ($r > 0.8$) at presence locations for each species were clustered. Within each cluster, the most relevant variable per species was retained based on a predefined ecological priority list (Appendix A), which favors variables that are easier to interpret ecologically (e.g. mean annual temperature), followed by variables describing climatic extremes during cold and warm seasons, and finally more derived or complex indices (e.g. temperature seasonality, isothermality). Negatively correlated variables, reflecting complementary environmental gradients, were not excluded. Depending on species, models include between three and eight bioclimatic variables after the selection process. Selected variables for each species are listed in the Appendix A.

Bioclimatic suitability modeling

Model development

A presence/pseudo-absence approach was employed to improve SDMs reliability when true absences are

unavailable (MATEO *et al.*, 2010). Following recommendations from Barbet-Massin et al., 2012 and Biomod2

team (GUEGUEN *et al.*, 2025), we tested different pseudo-absence configurations they outlined and retained

the one that, in our analyses, consistently improved model evaluation scores (Appendix B). The number of

pseudo-absences was set equal to the number of presences and sampled within the environmental space,

using environmental constraints to restrict sampling to unsuitable conditions (BROUSSIN *et al.*, 2024). In line

with Biomod2 guidelines, each algorithm was run 10 times with distinct pseudo-absence sets generated

separately for each species, thereby reducing background dependence. Each set is spatially thinned to retain

a single occurrence per environmental grid cell.

As modeling algorithms significantly impact SDMs outcomes (BUISSON *et al.*, 2010 ; THUILLER *et al.*, 2019) and

predictive performance varying across approaches (NORBERG *et al.*, 2019 ; VALAVI *et al.*, 2022), bioclimatic

suitability was modeled using five algorithms : Maxent (MXT) Generalized Linear Models (GLM) Generalized

Additive Models (GAM), Gradient Boosting Machines (GBM) and RandomForest (RF). To ensure comparability

across algorithms, probability outputs from GLM, GAM, GBM, and RF were converted into favorability values

using the Fav function from the fuzzySim package (BARBOSA, 2015), which corrects sample prevalence

(presence/absence ratio). This transformation was not applied to MXT outputs, as it generates suitability

scores rather than probabilities.

Parameters tuning, evaluation and ensemble modeling

An automated workflow tuned model parameters for each iteration and algorithm. For each case, a parameter

combination was tested (see Appendix C), and the combination yielding the highest performance score was

retained. This score was calculated as a weighted average of three evaluation metrics: the Boyce Index (weight

0.5), AUC-PR (0.25), and Sensitivity (0.25). Greater weight was given to the Boyce Index as it best captures

how predicted suitability aligns with observed occurrences, ensuring more reliable projections from sparse

data. Each metric captured distinct aspect of model performance : discrimination, classification, and calibration

metrics (SILLERO *et al.*, 2021). Discrimination was assessed via Area Under the Precision-Recall Curve (AUC-

PR, 0-1 scale, with 0.5 indicating random performance), balancing precision and recall, making it particularly

relevant for rare species, as it focuses on correctly predicting presences without being inflated by abundance

of absences like AUC (MCDERMOTT *et al.*, 2025 ; SOFAER *et al.*, 2019). Classification ability was estimated

through Sensitivity, measuring true positive rates (0 for no true positives, 1 for perfect classification), crucial

given limited occurrences, where any omission of known occurrences would disproportionately affect model

reliability. Calibration was quantified using Boyce Index, assesses alignment between predictions and

observed presences (-1 for counter-prediction, 0 for random, and 1 for optimal prediction) (Liu *et al.*, 2025 ;

Sillero *et al.*, 2021). This metric relies solely on presence data, making it particularly advantageous in

presence-pseudoabsence frameworks where absence information is artificially generated.

For each pseudo-absence set, block cross-validation was used to identify the best-performing parameter set,

and the corresponding model was then run and its evaluation retained for subsequent analyses. A four-fold

(k=4) spatial cross-validation was used, training models on k-1 blocks and testing on the remaining one. Spatial

block size was derived from the range of environmental spatial autocorrelation, estimated using the

cv_spatial_autocor() function from the blockCV package (Valavi *et al.*, 2019), based on 65,496 randomly

sampled cells (subsampled by the function from the 1,244,424 cells available in the study area). Given limited

occurrence data, final block size was set to two thirds of the median autocorrelation range. This configuration

of k and block size balances environmental independence and data sufficiency that ensure presence records

in each fold. Larger blocks improve spatial independence but constrain k to avoid empty folds; conversely,

higher k increases replication but may reduce independence. A script is available (Appendix C) to explore

multiple k and block size combinations, aiming to retain as many species as possible. Once the optimal

parameters were selected per species and algorithm, models were refitted, and only those with Boyce Index

> 0.5 were retained for ensemble modeling. This threshold ensures minimal calibration quality.

Ensemble modeling enhances SDMs robustness by combining multiple model outputs, effectively improving

predictive accuracy by better capturing species distribution patterns (Marmion *et al.*, 2009 ; Zhu & Peterson,

2017). However, ensemble models do not always outperform individual ones in transferability (Zhu *et al.*, 2021

; Zhu & Peterson, 2017). For this reason, ensemble modeling was applied at two distinct stages: (i) for current

conditions, based on model runs calibrated and validated (Boyce Index > 0.5, maximum 5 algorithms x 10

pseudo absence set = 50 models) under present-day climate; and (ii) for future projections, by projecting these

same validated models to every combination of future periods and scenarios, and then creating an ensemble

for each period-scenario combination. Consensus ensemble maps were computed as pixel-wise median

across all algorithms and replicates, minimizing the influence of outliers and aligning with model central

tendencies (Marmion *et al.*, 2009 ; Zhu *et al.*, 2021). The entire pipeline is independently applied to each

species, resulting in 17 continuous distribution maps per species (1 under current conditions, 16 under future).

Ensemble models assessments were calculated by averaging metrics from models used to build the ensemble

(Boyce Index > 0.5). Algorithm performances were compared using Kruskal-Wallis, with Dunn's post hoc test

(Bonferroni correction) identifying pairwise differences between algorithms for each evaluation metric.

Assessing uncertainty and spatio-temporal dynamics

Uncertainty was quantified using standard deviation across predictions. Pixels with >5 % divergence, indicating

inconsistent predictions across algorithms, were masked for visualization on the current distribution map to

highlight most reliable regions. To assess SDMs reliability under novel conditions (i.e. conditions in future

periods deviating from the calibration range), the Multivariate Environmental Similarity Surface (MESS) index

(ELITH *et al.*, 2010) was applied. Negative MESS values, indicating extrapolation, were masked for visualization

on future distribution maps to highlight most reliable projections.

To quantify extent changes in suitable bioclimatic areas, maps were binarized for each period and SSP using

the MaxSSS threshold, appropriate for presence-only data (LIU *et al.*, 2016, 2013). Spatial overlap between

current and future suitable areas was also computed on binarized maps, yielding the proportion of the present

distribution projected to persist under each SSP. Spatial shifts between the current period and 2081-2100 were

assessed by overlaying binarized projections with elevation data (CGIAR-CSI, 2018, ~1x1 km at the equator)

and latitude. Mean latitude and elevation of suitable pixels were calculated for the current period and for each

SSP in 2081-2100 period. Values were averaged across the four scenarios, and only species retaining suitable

areas in all SSPs were used to estimate shifts between the present and future. Continuous suitability maps for

the current and 2081-2100 periods and all SSPs were averaged across species, with persistent suitable pixels

identified as bioclimatic hotspots (areas where the highest number of species could find suitable bioclimatic

conditions). Results are measured on a continuous scale to emphasize relative hotspot importance.

Framework and model characteristics were documented following the ODMAP protocol (Zurell *et al.*, 2020,

Appendix C) and summarized in Figure 3.

3 | Results

Models performances

Between 19 and 50 models per species were retained for ensemble modeling based on the Boyce Index

criterion (Boyce Index > 0.5). Significant differences in model performance across the five algorithms were

detected for the Boyce Index and Sensitivity (Kruskal–Wallis, $p < 0.001$ and 0.01 respectively; see Figure 4a

for p-values), while no significant difference was found for AUC-PR ($p = 0.07$). Dunn's post hoc test revealed

significant pairwise differences between algorithms. For the Boyce Index, MXT and GBM outperformed GAM,

GLM and RF. Regarding Sensitivity, MXT outperformed GLM and GAM, while GBM and RF outperformed

GLM. Other comparisons were not significant (Figure 4a). This highlights variation in predictive performance,

with GBM and MXT generally performing best, and RF consistently performed well in identifying suitable areas

where species are present. GLM exhibited the lowest Boyce Index values, falling under 0.5 for multiple species,

indicating weaker prediction consistency.

Ensemble model performance ranged from 0.67-0.98 for AUC-PR, 0.62-0.92 for the Boyce index and 0.87-

0.98 for sensitivity (Figure 4b). Poor evaluation metrics were not consistently linked to species with fewer

presence points, suggesting additional factors influencing performance.

Projected bioclimatic niche loss and species vulnerability

The spatial extent of bioclimatic suitable areas for endemic Pyrenean plant species is projected to decline

under climate change (Figure 5). By the end of the century, on average across SSPs, 57 out of the 59 species

are projected to lose a significant part of their suitable bioclimatic niche, with 41 species losing >75 %, 11

losing 50-75 % and 7 experiencing a more moderate loss of <50 %. These 57 species are expected to lose on

average between -62.35 % (min: +31.9 %, max: -100 %) and -95.21 % (min: -19.8 %, max: -100 %) of their

suitable bioclimatic niche by 2081-2100 period under the most optimistic (SSP126) and the worst-case

(SSP585) climate scenario, respectively. The severity of loss varies across species. More than half (32

species) are projected to experience a complete loss (100 %) of their suitable bioclimatic areas by the end of

the century for at least one of the SSP, putting them at elevated risk of extinction. *Delphinium montanum*,

Endressia pyrenaica, Silene borderei and *Onobrychis pyrenaica* appear particularly vulnerable, as models

project a total disappearance of their bioclimatic suitable areas as early as the 2021-2040 period, under all

SSPs, suggesting that these species are already present in unsuitable bioclimatic conditions. In contrast,

Pinguicula longifolia and *Linaria Bubanii* emerged as exceptions, showing consistent increase in suitable areas

with an average gain of +127.4 % (min: +95.5%, max: +181.63%) by 2081-2100 period. This suggests that

shifting climatic conditions may benefit some species, including endemic and threatened ones. The analysis

also highlights an accelerating trend in suitable area loss over time, particularly in more extreme scenarios

such as SSP370 and SSP585. Most species experience sharp declines after the 2041-2060 period, reflecting

the escalating impact of climate change as the century progresses (Figure 5). All suitability maps, covering

both current and future projections for all species, are available in Appendix D.

Spatial dynamics in bioclimatic niche suitability and upslope shift of bioclimatic hotspots

303   The spatial distribution of bioclimatic niche suitability for endemic Pyrenean plant species is projected to

304   undergo significant changes by the 2081-2100 period under climate change (Figure 6). Currently, bioclimatic

305   hotspots, where most species could find suitable bioclimatic conditions, are mainly above 1000 m, with a

306   maximum favorability of 0.73 (Figure 6a, top panel). By the 2081-2100 period, these hotspots are projected to

shrink in size and shift upward, with the extent of this change strongly linked to the emission scenario. Climate scenarios such as SSP126 show a relatively moderate reduction, with some areas persisting near current locations. In contrast, more extreme scenarios projected a pronounced contraction of suitable areas, becoming increasingly fragmented. Under SSP585, the maximum favorability value drops to 0.29, with suitable conditions restricted to areas above 2000 m in elevation (Figure 6a, lower panels).

The projected mean elevation of bioclimatic suitable areas for endemic Pyrenean plant species shows a marked upward shift by the end of the century under climate change (Figure 6b). Currently, mean elevation of suitable bioclimatic area ranges between 748 and 2269 m, depending on the species. By 2081-2100, the average elevation across species is projected to increase by +182.58 m. While the lower margin remains relatively stable (671 m), the upper margin is expected to reach 2779 m. Species exhibit variable projected elevation trajectories. By the 2081-2100 period, 20 of the 27 species that still have suitable bioclimatic conditions across all SSPs display a clear upward projected trend, with mean increases ranging from +0.26 % to +38.62 %. Subsets of species that may experience a reduction in their mean elevation could see decreases from -8.93% to -41.51 %. On average, species show a +19.49 % increase in elevation, while the decrease is about -23.44 %. In addition to these vertical shifts, species also display heterogeneous latitudinal responses by 2081-2100 period. Among the 27 species included, 13 are projected to shift northward (+0.2 to +47.3 km, mean +11.1 km), while 14 shift southward (-11.3 to -0.6 km, mean -4.7 km), resulting in an overall average shift of +2.91 km northward. These results emphasize the significant influence of warming climate on the spatial distribution of suitable bioclimatic areas, underscoring the increasing vulnerability of endemic Pyrenean flora to rising temperatures.

By 2081–2100, overlap between current and future suitable bioclimatic areas is generally low. Out of 236 species x scenario combinations, 177 (75%) fall in the lowest overlap class (0-25%). At the species level, 55 of the 59 endemics experience at least one scenario with less than 25% overlap. Notably, 38 species show complete loss of overlap (0%) under the most severe scenario (SSP585). This means that while 32 species entirely disappear from the projections, 6 still retain suitable climatic conditions, but in areas completely distinct from their current distribution. In contrast, only 7 species retain relatively high stability (≥75% overlap) under at least one scenario. These results highlight the drastic spatial reshuffling of suitable areas and the limited persistence of current bioclimatic ranges under future climates.

4 | Discussion

Trends in bioclimatic niche suitability under climate change

This study paints a concerning picture of the future for Pyrenean endemic flora under ongoing and projected climate change. By the 2081-2100 period, 57/59 of the studied species are projected to experience substantial reduction in suitable bioclimatic areas (on average -62.35 % to -95.21 % for SSP126 and SSP585 respectively), ranging from moderate contraction in some species to complete losses in others. These findings underline the severe threat posed by changing climate to narrowly distributed species and align with patterns projected in similar mountain regions, such as the Alps (DAGNINO *et al.*, 2020 ; DIRNBÖCK *et al.*, 2011 ; DULLINGER *et al.*, 2012 ; HÜLBER *et al.*, 2016 ; ROTA *et al.*, 2022), where endemics are acutely sensitive to rapid warming. Parallels with other mountain systems (MANES *et al.*, 2021) underscore the universal challenge that climate change poses to mountain endemism which is, moreover, amplified by topography and human pressures (Elsen et al., 2020). Interestingly, not all species follow the same trajectory. While most of them show severe bioclimatic niche contractions, *Pinguicula longifolia* and *Linaria bubanii* are the only endemics projected to consistently gain suitable bioclimatic areas under all scenarios, mainly on the northern slopes and in the southeastern Pyrenees. For *P. longifolia*, this may reflect preadaptation to warmer climates, as suggested by the occurrence of a related subspecies in the drier Massif Central. For *L. bubanii*, a scree-dwelling species of high altitudes, the strong microclimatic heterogeneity of its habitat may facilitate persistence across diverse conditions and promote expansion. Despite these projected gains, both species may remain vulnerable as establishment depends on strict ecological requirements and interactions with other biotic components under climate change. These outliers highlight that while some species face heightened risks of extinction, others may find new opportunities under climate change. It also underlines the need for species-specific conservation strategies that account for this ecological variability.

Beyond species-specific patterns, this study revealed notable upward shifts in mean elevations of suitable bioclimatic niches, exceeding 180 m by the 2081-2100 period. If such a trend had been consistent over the past century, it could help explain a part of the upward plant movement already observed in the Pyrenees (AMEZTEGUI *et al.*, 2016 ; AULLÓ-MAESTRO *et al.*, 2023 ; DELPOUVE *et al.*, 2025 ; LENOIR *et al.*, 2008 ; MARSHALL *et al.*, 2020 ; PAULI *et al.*, 2012) and is consistent with documented upward trends in other various mountain ecosystems (CHAN *et al.*, 2024 ; ISELI *et al.*, 2023). In parallel for the same period, latitudinal dynamics indicate an overall northward displacement of about +3 km. Overlap with current ranges is also very limited: 55 of the 59 endemic species retain less than 25% of their present distribution under at least one scenario, implying that many future suitable areas will occur entirely outside their current ranges. Combined with the projected upward shift, it suggests a profound spatial displacement of endemic species under ongoing warming. However, the pace at which species can track these changes remains uncertain, as local-scale factors, such as microclimatic

buffering, habitat discontinuities, or dispersal limitations, may slow or constrain their responses (MACLEAN & EARLY, 2023). Hotspots of suitable bioclimatic niches, currently located above 1000 m, are projected to shift above 2000 m. This shift drastically reduces bioclimatic suitable space available for endemic flora, as high elevations cover smaller geographic extent in the Pyrenees (ELSEN & TINGLEY, 2015). Many species risk being squeezed between warming lower slopes and shrinking refugia, exacerbating their risk of extinction. Bioclimatic hotspot maps revealed spatial patterns suggesting a future fragmentation of suitable areas, with favorable habitats increasingly confined to isolated mountaintops separated by valleys. This fragmentation likely reduces connectivity across the landscape, limiting the ability of many species to track climate change. In the Pyrenees, this issue is compounded by regional warming and aridification, already documented at a rate of 30 % faster than the global average over the past 50 years (OPCC, 2018). Combined with challenges already posed by dispersal barriers, topographic constraints, and competition from upslope-moving species, these factors intensify pressure on the native flora. The severe suitable bioclimatic area reductions projected under SSP370 and SSP585 scenarios after the 2041-2060 period is particularly alarming and suggests that environmental pressures will intensify in the second half of the century. The urgency of monitoring is reinforced by the identification of high vulnerability of northern Spanish flora, to the south of the Pyrenees, which is among the most threatened in Europe (ENGLER *et al.*, 2011).

Strengths and limitations of the modeling framework

The pipeline developed in this study represents a step forward in addressing mountain biodiversity questions at a broad geographic scale, particularly for analyzing climate-driven trends across multiple species. By integrating five algorithms (GLM, GAM, GBM, RF, and MXT) into an ensemble modeling framework, this approach minimizes model-specific biases and improves the reliability of projected species distributions. Moreover, the agreement with previous projections for an endemic species also analyzed here (e.g. *D. montanum*, Salvado *et al.*, 2022) suggests that our ensemble approach produces ecologically consistent outcomes. RF, MXT and GBM emerged as particularly strong performers, likely due to their capacity to handle small sample sizes and capture non-linear relationships between species and environmental variables (VALAVI *et al.*, 2022). The pipeline addresses challenges posed by sparse data, providing consistent and interpretable results for narrowly distributed species, as demonstrated by its ability to model 75% of the endemic species in a mountainous region. It includes tools to adjust pseudo-absence sampling to obtain stable predictions despite limited occurrence data. Another component helps optimize spatial cross-validation settings ensuring robust as possible model evaluation while maximizing the number of species that can be assessed and thus studied. These features make the approach suitable for rare or endemic species, whose restricted ranges and small

sample sizes could limit the applicability of standard SDMs procedures. To account for endemic Pyrenean

plant species excluded from this study, the pipeline could be applied case-by-case at a smaller spatial scale

(in regions of recorded occurrences). As such, it provides a valuable and adaptable framework for informing

conservation strategies across a diversity of endemic taxa. Moving forward, our pipeline could be extended to

species with broader distributions, adapted to biodiversity hotspots worldwide, and optimized for a more

extensive analysis across multiple species. It would provide deeper insights into species and communities'

response to climate change, focusing on richness patterns and diversity dynamics.

Despite these strengths, several SDMs limitations and considerations warrant caution when interpreting the

results (Collette et al., in press), particularly regarding methodological biases and decision-making processes

(see Sillero *et al.*, 2021; Bryn *et al.*, 2021; Leroy, 2023). This study presents three main considerations. First,

in large-scale studies where fine-resolution data are lacking, coarse-resolution bioclimatic data can still capture

general environmental trends, reduce computational costs, and minimize biases from inaccuracies in

occurrence data. However, by failing to capture fine-scale microhabitat heterogeneity in mountainous

environments, they may misestimate the estimated extent of habitat suitability (FRANKLIN *et al.*, 2013 ;

GOTTFRIED *et al.*, 2012 ; NADEAU *et al.*, 2017 ; RANDIN *et al.*, 2009 ; TRIVEDI *et al.*, 2008), although the overall

trends remain consistent (TSIFTSIS *et al.*, 2024). Second, projections beyond 2041-2060 period remain

uncertain due to increasing divergence between climate models, particularly for worst climate scenarios

(SSP370, SSP585) (GOBERVILLE *et al.*, 2015 ; TEBALDI *et al.*, 2021). Although the trajectory for most species

aligns across scenarios, notable variation exists in whether suitable bioclimatic niche disappears or persists to

a lesser extent, depending on pseudo-absence sets. Long-term monitoring and high-resolution climate data

are essential to validate projections and improve predictive tools. Third, the exclusion of biotic interactions

(e.g. competition, mutualism, pollination), and important abiotic variables (e.g. soil, hydrology, radiation, land

use) reduces the ecological realism of projections. Integrating such dynamics into SDMs would enhance the

accuracy of projections and offer a more nuanced understanding of species' responses to climate change

(COSENTINO *et al.*, 2023 ; LEE-YAW *et al.*, 2022 ; PREUSS & PADIAL, 2021).

Beyond these methodological considerations, SDMs also fail to capture complex ecological and evolutionary

processes shaping species' responses to environmental changes (WILLIAMS *et al.*, 2008). Expansion of

bioclimatic suitability areas does not guarantee population growth, as dispersal barriers, establishment

limitations, and biotic interactions may restrict species from fully occupying suitable areas. Likewise, suitability

contraction does not imply imminent range collapse, as species can persist temporarily in suboptimal

conditions before extinction, known as "extinction debt" (COTTO *et al.*, 2017 ; NOMOTO & ALEXANDER, 2021).

On the adaptation perspective, rapid environmental changes may outpace the rate at which species can adapt, facing the disappearance of their suitable habitat (COTTO *et al.*, 2017 ; JUMP & PEÑUELAS, 2005 ; SHAW & ETTERSON, 2012). Moreover, fragmented populations may experience divergent selective pressures, fostering local adaptation, often overlooked in SDMs. Integrating local adaptation, phenotypic plasticity (BENITO-GARZON *et al.*, 2019 ; VALLADARES *et al.*, 2014), and genomic data (HÄLLFORS *et al.*, 2016 ; KORT *et al.*, 2024) could improve projections by better accounting for populations' adaptive capacity under environmental stress. Despite these limitations, correlative SDMs based on bioclimatic variables remain valuable tools for generating insights and understanding species' ecological niches, anticipating distributional changes, and informing conservation strategies.

Conservation implications and Policy recommendation

From conservation and management perspectives, our findings underscore the urgent need for conservation strategies that address specific challenges faced by mountain flora in the Pyrenees. High altitude environments, where suitable bioclimatic conditions are expected to persist, represent critical bulwarks against climate change. These areas must be prioritized for legal protection as critical refugia for biodiversity preservation, since they are projected to concentrate the remaining favorable conditions for species by 2081-2100. To ensure the long-term relevance of protected areas, conservation planning must integrate future suitability and regulate potentially harmful activities in these refugia, such as infrastructure development or unregulated tourism, particularly above 2000 m. However, traditional grazing should be promoted, as it maintains open habitats that support many endemic species adapted to these environments. In parallel, the consistent upward shift of bioclimatic hotspots and their fragmentation call for the maintenance of altitudinal corridors to support species migration and gene flow, thereby mitigating the adverse effects of isolation. Mapping these corridors based on projected trajectories could help identify priority areas for land acquisition or land-use planning. For species projected to experience a partial lose their suitable bioclimatic niche, conservation interventions such as assisted migration and translocation may provide options to prevent extinction (FERRARINI *et al.*, 2016 ; GALLAGHER *et al.*, 2014). Evaluating candidate sites for reintroduction in newly suitable areas would facilitate the implementation of these actions. Moreover, integrating predictive modeling tools with field monitoring could enhance the precision of these interventions, ensuring that resources are allocated to areas with the greatest potential for success (ELLIOTT *et al.*, 2024 ; EYRE *et al.*, 2022). Critically vulnerable taxa such as *Delphinium montanum*, *Silene borderei*, and *Onobrychis pyrenaica*, for which models predict the disappearance of suitable conditions as early as 2021-2040 under all SSPs, should be urgently targeted. For these species, in situ strategies may need to be complemented by ex situ measures, such as

seed banking or cultivation in botanical gardens, which represent last-resort solution to preserve genetic material and avoid extinction. Developing emergency conservation plans for such high-risk taxa would improve coordination across institutions and ensure readiness for rapid intervention. In parallel, identifying microrefugia through fine-scale topographic analysis may uncover overlooked zones of persistence and bring new areas for assisted migration. Similarly, intensified demographic and phenological monitoring within residual populations would help assess whether remnant individuals are viable or demographically declining, thereby informing prioritization. Re-evaluating conservation status in light of projected niche loss would better guide adaptive management priorities. Many endemic species are likely to face imminent extinction risks that are not yet recognized in current conservation assessments. Including climate-driven projections in red list criteria could improve the alignment between legal protection and future vulnerability. National and regional conservation strategies should be updated to incorporate forward-looking criteria, ensuring that legal frameworks remain dynamic in the face of environmental change.

The transboundary distribution of Pyrenean species across France, Spain, and Andorra requires coordinated cross-border efforts to avoid fragmented conservation initiatives, such as inconsistent Natura 2000 habitat interpretation along this frontier (PRUD'HOMME & NINOT, 2018). Adaptive legal frameworks are essential to address these shifts, supported by collaborative efforts that align research, policy, and community engagement for enhanced conservation outcomes. Establishing a transnational conservation platform for endemic mountain species could help harmonize monitoring, protection, and restoration efforts across borders, and reduce disparities in species management. Translating scientific evidence into clear and actionable policy briefs can bridge the gap between research and implementation, fostering informed and, consequently, more effective decision-making (GUISAN et al., 2013). Understanding how climate change affects extinction risks and habitat availability is key to prioritizing conservation actions, especially as pressures intensify. This also requires combining ecological modeling, field experimentation, and long-term monitoring to refine conservation strategies and better address species redistribution and their effects on human well-being (BONEBRAKE et al., 2018 ; PECL et al., 2017). Ultimately, preserving mountain biodiversity in an era of rapid change will require collaboration among scientists, resource managers, and local communities, supported by predictive tools to capture the complex realities of a rapidly changing world.

Funding statement: This study received financial support from the Occitanie Region (Emergence 2024) and the Commissariat du Massif des Pyrénées (FNADT 2023), as well as from the Fédération de Recherche 2043, Energie & Environnement (UPVD/CNRS). It also received technical and financial support from the Interreg-POCTEFA program as part of the Floralab+ project (EFA024/01).

Acknowledgements: We thank Romain BERTRAND (Toulouse III University) for valuable comments that contributed to shaping our methodology.

Author Contributions : Conceptualization, J.A.M.B., N.C., V.H. ; **Data curation**, N.C. ; **Formal analysis**, N.C. ; **Funding Acquisition**, J.A.M.B., S.P., V.H. ; **Investigation**, N.C. ; **Methodology**, J.A.M.B., N.C., S.P. V.H.; **Project administration**, J.A.M.B., S.P., V.H. ; **Software**, N.C. ; **Supervision**, J.A.M.B., S.P., V.H ; **Validation**, J.A.M.B., N.C., S.P., V.H. ; **Visualization**, J.A.M.B., N.C., S.P., V.H. ; **Writing – original draft**, J.A.M.B., N.C., S.P., V.H. ; **Writing – review & editing**, J.A.M.B., N.C., S.P., V.H..

Data and code availability statement: The original data supporting the findings of this study are openly available from public domain resources, including GBIF, WorldClim, and iNaturalist. The code and data required to reproduce the results are accessible in the Zenodo repository at [DOI : 10.5281/zenodo.17412264] and DRYAD Further supporting details are available in the Supplementary Information.

Conflict of interest statement: The authors declare no conflicts of interest.

References

ALEXANDER J.M., CHALMANDRIER L., LENOIR J., BURGESS T.I., ESSL F., HAIDER S., KUEFFER C., MCDOUGALL K., MILBAU A., NUÑEZ M.A., PAUCHARD A., RABITSCH W., REW L.J., SANDERS N.J., PELLISSIER L. 2018. Lags in the response of mountain plant communities to climate change. *Global Change Biology* 24, 563-579. https://doi.org/10.1111/gcb.13976

ALEXANDER J.M., DIEZ J.M., LEVINE J.M. 2015. Novel competitors shape species' responses to climate change. *Nature* 525, 515-518. https://doi.org/10.1038/nature14952

AMEZTEGUI A., COLL L., BROTONS L., NINOT J.M. 2016. Land-use legacies rather than climate change are driving the recent upward shift of the mountain tree line in the Pyrenees. *Global Ecology and Biogeography* 25, 263-273. https://doi.org/10.1111/geb.12407

AULLÓ-MAESTRO I., GÓMEZ C., HERNÁNDEZ L., CAMARERO J.J., SÁNCHEZ-GONZÁLEZ M., CAÑELLAS I., VÁZQUEZ DE LA CUEVA A., MONTES F. 2023. Monitoring montane-subalpine forest ecotone in the Pyrenees through sequential forest inventories and Landsat imagery. *Annals of Forest Science* 80, 32. https://doi.org/10.1186/s13595-023-01198-4

BARBET-MASSIN M., JIGUET F., ALBERT C.H., THUILLER W. 2012. Selecting pseudo-absences for species distribution models: how, where and how many? *Methods in Ecology and Evolution* 3, 327-338. https://doi.org/10.1111/j.2041-210X.2011.00172.x

BARBOSA A.M. 2015. fuzzySim: applying fuzzy logic to binary similarity indices in ecology. *Methods in Ecology and Evolution* 6, 853-858. https://doi.org/10.1111/2041-210X.12372

BENITO-GARZON M., ROBSON T., HAMPE A. 2019. ΔTraitSDMs: species distribution models that account for local adaptation and phenotypic plasticity. *New Phytologist* 222, 1757-1765. https://doi.org/10.1111/nph.15716

BIDEGARAY-BATISTA L., SÁNCHEZ-GRACIA A., SANTULLI G., MAIORANO L., GUISAN A., VOGLER A.P., ARNEDO M.A.
2016. Imprints of multiple glacial refugia in the Pyrenees revealed by phylogeography and
palaeodistribution modelling of an endemic spider. *Molecular Ecology* 25, 2046-2064.
https://doi.org/10.1111/mec.13585

BIJLSMA R., LOESCHCKE V. 2012. Genetic erosion impedes adaptive responses to stressful environments.
Evolutionary Applications 5, 117-129. https://doi.org/10.1111/j.1752-4571.2011.00214.x

BONEBRAKE T.C., BROWN C.J., BELL J.D., BLANCHARD J.L., CHAUVENET A., CHAMPION C., CHEN I., CLARK T.D.,
COLWELL R.K., DANIELSEN F., DELL A.I., DONELSON J.M., EVENGÅRD B., FERRIER S., FRUSHER S.,
GARCIA R.A., GRIFFIS R.B., HOBDAY A.J., JARZYNA M.A., LEE E., LENOIR J., LINNETVED H., MARTIN V.Y.,
MCCORMACK P.C., MCDONALD J., MCDONALD-MADDEN E., MITCHELL N., MUSTONEN T., PANDOLFI J.M.,
PETTORELLI N., POSSINGHAM H., PULSIFER P., REYNOLDS M., SCHEFFERS B.R., SORTE C.J.B.,
STRUGNELL J.M., TUANMU M., TWINAME S., VERGÉS A., VILLANUEVA C., WAPSTRA E., WERNBERG T.,
PECL G.T. 2018. Managing consequences of climate-driven species redistribution requires integration
of ecology, conservation and social science. *Biological Reviews* 93, 284-305.
https://doi.org/10.1111/brv.12344

BROUSSIN J., MOUCHET M., GOBERVILLE E. 2024. Generating pseudo-absences in the ecological space
improves the biological relevance of response curves in species distribution models. *Ecological*
Modelling 498, 110865. https://doi.org/10.1016/j.ecolmodel.2024.110865

BRYN A., BEKKBY T., RINDE E., GUNDERSEN H., HALVORSEN R. 2021. Reliability in Distribution Modeling—A
Synthesis and Step-by-Step Guidelines for Improved Practice. *Front. Ecol. Evol.* 9, 658713.
https://doi.org/10.3389/fevo.2021.658713

BUENO DE MESQUITA C.P., TILLMANN L.S., BERNARD C.D., ROSEMOND K.C., MOLOTCH N.P., SUDING K.N. 2018.
Topographic heterogeneity explains patterns of vegetation response to climate change (1972–2008)
across a mountain landscape, Niwot Ridge, Colorado. *Arctic, Antarctic, and Alpine Research* 50,
e1504492. https://doi.org/10.1080/15230430.2018.1504492

BUISSON L., THUILLER W., CASAJUS N., LEK S., GRENOUILLET G. 2010. Uncertainty in ensemble forecasting of
species distribution. *Global Change Biology* 16, 1145-1157. https://doi.org/10.1111/j.1365-
2486.2009.02000.x

CAMARERO J.J., GUTIERREZ E., FORTIN M. 2006. Spatial patterns of plant richness across treeline ecotones in
the Pyrenees reveal different locations for richness and tree cover boundaries. *Global Ecology and*
Biogeography 15, 182-191. https://doi.org/10.1111/j.1466-822X.2006.00211.x

CARROLL C., LAWLER J.J., ROBERTS D.R., HAMANN A. 2015. Biotic and Climatic Velocity Identify Contrasting
Areas of Vulnerability to Climate Change. *PLOS ONE* 10, e0140486.
https://doi.org/10.1371/journal.pone.0140486

CARROLL C., MAHONY C.R. 2025. Sources of uncertainty in estimation of climate velocity and their implications
for ecological and conservation applications. *Conservation Science and Practice* 7, e13296.
https://doi.org/10.1111/csp2.13296

CGIAR-CSI 2018. SRTM 90m DEM Digital Elevation Database [WWW Document]. URL
https://srtm.csi.cgiar.org/.

CHAN W.-P., LENOIR J., MAI G.-S., KUO H.-C., CHEN I.-C., SHEN S.-F. 2024. Climate velocities and species
tracking in global mountain regions. *Nature* 629, 114-120. https://doi.org/10.1038/s41586-024-07264-
9

CHARBONNEL A., LAFFAILLE P., BIFFI M., BLANC F., MAIRE A., NEMOZ M., SANCHEZ-PEREZ J.M., SAUVAGE S.,
BUISSON L. 2016. Can Recent Global Changes Explain the Dramatic Range Contraction of an
Endangered Semi-Aquatic Mammal Species in the French Pyrenees? *PLOS ONE* 11, e0159941.
https://doi.org/10.1371/journal.pone.0159941

COLLETTE N., FRAISSE J., BERTRAND R., PINEL S., DELORME-HINOUX V., MURIENNE J., BERTRAND A.M.J. in press.
Modèles de distribution d'espèces : enjeux, défis et perspectives pour une conservation éclairée. *Le*
Monde des Plantes.

COSENTINO F., SEAMARK E.C.J., VAN CAKENBERGHE V., MAIORANO L. 2023. Not only climate: The importance
of biotic interactions in shaping species distributions at macro scales. *Ecology and Evolution* 13,
e9855. https://doi.org/10.1002/ece3.9855

COTTO O., WESSELY J., GEORGES D., KLONNER G., SCHMID M., DULLINGER S., THUILLER W., GUILLAUME F. 2017.
A dynamic eco-evolutionary model predicts slow response of alpine plants to climate warming. *Nat*
Commun 8, 15399. https://doi.org/10.1038/ncomms15399

CUADRAT J.M., SERRANO-NOTIVOLI R., PROHOM M., CUNILLERA J., TEJEDOR E., SAZ M.Á., DE LUIS M., LLABRES-
BRUSTENGA A., SOUBEYROUX J.-M. 2024. Climate of the Pyrenees: Extremes indices and long-term
trends. *Science of The Total Environment* 933, 173052.
https://doi.org/10.1016/j.scitotenv.2024.173052

DAGNINO D., GUERRINA M., MINUTO L., MARIOTTI M.G., MÉDAIL F., CASAZZA G. 2020. Climate change and the
future of endemic flora in the South Western Alps: relationships between niche properties and
extinction risk. *Reg Environ Change* 20, 121. https://doi.org/10.1007/s10113-020-01708-4

DAINESE M., CREPAZ H., BOTTARIN R., FONTANA V., GUARIENTO E., HILPOLD A., OBOJES N., PANICCIA C., SCOTTI
A., SEEBER J., STEINWANDTER M., TAPPEINER U., NIEDRIST G. 2024. Global change experiments in
mountain ecosystems: A systematic review. *Ecological Monographs* 94, e1632.
https://doi.org/10.1002/ecm.1632

DALIBARD M., LAFFAILLE P., SANCHEZ-PEREZ J.-M., SAUVAGE S., BUISSON L. 2021. Accounting for flow
intermittence in freshwater species distribution modelling. *Ecohydrology* 14, e2346.
https://doi.org/10.1002/eco.2346

DELPOUVE N., BERGÈS L., DUPOUEY J.-L., CHAUCHARD S., LEROY N., THIRION E., RATHGEBER C.B.K. 2025. Long-
term forest-line dynamics in the French Pyrenees: an accelerating upward shift related to forest
context, global warming and pastoral abandonment. https://doi.org/10.5194/egusphere-2024-4099

DE POUS P., MONTORI A., ORRIOLS F., SANUY D. 2015. Range contraction and loss of genetic variation of the
Pyrenean endemic newt *Calotriton asper* due to climate change. *Regional Environmental Change* 16,
995-1009. https://doi.org/10.1007/s10113-015-0804-3

DI MUSCIANO M., DI CECCO V., BARTOLUCCI F., CONTI F., FRATTAROLI A.R., DI MARTINO L. 2020. Dispersal ability
of threatened species affects future distributions. *Plant Ecol* 221, 265-281.
https://doi.org/10.1007/s11258-020-01009-0

DIRNBÖCK T., ESSL F., RABITSCH W. 2011. Disproportional risk for habitat loss of high-altitude endemic species
under climate change. *Global Change Biology* 17, 990-996. https://doi.org/10.1111/j.1365-
2486.2010.02266.x

DULLINGER S., GATTRINGER A., THUILLER W., MOSER D., ZIMMERMANN N.E., GUISAN A., WILLNER W., PLUTZAR
C., LEITNER M., MANG T., CACCIANIGA M., DIRNBÖCK T., ERTL S., FISCHER A., LENOIR J., SVENNING J.-
C., PSOMAS A., SCHMATZ D.R., SILC U., VITTOZ P., HÜLBER K. 2012. Extinction debt of high-mountain

plants under twenty-first-century climate change. *Nature Clim Change* 2, 619-622.
https://doi.org/10.1038/nclimate1514

EICHEL J., STOFFEL M., WIPF S. 2023. Go or grow? Feedbacks between moving slopes and shifting plants in
high mountain environments. *Progress in Physical Geography: Earth and Environment* 47, 967-985.
https://doi.org/10.1177/03091333231193844

ELITH J., KEARNEY M., PHILLIPS S. 2010. The art of modelling range-shifting species. *Methods in Ecology and*
Evolution 1, 330-342. https://doi.org/10.1111/j.2041-210X.2010.00036.x

ELLIOTT C.P., TOMLINSON S., LEWANDROWSKI W., MILLER B.P. 2024. Species distribution and habitat attributes
guide translocation planning of a threatened short-range endemic plant. *Global Ecology and*
Conservation 51, e02915. https://doi.org/10.1016/j.gecco.2024.e02915

ELSEN P.R., MONAHAN W.B., MERENLENDER A.M. 2020. Topography and human pressure in mountain ranges
alter expected species responses to climate change. *Nat Commun* 11, 1974.
https://doi.org/10.1038/s41467-020-15881-x

ELSEN, TINGLEY M. 2015. Global mountain topography and the fate of montane species under climate change.
Nature Climate Change 5, 772-776. https://doi.org/10.1038/nclimate2656

ENGLER R., RANDIN C.F., THUILLER W., DULLINGER S., ZIMMERMANN N.E., ARAÚJO M.B., PEARMAN P.B., LE LAY
G., PIEDALLU C., ALBERT C.H., CHOLER P., COLDEA G., DE LAMO X., DIRNBÖCK T., GÉGOUT J.-C.,
GÓMEZ-GARCÍA D., GRYTNES J.-A., HEEGAARD E., HØISTAD F., NOGUÉS-BRAVO D., NORMAND S.,
PUŞCAŞ M., SEBASTIÀ M.-T., STANISCI A., THEURILLAT J.-P., TRIVEDI M.R., VITTOZ P., GUISAN A. 2011.
21st century climate change threatens mountain flora unequally across Europe. *Global Change*
Biology 17, 2330-2341. https://doi.org/10.1111/j.1365-2486.2010.02393.x

EYRE A.C., BRISCOE N.J., HARLEY D.K.P., LUMSDEN L.F., MCCOMB L.B., LENTINI P.E. 2022. Using species
distribution models and decision tools to direct surveys and identify potential translocation sites for a
critically endangered species. *Diversity and Distributions* 28, 700-711.
https://doi.org/10.1111/ddi.13469

FERRARINI A., SELVAGGI A., ABELI T., ALATALO J.M., ORSENIGO S., GENTILI R., ROSSI G. 2016. Planning for
assisted colonization of plants in a warming world. *Sci Rep* 6, 28542.
https://doi.org/10.1038/srep28542

FICK S.E., HIJMANS R.J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas.
International Journal of Climatology 37, 4302-4315. https://doi.org/10.1002/joc.5086

FRANKHAM R. 2005. Genetics and extinction. *Biological Conservation* 126, 131-140.
https://doi.org/10.1016/j.biocon.2005.05.002

FRANKLIN J. 2023. Species distribution modelling supports the study of past, present and future
biogeographies. *Journal of Biogeography* 50, 1533-1545. https://doi.org/10.1111/jbi.14617

FRANKLIN J., DAVIS F.W., IKEGAMI M., SYPHARD A.D., FLINT L.E., FLINT A.L., HANNAH L. 2013. Modeling plant
species distributions under future climates: how fine scale do climate projections need to be? *Global*
Change Biology 19, 473-483. https://doi.org/10.1111/gcb.12051

FREEMAN B.G., LEE-YAW J.A., SUNDAY J.M., HARGREAVES A.L. 2018. Expanding, shifting and shrinking: The
impact of global warming on species' elevational distributions. *Global Ecol Biogeogr* 27, 1268-1276.
https://doi.org/10.1111/geb.12774

GALLAGHER R., HOGBIN P., HANCOCK N. 2014. Assisted colonization as a climate change adaptation tool.
Austral Ecology 40, 12-20. https://doi.org/10.1111/aec.12163

GARCIA M.B., MIRANDA H., PIZARRO M., FONT X., ROQUET C., GONZALEZ-SAMPERIZ P. 2022. Habitats hold an
evolutionary signal of past climatic refugia. *Biodivers Conserv* 31, 1665-1688.
https://doi.org/10.1007/s10531-022-02419-4

GBIF.ORG 2025. Occurrence Download. https://doi.org/10.15468/DL.R84YM3

GEPPERT C., BERTOLLI A., PROSSER F., MARINI L. 2023. Red-listed plants are contracting their elevational range
faster than common plants in the European Alps. *Proceedings of the National Academy of Sciences*
120, e2211531120. https://doi.org/10.1073/pnas.2211531120

GLAD A., MALLARD F. 2022. Alpine marmot (*Marmota marmota*) distribution evolution under climate change:
The use of species distribution models at a local scale in the western Pyrenees massif (France).
Ecological Informatics 69, 101646. https://doi.org/10.1016/j.ecoinf.2022.101646

GOBERVILLE E., BEAUGRAND G., HAUTEKÈETE N.-C., PIQUOT Y., LUCZAK C. 2015. Uncertainties in the projection
of species distributions related to general circulation models. *Ecology and Evolution* 5, 1100-1116.
https://doi.org/10.1002/ece3.1411

GÓMEZ D., GARCÍA M.B., FONT CASTELL X., AIZPURU OIARBIDE I. 2017. Distribución espacial y análisis ambiental
de la flora vascular de los Pirineos. *Pirineos* 172, 028. https://doi.org/10.3989/pirineos.2017.172003

GOODWIN K., CHARDON N., PRADHAN K., LAMBERS J., ANGERT A. 2025. Lagged climate-driven range shifts at
species' leading, but not trailing, range edges revealed by multispecies seed addition experiment.
Ecography e07331. https://doi.org/10.1111/ecog.07331

GOTTFRIED M., PAULI H., FUTSCHIK A., AKHALKATSI M., BARANČOK P., BENITO ALONSO J.L., COLDEA G., DICK J.,
ERSCHBAMER B., FERNÁNDEZ CALZADO M.R., KAZAKIS G., KRAJČI J., LARSSON P., MALLAUN M.,
MICHELSEN O., MOISEEV D., MOISEEV P., MOLAU U., MERZOUKI A., NAGY L., NAKHUTSRISHVILI G.,
PEDERSEN B., PELINO G., PUSCAS M., ROSSI G., STANISCI A., THEURILLAT J.-P., TOMASELLI M., VILLAR
L., VITTOZ P., VOGIATZAKIS I., GRABHERR G. 2012. Continent-wide response of mountain vegetation to
climate change. *Nature Clim Change* 2, 111-115. https://doi.org/10.1038/nclimate1329

GUEGUEN M., BLANCHETEAU H., THUILLER W. 2025. biomod2: Ensemble Platform for Species Distribution
Modeling https://biomodhub.github.io/biomod2/.

GUISAN A., TINGLEY R., BAUMGARTNER J.B., NAUJOKAITIS-LEWIS I., SUTCLIFFE P.R., TULLOCH A.I.T., REGAN T.J.,
BROTONS L., MCDONALD-MADDEN E., MANTYKA-PRINGLE C., MARTIN T.G., RHODES J.R., MAGGINI R.,
SETTERFIELD S.A., ELITH J., SCHWARTZ M.W., WINTLE B.A., BROENNIMANN O., AUSTIN M., FERRIER S.,
KEARNEY M.R., POSSINGHAM H.P., BUCKLEY Y.M. 2013. Predicting species distributions for
conservation decisions. *Ecology Letters* 16, 1424-1435. https://doi.org/10.1111/ele.12189

GUTIERREZ F., GARCIA M., LEO MONTES M., FERNANDEZ-MAZUECOS M., LAMRANI ALAOUI M., TERRAB A., VARGAS
P. 2022. The Pyrenees as a cradle of plant diversity: phylogeny, phylogeography and niche modeling
of *Saxifraga longifolia*. *Journal of Systematics and Evolution* 61, 253-272.
https://doi.org/10.1111/jse.12917

HÄLLFORS M.H., LIAO J., DZURISIN J., GRUNDEL R., HYVÄRINEN M., TOWLE K., WU G.C., HELLMANN J.J. 2016.
Addressing potential local adaptation in species distribution models: implications for conservation
under climate change. *Ecological Applications* 26, 1154-1169. https://doi.org/10.1890/15-0926

HÜLBER K., WESSELY J., GATTRINGER A., MOSER D., KUTTNER M., ESSL F., LEITNER M., WINKLER M., ERTL S.,
WILLNER W., KLEINBAUER I., SAUBERER N., MANG T., ZIMMERMANN N.E., DULLINGER S. 2016.

Uncertainty in predicting range dynamics of endemic alpine plants under climate warming. *Global*
Change Biology 22, 2608-2619. https://doi.org/10.1111/gcb.13232

INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE 2022. Climate Change 2022 – Impacts, Adaptation and
Vulnerability: Working Group II Contribution to the Sixth Assessment Report of the Intergovernmental
Panel on Climate Change, 1re ed. Cambridge University Press.
https://doi.org/10.1017/9781009325844

INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE (IPCC) (Éd.) 2023. Mountains, in: Climate Change 2022 –
Impacts, Adaptation and Vulnerability: Working Group II Contribution to the Sixth Assessment Report
of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, p.
2273-2318. https://doi.org/10.1017/9781009325844.022

ISELI E., CHISHOLM C., LENOIR J., HAIDER S., SEIPEL T., BARROS A., HARGREAVES A.L., KARDOL P., LEMBRECHTS
J.J., MCDOUGALL K., RASHID I., RUMPF S.B., ARÉVALO J.R., CAVIERES L., DAEHLER C., DAR P.A.,
ENDRESS B., JAKOBS G., JIMÉNEZ A., KÜFFER C., MIHOC M., MILBAU A., MORGAN J.W., NAYLOR B.J.,
PAUCHARD A., RATIER BACKES A., RESHI Z.A., REW L.J., RIGHETTI D., SHANNON J.M., VALENCIA G.,
WALSH N., WRIGHT G.T., ALEXANDER J.M. 2023. Rapid upwards spread of non-native plants in
mountains across continents. *Nat Ecol Evol* 7, 405-413. https://doi.org/10.1038/s41559-022-01979-6

JUMP A.S., PEÑUELAS J. 2005. Running to stand still: adaptation and the response of plants to rapid climate
change. *Ecology Letters* 8, 1010-1020. https://doi.org/10.1111/j.1461-0248.2005.00796.x

KASS J.M., SMITH A.B., WARREN D.L., VIGNALI S., SCHMITT S., AIELLO-LAMMENS M.E., ARLÉ E., MÁRCIA BARBOSA
A., BROENNIMANN O., COBOS M.E., GUÉGUEN M., GUISAN A., MEROW C., NAIMI B., NOBIS M.P., ONDO
I., OSORIO-OLVERA L., OWENS H.L., PINILLA-BUITRAGO G.E., SÁNCHEZ-TAPIA A., THUILLER W., VALAVI
R., VELAZCO S.J.E., ZIZKA A., ZURELL D. 2024. Achieving higher standards in species distribution
modeling by leveraging the diversity of available software. *Ecography* e07346.
https://doi.org/10.1111/ecog.07346

KELLNER J.R., KENDRICK J., SAX D.F. 2023. High-velocity upward shifts in vegetation are ubiquitous in
mountains of western North America. *PLOS Climate* 2, e0000071.
https://doi.org/10.1371/journal.pclm.0000071

KNIGHT J. 2022. Scientists' warning of the impacts of climate change on mountains. *PeerJ* 10, e14253.
https://doi.org/10.7717/peerj.14253

KÖRNER C., HILTBRUNNER E. 2021. Why Is the Alpine Flora Comparatively Robust against Climatic Warming?
Diversity 13, 383. https://doi.org/10.3390/d13080383

KÖRNER C., PAULSEN J., SPEHN E.M. 2011. A definition of mountains and their bioclimatic belts for global
comparisons of biodiversity data. *Alp Botany* 121, 73-78. https://doi.org/10.1007/s00035-011-0094-4

KORT H.D., BLANCHET S., LENOIR J., MANEL S., MEERBEEK K.V. 2024. Species distribution models: a genomic
perspective without limits. Authorea. https://doi.org/10.22541/au.172462796.66134085/v1

KOUBA Y., ALADOS C.L., BUENO C.G. 2011. Effects of abiotic and anthropogenic factors on the spatial
distribution of *Quercus faginea* in the Spanish Central Pyrenees. *Plant Ecol* 212, 999-1007.
https://doi.org/10.1007/s11258-010-9880-0

LAMOREUX J.F., MORRISON J.C., RICKETTS T.H., OLSON D.M., DINERSTEIN E., MCKNIGHT M.W., SHUGART H.H.
2006. Global tests of biodiversity concordance and the importance of endemism. *Nature* 440, 212-214.
https://doi.org/10.1038/nature04291

LAWLOR J.A., COMTE L., GRENOUILLET G., LENOIR J., BAECHER J.A., BANDARA R.M.W.J., BERTRAND R., CHEN
I.-C., DIAMOND S.E., LANCASTER L.T., MOORE N., MURIENNE J., OLIVEIRA B.F., PECL G.T., PINSKY M.L.,
ROLLAND J., RUBENSTEIN M., SCHEFFERS B.R., THOMPSON L.M., VAN AMEROM B., VILLALOBOS F.,
WEISKOPF S.R., SUNDAY J. 2024. Mechanisms, detection and impacts of species redistributions under
climate change. *Nat Rev Earth Environ* 5, 351-368. https://doi.org/10.1038/s43017-024-00527-z

LEE-YAW J.A., MCCUNE J.L., PIRONON S., SHETH S.N. 2022. Species distribution models rarely predict the
biology of real populations. *Ecography* 45, e05877. https://doi.org/10.1111/ecog.05877

LENOIR J., GEGOUT J.-C., MARQUET P., RUFFRAY P., BRISSE H. 2008. A Significant Upward Shift in Plant Species
Optimum Elevation During the 20th Century. *Science* 320, 1768-1771.
https://doi.org/10.1126/science.1156831

LEROY B. 2023. Choosing presence-only species distribution models. *Journal of Biogeography* 50, 247-250.
https://doi.org/10.1111/jbi.14505

LI D., WU S., LIU L., ZHANG Y., LI S. 2018. Vulnerability of the global terrestrial ecosystems to climate change.
Global Change Biology 24, 4095-4106. https://doi.org/10.1111/gcb.14327

LIU C., NEWELL G., WHITE M. 2016. On the selection of thresholds for predicting species occurrence with
presence-only data. *Ecology and Evolution* 6, 337-348. https://doi.org/10.1002/ece3.1878

LIU C., NEWELL G., WHITE M., MACHUNTER J. 2025. Improving the estimation of the Boyce index using statistical
smoothing methods for evaluating species distribution models with presence-only data. *Ecography*
2025, e07218. https://doi.org/10.1111/ecog.07218

LIU C., WHITE M., NEWELL G. 2013. Selecting thresholds for the prediction of species occurrence with presence-
only data. *Journal of Biogeography* 40, 778-789. https://doi.org/10.1111/jbi.12058

MACLEAN I.M.D., EARLY R. 2023. Macroclimate data overestimate range shifts of plants in response to climate
change. *Nat. Clim. Chang.* 13, 484-490. https://doi.org/10.1038/s41558-023-01650-3

MANES S., COSTELLO M.J., BECKETT H., DEBNATH A., DEVENISH-NELSON E., GREY K.-A., JENKINS R., KHAN T.M.,
KIESSLING W., KRAUSE C., MAHARAJ S.S., MIDGLEY G.F., PRICE J., TALUKDAR G., VALE M.M. 2021.
Endemism increases species' climate change risk in areas of global biodiversity importance. *Biological*
Conservation 257, 109070. https://doi.org/10.1016/j.biocon.2021.109070

MARMION M., PARVIAINEN M., LUOTO M., HEIKKINEN R.K., THUILLER W. 2009. Evaluation of consensus methods
in predictive species distribution modelling. *Diversity and Distributions* 15, 59-69.
https://doi.org/10.1111/j.1472-4642.2008.00491.x

MARSHALL L., PERDIJK F., DENDONCKER N., KUNIN W., ROBERTS S., BIESMEIJER J. 2020. Bumblebees moving
up: Shifts in elevation ranges in the Pyrenees over 115 years. *Proceedings of the Royal Society B:*
Biological Sciences 287, 20202201. https://doi.org/10.1098/rspb.2020.2201

MARTíNEZ I., GONZÁLEZ-TABOADA F., WIEGAND T., CAMARERO J.J., GUTIÉRREZ E. 2012. Dispersal limitation and
spatial scale affect model based projections of *Pinus uncinata* response to climate change in the
Pyrenees. *Global Change Biology* 18, 1714-1724. https://doi.org/10.1111/j.1365-2486.2012.02660.x

MATEO R.G., CROAT T.B., FELICíSIMO Á.M., MUÑOZ J. 2010. Profile or group discriminative techniques?
Generating reliable species distribution models using pseudo-absences and target-group absences
from natural history collections. *Diversity and Distributions* 16, 84-94. https://doi.org/10.1111/j.1472-
4642.2009.00617.x

MATHUR S., TOMEČEK J.M., TARANGO-ARÁMBULA L.A., PEREZ R.M., DEWOODY J.A. 2023. An evolutionary perspective on genetic load in small, isolated populations as informed by whole genome resequencing and forward-time simulations. *Evolution* 77, 690-704. https://doi.org/10.1093/evolut/qpac061

MCDERMOTT M.B.A., ZHANG H., HANSEN L.H., ANGELOTTI G., GALLIFANT J. 2025. A Closer Look at AUROC and AUPRC under Class Imbalance. https://doi.org/10.48550/arXiv.2401.06091

MORGAN J.W., VENN S.E. 2017. Alpine plant species have limited capacity for long-distance seed dispersal. *Plant Ecology* 218, 813-819. https://doi.org/10.1007/s11258-017-0731-0

NADEAU C.P., URBAN M.C., BRIDLE J.R. 2017. Coarse climate change projections for species living in a fine-scaled world. *Global Change Biology* 23, 12-24. https://doi.org/10.1111/gcb.13475

NI M., VELLEND M. 2024. Soil properties constrain forest understory plant distributions along an elevation gradient. *Philosophical Transactions of the Royal Society B* 379. https://doi.org/10.1098/rstb.2023.0373

NOMOTO H.A., ALEXANDER J.M. 2021. Drivers of local extinction risk in alpine plants under warming climate. *Ecology letters* 24, 1157. https://doi.org/10.1111/ele.13727

NORBERG A., ABREGO N., BLANCHET F.G., ADLER F.R., ANDERSON B.J., ANTTILA J., ARAÚJO M.B., DALLAS T., DUNSON D., ELITH J., FOSTER S.D., FOX R., FRANKLIN J., GODSOE W., GUISAN A., O'HARA B., HILL N.A., HOLT R.D., HUI F.K.C., HUSBY M., KÁLÁS J.A., LEHIKOINEN A., LUOTO M., MOD H.K., NEWELL G., RENNER I., ROSLIN T., SOININEN J., THUILLER W., VANHATALO J., WARTON D., WHITE M., ZIMMERMANN N.E., GRAVEL D., OVASKAINEN O. 2019. A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecological Monographs* 89, e01370. https://doi.org/10.1002/ecm.1370

O'NEILL B.C., KRIEGLER E., EBI K.L., KEMP-BENEDICT E., RIAHI K., ROTHMAN D.S., VAN RUIJVEN B.J., VAN VUUREN D.P., BIRKMANN J., KOK K., LEVY M., SOLECKI W. 2017. The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Global Environmental Change* 42, 169-180. https://doi.org/10.1016/j.gloenvcha.2015.01.004

OPCC 2018. Le changement climatique dans les Pyrénées : impact, vulnérabilités et adaptation : bases de connaissances pour la future stratégie d'adaptation au changement climatique dans les Pyrénées. ISBN: 978-84-09-06268-3

PAULI H., GOTTFRIED M., DULLINGER S., ABDALADZE O., AKHALKATSI M., BENITO ALONSO J.L., COLDEA G., DICK J., ERSCHBAMER B., CALZADO F., GHOSN D., HOLTEN J., KANKA R., KAZAKIS G., KOLLAR J., LARSSON P., MOISEEV P., MOISEEV D., MOLAU U., GRABHERR G. 2012. Recent Plant Diversity Changes on Europe's Mountain Summits. *Science (New York, N.Y.)* 336, 353-5. https://doi.org/10.1126/science.1219033

PEARSE W.D., DAVIES T.J. 2019. The realised velocity of climate change reveals remarkable idiosyncrasy of species' distributional shifts. https://doi.org/10.1101/823930

PECL G.T., ARAÚJO M.B., BELL J.D., BLANCHARD J., BONEBRAKE T.C., CHEN I.-C., CLARK T.D., COLWELL R.K., DANIELSEN F., EVENGÅRD B., FALCONI L., FERRIER S., FRUSHER S., GARCIA R.A., GRIFFIS R.B., HOBDAY A.J., JANION-SCHEEPERS C., JARZYNA M.A., JENNINGS S., LENOIR J., LINNETVED H.I., MARTIN V.Y., MCCORMACK P.C., MCDONALD J., MITCHELL N.J., MUSTONEN T., PANDOLFI J.M., PETTORELLI N., POPOVA E., ROBINSON S.A., SCHEFFERS B.R., SHAW J.D., SORTE C.J.B., STRUGNELL J.M., SUNDAY J.M., TUANMU M.-N., VERGÉS A., VILLANUEVA C., WERNBERG T., WAPSTRA E., WILLIAMS S.E. 2017. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* 355, eaai9214. https://doi.org/10.1126/science.aai9214

PEPIN, ARNONE E., GOBIET A., HASLINGER K., KOTLARSKI S., SERAFIN S., TERZAGO S., THORNTON J.M., VUILLE
M., ADLER C. 2022. Climate Changes and Their Elevational Patterns in the Mountains of the World.
Reviews of Geophysics 60, e2020RG000730. https://doi.org/10.1029/2020RG000730

PEPIN, BRADLEY R., DIAZ H., BARAER M., CÁCERES B., FORSYTHE N., FOWLER H., GREENWOOD G., HASHMI M.,
LIU X., MILLER J., NING L., OHMURA A., PALAZZI E., RANGWALA I., SCHÖNER W., SEVERSKIY I.,
SHAHGEDANOVA M., WANG M., YANG D. 2015. Elevation-dependent warming in mountain regions of
the world. *Nature Climate Change* 5, 424-430. https://doi.org/10.1038/nclimate2563

PEREZ-GARCIA N., CASTELL X., FERRE A., CARRERAS J. 2013. Drastic reduction in the potential habitats for
alpine and subalpine vegetation in the Pyrenees due to twenty-first-century climate change. *Regional
Environmental Change* 13, 1157-1169. https://doi.org/10.1007/s10113-013-0427-5

PINTO A.V., HANSSON B., PATRAMANIS I., MORALES H.E., VAN OOSTERHOUT C. 2024. The impact of habitat loss
and   population   fragmentation   on   genomic   erosion.   *Conserv   Genet*   25,   49-57.
https://doi.org/10.1007/s10592-023-01548-9

PIRONON S., GOMEZ D., FONT X., GARCIA M.B. 2022. Living at the limit in the Pyrenees: Peripheral and endemic
plants are rare but underrepresented in protection lists. *Diversity and Distributions* 28, 930-942.
https://doi.org/10.1111/ddi.13487

PREUSS G., PADIAL A.A. 2021. Increasing reality of species distribution models of consumers by including its
food   resources.   *Neotropical   Biology   and   Conservation*   16,   411-425.
https://doi.org/10.3897/neotropical.16.e64892

PRUD'HOMME F., NINOT J.M. 2018. Interprétation de la Directive habitats sur des écosystèmes pyrénéens
emblématiques : les lacs de montagne. Conservatoire Botanique National des Pyrénées et de Midi-
Pyrénées & Universitat de Barcelona.

QAZI A.W., SAQIB Z., ZAMAN-UL-HAQ M. 2022. Trends in species distribution modelling in context of rare and
endemic plants: a systematic review. *Ecological Processes* 11, 40. https://doi.org/10.1186/s13717-
022-00384-y

QIXIANG W., WANG M., FAN X. 2018. Seasonal patterns of warming amplification of high-elevation stations
across the globe. *International Journal of Climatology* 38, 3466-3473. https://doi.org/10.1002/joc.5509

RAHBEK C., BORREGAARD M.K., ANTONELLI A., COLWELL R.K., HOLT B.G., NOGUES-BRAVO D., RASMUSSEN
C.M.Ø., RICHARDSON K., ROSING M.T., WHITTAKER R.J., FJELDSÅ J. 2019a. Building mountain
biodiversity: Geological   and   evolutionary   processes.   *Science*   365,   1114-1119.
https://doi.org/10.1126/science.aax0151

RAHBEK C., BORREGAARD M.K., COLWELL R.K., DALSGAARD B., HOLT B.G., MORUETA-HOLME N., NOGUES-BRAVO
D., WHITTAKER R.J., FJELDSÅ J. 2019b. Humboldt's enigma: What causes global patterns of mountain
biodiversity? *Science* 365, 1108-1113. https://doi.org/10.1126/science.aax0149

RANDIN C.F., ENGLER R., NORMAND S., ZAPPA M., ZIMMERMANN N.E., PEARMAN P.B., VITTOZ P., THUILLER W.,
GUISAN A. 2009. Climate change and plant distribution: local models predict high-elevation
persistence.   *Global   Change   Biology*   15,   1557-1569.   https://doi.org/10.1111/j.1365-
2486.2008.01766.x

ROGORA M., FRATE L., CARRANZA M.L., FREPPAZ M., STANISCI A., BERTANI I., BOTTARIN R., BRAMBILLA A.,
CANULLO R., CARBOGNANI M., CERRATO C., CHELLI S., CREMONESE E., CUTINI M., DI MUSCIANO M.,
ERSCHBAMER B., GODONE D., IOCCHI M., ISABELLON M., MAGNANI A., MAZZOLA L., MORRA DI CELLA U.,
PAULI H., PETEY M., PETRICCIONE B., PORRO F., PSENNER R., ROSSETTI G., SCOTTI A., SOMMARUGA R.,

TAPPEINER U., THEURILLAT J.-P., TOMASELLI M., VIGLIETTI D., VITERBI R., VITTOZ P., WINKLER M.,
MATTEUCCI G. 2018. Assessment of climate change effects on mountain ecosystems through a cross-
site analysis in the Alps and Apennines. *Science of The Total Environment* 624, 1429-1442.
https://doi.org/10.1016/j.scitotenv.2017.12.155

ROTA F., CASAZZA G., GENOVA G., MIDOLO G., PROSSER F., BERTOLLI A., WILHALM T., NASCIMBENE J.,
WELLSTEIN C. 2022. Topography of the Dolomites modulates range dynamics of narrow endemic
plants under climate change. *Sci Rep* 12, 1398. https://doi.org/10.1038/s41598-022-05440-3

RUBENSTEIN M.A., WEISKOPF S.R., BERTRAND R., CARTER S.L., COMTE L., EATON M.J., JOHNSON C.G., LENOIR
J., LYNCH A.J., MILLER B.W., MORELLI T.L., RODRIGUEZ M.A., TERANDO A., THOMPSON L.M. 2023.
Climate change and the global redistribution of biodiversity: substantial variation in empirical support
for expected range shifts. *Environmental Evidence* 12, 7. https://doi.org/10.1186/s13750-023-00296-
0

RUSSO L.F., FERNÁNDEZ-GONZÁLEZ Á., PENTERIANI V., DELGADO M. del M., PALAZÓN S., LOY A., DI FEBBRARO
M. 2023. The Different Fate of the Pyrenean Desman (*Galemys pyrenaicus*) and the Eurasian Otter
(*Lutra lutra*) under Climate and Land Use Changes. *Animals* 13, 274.
https://doi.org/10.3390/ani13020274

SALVADO P., AYMERICH BOIXADER P., PARERA J., VILA BONFILL A., MARTIN M., QUELENNEC C., LEWIN J.-M.,
DELORME-HINOUX V., BERTRAND J.A.M. 2022. Little hope for the polyploid endemic Pyrenean Larkspur
(*Delphinium montanum*): Evidences from population genomics and Ecological Niche Modeling.
Ecology and Evolution 12, e8711. https://doi.org/10.1002/ece3.8711

SALVADO P., COLLETTE N., GIBERT A., MARTIN M., QUÉLENNEC C., LEWIN J.-M., GARRIGUE J., MAGDALOU J.,
HURSON C., SOREL D., DELORME-HINOUX V., BERTRAND J.A.M. 2025. Species distribution modelling
predicts the extinction of the East-pyrenean local endemic form of *Pedicularis comosa : P. c.
asparagoides* in a near future (in press). *Vie et Milieu* 75, 1.

SCHMITT T. 2009. Biogeographical and evolutionary importance of the European high mountain systems.
Frontiers in Zoology 6, 9. https://doi.org/10.1186/1742-9994-6-9

SCHUCHARDT M.A., BERAUER B.J., DUC A.L., INGRISCH J., NIU Y., BAHN M., JENTSCH A. 2023. Increases in
functional diversity of mountain plant communities is mainly driven by species turnover under climate
change. *Oikos* 2023, e09922. https://doi.org/10.1111/oik.09922

SHAW R.G., ETTERSON J.R. 2012. Rapid climate change and the rate of adaptation: insight from experimental
quantitative genetics. *New Phytologist* 195, 752-765. https://doi.org/10.1111/j.1469-
8137.2012.04230.x

SILLERO N., ARENAS-CASTRO S., ENRIQUEZ-URZELAI U., VALE C.G., SOUSA-GUEDES D., MARTÍNEZ-FREIRÍA F.,
REAL R., BARBOSA A.M. 2021. Want to model a species niche? A step-by-step guideline on correlative
ecological niche modelling. *Ecological Modelling* 456, 109671.
https://doi.org/10.1016/j.ecolmodel.2021.109671

SNETHLAGE M.A., GESCHKE J., RANIPETA A., JETZ W., YOCCOZ N.G., KÖRNER C., SPEHN E.M., FISCHER M.,
URBACH D. 2022. A hierarchical inventory of the world's mountains for global comparative mountain
science. *Sci Data* 9, 149. https://doi.org/10.1038/s41597-022-01256-y

SOFAER H.R., HOETING J.A., JARNEVICH C.S. 2019. The area under the precision-recall curve as a performance
metric for rare binary events. *Methods in Ecology and Evolution* 10, 565-577.
https://doi.org/10.1111/2041-210X.13140

SONG Y., XU X., ZHANG S., CHI X. 2024. Uncertainty Assessment of Species Distribution Prediction Using
Multiple Global Climate Models on the Tibetan Plateau: A Case Study of *Gentiana yunnanensis* and
Gentiana siphonantha. *Land* 13, 1376. https://doi.org/10.3390/land13091376

STEINBAUER M., GRYTNES J.A., JURASINSKI G., KULONEN A., LENOIR J., PAULI H., RIXEN C., WINKLER M., BARDY-
DURCHHALTER M., BARNI E., BJORKMAN A., BREINER F., BURG S., CZORTEK P., DAWES M., DELIMAT A.,
DULLINGER S., ERSCHBAMER B., ASTRUP FELDE V., WIPF S. 2018. Accelerated increase in plant species
richness on mountain summits is linked to warming. *Nature* 556, 231-234.
https://doi.org/10.1038/s41586-018-0005-6

TEBALDI C., DEBEIRE K., EYRING V., FISCHER E., FYFE J., FRIEDLINGSTEIN P., KNUTTI R., LOWE J., O'NEILL B.,
SANDERSON B., VAN VUUREN D., RIAHI K., MEINSHAUSEN M., NICHOLLS Z., TOKARSKA K.B., HURTT G.,
KRIEGLER E., LAMARQUE J.-F., MEEHL G., MOSS R., BAUER S.E., BOUCHER O., BROVKIN V., BYUN Y.-H.,
DIX M., GUALDI S., GUO H., JOHN J.G., KHARIN S., KIM Y., KOSHIRO T., MA L., OLIVIÉ D., PANICKAL S.,
QIAO F., RONG X., ROSENBLOOM N., SCHUPFNER M., SÉFÉRIAN R., SELLAR A., SEMMLER T., SHI X., SONG
Z., STEGER C., STOUFFER R., SWART N., TACHIIRI K., TANG Q., TATEBE H., VOLDOIRE A., VOLODIN E.,
WYSER K., XIN X., YANG S., YU Y., ZIEHN T. 2021. Climate model projections from the Scenario Model
Intercomparison Project (ScenarioMIP) of CMIP6. *Earth Syst. Dynam.* 12, 253-293.
https://doi.org/10.5194/esd-12-253-2021

THUILLER W., GUÉGUEN M., RENAUD J., KARGER D.N., ZIMMERMANN N.E. 2019. Uncertainty in ensembles of
global biodiversity scenarios. *Nat Commun* 10, 1446. https://doi.org/10.1038/s41467-019-09519-w

TRIVEDI M., BERRY P., MORECROFT M., DAWSON T. 2008. Spatial Scale Affects Bioclimate Model Projections of
Climate Change Impacts on Mountain Plants. *Global Change Biology* 14, 1089-1103.
https://doi.org/10.1111/j.1365-2486.2008.01553.x

TRUNSCHKE J., JUNKER R.R., KUDO G., ALEXANDER J.M., RICHMAN S.K., TILL-BOTTRAUD I. 2024. Effects of
climate change on plant-pollinator interactions and its multitrophic consequences. *Alp Botany* 134,
115-121. https://doi.org/10.1007/s00035-024-00316-w

TSIFTSIS S., ŠTÍPKOVÁ Z., REJMÁNEK M., KINDLMANN P. 2024. Predictions of species distributions based only
on models estimating future climate change are not reliable. *Sci Rep* 14, 25778.
https://doi.org/10.1038/s41598-024-76524-5

TURNOCK M.F., TEISBERG J.E., KASWORM W.F., FALCY M.R., PROCTOR M.F., WAITS L.P. 2024. Gene flow
prevents genetic diversity loss despite small effective population size in fragmented grizzly bear (*Ursus
arctos*) populations. *Conserv Genet*. https://doi.org/10.1007/s10592-024-01666-y

VALAVI R., ELITH J., LAHOZ-MONFORT J.J., GUILLERA-ARROITA G. 2019. BLOCK CV : An R package for generating
spatially or environmentally separated folds for *k* -fold cross-validation of species distribution models.
Methods Ecol Evol 10, 225-232. https://doi.org/10.1111/2041-210X.13107

VALAVI R., GUILLERA-ARROITA G., LAHOZ-MONFORT J.J., ELITH J. 2022. Predictive performance of presence-
only species distribution models: a benchmark study with reproducible code. *Ecological Monographs*
92, e01486. https://doi.org/10.1002/ecm.1486

VALLADARES F., MATESANZ S., GUILHAUMON F., ARAÚJO M.B., BALAGUER L., BENITO-GARZÓN M., CORNWELL W.,
GIANOLI E., VAN KLEUNEN M., NAYA D.E., NICOTRA A.B., POORTER H., ZAVALA M.A. 2014. The effects
of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change.
Ecology Letters 17, 1351-1364. https://doi.org/10.1111/ele.12348

VASCONCELOS R.N., CANTILLO-PÉREZ T., FRANCA ROCHA W.J.S., AGUIAR W.M., MENDES D.T., DE JESUS T.B.,
941  DE SANTANA C.O., DE SANTANA M.M.M., OLIVEIRA R.P. 2024. Advances and Challenges in Species
942  Ecological Niche Modeling: A Mixed Review. *Earth* 5, 963-989. https://doi.org/10.3390/earth5040050
943 VITASSE Y., URSENBACHER S., KLEIN G., BOHNENSTENGEL T., CHITTARO Y., DELESTRADE A., MONNERAT C.,
944  REBETEZ M., RIXEN C., STREBEL N., SCHMIDT B.R., WIPF S., WOHLGEMUTH T., YOCCOZ N.G., LENOIR J.
945  2021. Phenological and elevational shifts of plants, animals and fungi under climate change in the
946  European Alps. *Biological Reviews* 96, 1816-1835. https://doi.org/10.1111/brv.12727
947 WELCH I. 2024. The IPCC Shared Socioeconomic Pathways (SSPs): Explained, Evaluated, Replaced (No.
948  w32178). National Bureau of Economic Research, Cambridge, MA. https://doi.org/10.3386/w32178
949 WILLIAMS S.E., SHOO L.P., ISAAC J.L., HOFFMANN A.A., LANGHAM G. 2008. Towards an Integrated Framework
950  for Assessing the Vulnerability of Species to Climate Change. *PLOS Biology* 6, e325.
951  https://doi.org/10.1371/journal.pbio.0060325
952 WILLIAMS-TRIPP M., D'AMICO F.J.N., PAGÉ C., BERTRAND A., NÉMOZ M., BROWN J.A. 2012. Modeling Rare
953  Species Distribution at the Edge: The Case for the Vulnerable Endemic Pyrenean Desman in France.
954  *The Scientific World Journal* 2012, e612965. https://doi.org/10.1100/2012/612965
955 ZANGO-PALAU A., JOLIVET A., LURGI M., CLARAMUNT-LÓPEZ B. 2024. A quantitative approach to the
956  understanding of social-ecological systems: a case study from the Pyrenees. *Reg Environ Change* 24,
957  9. https://doi.org/10.1007/s10113-023-02177-1
958 ZHU G., FAN J., PETERSON A.T. 2021. Cautions in weighting individual ecological niche models in ensemble
959  forecasting. *Ecological Modelling* 448, 109502. https://doi.org/10.1016/j.ecolmodel.2021.109502
960 ZHU G.-P., PETERSON A.T. 2017. Do consensus models outperform individual models? Transferability
961  evaluations of diverse modeling approaches for an invasive moth. *Biol Invasions* 19, 2519-2532.
962  https://doi.org/10.1007/s10530-017-1460-y
963 ZURELL D., FRANKLIN J., KÖNIG C., BOUCHET P.J., DORMANN C.F., ELITH J., FANDOS G., FENG X., GUILLERA-
964  ARROITA G., GUISAN A., LAHOZ-MONFORT J.J., LEITÃO P.J., PARK D.S., PETERSON A.T., RAPACCIUOLO
965  G., SCHMATZ D.R., SCHRÖDER B., SERRA-DIAZ J.M., THUILLER W., YATES K.L., ZIMMERMANN N.E.,
966  MEROW C. 2020. A standard protocol for reporting species distribution models. *Ecography* 43,
967  1261-1277. https://doi.org/10.1111/ecog.04960

    Figures for:

    Projecting spatiotemporal bioclimatic niche dynamics of endemic Pyrenean plant species under

    climate change: how many will we lose?

Figure 1. Geographic extent and topography of the Pyrenees. (a) The Pyrenees Mountain range (red

outline) spans the border regions of France, Spain, and Andorra. Its delimitation follows the definition set out

by Global Mountain Biodiversity Assessment (Körner *et al.*, 2011, updated by Snethlage *et al.*, 2022). The

inset map shows the location of the Pyrenees in Europe. Satellite imagery: Google Earth (© Google, Image

Landsat/Copernicus, Data SIO, NOAA, U.S. Navy, NGA, GEBCO). **(b)** Elevation gradient from a 90m-

resolution digital elevation model (CGIAR-CSI, 2018), showing the region's topographic variation from

lowlands (green) to high peaks (brown to white).

Figure 2. Projected changes in temperature and precipitation in the Pyrenees under Shared

Socioeconomic Pathways (SSP) by 2081-2100 period from WorldClim 2.1. (a) Mean annual temperature

(°C) and **(b)** annual precipitation (mm) projections for 2021-2100 under SSP126, SSP245, SSP370, and

SSP585. Values (+/-) reflect differences from the 1970-2000 baseline (current period) for the 2081-2100

period. 2021-2040, 2041-2060, 2061-2080, 2081-2100 periods designated as 2030, 2050, 2070, 2090.

1000

1001

1002

1003

Figure 3: Overview of the species distribution modeling pipeline. The framework consists of three main stages: (**1**) Input data, including occurrence records, environmental data processing, and background sampling; (**2**) Modeling, encompassing model building, performance assessment, projection, and ensemble modeling; (**3**) Output processing, manipulating species projections to assess spatiotemporal dynamics of suitable bioclimatic niches.

1033

1034

Figure 4: Endemic Pyrenean plant species model performances. (a) Boxplot of three performance metrics, AUC-PR (discrimination), Boyce Index (calibration), and Sensitivity (classification), for the five modeling algorithms (GAM, GBM, GLM, MAXENT, RF) for all species. Statistical significance is denoted as: *p < 0.05, **p < 0.01, ***p < 0.001. **(b)** Performance of ensemble models across species, computed by averaging evaluation metrics from the validated models (Boyce Index > 0.5) across the five algorithms used to build the ensemble consensus. Darker shades indicate higher performance.

Figure 5: Projected extent changes in bioclimatic suitability for endemic Pyrenean plant species under Shared Socioeconomic Pathways (SSP) for the 2021-2100 period. Extent changes in suitable areas under SSP126, SSP245, SSP370, SSP585, relative to the baseline period (1970-2000). Each box represents the extent change in suitable area, with percentage values shown for the 2081-2100 period. Orange indicates a reduction in suitable areas, green expansion, and black a complete loss of suitability. 2021-2040, 2041-2060, 2061-2080, 2081-2100 periods designated as 2030, 2050, 2070, 2090.

Figure 6. Projected distribution changes in bioclimatic suitability for endemic Pyrenean plant species between current and 2081-2100 periods. (a) Current and projected (2081-2100) bioclimatic hotspots, i.e. areas where the highest number of species could find suitable bioclimatic conditions, under Shared Socioeconomic Pathways (SSPs). The top panel shows the current distribution, while lower panels display projections for SSP126, SSP245, SSP370, and SSP585. Darker colors indicate higher bioclimatic suitability. Contour lines mark the 1000- and 2000-meter elevation topography. **(b)** Projected latitudinal and elevational shifts in bioclimatic suitability for the 2081–2100 period, averaged across SSPs. Each dot represents one of the 27 species retaining suitable areas under all SSPs. Dot color reflects the relative change in suitable area (green = gain; red = loss). The black arrow indicates the overall mean displacement in latitude ($\triangle$ lat) and elevation ($\triangle$ elev) across species. **(c)** Projected spatial overlap (%) between current and future (2081-2100) distributions of bioclimatically suitable areas under SSP126, SSP245, SSP370, and SSP585. Each row represents one species, with colors indicating the proportion (%) of overlap with present-day suitable areas.