Assessing species-specific neonicotinoid toxicity using cross-species chimeric nicotinic

acetylcholine receptors in a *Drosophila* model

ANNA LASSOTA[1,2], JAMES J. L. HODGE[3] AND MATTHIAS SOLLER[1,2,4*]

[1]School of Biosciences, College of Life and Environmental Sciences, University of

Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[2]Birmingham Centre for Genome Biology, University of Birmingham, Edgbaston,

Birmingham, B15 2TT, United Kingdom

[3]School of Physiology, Pharmacology and Neuroscience, University of Bristol, Biomedical

Sciences building, University Walk, Bristol, BS8 1TD, United Kingdom

[4]Division of Molecular and Cellular Function, School of Biological Sciences, University of

Manchester, Oxford Road, Manchester M13 9PT, United Kingdom

[*]Corresponding author: matthias.soller@manchester.ac.uk

Running title: Chimeric nAChRs for neonicotinoid toxicity in *Drosophila*

Key words: nAChR, neonicotinoids, *Apis mellifera*, *Drosophila*, CRISPR-Cas9 genome

engineering, pesticides

Abstract

Nicotinic acetylcholine receptors (nAChRs) are ligand-gated ion channels and the main mediators of synaptic neurotransmission in the insect brain. In insects, nAChRs are pivotal for sensory processing, cognition and motor control, and are the primary target of neonicotinoid insecticides. Neonicotinoids are potent neurotoxins and pollinators such as honey bees are more sensitive and affected by extremely low sub-lethal doses. The pentameric nAChR channel is made up either of five α-subunits constituting five ligand-binding sites or a mixture of two to three α and β subunits constitute two to three ligand-binding sites. Of particular note, the honey bee nAChRα8 subunit is converted into a β subunit (nAChRβ2) in *Drosophila*, raising the question whether this α to β conversion makes flies less sensitive to neonicotinoids. To investigate species-specific aspects of neonicotinoid toxicity we CRISPR-Cas9 engineered a cross-species chimeric nAChR subunit by swapping the ligand-binding domain in *Drosophila* of nAChRβ2 with honey bee nAChRα8. Toxicity assessment by neonicotinoid thiamethoxam revealed significantly impaired motor functions in climbing and flight assays when comparing the α8/β2 chimeric channel to wild type or a β2 knock-out. However, both the α8/β2 chimeric channel and the β2 knock-out showed the same increased survival after neonicotinoid exposure compared to wild type flies. Combinatorial exposure to neonicotinoids also did not reveal differences. These findings highlight the critical role of nAChR subunit composition in motor control and demonstrate how subtle structural differences can profoundly impact motor function and pesticide response, offering new insights into the molecular mechanisms of neurotoxicity across species.

Introduction

Pesticides are indispensable for protecting global agricultural yields by reducing crop losses caused by pests[1]. Among these, neonicotinoids are highly effective due to their broad-spectrum efficacy and systemic activity, which enable integration into plant tissues for targeted pest control, and versatility in application methods, such as seed coating, to reduce environmental contamination[2,3]. Neonicotinoids exhibit high toxicity to insects, but not to vertebrates, making them safer alternatives to other classes of chemical[4,5]. However, neonicotinoids do not distinguish between pests and beneficial insects, and harm beneficial insects like honey bees, which are critical pollinators for both wild flora and agricultural crops[6–8]. The widespread decline of pollinators, partially attributed to pesticide exposure, poses significant ecological and economic challenges, highlighting the need for pest-control strategies with reduced non-target effects.

Honey bees, wild bees and other pollinators are adversely affected by extremely low sub-lethal doses of neonicotinoids (2 ng/ml compared to the LC50 dose of 4.28 µg/ml) impairing foraging behaviour, cognition, navigation and weakening of the immune system, and reducing reproductive success and genetic diversity within a colony[6,9–20]. Neonicotinoids, which target cholinergic neurotransmission, inhibit mushroom body Kenyon cell activity[21] and disrupt memory and olfactory sensory neuron activity in honey bees (*Apis mellifera*) and fruit flies (*Drosophila melanogaster*)[21–24].

The sensitivity of insects to neonicotinoids varies considerably between species, with honey bees being highly susceptible and *Drosophila* more resistant. For example, the 24-hour lethal concentration 50% (LC50) of thiamethoxam (TMX) is 3.13 µg/ml for adult *Drosophila*[25], whereas for honey bees it is 4.28 µg/ml[20]. This variation arises from differences in the repertoire of cytochrome P40 detoxification genes. *Drosophila* encodes 85 of these neonicotinoid-metabolising enzymes, whereas honey bees have only 46[26,27]. Additionally, honey bees have

fewer antioxidant gene paralogs than *Drosophila*, which has an expanded antioxidant defence

system. Since neonicotinoids can induce oxidative stress, this weaker antioxidant capacity may

contribute to the greater sensitivity of bees compared to *Drosophila*[28].

Another reason contributing to this species-specific susceptibility could be the

composition of nicotinic acetylcholine receptors (nAChRs). nAChRs are essential for

neurotransmission, synaptic plasticity, and neuronal development, making prominent targets

for neonicotinoid pesticides[29–32]. These pentameric cys-loop ligand-gated cation channels

consist of α and β subunits forming homo- or heteropentamers. Ligand binding pockets, located

at α-α or α-β interfaces, consist of loops A, B, and C from an α subunit and loops D, E, and F

from the adjacent α or β subunit[33–35]. Importantly, β subunits lack the ability to form functional

homopentamers due to the absence of loops A, B, and C[33,35]. Consequently, the subunit

composition of nAChRs receptor affect their properties and determines susceptibility to

agonists, such as acetylcholine or neonicotinoids (Fig. 1a).

Insects have relatively small nAChR gene families. *Drosophila melanogaster* encodes

10 subunits (Dα1-Dα7 and Dβ1-Dβ3), while the honey bee *Apis mellifera* encodes 11 (Amelα1-

Amelα9 and Amelβ1-Amelβ2)[36]. α subunits are characterised by the presence of two adjacent

cysteines in loop C of the ligand-binding domain (LBD), whereas β subunits lack these

residues[35]. Notably, *Drosophila* encodes fewer α subunits compared to *Apis mellifera*. Instead,

the *Drosophila* β2 subunit exhibits striking conservation with the honey bee Amelα8 subunit

indicating a subunit conversion (Fig. 1b). However, the absence of loop C in *Drosophila* β2,

which is essential for ion channel gating, highlights a structural divergence that may contribute

to species-specific functional properties and differences in neonicotinoid susceptibility. Further,

specific amino acid substitutions in nAChR subunits are known to impact insecticide

susceptibility. For instance, *Drosophila* mutants lacking certain nAChR subunits or carrying

the natural variant R81T in the β1 subunit from aphids exhibit a 10- to 100-fold increase in

resistance to neonicotinoids[37–39].

Here, we examine the evolutionary divergence and functional implications of nAChR

subunits in *Drosophila melanogaster* and *Apis mellifera*. Phylogenetic analysis reveals that the

Drosophila β2 subunit has evolved from an ancestral α subunit. To investigate whether this

subunit conversion contributes to neonicotinoids sensitivity, we employed CRISPR-Cas9 gene-

editing technology to generate chimeric *Drosophila* flies expressing a chimeric nAChR

receptor. Specifically, we replaced the ligand-binding domain (LBD) of the *Drosophila* β2

subunit with that of the honey bee α8 subunit, providing a model to explore cross-species

receptor functionality. Behavioural and survival assays revealed significant impairments in

motor functions and altered sensitivity to insecticides, highlighting how structural differences

in receptor subunits can influence pesticide response. Our results provide insights into the

evolutionary trajectories of nAChR subunits and their roles in mediating neurotoxic compound

interactions, establishing a valuable *Drosophila* model for investigating cross-species pesticide

toxicity.

Results

Species-specific specialization of *Drosophila* β2 subunit during evolution

Insect *nAChRs* are classified into divergent subunits without clear orthologous

relationships between species, alongside a subset of genes that are highly conserved across

taxa[40]. Among these conserved subunits is the nAChRα8 subunit; however, it is absent in

certain Dipteran species, including *D. melanogaster*. Instead, the *Drosophila* β2 subunit

exhibits striking similarity to the honey bee α8 subunit, with 75% overall sequence identity and

92% identity within the LBD indicating that α8 converted in to the β2 subunit (Fig. 1b and

Supplementary Fig. 1). Alphafold structural modelling further confirmed conservation of the

protein secondary structure of these two subunits, particularly in the LBD, with the exception

of the loop C (Fig. 1d)[41].

Next, to investigate whether *Drosophila* β2 represents a divergent lineage from other β

subunits or shares a closer evolutionary relationship with α subunits, we conducted a

phylogenetic analysis of all nAChR subunits in *D. melanogaster* and *A. mellifera*. Strikingly,

the analysis revealed that *Drosophila* β2 clusters within a clade predominantly composed of α

subunits, including α1–α4 and α8, indicating that *Drosophila* β2 is more closely related to α

subunits than to other β subunits (Fig. 1c). Interestingly, the honey bee β2 subunit exhibits

significant divergence and does not closely group with any other subunits, suggesting a

divergent evolutionary origin.

***nAChR* subunits exhibit developmental dynamics and cell-specific distribution**

To investigate the expression patterns of *nAChR* subunits, we analysed tissue specific

expression data in adults, taking into account the enrichment values providing the gene

abundance measure in a tissue relative to that in the whole fly[42]. The majority of *nAChR*

subunits exhibits higher expression levels in females than in males (Fig. 2a). In both sexes,

nAChR subunits are predominantly expressed in the head, brain and CNS, and thoraco-

abdominal ganglion, with *Drosophila β2* being the most highly expressed subunit in the brain

and CNS. In other tissues, however, the levels of expression differ between sexes, with

Drosophila β2 and *α7* being the most expressed subunits in the thoraco-abdominal ganglion of

males and females, respectively. Furthermore, only two subunits, *Drosophila α4* and *β3*, are

expressed across all tissues, with *Drosophila β3* showing notable expression in the heart.

Specific differences were observed between sexes. For instance, in the male salivary glands,

carcass, and crop, only *Drosophila α1*, *α2*, and *α3* were expressed, while *α4*, *β1*, and *β3* were

restricted to females.

Further, we investigated whether the expression of *nAChR* subunit changes during

development. Larval tissue analysis revealed that *Drosophila α3* is not expressed in any tissue

(Supplementary Fig. 2a), suggesting it is dispensable during development. In contrast,

Drosophila β3 was expressed in all larval tissues, with the garland cells showing the highest

expression. The larval brain and CNS were the only tissues expressing all *nAChR* subunits, with

Drosophila β3 showing the highest levels. Overall, *nAChR* subunit expression is relatively low

in most tissues in both larvae and adults (Fig. 2a and Supplementary Fig. 2a).

To refine the spatial expression of *β2* in *Drosophila* cells in more detail, we analysed

expression in single cells[43]. In whole adult flies, *β2* was found to be expressed in various cell

types, including neurons, sensory neurons, glial cells, as well as male germline and reproductive

system, however, its expression is not ubiquitous (Fig. 2b). Notably, a prominent cluster of

neurons classified as adult fly head CNS cells strongly expressed *β2*, suggesting it may have a

significant role in neuronal circuits[23,44]. Since *β2* displayed the highest expression levels in the

brain and CNS, we further analysed single cell expression data for the larval brain and whole

adult fly head. In adults, *β2* expression was observed in distinct neuronal populations,

contrasting with the broader expression of the neuronal marker *elav* (Fig. 2c-e)[45]. In contrast,

larval *β2* was colocalised with *elav* in the majority of CNS neurons (Supplementary Fig. 1b-

d)[46].

To validate the expression patterns, we used *nAChRβ2[2A-GAL4]* allele which has a *T2A-*

GAL4 sequence fused at the 3' end of the gene, resulting in GAL4 expression as a separate

protein under the control of *nAChRβ2* regulation[47]. Therefore, we analysed both larval and adult

brains using nuclear *UAS-Histone2B-YFP* or membrane-tethered *UAS-mCD8-GFP* reporters,

co-staining neurons with anti-*elav* antibody. Consistent with our observations from the single

cell expression analyses, *β2* expression was restricted to specific neuronal populations,

confirming its non-ubiquitous distribution (Fig. 2f-k and Supplementary Fig. 1e-j).

Furthermore, we noticed enhanced *β2* expression in Kenyon cells, particularly when a

membrane-bound GFP reporter was used (Fig. 2i-k)[44].

Together, these findings reveal that nAChR subunit expression is dynamically regulated

during development and exhibits sex- and cell-specific specialisation, highlighting the

functional diversity of these receptors in neural circuits and other tissues. Furthermore, the

observed developmental shift in β2 expression, from widespread larval CNS expression to more

specialised adult neuronal populations, reflects the transition from simpler larval neural circuits

to the more specialised and complex adult CNS.

177 Chimeric *Amelα8/Dβ2* ligand-binding domain flies are viable with impaired motor

178 functions

Since the two adjacent cysteines in the loop C are essential for ligand binding in

nAChRs, we hypothesised that this difference may influence neonicotinoid susceptibility,

making honey bees more vulnerable. Therefore, we generated chimeric *nAChRβ2/α8*

(*nAChRβ2^{Amα8LBD}*) flies, where the *β2* LBD coding region in *Drosophila* was swapped with the

Amelα8 LBD coding sequence using CRISPR-Cas9 genome engineering (Fig. 3a and

Supplementary Fig. 3). Successful recombinant flies were marked with *white+* selection

marker, disrupting the LBD, therefore resulting in a null allele (*nAChRβ2^{nullA}*). To restore the

open reading frame (ORF) to generate a fly stock expressing the chimeric *nAChRβ2/α8* subunit

(*nAChRβ2^{Amα8LBD}*), the selection marker was then excised by PiggyBac transposase. Flies were

validated by polymerase chain reaction (PCR), Sanger sequencing, and subsequently by whole

genome sequencing.

To normalise the genetic background, we used trans-heterozygotic flies for the

nAChRβ2^{attP} allele (*nAChRβ2^{nullB}*) in which the coding region was replaced with an *attP* site,

3xP3-RFP and a *loxP* site[47], together with the *nAChRβ2^{nullA}* allele. The chimeric

nAChRβ2^{Amα8LBD} mutants, as well as *nAChRβ2^{nullA}* and *nAChRβ2^{nullB}* flies are fully viable when

crossed with a chromosomal deficiency (n = 453, 275, 421, respectively).

To assess motor abilities, we performed negative geotaxis assays (Fig. 3b) and evaluated

flight ability (Fig 3c). In negative geotaxis assays the climbing ability of LBD swap mutant

(*nAChRβ2^{Amα8LBD}*/*nAChRβ2^{nullB}*) was greatly reduced ($p = 7.5*10^{-7}$) (Fig. 3b) compared to wild

type (WT). We did not detect significant differences in the *β2* null mutant

(*nAChRβ2^{nullA}*/*nAChRβ2^{nullB}*) nor the heterozygous mutant controls.

In flying ability assays, we observed the same, namely the LBD swap mutant

(*nAChRβ2^{Amα8LBD}*/*nAChRβ2^{nullB}*) were impaired ($p = 0.0066$) (Fig. 3c) compared to control flies,

but not the *β2* null mutant (*nAChRβ2^{nullA}*/*nAChRβ2^{nullB}*) or heterozygote mutant controls.

Chimeric *Amelα8/Dβ2* ligand-binding domain flies display resistance to insecticides

To investigate the impact of substituting the *Drosophila* β2 LBD with the honey bee

Amelα8 LBD on neonicotinoid susceptibility, we selected two well-characterised

neonicotinoids, TMX and imidacloprid (IMI), both of which exhibit high toxicity to

pollinators[5,39]. These compounds differ slightly in chemical structure and receptor specificity,

with TMX being one of the most potent full agonists of *Drosophila* and honey bee α1, α2, β1

and β2 combinations when heterologously expressed in frog oocytes and assayed by patch-

clamp. In contrast, IMI shows a stronger affinity for *Drosophila* β2-containing receptors

compared to TMX[39,48]. Both compounds target multiple nAChR subunits, enhancing their

versatility broad-spectrum toxicity[39].

In addition, we tested flupyradifurone (FPF), a systemic butanolide compound that is

chemically distinct from neonicotinoids and represents a novel alternative to these compounds

in pest management[49]. While FPF is not primarily targeted at *Drosophila* β2 subunits[39], its

broad-spectrum activity and demonstrated toxicity to non-target pollinators, including bees[50,51],

make it an important candidate for assessing cross-resistance and potential off-target effects

resulting from altered nAChR subunit composition.

First, we conducted survival analysis testing a range of TMX concentrations on WT

flies (Supplementary Fig. 4a, b). Fly viability was measured over 6 days. Based on the survival

curves, three different TMX concentrations were used for the toxicity experiments.

At 10 μM TMX, wild type flies had 43.75% survival after 144 h, while

nAChRβ2^{Amα8LBD}/+, *nAChRβ2^{nullA}/+*, *nAChRβ2^{nullB}/+*, *nAChRβ2^{nullA}/nAChRβ2^{nullB}*, and

nAChRβ2^{Amα8LBD}/nAChRβ2^{nullB} flies remained viable (Fig. 4a). At 20 μM, survival rates

declined progressively across all genotypes, with *nAChRβ2^{nullA}/nAChRβ2^{nullB}* mutants showing

the highest survival rates, followed by *nAChRβ2^{Amα8LBD}/nAChRβ2^{nullB}* flies retaining more than

90% viability on day 6. This contrasted with *nAChRβ2^{nullA}/+* (71.25%), *nAChRβ2^{nullB}/+* (60%),

nAChRβ2^{Amα8LBD}/+ (35%), and WT (11.25%) (Fig. 4b). At 40 μM, similar trend was observed,

although the viability of all flies was decreased below 50%. *nAChRβ2^{nullA}/nAChRβ2^{nullB}*

mutants retained by 48.75% day 6, followed by *nAChRβ2^{Amα8LBD}/nAChRβ2^{nullB}* (41.25%),

nAChRβ2^{nullA}/+ (35%), *nAChRβ2^{nullB}/+* (16.25%), *nAChRβ2^{Amα8LBD}/+* (3.33%), while WT flies

showed 0% survival by day 4 (Fig. 4c). Interestingly, apart from the trans-heterozygous

β^{nullA}/β^{nullB} flies, all genotypes experienced a sharp drop in viability between the 1st and 2nd day

of treatment, particularly *nAChRβ2^{Amα8LBD}/+* dropping from 62.5% to 30% and WT failing from

79% to 20%.

While both neonicotinoids and butenolides act as nAChR agonists, bind to the same site

and cause prolonged activation, they vary in the binding dynamics, receptor subtype

preferences, and potential to develop resistance[52]. To assess whether combining neonicotinoids

and butanolides results in additive or synergistic toxicity, we subjected flies to a mixture of

TMX (20 μM) with either FPF (20 μM) or IMI (20 μM). In nature, pollinators and insects are

frequently exposed to multiple insecticides rather than a single compound, making it crucial to

understand potential interactions and cross-resistance effects[53,54]. The TMX concentration was chosen based on prior survival analyses, where double mutant flies retained > 90% viability (Fig. 4b), while WT flies showed significant susceptibility. We tested chimeric $nAChR\beta2^{Am\alpha8LBD}/nAChR\beta2^{nullB}$ flies and $nAChR\beta2^{nullA}/nAChR\beta2^{nullB}$ mutants to determine whether altered receptor composition influences responses to dual insecticide exposure. Flies were treated for six days, and viability on day six was analysed as a log-fold change relative to WT control and TMX-only treatment (Fig. 4d). Notably, exposure to TMX in combination with either FPF or IMI resulted in a slight increase in survival rates. However, the combined treatment did not significantly alter overall insecticide susceptibility. As expected, $nAChR\beta2^{Am\alpha8LBD}/nAChR\beta2^{nullB}$ flies and $nAChR\beta2^{nullA}/nAChR\beta2^{nullB}$ mutants flies exhibited significantly higher resistance compared to WT across all conditions.

Taken together, our findings demonstrate that $nAChR\beta2^{nullA}/nAChR\beta2^{nullB}$ null mutants as well as the LBD swap mutants $nAChR\beta2^{Am\alpha8LBD}/nAChR\beta2^{nullB}$ flies exhibit higher survival rates upon exposure to TMX, reinforcing the role of *Drosophila* β2 in neonicotinoid susceptibility. Further, exposure to combined neonicotinoid and butanolide treatments did not significantly alter overall lethality compared to TMX treatment on its own, indicating that sub-lethal doses of these nAChR-targeting insecticides do not act synergistically in this context. The observed resistance of chimeric and $\beta2^{null}$ mutants further highlights the functional divergence of Amelα8 in modifying nAChR-mediated insecticide susceptibility.

Discussion

The agricultural industry relies on pesticides to protect crops and maintain a stable food supply for a growing population. Among these, neonicotinoids are the most widely used insecticides due to their high efficacy and systemic action, reducing the need for repeated applications. However, residues in nectar and pollen pose a significant risk to pollinators.

Despite their well-documented toxicity to bees and other beneficial insects, neonicotinoids remain the dominant insecticides worldwide[55], highlighting the urgent need to mitigate their ecological impact.

Beyond their effects on pollinators, neonicotinoids exhibit cross-species variability in toxicity, with *Drosophila melanogaster* showing notably higher resistance than honey bees, despite both species having homologous nAChRs[48]. The mechanisms underlying this differential sensitivity remain unclear but are likely influenced by differences in receptor composition, metabolism, or detoxification pathways[26,28]. To investigate species-specific differences in neonicotinoid sensitivity, we established a *Drosophila* model where we swapped the LBD of *D. melanogaster* β2 with honey bee Amelα8 subunit. The LBD swap mutant did not reveal increased sensitivity to neonicotinoids compared to the *Drosophila* β2 null mutant. However, the LBD swap mutant displayed motor function deficits indicating that this chimeric receptor is functional. Minor structural changes in receptor subunits can profoundly impact pharmacological properties[33,35,38,39] and could account for the altered neonicotinoid responses observed in LBD chimeric mutant flies.

Previous studies in honey bees revealed tissue-specific expression of nAChR subunits, including α2, α8 and β1 in Kenyon cells, while α7 was restricted to antennal lobe neurons[56]. A similar functional diversity is observed in *Drosophila*, where nAChR subunits like α1 and α6 contribute to dendrite morphogenesis and synaptic transmission within larval visual circuits[57]. The absence of α3 expression in *Drosophila* larvae, combined with reduced nAChR subunit expression during early developmental stages, suggests that receptor composition undergoes developmental reprogramming to support the formation of complex adult neural circuits. This dynamic tissue-, sex-, and stage-specific expression underscores the diverse roles of nAChRs, including potential adaptations for behaviours such as reproduction[58,59]. Interestingly, in the cockroach *Periplaneta americana*, subchronic exposure to sublethal doses of imidacloprid

induces a significant decrease in α2 mRNA expression, which correlates with a reduced

sensitivity to the insecticide[60], highlighting potential adaptive changes in nAChR subunit

composition even in fully developed insects.

Furthermore, high *Drosophila* β2 expression levels in CNS-specific neuronal

populations, including Kenyon cells, suggest its involvement in higher-order neuronal functions

such as learning and memory[44]. Similarly, honey bee α8 is enriched in honey bee mushroom

bodies[36,61,62], suggesting a conserved role for *Drosophila* β2 and honey bee α8 in regulating

mushroom body function across insect taxa, despite species-specific specialisations. Notably,

it has been shown that exposure to neonicotinoid insecticides affects cell activity in mushroom

body, and impairs learning and memory in insects, including *Drosophila* and honey bees[21–23,63],

further highlighting the critical role of nAChRs in cognitive functions and suggesting that

disruption of these receptors can have detrimental effects on neuronal processes related to

behaviour and learning.

Pharmacological and genetic studies have identified α1, α2, β1, and β2 as the primary

mediators of neonicotinoid toxicity[40,48,64]. Here, we generated a cross-species chimeric nAChR

using CRISPR-Cas9 genome engineering to evaluate neonicotinoid sensitivity in a *Drosophila*

model. Our study reveals that the *Drosophila* β2 conversion of honey bee α8 is not a main

determine of neonicotinoid sensitivity requiring to expand this approach to the other main

mediators of neonicotinoid toxicity. Our study establishes a framework for investigating

molecular determinants of neonicotinoid sensitivity of insects and vertebrates nAChR using a

cross-species chimeric nAChR *Drosophila* model. This model can contribute to the design of

next generation of insecticides for enhanced species-specific responses with implications for

pollinator conservation and pesticide safety for vertebrates including humans.

Materials and methods

Fly stocks, genetics, immunostaining of tissues and imaging

      D. melanogaster CantonS and w[1118] were used as the wild type control. *nAChRβ2[attP]* (BDSC 84545) and *nAChRβ2[2A-GAL4]* (BDSC 84666) stocks were described previously [47] and were together with the chromosomal deficiency (BDSC 24996) from Bloomington. Fly crosses were maintained at 25°C in plastic vials containing 10 ml of a standard cornmeal/yeast-rich medium (1% agar, 2% yeast, 7% dextrose, 8% cornmeal w/v and 2% Nipagin from a 10% solution in ethanol) with a 12:12 h light–dark cycle.

      Third instar wandering larvae and adult brains from the progeny of *UAS-Histone2B-YFP/+;nAChRβ2[2A-GAL4]/+* and *UAS-mCD8-GFP/+;nAChRβ2[2A-GAL4]/+* were dissected in phosphate buffered saline (PBS) and fixed in 4% paraformaldehyde in PBT (PBS with 0.1% TritonTM X-100 (Sigma-Aldrich, T8787)) for 30 min, followed by washes in PBT 3 × 15 min. Samples were incubated overnight at 4°C with primary mouse anti-*elav* antibodies (MAb 7D, 1:20)[65], followed by secondary antibodies conjugated with Alexa Fluor 546 again overnight at 4°C. Samples were counterstained with DAPI (1:1000), mounted in Vectashield (Vector Labs), scanned with Leica SP8, and processed using FIJI.

Sequence analysis and single cell expression data visualisation

      Amino acid sequences were aligned using ClustalW with Megalin (DNAstar) or with MAFFT[66] and phylogenetic trees were generated in NGPhylogeny.fr[67–69] and visualised using iTOL[70].

      Raw whole-genome sequencing reads were assessed for quality using FastQC, followed by adapter trimming and quality filtering with Trim Galore. High-quality reads were aligned to the *Drosophila melanogaster* (dm6) and *Apis mellifera* (Amel_HAv3.1) reference genomes using BBMap. Mapping quality and coverage were evaluated, and alignments were visualized using IGV (Integrative Genomics Viewer).

343   Single cell expression data was visualised as t-distributed stochastic neighbour

344   embedding (tSNE) from the 10x Stringent dataset in Scope and ASAP[71].

Generation of chimeric *nAChRα8β2* flies

Two single guide RNAs (sgRNAs) flanking the ligand binding domain *Dβ2* were

designed using PlatinumCRISPr[72] (Supplementary Fig. 3a, b) and cloned into *pUC-3GLA* using

the following primers: nACHr8 sgRNA left F1

(AAGATATCCGGGTGAACTTCGCTTATTGGAGCTAGGAAAGGTTTTAGAGCTAGA

AATAGC) and nACHr8 sgRNA right R1

(GCTATTTCTAGCTCTAAAACCACATGGCACAATCAAATTCGACGTTAAATTGAAA

ATAGG) as described previously[72]. Chimeric nAChR subunits were constructed by combining

sequences of *nAChRα8* from *A. mellifera* and *nAChRβ2* from *D. melanogaster* in three steps

(Supplementary Fig. 3c-e). Fragments used for generating the chimeric nAChR subunit were

synthesised using PCR with primers Bee alpha8 F2

(GGGAGTGGTCATTGCCATCTCAACCCTATATAAATTTG) and Bee alpha8 R2

(GGAATGTAATACCCACGCAAGGAATTATTAAG), followed by Bee alpha8 F1

(GCCAGTGAATTCGAGCTCGGTACCGCAGCTAGCGAAGCAAATCCTGACACAAAG

AGACTTTATGATGAC) and Bee alpha8 R1

(TTTACGCAGACTATCTTTCTAGGGTTAACCGTATAGAATAATGTTTTTCTTCGCAT

TG) for Amelα8 ligand binding domain, Dmel nAChRb left F1

(CGGCCAGTGAATTCGAGCTCGGTACCGATCCTTTTAGATAAAACATTTAGGAGCT

ATC) and nAChRsgmut left R1

(CTCTTTGTGTCAGGATTTGCTTCGAAACTCACTGGAGCCGTGAGAGGAGGGGTA

ATTCAGGGGAAAAACAGAGAAAAATGCC) for Dβ2 left homology arm, and

nAChRsgmut right F1

(ATTTTACGCATGATTATCTTTAACGTACGTCACAATATGATTATCTTTCTAGGGTT

AATCTGATTGTGCCCTGCGTAGCTTTAACATTCC) and Dmel alpha8 right R1

(GCATGCCTGCAGGTCGACTCTAGAGGATCCTGATAGTTGCTGCTGCGAATGCGG

AGCTG) for Dβ2 right homology arm, ensuring sgRNA sites in both homology arms are

mutated. Mutations were introduced based on conservation of the region between closely

related *Drosophila* species analysed using the UCSC genome browser

(https://genome.ucsc.edu) as described previously[73]. Fragments were then cloned stepwise into

the *pUC19 pBac w+* [accession number: PV267745] containing *white +* marker, disrupting the

LBD, therefore resulting in *nAChRβ2^{null}* flies. The selection marker was flanked with inverted

terminal repeats inserted at a TTAA PiggyBac motifs for the later scarless excision by PiggyBac

transposase.

The *pUC19 pBac w+ Amelα8/Dβ2* plasmid was treated with the nicking endonuclease

*Nb.Bts*I (NEB, R0707S) prior injection. Transgenic lines were generated by injection into a

GFP nosCas9 carrying flies (kindly provided by FlyORF). Positive transformants were

identified by red eye colour.

For *Drosophila* whole genome sequencing, large fragment DNA was extracted with the

Quick-DNA Tissue/Insect Miniprep Kit (Zymogene) according to the manufacturers'

instructions but replacing the BashingBead lysis step with cracking flies in liquid nitrogen

followed by gentle homogenization with a pestle in the provided BashingBead Buffer. Illumina

sequencing was done by Novogene and the sequence submitted to GEO [PRJNA1233772].

Locomotive assessment

Negative geotaxis experiments were done as previously described[74]. Briefly, two to five

day-old flies of both sexes were collected with CO_2 anaesthesia and grouped in sets of 20. Flies

were allowed to recover for a day and then placed in two inverted fly vials (19 cm). Flies were

tapped to the bottom, and their climbing behaviour was recorded on video. Every 5 s for a total

of 30 s, the distance climbed by each fly was measured and recorded. The data was collected

for four biological replicates.

For flying ability assessment, two to five day-old flies of both sexes were collected with

CO_2 anaesthesia and grouped in sets of 10. Flies were allowed to recover for a day before the

flying test. Then, flies were tapped to a flat surface, and their ability to fly was recorded on

video. Flies were observed over a 30 s period. At the end of 30 s, the number of flies that

successfully flew away was recorded. The data was collected for four biological replicates.

Insecticide toxicity assay

Two to five day-old flies of both sexes, grouped in sets of 20, were exposed to various

concentrations of thiamethoxam (Sigma-Aldrich, 37924), or a combination of 20 µM of

thiamethoxam with 20 µM imidacloprid (Thermo Fisher, 466752500) or 20 µM flupyradifurone

(MCE, HY-145295). The insecticides were dissolved in 0.01% acetone and added on top of the

standard fly food, and flies were added to the vials after 24 h. Exposure concentrations ranged

from 1.25 µM to 80 µM for survival curve analysis (Supplementary Fig. 4) and from 5 µM to

40 µM for testing all genotypes. Flies were exposed for 6 days, and survival was assessed every

24 h.

Statistical analysis

Statistical analyses were carried out using GraphPad Prism 9, applying one-way

ANOVA for behavioural assays or two-way ANOVA for the toxicity assays followed by

Tukey's test for pairwise comparisons. The data represent a minimum of three independent

replicates conducted on separate days.

Acknowledgments

We thank the Bloomington stock centre and FlyORF for fly lines, FlyORF for *Drosophila* injection service for embryo injections, and Al Handler for the PiggyBac plasmid.

Authors' contributions

A.L. performed all experiments and analysed data. M.S. and J.J.L.H. conceptualised the project. A.L. wrote the manuscript and M.S. and J.J.L.H. reviewed it. All authors read and approved the final manuscript.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council to M.S. and J.J.L.H.

Availability of data and materials

The data generated or analysed during this study are included in the supplementary information in the supplementary information files. Whole genome sequencing data have been deposited at GEO under the accession number PRJNA1233772. The sequence for the *pUC19 pBac w+* has been deposited in GenBank under the accession number PV267745 and the plasmid is available from Addgene and the European plasmid repository.

Declarations:

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1: Highly conserved *Drosophila* β2 subunit originated from an alpha subunit in

other insects

(a) Schematic representation of homo- and heteropentameric nAChR receptors in insects

exhibiting different compositions of α (blue) and β (pink) subunits, with their active ligand

binding sites represented by ligands (cyan).

(b) Amino acid alignment of *D. melanogaster* β2 and honey bee Amelα8 subunits. Identical

amino acids are marked as red, complementary amino acid substitutions are labelled orange,

and non-complementary amino acid substitutions remained unmarked. The ligand binding

domain is underlined.

(c) Molecular phylogenetic analysis of all nAChR subunits present in *Drosophila* and honey

bee with two major α subunits clades highlighted with light blue and pink (subunits highly

conserved with mammalian α7). Species codes are Dm: *D. melanogaster* Amel: *A. mellifera*.

(d) Structural alignment of *D. melanogaster* β2 (blue) and honey bee α8 (pink) with all amino

acid differences in *D. melanogaster* ligand binding domain indicated (cyan).

Figure 2: *Drosophila β2* is highly expressed in adult *Drosophila* brain and CNS but not in

every cell

(a) Expression levels of *nAChR* subunits in female and male adult fly tissues.

(b) Visualisation of single cell expression for the *nAChRβ2* gene indicated in red with

annotation of the major cell clusters exhibiting *Dβ2* expression.

(c-e) Visualisation of single cell expression for *nAChRβ2* (c) and neuronal marker *elav* (d) or

both (e).

(f-k) Expression of *Dβ2* (f, i), *elav* (g, j) and merged (h, k) in adult *Drosophila* brains with

nuclear (f, h) and membrane-bound (i, k) *UAS* reporters. Scale bars are 20 μm.

Figure 3: Chimeric *nAChRα8β2* flies exhibit locomotor defects

(a) Schematic *Dβ2* gene model with the ligand binding domain swap indicated with coding

exons marked in pink. The *white+* selection marker was incorporated into the *nAChRβ2* gene

along with the honey bee *nAChRα8* LBD encoding fragment, generating a *nAChRβ2^{null}* allele.

The reading frame can be restored by excising the PiggyBac transposon.

(b) Climbing activity was assessed by negative geotaxis and is shown as means with the

standard error from four biologically independent groups (n=20). Statistically significant

differences are indicated by asterisks (****$p \leq 0.0001$).

(c) Flies of the indicated genotypes were tested for their flight ability shown as means with the

standard error from four biologically independent groups (n=10). Statistically significant

differences are indicated by asterisks (**$p \leq 0.01$).

Figure 4: The α8 ligand binding domain does not sensitize flies to neonicotinoids

(a-c) Flies viability exposed to 10 μM (a), 20 μM (b) and 40 μM (c) thiamethoxam measured

every 24 h for 6 days is shown as mean with standard error from four biologically independent

groups (n = 20).

(d) Survival rate of *nAChRβ2^{Amα8LBD}/nAChRβ2^{nullA}* and *nAChRβ2^{nullA}/nAChRβ2^{nullB}* mutants

compared to WT flies subjected to different insecticides combinations. The log2 fold-change

shown as mean with standard error from four biologically independent groups (n = 20)

represents the viability of flies after six days of treatment with thiamethoxam (TMX) or TMX

mixed with imidacloprid (IMI) or flupyradifurone (FPF) in equimolar ratios. Statistically

significant differences are indicated with letters (a = $*p \leq 0.5$, b = $**p \leq 0.01$, c = $***p \leq 0.001$,

d = $****p \leq 0.0001$).

References

1. Fenik, J., Tankiewicz, M. & Biziuk, M. Properties and determination of pesticides in fruits and vegetables. *TrAC Trends Anal. Chem.* **30**, 814–826 (2011).

2. Elbert, D. L., Mawuenyega, K. G., Scott, E. A., Wildsmith, K. R. & Bateman, R. J. Stable Isotope Labeling Tandem Mass Spectrometry (SILT): Integration with Peptide Identification and Extension to Data-Dependent Scans. *J. Proteome Res.* **7**, 4546–4556 (2008).

3. Jeschke, P., Nauen, R., Schindler, M. & Elbert, A. Overview of the Status and Global Strategy for Neonicotinoids. *J. Agric. Food Chem.* **59**, 2897–2908 (2011).

4. Gibbons, D., Morrissey, C. & Mineau, P. A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. *Environ. Sci. Pollut. Res.* **22**, 103–118 (2015).

5. Casida, J. E. Neonicotinoids and Other Insect Nicotinic Receptor Competitive Modulators: Progress and Prospects. *Annu. Rev. Entomol.* **63**, 125–144 (2018).

6. Henry, M. *et al.* A Common Pesticide Decreases Foraging Success and Survival in Honey Bees. *Science* **336**, 348–350 (2012).

7. Pisa, L. W. *et al.* Effects of neonicotinoids and fipronil on non-target invertebrates. *Environ. Sci. Pollut. Res.* **22**, 68–102 (2015).

8. Woodcock, B. A. *et al.* Impacts of neonicotinoid use on long-term population changes in wild bees in England. *Nat. Commun.* **7**, 12459 (2016).

9. Crall, J. D. *et al.* Neonicotinoid exposure disrupts bumblebee nest behavior, social networks, and thermoregulation. *Science* **362**, 683–686 (2018).

10. Gill, R. J., Ramos-Rodriguez, O. & Raine, N. E. Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature* **491**, 105–108 (2012).

11. Muth, F. & Leonard, A. S. A neonicotinoid pesticide impairs foraging, but not learning, in free-flying bumblebees. *Sci. Rep.* **9**, 4764 (2019).

12. Rundlöf, M. *et al.* Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* **521**, 77–80 (2015).

524   13. Tasman, K., Rands, S. A. & Hodge, J. J. L. Using radio frequency identification and locomotor

525   activity monitoring to assess sleep, locomotor, and foraging rhythmicity in bumblebees. *STAR*

526   *Protoc.* **2**, 100598 (2021).

527   14. Tasman, K., Rands, S. A. & Hodge, J. J. L. The Neonicotinoid Insecticide Imidacloprid Disrupts

528   Bumblebee Foraging Rhythms and Sleep. *iScience* **23**, 101827 (2020).

529   15. Whitehorn, P. R., O'Connor, S., Wackers, F. L. & Goulson, D. Neonicotinoid Pesticide Reduces

530   Bumble Bee Colony Growth and Queen Production. *Science* **336**, 351–352 (2012).

531   16. Forfert, N. *et al.* Neonicotinoid pesticides can reduce honeybee colony genetic diversity. *PLOS*

532   *ONE* **12**, e0186109 (2017).

533   17. Annoscia, D. *et al.* Neonicotinoid Clothianidin reduces honey bee immune response and contributes

534   to Varroa mite proliferation. *Nat. Commun.* **11**, 5887 (2020).

535   18. Decio, P. *et al.* Thiamethoxam exposure deregulates short ORF gene expression in the honey bee

536   and compromises immune response to bacteria. *Sci. Rep.* **11**, 1489 (2021).

537   19. Di Prisco, G. *et al.* Neonicotinoid clothianidin adversely affects insect immunity and promotes

538   replication of a viral pathogen in honey bees. *Proc. Natl. Acad. Sci.* **110**, 18466–18471 (2013).

539   20. Oliveira, R. A., Roat, T. C., Carvalho, S. M. & Malaspina, O. Side-effects of thiamethoxam on the

540   brain andmidgut of the africanized honeybee *Apis mellifera* (Hymenopptera: Apidae). *Environ.*

541   *Toxicol.* **29**, 1122–1133 (2014).

542   21. Palmer, M. J. *et al.* Cholinergic pesticides cause mushroom body neuronal inactivation in

543   honeybees. *Nat. Commun.* **4**, 1634 (2013).

544   22. Smith, D. B. *et al.* Insecticide exposure during brood or early-adult development reduces brain

545   growth and impairs adult learning in bumblebees. *Proc. R. Soc. B Biol. Sci.* **287**, 20192442 (2020).

546   23. Tasman, K., Hidalgo, S., Zhu, B., Rands, S. A. & Hodge, J. J. L. Neonicotinoids disrupt memory,

547   circadian behaviour and sleep. *Sci. Rep.* **11**, 2061 (2021).

548   24. Tatarko, A. R., Leonard, A. S. & Mathew, D. A neonicotinoid pesticide alters Drosophila olfactory

549   processing. *Sci. Rep.* **13**, 10606 (2023).

550   25. Li, X., Liu, J. & Wang, X. Exploring the multilevel hazards of thiamethoxam using Drosophila

551   melanogaster. *J. Hazard. Mater.* **384**, 121419 (2020).

26. Berenbaum, M. R. & Johnson, R. M. Xenobiotic detoxification pathways in honey bees. *Curr. Opin.*

Insect Sci. **10**, 51–58 (2015).

27. Tasman, K., Rands, S. A. & Hodge, J. J. L. The Power of Drosophila melanogaster for Modeling

Neonicotinoid Effects on Pollinators and Identifying Novel Mechanisms. *Front. Physiol.* **12**,

659440 (2021).

28. Corona, M. & Robinson, G. E. Genes of the antioxidant system of the honey bee: annotation and

phylogeny. *Insect Mol. Biol.* **15**, 687–701 (2006).

29. Tomizawa, M. & Casida, J. E. Structure and diversity of insect nicotinic acetylcholine receptors.

Pest Manag. Sci. **57**, 914–922 (2001).

30. Role, L. W. & Berg, D. K. Nicotinic Receptors in the Development and Modulation of CNS

Synapses. *Neuron* **16**, 1077–1085 (1996).

31. Wessler, I. & Kirkpatrick, C. J. Acetylcholine beyond neurons: the non-neuronal cholinergic system

in humans. *Br. J. Pharmacol.* **154**, 1558–1571 (2008).

32. Tomizawa, M., Millar, N. S. & Casida, J. E. Pharmacological profiles of recombinant and native

insect nicotinic acetylcholine receptors. *Insect Biochem. Mol. Biol.* **35**, 1347–1355 (2005).

33. Matsuda, K. *et al.* Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors.

Trends Pharmacol. Sci. **22**, 573–580 (2001).

34. Karlin, A. Emerging structure of the Nicotinic Acetylcholine receptors. *Nat. Rev. Neurosci.* **3**, 102–

114 (2002).

35. Ihara, M. *et al.* Crystal structures of Lymnaea stagnalis AChBP in complex with neonicotinoid

insecticides imidacloprid and clothianidin. *Invert. Neurosci.* **8**, 71–81 (2008).

36. Dupuis, J., Louis, T., Gauthier, M. & Raymond, V. Insights from honeybee (Apis mellifera) and fly

(Drosophila melanogaster) nicotinic acetylcholine receptors: From genes to behavioral functions.

Neurosci. Biobehav. Rev. **36**, 1553–1564 (2012).

37. Bass, C. *et al.* Mutation of a nicotinic acetylcholine receptor β subunit is associated with resistance

to neonicotinoid insecticides in the aphid Myzus persicae. *BMC Neurosci.* **12**, 51 (2011).

38. Homem, R. A. *et al.* Evolutionary trade-offs of insecticide resistance — The fitness costs associated

with target-site mutations in the nAChR of *Drosophila melanogaster*. *Mol. Ecol.* **29**, 2661–2675

(2020).

39. Lu, W. *et al.* Nicotinic acetylcholine receptor modulator insecticides act on diverse receptor

subtypes with distinct subunit compositions. *PLOS Genet.* **18**, e1009920 (2022).

40. Jones, A. K. & Sattelle, D. B. Diversity of Insect Nicotinic Acetylcholine Receptor Subunits. in

Insect Nicotinic Acetylcholine Receptors (ed. Thany, S. H.) vol. 683 25–43 (Springer New York,

New York, NY, 2010).

41. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589

(2021).

42. Krause, S. A., Overend, G., Dow, J. A. T. & Leader, D. P. FlyAtlas 2 in 2022: enhancements to the

Drosophila melanogaster expression atlas. *Nucleic Acids Res.* **50**, D1010–D1015 (2022).

43. Li, H. *et al.* Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science* **375**,

eabk2432 (2022).

44. Barnstedt, O. *et al.* Memory-Relevant Mushroom Body Output Synapses Are Cholinergic. *Neuron*

89, 1237–1247 (2016).

45. Ustaoglu, P. *et al.* Dynamically expressed single ELAV/Hu orthologue elavl2 of bees is required

for learning and memory. *Commun. Biol.* **4**, 1234 (2021).

46. Brunet Avalos, C., Maier, G. L., Bruggmann, R. & Sprecher, S. G. Single cell transcriptome atlas

of the Drosophila larval brain. *eLife* **8**, e50354 (2019).

47. Deng, B. *et al.* Chemoconnectomics: Mapping Chemical Transmission in Drosophila. *Neuron* **101**,

876-893.e4 (2019).

48. Ihara, M. *et al.* Cofactor-enabled functional expression of fruit fly, honeybee, and bumblebee

nicotinic receptors reveals picomolar neonicotinoid actions. *Proc. Natl. Acad. Sci.* **117**, 16283–

16291 (2020).

49. Nauen, R. *et al.* Flupyradifurone: a brief profile of a new butenolide insecticide. *Pest Manag. Sci.*

71, 850–862 (2015).

50. Tosi, S. *et al.* Long-term field-realistic exposure to a next-generation pesticide, flupyradifurone, impairs honey bee behaviour and survival. *Commun. Biol.* **4**, 805 (2021).

51. Siviter, H. *et al.* A novel pesticide has lethal consequences for an important pollinator. *Sci. Total Environ.* **952**, 175935 (2024).

52. Crossthwaite, A. J. *et al.* The invertebrate pharmacology of insecticides acting at nicotinic acetylcholine receptors. *J. Pestic. Sci.* **42**, 67–83 (2017).

53. Johnson, R. M., Dahlgren, L., Siegfried, B. D. & Ellis, M. D. Acaricide, Fungicide and Drug Interactions in Honey Bees (Apis mellifera). *PLoS ONE* **8**, e54092 (2013).

54. Decio, P. *et al.* Acute thiamethoxam toxicity in honeybees is not enhanced by common fungicide and herbicide and lacks stress-induced changes in mRNA splicing. *Sci. Rep.* **9**, 19196 (2019).

55. Sparks, T. C. *et al.* Insecticides, biologics and nematicides: Updates to IRAC's mode of action classification - a tool for resistance management. *Pestic. Biochem. Physiol.* **167**, 104587 (2020).

56. Cens, T. *et al.* Molecular Targets of Neurotoxic Insecticides in *Apis mellifera. Eur. J. Org. Chem.* **2022**, e202101531 (2022).

57. Rosenthal, J. S. *et al.* Temporal regulation of nicotinic acetylcholine receptor subunits supports central cholinergic synapse development in *Drosophila. Proc. Natl. Acad. Sci.* **118**, e2004685118 (2021).

58. Komori, Y. *et al.* Functional impact of subunit composition and compensation on Drosophila melanogaster nicotinic receptors–targets of neonicotinoids. *PLOS Genet.* **19**, e1010522 (2023).

59. Yoshinari, Y. *et al.* Neuronal octopamine signaling regulates mating-induced germline stem cell increase in female Drosophila melanogaster. *eLife* **9**, e57101 (2020).

60. Bantz, A., Goven, D., Siegwart, M., Maugin, S. & Raymond, V. Exposure to a sublethal dose of imidacloprid induces cellular and physiological changes in Periplaneta americana: Involvement of α2 nicotinic acetylcholine subunit in imidacloprid sensitivity. *Pestic. Biochem. Physiol.* **181**, 105014 (2022).

61. Thany, S. H., Crozatier, M., Raymond-Delpech, V., Gauthier, M. & Lenaers, G. Apisα2, Apisα7-1 and Apisα7-2: three new neuronal nicotinic acetylcholine receptor α-subunits in the honeybee brain. *Gene* **344**, 125–132 (2005).

62. Thany, S. H., Lenaers, G., Crozatier, M., Armengaud, C. & Gauthier, M. Identification and

localization of the nicotinic acetylcholine receptor alpha3 mRNA in the brain of the honeybee, *Apis*

mellifera. *Insect Mol. Biol.* **12**, 255–262 (2003).

63. Ferreira, L. C. *et al.* Disruption of exploratory behavior and olfactory memory in cockroaches

exposed to sublethal doses of the neonicotinoid Thiamethoxam. *Pestic. Biochem. Physiol.* **205**,

106167 (2024).

64. Perry, T. *et al.* Role of nicotinic acetylcholine receptor subunits in the mode of action of

neonicotinoid, sulfoximine and spinosyn insecticides in Drosophila melanogaster. *Insect Biochem.*

Mol. Biol. **131**, 103547 (2021).

65. Haussmann, I. U., Li, M. & Soller, M. ELAV-Mediated 3′-End Processing of *ewg* Transcripts Is

Evolutionarily Conserved Despite Sequence Degeneration of the ELAV-Binding Site. *Genetics*

189, 97–107 (2011).

66. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

67. Criscuolo, A. & Gribaldo, S. BMGE (Block Mapping and Gathering with Entropy): a new software

for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evol.*

Biol. **10**, 210 (2010).

68. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies:

Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).

69. Lemoine, F. *et al.* NGPhylogeny.fr: new generation phylogenetic services for non-specialists.

Nucleic Acids Res. **47**, W260–W265 (2019).

70. Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree

display and annotation tool. *Nucleic Acids Res.* **52**, W78–W82 (2024).

71. David, F. P. A., Litovchenko, M., Deplancke, B. & Gardeux, V. ASAP 2020 update: an open,

scalable and interactive web-based portal for (single-cell) omics analyses. *Nucleic Acids Res.* **48**,

W403–W414 (2020).

72. Haussmann, I. U. *et al.* Structure-optimized sgRNA selection with PlatinumCRISPr for efficient

Cas9 generation of knockouts. *Genome Res.* **34**, 2279–2292 (2024).

73. McQuarrie, D. W. J. & Soller, M. Phylogenomic instructed target analysis reveals ELAV complex

binding to multiple optimally spaced U-rich motifs. *Nucleic Acids Res.* **52**, 12712–12726 (2024).

74. Haussmann, I. U. *et al.* CMTr cap-adjacent 2′-O-ribose mRNA methyltransferases are required for

reward learning and mRNA localization to synapses. *Nat. Commun.* **13**, 1209 (2022).

Figure 1

a

b

Climbing activity (cm/s)

c

Flying flies (%)

WT (+/+)
nAChRß2$^{Am\alpha8LBD}$/+
nAChRß2$^{Am\alpha8LBD}$/*nAChRß2*^{nullB}
nAChRß2^{nullA}/+
nAChRß2^{nullB}/+
nAChRß2^{nullA}/*nAChRß2*^{nullB}

Figure 4