Structure of a new capsid form and comparison with A-, B- and C-capsids clarify herpesvirus assembly

Alexander Stevens[a,b,c,d,*], Saarang Kashyap[a,b,*], Ethan Crofut[a,b,*], Ana Lucia Alverez-Cabrera[a,b], Jonathan Jih[a,b,c], Yun-Tao Liu[a], Z. Hong Zhou[a,b,c,d]

[a]Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

[b]California NanoSystems Institute, UCLA, Los Angeles, CA, USA

[c]Molecular Biology Institute, UCLA, Los Angeles, CA, USA

[d]Department of Chemistry and Biochemistry, UCLA, Los Angeles, CA, USA

*Contributed equally, order determined by drawing straws

Running Head: Atomic structures of HSV-1 A-, B-, C- and D-capsids

#Address correspondence to Z. Hong Zhou, hong.zhou@ucla.edu (phone 310-694-7527)

Abstract word count: 250

Significance word count: 150

Text word count: 4746

Abstract

Three capsid types have been recognized from the nuclei of herpesvirus-infected cells: empty A-capsids, scaffolding-containing B-capsids, and DNA-filled C-capsids. Despite progress in determining atomic structures of these capsids and extracellular virions in recent years, debate persists concerning the origins and temporal relationships among these capsids during capsid assembly and genome packaging. Here, we have imaged over 300,000 capsids of herpes simplex virus type 1 by cryogenic electron microscopy (cryoEM) and exhaustively classified them to characterize the structural heterogeneity of the DNA-translocating portal complex and their functional states. The resultant atomic structures reveal not only the expected A-, B-, and C-capsids, but also capsids with portal vertices similar to C-capsids but no resolvable genome in the capsid lumen, which we named D-capsids. The dodecameric dsDNA-translocating portal complex varies across these capsid types in their radial positions in icosahedral capsids and exhibits structural dynamics within each capsid type. In D-capsids, terminal DNA density exists in multiple conformations including one reminiscent to that in C-capsids, suggesting D-capsids are products of failed DNA retention. This interpretation is supported by varying amounts of DNA outside individual D-capsids and by correlation of capsid counts observed in situ of infected cell nuclei and those after purification. Additionally, an "anchoring" segment of the scaffold protein is resolved interacting with the portal baskets of A- and B-capsids but not D- and C-capsids. Taken together, our data indicate that A-capsids arise from failed DNA packaging and D-capsids from failed genome retention, clarifying the origins of empty capsids in herpesvirus assembly.

Significance

As the prototypical herpesvirus, herpes simplex virus 1 (HSV-1) exhibits a global seroprevalence of 67% and approaching 90% in some localities. Herpesvirus infections can cause devastating cancers and birth defects, with HSV-1 infections leading to cold sores among the general population worldwide and blindness in developing nations. Here, we present atomic structures of the capsids sorted out from the nuclear isolates of HSV-1 infected cells, including the previously recognized A-, B-, and C-capsids, as well as the newly identified D-capsid. The structures show the details of protein-protein and protein-DNA interactions within each capsid type and the positional and interactional variability of the viral DNA-translocating portal vertex among these capsids. Importantly, our findings suggest that A-capsids are products of failed dsDNA packaging and D-capsids of failed genome retention. Together, the high-resolution 3D structures clarify the processes of genome packaging, maintenance, and ejection during capsid assembly, which are conserved across all herpesviruses.

Introduction

First described by the Greek historian Herodotus over 2000 years ago, human herpes simplex virus 1 (HSV-1) is a highly adapted linear double-stranded DNA (dsDNA) virus which evolved alongside humans since our last common ancestor (1, 2). HSV-1 is a prototypical member of the *Herpesviridae* family and infects two-thirds of humans worldwide, with regional seroprevalence as high as 90% (3). Infection typically presents as mild symptoms in healthy individuals, such as cold sores in the orofacial region, but can lead to serious disease amongst the immunocompromised or immune naïve. Indeed, HSV-1 is the leading cause of lethal encephalitis in the United States and a major cause of blindness worldwide, and the lack of efficacious treatments or prophylactics makes HSV-1 a serious global health challenge (4, 5).

Ever since the first 30-40 Å resolution 3D reconstructions of HSV-1 capsids some 35 years ago (6, 7), significant progress has been made in determining and understanding the virion structure and genome packaging (8-10). HSV-1 packages its 152 kbp genome into an icosahedral T=16 nucleocapsid 125 nm in diameter, surrounded by a pleomorphic tegument layer and glycoprotein-decorated lipid envelope. The HSV-1 nucleocapsid is composed of 955 copies of the major capsid protein (MCP) arranged into 150 hexons and 11 pentons that decorate icosahedral 5-fold (I5) vertices. A unique portal vertex exists at a twelfth I5 vertex through which the genome is translocated during assembly. 320 triplex protein complexes (Tri), each composed of one Tri1 and two Tri2 (Tri2A & Tri2B) subunits rivet MCP capsomers (i.e. hexons and pentons) together, while small capsid proteins (SCP) decorate the top of each MCP tower. Capsid vertex specific components (CVSC) buttress pentonal MCPs and their neighboring Tri complexes and stabilize the capsid (10).

During infection, HSV-1's lipid envelope fuses with the host cell membrane and releases tegument proteins and the nucleocapsid into the cytosol (11, 12). The nucleocapsid travels along microtubules and docks at a nuclear pore complex, where the viral genome ejects into the

nucleus for transcription (13-16). Nascent structural proteins including Tri1, Tri2, MCP, SCP,

scaffolding protein (pUL26.5), protease-containing scaffolding protein (pUL26), and portal

protein (pUL6) are then translated in the cytosol and trafficked back to the nucleus. Here, portal

proteins assemble into a dodecameric basket-like structure that is thought to seed the assembly

of scaffolding proteins, in turn driving the formation of spherical procapsids composed of MCP,

SCP, and triplexes (17-22). A separate protein complex, the terminase, associates with the

dodecameric portal and initiates genome packaging and the concurrent autoproteolytic cleavage

of the scaffold, although the C-terminal 25 residues of the scaffolding proteins remain attached

to the luminal capsid surface (23-27). Subsequent capsid maturation steps result in a range of

outcomes: successful genome packaging produces C-capsids, while aberrant byproducts also

form, like B-capsids that have angularized but fail to release scaffolding (28) and A-capsids that

have released scaffolding but generally do not contain visible genome. All capsid types exit the

nucleus into the cytosol, where they undergo additional tegumentation and envelopment. C-

capsids ultimately become infectious virions, while A- and B-capsids are released as non-

infectious enveloped particles (NIEPs) (29). A-, B-, and C-capsids are readily distinguishable in

singular electron micrographs by their capsid lumens, as C-capsids have a fingerprint-like

appearance created by the genome, B-capsids have an obvious scaffolding core, and A-capsids

appear largely empty. Even with a high degree of speciation amongst different herpesviruses

subfamilies (30-32), capsid assembly leading to the formation of A-, B-, and C-capsids within

the host nucleus is largely conserved.

Despite the progress outlined above, debate persists concerning temporal origins and

relationships amongst the various capsid outcomes, reflecting a lack of granular understanding

of the capsid assembly pathway. And while previous structural studies of DNA-devoid capsids

from herpes simplex virus type 2 (HSV-2), varicella zoster virus (VZV), and human

cytomegalovirus (HCMV) have shown additional characteristics specific to different capsid

types—for example, that A- and B-capsid portals sit lower within the capsid vertex than the

"elevated" C-capsid portals (33-35)—little is known about the underlying conformational

changes and protein-protein interactions that contribute to such differences.

Here, we used extensive computational sorting with an unprecedentedly large dataset of

capsid images to resolve the first high-resolution asymmetric reconstructions of the HSV-1 A-,

B-, and C-capsids derived from the nuclei of infected cells. In addition to features consistent with

previous herpesvirus structures (33, 34, 36), our dataset reveals a previously unidentified capsid

end product. These new capsids feature C-capsid-like "elevated" portals and, in many cases,

terminal genome within the portal translocation channel, but no organized genome in the capsid

lumen. We refer to these capsids as "D-capsids" because their C-capsid-like portal suggests

they were originally C-capsids, but the absence of most of the genome indicates that they are

degraded. Meanwhile, in high-resolution reconstructions of "lowered" A- and B-capsid portal

baskets, we identified densities corresponding to scaffolding proteins where they bind to the

portal. Additionally, we resolved the global arrangement of the capsid-bound fragment of the

scaffolding protein attached to the inner walls of B-capsids. These results provide novel insights

into capsid assembly, genome maintenance, and viral maturation.

Results

3D classification of capsids from host nuclei and discovery of new D-capsids

We isolated HSV-1 capsids from the nuclei of infected Vero cells and subjected them to single-

particle analysis using cryoEM to determine their 3D structures. We recorded 33,399 movies

and picked 309,503 particles. We extracted the portal vertices as sub-particles, which were

classified into 30,298 A-, 66,591 B-, and 153,080 C-capsid vertex particles and refined by

imposing C5 symmetry to 3.8, 3.7, and 3.5 Å reconstructions, respectively (Fig. S1). Using an

established symmetry expansion and relaxation computational procedure (10), we resolved the

symmetry mismatch between the C5 symmetrical vertices and C12 symmetrical portal baskets

to generate C1 reconstructions at 4.3, 4.0, and 4.1 Å for A-, B-, and C-capsid portal vertices,

respectively. These orientations were used to generate asymmetric whole-capsid

reconstructions (Fig. 1A through D) for comparison with a previously resolved virion structure

(Fig. 1E) (10). To improve the resolution of the portal baskets, we carried out separate sub-

particle reconstructions and refined them while imposing C12 symmetry, leading to portal basket

structures at resolutions of 3.6, 3.5, and 3.7 Å for the A-, B-, and C-capsids, respectively (Fig.

S1).

Classification of the presumed C-capsid portals revealed that 88% of particles lacked

both capsid luminal DNA and ring DNA (Fig. 1K and S1). While the absence of ordered luminal

genome conflicts with the canonical definition of C-capsids, these unusual portals had been

grouped with C-capsids during classification due to having extended turret helices, an elevated

portal basket, a portal cap, and genome density inside the DNA-translocating channel of the

portal, in contrast to A- and B-capsids where the turret helices are retracted, the portal basket is

lower, the cap is not present, and genome density is lacking inside the portal channel (Fig. 1F

through J) (10, 37, 38). Surprised by this result, we manually curated whole C-capsid particles,

revealing that 135,916 of 153,080 capsid particles appeared in micrographs as partially or totally

devoid of luminal genome—like A-capsids—yet still contributed to reconstructions with C-like

portal features (Fig. 1I, 1K, S1, and S2). Symmetry relaxation of these C-like portals revealed

that ~30% of particles had toroidal genome organization in the portal channel equivalent to the

genome structure in virions and C-capsids, with the other ~70% harboring less distinct DNA

density in the portal channel, perhaps due to variable arrangements (Fig. 1I and K).

These observations led us to investigate the origins of these unusual capsids with C-like

portal vertices. Previous studies have suggested that herpesvirus C-capsids isolated from nuclei

are prone to degradation (29, 33), so we wondered whether capsids with C-like portals may

represent former C-capsids in a degraded state. Indeed, we see varying levels of genome

occupation in the micrographs of capsids with C-like portals, often with genome egressing from

the capsid shell (Fig. S2). The total proportion of capsids with empty lumens in our study is

66%, which is much higher than the proportion usually observed within cell nuclei infected with

herpesviruses (Fig. 1L) (39). This high proportion of empty capsids resembles that of HCMV,

which is known to have unstable C-capsids resulting in a higher empty capsid proportion

specifically in purified samples (Fig. 1L) (29, 34, 39). These findings suggest that capsids with

C-like portals are aberrant products that arise when C-capsids fail to retain viral DNA, and we

hereafter refer to these as degraded (D-) capsids.

Atomic structures of portal complexes and their contacts with the capsid shell

Our large dataset allowed us to generate reconstructions of the portal basket for all nucleus-

derived capsid types at far higher resolution than previous herpesvirus structures, enabling

modeling of the portal protein's basket (a.a. 26-307 and 494-623) and turret helices (a.a. 337-

473) (10, 33, 35). The basket is a 12-fold symmetrical complex consisting of five domains,

including the wing (a.a. 26-62, 150-174, 222-370), stem (a.a. 271-297, 517-541), clip (a.a. 298-

307, 494-516), β-hairpin (a.a. 542-557) and crown (a.a. 62-149, 175-221, 558-623) (Fig. 2A).

Ten of the twelve monomers from the basket are linked to the turret helices, each of which

consists of one long helix extending towards the portal cap and connecting to two smaller

helices arranged in a small helical bundle. The two non-contributing portal monomers are

presumably disordered or flexible beginning at their turret sequences (Fig. 2A). In C-capsids, D-

capsids, and virions, the turret helices are fully extended, whereas in A- and B-capsids, they are

retracted and positioned ~65Å lower (Fig. 1F through J). These features are consistent with

previous herpesvirus structures (35, 40, 41).

159   In addition to the basket and turrets, the portals of all capsid types feature 10 helical

densities—five helix pairs and five lone helices—originating from the N-terminal wing domains

of 10 of the 12 portal subunits (Figs. 1F through I and 2B through C). These appear to anchor

the portal basket to the capsid shell (Fig. 2B and C), forming an arrangement similar to the "10-

helix anchor" seen in HCMV (Li et al., 2023). The helix pairs interface with hydrophobic pockets

defined by MCP Johnson-folds and dimerization domains adjacent to the portal (a.a. 143-145,

165-176, 342-346), while the lone helices interface with the leucine-rich Tri1 N-terminal anchors

(a.a. 82-92) and MCP Johnson-fold domains (a.a. 160-170) (Fig. 2D through 2G). When viewed

from outside the capsid, two of the twelve portal subunits corresponding to the 5 and 10 o'clock

positions did not have N-terminal anchors (Fig. 2C). This 12-fold to 5-fold symmetry mismatch is

similar to that observed in the turret helices, where two portal subunits are conspicuously absent

from the turret arrangement.

Portal basket positioning and flexibility varies across capsid types

The flexibility of the portal basket is essential to accommodate the structural changes and

mechanical demands associated with genome packaging, ejection, and the transition between

capsid states. To characterize the portal's structural variability, we used the 3D flexible

refinement (3DFlex) tool from cryoSPARC to model the movement of the portal basket relative

to the capsid shell (42). All capsid types showed some variability in portal basket position and

rotation relative to the capsid shell. A- and B-capsids varied in portal height by ~2 and ~1 nm

respectively, could rotate up to 13 degrees relative to the capsid shell, and exhibited up to 1 nm

of portal basket "sway", due to flexibility between the clip and nearby wing and stem domains

(Fig. 3A, 3B, and Movie S1 through 4). C-capsid portals did not vary in height but were still

capable of 9-degree rotations and ~1 nm of portal basket sway (Fig. 3C and Movie S5 though

6). D-capsid portals had 4 Å variability in height but otherwise resembled C-capsid portals (Fig.

3D and Movie S7 through 8). These findings reveal distinct patterns of portal basket variability,

with A- and B-capsids exhibiting greater positional and rotational flexibility compared to the

more constrained movement observed in C- and D-capsids.

186 **Scaffolding protein is anchored to A- and B-capsid portal baskets and B-capsid shells via**

187 **conserved hydrophobic motifs**

188 The C12 reconstruction of both A- and B-capsid portal baskets revealed a small hook-like

189 density with large aromatic side chains and sharp turns wedged into a hydrophobic pocket

190 between the crown domains of each portal monomer (Fig 4A through B and 4F through G). This

191 density is weaker in A-capsids, but still conspicuous, whereas it is completely absent in C-

192 capsids, D-capsids, and mature virions. (Fig. 4C through E, 4H through J). It neighbors W90

193 and W127 in the portal monomers, known to be essential for portal-scaffold interactions (43).

194 Investigators previously showed that portal protein residues 449-457—particularly Y451, P452,

195 and E454 (43, 44)—are essential to the portal-scaffold interaction and thus efficient capsid

196 assembly (43). Indeed, we found the scaffolding protein residues 449-455 (PYYPGEA,

197 numbered relative to pUL26) to be a good fit for the small density (Fig. 4K and L). Based on our

198 model, P452 interacts with the essential residues W127 and L131 (both adjacent to W90) from

199 the first portal monomer, whereas Y451 forms a hydrophobic interface with Y83 and Y608 from

200 the second portal monomer (Fig. 4L and M). Further, E454 from the scaffolding appears to

201 stabilize Y608 via hydrogen bonding, buttressing the hydrophobic interactions between the

202 three aromatic residues (Fig. 4L). Sequence alignment of scaffolding and portal proteins

203 showed that Y608 and Y83 are highly conserved across herpesviruses, suggesting this motif is

204 essential for portal-scaffold association (Fig. 4N).

205 Despite the high-resolution details revealed in residues 449-455 of the scaffolding

206 protein, the rest of the scaffolding complex remained poorly resolved both around the portal

207 basket and capsid lumen (Fig. 1B, and 5A through C). Our asymmetric reconstruction of B-

208 capsids shows that the portal basket is encircled by large prominent densities (Fig. 1B), which

209 we attribute to scaffolding protein that remains attached to the capsid shell (Fig. 5A). Further,

210 these densities, and others similar, appear to originate from local 3-fold axes near the positions

of proximal Tri1 N-terminal anchors (Fig. 5A), consistent with observations of the scaffolding

anchor domain in HCMV (34) (Fig. 5C). These densities are noticeably absent underneath the

local 3-fold axes immediately adjacent to pentons (Fig. 5A through 5C).

Discussion

Characterizing herpesvirus capsid assembly requires thorough sampling to capture diverse

intermediates and end products in viral replication. Here, we used single-particle cryoEM to

resolve the first near-atomic resolution asymmetric structures of HSV-1 capsids derived from the

nuclei of infected cells. Our large dataset enabled the identification of a new capsid form, the D-

capsids, and provided the highest resolution detail of a herpesvirus portal to date, including its

interactions with the scaffold and dynamics with the capsid shell. These findings shed light on

longstanding questions about how the genome impacts portal basket position, the portal's role

in procapsid formation, the portal-scaffold interaction, and the origins of empty capsids.

The presence or absence of genome within the portal translocation channel appears to

be related to the position of the portal basket. C-capsid portal baskets are positioned higher at

the portal vertex compared to their A- and B-capsid counterparts, which has been attributed to

the densely packed genome pushing the portal up against the capsid shell (34). However, our

D-capsid portal structures have the same portal basket position as C-capsids, despite being

mostly devoid of genome (Fig. 1K). These observations together suggest that genome pressure

alone does not determine portal basket height, and the elevated portal position observed in C-

and D-capsids may instead be an irreversible step during capsid maturation.

The portal is important for both procapsid assembly and genome packaging, but the

structural basis enabling these dual roles had not been previously characterized. Our analyses

of portal basket variability revealed that non-mature A- and B-capsid portals permit more

rotational flexibility than mature-state C-capsid portals, which may be important for the dynamic

process of genome packaging, during which torque is applied to the genome substrate (34, 45,

46). Additionally, although A- and B-capsid portal baskets can vary in elevation by up to ~2 nm,

they cannot descend to the position of the procapsid portal basket, which is further situated 4.5

nm lower with respect to the capsid shell (Fig. 5D) (47). Indeed, the linker residues connecting

the A- and B-capsid portal baskets to their capsid-docked N-terminal anchors are too short to

permit such a 4.5-nm shift, suggesting that the procapsid basket must not be initially attached to

the capsid shell via N-terminal anchors. This suggests that during angularization, the portal

basket moves away from the scaffold core, while its N-terminal anchors attach irreversibly to the

capsid shell, limiting the basket's positional variability and preventing it from descending back to

its procapsid position. Altogether, the ability of the portal basket to begin in a lowered

conformation in procapsids may be critical for proper interaction with the scaffold and thus the

nucleation of capsid assembly, while the subsequent attachment of portal N-terminal anchors

post-angularization restrains the basket near the capsid shell, enabling it to sustain robust

terminase-driven forces during genome packaging.

Our structures unveil the scaffolding proteins' interactions with the portal and capsid

shell, which were previously suggested to be important for viral assembly (43, 48-50) but lacked

structural detail. Herpesvirus procapsid assembly is promoted by scaffold-scaffold interactions,

enabling the progressive addition of scaffold-capsid complexes into partially formed capsids

(19). Our B-capsid structure shows scaffolding densities emanating from under local 3-fold axes

in close proximity to Tri1 N-anchors, consistent with observations in protease-null mutants (51).

While their exact interface to the B-capsid luminal wall remains unclear, the binding pocket

harboring the Tri1 N-anchor is known to be promiscuous, variably binding the portal N-anchor in

our study and in HCMV (34) as well as isomeric forms of MCP Johnson-fold components at

penton locations (9, 10). This raises the possibility that during early procapsid assembly stages,

these pockets may bind scaffold protein C-terminus, which may later be displaced by Tri1 N-

terminal anchors after scaffolding protease activation (34, 51, 52). In addition to scaffold-capsid

interactions, our structure shows high-resolution details of the scaffold-portal interface, enabling

the first atomic model of a portion of the scaffolding protein and revealing its tightly packed,

hydrophobic nature nestled between portal monomers (Fig. 4F, G, and K through M). This

finding is likely relevant in other herpesviruses, as the relevant residues on the scaffolding and

portal proteins are conserved (Fig. 4N), in addition to direct structural evidence of a portal-

scaffold interaction at an equivalent motif in HCMV (34). Thus, this interface may serve as an

attractive target for the development of pan-herpesvirus inhibitors of capsid assembly.

Prior to the current study, A-capsids were thought to result from unsuccessful genome

packaging, but their exact origins were uncertain. Our analyses show that the portal structures

of A-capsids still contain scaffolding hook densities, suggesting that these capsids were likely

never packaged with a genome, as even some amount of genome content in the capsid lumen

appears to displace the portal-bound scaffold. However, because D-capsids generally appear

with empty lumens in micrographs, it is likely that D-capsids have been observed in past studies

of herpesviruses but were incorrectly grouped with A-capsids. Therefore, the "empty" capsids

may be formed at various stages of assembly, both before and after the completion of genome

packaging for A- and D-capsids respectively (Fig. 6), providing partial validity to the A-capsid

multi-origin model put forward in Kaposi's sarcoma-associated herpesviruses (53).

In summary, 3D structural classification enabled by an unprecedentedly large dataset of

HSV-1 capsid images has led to the discovery of a capsid representing a temporally distinct

state compared to previously described capsids, and alongside our high-resolution structures of

the portal vertices, provide new insight into herpesvirus capsid assembly. Our structures reveal

atomistic details of portal-type specific interactions and more clearly delineate A-, B-, C-, and

newly defined D-capsids. The identification of D-capsids as a distinct end product resulting from

C-capsid degradation clarifies the nature of observed empty capsids and conformational

changes underpinning herpesvirus genome packaging and ejection. A more complete

choreography of herpesvirus assembly inside host nucleus (Fig. 6) and their atomic structures

also offers a framework for the development of structure-based antiviral therapies targeting

portal function.

Methods

Virus strain and cell line

The HSV-1 strain KOS (VR-1493; ATCC, Manassas, VA, USA) was used in this study. African

green monkey kidney cells (Vero) (CCL-81; ATCC, Manassas, VA, USA) served as host for

virus propagation and were maintained in culture media DMEM (Cat No. MT10-017-CV;

CORNING, New York, NY, USA) supplemented with 10% heat-inactivated fetal bovine serum

(FBS) (Cat No. S11150H; R&D systems, Minneapolis, MN, USA), Penicillin (100 IU/mL) and

Streptomycin (100 µg/mL) (Pen/Strep) (Cat No. 30-002-Cl; CORNING, New York, NY, USA), at

37 ˚C in a humidified atmosphere of 5% carbon dioxide (CO_2) and 95% oxygen O_2).

Preparation of HSV-1 capsids

Vero cells at 90-95% confluence, grown in 18 T175 (175 cm^2 area) tissue culture flasks, were

infected with 0.4 mL of HSV-1 stock dispersed in 38 mL of freshly supplemented cell culture

media per flask. After 48 h post-infection the cells were harvested and centrifuged at 10,000 × g

for 20 min in a Fiberlite F14-6 × 250y Fixed-Angle Rotor (Thermo Fisher Scientific, Waltham,

MA, USA) at 4 ˚C. All subsequent sample purification steps were carried out at 4 ˚C. The cell

pellet was washed with phosphate buffered saline (PBS) of pH 7.4 (Cat No. 10010049; GIBCO,

Carlsbad, CA, USA), and centrifuged at 2,000 × g for 10 min in a SX4750 Swing Bucket Rotor

(Beckman Coulter Life Sciences, San Jose, CA, USA). To remove the cell membrane and

pelleting the nucleus, the cells were resuspended in 40 mL of membrane lysis buffer (PBS pH

7.4, 0.5 % NP40, 1 X Protease Inhibitor cocktail), kept in ice for 30 min (mixing by gentle

inversion of the sample a few times every 10 min) and centrifuged at 2,000 × g for 15 min in a

SX4750 swing bucket rotor (Beckman Coulter Life Sciences, San Jose, CA, USA); this step was

repeated twice. The nuclear pellet was resuspended in 25 mL of extraction buffer (PBS pH 7.4,

1 mM EDTA, 1X protease inhibitor cocktail), and placed on orbital shaker at low speed for 40

min. Nuclear envelope was lysed via mechanical sheering during 5 consecutive passes of the

suspension through a 23-gauge needle. The nuclear lysate was centrifuged at 10,000 × g in a

SW 28 Ti swing bucket rotor (Beckman Coulter, Brea, CA, USA) for 1 h, to remove membrane

components. The clarified supernatant was placed atop a 30-50% sucrose cushion and

centrifuged at 100,000 x g in a SW 28 Ti S swing bucket rotor (Beckman Coulter, Brea, CA,

USA) for 1 hr. 1 mL of solution was carefully drawn from the 30-50% sucrose interphase, diluted

in 10 mL of PBS pH 7.4 with 1 mM EDTA, and pelleted at 100,000 x g in a SW 28 Ti swing

bucket rotor (Beckman Coulter, Brea, CA, USA) for 1 hr. The pellet was then carefully dispersed

in 15 µL of of PBS pH 7.4 with 1 mM EDTA, preceding to cryoEM grid preparation.

CryoEM of HSV-1 capsids

HSV-1 capsids were prepared for cryoEM analysis by applying 2.5 µL of sample to 200 mesh

holey quantifoil carbon grids which had been glow discharged for 30 seconds. Each grid was

flash frozen in liquid ethane using a custom-made plunge-freezing device prior to transfer and

storage in liquid nitrogen. CryoEM movies were recorded using a Titan Krios G1 electron

microscope (FEI) upgraded with a K3 direct electron detector and Gatan Imaging Filter (GIF)

and operated at 300 keV. Movies were recorded in super-resolution mode at a nominal

magnification of 81,000 X with a calibrated pixel size of 0.55 Å at the specimen level and

estimated electron dose of ~45 e$^-$/Å^2. Using SerialEM, 33,399 TIF movies were collected.

CryoEM image processing

Image alignment, motion correction, and contrast transfer function (CTF) estimation were

performed using MotionCor2 and GCTF respectively on 2 X binned micrographs (54, 55). Topaz

automated particle picking was performed to pick all 309,503 HSV-1 nucleocapsids within the

dose-weighted micrographs, preceding import to RELION (56, 57). To reduce computation time,

capsids were boxed and extracted with 1/4 cropping in Fourier space before being subjected to

3-dimensional refinement with icosahedral symmetry enforced. This yielded an initial

reconstruction of all capsids at ~8.8 Å.

We next combined stepwise symmetry expansion and relaxation with sub-particle

reconstructions to both improve local resolution and identify the asymmetric portal vertices on

each capsid as described previously (10). Briefly, we used RELION's symmetry expansion utility

with I3 symmetry selected to generate 60 equivalent orientations for each capsid particle with

different combinations of Euler angles for rotation, tilt and psi. To remove redundant rotations

about the Z-axis, which have identical tilt and psi angles we kept only 1 of each set of 5 copies,

to yield 12 unique orientations for each particle. Each orientation could then be used to identify

the position of one 5-fold vertex in Cartesian coordinates, so that there are 12 in total for each

particle which can be reextracted for processing as sub-particles. Portal sub-particles identified

for particles whose portals were in frame during movie recording, using exhaustive 3D

classification in RELION. This yielded 249,897 portal particles, less than the number of total

capsids as some capsids were only partially in frame in the electron micrographs (Fig. S1).

Portal particles were then differentiated by their turret helices and separated into "C-like" and "B-

like" portal groups. Next the whole particles were manually sorted to identify A-capsids, and

these were separated from C-like and B-like portal groups, yielding C-portals, B-portals, B-like

A-portals, and C-like portals (D-portals).

Following sorting, sub-particle stacks were transferred to cryoSPARC for further

processing (58). Particles were initially subjected to 3D refinement with imposed C5 symmetry,

which yielded resolutions of 3.8, 3.7, and 3.5 Å respectively. We then continued with stepwise

symmetry expansion and relaxation to resolve each portal types 12-fold symmetrical basket to

3.6, 3.5, 3.7 Å respectively (10). In the case of C-particles we also employed this technique to

resolve the C1 reconstruction of the genome segment within the portal.

Scaffolding density reconstructions

During processing of complete B-capsids, cryoEM map density was observed along the luminal wall of the capsid which appeared to originate from the faces of the icosahedron and beneath the portal vertex. To better resolve these densities, we performed 3D classification of B-portal sub-particles which had been symmetry expanded using the *symmetry expansion* utility in cryoSPARC and a focused classification using a mask which included regions beneath the capsid shell. This yielded large densities which appeared to be protruded regions near the triplex N-terminal anchor and surround the portal baskets in portal sub-particles.

Model building

Atomic modeling of the HSV-1 peripentonal and portal asymmetric units has been described previously (10, 59). For both assemblies, we modeled previously unresolved loops of MCP, Tri1, Tri2A, and Tri2B combining the *Build Structure* function in ChimeraX and *real-space refinement* in Coot (60, 61). To model the complete asymmetric units, we utilized sub-particle maps from the icosahedral vertices and faces of the viral capsids to assign new helical densities beneath triplexes as the N-terminal anchor of Tri1. We resolved intermolecular clashing in our updated portal and penton asymmetric units using *ISOLDE* from ChimeraX and iterative refinement in Phenix (62).

We flexibly fitted published ab-initio models of the portal dodecamer (10) into our sub-particle portal vertex maps for A-, B-, and C-capsids in *ISOLDE* (63). To model structural flexibility, we fit these updated portal basket models into sub-particle 3D Flex maps of B- and C-baskets.

Data availability

The atomic models have been deposited in the PDB with accession codes: XXXX. The cryoEM map has been deposited in the EMDB with accession codes: EMD-XXXXX.

Acknowledgements

This research was supported in part by grants from the National Institutes of Health

(R01DE025567 to Z.H.Z and T.-T.W.). J.Z. acknowledges support from an Interdisciplinary

Training in Virology and Gene Therapy training grant (NIH 5T32AI060567). We acknowledge

the use of resources in the Electron Imaging Center for Nanomachines supported by UCLA,

Core Center of Excellence in Nano Imaging (CNI) at USC, and grants from NIH (S10RR23057,

S10OD018111, U24GM116792, and T32GM145388) and NSF (DBI-1338135). We also

acknowledge use of resources from the UCLA AIDS Institute, the James B. Pendleton

Charitable Trust, and the McCarthy Family Foundation.

394 Author contributions

395 Z.H.Z. conceived the project. A.C.A-C grew and isolated virus, prepared cryoEM samples and

396 obtained cryoEM images; A.S. processed the cryoEM images and determined the structures

397 with assistance from Y.-T.L. A.S., S.K., E.C. and Z.H.Z. interpreted the results and wrote the

398 manuscript; all authors reviewed and approved the manuscript.

399 Conflict of Interest

400 The authors declare that they have no conflicts of interest.

401 References

1.   Wertheim JO, Smith MD, Smith DM, Scheffler K, Kosakovsky Pond SL. 2014.
Evolutionary Origins of Human Herpes Simplex Viruses 1 and 2. *Mol Biol Evol* 31:2356-
2364.

2.   McGeoch DJ. 2018. Human Herpesviruses. *Adv Exp Med Biol* 1045:31-32.

3.   James C, Harfouche M, Welton NJ, Turner KM, Abu-Raddad LJ, Gottlieb SL, Looker KJ.
2020. Herpes simplex virus: global infection prevalence and incidence estimates, 2016.
Bull World Health Organ 98:315-329.

4.   Levitz RE. 1998. Herpes simplex encephalitis: A review. *Heart Lung* 27:209-212.

5.   Farooq AV, Shukla D. 2012. Herpes Simplex Epithelial and Stromal Keratitis: An
Epidemiologic Update. *Surv Opthalmol* 57:448-462.

6.   Schrag J, Prasad B, Rixon F, Chiu W. 1989. Three-dimensional structure of the HSV1
nucleocapsid. *Cell* 56:561.

7.   Booy FP, Newcomb WW, Trus BL, Brown JC, Baker TS, Steven AC. 1991. Liquid-
crystalline, phage-like packing of encapsidated DNA in herpes simplex virus. *Cell*
64:1007-15.

8.   Zhou ZH, Dougherty M, Jakana J, He J, Rixon FJ, Chiu W. 2000. Seeing the herpesvirus
capsid at 8.5 Å. *Science* 288:877-80.

9.   Dai X, Zhou ZH. 2018. Structure of the herpes simplex virus 1 capsid with associated
tegument protein complexes. *Science* 360.

10.   Liu Y-TT, Jih J, Dai X, Bi G-QQ, Zhou ZH. 2019. Cryo-EM structures of herpes simplex
virus type 1 portal vertex and packaged genome. *Nature* 570:257-261.

11.   Roizman B, Knipe DM, Whitley RJ. 2007. Herpes simplex viruses p2502-2601. *In* Knipe
DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE (ed), *Fields*
Virology 2.

12.   Liu F, Zhou ZH. 2007. Comparative virion structures of human herpesviruses. *In* Arvin A,
Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R, Yamanishi K (ed),
Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis, Cambridge.

13.   Sodeik B, Ebersold MW, Helenius A. 1997. Microtubule-mediated Transport of Incoming
Herpes Simplex Virus 1 Capsids to the Nucleus. *J Cell Biol* 136:1007-1021.

14.   Ojala PM, Sodeik B, Ebersold MW, Kutay U, Helenius A. 2000. Herpes Simplex Virus
Type 1 Entry into Host Cells: Reconstitution of Capsid Binding and Uncoating at the
Nuclear Pore Complex In Vitro. *Mol Cell Biol* 20:4922-4931.

15.   Döhner K, Cornelius A, Serrero MC, Sodeik B. 2021. The journey of herpesvirus capsids
and genomes to the host cell nucleus. *Curr Opin Virol* 50:147-158.

16.   Peng L, Ryazantsev S, Sun R, Zhou ZH. 2010. Three-dimensional visualization of
gammaherpesvirus life cycle in host cells by electron tomography. *Structure* 18:47-58.

17.   Brown JC, Newcomb WW. 2011. Herpesvirus capsid assembly: Insights from structural
analysis. *Curr Opin Virol* 1:142-149.

18.   Motwani T, Lokareddy RK, Dunbar CA, Cortines JR, Jarrold MF, Cingolani G, Teschke
CM. 2017. A viral scaffolding protein triggers portal ring oligomerization and
incorporation during procapsid assembly. *Science Advances* 3.

19.   Newcomb WW, Homa FL, Thomsen DR, Booy FP, Trus BL, Steven AC, Spencer JV,
Brown JC. 1996. Assembly of the Herpes Simplex Virus Capsid: Characterization of
Intermediates Observed During Cell-free Capsid Formation. *J Mol Biol* 263:432-446.

20.   Newcomb WW, Thomsen DR, Homa FL, Brown JC. 2003. Assembly of the Herpes
Simplex Virus Capsid: Identification of Soluble Scaffold-Portal Complexes and Their
Role in Formation of Portal-Containing Capsids. *J Virol* 77:9862-9871.

21.   McElwee M, Vijayakrishnan S, Rixon F, Bhella D. 2018. Structure of the herpes simplex
virus portal-vertex. *PLoS Biol* 16:e2006191.

22.   Gong D, Dai X, Jih J, Liu Y-T, Bi G-Q, Sun R, Zhou ZH. 2019. DNA-Packing Portal and
Capsid-Associated Tegument Complexes in the Tumor Herpesvirus KSHV. *Cell*
178:1329-1343.e12.

23.   Gao M, Matusick-Kumar L, Hurlburt W, Ditusa SF, Newcomb WW, Brown JC, McCann
PJ, Deckman I, Colonno RJ. 1994. The protease of herpes simplex virus type 1 is
essential for functional capsid formation and viral growth. *J Virol* 68:3702-3712.

24.   Preston VG, Coates JAV, Rixon FJ. 1983. Identification and Characterization of a
Herpes Simplex Virus Gene Product Required for Encapsidation of Virus DNA. *J Virol*
45:1056-1064.

25.   Thomsen DR, Newcomb WW, Brown JC, Homa FL. 1995. Assembly of the herpes
simplex virus capsid: requirement for the carboxyl-terminal twenty-five amino acids of
the proteins encoded by the UL26 and UL26.5 genes. *J Virol* 69:3690-3703.

26.   Newcomb WW, Juhas RM, Thomsen DR, Homa FL, Burch AD, Weller SK, Brown JC.
2001. The UL6 Gene Product Forms the Portal for Entry of DNA into the Herpes Simplex
Virus Capsid. *J Virol* 75:10923-10932.

27.   Draganova EB, Valentin J, Heldwein EE. 2021. The Ins and Outs of Herpesviral
Capsids: Divergent Structures and Assembly Mechanisms across the Three Subfamilies.
Viruses 13:1913.

28.   Zhou ZH, Chiu W, Haskell K, Spears H, Jr., Jakana J, Rixon FJ, Scott LR. 1998.
Refinement of herpesvirus B-capsid structure on parallel supercomputers. *Biophys J*
74:576-88.

29.   Tandon R, Mocarski E, Conway J. 2015. The A, B, Cs of Herpesvirus Capsids. *Viruses*
7:899-914.

30.   Zhou ZH, Chen DH, Jakana J, Rixon FJ, Chiu W. 1999. Visualization of tegument-capsid
interactions and DNA in intact herpes simplex virus type 1 virions. *J Virol* 73:3210-8.

31.   Yu X, Shah S, Lee M, Dai W, Lo P, Britt W, Zhu H, Liu F, Zhou ZH. 2011. Biochemical
and structural characterization of the capsid-bound tegument proteins of human
cytomegalovirus. *J Struct Biol* 174:451-60.

32.   Dai X, Gong D, Wu TT, Sun R, Zhou ZH. 2014. Organization of Capsid-Associated
Tegument Components in Kaposi's Sarcoma-Associated Herpesvirus. *J Virol* 88:12694-
702.

33.   Wang N, Chen W, Zhu L, Zhu D, Feng R, Wang J, Zhu B, Zhang X, Chen X, Liu X, Yan
R, Ni D, Zhou GG, Liu H, Rao Z, Wang X. 2020. Structures of the portal vertex reveal
essential protein-protein interactions for Herpesvirus assembly and maturation. *Protein*
Cell 11:366-373.

34. Li Z, Pang J, Gao R, Wang Q, Zhang M, Yu X. 2023. Cryo-electron microscopy
structures of capsids and in situ portals of DNA-devoid capsids of human
cytomegalovirus. *Nat Commun* 14:2025.

35. Cao L, Wang N, Lv Z, Chen W, Chen Z, Song L, Sha X, Wang G, Hu Y, Lian X, Cui G,
Fan J, Quan Y, Liu H, Hou H, Wang X. 2024. Insights into varicella-zoster virus
assembly from the B- and C-capsid at near-atomic resolution structures. *hLife* 2:64-74.

36. Yang Y, Yang P, Wang N, Chen Z, Su D, Zhou ZH, Rao Z, Wang X. 2020. Architecture
of the herpesvirus genome-packaging complex and implications for DNA translocation.
Protein Cell 11:339-351.

37. Newcomb WW, Cockrell SK, Homa FL, Brown JC. 2009. Polarized DNA Ejection from
the Herpesvirus Capsid. *J Mol Biol* 392:885-894.

38. Tang J, Olson N, Jardine PJ, Grimes S, Anderson DL, Baker TS. 2008. DNA Poised for
Release in Bacteriophage ø29. *Structure* 16:935-943.

39. Stevens A, Cruz-Cosme R, Armstrong N, Tang Q, Zhou ZH. 2024. Structure-guided
mutagenesis targeting interactions between pp150 tegument protein and small capsid
protein identify five lethal and two live-attenuated HCMV mutants. *Virology* 596:110115-
110115.

40. Wang N, Chen W, Zhu L, Zhu D, Feng R, Wang J, Zhu B, Zhang X, Chen X, Liu X, Yan
R, Ni D, Zhou GG, Liu H, Rao Z, Wang X. 2020. Structures of the portal vertex reveal
essential protein-protein interactions for Herpesvirus assembly and maturation. *Protein
Cell* 11:366-373.

41. Li Z, Pang J, Gao R, Wang Q, Zhang M, Yu X. 2023. Cryo-electron microscopy
structures of capsids and in situ portals of DNA-devoid capsids of human
cytomegalovirus. *Nat Commun* 14.

42. Punjani A, Fleet DJ. 2023. 3DFlex: determining structure and motion of flexible proteins
from cryo-EM. *Nat Methods* 20:860-870.

43. Yang K, Baines JD. 2009. Proline and Tyrosine Residues in Scaffold Proteins of Herpes
Simplex Virus 1 Critical to the Interaction with Portal Protein and Its Incorporation into
Capsids. *J Virol* 83:8076-8081.

44. Huffman JB, Newcomb WW, Brown JC, Homa FL. 2008. Amino Acids 143 to 150 of the
Herpes Simplex Virus Type 1 Scaffold Protein Are Required for the Formation of Portal-
Containing Capsids. *J Virol* 82:6778-6781.

45. Buch MHC, Newcomb WW, Winkler DC, Steven AC, Bernard Heymann J. 2021. Cryo-
electron tomography of the herpesvirus procapsid reveals interactions of the portal with
the scaffold and a shift on maturation. *mBio* 12:1-13.

46. Selvarajan Sigamani S, Zhao H, Kamau YN, Baines JD, Tang L. 2013. The Structure of
the Herpes Simplex Virus DNA-Packaging Terminase pUL15 Nuclease Domain

Suggests an Evolutionary Lineage among Eukaryotic and Prokaryotic Viruses. *J Virol*
87:7140-7148.

47.   Woodson M, Pajak J, Mahler BP, Zhao W, Zhang W, Arya G, White MA, Jardine PJ,
526   Morais MC. 2021. A viral genome packaging motor transitions between cyclic and helical
527   symmetry to translocate dsDNA. *Sci Adv* 7:eabc1955.

529   48.   Huffman JB, Newcomb WW, Brown JC, Homa FL. 2008. Amino Acids 143 to 150 of the
530   Herpes Simplex Virus Type 1 Scaffold Protein Are Required for the Formation of Portal-
531   Containing Capsids. *J Virol* 82:6778-6781.

532   49.   Singer GP, Newcomb WW, Thomsen DR, Homa FL, Brown JC. 2005. Identification of a
533   Region in the Herpes Simplex Virus Scaffolding Protein Required for Interaction with the
534   Portal. *J Virol* 79:132-139.

535   50.   Yang K, Baines JD. 2009. Tryptophan Residues in the Portal Protein of Herpes Simplex
536   Virus 1 Critical to the Interaction with Scaffold Proteins and Incorporation of the Portal
537   into Capsids. *J Virol* 83:11726-11733.

538   51.   Zhou ZH, Macnab SJ, Jakana J, Scott LR, Chiu W, Rixon FJ. 1998. Identification of the
539   sites of interaction between the scaffold and outer shell in herpes simplex virus-1
540   capsids by difference electron imaging. *Proc Natl Acad Sci USA* 95:2778-2783.

541   52.   Walters JN, Sexton GL, McCaffery JM, Desai P. 2003. Mutation of Single Hydrophobic
542   Residue I27, L35, F39, L58, L65, L67, or L71 in the N Terminus of VP5 Abolishes
543   Interaction with the Scaffold Protein and Prevents Closure of Herpes Simplex Virus Type
544   1 Capsid Shells. *J Virol* 77:4043-4059.

545   53.   Deng B, O'Connor CM, Kedes DH, Zhou ZH. 2008. Cryo-electron tomography of
546   Kaposi's sarcoma-associated herpesvirus capsids reveals dynamic scaffolding
547   structures essential to capsid assembly and maturation. *J Struct Biol* 161:419-427.

548   54.   Zheng SQ, Palovcak E, Armache J-P, Verba KA, Cheng Y, Agard DA. 2017.
549   MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
550   microscopy. *Nat Methods* 14:331-332.

551   55.   Zhang K. 2016. Gctf: Real-time CTF determination and correction. *J Struct Biol* 193:1-
552   12.

553   56.   Scheres SHW. 2013. Single-particle processing in RELION, p 1-21.

554   57.   Bepler T, Morin A, Rapp M, Brasch J, Shapiro L, Noble AJ, Berger B. 2019. Positive-
555   unlabeled convolutional neural networks for particle picking in cryo-electron
556   micrographs. *Nat Methods* 16:1153-1160.

557   58.   Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. 2017. cryoSPARC: algorithms for
558   rapid unsupervised cryo-EM structure determination. *Nat Methods* 14:290-296.

59.   Dai X, Zhou ZH. 2018. Structure of the herpes simplex virus 1 capsid with associated
tegument protein complexes. *Science* 360:eaao7298.

60.   Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr 66:486-501.

61.   Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE.
2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci 27:14-25.

62.   Afonine PV, Poon BK, Read RJ, Sobolev OV, Terwilliger TC, Urzhumtsev A, Adams PD.
2018. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta*
Crystallogr D Struct Biol 74:531-544.

63.   Croll TI. 2018. ISOLDE: A physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr D Struct Biol* 74:519-530.

Supplementary Information

Supplementary Table 1. CryoEM data collection

	WT HSV-1 A whole capsid (EMD-XXXXX)	WT HSV-1 B whole capsid (EMD-XXXXX)	WT HSV-1 C Whole capsid (EMD-XXXXX)	WT HSV-1 D whole capsid (EMD-XXXXX)	(EMD-XXXXX)
Data collection and processing					
Magnification	81,000	81,000	81,000	81,000	81,000
Voltage (kV)	300	300	300	300	300
Electron exposure (e–/Å^2)	45	45	45	45	45
Defocus range (μm)	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5
Pixel size (Å)	1.1	1.1	1.1	1.1	1.1
Symmetry imposed	C1	C1	C1	C1	C1
Particle images (no.)	23,307	38,494	12,839	11,054	92,533
Map resolution (Å) FSC threshold	6.7 0.143	8.71 0.143	8.91 0.143	10.1 0.143	6.64 0.143
Symmetry imposed	C1	C1	C1	C1	C1

Supplementary Table 1. CryoEM data collection

	WT HSV-1 A C5 Portal Vertex (EMD-XXXXX)	WT HSV-1 B C5 Portal Vertex (EMD-XXXXX)	WT HSV-1 C C5 Portal Vertex (EMD-XXXXX)	WT HSV-1 C5 Portal Vertex (EMD-XXXXX)	C-portal genome
Data collection and processing					
Magnification	81,000	81,000	81,000	81,000	81,000
Voltage (kV)	300	300	300	300	300
Electron exposure (e–/Å^2)	45	45	45	45	45
Defocus range (µm)	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5
Pixel size (Å)	1.1	1.1	1.1	1.1	1.1
Symmetry imposed	C1	C1	C1	C1	C1
Particle images (no.)	30,298	66,591	19,345	135,916	18,364
Map resolution (Å)	3.8	3.7	3.7	3.5	5.2
FSC threshold	0.143	0.143	0.143	0.143	0.143
Symmetry imposed	C5	C5	C5	C5	C1

Supplementary Table 1. CryoEM data collection

	WT HSV-1 A capsid basket (EMD-XXXXX)	WT HSV-1 B Capsid basket (EMD-XXXXX)	WT HSV-1 C all (C+D) Capsid basket (EMD-XXXXX)	WT HSV-1 C Capsid basket (EMD-XXXXX)	D-portal genome (EMD-XXXXX)
Data collection and processing					
Magnification	81,000	81,000	81,000	81,000	81,000
Voltage (kV)	300	300	300	300	300
Electron exposure (e–/Å^2)	45	45	45	45	45
Defocus range (µm)	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5	-1.5 to -2.5
Pixel size (Å)	1.1	1.1	1.1	1.1	1.1
Symmetry imposed	C1	C1	C1	C1	C1
Particle images (no.)	363,576	799,092	1,863,132	155,261	92,533
Map resolution (Å) FSC threshold	3.6 0.143	3.5 0.143	3.7 0.143	4.1 0.143	6.64 0.143
Symmetry imposed	C12	C12	C12	C12	C1

Figure 1: Asymmetric reconstructions and comparisons of HSV-1 A-, B-, C-, D-, and virion nucleocapsids.
(A)-(E) asymmetric reconstructions of A-, B-, C-, D-, and virion (EMDB: 9864) capsids showing empty (A,D), scaffold-containing (B), and genome-containing (C,E) capsid shells. The color bar on the right indicates radial distance from the capsid center in Å. (F)-(J) asymmetric reconstructions of the portal vertices from A-, B-, C-, D-, and virion (EMDB: 9860, 9862) capsids in A-E. (K) Comparison of genome organization in D-, C-, and virion portal vertices, showing consistent arrangement in virions, C-capsids, and some D-capsids, but symmetric, round density in another class of D-capsids. (L) graph of prevalence of empty capsids in our study of HSV-1, as well as nuclear thin-section versus purified samples for HCMV using data from (39).

Figure 2: High resolution structure of the portal basket.

(A) from left to right: resolved portions of a portal monomer colored by domain, full portal dodecamer including turret helices, and examples of model fit to cryoEM density in the clip (green), turret (cyan), and wing (blue green) domains. (B) cryoEM density of the portal vertex showing orientation of top (red) and bottom (blue) views in panels C-E. (C) Top view of the portal basket showing assignment of 10-helix anchors (smooth outer blobs) to portal subunits, with subunits missing an anchor labeled in red. (D) bottom view of MCP (lavender) and Tri1 (purple) fragments near the 10-helix anchors (dodger blue) with important domains of MCP and Tri1 labeled. (E) bottom view of MCP and Tri1 fragments near 10-helix anchors colored by hydrophobicity. (F)-(G) zoomed-in views of the lone helix (left) and helix pair (right) of the 10-helix anchors (blue outlines) overlayed with hydrophobic surface view of their binding pockets, with hydrophobic residues shown.

	Height variability	Rotation	Sway
A A-capsid portals	~2 nm	90° ~13°	~0.5 nm
B B-capsid portals	~1 Å	90° ~13°	~1 nm
C C-capsid portals		90° ~9°	~1 nm
D D-capsid portals	~4 Å	90° ~9°	~1 nm

█ Extreme conformation   █ Opposite extreme conformation

Figure 3: Portal basket flexibility and positional variability within each capsid type for HSV-1 A-, B-, C-, and D-capsids.
(A) – (D) Portal basket models fit into alternative conformations found via 3D flex analysis for each of the four capsid types, with the extreme edges of their conformational ranges represented as salmon and aqua structures. Multiple separate analyses show vertical translation (left), rotation (center), and a swaying motion of the top half of basket (right).

Figure 4: The scaffold protein interacts with the portal dodecamer. (A)-(E) ribbon diagram of modeled residues in the portal dodecamer for A-, B-, C-, D-, and virion capsids, including basket (salmon), turret helices (cyan), and where applicable, the associated genome (orange) and scaffolding protein loops (lime). (F)-(J) zoom-ins of the ribbon model and density of the portal's scaffold binding pocket for the five capsid types in (A)-(E), showing presence or absence of scaffold density (lime). (K) Residues 449-455 of the scaffolding protein fit into a density interacting with the portal. (L) Interatomic distances between atoms of scaffolding and portal at their interaction site, highlighting hydrophobic residues. (M) Scaffolding loop shown with a hydrophobicity map of the portal, showing hydrophobic spots near Y451 and P452. (N) Sequence alignment of portal and scaffolding proteins across a range of herpesviruses, demonstrating sequence conservation of residues important for portal-scaffold interactions.

Figure 5: Global arrangement of the scaffold in whole B-capsids. (A) Cut view of whole B-capsid reconstruction showing large internal scaffold densities (top left), with Tri1 and Tri2 labeled in purple and red, respectively. (B) Unwrapped view of the capsid floor showing global scaffold arrangement, with penton locations marked with black-outlined white pentagons. (C) Scaffold densities at portal (blue dashed outline) and antipodal (green dashed outline) poles. (D) Comparison of the portal basket in HSV-1 procapsids from Buch et al. with our B-capsid reconstruction, showing 4.5 nm difference in position.

Figure 6: Proposed model of herpesvirus maturation and assembly.
Virion capsids release the viral genome into the host cell nucleoplasm, where the genome is transcribed and translated to form viral proteins. These proteins are then imported into the nucleus and associate with a free portal complex via scaffold-mediated interactions, initiating assembly of the procapsid. If angularization occurs prematurely, B-capsids are formed. If angularization and scaffold digestion are completed but not genome packaging, A-capsids are formed. Successfully packaged capsids, or C-capsids, may either undergo further maturation into virions or fail to retain genome, resulting in the formation of D-capsids.

Figure S1: CryoEM image processing workflow for portal vertex reconstructions.

| A-capsid
(empty) | D-capsid
(empty) | D-capsid
(partially full) | C-capsid
(full and organized) |

Figure S2: D-capsids occur in a spectrum of genome occupancies.
(A) Typical A-capsid micrograph, appearing as an empty capsid. (B) D-capsid micrograph appearing empty, like an A-capsid. (D) Another D-capsid micrograph, appearing filled but not clearly organized. D-capsids occur as a continuum of capsid occupancies, with some empty and some partially filled, but lack the clear genome organization of C-capsids. White arrows indicate free nucleic acid strands, which are common in the sample and may be due to genome ejection in D-capsids. (D) Typical C-capsid micrograph, displaying clear organization of the genome.

Figure S3: Cross section of D-capsid portal vertex without terminal genome.

Movie S1. 3DFlex volume series of the A-capsid portal vertex, showing slight sway of the basket relative to the capsid shell, as well as substantial variability in height.

Movie S2. 3DFlex volume series of the A-capsid floor at the portal vertex, showing rotation of portal relative to the capsid shell.

Movie S3. 3DFlex volume series of the B-capsid portal vertex, showing sway of the basket relative to the capsid shell.

Movie S4. 3DFlex volume series of the B-capsid floor at the portal vertex, showing rotation of portal relative to the capsid shell.

Movie S5. 3DFlex volume series of the C-capsid portal vertex, showing sway of the basket relative to the capsid shell.

Movie S6. 3DFlex volume series of the C-capsid floor at the portal vertex, showing rotation of portal relative to the capsid shell.

Movie S7. 3DFlex volume series of the D-capsid portal vertex, showing sway of the basket relative to the capsid shell.

Movie S8. 3DFlex volume series of the D-capsid floor at the portal vertex, showing rotation of portal relative to the capsid shell.