The Silent Saboteur: How Mitochondria Shape the Long-Term Fate of the Injured Brain.

Sahan SB Kansakar[1], Sydney P Sterben[1], Charitha Anamala[1], Mitchel Thielen[1], Volha Liaudanskaya[1*]

[1]Biomedical Engineering Department, University of Cincinnati, Cincinnati, USA
[*] Corresponding author

Abstract

Traumatic brain injury (TBI) is a major risk factor for neurodegenerative diseases, including Alzheimer's disease (AD), yet the mechanistic link remains unclear. Here, we integrated human patient-derived transcriptomics with a 3D in vitro brain injury model to dissect cell-specific mitochondrial dysfunction as a driver of neurodegeneration. Comparative transcriptomic analysis at 6 and 48 hours post-injury revealed conserved mitochondrial impairments across excitatory neurons, interneurons, astrocytes, and microglia. Using a one-of-a-kind cell-specific mitochondria tracking system, we demonstrate prolonged neuronal mitochondrial fragmentation, bioenergetic failure, and metabolic instability, coinciding with the emergence of AD markers, including pTau, APP, and Aβ42/40 dysregulation. Glial mitochondria exhibited delayed but distinct metabolic dysfunctions, with astrocytes failing to maintain metabolic support and microglia sustaining chronic inflammation. These findings establish neuronal mitochondrial failure as an early trigger of injury-induced neurodegeneration, reinforcing mitochondrial dysfunction as a therapeutic target for preventing TBI-driven AD pathology.

Introduction

Traumatic Brain Injury (TBI) affects an estimated 69 million people annually [1]. TBI is a debilitating condition with multifaceted pathobiology that leads to long-lasting disabilities and morbidity. What begins as a mechanical injury quickly launches a complex cascade of poorly understood molecular and metabolic events, with intercellular interactions amplifying and progressing the injury [2,3]. Survivors of TBI face a spectrum of outcomes that range from short-term, non-incapacitating injuries to severe, debilitating sequelae. Although the primary insult may produce significant brain damage, the secondary injury cascades — including mitochondrial dysfunction, metabolic alterations, neuroinflammation, axonal degeneration, protein misfolding (alpha-synuclein (aS), amyloid precursor protein (APP), tubulin-associated unit (Tau), transactive response DNA binding protein of 43 (TDP43)), persistent inflammation, and cell death — play a critical role in injury outcomes. Chronic TBI pathology varies with patients, where some recover completely while others suffer physical and cognitive decline, leading to the onset of neurodegenerative diseases [4]. Numerous epidemiological studies have demonstrated that individuals who have sustained TBI have an elevated risk of developing AD and AD-related dementias (ADRD) [5-7]. TBI is the third strongest risk group for individuals to develop AD after age and APOE genotype [8]. While TBI-associated clinicopathological studies have demonstrated that amyloid beta (Ab) plaques and neurofibrillary tangles (NFTs) consisting of hyperphosphorylated Tau (p-tau) significantly contribute to AD/ADRD development, mechanisms are still poorly understood [8,9].

Mitochondria are a potential regulator of secondary injury progression and injury-induced AD-like neurodegeneration; they play a crucial role in controlling brain cells' activity and function through balanced mechanisms of mitochondrial fission and fusion that maintain their growth, shape, distribution, function, and structure [10]. Mitochondria dysfunction is common in many diseases, including neurodegenerative disorders, metabolic disorders, and cancer. Disruption in mitochondria function influences inflammatory pathway activation, energy production, metabolism, calcium regulation, transcription factor phosphorylation, and cell death activation [11,12]. Studies demonstrated that TBI alters mitochondria function and metabolism, implicating these alterations with secondary neurodegeneration and neuroinflammation [13,14]. Several human athletes and *in vivo* TBI studies have established alterations in mitochondria metabolism as a defining and pharmacologically targetable feature of injury [15-21]. Alterations in TCA cycle-related enzymes [22-24], mitochondria metabolism [15,17,19,25,26], fructose metabolism [27], and lipid peroxidation [28,29] were observed following injury. Studies also displayed mitochondrial dysfunction — particularly impaired mitochondrial respiration — and decreased adenosine triphosphate (ATP) production in early AD development [30,31]. In addition, mitochondria dysfunction has been linked to the accumulation of Aβ and p-Tau proteins, characteristic features of AD pathology [8,9], and is observed in early AD patients before any clinical symptoms. Various experimental models of the disease have shown that mitochondrial dysfunction plays a crucial role in initiating destructive processes, causing extensive oxidative damage to neuronal lipids, nucleic acids, and proteins in the brains of AD patients [32]. Despite overwhelming proof of injury-induced mitochondria dysfunction and its association with an AD-like neurodegenerative phenotype, some critical gaps remain: Is this mitochondria dysfunction specific or present in all cell types? Can mitochondria recover over time post-injury? Is there a correlation between cell-specific mitochondria dysfunction and AD-like neurodegenerative phenotype progression?

Although human studies are invaluable in providing information on the brain response to trauma, they struggle with causal validation of the molecular mechanisms associated with injury progression. Similarly, rodent TBI models present challenges in interrogating cell-specific contribution and molecular footprint due to the complexity and pricing of such studies [30]. Thus, there is a critical need for an alternative approach using human data to build a relevant hypothesis for validating cell-specific molecular mechanisms that drive the progression of brain injury-induced neurodegeneration and neuroinflammation.

Recent advances in biomedical engineering, with the development of protocols for induced pluripotent stem cell differentiation and spheroids/organoids generation, allowed scientific communities to study the genetic and molecular footprints of various human disorders. While organoids are exceptional in studying developmental or genetic disorders, bioengineered 3D *in vitro* models might be a better fit to study injury-associated molecular alterations as they provide mechanical strength to withstand the injury, allow manipulation of cellular composition, and can have a better resemblance of extracellular structure. Here, to study brain injury and the molecular mechanisms involved, we used an established human triculture 3D in-vitro silk scaffold as a brain model [33-37]. This model provides physiologically relevant anatomical, structural, and molecular responses that closely resemble those of the human brain, as observed through intercellular communication, extracellular matrix composition, and 3D tensile strength [33,37].

Based on human patients' data and extensive literature review, we hypothesize that injury-induced cell-specific dysregulation in neuronal mitochondria bioenergetic and metabolic functions transforms them into neurodegenerative factories that control the onset of acute neuroinflammation and neuronal damage and the widespread long-AD-like pathology. To address this hypothesis, we have integrated fluorescent tags with different wavelengths in brain cells (neurons (dsRED2), astrocytes (EGFP), and microglia (BFP2)) under mitochondria-specific promoter COX8A (cytochrome c Oxidase subunit 8A) and used these novel lines to triculture in the human 3D brain-like injury model, to unveil cell-specific alterations of mitochondria and their contribution to AD-like neurodegeneration progression post-TBI.

This study provides a comprehensive, cell-specific characterization of mitochondrial dysfunction following traumatic brain injury (TBI) and its potential role in driving Alzheimer's-like neurodegeneration. We first validated acute transcriptomic alterations by expanding our initial 24-hour analysis to 6 and 48 hours, revealing early mitochondrial stress responses, transient compensatory mechanisms, and subsequent metabolic shifts. To assess whether these alterations translate to human-relevant neurodegenerative processes, we identified cell-specific transcriptomic convergence between our 3D in vitro human brain injury model and TBI patient-derived datasets [38], demonstrating that mitochondrial impairments in neurons, astrocytes, and microglia align with those observed in human pathology. To further dissect mitochondrial dysfunction at a cellular level, we developed and validated brain cell lines with fluorescently labeled mitochondria, enabling precise visualization and functional assessment of mitochondria post-injury. Using this system, we uncovered distinct mitochondrial dysfunction trajectories across cell types, with neuronal mitochondria exhibiting persistent fragmentation and bioenergetic failure, astrocytes demonstrating delayed metabolic collapse, and microglia shifting toward sustained inflammatory activity. Notably, these mitochondrial impairments correlated with the progressive accumulation of AD-related markers, including pTau, APP, and Aβ42/40 dysregulation, reinforcing the idea that mitochondrial failure serves as an early driver of TBI-induced neurodegeneration. This study is the first to integrate cell-specific transcriptomics, live mitochondrial imaging, and bioenergetic and metabolic profiling in a human-relevant TBI model, providing unprecedented insight into how mitochondrial dysfunction may initiate and sustain long-term neurodegenerative processes.

Results

Acute mitochondria damage in human 3D in vitro brain tissue model.

A previous study [34] with a 24-hour transcriptomic dataset demonstrated detrimental alterations to molecular pathways associated with mitochondria bioenergetic and metabolic functions, specifically in Mitochondrial Complex I gene expression. Additionally, the significant upregulation of fatty acids oxidation, carnitine shuttle, and mitochondria central dogma genes was shown to be crucial in mitochondria maintenance and repair. While the original study was the foundation for providing the starting information on mitochondria alterations, it was limited to one-time points, thus not allowing for dynamic study. In this project, we first perform an extended analysis of responses at 6 and 48 hours to complete the transcriptomic profile of acute mitochondria dysfunction.

We used the CEMiTool R package [39,40] to group differentially expressed genes (DEGs) into modules at 6 and 48 hours post-injury, identifying five distinct modules. One module did not show significant changes in response to trauma and was excluded from further analysis. The remaining four modules displayed dynamic shifts in gene expression, indicating early metabolic and signaling alterations followed by activation of cell cycle regulation and inflammation (Figure 1 A-B).

Module 1, associated with the 6-hour time point, exhibited increased gene expression following injury, with COX6A2 (cytochrome c oxidase subunit 6A2) emerging as a key hub, directly linking this module to mitochondrial bioenergetic function. Module 2 showed a sharp decline in gene expression at 6 hours, primarily in genes related to cellular division, suggesting an early cell cycle arrest. However, by 48 hours, these genes were upregulated, indicating a shift toward cell proliferation and mitotic activity. Module 3 demonstrated a transient spike in transcriptional signaling pathways, including Nerve Growth Factor (NGF), Nuclear Factor kappa-light-chain-enhancer of activated B cells (NFkB), Neurotrophic Receptor Tyrosine Kinase 1 (NTRK1), and G Protein-Coupled Receptor (GPCR) - associated pathways, at 6 hours, with gene expression returning to baseline by 48 hours. Key mitochondrial hubs in this module, ATP5MGL (ATP Synthase Membrane Subunit g-Like) and NDUFC2-KCTD14 (a read-through fusion gene of NADH: Ubiquinone Oxidoreductase Subunit C2 and Potassium Channel Tetramerization Domain Containing 14), suggest a direct link between these early transcriptional responses and mitochondrial adenosine triphosphate (ATP) production. Lastly, module 5, which displayed downregulated gene expression at 48 hours, was enriched for inflammatory response pathways, with hubs including CCL3L3 (C-C Motif Chemokine Ligand 3-Like 3), Tumor Necrosis Factor (TNF), CXCL10 (C-X-C Motif Chemokine Ligand 10), CD80 (Cluster of Differentiation 80), CD86 (Cluster of Differentiation 86), Interleukin 10 (IL-10), and CCL22 (C-C Motif Chemokine Ligand 22) highlighting the activation of immune signaling at later time points.

Next, we used Gene Set Enrichment (GSEA) to analyze the MitoCarta3.0 dataset (Figure 1 C-D) to evaluate enriched metabolic pathways affected by acute post-trauma [41-44]. The overall cellular metabolic state was upregulated acutely after injury (including 6, 24, and 48 hours), alluding that injury induces hypermetabolism acutely after trauma, including pathways associated with OXPHOS, Mitochondria Maintenance, and repair, Metabolism, and Signaling. While overall cells demonstrated hypermetabolism, electron transport genes (ETC) at the acute 6-hour time point (6h CCI vs 6h Sham) demonstrated significant downregulation of Mitochondrial Complex I gene expression (Figure 1C), apart from the mitochondrially encoded NADH dehydrogenases (MT-ND) gene family (the gene family responsible for the onset of electron acceptance in the inner mitochondria membrane to initiate the proton gradient). Similar to Complex I, Complex II-IV also demonstrated downregulated genes. However, genes in Complex V were significantly upregulated.

48 hours after the injury, these transcriptomic alterations (48h CCI vs 48h Sham) return to baseline, with none of them significantly different. On the contrary, the 24-hour timepoint [34] showed

a drastic increase in the complex I genes, with a few genes being differentially expressed in the other complexes. Unlike in 6 hours, complex V was almost unchanged.

Figure 1. Transcriptomic profile demonstrated acute mitochondrial damage in human 3D *in vitro* brain tissues 6 and 48 hours after moderate injury inflicted by the controlled cortical impactor. The human 3D in vitro triculture model comprised 2 mln of neurons, 0.5 mln astrocytes, and 0.1 mln of HMC3 microglial cells. Sham (n = 5) and CCI (n = 6) at 6 hours and Sham (n = 4) and CCI (n = 4) were used to obtain data for this transcriptomic profile.

(A) Gene set enrichment analysis revealed 6 distinct expression modules associated with M1 – mitochondria function (positively associated with an injury); M2 – cell cycle progression (negatively associated at 6 hours); M3 – cellular signaling (positively associated with injury at 6 hours and negatively at 48 hours); M5 – neuroinflammation (negatively associated at 48 hours).

(B) Critical regulators of molecular mechanisms associated with M1, M2, M3, and M5 modules.

(C) Differentially expressed genes (DEGs) associated with mitochondria pathways and

(D) mitochondria electron transport chain complex genes were detected (abs(Log2(Fold Change))>0.585 and q<0.01).

Multicellular mitochondria dysfunction as the convergent signature of brain trauma between human patients and human 3D *in vitro* model.

In the previous section, we have shown that injury induces severe changes to mitochondrial function. In addition to our studies, there is a consensus in the field of brain trauma that mitochondria dysfunction is a potential regulator of injury-induced neurodegenerative alterations for disorders (Alzheimer's, Parkinson's, Chronic Traumatic Encephalopathy) [5-7].

While our study and many others suggest the involvement of mitochondria metabolism in injury-induced neurodegeneration, one question remains unknown: is this a general phenomenon, or does the mitochondria of different brain cells respond differently to brain trauma? To address whether human patients demonstrate cell-specific mitochondria dysfunction and how these responses differ, we focused on understanding cell-specific metabolic pathway alterations convergent between our bulk RNA-seq data (3D *in vitro* brain injury model) and single-nuclei RNA-seq (snRNA-seq) data (TBI human samples) [38]. We first used bulk RNA-seq data from our 3D *in vitro* brain injury model to compare with the Reactome Pathway dataset, MitoCarta 3.0 dataset, and Mitochondrial Complex Gene dataset to identify altered pathways post-injury. We then used cell-specific human patient single-nuclei RNA-Seq data to identify how similar DEGs between these two samples contribute to altered pathways.

We used a publicly available snRNA-seq dataset that collected samples from 12 critically injured or deceased patients from brain trauma and curated cell-specific (excitatory neurons, interneurons, astrocytes, and microglia) differentially expressed genes (DEGs) compared to 5 control patients [45]. Regardless of convergence or divergence, we selected similar DEGs across the datasets for STRING analysis to determine cell-specific pathways significantly altered in response to trauma (Figure 2A-B) 6 and 48 hours after injury. The results demonstrate pathways in excitatory neuronal and interneuron were mostly associated with neuronal and synaptic network repair and electrical activity 6 and 48 hours post-injury. In addition, the transcriptomic profile also demonstrated dysfunction in MAPK signaling, a pathway that is a critical regulator of mitochondria function, oxidative stress, inflammation, and apoptosis.

While neuronal cells attempt to restore their function, both glial cells demonstrated drastic changes to ATP production via the oxidative phosphorylation (OXPHOS) pathway. At 6 hours post-injury, astrocytes demonstrated alterations in the Mitogen-Activated Protein Kinase (MAPK) pathway and O-glycan biosynthesis, and microglia demonstrated ATP production-related mechanisms and potential epigenetic alterations (Figure 2A). At 48 hours post-injury, astrocytes demonstrated changes to pyrimidine, purine, and nicotinamide metabolism, as well as microglia inflammation activation. Both glial cells demonstrate an association with axonal guidance.

Next, we focused on convergent DEGs shared between human patients and our human 3D *in vitro* brain model. DEGs shared between excitatory neurons and our human 3D *in vitro* brain model demonstrate alterations in Calcium/Calmodulin-Dependent Protein Kinase II Delta (CAMK2D) [46-49] and Monoamine Oxidase B (MAOB) gene expression, with CAMK2D influencing synaptic plasticity and memory formation and MAOB contributing to oxidative stress and neurodegeneration through excessive production of hydrogen peroxide, both of which are disrupted in AD-phenotype. Though not directly connected to Alzheimer's disease (AD), some DEGs like CTSB (cathepsin B), Solute Carrier Family 7 Member 11 (SLC7A11) (glutathione metabolism), and B-cell Lymphoma 6 (BCL6) (inflammation) may facilitate neurodegeneration through oxidative stress, proteostasis, and immune responses [50]. Other convergent DEGs like Neuregulin 1 (NRG1), Low-Density Lipoprotein Receptor Class A Domain-Containing 4 (LDLRAD4), Spectrin Repeat Containing Nuclear Envelope Protein 1 (SYNE1), Kin of IRRE-Like Protein 1 (KIRREL1), Rho GTPase Activating Protein 22 (ARHGAP22), Nuclear Factor I X (NFIX), Ectodermal-Neural Cortex 1 (ENC1), SLIT-ROBO Rho GTPase Activating Protein 3 (SRGAP3), Zinc Finger Protein 365 (ZNF365), Mediator Complex Subunit 12 Like (MED12L), Neuronal Cell Adhesion Molecule (NRCAM), Tropomyosin 4 (TPM4), Sec1 Family Domain-Containing Protein 2 (SCFD2), HIVEP Zinc Finger 1 (HIVEP1), Aryl Hydrocarbon Receptor Nuclear Translocator 2 (ARNT2),

C-Type Lectin Domain Containing 16A (CLEC16A), Fos-Related Antigen 2 (FOSL2), Calcium Voltage-Gated Channel Auxiliary Subunit Beta 4 (CACNB4) may influence neurodegeneration through neuronal development, synaptic function, cytoskeletal organization, immune signaling, and transcriptional regulation.

Next, DEGs shared between interneurons and our human 3D *in vitro* brain model demonstrate alterations in HSP90AB1 and INSR that are directly linked to Alzheimer's disease (AD) [51,52]. HSP90AB1, a molecular chaperone, plays a role in protein folding and tau stabilization, which can influence tau aggregation in AD pathology. INSR (insulin receptor) drives brain glucose metabolism, which is associated with cognitive decline and amyloid-beta accumulation in AD.

Microglia cells had convergent DEGs directly linked to Alzheimer's disease, such as STIP1, DTL, and indirectly, CTSB [53-55]. Other DEGs commonly associated with epigenetic regulation of histones or DNA methylation are UHRF1, ATAD5, NSD2, and SMC4 [56-58]. Given that microglia are believed to be at the core of Alzheimer's disease onset and progression, these findings suggest that injury-induced microglial mitochondrial dysfunction and transcriptional alterations may contribute to the early priming of neuroinflammatory and neurodegenerative pathways, potentially accelerating disease progression in vulnerable individuals.

At last, some DEGs shared between astrocytes and our human 3D *in vitro* brain model play roles in transcription regulation (RFX2, BCL6, MED12L, KNDC1), cytoskeletal organization (SYNE1, BCAR3), ubiquitination (LRSAM1, UBE2H), DNA repair (BRIP1, FANCA, GINS3), metabolic transport (SLC7A11, SLC25A18, PSAT1), and signaling (ABR, ARHGAP39, ARHGAP26, LTBP1) [59-62]. Though not directly connected to Alzheimer's disease (AD), some DEGs like CTSB (cathepsin B), SLC7A11 (glutathione metabolism), and BCL6 (inflammation)[50] may facilitate neurodegeneration through oxidative stress, proteostasis, and immune responses.

While studying convergence between human patients and our human 3D *in vitro* brain tissue model, we have discovered two emerging patterns: cell-specific mitochondria dysfunction and a link to neurodegeneration.

Figure 2. Human TBI transcriptomics align with our 3D in vitro model, revealing conserved mitochondrial dysfunction. The transcriptomic profile of human TBI patients[38] revealed a cell-specific mitochondrial dysfunction signature that converges with our human 3D *in vitro* brain tissue model at 6 hours post-injury, highlighting conserved metabolic vulnerabilities across experimental and clinical contexts (comprised of 2 mln of neurons, 0.5 mln astrocytes, and 0.1 mln of HMC3 microglial cells; n=5 Shams and n=6 CCI samples at 6 hours and n=5 Shams and n=6 CCI samples were used to obtain data for this transcriptomic profile). Single-cell RNA sequencing from 12 critically injured or deceased TBI patients was analyzed to curate cell-specific differentially expressed genes (DEGs) across excitatory neurons, interneurons, astrocytes, and microglia, using data from five uninjured controls as a reference. This patient-derived transcriptomic dataset was then integrated with bulk RNA sequencing from human 3D in vitro brain injury samples at 6 and 48 hours post-injury,

enabling a comparative analysis of conserved molecular signatures between clinical and experimental models. Differentially expressed genes (abs(Log2(Fold Change))>0.585 and q<0.01).

(A) Regardless of convergence or divergence, similar DEGs across the datasets were run with STRING analysis to determine cell-specific pathways significantly altered in response to trauma.

(B) Convergent upregulated and downregulated DEGs were isolated and plotted for 4 distinct cell populations.

Fabrication and validation of fluorescently labeled mitochondria for cell-specific analysis in brain injury.

To dissect the cell-specific mitochondria role in injury-induced neurodegeneration, we inserted genetic information for the expression of three distinct fluorescent tags under COX8A mitochondrial complex IV protein promoter in human brain cell types – human-induced neuronal stem-derived neurons, primary astrocytes, and immortalized HMC3 microglia (Figure 3A-D).

Addgene provided two of the three fluorescent plasmids (EGFP and dsRED2) under the COX8A promoter (Figure 3B). We used VectorBuilder to implement a BFP2 fluorescent protein into the published backbone of mitochondria-targeted fluorescent plasmid under the COX8A promoter (Figure 3B) [63]. We selected COX8A for all cell types due to its continuous presence and abundance in the mitochondria.

We first performed fluorescent intensity vs. viability experiments through various multiplicities of infection (MOI) (Figure 3C). Cell lines with the best viability and signal were selected. MOI40 was selected for mtdsRED2-iNSCs, MOI40 was selected for mtEGFP-astrocytes, and MOI10 was selected from mtBFP2-microglia. After transduction and expansion, we selected cells positive with target fluorophore using the Fluorescent-Activated Cell Sorter (FACS) (Figure 3D) and finally expanded cell lines for storage and experimentation. Please see the detailed description of the transduction protocol in the methods section.

Figure 3. Design and validation of fluorescently labeled mitochondria.

(A) Schematic representation of mitochondria transduction experiments.

(B) dsRED2 Addgene plasmid used to transduce human-induced neural stem cells, EGFP Addgene plasmid used to transduce human primary astrocytes, and custom-designed BFP2 plasmid used to transduce HMC3 microglia cell line.

(C) Epifluorescent images of experimental multiplicities of infection (MOIs) in transduced fluorescent cell lines cultured in 2D, captured using a Cytation epifluorescent microscope.

(D) Fluorescent-activated cell sorter gating parameters for transduced cells with final MOIs.

Figure 4. Molecular characterization and validation of the lentiviral transduction of the brain cells.
(A) Confocal images of 2D monocultures of mtdsRED2-Neurons, mtEGFP-Astrocytes, and mtBFP2-Microglia counterstained with MTCO1 (mitochondrially encoded cytochrome C oxidase I). Scale bar: 10 mm.

(B) Mitochondria-specific markers analysis using Western blots of mitochondria isolated from 2D monocultures and NAM tricultures. pDRP1/DRP1 (Dynamin-related protein 1 phosphorylated at Ser 616 over total DRP1 protein) as a marker of mitochondria fission; MTCO1 as a marker of OXPHOS; VDAC1 (voltage-dependent anion channel 1) as a marker of glycolysis, and TOMM20 (translocase of outer mitochondria membrane 20) as a generic mitochondria marker. Data presented as mean±SEM of at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups in each independent experiment. * indicates a significant difference with p<0.05. One-way ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. Data were normalized to sham at each time point.

(C) Confocal images of novel fluorescent cell lines for 5 weeks in 3D silk scaffolds, stained with a neurons (Tuj1- β3 tubulin), astrocyte (ALDH1A1 - Aldehyde Dehydrogenase 1 Family Member A1), and microglia-specific markers (TMEM119 – transmembrane protein 119) to validate cell-specific staining. Scale bar: 20 mm.

Next, to validate the efficiency and specificity of the transduction, we co-stained 2D monocultures of the brain cells with mitochondria-specific marker Mitochondrially Encoded Cytochrome C Oxidase I (MTCO1) to ensure the fluorescent tags were mitochondria-specific. (Figure 4A). The images from panel A demonstrate precise co-localization of the mitochondria fluorescent tag with MTCO1 staining. Next, we isolated mitochondria from 2D monoculture dishes (using an Invitrogen mitochondria isolation kit from tissues) and human 3D *in vitro* brain models to validate mitochondria-specific protein markers via Western Blot analysis. (Figure 4B) displays cell-specific western blot stains of Phosphorylated Dynamin 1 Like / Dynamin 1 Like (pDRP1/DRP1), Mitochondrially Encoded Cytochrome C Oxidase I (MTCO1), Voltage Dependent Anion Channel 1 (VDAC), and Translocase Of Outer Mitochondrial Membrane 20 (TOMM20). The results demonstrate high OXPHOS activity in neuronal monocultures and tricultures (MTCO1 expression) and high glycolytic activity and mitochondrial fission in astrocytes (VDAC1 expression and pDRP1/DRP1). Next, we stained tricultures with cell-specific cytoplasmic marker Tubulin Beta 3 Class III (Tuj1) for neurons, Aldehyde Dehydrogenase 1 Family Member A1 (ALDH1A1) for astrocytes, and Transmembrane Protein 119 (TMEM119) for microglia to ensure the correct fluorescent colors were expressed in the corresponding cell type (Figure 4C). The fluorescent images demonstrated that integrated fluorescent tags were expressed in specific cell types.

Cell-specific mitochondria fragmentation reveals neuronal vulnerability post-injury.

We used our human 3D in vitro brain model with fluorescently labeled mitochondria to assess mitochondrial health and fission dynamics following injury over time (Figure 5A). We used a controlled cortical impact (CCI) machine to inflict a moderate injury (6 m/s with 3 mm tip and 0.6 mm penetration) on our human 3D in vitro brain models to observe cell-specific mitochondria alterations at 5-time points - 24h, 72h, 1wk, 4wk, and 8wk. Western blot analysis of bulk mitochondrial-specific markers at acute and chronic post-injury time points revealed that TOMM20 expression remained largely unchanged, except for a transient increase at 1-week post-injury. Similarly, MTCO1 expression remained stable throughout the study. However, VDAC1 exhibited a marked glycolytic depression at 24 hours, followed by a significant upregulation at 72 hours, aligning with previous reports of early metabolic suppression and compensatory glycolytic activation in TBI [64-66]. VDAC1 levels then remained stable until a significant increase at 8 weeks, suggesting a late-stage metabolic shift.

To further assess mitochondrial structural dynamics, we evaluated mitochondrial fragmentation across neuronal, astrocytic, and microglial populations in our 3D in vitro model using confocal imaging of dsRED2-labeled neurons, EGFP-labeled astrocytes, and BFP2-labeled microglia (Figure

5B). Mitochondrial aspect ratio analysis (Figure 5C) revealed that all cell types exhibited substantial mitochondrial fragmentation at 24 hours post-injury. However, the recovery patterns varied across cell types. Neuronal mitochondria remained highly fragmented for up to 8 weeks, with only partial recovery observed between 72 hours and 8 weeks. Astrocytic mitochondria exhibited progressive fragmentation up to 4 weeks post-injury, followed by complete recovery at 8 weeks, indicating a delayed but reversible structural response. In contrast, microglial mitochondria demonstrated only transient fragmentation at 24 and 72 hours, with full recovery by 1 week, suggesting a more rapid adaptation to injury. These findings highlight neuronal mitochondria as the most vulnerable to persistent structural deficits.

Figure 5. Acute and Chronic Cell-specific mitochondria fragmentation in response to moderate injury.

A human 3D in vitro model composed of mtDsRED2-Neurons, mtEGFP-Astrocytes, and mtBFP2-HMC3 microglia matured in 3D for 6 weeks prior to being exposed to controlled cortical impact injury of moderate severity (6 m/s speed, 0.6 mm penetration, 3 mm tip). At 24 hours, 72 hours, 1 week, 4 weeks and 8 weeks

(A) Mitochondria-specific markers were assessed using Western blots: TOMM20 (translocase of outer mitochondria membrane 20) was used as a generic mitochondria marker; MTCO1 as a marker of OXPHOS; VDAC1 (voltage-dependent anion channel 1) as a marker of glycolysis. Data presented as mean±SEM of at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups. * indicates a significant difference with $p<0.05$. One-way ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. Data were normalized to sham at each time point.

(B) Representative confocal images of mitochondria fluorescent 3D in vitro cultures. Scale bar: 20 mm.

(C) Frequency distribution (%) of cell-specific mitochondria fluorescent signal over time. Data for this plot were collected from at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups in each independent experiment. * indicates a significant difference with p<0.05. One-way ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. Data were normalized to sham at each time point.

Cell-specific mitochondrial bioenergetics reveal divergent metabolic responses to injury.

Mitochondrial fragmentation analysis provides critical insights into structural changes associated with bioenergetic function; however, it does not directly assess whether cells generate sufficient ATP to meet metabolic demands. To address this, we isolated mitochondria from scaffolds at five-time points—24h, 72h, 1w, 4w, and 8w—followed by Fluorescence-Activated Cell Sorting (FACS) to separate mitochondria based on cell type-specific fluorescence. Freshly sorted mitochondria were then assessed for bioenergetic function using the Seahorse ATP Rate Assay to measure oxygen consumption rate (OCR) and extracellular acidification rate (ECAR), followed by a Ray Biotech Metabolite Assay to analyze metabolic alterations (Figure 6A-B).

Neuronal mitochondria exhibited a significant drop in OCR by 72 hours, followed by partial recovery at 4 weeks before declining again at 8 weeks, indicating prolonged bioenergetic instability (Figure 6B). Astrocytic mitochondria showed an initial increase in OCR acutely post-injury, followed by a steady decline until 4 weeks, after which levels stabilized. In contrast, microglial mitochondria demonstrated a slight acute increase in OCR, remained stable until 1 week, and then exhibited a significant reduction by 8 weeks. ECAR measurements at 24 and 72 hours were below detection limits across all cell types (Figure 6D). Astrocytic mitochondria displayed a transient increase in ECAR at 1 week, which significantly declined at 8 weeks, whereas microglial mitochondria maintained stable proton pumping capacity until a marked increase at 8 weeks. These findings reveal cell-specific metabolic vulnerabilities, with neuronal mitochondria experiencing persistent energy deficits, astrocytes exhibiting delayed but progressive dysfunction, and microglia undergoing late-stage metabolic reprogramming. At 20 minutes, we added Oligomycin, we added Rotenone + Antimycin A at 40 minutes. P value: * = .05 | ** = .01 | *** = .001 |

Figure 6. Cell-specific mitochondrial bioenergetics reveal divergent metabolic responses to injury.

(A) Schematic representation of mitochondria sorting experiments followed by measurement of mitochondria bioenergetic function

(B) The total number of cell-specific mitochondria sorted at each time point after injury. Data presented as mean±SEM of at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups in each independent experiment. * indicates a significant difference with p<0.05. One-way ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. The ROUT outlier analysis method was used to exclude statistical outliers. Data were normalized to sham at each time point.

(C) Oxygen Consumption Rate (OCR) and

(D) Extracellular Acidification Rates were measured with Real-time ATP Rate Assay on freshly sorted mitochondria 24, 72 hours, 1, 4 and 8 weeks post-injury. Data presented as mean±SEM of at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups. *, **, *** indicates a significant difference with p<0.05, p<0.01, p<0.001 respectively. Repetitive measures ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. The ROUT outlier analysis method was used to exclude statistical outliers. Presented OCR and ECAR rates were normalized to mitochondria DNA content.

Divergent mitochondria metabolic profiles in brain cells following injury

Despite decades-long assumptions that mitochondria primarily function as cellular powerhouses, emerging evidence highlights their equally critical role in metabolic regulation, transcriptional control, and intracellular signaling, all of which are essential in the cellular response to injury and the progression of neurodegeneration. Thus, we next evaluated cell-specific metabolic alterations.

Figures 7A–C illustrate cell-type-specific metabolic alterations in neuronal, astrocytic, and microglial mitochondria at 24 hours and 8 weeks post-injury, as assessed by the Ray Biotech Metabolite Assay, with metabolite interaction networks analyzed via StringDB functional enrichment. These findings reveal distinct mitochondrial metabolic adaptations in response to injury, highlighting divergent mechanisms of recovery, dysfunction, and long-term metabolic reprogramming across brain cell types.

Neuronal mitochondria exhibit acute metabolic dysfunction at 24 hours post-injury, characterized by substantial alterations in pathways linked to cellular recovery, including nicotinate and nicotinamide, pyrimidine, purine, and riboflavin metabolism. By 8 weeks, these pathways return to baseline, suggesting metabolic resilience and recovery. However, early disruptions in sphingolipid metabolism and glycans degradation, key processes in post-translational protein modification, suggest transient impairments in protein homeostasis. Critical regulatory hubs identified in Figure 6C, including SIRT2, SIRT3, NAMPT, and ARSA, further emphasize mitochondrial involvement in coordinating metabolic recovery.

Astrocytic mitochondria show no significant metabolic alterations at 24 hours, but by 8 weeks, they exhibit a profound decline in metabolic activity, particularly in pathways essential for O-glycans, glycosaminoglycans, glycophospholipids, and glycosphingolipid biosynthesis. This mirrors the progressive impairment in astrocyte-mediated metabolic support, potentially affecting neuronal function and long-term homeostasis. Like neuronal mitochondria, astrocytic mitochondria exhibit decreased pantothenate and CoA biosynthesis, crucial for hexosamine biosynthetic pathways regulating protein glycosylation and sialylation. However, unlike neuronal mitochondria, the astrocytic mitochondrial enrichment network is fragmented, reflecting a loss of metabolic coordination that may contribute to sustained dysfunction.

Microglial mitochondria, in contrast, show a rapid metabolic upregulation at 24 hours, with several metabolites remaining elevated at 8 weeks, including TYMP, VISFATIN, SIRT1, and SIRT2. These findings indicate that microglial mitochondria sustain an altered metabolic state, likely linked to prolonged inflammatory activation. Similar to neuronal and astrocytic mitochondria, microglial mitochondria exhibit alterations in sphingolipid, glycosphingolipid, and glycophospholipid metabolism, which are critical for membrane homeostasis and post-translational modifications. The enrichment network reveals two distinct microglial mitochondrial clusters—one highly interconnected network centered around SIRT1, SIRT2, CD38, and BST1, and a second, more fragmented network with multiple disconnected metabolic markers. This suggests that microglial

mitochondrial metabolism remains in a prolonged state of dysregulation, likely contributing to chronic neuroinflammation and neurodegenerative processes following injury.

Figure 7. Divergent mitochondria metabolic profiles in brain cells following injury. Mitochondria that was stored after the bioenergetic assessment was used to determine metabolic alterations at acute (24 hours) and chronic (8 weeks) time points using the RayBiotech Metabolic panel.

(A) Differentially produced metabolites Mitochondria from 3 independent experiments were assessed, with n=3 for sham groups and n=4 scaffolds for injury groups in each independent experiment. Data presented as mean±SEM of at least three independent experiments. * indicates a significant difference with p<0.05. Two-way ANOVA with Tukey's post-hoc test determined the difference between the control and experimental groups. The ROUT outlier analysis method was used to exclude statistical outliers. Data were normalized to sham at each time point.

(B) KEGG Pathway enrichment analysis isolated in panel A metabolites that were differentially produced.

(C) Enrichment network corresponding to cell-specific metabolites from Panel A.

Temporal Progression of Mitochondrial Dysfunction and Alzheimer's-Like Pathology Following Injury

Figure 8. Mitochondria dysfunction in association with injury-induced AD-like neurodegenerative phenotype.

(A) Displays representative confocal images of the 3D triculture scaffold of Tuj1 neuronal membrane stain over time for the sham and CCI groups.

(B) Neurodegenerative markers were assessed by Western blot at acute (24 and 72 hours) and chronic (1, 4, and 8 weeks) post-injury time points. Phosphorylated Tau (AT8 at Ser202 and T205) presented over total Tau; TAR DNA Binding Protein (TDP43), amyloid precursor protein (APP), Gap Junction Protein Alpha 1 (Connexin43), Annexin A2 (ANXA2) as well as ELISA of Amyloid Beta 42 presented over Amyloid-Beta 40. Data presented as mean±SEM of at least three independent experiments with n=3 for sham groups and n=4 scaffolds for injury groups in each independent experiment. *, **, *** indicates a significant difference with p<0.05, p<0.01, p<0.001 respectively. One-way ANOVA with Tukey's post-hoc test was used to determine the difference between control and experimental groups. Data were normalized to sham at each time point.

The final question we investigated was whether a temporal link exists between injury-induced mitochondrial dysfunction and the progression of Alzheimer 's-like pathology. To address this, we assessed neuronal network integrity (Figure 8A) alongside well-established neurodegenerative markers associated with Alzheimer's disease (Figure 8B). The results indicate that key hallmarks of Alzheimer's pathology, including pTau (AT8), TDP-43, APP, connexin 43, annexin A2, and the Aβ42/40 ratio, exhibit injury-induced alterations at acute time points, with all markers beginning to increase by 72 hours post-injury.

Phospho-Tau (AT8) shows an initial increase at 72 hours, returns to baseline at 4 weeks, and becomes upregulated again at 8 weeks, indicating the presence of a chronic neurodegenerative process despite apparent neuronal network recovery. TDP-43 follows a similar pattern, with upregulation observed at 72 hours post-injury. APP, a key precursor of Aβ, increases significantly from 72 hours to 1 week before returning to baseline by 4 weeks. Connexin 43 and annexin A2, both of which are linked to Alzheimer's disease, exhibit a similar temporal pattern, with upregulation at 72 hours, 1 week, and 8 weeks, followed by a transient decline at 4 weeks. Lastly, the Aβ42/40 ratio, a hallmark of Alzheimer's pathology, shows a significant drop at 24 and 72 hours, recovers to baseline by 1 week, and then increases significantly at 4 and 8 weeks, suggesting a delayed shift toward amyloidogenic processing. These findings establish a clear timeline in which mitochondrial dysfunction precedes and potentially drives progressive neurodegenerative changes, reinforcing the role of energy failure as a key contributor to injury-induced Alzheimer 's-like pathology.

Discussion

Epidemiological studies demonstrated a strong link between traumatic brain injury (TBI) and an increased risk of neurodegenerative disorders, particularly Alzheimer's disease (AD) [5-8]. TBI ranks as the third most significant risk factor for AD, following aging and APOE genotype [8]. The likelihood of developing AD correlates with injury severity, with severe TBI increasing the risk by more than fourfold [8]. Clinicopathological studies have identified hallmark AD pathologies, such as amyloid-beta (Aβ) plaques and neurofibrillary tangles composed of hyperphosphorylated Tau, in the postmortem brains of TBI patients [8,9]. These findings suggest that a single TBI event may predispose individuals to AD, yet the underlying mechanisms remain unclear. Metabolic and bioenergetic dysfunction [30,31,67,68], protein aggregation [68-70], and chronic inflammation [71-74] have emerged as potential triggers of disease progression, but their precise role in driving neurodegenerative changes after TBI requires further investigation.

Mitochondrial dysfunction is one of the earliest and most critical regulators of injury-induced neurodegeneration, which disrupts brain energy homeostasis and cellular maintenance pathways [6,10,75-77]. Mitochondria generate ATP through oxidative phosphorylation, supporting essential neuronal functions such as synaptic activity, ion balance, and cytoskeletal dynamics [11,78]. TBI-induced mitochondrial damage leads to impaired ATP production, Ca^{2+} dysregulation, oxidative stress, and activation of cell death pathways [10,79-85]. These metabolic disturbances may set the stage for pathological protein aggregation, including Tau and Aβ misfolding, further exacerbating neurodegeneration [84,86-91]. Additionally, mitochondrial dysfunction affects lipid metabolism and glycosylation pathways, which are crucial for cellular recovery and signaling and were both implicated in AD patients [92,93]. Recent studies have established that mitochondria across different cell types possess distinct protein profiles [94]. However, whether these unique profiles translate to differential responses to brain trauma and their role in driving injury-induced neurodegeneration remains unexplored. Thus, we raised the question of whether the mitochondria of different brain cells exhibit distinct responses to brain trauma and how these differential responses shape the progression of injury-induced neurodegeneration.

Previous 24-hour transcriptomic analysis [34] showed extra and intracellular mitochondria dysfunction; here, we further corroborated these results with extended bulk transcriptomic analysis at 6- and 48-hours post-injury, indicating drastic changes to OXPHOS and mitochondria maintenance pathways (Figure 1). While previous studies, including our own, have established overall mitochondrial dysfunction, none have delineated cell-specific mitochondrial alterations. One study suggested acute, brain cell-specific mitochondrial dysfunction post-TBI using non-invasive metabolic imaging and secreted miRNA [95]; however, no prior human cell or patient-based brain injury study [15,25,96] has comprehensively characterized the cell-specific bioenergetic and metabolic dysfunction of mitochondria across both acute and chronic time points or directly linked these alterations to injury-induced neurodegeneration.

Our results demonstrated that following brain injury, mitochondrial gene expression undergoes a dynamic response indicative of acute stress, compensation, and recovery. At 6 hours post-injury, the downregulation of all mitochondrial complexes except Complex V, alongside selective upregulation of MT-ND genes in Complex I, suggests an early compensatory mechanism to sustain minimal ATP production while mitigating oxidative stress [97-99]. By 24 hours, the drastic increase in Complex I gene expression and moderate upregulation of other complexes indicate a metabolic shift, likely driven by PGC-1α-mediated mitochondrial biogenesis and neuroinflammatory activation, particularly in microglia. This phase reflects an attempt to restore oxidative phosphorylation as cells transition from an acute to a subacute injury response. By 48 hours, the lack of altered gene expression suggests mitochondrial recovery or clearance of damaged mitochondria via mitophagy, marking the return to metabolic homeostasis. However, long-term bioenergetic assessments challenge this assumption of recovery. Despite the initial transcriptomic normalization, by 72 hours, mitochondrial function declined across all cell types, with progressive worsening at chronic time points. This persistent bioenergetic dysfunction, in contrast to the early transcriptional recovery, suggests that mitochondrial impairment is not transient but instead becomes a sustained pathological feature, potentially contributing to long-term neurodegeneration rather than returning to a true homeostatic state.

Our further cell-specific transcriptomic analysis between human TBI patients and our 3D in vitro brain injury model reveals a striking convergence in cell-specific mitochondrial dysfunction. This supports the hypothesis that metabolic disturbances are central to injury-induced neurodegeneration. Showing that while neurons primarily attempt to restore synaptic function post-injury, they also exhibit dysregulation of MAPK signaling, a key regulator of mitochondrial activity, oxidative stress, and apoptosis. In contrast, astrocytes and microglia demonstrate profound metabolic shifts, with early disruptions in OXPHOS and ATP production, followed by inflammatory activation and alterations in lipid metabolism. In addition to mitochondria dysfunction markers, we identified multiple genes linked to Alzheimer's disease (AD) across different brain cell types, reinforcing the potential role of mitochondrial impairment in driving injury-related neurodegeneration. Microglia exhibited alterations in STIP1, DTL, and CTSB, genes associated with AD [53-55], alongside changes in epigenetic regulators [56-58], potentially influencing long-term neuroinflammatory responses. Astrocytes displayed metabolic and transcriptional dysregulation, while excitatory neurons showed disruptions in CAMK2D and MAOB, affecting synaptic plasticity and oxidative stress, both hallmarks of AD [46,47,49,100,101]. The convergence of mitochondrial dysfunction, metabolic stress, and neurodegenerative signatures across human patients and our in vitro model underscores the importance of mitochondrial health in determining long-term injury outcomes. These findings suggest that mitochondrial impairment may serve as an early driver of injury-induced neurodegeneration, linking TBI to the onset of AD-like pathology. Additionally, this analysis enforced the physiological relevance of our in vitro model.

Armed with the initial analysis between human data and our bulk transcriptomics, we hypothesized that neuronal mitochondria undergo devastating damage and require immediate assistance by glial cells to restore their function. To address this fundamental question, we have developed a one-of-a-kind model of brain injury that allows for cell-specific live mitochondria research (Figure 3-8).

Our integrated analysis of cell-specific transcriptomics, mitochondrial fragmentation, bioenergetics, and metabolic shifts provides a clear mechanistic link between injury-induced mitochondrial dysfunction and the emergence of neurodegenerative markers. Confocal imaging of mitochondrial health showed that while outer membrane integrity (TOMM20) remained stable, VDAC1 expression exhibited an acute drop at 24 hours, aligning with known glycolytic suppression in TBI patients [64-66,102]. Mitochondria bioenergetic assessments further supported this, where neuronal OCR dropped substantially by 72 hours, marking the first significant decline in mitochondrial respiration. This coincided with a sharp increase in neurodegenerative markers, including pTau, APP, and TDP-43, suggesting that mitochondrial dysfunction may be directly contributing to early neurodegenerative processes. The sustained neuronal mitochondrial fragmentation for up to 8 weeks, alongside fluctuations in OCR recovery and subsequent decline, indicates that neurons fail to fully restore metabolic homeostasis, making them highly susceptible to long-term degeneration.

Astrocytic and microglial mitochondria displayed distinct metabolic trajectories, with astrocytes showing minimal acute metabolic disruptions but developing chronic impairments in glycan biosynthesis and lipid metabolism by 8 weeks, as seen in our metabolic panels. Conversely, microglial mitochondria initially upregulated metabolic pathways, including markers such as SIRT1 and SIRT2, but displayed persistent inflammation-associated metabolic activity at chronic time points. Notably, at 72 hours, microglial OCR remained stable while astrocytes showed an increasing trend, contrasting with neuronal OCR decline, suggesting that while glial cells actively respond to metabolic demands, neurons experience an early energy crisis that coincides with neurodegenerative marker upregulation.

The Aβ42/40 ratio, a key hallmark of AD pathology [101,103,104], initially dropped at 24 hours, remained suppressed for up to 1 week, and then significantly increased at chronic time points, mirroring mitochondrial metabolic instability. This pattern is highly relevant to Alzheimer's disease (AD) pathology, where disruptions in Aβ metabolism contribute to synaptic toxicity and neuronal dysfunction. The initial decrease in Aβ42/40 levels may indicate an acute impairment in amyloid processing or clearance, a phenomenon observed in early TBI pathology. However, the subsequent increase at chronic time points parallels AD-like progression, where a sustained metabolic dysfunction leads to excessive Aβ accumulation and aggregation. This shift, combined with sustained pTau upregulation and persistent neuronal OCR instability, suggests that mitochondrial dysfunction after injury may create a permissive environment for AD-like neurodegeneration, potentially accelerating disease onset in predisposed individuals. These findings reinforce the growing body of evidence that TBI serves as a trigger for AD pathology, with mitochondrial failure playing a central role in the transition from acute injury to chronic neurodegeneration.

In summary, our study integrated cell-specific mitochondrial profiling with patient-derived transcriptomic data, providing unprecedented insight into how mitochondrial dysfunction progresses following TBI, and provided the first comprehensive evidence that mitochondrial dysfunction is not a uniform event but instead manifests in a cell-type-specific manner, with neuronal mitochondria exhibiting the most prolonged metabolic instability and direct correlation with neurodegenerative marker upregulation. Our cell-specific mitochondrial analyses aligned with transcriptomic findings, reinforcing that neurons struggle with energy production, astrocytes fail to maintain metabolic support long-term, and microglia sustain chronic inflammatory activity. This confirms that neuronal bioenergetic failure is an early driver of neurodegeneration, while astrocytes

and microglia contribute to disease progression over time. Our approach proved helpful in identifying mitochondrial dysfunction as a therapeutic target, emphasizing the need for interventions that stabilize neuronal metabolism early after injury to prevent long-term neurodegeneration.

Limitations to the study
While our study provides unprecedented insights into cell-specific mitochondrial dysfunction and its role in injury-induced neurodegeneration, several limitations should be considered. First, our model lacks a vascular component and neuronal myelination, limiting our ability to assess how blood-brain barrier integrity and white matter injury contribute to metabolic dysfunction. The absence of these key elements may underestimate the impact of vascular and myelination deficits on long-term neurodegeneration. Second, our study utilized a single genetic background, restricting our ability to account for individual variability in mitochondrial resilience, immune responses, and neurodegenerative susceptibility. Future studies incorporating multiple donor-derived cell lines will be necessary to determine whether genetic differences influence mitochondrial responses to injury. Lastly, our study is limited to an 8-week post-injury period, providing insight into early and intermediate stages of injury progression but not capturing long-term neurodegenerative changes beyond this window. Extending the study duration would allow us to determine whether mitochondrial dysfunction persists or worsens over time, particularly in relation to Alzheimer's-like pathology. Addressing these limitations in future research will enhance our understanding of how mitochondrial dysfunction contributes to chronic neurodegeneration following TBI.

Authors contribution
VL conceived the project. SBSK and VL designed and interpreted the experiments and wrote the manuscript. SBSK and SPS performed scaffold seeding and sample maintenance (media changes). SBSK performed viral transductions, all in vitro experiments, injuries, and Raybiotech assays. SBSK and VL performed mitochondria isolation and FACS sorting. SPS and SBSK performed mitochondria bioenergetic function assessment. SPS, VL, and SBSK performed confocal imaging. SBSK, SPS, CA, and MT conducted Western Blots. VL performed ELISA. CA performed an aspect ratio analysis. SBSK and SPS prepared silk materials. SBSK performed RNA sequencing analysis.

Acknowledgments
The authors thank the University of Cincinnati Start-Up Funds and Research Scholar Award, NIH (1R21AG085052-01A1), Cincinnati Children's Hospital Bioimaging and Analysis Facility, and Research Flow Cytometry Facility.

Methods:

Fabrication of 3D Silk Scaffolds

3D silk scaffolds were fabricated following established protocols [34,105,106]. Bombyx mori cocoons were boiled in 0.02 M sodium carbonate for 30 minutes, then dried overnight under a fume hood. The dried silk fibers were dissolved in 9.3 M lithium bromide at 60 °C for 4 hours. To remove lithium bromide, the silk solution underwent dialysis in deionized water at room temperature for 72 hours using 3,500 molecular weight cut-off dialysis tubing (Invitrogen), with water changes every 12 hours. The dialyzed silk solution was centrifuged at 9,000 rpm for 20 minutes twice, and debris was removed using a 100 µm pore strainer. The concentration was adjusted to 6 mg/mL before casting into 10 cm dishes. To induce porosity, 400–500 µm sodium chloride (Sigma) particles were applied at a 1:2 (v/w) ratio. The solution was left to solidify at room temperature for 48 hours, followed by incubation at 60 °C for 1 hour to form silk sponges. Silk sponges underwent further dialysis in deionized water at room temperature for 48 hours, with six total water changes.

Scaffolds were cut into a 6 mm outer diameter and 2 mm inner diameter using biopsy punches (Integra) and trimmed to a height of 1.5 mm. Prior to cell seeding, scaffolds were autoclaved in deionized water for 20 minutes on a liquid cycle and equilibrated to room temperature. Scaffolds (n = 50 per condition) were placed in 6-well plates and incubated overnight at 37 °C in 7 mL of 10 µg/mL poly-L-ornithine (PLO) solution. Following three washes in distilled water at 5-minute intervals, scaffolds were incubated at 4 °C overnight in 0.5 mg/mL laminin (ThermoFisher 50-100-3381) prepared in phenol red-free Dulbecco's Modified Eagle Medium/Nutrient Mixture (DMEM/F12, Invitrogen). Before use, scaffolds were equilibrated at 37 °C.

Mycoplasma testing

All cell lines used in this project were either checked for mycoplasma before starting or were directly purchased from established vendors, that provided certification for Mycoplasma testing.

Mouse embryonic fibroblasts (MEFs)

Mouse embryonic fibroblasts (MEFs) were cultured as a feeder layer for human-induced neuronal stem cells (hiNSCs) following ATCC guidelines. MEFs at passages 1–4 were maintained in Dulbecco's Modified Eagle Medium (DMEM) (1X) + GlutaMAX (ThermoFisher, 10569010) supplemented with 10% fetal bovine serum (FBS) (Gibco, 10-438-026) and 1% antibiotic-antimycotic (ThermoFisher, 15240062). Cells were incubated at 37 °C with 5% CO_2, and media was changed every 3–4 days.

For seeding, 15 cm culture dishes were coated with 0.1% gelatin for 1 hour at room temperature before plating MEFs in 20 mL of MEF media. For expansion, cells were washed with phosphate-buffered saline (PBS), detached using 0.25% Trypsin-EDTA for 3 minutes at 37 °C, neutralized with MEF media, and centrifuged at 1,000 rpm for 5 minutes. Cells were split at a 1:3 ratio and passaged at 70–80% confluency.

To prepare MEFs as a feeder layer, cells were grown to 90–100% confluency and inactivated by incubation with 10 µg/mL Mitomycin C (Sigma, M4287-5X2MG) in MEF media at 37 °C for 3 hours. Following inactivation, cells were washed three times with PBS. If hiNSCs were seeded immediately, 20 mL of hiNSC media was added. If not used immediately, inactivated MEFs were maintained in 20 mL of MEF media with media changes every 3–4 days for up to 7 days.

Human induced neural stem cells (iNSCs) and mtRFP-iNSCs

Induced neural stem cells (iNSCs) were generated from dermis-derived human fibroblasts, originally derived from neonatal foreskin fibroblasts. Cells were maintained in KnockOut DMEM (ThermoFisher, 10829018) supplemented with 1% GlutaMAX (ThermoFisher, 35050061), 1% antibiotic-antimycotic

(ThermoFisher, 15240062), 0.2% 2-Mercaptoethanol (ThermoFisher, 21985023), and 20% KnockOut Serum Replacement (KNSR) (ThermoFisher, 12-618-013). Cultures were incubated at 37°C with 5% CO_2, and media was refreshed every 1–2 days.

For expansion, complete iNSC media was prepared by supplementing 50 mL of iNSC media with 800 μL of 10 μg/mL basic fibroblast growth factor (bFGF) (ThermoFisher, PHG0261). Prior to seeding, MEF feeder plates were cultured until 90–100% confluency and inactivated with Mitomycin C. iNSCs were plated on inactivated MEF feeders in 20 mL of iNSC media.

For passaging, iNSCs were maintained until reaching 70–80% confluency. Cells were washed with PBS, detached using TrypLE at 37°C for 1 minute, neutralized with iNSC media, and centrifuged at 3,000 rpm for 2 minutes. The cell pellet was resuspended by gentle pipetting (5 times) and split at a 1:20 ratio for further expansion. The same protocol was used for culturing fluorescently labeled iNSC lines.

Human primary astrocytes and mtGFP-Astrocytes

Primary astrocytes were cultured in astrocyte growth media containing fetal bovine serum (FBS), astrocyte growth supplement, and Penicillin-Streptomycin (Pen/Strep) solution (ScienCell Research Laboratories, NC9707094). Cells were maintained at 37°C with 5% CO_2, and media was refreshed every 3–4 days.

For seeding, 15 cm culture dishes were coated with either 2.5 μL Poly-L-Lysine (Sigma, P4832) in 15 mL sterile distilled water for 1 hour (normal astrocytes) or 25 μL Poly-L-Lysine, 150 μL Laminin (ThermoFisher, 50-100-3381), and 15 mL sterile distilled water for 1 hour (mtGFP-labeled astrocytes). After coating, astrocytes were plated in 20 mL of astrocyte media.

For expansion, cells were washed with PBS, detached using 0.25% Trypsin-EDTA for 3 minutes at 37°C, neutralized with astrocyte media, and centrifuged at 1,000 rpm for 5 minutes. Cells were resuspended and re-seeded at a density of 350,000 cells per 15 cm dish. Passaging was performed at 70–80% confluency.

HMC3 microglia and mtBFP-Microglia

HMC3 human microglial cells were cultured in Eagle's Minimum Essential Medium (EMEM) (ThermoFisher, 50-188-268FP) supplemented with 10% fetal bovine serum (FBS) (Gibco, 10-438-026) and 1% antibiotic-antimycotic (ThermoFisher, 15240062). Cultures were maintained at 37°C with 5% CO_2, and media was changed every 3–4 days.

For seeding, HMC3 cells were plated in 20 mL of HMC3 media. For expansion, cells were washed with PBS, detached using 0.25% Trypsin-EDTA for 3 minutes at 37°C, neutralized with HMC3 media, and centrifuged at 1,000 rpm for 5 minutes. Cells were resuspended and re-seeded at a density of 350,000 cells per 15 cm dish. Passaging was performed at 70–80% confluency. The same protocol was used for subculturing fluorescently labeled cell lines.

Lentivirus Transduction Protocol

Seeding Density:
In order to quantify optimal confluency and cell growth of each cell type, seeding cells at different densities is required.

hiNSC: Coat plates with 250 microL of Matrigel diluted in 24 mL of DMEM F12. Reconstitute, add 2 mL per plate, and wait 1 hour. Prepare hiNSC cells from cryovial or dishes. Count cells. Centrifuge cells for 2 min at 3000 rpm. Remove supernatant and add hiNSC media. Resuspend cell pellet 5 times. Add number of cells based on map and add sufficient media of each well to 2 mL. Wait 24 hours, image wells, and choose optimal seeding density.

Astrocyte: Coat plates with 2.5 microL of Poly-L-Lysine diluted in 12 mL of sterile UltraPure water. Add 2 mL per well and wait 1 hour. Prepare astrocyte cells from cryovial or dishes. Count cells. Centrifuge cells for 5 min at 1000 rpm. Remove supernatant and add astrocyte media. Resuspend cell pellet 5 times. Add number of cells based on map and add sufficient media of each well to 2 mL. Wait 24 hours, image wells, and choose optimal seeding density.

HMC3: Prepare HMC3 cells from cryovial or dishes. Count cells. Centrifuge cells for 5 min at 1000 rpm. Remove supernatant and add HMC3 media. Resuspend cell pellet 5 times. Add number of cells based on map and add sufficient media of each well to 2 mL. Wait 24 hours, image wells, and choose optimal seeding density.

Lentiviral Transduction:
In order to quantify optimal transduction efficiency and cell growth of each cell type, transducing cells at different MOIs is required. Abided to BSL 2 Protocol link protocol

Lentivirus Transduction

hiNSC: The hSYN-mito-dsRED2 plasmid (Addgene, #173069) was used to generate dsRED2-expressing iNSCs via lentiviral transduction.

Day 1: Corning 6-well plates were coated with Matrigel by adding 20 μL of Matrigel diluted in 2 mL of DMEM/F12 per well. After 1 hour at room temperature, the Matrigel solution was aspirated, and iNSCs were seeded at a density of 750,000 cells per well. Sufficient hiNSC media was added to achieve a final well volume of 2 mL. Plates were incubated overnight at 37°C with 5% CO_2. *Day 2*: Lentiviral cryovials were transferred from −80°C to an ice bucket and thawed completely. hiNSC media was aspirated from each well, and lentiviral particles were added according to the viral titer and the desired multiplicity of infection (MOI). A fresh micropipette tip was used for each well. Media was adjusted to maintain a final volume of 2 mL per well. Control wells received 2 mL of hiNSC media without lentivirus. Plates were incubated overnight at 37°C with 5% CO_2. All equipment that came into contact with lentiviral particles was treated with bleach for 30 minutes before disposal. *Day 3*: After 24 hours, media was aspirated from each well using a new pipette tip for each well and disposed of in bleach. Wells were washed with 2 mL of PBS, and the wash solution was removed. Fresh iNSC media (2 mL) was added to each well, and plates were returned to the incubator at 37°C with 5% CO_2. Media changes were performed every 3–4 days.

Astrocyte: The pLV-mitoGFP plasmid (Plasmid #44385) was used to generate GFP-expressing astrocytes via lentiviral transduction.

Day 1: Corning 6-well plates were coated with Poly-L-Lysine (.333 microL of Poly-L-Lysine and 2 mL of UltraPure sterile water). After 1 hour at room temperature, Poly-L-Lysine solution was aspirated, and astrocytes were seeded at a density of 300,000 cells per well. Sufficient astrocyte media was added to achieve a final well volume of 2 mL. Plates were incubated overnight at 37°C with 5% CO_2. *Day 2*: The same procedure was conducted as Day 2 for iNSC transduction, but used EGFP lentivirus and astrocyte media instead of DsRed2 lentivirus and iNSC media. *Day 3*: The same procedure was conducted as Day 2 for iNSC transduction, but used astrocyte media instead of iNSC media.

HMC3: The custom plasmid used VectorBuilder to implement a double COX8A promoter and EBFP2 into the plasmid backbone published by Isamu Taiko and his group [14].

Day 1: Corning 6-well plates were not coated. HMC3 were seeded at a density of 300,000 cells per well. Sufficient astrocyte media was added to achieve a final well volume of 2 mL. Plates were incubated overnight at 37°C with 5% CO_2. *Day 2*: The same procedure was conducted as Day 2 for iNSC transduction, but used EBFP2 lentivirus and HMC3 media instead of DsRed2 lentivirus and

iNSC media. *Day 3*: The same procedure was conducted as Day 2 for iNSC transduction, but used HMC3 media instead of iNSC media.

3D in vitro human triculture brain tissue model fabrication

3D *in vitro* human triculture brain tissue models were produced following the same protocol established in. RFP-expressing induced neural stem cells (RFP-INSCs), GFP-expressing astrocytes (GFP-Astros), and BFP-expressing human microglial clone 3 cells (BFP-HMC3) were cultured to near 100% confluency. Cells were lifted using monoculture methods and combined at a ratio of 2:0.5:0.1 million (RFP-INSCs:GFP-Astros:BFP-HMC3). The cell suspension was centrifuged at 1000 rpm for 5 minutes, and the supernatant was aspirated. Cells were resuspended in neurobasal medium (ThermoFisher 21-103-049) supplemented with 2% B-27 (ThermoFisher 17-504-044), 1% Anti-Anti (ThermoFisher 15240062), 1% GlutaMAX (ThermoFisher 35050061), and 1% astrocyte growth factors (ThermoFisher NC9746822).

Poly-L-ornithine (PLO)- and laminin-coated silk scaffolds were prepared in 96-well plates, and excess liquid was removed using a vacuum line. A 40 μL cell suspension (2:0.5:0.1 million cells) was seeded onto each 3D silk scaffold. Scaffolds were incubated at 37℃ for 30 minutes to allow cell attachment, followed by the addition of 150 μL neurobasal medium per well. Scaffolds were incubated overnight at 37°C with 5% CO2.

The following day, cell-seeded scaffolds were transferred to new 96-well plates. A 100 μL collagen type I solution (ThermoFisher 344702001) was added to each scaffold, and the plates were incubated for 30 minutes at 37°C with 5% CO2. Subsequently, 150 μL neurobasal medium was added to each well, and the scaffolds were incubated overnight at 37°C with 5% CO2.

The next day, brain-like tissues were transferred to 48-well plates, each containing 1 mL neurobasal medium. The medium was changed every 3-4 days, and the plates were incubated at 37°C with 5% CO2. 3D in vitro silk brain models were ready 6 weeks post-seeding.

Mitochondria Isolation

Mitochondria were isolated from 2D cell lines using the Mitochondria Isolation Kit for Cultured Cells (Thermo Fisher Scientific). Human microglial clone 3 cells (HMC3), astrocytes, and induced neural stem cells (hiNSC) were seeded independently in 6-well plates at densities of 350,000, 350,000, and 1,000,000 cells per well, respectively, and cultured for 3 days.

Cells were harvested and mitochondria were isolated according to the manufacturer's Option A protocol. Briefly, cells were centrifuged at 850 × g for 2 minutes. The supernatant was removed, and cells were resuspended in 800 μL Reagent A. The suspension was transferred to a 2.0 mL microcentrifuge tube, vortexed at medium speed for 5 seconds, and incubated on ice for exactly 2 minutes. 10 μL Reagent B was added, and the sample was vortexed at maximum speed for 5 seconds. After a 5-minute incubation on ice, with vortexing at maximum speed every minute, 800 μL Reagent C was added by gentle inversion. The tube was centrifuged at 700 × g for 10 minutes at 4°C with low braking. The supernatant was transferred to a 1.0 mL microcentrifuge tube and centrifuged at 12,000 × g for 15 minutes at 4°C. The supernatant was aspirated, and the mitochondrial pellet was resuspended in 100 μL RIPA buffer. Samples were stored at -80°C until Western blot analysis.

Use tweezers to transfer scaffolds to 2mL eppendorfs. Add 800μL of Mito Iso Reagent A. Squish and open scaffold to remove cells. Vortext samples for 5 seconds on medium speed and incubate on ice for 2 minutes. Add 10μL of Mito Iso Reagent B and vortex on medium speed for 5 seconds. Incubate on ice. Vortex every minute for 5 minutes. Add 800μL of Mito Iso Reagent C to each sample. Invert 10 times (do not vortex). Remove scaffold. Centrifuge eppendorfs at 1000xg for 10 minutes at 4°C. Transfer supernatant to new 2mL eppendorfs. Throw out eppendorfs that only contain pellet (cell debris). Centrifuge supernatant eppendorf at 12,000xg for 25 minutes at 4°C.

Transfer supernatant to new 2mL eppendorf (this is the protein fraction). Resuspend pellet in 300μL of Seahorse Buffer (this is the mitochondria pellet).

Seahorse ATP Rate Assay

Day prior to assay: open the Extracellular Flux Assay Pack and separate the utility plate (96-well plate) and Agilent Seahorse Sensor Cartridge. Fill each well of the utility plate with 200μL of sterile water using a multichannel pipette. Place the pink guard on the utility plate and insert the XF Sensor Cartridge. Incubate the cartridge/utility plate assembly in a non-CO_2 37°C incubator overnight.

Day of assay: remove sterile water from the 96-well utility plate and add 200μL of calibrant to each well. Place the cartridge/utility plate assembly back in a non-CO_2 37°C incubator until ready for the assay. 50mL DMEM F12 Phenol Red-free media, 0.5mL pyruvate, 0.5mL glucose, and 0.5mL glutamine is used for Seahorse media. Add 420 μL Seahorse media to Oligomycin. Take 300 μL of solution, and add 2.7 mL of Seahorse media. Add 540 μL Seahorse media to Rotenone + Antimycin A. Take 300 μL of solution and add 2.7 mL of Seahorse media. When samples are ready, remove calibrant from Seahorse cartridge. add 180 μL of Seahorse media to mitochondria pellet and put into Seahorse cartridge. Go to Seahorse plate reader and open Wave Desktop Software. Designate seahorse cartridge wells to sample groups. Calibrate the cartridge and insert mitochondria plate. Wait 1 hour. Seahorse data is normalized to DNA quantification.

DNA Quantification

After Seahorse ATP Rate Assay, 20 μL of mitochondria sample is used for DNA quantification. 20 μL of Triton-X solution is added to each mitochondria sample, followed by incubation at room temperature (RT) for 10 minutes. Samples are then diluted 1:5 in TE buffer to optimize fluorescence signal. A standard curve is generated using DNA lambda standards (2000–0 ng/mL). Samples and standards are mixed with 100μL of PicoGreen reagent in black 96-well plates and incubated in the dark for 5 minutes at RT. Fluorescence intensity is measured using a microplate reader (Excitation: 485nm, Emission: 538nm). DNA concentration is determined by interpolation from the standard curve and used to normalize mitochondrial bioenergetics data to DNA content (pg DNA/cell).

Immunofluorescence

Transfer scaffold to 48 well plate. Add 300μL of 4% PFA / 4% Sucrose and wait for 1 hour. Aspirate solution. Add 1mL of PBS and do 4 10-minute washes put on shaker (RT). Aspirate PBS and Add 300μL of IF blocking solution and incubate on shaker (RT) for 1 hour. Aspirate IF blocking solution. Add 300μL of primary antibody (1:1000 dilution) and incubate on rocker (4°C) overnight . Aspirate primary antibody. Add 1mL of PBS and put on shaker (RT) for 10 minutes. Repeat 2 more times. Aspirate PBS. Cover plate with aluminum foil (secondary is light sensitive). Add 300μL of secondary antibody (1:500 dilution) and incubate on shaker (RT) at slow speed for 1 hour. Aspirate secondary antibody. Add 1mL of PBS and put on shaker (RT) for 10 minutes. Repeat 2 more times. Aspirate PBS. Add .5mL PBS. Store in 4°C or proceed with imaging.
IF Primary: MTCO1 Ms (Invitrogen REF 459600), Tuj1 (Anti-beta III Tubulin) Ms (Abcam ab78078), ALDH1A1 Rb (Invitrogen REF MA5-42586), TMEM119 Rb (Invitrogen REF PA5-119902). *IF Secondary*: Goat anti-Chicken IgY (H+L) (Invitrogen REF A32933), Goat anti-Mouse IgG (H+L) (Invitrogen REF A32728), Goat anti-Rabbit IgG (H+L) (Invitrogen A32733)

RayBio Human Metabolic Pathways Antibody Array 1:

Remove kit from −20°C and let thaw. Remove Antibody Arrays from plastic packaging. Place each membrane face-up into 6 well incubation tray (ensure proper orientation). Add 2mL of RayBio Blocking Buffer into each well and incubate on shaker (RT) at slow speed for 30 minutes. Aspirate RayBio Blocking Buffer.

Pipette 1 mL of sample into each well and incubate on shaker (RT) at slow speed overnight. Aspirate sample from each well. Add 2mL of 1x Wash Buffer I to each well and incubate on shaker (RT) at slow speed for 5 minutes. Repeat 2 more times. Aspirate 1x Wash Buffer I. Add 2mL of 1x Wash Buffer II to each well and incubate on shaker (RT) at slow speed. Repeat 1 more time. Aspirate 1x Wash Buffer II.

Add Biotinylated Antibody Cocktail into each well and incubate on shaker (RT) at slow speed for 2 hours. Aspirate Biotinylated Antibody Cocktail. Add 2mL of 1x Wash Buffer I to each well and incubate on shaker (RT) at slow speed for 5 minutes. Repeat 2 more times. Aspirate 1x Wash Buffer I. Add 2mL of 1x Wash Buffer II to each well and incubate on shaker (RT) at slow speed. Repeat 1 more time. Aspirate 1x Wash Buffer II.

Add HRP-Streptavidin into each well and incubate on shaker (RT) at slow speed for 2 hours. Aspirate HRP-Streptavidin. Add 2mL of 1x Wash Buffer I to each well and incubate on shaker (RT) at slow speed for 5 minutes. Repeat 2 more times. Aspirate 1x Wash Buffer I. Add 2mL of 1x Wash Buffer II to each well and incubate on shaker (RT) at slow speed. Repeat 1 more time. Aspirate 1x Wash Buffer II. Pick up membrane and remove excess solution by touching kimwipe. Lay on plastic sheet. Add as many membranes as needed. Add 400μL of Detection Buffer onto each membrane and wait 2 minutes. Carefully sandwich membranes with another plastic sheet. Image.

Western blot

For the analysis, sample from protein fraction (separated from mitochondria) was separated on a 4-12% Bolt Bis-Tris gel (Invitrogen, NW04125BOX) and transferred using an iBlot2 Transfer system (Invitrogen) to PVDF membranes (Invitrogen, IB242002) at 23 V for 3 minutes. Membranes were blocked for 5 minutes with blocking buffer then incubated with primary antibodies diluted in blocking buffer on a rocker 4 °C overnight. The following day, membranes were washed 4 10-minute washes in TBST at room temperature. Membranes were incubated in a secondary antibody solution for 1 h in blocking buffer on a rocker 4 °C. membranes were washed 4 10-minute washes in TBST at room temperature, the membranes were incubated for 5 minutes in the dark with SuperSignal™ West Pico Plus substrate ((Invitrogen, 34577) used for actin and Tuj1 detection) or SuperSignal™ West Atto Ultimate Sensitivity substrate ((Invitrogen, A38555) used for detection of the rest of the proteins) to identify the proteins of interest. The membranes were serially imaged using a BioRad ChemiDoc MP. The images were analyzed using ImageJ Software.

Western blot Primary: pDRP1 (S616) Rb (Cell Signaling 3455S), DRP1 (DNM1L) Ms (Invitrogen REF MA5-26255), TOMM20 Rb (Invitrogen REF MA5-24859), MTCO1 Ms (Invitrogen REF 459600), VDAC1 Rb (Cell Signalling D73D12), pTau (Ser202 & Thr205) Ms (Invitrogen REF MN1020), TDP43 Rb (Abcam ab109535), APP Rb (Invitrogen REF PA5-17829), Connexin 43 Rb (Cell Signal 3512), Annexin A2 Ms (Invitrogen REF 03-4400). Western Blot Secondary: Anti-rabbit IgG, HRP-linked (Cell Signaling 7074), Anti-mouse IgG, HRP-linked (Cell Signaling 7076)

FAC Sorting for 2D transduced cell lines

Before FAC Sorting, 4 different samples must be prepared: control cells without Propidium Iodide (PI), control cells with PI, transduced cells without PI, and transduced cells with PI. Collect all MOIs of cells (including control) by aspirating media, adding 3 mL of trypsin (microglia and astrocytes) for 3

min or Trip-LE (hi-NSC) for 1 min in 37 C incubator, hit dish and add 3 mL of corresponding cell type media, collect in individual tubes. Count cells. For control cells, split evenly into two groups. For the transduced cell line with the brightest fluorescence, separate 10% of cells to a separate tube. Centrifuge all cells at 1000 rpm for 5 min (microglia and astrocyte) or 3000 rpm for 2 min (hi-NSC). Aspirate supernatant. For all tubes, add cell sorting buffer(composed of 1x PBS, 1% Bovine Serum Albumin (ThermoFisher NC9871802), and 25 mM HEPES buffer (ThermoFisher 11-039-021)) based on 20,000,000 cells/mL. For one of the control tubes and the separated number of cells from the brightest fluorescent cell line, don't add PI. For the rest of the tubes, add PI based on 1 microL/mL. Filter cells with 35 µm strainer cap on 5 mL polystyrene FACS tubes. The FACS machines at the Cincinnati Children's Hospital Medical Center included BD FACSAria II.

FACS for cell-specific mitochondria sorting

Before FAC Sorting, 9 different samples must be prepared DNA Quantification: sham scaffolds, CCI scaffolds, NAM control scaffolds, monoculture RFP-iNSC, monoculture GFP-astrocytes, monoculture BFP-HMC3, monoculture iNSC, monoculture astrocytes, and monoculture microglia. We will use Seahorse Media (50mL F12 Phenol Red-free media, 0.5mL pyruvate, 0.5mL glucose, 0.5mL glutamine) previously mentioned to resuspend our samples. The MitoTracker Stock Solution is composed of 92µL DMSO + 50µg DeepRed MitoTracker while the MitoTracker Working Solution is made of 0.5µL of MitoTracker stock + 250µL SH buffer. 5% Triton X: 2.5mL Triton X + 47.5mL PBS (vortex until mixed). Thaw cryovials of cells, centrifuge according to cell culture protocol, aspirate supernatant and add to 2 mL tube. Use tweezers to transfer designated scaffolds to 2 mL tubes. Add 800 µL of Reagent A to all tubes. Use a P1000 pippette tip to release cells in scaffolds. Proceed with mitochondria isolation protocol. Split NAM control solution. Add 15 µL of MitoTracker working solution to all samples except one of the NAM control solutions. Take samples to facility and proceed with sorting.

Aspect Ratio

Mitochondrial aspect ratio (length-to-width ratio) was measured using a macro available in NIH ImageJ software, following a previously published protocol [cite here]. Maximum projection images of mitochondria, labeled with cell-specific fluorescent markers (red for neurons, blue for microglia, and green for astrocytes), were acquired using a [insert Nikon model here] confocal microscope and exported prior to analysis. The exported images were processed using the ImageJ macro to calculate the aspect ratio (AR) for each mitochondrion. Measurements were obtained from at least four images per sample, and the AR values were plotted as a distribution graph for comparative analysis across experimental groups.

ELISA

Amyloid B 40 and Amyloid B 42 were measured using previously established protocols with slight optimization to better fit our system. A serial dilution of a prepared 2000pg/mL human amyloid beta standard was used to generate a standard curve. 50uL each of either sample or standard was added to the appropriate wells, followed by 50uL of Detection antibody. The plate was gently agitated to mix, and the incubated overnight on a rocker at 4* Celsius. After incubation, 4 instant washes were conducted using 1x wash buffer. Following the washes, 100ul of Anti-Rabbit HRP was added to each well and incubated at room temperature for 30 minutes on a shaker. Afterwards 100ul of stabilized chromogen was added, and the plate was placed in a dark environment and periodically observed to ensure over-saturation does not occur. Once optimal color development was achieved, 100ul of stop

solution was added to each well. The plate was then read at 450nm, and a standard curve was applied to normalize and extract data. Data were reported as the ratio of Ab42 to Ab40.

Messenger RNA (mRNA) sequencing and bioinformatic analysis

Approximately 1 µg of total RNA for each sample underwent RNA quality control analyses before library preparation and bulk mRNA sequencing, as previously described [50, 51]. RNA integrity numbers (RIN) were determined with a Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA), which indicated minimal RNA degradation. Illumina Novaseq sequencers generated paired-end reads (150 bp) at a read depth of 20 million reads [50].

Azenta processed raw .fastq files and returned counts with featureCounts. DESeq2 R package was used to normalize and derive differentially expressed genes by comparing CCI and sham groups, 6 hours and 48 hours post injury. comparing injury and Sham conditions, with or without P110 treatment. Gene set enrichment analyses (GSEA) were performed using the desktop software (https://www.gsea-msigdb.org/gsea/index.jsp) [56, 57] with MitoCarta3.0 [58]. Heatmaps were generated using ggplot and Illustrator. Gene module co-expression analyses were performed with the CEMiTool R package [60, 61], with the Reactome pathway database (https://reactome.org/), and signed interaction network analysis utilizing high confidence STRING protein-protein interactions (STRING score > 700). Oxidative phosphorylation complex gene set used [59] dataset.

References:

Dewan, M. C. *et al.* Estimating the global incidence of traumatic brain injury. *J Neurosurg*, 1-18 (2018). https://doi.org/10.3171/2017.10.JNS17352

Osier, N. D., Carlson, S. W., DeSana, A. & Dixon, C. E. Chronic Histopathological and Behavioral Outcomes of Experimental Traumatic Brain Injury in Adult Male Animals. *J Neurotrauma* **32**, 1861-1882 (2015). https://doi.org/10.1089/neu.2014.3680

Villapol, S., Byrnes, K. R. & Symes, A. J. Temporal dynamics of cerebral blood flow, cortical damage, apoptosis, astrocyte-vasculature interaction and astrogliosis in the pericontusional region after traumatic brain injury. *Front Neurol* **5**, 82 (2014). https://doi.org/10.3389/fneur.2014.00082

Ladak, A. A., Enam, S. A. & Ibrahim, M. T. A Review of the Molecular Mechanisms of Traumatic Brain Injury. *World Neurosurg* **131**, 126-132 (2019). https://doi.org/10.1016/j.wneu.2019.07.039

Johnson, V. E., Stewart, W. & Smith, D. H. Traumatic brain injury and amyloid-beta pathology: a link to Alzheimer's disease? *Nat Rev Neurosci* **11**, 361-370 (2010). https://doi.org/10.1038/nrn2808

Ramos-Cejudo, J. *et al.* Traumatic Brain Injury and Alzheimer's Disease: The Cerebrovascular Link. *EBioMedicine* **28**, 21-30 (2018). https://doi.org/10.1016/j.ebiom.2018.01.021

Wu, D. *et al.* Traumatic Brain Injury Accelerates the Onset of Cognitive Dysfunction and Aggravates Alzheimer's-Like Pathology in the Hippocampus by Altering the Phenotype of Microglia in the APP/PS1 Mouse Model. *Front Neurol* **12**, 666430 (2021). https://doi.org/10.3389/fneur.2021.666430

Srinivasan, G. & Brafman, D. A. The Emergence of Model Systems to Investigate the Link Between Traumatic Brain Injury and Alzheimer's Disease. *Front Aging Neurosci* **13**, 813544 (2021). https://doi.org/10.3389/fnagi.2021.813544

Cherry, J. D. *et al.* Microglial neuroinflammation contributes to tau accumulation in chronic traumatic encephalopathy. *Acta Neuropathol Commun* **4**, 112 (2016). https://doi.org/10.1186/s40478-016-0382-8

Bader, V. & Winklhofer, K. F. Mitochondria at the interface between neurodegeneration and neuroinflammation. *Semin Cell Dev Biol* **99**, 163-171 (2020). https://doi.org/10.1016/j.semcdb.2019.05.028

Herst, P. M., Rowe, M. R., Carson, G. M. & Berridge, M. V. Functional Mitochondria in Health and Disease. *Front Endocrinol (Lausanne)* **8**, 296 (2017). https://doi.org/10.3389/fendo.2017.00296

Norat, P. *et al.* Mitochondrial dysfunction in neurological disorders: Exploring mitochondrial transplantation. *NPJ Regen Med* **5**, 22 (2020). https://doi.org/10.1038/s41536-020-00107-x

Vike, N. L. *et al.* A preliminary model of football-related neural stress that integrates metabolomics with transcriptomics and virtual reality. *iScience* **25**, 103483 (2022). https://doi.org/10.1016/j.isci.2021.103483

Bari, S. *et al.* Integrating multi-omics with neuroimaging and behavior: A preliminary model of dysfunction in football athletes. *Neuroimage: Reports* **1** (2021). https://doi.org/10.1016/j.ynirp.2021.100032

Hubbard, W. B. *et al.* Acute Mitochondrial Impairment Underlies Prolonged Cellular Dysfunction after Repeated Mild Traumatic Brain Injuries. *J Neurotrauma* **36**, 1252-1263 (2019). https://doi.org/10.1089/neu.2018.5990

Song, H. *et al.* Proteomic Analysis and Biochemical Correlates of Mitochondrial Dysfunction after Low-Intensity Primary Blast Exposure. *J Neurotrauma* **36**, 1591-1605 (2019). https://doi.org/10.1089/neu.2018.6114

Lyons, D. N. *et al.* A Mild Traumatic Brain Injury in Mice Produces Lasting Deficits in Brain Metabolism. *J Neurotrauma* **35**, 2435-2447 (2018). https://doi.org/10.1089/neu.2018.5663

Robertson, C. L., Saraswati, M. & Fiskum, G. Mitochondrial dysfunction early after traumatic brain injury in immature rats. *J Neurochem* **101**, 1248-1257 (2007). https://doi.org/10.1111/j.1471-4159.2007.04489.x

Yonutas, H. M., Vekaria, H. J. & Sullivan, P. G. Mitochondrial specific therapeutic targets following brain injury. *Brain Res* **1640**, 77-93 (2016). https://doi.org/10.1016/j.brainres.2016.02.007

Ahluwalia, M. *et al.* Rescuing mitochondria in traumatic brain injury and intracerebral hemorrhages - A potential therapeutic approach. *Neurochem Int* **150**, 105192 (2021). https://doi.org/10.1016/j.neuint.2021.105192

Lazzarino, G. *et al.* Pyruvate Dehydrogenase and Tricarboxylic Acid Cycle Enzymes Are Sensitive Targets of Traumatic Brain Injury Induced Metabolic Derangement. *Int J Mol Sci* **20** (2019). https://doi.org/10.3390/ijms20225774

Xing, G., Ren, M., Watson, W. D., O'Neill, J. T. & Verma, A. Traumatic brain injury-induced expression and phosphorylation of pyruvate dehydrogenase: a mechanism of dysregulated glucose metabolism. *Neurosci Lett* **454**, 38-42 (2009). https://doi.org/10.1016/j.neulet.2009.01.047

Huynh, L. M., Burns, M. P., Taub, D. D., Blackman, M. R. & Zhou, J. Chronic Neurobehavioral Impairments and Decreased Hippocampal Expression of Genes Important for Brain Glucose Utilization in a Mouse Model of Mild TBI. *Front Endocrinol (Lausanne)* **11**, 556380 (2020). https://doi.org/10.3389/fendo.2020.556380

Xing, G. *et al.* Pyruvate dehydrogenase phosphatase1 mRNA expression is divergently and dynamically regulated between rat cerebral cortex, hippocampus and thalamus after traumatic brain injury: a potential biomarker of TBI-induced hyper- and hypo-glycaemia and neuronal vulnerability. *Neurosci Lett* **525**, 140-145 (2012). https://doi.org/10.1016/j.neulet.2012.07.055

Hubbard, W. B., Harwood, C. L., Geisler, J. G., Vekaria, H. J. & Sullivan, P. G. Mitochondrial uncoupling prodrug improves tissue sparing, cognitive outcome, and mitochondrial bioenergetics after traumatic brain injury in male mice. *J Neurosci Res* **96**, 1677-1688 (2018). https://doi.org/10.1002/jnr.24271

Arun, P. *et al.* Acute mitochondrial dysfunction after blast exposure: potential role of mitochondrial glutamate oxaloacetate transaminase. *J Neurotrauma* **30**, 1645-1651 (2013). https://doi.org/10.1089/neu.2012.2834

Agrawal, R. *et al.* Dietary fructose aggravates the pathobiology of traumatic brain injury by influencing energy homeostasis and plasticity. *J Cereb Blood Flow Metab* **36**, 941-953 (2016). https://doi.org/10.1177/0271678X15606719

Hill, R. L., Singh, I. N., Wang, J. A., Kulbe, J. R. & Hall, E. D. Protective effects of phenelzine administration on synaptic and non-synaptic cortical mitochondrial function and lipid peroxidation-mediated oxidative damage following TBI in young adult male rats. *Exp Neurol* **330**, 113322 (2020). https://doi.org/10.1016/j.expneurol.2020.113322

Lamade, A. M. *et al.* Mitochondrial damage & lipid signaling in traumatic brain injury. *Exp Neurol* **329**, 113307 (2020). https://doi.org/10.1016/j.expneurol.2020.113307

Ryu, W. I. *et al.* Brain cells derived from Alzheimer's disease patients have multiple specific innate abnormalities in energy metabolism. *Mol Psychiatry* **26**, 5702-5714 (2021). https://doi.org/10.1038/s41380-021-01068-3

Yu, L., Jin, J., Xu, Y. & Zhu, X. Aberrant Energy Metabolism in Alzheimer's Disease. *J Transl Int Med* **10**, 197-206 (2022). https://doi.org/10.2478/jtim-2022-0024

Bhatti, J. S., Bhatti, G. K. & Reddy, P. H. Mitochondrial dysfunction and oxidative stress in metabolic disorders - A step towards mitochondria based therapeutic strategies. *Biochim Biophys Acta Mol Basis Dis* **1863**, 1066-1077 (2017). https://doi.org/10.1016/j.bbadis.2016.11.010

Liaudanskaya, V. *et al.* Modeling Controlled Cortical Impact Injury in 3D Brain-Like Tissue Cultures. *Adv Healthc Mater* **9**, e2000122 (2020). https://doi.org/10.1002/adhm.202000122

Liaudanskaya, V. *et al.* Mitochondria dysregulation contributes to secondary neurodegeneration progression post-contusion injury in human 3D in vitro triculture brain tissue model. *Cell Death Dis* **14**, 496 (2023). https://doi.org/10.1038/s41419-023-05980-0

Rouleau, N. *et al.* A Long-Living Bioengineered Neural Tissue Platform to Study Neurodegeneration. *Macromol Biosci* **20**, e2000004 (2020). https://doi.org/10.1002/mabi.202000004

Snapper, D. M. *et al.* Development of a novel bioengineered 3D brain-like tissue for studying primary blast-induced traumatic brain injury. *J Neurosci Res* **101**, 3-19 (2023). https://doi.org/10.1002/jnr.25123

Tang-Schomer, M. D. *et al.* Bioengineered functional brain-like cortical tissue. *Proc Natl Acad Sci U S A* **111**, 13811-13816 (2014). https://doi.org/10.1073/pnas.1324214111

Garza, R. *et al.* Single-cell transcriptomics of human traumatic brain injury reveals activation of endogenous retroviruses in oligodendroglia. *Cell Rep* **42**, 113395 (2023). https://doi.org/10.1016/j.celrep.2023.113395

Russo, P. S. T. *et al.* CEMiTool: a Bioconductor package for performing comprehensive modular co-expression analyses. *BMC Bioinformatics* **19**, 56 (2018). https://doi.org/10.1186/s12859-018-2053-1

Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008). https://doi.org/10.1186/1471-2105-9-559

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005). https://doi.org/10.1073/pnas.0506580102

Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267-273 (2003). https://doi.org/10.1038/ng1180

Rath, S. *et al.* MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic Acids Res* **49**, D1541-D1547 (2021). https://doi.org/10.1093/nar/gkaa1011

Guarnieri, J. W. *et al.* Core mitochondrial genes are down-regulated during SARS-CoV-2 infection of rodent and human hosts. *Sci Transl Med* **15**, eabq1533 (2023). https://doi.org/10.1126/scitranslmed.abq1533

Abu Hamdeh, S. *et al.* Differential DNA Methylation of the Genes for Amyloid Precursor Protein, Tau, and Neurofilaments in Human Traumatic Brain Injury. *J Neurotrauma* **38**, 1679-1688 (2021). https://doi.org/10.1089/neu.2020.7283

Sabbir, M. G. Loss of calcium/calmodulin-dependent protein kinase kinase 2, transferrin, and transferrin receptor proteins in the temporal cortex of Alzheimer's patients postmortem is associated with abnormal iron homeostasis: implications for patient survival. *Front Cell Dev Biol* **12**, 1469751 (2024). https://doi.org/10.3389/fcell.2024.1469751

Ghosh, A. & Giese, K. P. Calcium/calmodulin-dependent kinase II and Alzheimer's disease. *Mol Brain* **8**, 78 (2015). https://doi.org/10.1186/s13041-015-0166-2

Abiose, O. *et al.* Post-translational modifications linked to preclinical Alzheimer's disease-related pathological and cognitive changes. *Alzheimers Dement* **20**, 1851-1867 (2024). https://doi.org/10.1002/alz.13576

Jaisa-Aad, M., Munoz-Castro, C., Healey, M. A., Hyman, B. T. & Serrano-Pozo, A. Characterization of monoamine oxidase-B (MAO-B) as a biomarker of reactive astrogliosis in Alzheimer's disease and related dementias. *Acta Neuropathol* **147**, 66 (2024). https://doi.org/10.1007/s00401-024-02712-2

Escartin, C. *et al.* Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci* **24**, 312-325 (2021). https://doi.org/10.1038/s41593-020-00783-4

Wei, Z., Koya, J. & Reznik, S. E. Insulin Resistance Exacerbates Alzheimer Disease via Multiple Mechanisms. *Front Neurosci* **15**, 687157 (2021). https://doi.org/10.3389/fnins.2021.687157

Gonzalez-Rodriguez, M. *et al.* Neurodegeneration and Astrogliosis in the Human CA1 Hippocampal Subfield Are Related to hsp90ab1 and bag3 in Alzheimer's Disease. *Int J Mol Sci* **23** (2021). https://doi.org/10.3390/ijms23010165

Lackie, R. E. *et al.* Increased levels of Stress-inducible phosphoprotein-1 accelerates amyloid-beta deposition in a mouse model of Alzheimer's disease. *Acta Neuropathol Commun* **8**, 143 (2020). https://doi.org/10.1186/s40478-020-01013-5

Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* **153**, 707-720 (2013). https://doi.org/10.1016/j.cell.2013.03.030

Cermak, S. *et al.* Loss of Cathepsin B and L Leads to Lysosomal Dysfunction, NPC-Like Cholesterol Sequestration and Accumulation of the Key Alzheimer's Proteins. *PLoS One* **11**, e0167428 (2016). https://doi.org/10.1371/journal.pone.0167428

Kim, A. & Benavente, C. A. Oncogenic Roles of UHRF1 in Cancer. *Epigenomes* **8** (2024). https://doi.org/10.3390/epigenomes8030026

Miyashita, R. *et al.* The termination of UHRF1-dependent PAF15 ubiquitin signaling is regulated by USP7 and ATAD5. *Elife* **12** (2023). https://doi.org/10.7554/eLife.79013

Wei-Shan, H., Amit, V. C. & Clarke, D. J. Cell cycle regulation of condensin Smc4. *Oncotarget* **10**, 263-276 (2019). https://doi.org/10.18632/oncotarget.26467

Ramachandran, K. *et al.* Transcriptional programming of translation by BCL6 controls skeletal muscle proteostasis. *Nat Metab* **6**, 304-322 (2024). https://doi.org/10.1038/s42255-024-00983-3

Baskin, K. K. *et al.* MED12 regulates a transcriptional network of calcium-handling genes in the heart. *JCI Insight* **2** (2017). https://doi.org/10.1172/jci.insight.91920

Dabovic, B. *et al.* Control of lung development by latent TGF-beta binding proteins. *J Cell Physiol* **226**, 1499-1509 (2011). https://doi.org/10.1002/jcp.22479

Moon, D. O. Deciphering the Role of BCAR3 in Cancer Progression: Gene Regulation, Signal Transduction, and Therapeutic Implications. *Cancers (Basel)* **16** (2024). https://doi.org/10.3390/cancers16091674

Taiko, I. *et al.* Selection of red fluorescent protein for genetic labeling of mitochondria and intercellular transfer of viable mitochondria. *Sci Rep* **12**, 19841 (2022). https://doi.org/10.1038/s41598-022-24297-0

Carpenter, K. L., Jalloh, I. & Hutchinson, P. J. Glycolysis and the significance of lactate in traumatic brain injury. *Front Neurosci* **9**, 112 (2015). https://doi.org/10.3389/fnins.2015.00112

Stovell, M. G. *et al.* Assessing Metabolism and Injury in Acute Human Traumatic Brain Injury with Magnetic Resonance Spectroscopy: Current and Future Applications. *Front Neurol* **8**, 426 (2017). https://doi.org/10.3389/fneur.2017.00426

Xu, X. J. *et al.* Glucose metabolism: A link between traumatic brain injury and Alzheimer's disease. *Chin J Traumatol* **24**, 5-10 (2021). https://doi.org/10.1016/j.cjtee.2020.10.001

Cleland, N. R. W., Al-Juboori, S. I., Dobrinskikh, E. & Bruce, K. D. Altered substrate metabolism in neurodegenerative disease: new insights from metabolic imaging. *J Neuroinflammation* **18**, 248 (2021). https://doi.org/10.1186/s12974-021-02305-w

Sundman, M. H., Hall, E. E. & Chen, N. K. Examining the relationship between head trauma and neurodegenerative disease: A review of epidemiology, pathology and neuroimaging techniques. *J Alzheimers Dis Parkinsonism* **4** (2014). https://doi.org/10.4172/2161-0460.1000137

Gardner, R. C. *et al.* Mild TBI and risk of Parkinson's disease: A chronic effect of Neurotrauma Consortium Study. *Neurology* **90** (2018).

Lizhnyak, P. N. & Ottens, A. K. Proteomics: in pursuit of effective traumatic brain injury therapeutics. *Expert Review of Proteomics* **12** (2015).

Borst, K., Schwabenland, M. & Prinz, M. Microglia metabolism in health and disease. *Neurochem Int* **130**, 104331 (2019). https://doi.org/10.1016/j.neuint.2018.11.006

Brett, B. L., Gardner, R. C., Godbout, J., Dams-O'Connor, K. & Keene, C. D. Traumatic Brain Injury and Risk of Neurodegenerative Disorder. *Biol Psychiatry* **91**, 498-507 (2022). https://doi.org/10.1016/j.biopsych.2021.05.025

Guerrero, A., De Strooper, B. & Arancibia-Carcamo, I. L. Cellular senescence at the crossroads of inflammation and Alzheimer's disease. *Trends Neurosci* **44**, 714-727 (2021). https://doi.org/10.1016/j.tins.2021.06.007

Haukedal, H. & Freude, K. K. Implications of Glycosylation in Alzheimer's Disease. *Front Neurosci* **14**, 625348 (2020). https://doi.org/10.3389/fnins.2020.625348

Lauro, C. & Limatola, C. Metabolic Reprograming of Microglia in the Regulation of the Innate Inflammatory Response. *Front Immunol* **11**, 493 (2020). https://doi.org/10.3389/fimmu.2020.00493

Strope, T. A., Birky, C. J. & Wilkins, H. M. The Role of Bioenergetics in Neurodegeneration. *Int J Mol Sci* **23** (2022). https://doi.org/10.3390/ijms23169212

Zong, Y. *et al.* Mitochondrial dysfunction: mechanisms and advances in therapy. *Signal Transduct Target Ther* **9**, 124 (2024). https://doi.org/10.1038/s41392-024-01839-8

Picard, M. & Shirihai, O. S. Mitochondrial signal transduction. *Cell Metab* **34**, 1620-1653 (2022). https://doi.org/10.1016/j.cmet.2022.10.008

Amorim, J. A. *et al.* Mitochondrial and metabolic dysfunction in ageing and age-related diseases. *Nat Rev Endocrinol* **18**, 243-258 (2022). https://doi.org/10.1038/s41574-021-00626-7

Belenguer, P., Duarte, J. M. N., Schuck, P. F. & Ferreira, G. C. Mitochondria and the Brain: Bioenergetics and Beyond. *Neurotox Res* **36**, 219-238 (2019). https://doi.org/10.1007/s12640-019-00061-7

Calvo-Rodriguez, M. & Bacskai, B. J. Mitochondria and Calcium in Alzheimer's Disease: From Cell Signaling to Neuronal Cell Death. *Trends Neurosci* **44**, 136-151 (2021). https://doi.org/10.1016/j.tins.2020.10.004

Chen, L. *et al.* Mitochondrial heterogeneity in diseases. *Signal Transduct Target Ther* **8**, 311 (2023). https://doi.org/10.1038/s41392-023-01546-w

Chen, W., Zhao, H. & Li, Y. Mitochondrial dynamics in health and disease: mechanisms and potential targets. *Signal Transduct Target Ther* **8**, 333 (2023). https://doi.org/10.1038/s41392-023-01547-9

Cheng, Y. & Bai, F. The Association of Tau With Mitochondrial Dysfunction in Alzheimer's Disease. *Front Neurosci* **12**, 163 (2018). https://doi.org/10.3389/fnins.2018.00163

Collier, J. J., Olahova, M., McWilliams, T. G. & Taylor, R. W. Mitochondrial signalling and homeostasis: from cell biology to neurological disease. *Trends Neurosci* **46**, 137-152 (2023). https://doi.org/10.1016/j.tins.2022.12.001

Chakraborty, R., Nonaka, T., Hasegawa, M. & Zurzolo, C. Tunnelling nanotubes between neuronal and microglial cells allow bi-directional transfer of alpha-Synuclein and mitochondria. *Cell Death Dis* **14**, 329 (2023). https://doi.org/10.1038/s41419-023-05835-8

Pan, K. H., Chang, H. & Yang, W. Y. Extracellular release in the quality control of the mammalian mitochondria. *J Biomed Sci* **30**, 85 (2023). https://doi.org/10.1186/s12929-023-00979-3

Pinto, G. *et al.* Patient-derived glioblastoma stem cells transfer mitochondria through tunneling nanotubes in tumor organoids. *Biochem J* **478**, 21-39 (2021). https://doi.org/10.1042/BCJ20200710

Scheiblich, H. *et al.* Microglia jointly degrade fibrillar alpha-synuclein cargo by distribution through tunneling nanotubes. *Cell* **184**, 5089-5106 e5021 (2021). https://doi.org/10.1016/j.cell.2021.09.007

Zampieri, L. X., Silva-Almeida, C., Rondeau, J. D. & Sonveaux, P. Mitochondrial Transfer in Cancer: A Comprehensive Review. *Int J Mol Sci* **22** (2021). https://doi.org/10.3390/ijms22063245

Zhao, Z. *et al.* Extracellular mitochondria released from traumatized brains induced platelet procoagulant activity. *Haematologica* **105**, 209-217 (2020). https://doi.org/10.3324/haematol.2018.214932

Lin, A. P. *et al.* MYC, mitochondrial metabolism and O-GlcNAcylation converge to modulate the activity and subcellular localization of DNA and RNA demethylases. *Leukemia* **36**, 1150-1159 (2022). https://doi.org/10.1038/s41375-021-01489-7

Tracey, T. J., Steyn, F. J., Wolvetang, E. J. & Ngo, S. T. Neuronal Lipid Metabolism: Multiple Pathways Driving Functional Outcomes in Health and Disease. *Front Mol Neurosci* **11**, 10 (2018). https://doi.org/10.3389/fnmol.2018.00010

Fecher, C. *et al.* Cell-type-specific profiling of brain mitochondria reveals functional and molecular diversity. *Nat Neurosci* **22**, 1731-1742 (2019). https://doi.org/10.1038/s41593-019-0479-z

Zhang, Y. *et al.* Synergistic label-free fluorescence imaging and miRNA studies reveal dynamic human neuron-glial metabolic interactions following injury. *Sci Adv* **10**, eadp1980 (2024). https://doi.org/10.1126/sciadv.adp1980

Hubbard, W. B. *et al.* Mitochondrial Dysfunction After Repeated Mild Blast Traumatic Brain Injury Is Attenuated by a Mild Mitochondrial Uncoupling Prodrug. *J Neurotrauma* **40**, 2396-2409 (2023). https://doi.org/10.1089/neu.2023.0102

Vafai, S. B. & Mootha, V. K. Mitochondrial disorders as windows into an ancient organelle. *Nature* **491**, 374-383 (2012). https://doi.org/10.1038/nature11707

Mattson, M. P., Gleichmann, M. & Cheng, A. Mitochondria in neuroplasticity and neurological disorders. *Neuron* **60**, 748-766 (2008). https://doi.org/10.1016/j.neuron.2008.10.010

Cherry, J. D., Olschowka, J. A. & O'Banion, M. K. Neuroinflammation and M2 microglia: the good, the bad, and the inflamed. *J Neuroinflammation* **11**, 98 (2014). https://doi.org/10.1186/1742-2094-11-98

Cheng, A., Hou, Y. & Mattson, M. P. Mitochondria and neuroplasticity. *ASN Neuro* **2**, e00045 (2010). https://doi.org/10.1042/AN20100019

Lee, A. *et al.* Abeta42 oligomers trigger synaptic loss through CAMKK2-AMPK-dependent effectors coordinating mitochondrial fission and mitophagy. *Nat Commun* **13**, 4444 (2022). https://doi.org/10.1038/s41467-022-32130-5

Devanney, N. A., Stewart, A. N. & Gensel, J. C. Microglia and macrophage metabolism in CNS injury and disease: The role of immunometabolism in neurodegeneration and neurotrauma. *Exp Neurol* **329**, 113310 (2020). https://doi.org/10.1016/j.expneurol.2020.113310

Ashton, N. J. *et al.* Differential roles of Abeta42/40, p-tau231 and p-tau217 for Alzheimer's trial selection and disease monitoring. *Nat Med* **28**, 2555-2562 (2022). https://doi.org/10.1038/s41591-022-02074-w

Karran, E. & De Strooper, B. The amyloid hypothesis in Alzheimer disease: new insights from new therapeutics. *Nat Rev Drug Discov* **21**, 306-318 (2022). https://doi.org/10.1038/s41573-022-00391-w

Rockwood, D. N. *et al.* Materials fabrication from Bombyx mori silk fibroin. *Nat Protoc* **6**, 1612-1631 (2011). https://doi.org/10.1038/nprot.2011.379

Dingle, Y. L., Bonzanni, M., Liaudanskaya, V., Nieland, T. J. F. & Kaplan, D. L. Integrated functional neuronal network analysis of 3D silk-collagen scaffold-based mouse cortical culture. *STAR Protoc* **2**, 100292 (2021). https://doi.org/10.1016/j.xpro.2020.100292