μCeta: a set of cetacean-specific primers for environmental DNA metabarcoding with minimal amplification of non-target vertebrates

Masayuki Ushio[1,*], Sachie Ozawa[2], Shin-ichiro Oka[2], Tetsuya Sado[3], Robinson O. Kisero[1], Lindsay Porter[4], Eszter Matrai[5], Masaki Miya[3]

[1]Department of Ocean Science, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR, China, [2]Okinawa Churashima Foundation, 888 Aza Ishikawa, Motobu-cho, Okinawa 905-0206, Japan, [3]Natural History Museum and Institute, Chiba, Chiba 260-8682, Japan, [4]Southeast Asia Marine Mammal Research, Lippo Centre, Tower 1, 89, Queensway, Hong Kong SAR, China, [5]Ocean Park Hong Kong, 189 Wong Chuk Hang Road, Aberdeen, Hong Kong SAR, China

ABSTRACT

Biodiversity monitoring is crucial for understanding ecosystem dynamics and species distributions, particularly in the context of anthropogenic impacts and climate change. Cetaceans, as key indicator species of marine ecosystems, face increasing threats from human activities, highlighting the need for effective, non-invasive monitoring methods. In the present study, we developed novel Cetacea-specific primer sets to enhance the detection efficiency of cetacean species through environmental DNA (eDNA) metabarcoding, while minimizing the amplification of non-target vertebrates, such as fish and humans. We retrieved 71 cetacean mitochondrial genomes from a public database and designed 20 candidate primer sets, which were assessed *in silico* for their specificity and capacity to differentiate cetacean sequences. Four primer sets with the best *in silico* performance were selected for empirical validation using DNA from tissue samples and eDNA from seawater collected from aquarium pools and Hong Kong coastal waters. All four primer sets effectively amplified cetacean DNA from tissue samples. However, in the aquarium pool tests, three primer sets failed to accurately identify one or more cetacean species due to a lack of interspecific variation within the amplified region. From these, we selected one primer set targeting 268 bp insert length in mitochondrial 12S rRNA region, named μCeta, and applied it to water samples collected from Hong Kong coastal waters, where the iconic Indo-Pacific humpback dolphin (*Sousa chinensis*) was observed. μCeta successfully detected *S. chinensis* eDNA while avoiding amplification of non-target species such as fish or humans. Our results demonstrate that μCeta is a reliable tool for cetacean eDNA detection, contributing to cetacean conservation and enhancing our understanding of marine biodiversity.

Keywords: Biodiversity monitoring, Cetaceans, Conservation, Environmental DNA, Metabarcoding, Primer design

1. Introduction

Biodiversity monitoring is crucial for conservation and ecological research, offering critical

insights into ecosystem status, species distributions, and the impacts of ongoing human

activities and climate change on ecosystem functioning (Cardinale et al., 2012). Ecosystem

functions—such as biomass production, nutrient cycling, and sustainable food production—

are directly linked to biodiversity (e.g., Hooper et al., 2005; Mace et al., 2012), underscoring

the importance of maintaining diverse biological communities. Traditional monitoring

methods, including direct visual censuses and camera and video traps, have long been used to

assess species presence and abundance (Masuda et al., 2010; Nakagawa, 2019; Samejima et

al., 2012). These methods provide invaluable data for enhancing our understanding,

conservation, and management of ecosystems under increasing anthropogenic pressures.

However, they are often labor-intensive and costly, requiring taxa-specific expertise, which

limits their scalability and the monitoring frequency of biodiversity. Consequently, there is an

increasing demand for complementary tools that provide more efficient and cost-effective

approaches for biodiversity monitoring.

Environmental DNA (eDNA), defined as extra-organismal DNA directly extracted

from environmental samples, has been utilized to detect the presence of macro-organisms

(Ficetola et al., 2008; Miya et al., 2015; Taberlet et al., 2012, 2018). In macro-organisms,

eDNA originates from various sources, including metabolic waste and damaged tissue (Kelly

et al., 2014), and exists in different physical forms in the environment (Power et al., 2023).

eDNA carries species-specific genetic information from the organisms that produced it, and

eDNA analysis, such as metabarcoding, enables detection of multiple species from a single

environmental sample. This approach provides a non-invasive, cost-effective, and highly

sensitive method for biodiversity monitoring in both aquatic and terrestrial environments

(Bista et al., 2017; Bohmann & Lynggaard, 2022; Lynggaard et al., 2022; Ushio et al., 2017;

Yamamoto et al., 2017).

Cetaceans, a term that includes all species of whales, dolphins, and porpoises, are

iconic megafauna that inhabit all marine environments and serve as important indicators of

the status of ocean ecosystems (Plön et al., 2024). However, many cetacean species are

increasingly threatened by human activities and climate change (Li et al., 2023; van Weelden

et al., 2021; Weir & Pierce, 2013). Their conservation is crucial not only for protecting

marine biodiversity, but also for maintaining marine ecosystem functions and services, as

they play critical roles in carbon sequestration, nutrient cycling, trophic cascades and eco-

tourism (Estes et al., 2011; Kiszka et al., 2022; Roman & McCarthy, 2010). For instance,

large, deep-diving cetaceans, such as baleen and sperm whales, contribute to vertical carbon

and nitrogen cycling through their foraging behavior (Pearson et al., 2023). Smaller cetaceans

also contribute to both horizontal (e.g., from coastal areas to the open ocean) and vertical

transport of carbon and nutrients (Kiszka et al., 2022). Given their ecological significance

and vulnerability (more than a quarter of all cetacean species are listed as threatened; *The*

IUCN Red List of Threatened Species, 2024), reliable and efficient monitoring tools are

urgently needed to support the conservation of cetacean diversity and their functional roles in

marine ecosystems.

Cetacean distribution and occurrence is often determined through interviews with sea-

faring communities and their traditional knowledge, visual and acoustic surveys, and, more
recently, satellite image analysis (Bamford et al., 2020; Frasier et al., 2021; Liu et al., 2017).
Visual monitoring methods, including the use of vessel and aerial survey platforms, are
widely used to collect data suitable for various purposes such as species identification,
abundance and distribution estimation, individual recognition, and estimation of body length
(Boyd et al., 2019; Hammond et al., 2017; Thomas et al., 2010; Urian et al., 2015). However,
such surveys are often expensive, rely on good weather windows and are labour intensive.
Also, all monitoring techniques that rely on visual detection require the cetacean species
under study to surface frequently and remain the surface long enough to gather accurate data.
Acoustic monitoring methods are also widely used as nearly all cetacean species produce a
variety of vocalisations, often unique to that species and even to a population, for
communication, navigation, and prey detection (Janik, 2014; Madsen et al., 2023). Passive
acoustic monitoring does not rely on good visibility and therefore, can be used effectively at
night and/or in all weather conditions. In addition, the highly automated nature of acoustic
detection software makes data analysis efficient and timely (Gillespie et al., 2009) and is
particularly effective when combined with vessel survey data (Duporge et al., 2024; Frasier et
al., 2021). Acoustic data, like visual data, can be used to identify species, estimate population
abundance and distribution, but if the species is not very vocal or its habitat is noisy, acoustic
detection may be missed. Satellite imagery analysis is an emerging technique that has the
potential to monitor large areas and remote habitats, however, at this time, standard analytical
methodology is still being developed and the cost of purchasing satellite images or even
tasking a satellite remains prohibitively expensive (Bamford et al., 2020). Further, weather
conditions must be good, for example, low cloud cover and low wind speeds, for clear
images to be obtained (Höschle et al., 2021).
Given these challenges, eDNA analysis offers an efficient new tool for detecting
cetaceans. While eDNA analysis does not cover large spatial areas like satellite imagery
analysis, it has several advantages over other methods: it can provide detailed genetic
information, including intra-specific variation (Sigsgaard et al., 2017; Tsuji et al., 2023),
requires relatively low on-site labor, and can detect cetaceans even when they are submerged
or that have already left the immediate survey area. Additionally, eDNA concentration may
serve as an estimate of biomass or abundance of target species (Rourke et al., 2022; Takahara
et al., 2012), although careful study design and interpretation are essential (Rourke et al.,
2022). Therefore, eDNA-based cetacean monitoring has the potential to enhance both the
efficiency and effectiveness of cetacean monitoring.
Detecting cetaceans using eDNA analysis is a relatively new field. The widely used
fish-targeting universal primer set, MiFish, which amplifies mitochondrial 12S rRNA gene
(Miya et al., 2015), has been shown to detect cetacean eDNA (Closek et al., 2019).
Additionally, cetacean eDNA has been detected using other primers targeting the
mitochondrial genome (Alter et al., 2022; Robinson et al., 2023; Zhang et al., 2023) , such as
the Riaz 12S primers (Riaz et al., 2011), MarVer primers (Valsecchi et al., 2020), and
quantitative PCR approaches (Foote et al., 2012; Hashimoto et al., 2023). Valsecchi et al.
(2020) developed primer sets targeting marine mammals and other vertebrates, including
cetaceans, pinnipeds, elasmobranchs, and bony fish. More recently, Deng et al. (2024)

modified a vertebrate universal primers and detected 18 cetacean species in the South China
Sea. However, these primers were not specifically designed for cetaceans, leading to the
amplification of more abundant non-cetacean eDNA, such as that from fish and common
contaminants (e.g., human eDNA). This non-specific amplification can suppress the detection
of target eDNA (Collins et al., 2019), and this issue is particularly important for cetacean
eDNA metabarcoding because cetacean eDNA is typically present at low concentrations (e.g.,
Baker et al., 2018). These challenges underscore the need for novel primer sets that
selectively amplify cetacean DNA while minimizing the amplification of other marine
organisms, including fish, humans, and other vertebrates. Developing such primers is critical
for establishing effective eDNA-based cetacean monitoring in marine ecosystems.
This study addresses the gap in cetacean-specific eDNA detection by developing and
validating novel cetacean-targeting metabarcoding primer sets with minimal or no
amplification of non-cetacean species. Our objective is to design primers that specifically
target cetacean mitochondrial DNA while minimizing the amplification of non-target species.
To achieve this, we collected mitochondrial genome sequences from cetaceans and related
mammals from public databases, conducted *in silico* analyses to evaluate primer
performance, and empirically tested multiple primer sets in both controlled and natural
environments. Finally, we applied our newly developed primer sets to water samples
collected from Hong Kong waters to detect the 'vulnerable' the Indo-Pacific humpback
dolphin (*Sousa chinensis*) (Jefferson & Smith, 2016). By achieving these objectives, we aim
to establish a robust experimental protocol for cetacean eDNA monitoring, thereby
contributing to the conservation of cetacean species.
149

2. Materials and Methods

2.1 Ethics statement

This study used cetacean tissue samples from stranded individuals found in fields or dead
individuals in an aquarium to validate cetacean-specific primers for eDNA metabarcoding.
All tissue samples were obtained with the necessary permissions from the Okinawa
Churashima Foundation (for all cetacean tissues except *Sousa chinensis*), the Agriculture,
Fisheries and Conservation Department (AFCD) of Hong Kong, and Ocean Park Hong Kong
(for *S. chinensis* samples). Seawater samples from dolphin pools were collected with
permission from the Okinawa Churashima Foundation. No direct interaction with or
sampling from live cetaceans occurred during this research.
160

2.2 Primer design

To design candidate primers, we retrieved 71 cetacean mitochondrial genomes from the
NCBI Reference Sequence Database (https://www.ncbi.nlm.nih.gov/refseq/; accessed 11 July
2023). Additionally, mitochondrial genomes of five mammal species (cattle [*Bos taurus*],
wild boar [*Sus scrofa*], mouse [*Mus musculus*], cat [*Felis catus*], dog [*Canis lupus*], and
human [*Homo sapiens*]) as well as other related taxa (dugong, manatee, and other marine
mammals) were retrieved to assess potential amplification of non-target species (Table S1).

Retrieved sequences were aligned using the online version of MAFFT with default
settings (Katoh & Standley, 2013; https://mafft.cbrc.jp/alignment/server/). Based on these
alignments, 20 candidate regions were identified using the DECIPHER package in R
(Wright, 2016) and visual inspection. Suitable regions for cetacean eDNA metabarcoding
were found in the 12S rRNA, 16S rRNA, and NAD3 genes (Fig. 1 and Table S2).
We designed the primer sets to amplify cetacean DNA while minimizing amplification
of non-cetacean DNA (e.g., other marine mammals and common terrestrial species such as
dogs and cats). In particular, we ensured that the primers do not amplify human DNA, as co-
amplification of human DNA—a common contaminant, especially in coastal regions—could
obscure the detection of low-concentration cetacean DNA. During primer design, we
considered several technical tips: primer length between 19 and 28 bp, GC% of 40–60%, T_m
values of forward and reverse primers are as close as possible, and an insert length of less
than 450 bp to achieve high sensitivity and compatibility with short-read sequencing.

Figure 1 | Candidate primers for Cetacea-specific eDNA metabarcoding designed from the mitochondrial genome. Previously reported mammal and vertebrate-targeting primers and our candidates primers for Cetacea-specific eDNA metabarcoding "MiMammal," "MarVer1," "Riaz12S," and "Ceto2" are previously reported primers, while all other primers were designed in this study. Primers shown in bold face (either in red or black) were empirically validated in this study, and primers in red bold face are newly designed. Mu31F/Dc320R was later named "µCeta." The prefixes "Mu" and "Dc" denote primers designed manually (by M. Ushio) and using DECIPHER, respectively. Each number between the primer regions indicates the insert length in base pairs (bp). The base numbers are derived from the mitochondrial genome sequence of *Delphinus capensis*.

2.3 *In silico* evaluation of the candidate primers

The performance of the candidate primers was evaluated *in silico* alongside other vertebrate-,
mammal-, and cetacean-specific primers, including MiMammal (Ushio et al., 2017),
MarVer1, MarVer2, MarVer3, Ceto2 (Valsecchi et al., 2020), and Riaz12S (Riaz et al., 2011).
Ceto2 was included in our analysis as it was specifically designed as a cetacean primer set.
The characteristics and performance of the candidate primers were assessed using

mitochondrial genomes of vertebrates retrieved from NCBI as of 19 July 2023. A total of
36,250 sequences from 10,855 vertebrate taxa were used for the *in silico* PCR which included
76 cetacean taxa, 2,378 non-cetacean mammal taxa, 1,556 bird taxa, 700 reptile taxa, 733
amphibian taxa, 5,411 fish taxa, and 1 additional taxon (*Branchiostoma belcheri*) (see "The
number of taxa in *in silico* analysis" and Table S3 for details).

In silico PCR was conducted using "search_pcr" and "search_oligodb" commands
implemented in USEARCH (Edgar, 2010). Specifically, we employed the following
commands: `usearch -search_oligodb vertebrate_db.fa -db primer.fa -maxdiffs`
`x -strand both -userout output.txt` (for each primer; `-maxdiffs` was examined up to
4) and `usearch -search_pcr vertebrate_db.fa -db primer_set.fa -maxdiffs x -`
`strand both -minamp 30 -maxamp 2000 -pcrout output.txt -ampout ampout.txt.`
(for each primer set; `-maxdiffs` was examined up to 3 for each primer set).

Based on the results of the *in silico* PCR results, we evaluated several key features of
the primer sets: predicted amplicon lengths, the number of species and sequences potentially
amplified, and the ability to distinguish closely related taxa. The predicted amplicon lengths
and the number of amplified species and sequences were visualized using box and jitter plots.
The ability to distinguish closely related taxa was assessed by calculating the edit distance
between pairs of amplified species. If the edit distance between the amplicons of two
different species was zero, the sequences were identical, making it impossible to distinguish
between them.

210 2.4 Amplification of tissue-extracted DNA

Based on the *in silico* evaluation of the candidate primer sets (see Results), we selected four
primer sets for empirical validation (Table 1). Mu31F/Dc320R and Dc671F/Dc1015R target
the 12S rRNA gene, Mu2084F/Dc2438R targets the 16S rRNA gene, and
Mu9459F/Mu9822R targets the NAD3 gene. Mu31F/Dc320R was later renamed "μCeta"
(mάɪkroʊ-sɪtə) due to its superior performance in detecting cetacean species (see Results).

The amplification capacity of the four primer sets was initially tested using tissue-
extracted DNA from 21 cetacean species. Tissue (muscle) samples from all cetacean species,
except the Indo-Pacific humpback dolphin (*S. chinensis*), were obtained from the Okinawa
Churashima Foundation, where individual samples are preserved. Tissue samples of *S.*
chinensis were archived by OPCFHK at Ocean Park Hong Kong. Both institutes collect and
preserve stranded cetaceans for scientific research. DNA extraction protocols for the tissue
samples and PCR conditions are described in the Supplementary Methods.

Table 1. Cetacean-specific primers for eDNA metabarcoding tested empirically

Primer set ID	Region	Forward primer sequence	Reverse primer sequence
Mu31F/Dc320R (μCeta[*])	12S rRNA	GACACTGAAAATGTCTAGATGG	**TY**AATCGTATGACCGCGGTG
Dc671F/Dc1015R	12S rRNA	GCTACT**Y**CAGTCTATATACC	CACAC**YTTCCR**GTA**Y**GCTTACC
Mu2084F/Dc2438R	16S rRNA	ATGAA**Y**GGCCACACGAGGGTTTTA	TGTCCTGATCCAACATCGAGG
Mu9459F/Mu9822R	NAD3	CTGACTTCCAATCAGTT**R**GTTTCGG	CATTCTA**RR**CC**Y**T**Y**TTG**R**G

2.5 Water samples from aquarium and Hong Kong coastal regions

To evaluate the performance of the primer sets on water samples, we collected two types of
water samples: Aquarium pool samples and Hong Kong coastal water samples. The first set
of samples was collected from dolphin pools at the Okinawa Churaumi Aquarium using
Sterivex filter cartridges (φ0.45-µm, SVHV010RS; Merck Millipore, Darmstadt, Germany)
and the gravity filtration method (Oka et al., 2022). These water samples are hereafter
referred to as "aquarium pool samples." The aquarium has seven dolphin pools divided into
two compartments, with water circulating within each compartment. The first compartment
includes the "Main pool" (1,200 m^3), "North Lagoon pool" (500 m^3), and "South Lagoon
pool" (500 m^3). The second compartment consists of the "Main Show pool" (1,750 m^3),
"Underwater Show pool" (500 m^3), "Medical Treatment pool" (500 m^3), and "Reproduction
pool" (400 m^3). Detailed information about the dolphin species reared in the pools is
provided in Table S4.
From each pool, one or two 1 L seawater samples were collected (Table S4).
Additionally, 1 L of MilliQ water was filtered as a field negative control. After filtration, 2
mL of RNAlater (Thermo Fisher Scientific, Waltham, MA, USA) was added, and samples
were stored in a cooler box to preserve eDNA before transfer to a –20˚C freezer in the
laboratory. In total, we collected 10 seawater samples from the dolphin pools (nine pool
samples and one field negative control).
Natural seawater samples were collected from Hong Kong coastal waters
(approximately 22˚10′N–22˚30′N, 113˚50′E–114˚25′E). Three seawater samples, hereafter
referred to as "preliminary Hong Kong water samples," were taken from different locations to
assess the amplification of target and non-target species. These samples were filtered using
0.45-µm Sterivex filter cartridges and plastic syringes: 500 ml of surface seawater at the first
location (P1; 22˚22′07″N, 114˚25′55″E) in February 2023 and 1 L of surface and subsurface
seawater at the second location (P2; 22˚20′27″N, 114˚17′42″E) in July 2023. As a field
negative control, 500-ml MilliQ water was filtered at the second location. After filtration,
samples were stored in a cooler box to preserve eDNA before transfer to a –20˚C freezer in
the laboratory.
In addition to these preliminary seawater samples, we collected 500–1,500 mL of
seawater samples in July 2024 from areas where Indo-Pacific humpback dolphins (*S.
chinensis*) were sighted. These samples served as field positive controls, with two samples on
16 July 2024 (22˚11′32″N, 113˚50′57″E). The distance between *S. chinensis* individuals
sighted and the sampling locations was approximately 30, leading us to expect that the
seawater samples contained detectable amounts of *S. chinensis* eDNA. Hereafter, we refer to
these as "natural positive samples." To ensure contamination control, we also collected a
MilliQ water sample as a field negative control. For all field samples, 2 ml of RNAlater was
added and/or the samples were stored in a cooler box to preserve eDNA before transfer to a –
20˚C freezer in the laboratory.

2.6 eDNA extraction from seawater samples

Detailed eDNA extraction protocols are provided in the Supplementary Methods. Briefly,
eDNA was extracted from Sterivex filter cartridges using a modified protocol based on a
previous study (Fukuzawa et al., 2023). We used a DNeasy Blood & Tissue kit (Qiagen,
Hilden, Germany), with the initial lysis step performed using Buffer ATL and Proteinase K.
The extracted DNA was then purified following the manufacturer's instructions and eluted in
100 µL of Buffer AE. To monitor contamination, at least one DNA extraction negative
control (i.e., an empty filter cartridge) was included in each extraction process. Two replicate
eDNA samples from the "Main Pool" and the "Main Show Pool" were pooled to generate a
single eDNA sample for each pool, resulting in a total of seven aquarium pool eDNA
samples. All eluted DNA samples were stored at −20°C until further processing.

2.7 Library preparation and sequencing

Detailed protocols for library preparation and sequencing are provided in the Supplementary
Methods. For both aquarium pool samples and preliminary Hong Kong water samples,
libraries were prepared using four candidate primer sets (Mu31F/Dc320R, Dc671F/Dc1015R,
Mu2084F/Dc2438R, and Mu9459F/Mu9822R).
The first-round PCR (1st PCR) was conducted with a 20-µL reaction volume
containing 10.0 µL of 2 × Platinum SuperFi II PCR Master Mix (Thermo Fisher Scientific,
Waltham, MA, USA), 2.0 µL of each 5.0 µM forward and reverse primer with the Illumina
sequencing primer (Table S5), 2.0 µL of MilliQ water, and 4.0 µL of template DNA. We used
4.0 µL of template DNA to enhance the detection probability of low-concentration eDNA
(Ushio et al., 2022). Two PCR negative controls (using MilliQ water instead of template
DNA) were included to monitor potential contamination during the library preparation
process. After the 1st PCR, the PCR products for each sample were purified using AMPure
XP (Beckman Coulter, Brea, CA, USA). The second-round PCR (2nd PCR) was then
performed to append the Illumina adapters. Following the second PCR, the indexed PCR
products were pooled, and the pooled product was purified again using AMPure XP. The
target-sized DNA of the purified library was excised, and the double-stranded DNA
concentration was adjusted before sequencing on the Illumina NovaSeq platform (Illumina,
San Diego, CA, USA) using a 2 × 250 PE reagent kit.
For natural positive samples, only the Mu31F/Dc320R primer set was used due to its
superior performance (see Results and Discussion). In the library preparation process, we
generally followed the same thermal cycler profile as that used for the aquarium samples.
However, instead of using the 1st PCR primers with the Illumina sequencing primer, we
employed "tagged" 1st PCR primers for more efficient library preparation (referred to as the
"early-pooling protocol" in Ushio et al., 2022). The 1st PCR products were purified using
ExoSAP-IT Express (Thermo Fisher Scientific, Waltham, MA, USA) and pooled for the 2nd
PCR. The remaining processes including sequencing were identical to those described above.
In addition to the four candidate primer sets, we briefly tested the performance of

previously developed primers, MiMammal (Ushio et al., 2017) and Ceto2 (Valsecchi et al.,
2020), using aquarium and natural seawater samples. During library preparation, only the
aquarium and natural seawater samples, along with PCR negative controls, were included.
The library preparation method followed the standard 2-step PCR protocol, and the library
was sequenced on the Illumina iSeq 100 system using iSeq 100 Reagent v2 (2 × 150 bp PE).
For iSeq sequencing, 30% PhiX was spiked in to enhance sequencing quality.
In total, for testing the four primer sets, we had seven aquarium pool samples, one
aquarium field negative control, three Hong Kong water samples, one Hong Kong water
negative control sample, two DNA extraction negative controls, and two PCR negative
controls (a total of 16 samples for each primer set). For testing the Mu31F/Dc320R primer set
using natural positive samples, we had three Hong Kong water samples, two Hong Kong
water negative controls, one DNA extraction negative control, and two PCR negative controls
(a total of eight samples). For the MiMammal and Ceto2 primer sets, we had seven aquarium
pool samples, three Hong Kong water samples, and two PCR negative controls (a total of 12
samples for each primer set).
320

2.8 Sequence data analysis

Detailed procedures for sequence data processing are available on Github and were archived
on Zenodo (see "*Data and code availability statement*"). When the standard two-step PCR
was used for library preparation, demultiplexing was performed using Illumina BaseSpace.
When the early-pooling method was employed, demultiplexing was conducted using a
custom script with Cutadapt v4.7 (Martin, 2011).
The demultiplexed sequences were then quality-filtered using fastp v0.23.4 (Chen et
al., 2018), and primer sequences were trimmed using Cutadapt. The processed sequences
were analyzed using DADA2 (Callahan et al., 2016), an amplicon sequence variant (ASV)
approach implemented in R (R Core Team, 2023). During quality filtering, low-quality and
unexpectedly short reads were removed. Error rates were estimated, and sequences were
dereplicated, error-corrected, and merged to generate an ASV-sample matrix. Chimeric
sequences were subsequently removed using the `removeBimeraDenovo` function in dada2
package. If multiple sequencing runs were conducted for the same library,
`mergeSequenceTables` function was used after DADA2 processing to merge results. For the
aquarium and preliminary Hong Kong water samples, we performed taxonomic identification
using the ASV sequences, as these samples may contain eDNA from closely related species
or hybrid species. For the natural positive samples, ASVs were clustered into operational
taxonomic units (OTUs) at 97% similarity using the DECIPHER package in R (Wright,
2016).
Taxonomic identification of ASVs or OTUs was performed using two methods: the
Query-centric auto-k-nearest-neighbor (QCauto) method (Tanabe & Toju, 2013) implemented
in Claident v0.9.2021.10.22 (https://www.claident.org/) and `assignSpecies()` function
implemented in the dada2 package, supplemented with a manual BLAST search
(https://blast.ncbi.nlm.nih.gov/Blast.cgi). QCauto method is conservative; all nearest
neighbors of a query sequence must share the same taxonomic information for taxa

assignment. For example, if the QCauto method assigns a genus name to an ASV or OTU, all
nearest neighbors surrounding the ASVs or OTUs must have the same genus name. This
method increases the number of ASVs or OTUs without species names but reduces the
likelihood of misassignments in species identification compared to other common methods
(Sato et al., 2018; Q. Wang et al., 2007).
Given that we knew exactly which species were reared in the aquarium pools and that
our target species included closely related species (e.g., *Tursiops truncatus*, *Tursiops*
aduncus, and their hybrid *Tursiops truncatus × aduncus*), we also utilized the
`assignSpecies()` function, using cetacean mitochondrial genomes retrieved from NCBI as a
reference database. If the two approaches assigned different species names to an ASV or
OTU, we adopted the result from the `assignSpecies()` function. In addition, sequences
assigned to the genus "*Stenella*" were manually checked using BLAST, due to the presence
of multiple closely related species. If species-level identification failed, sequences were
labeled as "Unidentified Delphinidae" or "Unidentified Cetacea." If an identified species was
not reared in the aquarium pools, it was classified as "Non-present species."
362

2.9 Statistical analysis

All statistical analyses were performed using R (R Core Team, 2023). For the aquarium
pool and preliminary Hong Kong water samples, ASV matrices were rarefied using the
coverage-based rarefaction (Chao & Jost, 2012) at a minimum coverage of 99.9%. The
rarefied matrices were then converted to relative abundance (*i.e.*, percentage data), and rare
ASVs (< 0.5%) were excluded from the analysis. Samples with fewer than 10 original
sequences were excluded (i.e., most negative controls and some field samples). The relative
abundance data were visualized using the ggplot2 package in R (Wickham, 2009). For the
natural positive samples, only sequence reads were reported, as only one OTU was detected.
372

3. **Results and Discussion**

3.1 *In silico* evaluation: Amplicon lengths

A total of 20 candidate primer sets were identified using the DECIPER package and manual
inspection (Fig. 1 and Table S2). Most primer sets target the mitochondrial 12S rRNA and
16S rRNA gene regions, while two sets target the NAD3 gene region. The performance of the
candidate primers was compared against other universal primers listed in Table S2.
Expected amplicon lengths for Cetacea (74 species), fish (5,411 species), and other
vertebrates (5,368 species) were analyzed using *in silico* PCR, revealing median amplicon
lengths ranging from approximately 100–500 bp (Fig. S1). All primer sets generated
amplicons suitable for sequencing on short-read sequencers (e.g., Illumina). However, certain
primer sets, such as Mu31F/MiMammalR, require 300 bp paired-end sequencing, which
increases sequencing costs.
385

3.2 *In silico* evaluation: primer mismatches in target/non-target taxa

All primer sets exhibited no more than one mismatch with the majority of cetacean DNA, indicating their strong amplification capability for most cetacean sequences (Fig. 2a; see individual primer results in Fig. S2).

The MarVer primes and Riaz12S showed few mismatches with fish and non-fish vertebrate DNA (Fig. 2b,c), consistent with their design as vertebrate-specific primer sets (Riaz et al., 2011; Valsecchi et al., 2020). The MiMammal (mammal-targeted primer set) and Ceto2 (cetacean-targeted primer set) exhibited three or more mismatches with most fish species, but could still amplify a significant number of non-fish vertebrates (Fig. 2b,c).

To further assess primer specificity, we analyzed the number of sequences amplified (Fig. S3) and the mismatch patterns between the forward/reverse primers and different taxa (Fig. S4). These analyses revealed similar patterns to those observed in Fig. 2. Importantly, the candidate primer sets we designed exhibited greater mismatches with human DNA and common mammal DNA (e.g., dogs, cats, and others) as well as with fish, birds, reptiles, and amphibians compared to previously reported primers (Fig. S4). This suggests that our candidate primer sets have a lower likelihood of amplifying non-cetacean vertebrates. In this *in silico* evaluation, we regarded the non-amplification of human DNA as more important than that of other non-target DNA. Among the 20 candidate primer sets, 10 exhibited at least four mismatches with human DNA (i.e., Mu31F/Dc320R, Mu642F/Dc1015R, Mu642F/Mu1021R, Dc671F/Dc1015R, Dc671F/Mu1021R, Mu2084F/Dc2438R, Mu2187F/Mu2563R, Dc2438F/Mu2563R, Mu9459F/Mu9822R, and Mu9459F/Mu9834R).

3.3 *In silico* evaluation: The capacity to differentiate cetacean species

The capacity of each primer set to differentiate cetacean species was evaluated by calculating the edit distance between their amplified products (Fig. 3). As expected, primer sets that amplify shorter regions, such as MarVer2, Ceto2, Riaz12S, Dc494F/Mu643R, and Dc2505/Dc2580, exhibited a reduced capacity to distinguish cetacean species. Specifically, more than 30–100 cetacean amplicon pairs generated by these primers shared identical sequences.

Among the 10 primer sets with a low likelihood of amplifying human DNA, Dc2438F/Mu2563R exhibited a limited capacity to differentiate cetacean species, whereas the other primer sets resulted in fewer cetacean species pairs with identical sequences. From the nine primer sets that demonstrated high specificity for cetacean DNA and a superior capacity to differentiate cetacean species in the *in silico* analysis, we selected several primer sets for empirical validation. Preference was given to primers with fewer degenerate bases, as highly degenerate primers may exhibit lower amplification efficiency. The selected primers included: Mu31F/Dc320R, Dc671F/Dc1015R, and Mu2084F/Dc2438R. Additionally, despite containing seven degenerate bases, we selected Mu9459F/Mu9822R for empirical validation due to its strong cetacean specificity and superior capacity to distinguish cetacean species. In total, we selected four primer sets for empirical validation (Table 1; see Fig. S5 for their primer logos).

Figure 2 | The number of species amplified by *in silico* PCR. The number of cetacean (**a**), fish (**b**), and non-fish vertebrate (**c**) species that can potentially be amplified using the previously reported and candidate primers. Colors represent the total number of mismatches between each primer set (forward and reverse primers) and the template DNA (more detailed results are provided in Figures S2—S4). The dashed horizontal line in each panel indicates the total number of species examined. Primer names shown in bold face are new primers that were empirically validated in this study.

3.4 Amplification of tissue-extracted cetacean DNA

We prepared tissue-extracted DNA from 19 cetacean species. The primer sets

Mu31F/Dc320R and Mu2084F/Dc2438R successfully amplified all tissue-extracted DNA

samples, while Dc671F/Dc1015R failed to amplify one species (*Globicephala*

macrorhynchus), and Mu9459F/Mu9822R did not amplify four cetacean species

(*Globicephala macrorhynchus*, *Kogia sima*, *Kogia breviceps*, and *Mesoplodon ginkgodens*)

(Table 2). These results suggest that Mu9459F/Mu9822R may have lower amplification

efficiency, potentially due to its higher number of degenerate bases. Another possible

explanation for the unsuccessful amplifications is DNA degradation, as the tissue samples of
G. macrorhynchus, *K. sima*, and *M. ginkgodens* had been stored for 14, 12, and 14 years,
respectively. Prolonged storage may have contributed to DNA degradation, reducing
amplification success.
441

Figure 3 | Capacity of previously reported and candidate primers to distinguish among different cetacean species, genera, or families. The capacity to distinguish different cetaceans was assessed by calculating the edit distances between amplicons through *in silico* PCR. The x-axis represents the edit distance, and the y-axis represents the number of species pairs corresponding to each edit distance. Different colors represent different taxonomic resolutions. The number of species pairs in the "0" edit distance category indicates the number of species pairs with identical sequences in the target region. Primer names shown in bold face are new primers that were empirically validated in this study.

Table 2. List of cetacean species for testing cetacean primers using tissue-extracted DNA

Species	Family	Accession No. (if amplified)			
		Mu31F /Dc320R	Dc671F /Dc1015R	Mu2084F /Dc2438R	Mu9459F /Mu9822R
Megaptera novaeangliae	Balaenopteridae	✓	✓	✓	✓
Globicephala macrorhynchus	Delphinidae	✓	No	✓	No

Grampus griseus	Delphinidae	✓	✓	✓	✓
Lagenodelphis hosei	Delphinidae	✓	✓	✓	✓
Peponocephala electra	Delphinidae	✓	✓	✓	✓
Pseudorca crassidens	Delphinidae	✓	✓	✓	✓
Sousa chinensis	Delphinidae	✓	✓	✓	✓
Stenella attenuata	Delphinidae	✓	✓	✓	✓
Stenella coeruleoalba	Delphinidae	✓	✓	✓	✓
Steno bredanensis	Delphinidae	✓	✓	✓	✓
Tursiops aduncus	Delphinidae	✓	✓	✓	✓
Tursiops truncatus	Delphinidae	✓	✓	✓	✓
Kogia breviceps	Kogiidae	✓	✓	✓	No
Kogia sima	Kogiidae	✓	✓	✓	✓
Physeter macrocephalus	Physeteridae	✓	✓	✓	✓
Indopacetus pacificus	Ziphiidae	✓	✓	✓	✓
Mesoplodon densirostris	Ziphiidae	✓	✓	✓	✓
Mesoplodon ginkgodens	Ziphiidae	✓	✓	✓	No
Ziphius cavirostris	Ziphiidae	✓	✓	✓	✓

*Only partial sequence was obtained.

3.5 eDNA metabarcoding for the aquarium pool and preliminary Hong Kong water samples

3.5.1 Sequencing summary

We conducted eDNA metabarcoding on seawater samples collected from the aquarium pools and coastal regions of Hong Kong (Fig. 4). A total of 3,332,270 sequence reads were generated from 64 samples (16 libraries × four candidate primers; Table S6) by three NovaSeq runs, with an overall Q30 score of 92.08%. The four primer sets, Mu31F/Dc320R, Dc671F/Dc1015R, Mu2084F/Dc2438R, and Mu9459F/Mu9822R produced 902,908 reads, 781,005 reads, 782,716 reads, and 865,741 reads, respectively. After quality filtering and DADA2 denoising, 2,572,852 reads remained (77.21% of the total reads). Overall, sequencing quality was sufficiently high.

Negative control samples yielded negligible sequence reads. The highest read count among negative controls was from the filed negative control collected from the Churaumi aquarium pools, but it accounted for at most 0.7% of the average reads from the pool samples. PCR negative controls had only a few reads (<0.1% of average sequence reads). These findings indicate that cross-contamination was negligible, and thus, no post-hoc decontamination analysis (e.g., using *decontam*; Davis et al., 2018) was performed.

Figure 4 | Collection of water samples at dolphin pools in Okinawa Churaumi Aquarium, field site locations in Hong Kong waters, and an Indo-Pacific Humpback Dolphin (*Sousa chinensis*) sighted during the field survey. (**a**) Water sampling at a dolphin pool in Okinawa Churaumi Aquarium for the "aquarium pool" experiment. (**b**) Water sampling locations in Hong Kong waters. (**c**) an Indo-Pacific Humpback Dolphins (*Sousa chinensis*) sighted during the field survey at site P3 (Photo by E. Matrai).

463
464

3.5.2 Detection of cetacean eDNA in the aquarium ponds

First, we tested whether the four primer sets could detect cetacean species reared in the

aquarium pools (Figs. 4a and 5; Table S4). In the Churaumi Aquarium pool test, *Pseudorca*

crassidens and *Tursiops truncatus* (or *T. truncatus × aduncus*) were reared in four pools. All

four primer sets successfully detected *P. crassidens* and *T. truncatus* in these pools (Fig. 5a).

Tursiops aduncus was reared in two pools, where its eDNA was detected using the

Mu31F/Dc320R and Mu9459F/Mu9822R primer sets. The majority of sequences identified

as "Unidentified Delphinidae" in the North Lagoon and the Reproduction Pool, detected by

Dc671F/Dc1015R and Mu2084F/Dc2438R, were likely *T. aduncus*; however, the sequences

could not be assigned to *T. aduncus* due to a lack of resolution in differentiating closely

related species. eDNA from *Steno bredanensis* was detected by all candidate primer sets in

the pool where *S. bredanensis* was reared. eDNA of *Feresa attenuata* was detected by all

primer sets except Mu9459F/Mu9822R in the pool where *F. attenuata* was reared. Many

sequences classified as "Unidentified Delphinidae" in the Reproduction Pool detected by

Mu9459F/Mu9822R were likely *F. attenuata*, yet these sequences could not be assigned to *F.*

attenuata. These results suggest that all reared cetacean species were likely amplified by the

four primer sets. However, only Mu31F/Dc320R was able to assign species names to all
detected sequences.
483

3.5.3 Detection of non-present cetacean eDNA

We detected several species that were not present in the dolphin pools (see the enlarged figure
in Fig. 5b). For instance, Mu31F/Dc320R detected eDNA of *T. aduncus* in the Main pool and
the Main Show pool, and Dc671F/Dc1015R detected eDNA of *F. attenuata* in the
Underwater Show pool. The Main, North Lagoon, and South Lagoon pools belong to the
same compartment, whereas the remaining pools belong to a separate compartment. Since
water circulates within each compartment, cetacean eDNA in may have been transported
between pools. For example, *F. attenuata* eDNA could have moved from Reproduction pool
to the Underwater Show pool, resulting in detection via eDNA metabarcoding. Similar
observations have been reported in other eDNA studies using aquariums (Kelly, Port,
Yamahara, & Crowder, 2014; Miya et al., 2015). In some cases, the source of detection for
non-present (referred to as "non-tank" in the papers) species was unknown; however, this
underscores the sensitivity of eDNA metabarcoding and emphasizes the need for meticulous
water sampling and library preparation to minimize cross-contamination.
The category of "Non-present species" includes *Balaenoptera acutorostrata*, which
was detected only with Dc671F/Dc1015R, with a maximum proportion of 0.76%. This is
negligible and does not impact our conclusions. Rare ASVs removed prior to this analysis
included other non-present species such as *Tursiops australis* (*T. australis* is not recognized
as a species, but here we follow the description of NC_022805 in the NCBI), *Globicephala*
macrorhynchus, *Stenella longirostris*, and *Peponocephala electra*. The sources of these
eDNA detections also remain unknown, and some may be attributed to contamination from
tissue samples.
506

3.5.4 Supplemental eDNA metabarcoding: Detection of cetacean eDNA using MiMammal
and Ceto2

To further assess species detection, we conducted eDNA metabarcoding using the
MiMammal and Ceto2 primer sets (Fig. S6). A total of 2,406,398 sequence reads were
generated from 24 samples (12 libraries using MiMammal and Ceto2 primers) in a single
iSeq run, with an overall Q30 score of 95.80%. MiMammal and Ceto2 generated 1,078,756
and 1,327,642 reads, respectively. After quality filtering and DADA2 denoising, 2,212,982
reads (91.96%) remained for MiMammal and Ceto2. Both primer sets detected several
cetacean species reared in the pools. However, some species could not be assigned species
names, indicating that the capacity of these primer sets had lower species resolution than our
candidate primer sets. Ceto2 exhibited particularly low taxonomic resolution, assigning
species names to only two species, likely due to its short insert length (74 bp), consistent with
our *in silico* analysis (Fig. 3).
520

Figure 5 | Cetacean eDNA metabarcoding results from the aquarium pool samples using the candidate primers. (**a**) Each panel indicates the results for four candidate primer sets. The y-axis represents seven dolphin pools from which seawater samples were collected, with species names and filled squares indicating the dolphin species reared in each pool. The x-axis represents the relative abundance of detected species eDNA. Colors represent different dolphin species or taxonomic categories when species identification is not possible. "Non-present species" denotes species detected that are not reared in the pool. A red tick (✓) indicates successful detection of target species. A black cross mark (×) indicates that, although the target species' eDNA was seemingly amplified, its species name was not correctly assigned to the detected eDNA. Photo credits: *T. truncatus* (Public domain), *T. aduncus* (CC BY-SA 1.0, Aude Steiner), *P. crassidens* (Public domain), *F. attenuata* (Public domain), and *S. bredanensis* (CC BY-SA 3.0, Gustavo Pérez). All photos are sourced from Wikipedia. (**b**) The range from 0 to 0.1 is enlarged for clarity.

3.5.5 Detection of non-target eDNA

We analyzed the detection of non-cetacean species from the aquarium pool samples and preliminary Hong Kong water samples (Fig. 4b) using the four candidate primer sets as well as the MiMammal and Ceto2 primer sets (Fig. S7).

Mu31F/Dc320R, Dc671F/Dc1015R, and Mu9459F/Mu9822R detected almost exclusively cetacean eDNA. Mu31F/Dc320R and Dc671F/Dc1015R detected eDNA of the

Indo-Pacific finless porpoise (*Neophocaena phocaenoides*) in the P1 surface water sample, a
resident cetacean in Hong Kong waters (Jefferson & Moore, 2020). MiMammal primer set
detected finless porpoise eDNA in P1, though at a relatively low proportion (Fig. S7). Despite
our efforts to minimize amplification of non-cetacean eDNA, the Mu2084F/Dc2438R primer
set detected a substantial amount of eDNA from common mammals (e.g., cows, pigs, and
humans).
The MiMammal primer set detected a substantial number of human eDNA reads in all
Hong Kong water samples, as expected. The Ceto2 primer set detected a large proportion of
non-cetacean eDNA, including bacteria, non-cetacean mammals, and other eukaryotes (Fig.
S7). The Ceto2 primer set was originally developed as a cetacean-specific primer set.
However, its empirical validation was not conducted in Valsecchi et al. (2020), as it was not
the primary objective of the study. Our empirical test suggests that Ceto2 may not be suitable
for cetacean-targeting eDNA metabarcoding.
542

3.5.6 Selection of μCeta (Mu31F/Dc320R) as the optimal primer set

Our empirical validation using the aquarium pool and preliminary Hong Kong water samples
demonstrated that Mu31F/Dc320R effectively amplifies and differentiates most cetacean
species (Fig. 5a) while exhibiting a low likelihood of amplifying non-cetacean species.
Consequently, we designated Mu31F-Dc320R as μCeta (pronounced as mάɪkroʊ-sɪtə) and
selected it for analyzing seawater samples collected from Hong Kong coastal areas where the
Indo-Pacific humpback dolphin (*S. chinensis*) was sighted.
550

3.6 Detection of *Sousa chinensis* in Hong Kong waters

Two seawater samples were collected from area where *S. chinensis* were sighted (Fig. 4c),
with the distance between sightings and the sampling location of approximately 30 m. We
then applied the μCeta primer set to these samples. A total of 3,539,281 sequence reads with
an overall Q30 score of 92.48% were generated across three NovaSeq runs (Table S6). After
quality filtering and DADA2 denoising, 3,246,690 reads remained (91.73% of the total
reads). The substantial number of reads originated from seawater samples, while negative
control samples for DNA extraction, PCR, and for field surveys showed negligible read
counts (0 or 1 read per negative control).
Only one OTU was detected from the three seawater samples, and it was assigned to *S.*
chinensis. From the two seawater samples collected on 16 July 2024, 1,799,357 and
1,446,733 reads of *S. chinensis* eDNA were identified, demonstrating that detection of *S.*
chinensis is indeed possible using our eDNA-based approach. Importantly, no non-cetacean
eDNA was detected in any of the three samples, suggesting that μCeta eDNA metabarcoding
is highly sensitive for detecting cetaceans in natural seawater samples. Unfortunately,
quantitative analysis of the detection sensitivity of μCeta primers is beyond the scope of this
study (e.g., how close we should be to detect *S. chinensis* eDNA from a water sample), which
should be addressed in future research.
569

3.7 Future direction of cetacean eDNA metabarcoding

Our *in silico* and empirical tests demonstrated that μCeta primers are well-suited for detecting cetacean eDNA from seawater samples. Although a quantitative evaluation of detection sensitivity is beyond the scope of this study, our results suggest that μCeta primers would detect cetacean eDNA from individuals located within 30–200 m at least in Hong Kong waters. Passive acoustic monitoring has suggested that *S. chinensis* and *N. phocaenoides* inhabiting Hong Kong waters may be more active at night, when visual survey methodology is inappropriate (Wang & Hung, 2022). Consequently, μCeta metabarcoding could serve as an alternative method to study cetacean occurrence during periods of low visibility.

The abundance of cetacean species is critical information for conservation; however, this cannot be directly obtained from traditional eDNA metabarcoding. Species-specific qPCR can quantify cetacean eDNA (Hashimoto et al., 2023). Additionally, eDNA metabarcoding using spike-in DNA (Ushio et al., 2018) can provide an estimate of eDNA concentrations, serving as a rough index of cetacean abundance. Alternatively, eDNA metabarcoding with long-read sequencing could facilitate individual identifications based on intra-specific variation (Urban et al., 2023), which could enable direct estimation of the number of individuals under natural conditions. An intriguing direction for eDNA applications is the integration with DNA methylation analysis, which could enable the estimation of the age of target species (Hirayama et al., 2024; Lu et al., 2023; Mayne et al., 2021; Yagi et al., 2024).

Lastly, given that cetacean eDNA concentrations are expected to be lower than those of other marine vertebrates, such as fish, multiplexing μCeta primers with fish- or vertebrate-specific primers (Miya et al., 2015; Riaz et al., 2011; Valsecchi et al., 2020) would enhance the efficiency of marine vertebrate monitoring using eDNA metabarcoding, while circumventing the adverse effects of amplification of non-target eDNA.

4. Conclusions

In this study, we developed cetacean-specific primers for eDNA metabarcoding and conducted *in silico* analysis to select four promising primer sets. These were empirically validated using aquarium pools and Hong Kong seawater samples, leading to the identification of μCeta—a primer set with high detection sensitivity, strong species differentiation capacity, and minimal non-target amplification.

The μCeta primer successfully detected the Indo-Pacific humpback dolphin (*S. chinensis*), a vulnerable resident cetacean in Hong Kong waters. Additionally, μCeta detected Indo-Pacific finless porpoise (*N. eophocaena phocaenoides*), also classified as 'vulnerable' (*The IUCN Red List of Threatened Species*, 2024), in a seawater sample from the eastern Hong Kong waters. These findings highlight the potential of μCeta eDNA metabarcoding as a non-invasive and highly effective tool for cetacean monitoring sparsely occurring and cryptic cetacean species.

Since μCeta was designed to detect a wide taxonomic range of cetacean species, its application can enhance cetacean monitoring efforts globally, supporting conservation

612 initiatives and ecosystem management across various marine environments. Future studies
613 should validate the effectiveness of μCeta using a broader range of field samples across
614 diverse marine environments to further assess its applicability for global cetacean monitoring.
615
616

617 **Author contributions:** MU conceived and designed research; MU and MM designed
618 primers; MU, SO, SIO, TS, ROK, and MM collected tissues samples; MU, SO, and SIO
619 collected aquarium water samples; MU, ROK, LP, and EM collected natural seawater
620 samples; MU, SO, ROK, and TS performed experiments; MU analyzed data; MU and MM
621 wrote the draft; All authors discussed the results and completed the manuscript.
622

623 **Corresponding author: *Masayuki Ushio**, Department of Ocean Science, Hong Kong
624 University of Science and Technology, Hong Kong SAR, Email: ong8181@gmail.com, Tel:
625 +852-3469-2938 (ORCID iD = 0000-0003-4831-7181)
626

627 **Keywords:** Biodiversity monitoring, Cetaceans, Conservation, Environmental DNA,
628 Metabarcoding, Primer design
629

630 **Data and code availability statements**: All scripts used in the present study are available on
631 Github (https://github.com/ong8181/micro-ceta). Sequence data will be deposited in DDBJ
632 Sequence Read Archives (DRA).
633

Acknowledgements: We thank Suixuan Huang, Ming-Wai Li, and Takamitsu Ohigashi for
their assistance in the field work and Leung Kwan Chak for his assistance in the tissue
sample collection and management. We thank the staff at the Okinawa Churashima
Foundation, Okinawa Churaumi Aquarium, the Agriculture, Fisheries and Conservation
Department (AFCD) of Hong Kong and the Ocean Park Hong Kong for their permission for
using cetacean tissue samples and assistance in collecting them. This research was supported
by The Hong Kong University of Science and Technology Startup Fund to MU, and
GRF16100724 from the Research Grants Council of the Hong Kong SAR, China to MU.
642

643 **Conflict of Interest Statements**: The authors declare no conflict of interest.
644
645
646

References:

Alter, S. E., King, C. D., Chou, E., Chin, S. C., Rekdahl, M., & Rosenbaum, H. C. (2022). Using Environmental DNA to Detect Whales and Dolphins in the New York Bight. *Frontiers in Conservation Science*, *3*. https://www.frontiersin.org/articles/10.3389/fcosc.2022.820377

Baker, C. S., Steel, D., Nieukirk, S., & Klinck, H. (2018). Environmental DNA (eDNA) From the Wake of the Whales: Droplet Digital PCR for Detection and Species Identification. *Frontiers in Marine Science*, *5*. https://doi.org/10.3389/fmars.2018.00133

Bamford, C. C. G., Kelly, N., Dalla Rosa, L., Cade, D. E., Fretwell, P. T., Trathan, P. N., Cubaynes, H. C., Mesquita, A. F. C., Gerrish, L., Friedlaender, A. S., & Jackson, J. A. (2020). A comparison of baleen whale density estimates derived from overlapping satellite imagery and a shipborne survey. *Scientific Reports*, *10*(1), 12985. https://doi.org/10.1038/s41598-020-69887-y

Bista, I., Carvalho, G. R., Walsh, K., Seymour, M., Hajibabaei, M., Lallias, D., Christmas, M., & Creer, S. (2017). Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics of lake ecosystem biodiversity. *Nature Communications*, *8*, 14087. https://doi.org/10.1038/ncomms14087

Bohmann, K., & Lynggaard, C. (2022). Transforming terrestrial biodiversity surveys using airborne eDNA. *Trends in Ecology & Evolution*, *0*(0). https://doi.org/10.1016/j.tree.2022.11.006

Boyd, C., Hobbs, R. C., Punt, A. E., Shelden, K. E. W., Sims, C. L., & Wade, P. R. (2019). Bayesian estimation of group sizes for a coastal cetacean using aerial survey data. *Marine Mammal Science*, *35*(4), 1322–1346. https://doi.org/10.1111/mms.12592

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Meth*, *13*(7), 581–583.

Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., Narwani, A., Mace, G. M., Tilman, D., Wardle, D. A., Kinzig, A. P., Daily, G. C., Loreau, M., Grace, J. B., Larigauderie, A., Srivastava, D. S., & Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature*, *486*(7401), 59–67. https://doi.org/10.1038/nature11148

Chao, A., & Jost, L. (2012). Coverage-based rarefaction and extrapolation: Standardizing samples by completeness rather than size. *Ecology*, *93*(12), 2533–2547. https://doi.org/10.1890/11-1952.1

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*(17), i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Closek, C. J., Santora, J. A., Starks, H. A., Schroeder, I. D., Andruszkiewicz, E. A., Sakuma, K. M., Bograd, S. J., Hazen, E. L., Field, J. C., & Boehm, A. B. (2019). Marine Vertebrate Biodiversity and Distribution Within the Central California Current Using Environmental DNA (eDNA) Metabarcoding and Ecosystem Surveys. *Frontiers in Marine Science*, *6*. https://doi.org/10.3389/fmars.2019.00732

Collins, R. A., Bakker, J., Wangensteen, O. S., Soto, A. Z., Corrigan, L., Sims, D. W., Genner,
M. J., & Mariani, S. (2019). Non-specific amplification compromises environmental
DNA metabarcoding with COI. *Methods in Ecology and Evolution*, *10*(11), 1985–2001.
https://doi.org/10.1111/2041-210X.13276

Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018). Simple
statistical identification and removal of contaminant sequences in marker-gene and
metagenomics data. *Microbiome*, *6*(1), 226. https://doi.org/10.1186/s40168-018-0605-
2

Deng, S., Zhang, X., Liu, M., Lin, B., Zhou, Y., Liu, M., Lin, W., Lin, M., Dong, L., Kang, H.,
Liu, B., Chen, S., Ouyang, M., Jiang, S., Li, J., & Li, S. (2024). Distribution pattern of
cetaceans in the South China Sea based on visual surveys and environmental DNA
metabarcoding. *Authorea*. https://doi.org/10.22541/au.172520933.38586687/v1

Duporge, I., Lee, R., Eweida, A., Mackelworth, P., Ten, S., Alghamdi, A., Alkhamis, R.,
Cochran, J., Lee, S., & Klinck, H. (2024). Passive acoustic monitoring and visual
sighting survey of cetacean occurrence patterns in the Kingdom of Saudi Arabia.
Marine Mammal Science, *40*(3), e13113. https://doi.org/10.1111/mms.13113

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.
Bioinformatics, *26*(19), 2460–2461.

Estes, J. A., Terborgh, J., Brashares, J. S., Power, M. E., Berger, J., Bond, W. J., Carpenter, S.
R., Essington, T. E., Holt, R. D., Jackson, J. B. C., Marquis, R. J., Oksanen, L., Oksanen,
T., Paine, R. T., Pikitch, E. K., Ripple, W. J., Sandin, S. A., Scheffer, M., Schoener, T.
W., … Wardle, D. A. (2011). Trophic Downgrading of Planet Earth. *Science*, *333*(6040),
301–306. https://doi.org/10.1126/science.1205106

Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using
environmental DNA from water samples. *Biology Letters*, *4*(4), 423–425.
https://doi.org/10.1098/rsbl.2008.0118

Foote, A. D., Thomsen, P. F., Sveegaard, S., Wahlberg, M., Kielgast, J., Kyhn, L. A., Salling,
A. B., Galatius, A., Orlando, L., & Gilbert, M. T. P. (2012). Investigating the Potential
Use of Environmental DNA (eDNA) for Genetic Monitoring of Marine Mammals.
PLOS ONE, *7*(8), e41781. https://doi.org/10.1371/journal.pone.0041781

Frasier, K. E., Garrison, L. P., Soldevilla, M. S., Wiggins, S. M., & Hildebrand, J. A. (2021).
Cetacean distribution models based on visual and passive acoustic data. *Scientific
Reports*, *11*(1), 8240. https://doi.org/10.1038/s41598-021-87577-1

Fukuzawa, T., Shirakura, H., Nishizawa, N., Nagata, H., Kameda, Y., & Doi, H. (2023).
Environmental DNA extraction method from water for a high and consistent DNA yield.
Environmental DNA, *5*(4), 627–633. https://doi.org/10.1002/edn3.406

Gillespie, D., Mellinger, D. K., Gordon, J., McLaren, D., Redmond, P., McHugh, R., Trinder,
P., Deng, X., & Thode, A. (2009). PAMGUARD: Semiautomated, open source software
for real-time acoustic detection and localization of cetaceans. *The Journal of the
Acoustical Society of America*, *125*(4_Supplement), 2547.
https://doi.org/10.1121/1.4808713

Hammond, P. S., Lacey, C., Gilles, A., Viquerat, S., Boerjesson, P., Herr, H., Macleod, K.,

Ridoux, V., Santos, M., Scheidat, M., Teilmann, J., Vingada, J., & Oeien, N. (2017).
Estimates of cetacean abundance in European Atlantic waters in summer 2016 from
the SCANS-III aerial and shipboard surveys (p. 40). Wageningen Marine Research.
https://research.wur.nl/en/publications/estimates-of-cetacean-abundance-in-european-
atlantic-waters-in-su

Hashimoto, N., Iwata, T., Kihara, N., Nakamura, K., Sakata, M., & Minamoto, T. (2023).
Detection of environmental DNA of finless porpoise (Neophocaena asiaeorientalis) in
Osaka Bay, Japan. https://doi.org/10.21203/rs.3.rs-3656536/v1

Hirayama, I. T., Wu, L., & Minamoto, T. (2024). Stability of environmental DNA methylation
and its utility in tracing spawning in fish. *Molecular Ecology Resources*, *24*(8), e14011.
https://doi.org/10.1111/1755-0998.14011

Hooper, D. U., Chapin Iii, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H.,
Lodge, D. M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J.,
Vandermeer, J., & Wardle, D. A. (2005). Effects of biodiversity on ecosystem
functioning: A consensus of current knowledge. *Ecological Monographs*, *75*(1), 3–35.

Höschle, C., Cubaynes, H. C., Clarke, P. J., Humphries, G., & Borowicz, A. (2021). The
Potential of Satellite Imagery for Surveying Whales. *Sensors*, *21*(3), Article 3.
https://doi.org/10.3390/s21030963

Janik, V. M. (2014). Cetacean vocal learning and communication. *Current Opinion in*
Neurobiology, *28*, 60–65. https://doi.org/10.1016/j.conb.2014.06.010

Jefferson, T. A., & Moore, J. E. (2020). Abundance and Trends of Indo-Pacific Finless
Porpoises (Neophocaena phocaenoides) in Hong Kong Waters, 1996–2019. *Frontiers*
in Marine Science, *7*. https://www.frontiersin.org/articles/10.3389/fmars.2020.574381

Jefferson, T. A., & Smith, B. D. (2016). Re-assessment of the Conservation Status of the Indo-
Pacific Humpback Dolphin (Sousa chinensis) Using the IUCN Red List Criteria.
Advances in Marine Biology, *73*, 1–26. https://doi.org/10.1016/bs.amb.2015.04.002

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. *Molecular Biology and Evolution*,
30(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kelly, R. P., Port, J. A., Yamahara, K. M., & Crowder, L. B. (2014). Using environmental DNA
to census marine fishes in a large mesocosm. *PloS One*, *9*(1), e86175.
https://doi.org/10.1371/journal.pone.0086175

Kelly, R. P., Port, J. A., Yamahara, K. M., Martone, R. G., Lowell, N., Thomsen, P. F., Mach,
M. E., Bennett, M., Prahler, E., Caldwell, M. R., & Crowder, L. B. (2014). Harnessing
DNA to improve environmental management. *Science*, *344*(6191), 1455–1456.
https://doi.org/10.1126/science.1251156

Kiszka, J. J., Woodstock, M. S., & Heithaus, M. R. (2022). Functional Roles and Ecological
Importance of Small Cetaceans in Aquatic Ecosystems. *Frontiers in Marine Science*, *9*.
https://www.frontiersin.org/articles/10.3389/fmars.2022.803173

Li, X., Wang, H., McCauley, D. J., Altieri, A. H., Silliman, B. R., Lefcheck, J. S., Wu, J., Li,
B., & He, Q. (2023). A wide megafauna gap undermines China's expanding coastal
ecosystem conservation. *Science Advances*, *9*(32), eadg3800.

773   https://doi.org/10.1126/sciadv.adg3800

774   Liu, M., Lin, M., Turvey, S. T., & Li, S. (2017). Fishers' knowledge as an information source
775   to investigate bycatch of marine mammals in the South China Sea. *Animal*
776   *Conservation*, *20*(2), 182–192. https://doi.org/10.1111/acv.12304

777   Lu, A. T., Fei, Z., Haghani, A., Robeck, T. R., Zoller, J. A., Li, C. Z., Lowe, R., Yan, Q., Zhang,
778   J., Vu, H., Ablaeva, J., Acosta-Rodriguez, V. A., Adams, D. M., Almunia, J., Aloysius,
779   A., Ardehali, R., Arneson, A., Baker, C. S., Banks, G., … Horvath, S. (2023). Universal
780   DNA methylation age across mammalian tissues. *Nature Aging*, *3*(9), Article 9.
781   https://doi.org/10.1038/s43587-023-00462-6

782   Lynggaard, C., Bertelsen, M. F., Jensen, C. V., Johnson, M. S., Frøslev, T. G., Olsen, M. T., &
783   Bohmann, K. (2022). Airborne environmental DNA for terrestrial vertebrate
784   community monitoring. *Current Biology*, *0*(0).
785   https://doi.org/10.1016/j.cub.2021.12.014

786   Mace, G. M., Norris, K., & Fitter, A. H. (2012). Biodiversity and ecosystem services: A
787   multilayered relationship. *Trends in Ecology & Evolution*, *27*(1), 19–26.
788   https://doi.org/10.1016/j.tree.2011.08.006

789   Madsen, P. T., Siebert, U., & Elemans, C. P. H. (2023). Toothed whales use distinct vocal
790   registers for echolocation and communication. *Science*, *379*(6635), 928–933.
791   https://doi.org/10.1126/science.adc9570

792   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
793   reads. *EMBnet.Journal*, *17*(1), 10–12. https://doi.org/10.14806/ej.17.1.200

794   Masuda, R., Shiba, M., Yamashita, Y., Ueno, M., Kai, Y., Nakanishi, A., Torikoshi, M., &
795   Tanaka, M. (2010). Fish assemblages associated with three types of artificial reefs:
796   Density of assemblages and possible impacts on adjacent fish abundance. *Fishery*
797   *Bulletin*, *108*(2), 162–173.

798   Mayne, B., Espinoza, T., Roberts, D., Butler, G. L., Brooks, S., Korbie, D., & Jarman, S. (2021).
799   Nonlethal age estimation of three threatened fish species using DNA methylation:
800   Australian lungfish, Murray cod and Mary River cod. *Molecular Ecology Resources*,
801   *21*(7), 2324–2332. https://doi.org/10.1111/1755-0998.13440

802   Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto,
803   S., Yamanaka, H., Araki, H., Kondoh, M., & Iwasaki, W. (2015). MiFish, a set of
804   universal PCR primers for metabarcoding environmental DNA from fishes: Detection
805   of more than 230 subtropical marine species. *Royal Society Open Science*, *2*(7), 150088.
806   https://doi.org/10.1098/rsos.150088

807   Nakagawa, H. (2019). A data set of long-term observation of aquatic vertebrates and associate
808   environmental factors in a temperate stream, the Yura River, Kyoto, Japan. *Ecological*
809   *Research*, *34*(1), 8–8. https://doi.org/10.1111/1440-1703.1059

810   Oka, S.-I., Miya, M., & Sado, T. (2022). Gravity filtration of environmental DNA: A simple,
811   fast, and power-free method. *MethodsX*, *9*, 101838.
812   https://doi.org/10.1016/j.mex.2022.101838

813   Pearson, H. C., Savoca, M. S., Costa, D. P., Lomas, M. W., Molina, R., Pershing, A. J., Smith,
814   C. R., Villaseñor-Derbez, J. C., Wing, S. R., & Roman, J. (2023). Whales in the carbon

cycle: Can recovery remove carbon dioxide? *Trends in Ecology & Evolution*, *38*(3),
238–249. https://doi.org/10.1016/j.tree.2022.10.012

Plön, S., Andra, K., Auditore, L., Gegout, C., Hale, P. J., Hampe, O., Ramilo-Henry, M.,
Burkhardt-Holm, P., Jaigirdar, A. M., Klein, L., Maewashe, M. K., Müssig, J.,
Ramsarup, N., Roussouw, N., Sabin, R., Shongwe, T. C., & Tuddenham, P. (2024).
Marine mammals as indicators of Anthropocene Ocean Health. *Npj Biodiversity*, *3*(1),
1–9. https://doi.org/10.1038/s44185-024-00055-5

Power, H., Takahashi, M., Jarman, S., & Berry, O. (2023). What is environmental DNA?
Environmental DNA, *5*(6), 1743–1758. https://doi.org/10.1002/edn3.497

R Core Team. (2023). *R: A language and environment for statistical computing* [Computer
software]. https://www.R-project.org/

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers:
Inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic
Acids Research*, *39*(21), e145. https://doi.org/10.1093/nar/gkr732

Robinson, C. V., Migneault, A., Dracott, K., & Glover, R. D. (2023). Seas the DNA? Limited
detection of cetaceans by low-volume environmental DNA transect surveys.
Environmental DNA, *5*(6), 1641–1651. https://doi.org/10.1002/edn3.485

Roman, J., & McCarthy, J. J. (2010). The Whale Pump: Marine Mammals Enhance Primary
Productivity in a Coastal Basin. *PLOS ONE*, *5*(10), e13255.
https://doi.org/10.1371/journal.pone.0013255

Rourke, M. L., Fowler, A. M., Hughes, J. M., Broadhurst, M. K., DiBattista, J. D., Fielder, S.,
Wilkes Walburn, J., & Furlan, E. M. (2022). Environmental DNA (eDNA) as a tool for
assessing fish biomass: A review of approaches and future considerations for resource
surveys. *Environmental DNA*, *4*(1), 9–33. https://doi.org/10.1002/edn3.185

Samejima, H., Ong, R., Lagan, P., & Kitayama, K. (2012). Camera-trapping rates of mammals
and birds in a Bornean tropical rainforest under sustainable forest management. *Forest
Ecology and Management*, *270*, 248–256. https://doi.org/10.1016/j.foreco.2012.01.013

Sato, Y., Miya, M., Fukunaga, T., Sado, T., & Iwasaki, W. (2018). MitoFish and MiFish
Pipeline: A Mitochondrial Genome Database of Fish with an Analysis Pipeline for
Environmental DNA Metabarcoding. *Molecular Biology and Evolution*, *35*(6), 1553–
1555. https://doi.org/10.1093/molbev/msy074

Sigsgaard, E. E., Nielsen, I. B., Bach, S. S., Lorenzen, E. D., Robinson, D. P., Knudsen, S. W.,
Pedersen, M. W., Jaidah, M. A., Orlando, L., Willerslev, E., Møller, P. R., & Thomsen,
P. F. (2017). Population characteristics of a large whale shark aggregation inferred from
seawater environmental DNA. *Nature Ecology & Evolution*, *1*(1), Article 1.
https://doi.org/10.1038/s41559-016-0004

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For Biodiversity
Research and Monitoring*. OUP Oxford.

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-
generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*,
21(8), 2045–2050. https://doi.org/10.1111/j.1365-294X.2012.05470.x

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of fish

biomass using environmental DNA. *PLOS ONE*, *7*(4), e35868.
https://doi.org/10.1371/journal.pone.0035868

Tanabe, A. S., & Toju, H. (2013). Two new computational methods for universal DNA
barcoding: A benchmark using barcode sequences of bacteria, archaea, animals, fungi,
and land plants. *PloS One*, *8*(10), e76910.
https://doi.org/10.1371/journal.pone.0076910

The IUCN Red List of Threatened Species. (2024, February). IUCN Red List of Threatened
Species. https://www.iucnredlist.org/en, accessed on 11 March 2025

Thomas, L., Buckland, S. T., Rexstad, E. A., Laake, J. L., Strindberg, S., Hedley, S. L., Bishop,
J. R. B., Marques, T. A., & Burnham, K. P. (2010). Distance software: Design and
analysis of distance sampling surveys for estimating population size. *Journal of Applied*
Ecology, *47*(1), 5–14. https://doi.org/10.1111/j.1365-2664.2009.01737.x

Tsuji, S., Shibata, N., Inui, R., Nakao, R., Akamatsu, Y., & Watanabe, K. (2023). Environmental
DNA phylogeography: Successful reconstruction of phylogeographic patterns of
multiple fish species from cups of water. *Molecular Ecology Resources*, *23*(5), 1050–
1065. https://doi.org/10.1111/1755-0998.13772

Urban, L., Miller, A. K., Eason, D., Vercoe, D., Shaffer, M., Wilkinson, S. P., Jeunen, G.-J.,
Gemmell, N. J., & Digby, A. (2023). Non-invasive real-time genomic monitoring of the
critically endangered kākāpō. *eLife*, *12*, RP84553. https://doi.org/10.7554/eLife.84553

Urian, K., Gorgone, A., Read, A., Balmer, B., Wells, R. S., Berggren, P., Durban, J., Eguchi, T.,
Rayment, W., & Hammond, P. S. (2015). Recommendations for photo-identification
methods used in capture-recapture models with cetaceans. *Marine Mammal Science*,
31(1), 298–321. https://doi.org/10.1111/mms.12141

Ushio, M., Fukuda, H., Inoue, T., Makoto, K., Kishida, O., Sato, K., Murata, K., Nikaido, M.,
Sado, T., Sato, Y., Takeshita, M., Iwasaki, W., Yamanaka, H., Kondoh, M., & Miya, M.
(2017). Environmental DNA enables detection of terrestrial mammals from forest pond
water. *Molecular Ecology Resources*. https://doi.org/10.1111/1755-0998.12690

Ushio, M., Furukawa, S., Murakami, H., Masuda, R., & Nagano, A. J. (2022). An efficient
early-pooling protocol for environmental DNA metabarcoding. *Environmental DNA*,
4(6), 1212–1228. https://doi.org/10.1002/edn3.337

Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H.,
Minamoto, T., & Kondoh, M. (2018). Quantitative monitoring of multispecies fish
environmental DNA using high-throughput sequencing. *Metabarcoding and*
Metagenomics, *2*, e23297. https://doi.org/10.3897/mbmg.2.23297

Valsecchi, E., Bylemans, J., Goodman, S. J., Lombardi, R., Carr, I., Castellano, L., Galimberti,
A., & Galli, P. (2020). Novel universal primers for metabarcoding environmental DNA
surveys of marine mammals and other marine vertebrates. *Environmental DNA*, *2*(4),
e72. https://doi.org/10.1002/edn3.72

van Weelden, C., Towers, J. R., & Bosker, T. (2021). Impacts of climate change on cetacean
distribution, habitat and migration. *Climate Change Ecology*, *1*, 100009.
https://doi.org/10.1016/j.ecochg.2021.100009

Wang, J. Y., & Hung, S. K. (2022). *Passive acoustic monitoring of marine mammals in Hong*

Kong waters (2021–2022): Final report (An Unpublished Report Submitted to the
Agriculture, Fisheries and Conservation Department of Hong Kong SAR Government,
p. 131).

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naïve Bayesian Classifier for
Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Applied and*
Environmental Microbiology, *73*(16), 5261–5267.
https://doi.org/10.1128/AEM.00062-07

Weir, C. R., & Pierce, G. J. (2013). A review of the human activities impacting cetaceans in the
eastern tropical Atlantic. *Mammal Review*, *43*(4), 258–274.
https://doi.org/10.1111/j.1365-2907.2012.00222.x

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
http://ggplot2.org

Wright, E. S. (2016). Using DECIPHER v2.0 to analyze big biological sequence data in R. *The*
R Journal, *8*(1), 352–359.

Yagi, G., Qi, H., Arai, K., Kita, Y. F., Kogi, K., Morisaka, T., Yoshioka, M., & Inoue-Murayama,
M. (2024). Non-invasive age estimation based on faecal DNA using methylation-
sensitive high-resolution melting for Indo-Pacific bottlenose dolphins. *Molecular*
Ecology Resources, *24*(2), e13906. https://doi.org/10.1111/1755-0998.13906

Yamamoto, S., Masuda, R., Sato, Y., Sado, T., Araki, H., Kondoh, M., Minamoto, T., & Miya,
M. (2017). Environmental DNA metabarcoding reveals local fish communities in a
species-rich coastal sea. *Scientific Reports*, *7*, 40368. https://doi.org/10.1038/srep40368

Zhang, S., Cao, Y., Chen, B., Jiang, P., Fang, L., Li, H., Chen, Z., Xu, S., & Li, M. (2023).
Assessing the potential use of environmental DNA for multifaceted genetic monitoring
of cetaceans: Example of a wandering whale in a highly disturbed bay area. *Ecological*
Indicators, *148*, 110125. https://doi.org/10.1016/j.ecolind.2023.110125

924
925