Title

Annotating Hypothetical Proteins in *Escherichia coli* K-12

Author information

Sagarika Chakraborty[1], Zachary Ardern[1,2], Habibu Aliyu[1] and Anne-Kristin Kaster[1,3]*

[1] Institute for Biological Interfaces 5 (IBG-5), Biotechnology and Microbial Genetics, Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

[2] Wellcome Trust Sanger Institute, Hinxton, Saffron Walden CB10 1RQ, United Kingdom

[3] Institute for Applied Biosciences (IAB), Karlsruhe Institute of Technology (KIT), Kaiserstraße 12, 76131 Karlsruhe, Germany

* To whom correspondence should be addressed: Email: kaster@kit.edu

Graphical Abstract

Abstract

Omics technologies have led to the discovery of a vast number of proteins that are expressed

but have no functional annotation - so called hypothetical proteins (HPs). Even in the best-

studied model organism *Escherichia coli* K-12, over 2% of the proteome remains

uncharacterized. This knowledge gap becomes even worse when looking at microbial dark

matter. However, knowing the functions of proteins is crucial for elucidating cellular and

metabolic processes and harnessing biotechnological potentials. Here, we employed machine

learning to decipher the transcriptional regulatory network of *E. coli* K-12, as well as other *in*

silico tools to assign functions to uncharacterized HPs. We further provide experimental

validation of *in silico* predicted functions for three HP-encoding genes (*yhdN*, *yeaC* and *ydgH*)

as proof of concept, by analyzing growth patterns of deletion mutants compared to the wild

type, as well as their transcriptional responses to specific conditions. This study demonstrates

that the use of Big Omics Data in combination with Artificial Intelligence and experimental

controls is a powerful approach to illuminate functional dark matter.

Keywords: Artificial Intelligence / Big Omics Data / Functional Annotation of Proteins

/ Functional Dark Matter / Independent Component Analysis (ICA)

Introduction

Due to the advents in Next Generation Sequencing technologies, the volume of omics data has massively increased over the past years. As of today, gold-standard knowledgebases like the National Center for Biotechnology Information (NCBI)'s Reference Sequences Database (RefSeq) and the Universal Protein Resource Database (UniProt) [1] provide freely accessible sequences and information for over 163 million unique prokaryotic proteins **(Figure 1)**. This rapid accumulation of sequencing data has long surpassed the rate of possible experimental characterization *in vivo* and *in vitro*. Hence, researchers often rely on functional analysis of proteins from model organisms. These findings are then extrapolated *in silico* to closely related sequence homologues. Consequently, the definition of cut-off values plays a crucial role in protein annotation. To manage this challenge, automated annotation pipelines, such as InterPro have become a valuable tool [2]. Here, analysis of protein sequences is provided by classifying them into families and integrating predictive models from multiple protein databases such as e.g. Pfam [3], ProSite [4], SMART [5] and/or CDD [6] to infer functions for experimentally uncharacterized proteins. However, only 83% of all UniProt sequences have so far been functionally annotated by InterPro [7] The remaining 17% of proteins are designated with terms such as 'putative uncharacterized', 'uncharacterized', 'unknown protein families (UPFs)' or proteins bearing 'domains of unknown functions' (DUFs). These proteins are summarized by the term hypothetical proteins (HPs), since they are expressed but could so far not be functionally characterized by classical *in vivo*, *in vitro*, and/or *in silico* methods (8,9). In addition, there is also no information on these proteins available in widely-used knowledge-databases such as BioCyc [10], RegulonDB [11], and EggNOG [12] **(Figure 1).**

While there have been notable advancements in methodologies for protein function prediction in the recent past [13–16] several challenges persist [17,18]: These include different functionalities across homologous proteins, proteins performing different functions in distinct cellular locations, proteins with multiple three-dimensional structures, the lack of conserved proteins across different species to identify shared functional relationships and a heavy

reliance on pre-existing data for inferential annotations [19]. Even in the best studied model

organisms *Escherichia coli* K-12, 31.8% of the genome comprises of protein encoding genes

that lack experimental validation, commonly referred to as 'Putative HPs' [8], and 2.1% that

even lack sequence homologies (this study) (**Figure 1**). The problem of annotating HPs

becomes even larger when looking at microbial dark matter(9, 20).In uncultivated

microorganisms, the proportions of genes with unknown functions can comprise up to 60% in

bacterial and 80% in archaeal genomes [21]. However, elucidating the function of these

proteins is essential for understanding cellular processes, metabolism and evolution, as well

as for harnessing their biotechnological potential [22].

Transcriptional profiling has helped to gain insights on protein functions by monitoring gene

expressions and clustering genes based on their responses to varying environmental

conditions and stimuli [23]. Here, the entire set of RNA transcripts produced by a genome

under specific conditions provides a dynamic representation of the organisms' operational

state. However, standard bioinformatics-based methods for transcriptomic data analysis are

low-throughput and require quite extensive computing power and time due to the high degrees

of complexity and heterogeneity of the data [24]. Today, artificial intelligence (AI)-based

methodologies have made significant advancements over traditional bioinformatics

approaches, offering the capacity to process and analyze data at a scale and speed that were

previously unattainable [24, 25]. They are particularly suited for deciphering complex

interactions among genes and transcription factors or proteins that regulate them, as well as

revealing the regulation of gene expression in response to diverse environmental conditions

[27]. Among various techniques for analysis, module-detection methods have proven to

efficiently predict the functions of co-regulated groups of genes from large gene expression

datasets [28]. Independent Component Analysis (ICA), introduced by Comon in 1994 [29],

employs an unsupervised machine learning (ML) approach, categorizing the input data without

any prior information. This technique is especially relevant in the context of transcriptomics,

where the goal is to understand the complex interplay of gene expression signals and to

decipher co-regulated gene sets. ICA therefore allows to reveal the regulatory patterns and

activity levels of genes governed by specific regulators (i.e. transcription factors controlling the

expression of genes) across diverse experimental conditions, so called transcriptional

regulatory networks (TRNs), thereby disclosing the organizational and functional architecture

of genes. Expanding upon this, McConn et al. developed a variant of ICA, termed OptICA,

which can be applied to large TRNs, analyzing interactions between transcription factors and

their target genes. Here, discrete groups of genes are clustered into robust independent

components by avoiding over-clustering of datasets (leading to loss of biological information)

or under-clustering (leading to too few clusters and a loss of output resolution), and therefore

providing an optimal representation of the organism's underlying TRN [27]. This results in

revealing independently regulated gene groups, so called iModulon from transcriptomics

datasets [29, 30]. iModulons are therefore the data-driven analogs of so-called regulons - sets

of genes governed by regulators [30]. The OptICA algorithm along with high-quality RNA-seq

data has recently been used to decipher the TRN of *Bacillus subtilis* [32] and *E. coli* [27] to

uncover the detailed responses to environmental conditions and genetic perturbations, e.g.

deletion mutants. However, these studies focused on the identification of regulons that were

not experimentally described before rather than the functional characterization of HPs.

In order to decipher the function of HP-encoding genes in the *E. coli* K-12 proteome, we

adapted the workflow originally designed for *Bacillus subtilis* [32] and analyzed the gene

expression data of the *E. coli* K-12 sub-strains MG1655 and BW25113. Functional

characterization of HP-encoding genes was inferred by identifying co-regulated genes under

the influence of a common regulator from publicly available transcriptomic datasets. This

characterization was then further validated by annotating these genes using specific Gene

Ontology (GO) categories through the PANTHER tool (Protein ANalysis THrough Evolutionary

Relationships) [33]. Additionally, metadata conditions for the transcriptomic datasets were

extracted. Furthermore, other *in silico* methods were employed to determine the sub-cellular

localization [34], protein-protein interactions (PPIs) [35], genomic context [36] and sequence-

based remote homologies [37]. Structural homology information was obtained by using the

AlphaFold Protein Structure Database (AFDB) clusters webserver, which is based on a deep

learning (DL) method [38].

Since AI-predicted functions are usually not experimentally verified, raising the possibility of

untrue annotations [9], we also provide an experimental validation of three candidate HP-

encoding genes (*yhdN*, *yeaC* and *ydgH*). The growth curves and transcriptional responses of

E. coli K-12 deletion mutants in comparison to the wild-type strain under specific conditions

were analyzed, showing that the *in silico* predictions were indeed true. Hence, the here

presented pipeline not only facilitates the discovery of previously unidentified regulators, but

also sheds light on our understanding of the function of HPs **(Figure 1).**

Figure 1. Total number of sequences for all unique prokaryotic and *Escherichia coli* proteins deposited in the National Center for Biotechnology Information (NCBI) as of April 2024 and methodological set-up of this study. Of the 4403 *E. coli* K-12 protein encoding genes, 1403 (31.8%) encode for unique proteins with functions predicted only *in silico* based on homologous sequences but lacking *in vivo* or *in vitro* experimental evidence (termed "putative hypothetical proteins"). 95 protein encoding genes (2.1%) of *E. coli* K-12 are completely uncharacterized with no sequence homologues according to the four knowledge databases - EcoCyc [39], RegulonDB [11], EggNOG [12] and UniProt (1) (termed "hypothetical proteins"). Transcriptomic datasets from NCBI were filtered and processed using the OptICA approach to generate iModulons [32]. Metadata information was curated in parallel using manual or semi-automated approaches(39–41).Bioinformatics, machine learning and deep learning tools along with the presence of relevant metadata then resulted in potential functions for HP candidates for *in vitro* testing. HPs, hypothetical proteins; ICA, independent component analysis; ML, machine learning.

Results and Discussion

Identification of uncharacterized HPs in *E. coli*

In 2019, Ghatak et al. had identified HP-encoding genes in *E. coli* K-12, however only by employing keywords such as 'possibly', 'predicted' or 'hypothetical' from the EcoCyc database, or annotation scores of two or below in UniProt [8]. This approach did, however, not establish a stringent threshold for accurate identification of completely uncharacterized HP-encoding genes. Hence, a more rigorous approach was used in this study, which included additional keywords for HPs (such as 'Uncharacterized', 'Putative uncharacterized', 'DUF' etc.) and information on the absence of any functional characterization using multiple gold-standard databases (EcoCyc, UniProt , RegulonDB and EggNOG). 158 of the 1600 HP-encoding genes previously listed by Ghatak et al. have now been functionally characterized with experimental evidence in the EcoCyc database. Notably, the HP-encoding gene *ydfX*, curated as one of the HPs in this study, was missing from Ghatak et al.'s list since it was previously categorized as a pseudogene/phantom gene. Its exclusion, however, could not be verified in the current EcoCyc database. We identified 1403 of 4403 genes (31.8%) of the *E. coli* K-12 genome as 'putative' HPs, implying that their functions were inferred based on sequence homologies in the gold-standard databases, but lacking an experimental evidence for function. A recent study using the EggNOG tool also identified about 30% in a different *E.* strain (O157:H7 strain Sakai) as putative HPs [18].

95 HP-encoding genes (2.1%) could be identified with no sequence homologies in the *E. coli* K-12 genome. These could be further sub-categorized in 'uncharacterized proteins' (53), 'putative uncharacterized' (5), '(DUF)' domain containing proteins (21) and 'UPF' (HPs with unknown protein families, 3). 13 HPs had the characteristic 'Protein Y…' naming scheme. The distribution of those 95 HP-encoding genes within the genome can be found in the Supplementary (*Supplementary Figure 1)*.

iModulon generation and analysis

In order to investigate the functions for the 95 uncharacterized HP-encoding genes, publicly

available RNAseq datasets from *E. coli* K-12 substrain MG1655 were downloaded from NCBI

and quality filtered. 779 high-quality datasets from the MG1655 substrain were then used as

input for the OptICA algorithm [27]. For the substrain BW25113, a separate OptICA analysis

was not conducted due to the very small size of the dataset. The iModulons were identified by

utilizing the PyModulon package, specifically employing the *Inferring iModulon Activities*

function [32] for the BW25133 substrain. In total, 131 iModulons were obtained for the entire

E. coli K-12 transcriptome. Three single gene iModulons were discarded from the analyses,

since they were considered a result of artificial knockouts or overexpression of single genes

in the dataset. Therefore, a total of 128 iModulons were selected for further analysis.

Functional information on the HP-encoding genes was obtained by mapping the iModulons to

existing experimentally derived regulons (groups of genes regulated by the same transcription

factor), using data from the EcoCyc and RegulonDB databases. Additionally, Gene Ontology

(GO)-based functional annotations for member genes of an iModulon (co-regulated gene sets)

were identified using the PANTHER tool (Protein ANalysis THrough Evolutionary

Relationships) [33]. A total of 51 HP-encoding genes were clustered into iModulons using

OptICA. Of these, 46 genes were assigned putative functions based on information from

regulator proteins and/or Gene Ontology (GO) categories. (**Figure 2**). Two genes, *yfeS* and

yffL had information on their regulators (RpoE and NarL, respectively) but no assigned GO

categories, while five genes (*ybaA, yfaP, yjgZ, ymgI and ymgJ*) (white) could not be associated

with a regulator or a GO category. 44 of the HP-encoding genes could not be clustered into

distinct iModulons (grey) (**Figure. 2**).

Figure 2. Regulator and functional classification for 95 HP-encoding genes in *Escherichia coli*
K-12. 44 HP-encoding genes could not be clustered by the OptICA method [27] (grey), since they did
not co-regulate with any other genes. 22 genes were characterized based on information on their
regulators as obtained from RegulonDB and/or EcoCyc, as well as by Gene Ontology (GO) categories
based on the co-regulating genes using the PANTHER tool (Protein ANalysis THrough Evolutionary
Relationships) (orange) [33]. 29 genes co-regulated with other genes, but no information on their
regulators could be obtained (blue). 24 out of these 29 genes could be assigned putative functions
based on GO annotations of co-regulating genes derived from the PANTHER tool. The striped regions
denote genes where the regulator-associated function from EcoCyc/RegulonDB did not match a GO-
derived functional category obtained from PANTHER. Highlighted in bold are the three HP-encoding
genes which were selected for *in vitro* testing.

208
In addition to the 95 HP-encoding genes, all other *E. coli K-12* genes were also analysed using

this method. 2630 of the 2905 (91%) genes which have experimental evidence of function

could also be characterized into iModulons. 1697 (64%) co-regulated with other genes, i.e.

they were part of characterized iModulons with information on their regulator on RegulonDB

and/or EcoCyc, while 933 of the genes were part of uncharacterized iModulons. For genes

with known regulators, only 121 (5%) had either a non-matching functional category when

compared with the PANTHER categories or a non-significant (*p*>0.05) functional category,

and 95% had a matching PANTHER GO category. For genes which were part of

uncharacterized iModulons, with no regulator information, 546 (58%) could be functionally

classified using PANTHER. 1061 of the 1403 putative HP-encoding genes (75%), i.e. genes

with functional characterisation based on sequence homology information, could be

characterized into iModulons. 342 genes (25%) did not co-regulate with other genes. For

genes that had co-regulation, 539 (51%) had information on their regulator on RegulonDB

and/or EcoCyc. The other 522 (49%) were part of uncharacterized iModulons. For genes with

known regulators, only 20 genes (4%) had either a non-matching function when compared

with the PANTHER categories or a non-significant (p>0.05) functional category in PANTHER.

For genes with no information on their regulator, 366 (67.9%), could however be characterized

by PANTHER. Hence, about 95% of the iModulon functional categories matched the

PANTHER GO categories, demonstrating the robustness of the method. Moreover, the

inclusion of experimentally characterized genes within uncharacterized iModulons shows

potential for the discovery of new regulons or co-regulated genes.

Functional analysis of HP-encoding genes using *in silico* tools

All 95 HP-encoding genes were analysed based on multiple criteria including the generated

iModulons from the OptICA algorithm, functional information based on the regulators of the

co-regulated gene sets obtained *via* RegulonDB/EcoCyc and their GO categories from

PANTHER. Information on metadata conditions pertaining to the highest level of gene

expression relative to the reference gene *frr* (encoding the ribosome recycling factor) [42] and

annotation of the most significant co-expressed gene were also extracted. Additionally, the

sub-cellular localization, protein-protein interactions, genomic context, remote sequence

homology as well as structural homology aided in forming hypothesises for assigning functions

to the respective HPs. 33 of the protein encoding genes (highlighted in green) could be

assigned functions with a high confidence. 28 of the genes were assigned functions based on

lower confidence, since only information from a maximum of two sources could be well-

correlated (highlighted in yellow). A clear inference could not be made for 34 of the genes

(highlighted in red). Information retrieved for each gene and functional inference (if applicable)

is detailed in **Table1**. A pictorial overview of Table 1 based on the functional annotations for

the 95 HPs can be found in *Supplementary Figure 2*.

Table1. A comprehensive functional interference of HPs based on different *in silico* tools. Annotations are based on regulator information from RegulonDB/EcoCyc and/or GO category based on co-regulated genes obtained from PANTHER) [33](as shown in Fig. 2). The highest gene expression condition, obtained from the Tjaden dataset [43], with metadata conditions categorized by the type of medium, oxygen availability (normal O_2 conditions unless stated otherwise), temperature etc. in comparison to the reference gene 'frr' are described. Functional annotations of co-expressed genes were retrieved from EcoCyc [36]. 'SCL' or sub-cellular localization inferred from the BUSCA tool [34] are abbreviated as either 'NUC' for nuclear 'CYT' for cytoplasmic, 'ECS' for extra cellular space or 'PM' for plasma membrane. Functional information from protein-protein interaction (PPI) analyses was obtained from STRINGdb [35]. Annotations for neighbouring upstream (←) and downstream (→) genes in context of the HP-encoding gene were retrieved from the EcoCyc database. Additionally, remote homology was assessed using the Uniref30 database using the Hhblits tool [37], and structural homology was inferred from the AlphaFold Protein Structure Database [38]. A functional inference can be made for genes supported by multiple lines of evidence when similar functional information was provided by at least two sources. Proteins with well correlated information from at least three sources are highlighted in green (with higher confidence) and proteins with information that can be correlated using at least two sources are highlighted in yellow (with lower confidence). Protein encoding genes highlighted in grey were selected for *in vitro* testing. An unclear inference suggests that, based on the different knowledge bases, the extracted information did not provide enough confidence or could not be well correlated to assign a function (red).

Gene	Function based on regulator and/or co-regulated genes	Condition with highest gene expression (growth medium)	Annotation of most significant co-expressed gene	SCL	PPI: function of partner protein	Annotation of neighbouring genes			Remote homology	Structural homology	Overall functional inference
1. yabP	Anaerobic respiration	M9, 37°C; 100 µg/ml Ampicillin (inhibits cell wall synthesis) [44]	Growth related protein (yabQ)	CYT	No interaction partner	djlA ← / Co-chaperone protein	**yabP** →	yabQ / Growth related protein	Putative Undecaprenyl pyrophosphate phosphatase (cell wall biosynthesis) [45]	Type III secretion system (bacterial pathogenecity) [46]	Cell wall biosynthesis
2. yagM	Cell adhesion	LB, 37°C; 20 µg/ml Bicyclomycin (produces ROS, blocks cell division) [47]	Transcriptional regulator (fimZ) (involved in fimbriae production) [48]	CYT	DNA-binding protein (YagL)	yagL ← / resolvase like prophage	**yagM** ←	ykgV ← / Growth related protein	Unknown function	Unknown function	Cell adhesion under antibiotic stress
3. yagK	Glycine betaine bio-synthesis from choline (osmoprotectant) [49]	Na-Phosphate buffer, 37°C; Δthil (thil is responsive to UV) [50]	Putative fimbrial usher protein (yhcD)	NUC	Putative ligase (GtrS)	yagJ → / Response to X-radiation	**yagK** ←	yagL ← / resolvase like prophage	CP4-57 prophage protein (cell growth/biofilm formation/ motility/ transport) [51]	Unknown function	Protection against radiation-induced stress

Gene	Co-regulation	Conditions	Function	Loc	Function	Gene (left)	Gene (center)	Gene (right)			Category
4. yahE	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (cause cellular stress) [52]	Putative acyl-CoA synthetase (yahF) (energy metabolism)	CYT	Putative NAD(P)-binding succinyl-CoA synthase (SucD) (energy metabolism)	yahD → Response to UV	**yahE** →	yahF → Putative acyl-CoA synthetase	Unknown function	Unknown function	Energy metabolism under cellular stress
5. yahL	Cell adhesion	LB, 37°C	Membrane protein (yahC)	NUC	Unknown function (YahE)	yahK → Aldehyde reductase (involved in biofilm formation) [53]	**yahL** →	xtpA → Putative transcr. regulator	Unknown function	Unknown function	Cell adhesion
6. yahM	Glycine betaine bio-synthesis from choline (osmoprotection) [49]	M9, 37°C; 25 µg/ml Kanamycin	Putative amino acid exporter (yahN) (Efllux of toxic sunstances) [54]	CYT	Putative replicase (YidX)	xtpA → Putative transcr. regulator (confers antibiotic resistance) [55]	**yahM** →	yahN ← Putative amino acid exporter	Unknown function	Unknown function	Protection against stress
7. ybaA	Co-regulated (unknown regulator or function)	LB, 37°C	Putative oxido-reductase (yohF)	CYT	Double strand break repair protein (YegP)	ffs → Signal recognition particle	**ybaA** →	pdeB ← Cyclic di-GMP phospho-diesterase	RNA signal recognition protein [56]	Unknown function	Oxidoreduction/ DNA damage repair
8. ybaP	Co-regulated (unknown regulator or function)	LB, 37°C	DNA binding transcriptional activator (AaeR) (role in biofilm formation) [57]	CYT	Cysteine-tRNA deacylase (YbaK)	chiX → Small regulatory RNA (role in biofilm formation) [58]	**ybaP** ←	ybaQ → DNA binding transcr. regulator (regulates biofilm based response) [59]	Conjugal transfer protein	Unknown function	Biofilm formation
9. ybdJ	Not co-regulated	LB, 37°C	Putative transporter (yeeA) (aromatic acid exporter) [60]	PM	DUF419 family protein (YbdF)	ybdF ← Defense mechanism	**ybdJ** ←	ybdK ← Amino acid bio-synthesis	Unknown function	Unknown function	Stress response

10. ybeR	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (cause cellular stress) [52]	Putative oxamate carbomyl-transferase (*fdrA*)	CYT	Putative co-chaperone (DjlB) (heat stress response)	*ybeQ* ← Oxidative stress response	**ybeR** →	*djlB* → J-domain type protein	Unknown function	Unknown function	Multi-stress response
11. yceQ	Cell death	LB, 37°C	DNA binding transcriptional repressor (*acrR*) – involved in efflux of toxic substances [61]	CYT	Endoribonuclease (RNE) – adaptation to different growth conditions [62]	*Rne* ← Ribosomal nuclease	**yceQ**	*rluC* → 23S rRNA pseudo-uridine synthase	Unknown function	Unknown function	Cell death prevention
12. ycgX	Not co-regulated	LB, no O₂, 37°C	Membrane protein (*yigF*)	CYT	Repressor of blue light-responsive genes (BluR)	*ymgL* ← Consti-tutively expressed protein in rich medium	**ycgX** ←	*bluR* ← DNA binding transcr. regulator	Prophage protein	Unknown function	Unclear
13. ycgY	Stress response in stationary phase	LB, no O₂, 37°C	Outer membrane protein (*yaiO*)	CYT	Uncharacterized protein (YncH)	*ymgE* → DNA Recomb-ination	**ycgY** →	*treA* ← Trehalase protein (nutrient stress in stationary phase) [63]	Unknown function	Unknown function	Growth-associated response during stationary phase
14. ychS	Not co-regulated	LB, 37°C; 1 ug/mL of Ciprofloxacin+ 500 µM of ET2DA (cause oxidative/ cellular stress) [64]	Putative amino acid de-hydrogenase (*ykfG*) – role in oxidative stress [65]	CYT	Not found	*narI* → Nitrate reductase (oxido-reductase activity) (UniProt)	**ychS** →	*tpr* ← Protamine like protein	Beta-galactosidase	Protein YchS (Unknown function)	Oxidative stress response

Gene	Co-regulation	Condition	Protein	Loc.	Protein	Gene context	Left function	Right function	Conclusion
15. yciX	Not co-regulated	M9, 37°C; *yqjD* over-expression (inhibits cell growth) [66]	Membrane-associated regulatory protein (*ymiA)* (in exponential or tationary phase) [67]	CYT	Membrane-associated regulatory protein (YmiA)	*ymiA* → **yciX** → *ymiC* → Membrane-associated regulatory protein — Stationary phase adaptation protein	Unknown function	Unknown function	Growth-associated response during stationary phase
16. ydaF	Fucose metabolism/DNA damage response (Fucose metabolism is disrupted by prophage) [68]	LB, no O₂, 37°C	Prophage protein (*ydaG)*	CYT	Prophage protein (YdaG)	*sieB* → **ydaF** ← *ydaG* ← Phage infection exclusion protein — Prophage protein	Prophage protein	Unknown function	Prophage protein involved in disruption of fucose metabolism
17. ydcD	Phenylacetate catabolic process (prevalant under anaerobic conditions) [69]	LB, no O₂, 37°C	Cytoplasmic trehalase (*treF)* (osmotic stress) [70]	CYT	Putative peri-plasmic protein (YjbT)	*yncH* → **ydcD** → *yncO* ← growth-related protein expressed in rich medium)	dCTP de-aminase	Unknown function	Anaerobic respiration
18. ydcH	Phenyl acetate catabolic process (prevalant under anaerobic conditions) [71]	LB, no O₂, 37°C	Transcription (*ygeV)*	CYT	Putative structural protein (YaeH)	*opgD* → **ydcH** → *rimL* → Glucans synhesis protein — Ribosomal transferase	Unknown function	Unknown function	Anaerobic respiration
19. ydfA	Not co-regulated	LB, no O₂, 37°C	Putative transcriptional regulator (*ygeV)*	CYT	Qin prophage (constitutively expressed, unknown function) (YdfB)	*dicA* → **ydfA** → *ydfB* → DNA binding transcr. regulator — Consti-tutively expressed protein	Phage protein	Unknown function	Unclear
20. ydfC	Not co-regulated	M9, 37°C; 0.3 M NaCl, + 0.6% n-butanol (causes osmotic stress) [52]	Putative carbamate kinase (*ycbF)* (biofilm formation) [72]	CYT	Qin prophage (constitutively expressed, unknown function) (YdfB)	*ydfB* → **ydfC** → *dicB* → Consti-tutively expressed protein — Cell division inhibition protein	Putative phage protein	Unknown function	Cellular stress response

No. / Gene	Co-regulation	Conditions	Gene function	Loc.	Neighbor function	Left neighbor	Query gene	Right neighbor	Protein description	Predicted function	Conclusion
21. *ydfX*	Not co-regulated	LB, 37°C	Putative glycosyl-transferase (*yefI*)	CYT	Qin prophage (transcriptional regulator) (DicA)	*ydfW* ← Response to antibiotics	**ydfX** ←	*dicC* ← DNA-binding transcr. regulator	Bacterial toxin protein	Unknown function	Unclear
22. *ydgH*	Oxidoreduction (Respiratory complex I) (NADH: ubiquinone oxidoreductase, oxidative stress protection) [73]	LB, 37°C; 1ug/ml of Ciprofloxacin (FC=0.38, relative to *frr* gene)	Beta-glucosidase (*bglA)* (Bgl operon confers fitness advantage under stress) [74]	ECS	Dihydro-monapterin reductase (FolM) (role in oxidative stress) [75]	*pntA* ← Pyrimidine nucleotide hydro-genase – role in oxidative stress [76]	**ydgH** →	*ydgI* → Putative Arginine: ornithine antiporter	YdgH/BhsA/McbA-like domain-containing protein (involved in stress response/biofilm formation and patho-genesis)	Unknown function	Oxidative stress response
23. *ydiH*	Co-regulated (Unknown/ regulator or function)	M9, 37°C; 10% n-hexane (oxidative stress) [77]	Cyclopropane fatty acyl phospholipid synthase (*cfa)*	CYT	Putative ferrodoxin-like lipoprotein (YdjY) (oxidative stress response) [78]	*menI* ← Mena-quinone bio-synthesis	**ydiH** ←	*ydiK* ← Putative trans-porter	Unknown function	Unknown function	Oxidative stress response
24. *ydjO*	Response to acidic pH (upregulation of oxidative stress response genes) [79]	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	Cysteine/ selenocysteine cation transporter (*tcyP)* [80]	CYT	Cell division modulator (CedA) (induced in oxidative stress)	*tcyP* → Cation transporter	**ydjO** ←	*cedA* ← Cell division modulator	Unknown function	Unknown function	Oxidoreduction under acidic pH conditions
25. *yeaC*	Aerobic respiration	LB, 37°C	Methionine sulphoxide reductase (*msrB)* (oxidative stress) response [81]	CYT	Methionine sulfoxide reductase B (MsrB)	*ydjL* → Putative Zn-binding dehydro-genase (oxidative stress response) (UniProt)	**yeaC** ←	*msrB* ← Meth-ionine sulphoxide reductase	Transcriptional regulator	Unknown function	Oxidative stress response during aerobic conditions

# Gene	Process	Conditions	Interacting protein	Loc.	Description	Left gene	Center gene	Right gene	Function A	Function B	Conclusion
26. yebO	Cellular response to Magnesium ion (during aerobic respiration) [82]	M9, 37°C; double mutant for genes involved in aerobic respiration *(Δndh ΔcyoB)* [83]	PhoQ kinase inhibitor (*mgrB*)	PM	PhoP/PhoQ signaling protein (MgrB) (induced by low Mg2+) [84]	*yobF* ← General stress response protein	**yebO** ←	*mgrB* ← Kinase inhibitor	Unknown function	Unknown function	Response to magnesium ions during aerobic respiration.
27. yebW	Not co-regulated	LB, 37°C	Phospho-protein phosphatase (*pphA*)	CYT	Lipopolysaccharide biosynthesis protein (WbbK)	*yebV* → Protein binding	**yebW** ←	*pphA* → Phospho-protein phos-phatase	Lipoprotein	DUF1482 family protein	Unclear
28. yecH	Anaerobic respiration	M9, 37°C; triple mutant of genes involved in aerobic respiration *(ΔndH, ΔcydB, ΔappC)* [83]	DNA gyrase inhibitor (*sbmC*) (involved in SOS response) [85]	CYT	Putative nucleoside tri-phosphatase (SgcQ)	*ftnA* → Ferritin iron storage protein (oxidative stress response) [86]	**yecH** ←	*tyrP* → Tyrosine symporter	Metal-binding protein	YecH family protein	Switching from aerobic to anaerobic respiration under stress
29. yecT	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (causes osmotic stress) [52]	Multidrug/cho-line/betaine efflux transporter (*emrE*) (osmotic stress response) [87]	CYT	Putative DNA-binding transcriptional regulator (YhfZ)	*yecV* ← Expo-nential phas protein (in rich medium)	**yecT** →	*flhE* ← Flagellar protein	Lysozyme inhibitor	LprI domain-containing protein (lipo-protein)	Osmotic stress response
30. yegJ	Cellulose biosynthetic process	LB, no O₂, 37°C	Reactive chlorine species responsive protein (*rclC*)	ECS	Putative non-specific DNA-binding protein (YegI)	*yegI* → Ser/Thr Phos-phatase	**yegJ** →	*pphC* → Ser/Thr Phosp-hatase	Unknown function	Unknown function	Unclear
31. yegL	Not co-regulated	LB, no O₂, 37°C	Zinc-finger domain protein (*yehQ*)	CYT	Ser/thr phosphatase realted protein (YegK)	*pphC* → Ser/Thr phos-phatase	**yegL** ←	*yegR* →	Homeostasis protein	Mediator of transcription	Unclear
32. yegR	Phosphorelay signal transduction system / Oxalate catabolic	M9, 37°C; double mutant for genes involved in aerobic respiration *(Δndh ΔcyoB)* [83]	Putative oxido-reductase (*ydeP*)	ECS	Orphan protein (OrgK)	*ogrK* ← Trans-cription regulation	**yegR** ←	*yegS* Lipid kinase	Lipoprotein	Unknown function	Oxidoreduction

	process (involved in acid stress resistance) [88]										
33. yehI	Not co-regulated	LB, no O₂, 37°C	ATP-binding lipopolysaccharide transport protein (*msbA*)	CYT	Putative ATPase (YehL)	*yehF* → Molybdate transport regulator	**yehI** →	*yehK* →	Putative regulator	DUF4132 domain-containing protein	Transport protein
34. yehK	Not co-regulated	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	Putative fimbrial adhesin protein (*sfmH*)	CYT	Putative ATPase (YehL)	*yehI* →	**yehk** →	*orzP* → Small RNA	Unknown function	Unknown function	Unclear
35. yeiS	Not co-regulated	LB, 37°C	Vanomycin responsive protein (*sanA*)	CYT	Cell membrane barrier (SanA)	*sanA* → Response to Vanomycin	**yeiS** →	*preT* → dehydro-genase	Vancomycin high temperature exclusion protein	Unknown function	Maintaining cell membrane integrity against drugs (e.g. Vanomycin)
36. yfbM	Aerobic respiration	M9, no O₂, 37°C	Hemolysin E (*hlyE*) (pore forming protein) [89]	CYT	Unknown function (YfbN)	*yfbL* → Putative peptidase (proteo-lysis) (EcoCyc)	**yfbM** →	*yfbN* ←	Unknown function	DUF1877 family protein	Protein lysis
37. yfbN	Response to acidic pH	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	DNA binding transcriptioal regulator (*ydeO*)	CYT	Unknown function, YfbO	*yfbM* →	**yfbN** ←	*yfbO* →	Unknown function	Unknown function	Unclear
38. yfbO	Not co-regulated	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	Function unknown (*yfbP*)	ECS	Unknown function (YfbP)	*yfbN* ←	**yfbO** →	*yfbP* →	Immunity protein (cyto-toxicity)	Unknown function	Unclear
39. yfbP	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol	Inner membrane protein (*ynjI*)	CYT	Putative amino-peptidase (YfbL) (Proteolysis) (EcoCyc)	*yfbO* ←	**yfbP** →	*nuoN* ← Oxido-reductase	Protein folding and transport	Unknown function	Protein quality control under stress
40. yfdN	Not co-regulated	LB, no O₂, 37°C	Oxidative stress response protein (*yfdO*)	CYT	Prophage protein (phage-related function) (YfdT)	*yfdM* → Putative methyl-transferase	**yfdN** ←	*yfdO* ← Oxidative stress response	Transcriptional activator family protein	Unknown function	Oxidative stress response

41. yfdP	Not co-regulated	LB, no O₂, 37°C	Putative auto-transporter adhesin (*ycgV*)	CYT	Prophage protein (response to oxidative stress) (YfdS)	*yfdO* ← Oxidative stress response **yfdP** →	*yfdQ* → Phage-related function	Prophage protein	Unknown function	Oxidative stress response
42. yfeS	Cellular response to envelope stress	MOPS, 37°C	Putative lipoprotein (*yidX*)	CYT	Putative efflux protein (YdhK)	*yfeK* → Transcr. regulation **yfeS** →	*cysM* ← Cysteine synthase	Molybdate metabolism protein	Unknown function	Envelope stress response
43. yffL	Response to Nitrate/ Nitrite (anaerobic respiration)	M9, 37°C	Anaerobic glycerol-3-phosphate dehydro-genase (*glpB*) [89]	CYT	Prophage protein (Integrase) (IntZ)	*intZ* → Putative integrase **yffL** →	*yffM* →	Prophage protein	CPZ-55 prophage (involved in general stress response) [90]	Anaerobic respiration
44. yffM	Not co-regulated	LB, no O₂, 37°C	Alpha-mannosidase (*mngB*)	CYT	Prophage (unknown function) (YffN)	*yffL* → **yffM** →	*yffN* →	Unknown function	Unknown function	Unclear
45. yffN	Not co-regulated	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	Tail fiber protein (*stfR*)	CYT	Prophage (unknown function) (YffM)	*yffM* → **yffN** →	*yffO* →	Unknown function	Unknown function	Unclear
46. yffP	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol	Function unknown (*yffO*)	CYT	Prophage (unknown function) (YffO)	*yffO* → **yffP** →	*yffQ* →	Unknown function	Unknown function	Unclear
47. yffQ	Not co-regulated	MOPS, 37°C; 0.4% Glycerol	CRISPR-associated endo-ribonuclease (*cas2*) (defense response to virus) (EcoCyc)	CYT	Prophage (unknown function) (YffP)	*yffP* → **yffQ** →	*yffR* ←	Unknown function	CPZ-55 prophage (involved in general stress response) [90]	Stress response
48. yffR	Not co-regulated	M9, 37°C; triple mutant of genes involved in aerobic respiration (*ΔndH, ΔcydB, ΔappC*) [83]	Bacterial adhesion/ biofilm formation (*yhjY*)	CYT	Prophage (unknown function) (YffQ)	*yffQ* → **yffR** →	*yffS* → DNA bindin	Unknown function	Unknown function	Stress response

# Gene	Co-regulation	Condition	Protein	Loc.	Function	Gene neighbourhood			Product	Function	Conclusion
49. *yfiM*	Cell death	LB, 37°C	Alpha-ketoglutarate: H^+ symporter (*kgtP*) (efflux transporter for sugar-phosphate stress) [91]	ECS	Phospholipid biosynthesis protein (PssA)	*pssA* → Phos-phatidyl-serine synthase	**yfiM** →	*kgtP* ← H^+ symporter	Lipoprotein	Unknown function	Lipoprotein with a role in mitigating the effects of cellular stress
50. *yfjH*	Not co-regulated	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose	Growth related protei (*htrL*)	CYT	Prophage (transcriptional activation) (AlpA)	*intA* → Integrase protein	**yfjH** ←	*alpA* → Transcr. regulator	Unknown function	Unknown function	Unclear
51. *yfjJ*	Response to acidic pH (upregulation of genes involved in cell adhesion) [92]	MOPS, 37°C; 0.4% Glycerol and 0.5% 2-Deoxy-D-glucose	Putative fimbrial usher protein (*htrE*) – cell adhesion or pilus assembly protein [92]	CYT	Prophage (cytosolic protein) (YfjI)	*yfjI* → Prophage protein	**yfjJ** →	*abpB* ← Putative helicase	Inovirus-type Gp2 protein (viral DNA replication)	Unknown function	Cell adhesion
52. *ygdT*	Not co-regulated	LB, no O_2, 37°C	Putative lipoprotein (*yidX*)	CYT	Not found	*rppH* ← RNA phospho-hydrolase	**ygdT** ←	*mutH* → DNA repair protein	Transposase protein	Unknown function	Unclear
53. *yhaL*	Not co-regulated	LB, 37°C; *Δhns* (stress adaptation)	DNA binding transcriptional activator (*fucR*)	CYT	Ser/thr phosphatase-related protein (Yegk)	*yhaK* → bicupin related protein (acid stress/oxidative stress response) [93]	**yhaL** →	*cyuA* ← L-cysteine di-sulphidase	Cytoplasmic protein	Unknown function	Stress response

54. yhdN	Response to heat	LB, 30°C (stationary phase)	DNA binding transcriptional activator (*zntR*) (stress responsive transcription factor) [94]	CYT	Transcriptional activator (ZntR)	*zntR* ← DNA binding transcr. regulator	**yhdN** ←	*rplQ* ← 50S ribosome subunit protein (protein bio-genesis) [95]	Conserved protein	DUF1992 domain-containing protein (conserved protein belonging to the DNAJ homolog family, heat shock response) (UniProt)	Regulation of heat shock response, expressed during stationary phase
55. yhfU	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (osmotic stress) [53]	Aromatic carboxylic acid efflux pump (*aaeA)* (involved in defense mechanisms) [96]	CYT	Putative transport protein (YhfT)	*yhfT* ← Membrane protein	**yhfU** ←	*php* ← Putative hydrolase	Unknown function	Unknown function	Osmotic stress response
56. yhhL	Not co-regulated	LB, 37°C; *ΔdsrA* (gene involved in stress response, biofilm formation) [97]	16S rRNA methyl-transferase (*rsmD)*	CYT	16S rRNA methyl-Transferase (RsmD)	*rsmD* → 16S rRNA methyl transferase	**yhhL** xs →	*yhhM* Quorum sensing protein – (stress response, biofilm formation) [98]	rRNA small subunit methyl-transferase protein	Unknown function	Biofilm formation
57. yibI	Response to nitrosative stress (redox stress) [99]	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (osmotic stress) [52]	Inner membrane protein (*yibH)* (stress response) [100]	PM	Inner membrane protein (YibH)	*yibH* ← Inner membrane protein	**yibI** ←	*mtlA* → mannitol-specific PTS enzyme	Inner membrane protein	Unknown function	Stress response protein.
58. yicN	Not co-regulated	LB, 37°C; *Δhns* (stress adaptation) [101]	DNA binding transcriptional regulator (*frlR)*	CYT	Putative Transporter (NepI)	*nepI* ← Purine ribo-nucleoside exporter	**yicN** ←	*adeQ* Adenine trans-porter	Unknown function	Unknown function	Nucleotide transport

Gene	Co-regulation	Conditions	Protein (gene)	Loc.	Characterized protein	Upstream gene	Gene	Downstream gene	Protein	Function	Conclusion
59. *yidB*	Not co-regulated	LB, 37°C; triple mutant of genes involved in respiration *(ΔndH, ΔcydB, ΔappC)* [83]	Bitopic inner membrane protein (*ypfJ*)	CYT	Sugar phosphate phosphatas (YidA) (phosphate metabolism) [102]	*yidA* ← Sugar phos-phatase	**yidB** ←	*gyrB* ← DNA gyrase	Radical SAM domain-containing protein	Unknown function	Unclear
60. *yigA*	Not co-regulated	LB, 37°C	Diamino-pimelate epimerase (*dapF)* (cell wall biosynthesis) [103]	CYT	Diamino-pimelate epimerase (DapF) (cell wall structure) [104]	*dapF* → diamino-pimelate epimerase	**yigA** →	*xerC* → Tyrosine recombi-nase	Diamino-pimelate epimerase protein	Not found	Cell wall biosynthesis
61. *yigE*	Not coregulated	LB, no O₂, 37°C	Putative magnesium chelatase (*yifB*) (chaperonic activity) [105]	CYT	Inner membrane protein (YedI)	*uvrD* → DNA helicase	**yigE** ←	*ysgD* → Consti-tutively expressed protein (in rich medium)	Phospho-diester glycosidase family protein	Anaerobic nucleoside-tri-phosphate reductase-activating protein (DNA synthesis and repair) [106]	Stress response during anaerobic conditions
62. *yjbD*	Not coregulated	M9, 37°C; 10% n-hexane	Putative methyl-transferase (*yafS*)	CYT	Cysteine cluster protein (YcgN)	*rluF* → Pseudo-uridine synthase	**yjbD** ←	*lysC* ← Aspartate kinase	Unknown function	Unknown function	Unclear
63. *yjbL*	Response to acidic pH	LB, no O₂, 37°C	Cell envelope stress response protein (*yqcG*)	CYT	Uncharacterized protein (YjbM)	*zur* → Transcr. regulator	**yjbL** →	*yjbM* → Cell-envelope related protein	Unknown function	Unknown function	Envelope stress response under stress
64. *yjbS*	Response to acidic pH (upregulation of oxidative stress response genes) [107]	M9, 37°C; 10µg/ml Mitomycin-C (genotoxic agent, ROS producing) [108]	Pathogenicity-related (*yqeK*)	CYT	Putative oxido-Reductase (YqiJ) ([109]	*tyrB* → Tyrosine amino-transferase	**yjbS** ←	*aphA* → Acid phos-phatase	Unknown function	Unknown function	Oxidative stress response.
65. *yjcB*	Carbohydrate transport	M9, no phosphate, 37°C; *ΔphoB* (phosphate transport gene)	Capsule synthesis regulator (*rcsB*)	CYT	Putative peri-plasmic protein (YpeC)	*ssb* → ss-DNA binding protein	**yjcB** ←	*pdeC* → Cyclic di-GMP Phospho-diesterase	Membrane protein	Not found	Unclear

# Gene	Function	Condition		Loc		Gene neighborhood				Conclusion
66. yjcF	Response to acidic pH	LB, 37°C	Pathogenicity-related (*ygeH*)	CYT	Ribosomal silencing factor (RsfS)	*yjcE* → Putative transporter	**yjcF** ←	*actP* ← Cation trans-porter	Unknown function / Unknown function	Unclear
67. yjdI	Oxidoreduction (Respiratory complex I)	LB, 37°C; triple mutant of genes involved in respiration *(ΔndH, ΔcydB, ΔappC)*	Putative N-acetyl-Transferase (*yjdJ*)	CYT	Putative N-acetyl-Transferase (YjdJ)	*dcuS* ← Sensor histidine kinase	**yjdI** →	*yjdJ* → Putative acetyl-trans-ferase	Unknown function / Fer4_19 domain-containing protein (involved in iron-sulphur binding)	Aerobic respiration
68. yjeJ	Cell adhesion	LB, 37°C; HS, 50°C for 7'	6-phospho-gluconate phosphatase (*yieH*)	CYT	Lysylation protein (EpmB)	*yejI* → Lipoprotein (heat shock response) [110]	**yjeJ** ←	*epmB* ← Lysine amino-mutase	Unknown function / Unknown function	Cell adhesion under heat shock response
69. yjeT	Not coregulated	LB, 37°C; HS, 50°C for 15'	Lipo-polysaccharide core heptose kinase *(waaP)*	CYT	HflB protease modulator (HflC) (in heat shock response) [111]	*hflC* → ftsH protease regulator (heat shock response) [111]	**yjeT** →	*purA* → Adenylo-suucinate syn-thetase	Unknown function / Unknown function	Heat shock response
70. yjfI	Coregulated (unknown regulator or function)	LB, 37°C, reaction buffer (with Lysine)	Inner membrane protein (*yjgN*)	CYT	Membrane chaperone family protein (YjfJ) [112]	*rlmB* → Methyl-transferase	**yjfI** →	*yjfJ* → Transcr. regulator	Cytoplasmic protein / Unknown function	Unclear
71. yjfK	Not coregulated	LB, no O$_2$, 37°C	Membrane chaperone family protein (*yjfJ*)	CYT	Inner membrane protein (YjfL)	*yjfJ* → Transcr. regulator	**yjfK** →	*yjfL* → Inner membrane protein	Unknown function / Unknown function	Unclear
72. yjfZ	Not coregulated	LB, 37°C	Acyl transferase (*yihG*)	CYT	Putative inner membrane poly-saccharide flippase (YghQ)	*rpiI* → Ribosomal subunit protein	**yjfZ** ←	*ytfB* ← Cell division	Unknown function / Unknown function	Unclear

73. *yjgR*	Not coregulated	LB, no O₂, 37°C	NADPH dependent aldehyde reductase (*ahR)*	PM	L-idonate and D-gluconate transporter (IdnT)	*iptG* → LPS Transport system	**yjgR** ←	*idnR* ← DNA binding transcr. regulator	Ornithine amino-transferase	Conjugal transfer protein	Transport
74. *yjgZ*	Coregulated (unknown regulator or function)	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol	Putative multidrug efflux pump (*mdtO*)	CYT	Putative receptor protein (YhhM)	*insD6* → Trans-posase	**yjgZ** →	*insG* ← Putative transposase	Transposase	Unknown function	Unclear
75. *yjjY*	Cell death	LB, 37°C	DNA binding transcriptional regulator (*arcA*)	CYT	Uncharacterized protein YniD	*arcA* ← DNA binding transcr. regulator	**yjjY** →	*ytjE* → Growth related protein	Unknown function	Unknown function	Unclear
76. *ykgL*	Glycine betaine biosynthesis from choline [49]	M9, 37°C; *ΔkdpE*; Tris-maleic acid+115 mM KCl (disrupts potassium uptake under osmotic stress) [113]	50S ribosomal subunit protein (*ykgM*)	CYT	Carnitinyl-CoA dehydratase (CaiD) – osmoprotectant protein [114]	*matA* ← DNA binding transcr. regulator	**ykgL** →	*ykgO* ← 50S ribosomal subunit protein	Unknown function	Unknown function	Osmotic stress response.
77. *ylcH*	Trans-criptional attenuation	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol	Ionic detergents response protein (*yfbK*)	No inf.	Uncharacterized protein (YlcG)	*ybcM* → DNA binding transcr. regulator	**ylcH** →	*ybcN* → DNA base-flipping protein	Unknown function	Unknown function	Unclear
78. *ylcI*	Aerobic respiration	LB, 37°C; double mutant of genes involved in respiration (*ΔndH, ΔcyoB*)	Phage related protein (*ybcW*)	CYT	Oxidoreductase (NuoM) – involved in aerobic respiration [115]	*ybcW* → Prophage protein	**ylcI** ←	*nohD* → DNA packing protein	Unknown function	DUF3950 domain-containing protein	Aerobic respiration
79. *ymfI*	DNA restriction/ modification system	LB, 37°C; 1 ug/mL of Ciprofloxacin+ 500 μM of ET2DA	Phage related protein (*ymfJ*)	CYT	Uncharacterized protein (*ymfJ)*	*ymfH* ← Prophage protein	**ymfI** →	*ymfJ* ← Prophage protein	Phage protein	Unknown function	Unclear
80. *ymgI*	Not co-regulated	M9, no phosphate, 37°C; *ΔphoB* (phosphate transport gene)	Heat shock responsive protein (*ymgG*)	CYT	Cell division topological specificity protein (MinE)	*ymgG* ← Heat shock response protein	**ymgI** ←	*ymgJ* →	Unknown function	Unknown function	Unclear

Gene	Co-regulation	Condition	Related protein	Loc.	Function	Left gene	Center gene	Right gene			Category
81. ymgJ	Not co-regulated	M9, 37°C; 0.3 M NaCl + 0.6% n-butanol	Putative fimbrial usher protein (*htrE*) (heat shock response) [116]	CYT	Heat shock response protein (YmgG)	*ymgI* ←	***ymgJ*** →	*minE* ← / Cell division	Transposase protein	Unknown function	Heat shock response
82. ynaJ	Aerobic respiration	LB, 37°C	ADP-sugar diphosphatase (*nudE*)	CYT	Translation-realted protein (YciU)	*insH4* → / Trans-posase	***ynaJ*** →	*uspE* ← / Universal stress protein	Inner membrane protein	Unknown function	Unclear
83. yncH	Response to acidic pH	LB, 37°C; *rpoD* overexpression (response to heat) [117]	Cell envelope stress response protein (*yqcG*)	CYT	Glutathione S-transferase homolog (YncG)	*yncG* → / Putative Glutata-thione-S transferase	***yncH*** →	*yncO* ← / Consti-tutively expressed protein (in rich medium)	Unknown function	Unknown function	Stress response
84. yneG	Not co-regulated	M9, no ammonium, 37°C	Glutaminase (*glsB*) Ammonium assimilation [118]	CYT	Putative Glutaminase (GlsB) (amino acid metabolism) [119]	*dgcF* ← / Putative Di-guanylate cyclase	***yneG*** ←	*glsB* ← / Glutaminase	Cytoplasmic protein	Unknown function	Nitrogen metabolism during ammonium limitation
85. yneK	Not co-regulated	MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose (metabolic stress) [120]	Putative fimbrial usher protein (*htrE*) (heat shock response) [116]	CYT	Putative oxido-reductase (YdhF)	*ptrR* → / L-glutamate meta-bolism	***yneK*** →	*ydeA* → / L-arabinose exporter	Unknown function	Unknown function	Stress response
86. ynfO	Anaerobic respiration	LB, 37°C; 100 µg/ml Ampicillin	Function unknown (*ylcI*)	CYT	Putative (D)-galactarate transporter (GarP)	*nohA* ← / Prophage protein	***ynfO*** →	*ydfO* → / Prophage protein	Unknown function	DUF3950 domain-containing protein	Unclear
87. yoaC	Not co-regulated	LB, 37°C	Lipoprotein (*yfiL*)	CYT	Protein export (MsyB)	*yoaB* → / Biofilm based activity	***yoaC*** →	*yoaH* ←	Unknown function	Unknown function	Unclear

Gene	Co-regulation	Condition	Function	Loc.	Domain / description	Gene context		Left neighbor	Right neighbor	Category
88. *yoaH*	Not co-regulated	LB, 37°C; *Δhns* (stress adaptation)	DNA-binding transcriptional repressor (*acrR*)	CYT	DUF1889 family protein (YoaC)	*yoaC* → **yoaH** ← *pabB* → Amino-deoxy-choris-mate synthase		Unknown function	Unknown function	Unclear
89. *yodC*	Oxidoreduction (Respiratory complex I)	LB, 37°C	Putative adenosine mono-phosphate (*fic*)	CYT	Ribosome modulation protein (RmF) (response to oxidative stress) [121]	*dgcQ* ← **yodC** ← *yedI* ← Di-guanylate cyclase / Inner mem-brane protein		Cytoplasmic protein	Unknown function	Oxidoreduction
90. *ypaB*	Not coregulated	M9, 37°C; 0.3 M NaCl +0.6% n-butanol	Putative fimbrial chaperone (*yadV*)	CYT	Not found	*yfaL* ← **ypaB** ← *nrdA* → Putative auto-transporter adhesin / Ribo-nucleoside reductase		Transposase	Unknown function	Unclear
91. *ypeB*	Ribosome structure	M9, 37°C; 50 mM isoprenol	Tetraacyl-disaccharide 4'-kinase (*lpxK*)	CYT	DNA ligase (LigA)	*yfeH* → **ypeB** ← *ligA* ← Putative Na⁺ symporter / DNA ligase		Cytoplasmic protein	Cyto-plasmic protein	Unclear
92. *ypjB*	Not coregulated	LB, no O₂, 37°C	Putative type III secreted effector (*yqeK*) (bacterial patho-genecity) [122]	CYT	Not found	*ypjA* ← **ypjB** ← *ypjC* → auto-transporter / Patho-genicity		Uncharacterized protein	Unknown function	Pathogenecity
93. *yrdB*	Oxidoreduction (Respiratory complex I)	LB, 37°C	Shikimate dehydro-genase (*aroE*)	CYT	Dehydro-shikimate reductase (AroE) - oxidative stress response) [123]	*yrdA* → **yrdB** ← *aroE* ← carbonic anhydrase / shikimate anhydrase		DUF1488 domain-containing protein	Unknown function	Oxidoreduction
94. *yrhD*	Response to acidic pH	LB, no O₂, 37°C	Putative type III secreted effector (*yqeK*) (bacterial patho-genecity) [122]	CYT	Not found	IS1E → **yrhD** → *yrhB* → Insertion sequence / Putative HS chaperone		Unknown function	Unknown function	Unclear
95. *ytiA*	Not co-regulated	LB, no O₂, 37°C	Putative transporter (*yjiJ*)	CYT	Not found	*yjiK* ← **ytiA** ← *yjiL* ← Membrane / Putative de-		DUF3343 domain-containing	Unknown function	Unclear

						protein	hydratase	protein		

Abbreviations: ATP, Adenosine triphosphate; dCTP, Deoxycytidine triphosphate; ET2DA, Ethyl trans-2-decenoic acid; HS, Heat shock; KCl, Potassium chloride; LB,

Luria Bertani; LPS, Lipopolysaccharide; M9, Minimal medium; MOPS, 3-(N-morpholino) propanesulfonic acid; NADP, Nicotinamide dinucleotide phosphate; PTS,

phosphotransferase system; ROS, Reactive oxygen species; SAM, S-adenosylmethionine; Ser/Thr, Serine/Threonine; Transcr. , Transcription

Genes with a higher confidence in the inference section (green) were analysed for candidates

to be used for *in vitro* testing. Three HP-encoding genes, namely *yhdN*, *yeaC*, and *ydgH*, were

selected for further analysis (highlighted in grey) (**Table 1).**

Protein encoding gene *yhdN*

Protein YhdN, characterized by the presence of a domain of unknown function (DUF1992),

was predicted to be involved in heat shock response. The gene encoding *yhdN* is regulated

by the RNA polymerase sigma factor RpoH (σ32), which serves as the primary heat shock

transcriptional regulator. Under normal conditions, RpoH levels are kept low due to chaperone-

mediated degradation; however, these levels increase significantly upon heat shock [124].

Gene Ontology (GO) annotation for this cluster is categorized under 'response to heat'. Based

on publicly available transcriptomic data [43], the *yhdN* gene exhibited significant expression

in cells grown to the stationary phase, with a Log_2FC of 5.5 relative to the reference gene *frr*,

signifying a 45 times higher expression of *yhdN* gene. This observation suggests that

the *yhdN* gene is highly expressed during the stationary phase and therefore might have a

role in nutrient stress response. Additionally, *yhdN* expression was upregulated following

exposure to a transient heat shock at 50°C for 15 minutes, with a Log_2FC of 4.8 and 27 times

higher expression, suggesting that the gene might also play a role in heat shock response.

Co-expression analysis using Spearman correlation of the gene resulted in a high correlation

with the gene *zntR* (0.93) encoding an HTH (helix-turn-helix)-type transcriptional regulator

[125] which is involved in protein stability and prevention of aggregates [126,127]. Genes *yhdN*

and *zntR* are also part of the same transcription unit (EcoCyc database). Other genes with

high correlations to *yhdN* were *relB* (0.84)*, pspB* (0.84) *and pspC* (0.84), all of which are

involved in stress response [128] (*Supplementary Figure 3A.*). Gene *relB* encodes for an

antitoxin protein involved in regulation of growth [129] while genes *pspB* and *pspC* encode for

phage shock proteins that are known to be induced by infection with bacteriophages or

ethanol, osmotic and/or nutrient stress as well as heat shock conditions [130]. Protein-protein

interaction (PPI) analysis showed highest interaction confidence with ZntR. The local genomic

context analysis showed genes *zntR* and *rplQ* (a ribosomal subunit protein) as neighbouring

genes. Protein YhdN showed remote sequence homology to a protein cluster comprising of

uncharacterized conserved proteins and was structurally homologous to conserved proteins

with similarity to J-domains of chaperones, regulating the activity of heat-shock proteins [131].

To elucidate the potential function of *yhdN*, we hypothesized that its absence would affect

bacterial growth under heat shock conditions and possibly the cells stationary phase under

nutrient limitation. This hypothesis was tested by comparing the growth patterns of the

E. coli K-12 substrain BW25113 wild type and its isogenic *ΔyhdN* knockout mutant under a

transient heat shock condition (50°C) and recovery at 37°C in LB and nutrient-limited M9

medium.

Protein encoding gene *yeaC*

Protein YeaC is characterized by the presence of a DUF1315 domain. Gene *yeaC* is predicted

to be governed by the regulator ArcA, which regulates a group of proteins involved in redox

homeostasis and aerobic respiration [132]. The Gene Ontology (GO) annotation for this cluster

was predicted as 'aerobic respiration'. Analysis of gene *yeaC* in publicly available

transcriptomic datasets showed slightly higher expression levels against gene *frr* in LB

medium at 37°C under normal O_2 conditions ($\log_2$ FC 1.9) (**Table1**) and in knock-out mutants

for genes encoding type II NADH:quinone oxidoreductase (*ndh*), Cytochrome bd-I ubiquinol

oxidase (*cydB*) and Cytochrome bd-II ubiquinol oxidase (*appC*), involved in the Tricarboxylic

acid cycle or oxidative phosphorylation [133] ($\log_2$ FC 1.7). Based on the co-expression

analysis of public data, the highest correlation was observed with a gene encoding for a

peptide methionine sulfoxide reductase (*msrB),* which is an oxidoreductase and known to be

highly expressed during oxidative stress to maintain the cell-redox homeostasis [134]

(*Supplementary Figure3B.*). Other highly correlated genes involved succinate dehydrogenase

cytochrome b556 subunit (*sdhC*), succinate dehydrogenase hydrophobic membrane anchor

subunit (*sdhD)* and Fumarate hydratase class I, aerobic (*fumA),* which are all involved in the

tricarboxylic acid or TCA cycle [135,136] (*Supplementary Figure 3B.*). PPI analysis also

showed highest interaction confidence with MsrB. Additionally, it neighbours the *msrB* gene

and a putative zinc-binding dehydrogenase, *ydjL* which is also involved in oxidoreduction

(UniProt) (**Table 1**). Protein YeaC shows remote sequence homology to other proteins

involved in transcriptional regulation. Structural homology revealed similarities to a group of

uncharacterized proteins with unknown functions. We hypothesized that *yeaC* affects the cell-

redox homeostasis and cellular growth under oxidative stress. This hypothesis was tested by

comparing the growth patterns of the *E. coli* K-12 substrain BW25113 wild type and its isogenic

ΔyeaC knockout mutant with a sub-lethal concentration of H_2O_2 (2.5 mM) [137].

Protein encoding gene *ydgH*

Protein YdgH has a DUF1471 domain of unknown function. It is predicted to be part of an

unknown cluster, with no information on its regulator. The GO annotation for this cluster,

however, is categorized under oxidoreduction (complex I). The energy-converting

NADH:ubiquinone oxidoreductase respiratory complex I, is the main entry point for electrons

from NADH into the respiratory chains in bacteria [138]. Based on the public datasets, we

could not obtain a gene-specific condition where gene *ydgH* had higher expression compared

to gene *frr.* Co-expression analysis of *ydgH* from the publicly available RNAseq datasets

revealed highest correlation with a 6-phospho-beta-glucosidase (*bglA*) encoding hydrolase,

involved in the hydrolysis of phosphorylated beta-glucosides [139]. Other highly correlated

genes were phosphoglycolate phosphatase *(gph)* [140] and ribulose-phosphate 3-epimerase

(rpe) [141], involved in oxidative stress response (*Supplementary Figure 3C.*). PPI analysis

showed highest interaction confidence with YjfY*,* a putative HP, with a possible role in stress

resistance (UniProt). Genomic context provided information on a neighbouring upstream gene

involved in $NAD(P)^+$ transhydrogenase activity, annotated as NAD(P) transhydrogenase,

pntA, which is known to play a role in oxidative stress [142]. The protein showed remote

sequence homology to a YdgH/BhsA/McbA-like domain involved in stress response/biofilm

formation and pathogenesis [143]. The structural similarity was only to uncharacterized

proteins with unknown function. Following a similar hypothesis to that proposed for *yeaC*, we

evaluated the growth patterns of the wild type and its isogenic *ΔydgH* knockout mutant under

oxidative stress.

In vitro analyses of candidate proteins

Effect of heat shock on growth

To determine the effect of a transient heat shock at 50°C (7 minutes), the wild type and the

mutant *ΔyhdN E. coli* BW25113 strains were grown in LB medium. The *yhdN* mutant was seen

to have a growth defect of ~18% after 6h of growth based on the OD values obtained (**Figure**

3A). It was observed that after 6h of growth, the mutant could not resume normal growth

compared to the wild type. Additionally, the effects of a transient heat shock in Nitrogen-limited

M9 medium showed a significant growth defect of ~50% (**Figure 3B**). Growth curves for WT

and *ΔyhdN* in different temperatures can be found in the supplementary (*Supplementary*

Figure 4A.). Growth defects were also observed in the *yhdN* mutant compared to the wild type

when both the strains were grown continuously at 50°C. Since growth at 50°C is however

severely inhibited and cell started dying after 3h (*Supplementary Figure 4B.*), it was difficult to

say whether this was solely based on the absence of the protein YhdN in the mutant cells.

Effect of sublethal H_2O_2 concentration on growth

To determine the effect of sublethal H_2O_2 concentration, the wild type (WT), *ΔydgH* and *ΔyeaC*

E. coli BW25113 strains were grown in LB medium until an OD of 0.2 and then exposed to an

H_2O_2 concentration of 2.5 mM, which is known to causes DNA damage and oxidative stress

[143] (**Figure 3C & D**). To study the responses due to oxidative stress, the absorbance values

(OD_{600}) after each hour over a 5h time period were measured. H_2O_2 was added to the cell

culture during the exponential phase of growth, which is characterized by rapid cell division

and metabolic activity and where the quorum-sensing signalling, i.e. process which allows the

bacteria to communicate and adjust gene expression according to cell density, is minimal.

Under non-stress conditions, the WT and mutant strains (*ΔydgH* and *ΔyeaC*) exhibited

comparable growth, with no significant differences observed. A growth lag was observed in

the mutants (*ΔydgH* and *ΔyeaC*) as compared to wild type when the medium was

supplemented with 2.5 mM H_2O_2. A growth defect of ~25% was observed in the mutant when

WT was compared to *ΔyeaC* at the 1h timepoint. This was reduced to ~7% after 5h of growth

underscoring the probability of the gene being expressed at this time point to counterfeit the

oxidative stress. Similarly, a growth defect of ~36% was observed in *ΔydgH* as compared to

WT at 1h, which decreased to ~2% after 5h, again probably due to stress adaptation and

expression of gene *ydgH*.

371
372
Figure 3. Growth curves of *E.* coli K-12 BW25113 wild type (WT) and isogenic deletion mutants.
(A, B) Effect of transient heat shock on WT and *ΔyhdN* cells. WT and the respective mutant strain were
exposed to a transient heat shock at 50°C for 7 minutes, (at OD_{600} nm = 0.04) and grown for 12 h in **A.**
LB medium and **B.** in Nitrogen-limited M9 medium. **(C, D)** Bacterial growth of WT and respective
mutants at 37°C exposed to sub-lethal concentration of 2.5 mM H_2O_2, added during the exponential
phase (OD_{600} = 0.2) and grown for 5h. **C.** WT and *ΔyeaC* and D. WT and *ΔydgH* cells. WT are indicated
by filled circles (●) and mutants by filled triangles (▲). Green lines represent controls, orange lines stress
conditions. Average of three independent readings taken for each specified condition. Error bars on the
graph indicate standard deviation from the mean.

Differential gene expression analysis

For a comparative analysis of differentially expressed genes (DEGs) and their functions in WT

vs. the mutant strains, all up- and down-regulated genes were analysed (*Figshare > DEGs >*

Table1). For positively regulated DEGs, representing genes upregulated in the WT strain, a

Log$_2$ fold change (Log$_2$FC) threshold of ≥1.5 was applied, corresponding to a minimum 2.8-

fold increase in expression levels. This cut-off ensured the inclusion of genes with significant

upregulation while minimizing noise [144]. Conversely, for negatively regulated DEGs,

corresponding to genes upregulated in the mutant strains, a Log2FC threshold of ≤-1.5 was

used.

WT *vs. ΔyhdN*

Analysis of the WT strain at the 2h and 6h time points showed that *yhdN* was only significantly

expressed after 6h, hence this time point was chosen. In total, 484 DEGs (p_{adj}≤0.1) were

identified, (181) 213 genes with a positive Log$_2$ FC and 271 with a negative Log$_2$FC of 1.5.

Within the positive DEGs, an analysis of the top 10 genes, resulted in one gene involved in

protein folding, five in stress response and four in motility/chemotaxis (**Figure 4A**).

Significantly upregulated genes in the WT strain were *tdcB* (Log$_2$ FC=7.37) and *pyrB* (Log$_2$

FC=6.14)*. tdcB*, encoding for threonine dehydratase is known for being expressed during

nutrient deprivation (EcoCyc) and gene *pyrB* is involved in pyrimidine metabolism and

associated with motility under stress [145]. Gene *tnaA* (Log$_2$ FC=6.02), which encodes for

tryptophanase, has been recently studied for its role in protein folding and dissolution of protein

aggregates [146]. Motility and chemotaxis genes such as *fliL* (Log$_2$ FC=6.24)*, flgD* (Log$_2$

FC=5.99) are also known to be involved in response to environmental stresses [147].

In the deletion mutant (*ΔyhdN*), the top ten negative DEGs included two genes involved in

heat shock, six in some type of stress response and interestingly two other were

uncharacterized hypothetical proteins. *bolA* (Log$_2$ FC=-5.53)*,* a DNA binding transcriptional

regulator, plays a role in cellular stress response including heat stress [148] and *mgtS* (Log$_2$

FC=-5.01), a small inner membrane protein encoding gene, is known to be expressed upon

heat shock [149] (**Figure 4A**). Other upregulated genes in *ΔyhdN* involved *glgS* (Log$_2$ FC=-

5.34), encoding for a surface composition regulator for biofilms and *ymgA* (Log$_2$ FC=-4.44)

encoding a putative two-component system connector protein, known to play an important role

in biofilm formation (EcoCyc). Upregulation of these genes in the deletion mutant suggests a

compensatory mechanism to mitigate stress. Additionally, a HP-encoding gene, *yodC* was

also upregulated (Log$_2$ FC=-6.39) in *ΔyhdN*. Interestingly, this gene was predicted to play a

role in oxidoreduction (**Table1)**. The results of this transcriptomic experiment could further aid

in forming a hypothesis of this gene for *in vitro* testing. Another HP-encoding gene *yffR* (Log$_2$

FC=-4.85)*,* was predicted to play a role in stress response (this study, Table 1) which aligns

with the fact that a majority of stress-response genes were upregulated in the WT, when gene

yhdN is present.

All observations highlight WT strain's prioritization of growth and motility, possibly facilitating

better environmental sensing and resource acquisition under stress while the absence of *yhdN*

gene seems to upregulate other heat shock and stress response genes which reorient

metabolic processes towards alternative survival strategies such as biofilm formation. Hence,

the protein encoding gene *yhdN* is predicted to be a stress response gene involved in the

regulation of heat shock or nutrient limitation response, possibly also playing a broader role in

general stress response.

WT *vs. ΔyeaC*

In WT strain, the *yeaC* gene exhibited a significant upregulation after the 5h, with a log$_2$ FC of

3 relative to the 1h time point, where its expression was undetectable. This log$_2$FC

corresponds to an 8-fold increase in expression. Hence, the 5h time point was chosen for the

analysis. A total of 156 DEGs for WT *vs. ΔyeaC* were obtained after filtering (p_{adj}≤0.1). In WT

vs. yeaC, 87 genes had a positive Log$_2$ FC while 69 had a negative Log$_2$ FC.

Analysis of the top ten upregulated DEGs in the WT strain resulted in three oxidative stress

response genes, two other stress response genes, two methionine biosynthesis genes and

three genes involved in bacterial secretion system and transposition (EcoCyc). The

upregulation of the latter three genes in WT was difficult to explain, since there was no

conclusive literature available (**Figure 4B**). Significantly upregulated DEGs involved *ymcE* (Log$_2$ FC=6.81) which provides tolerance to n-butanol [145], and is also known to provide protection against oxidative stress [150], *prpE* (Log$_2$ FC=4.75) which responds to hydrogen peroxide [151] and *iprA* (Log$_2$ FC=4.34), which encodes for an inhibitor of hydrogen peroxide [152]. *metF* (Log$_2$ FC=4.49), a methionine biosynthesis gene, is also known to provide protection against oxidative stress: Methionine is a precursor in the synthesis of cysteine, which is a component of glutathione—a major intracellular antioxidant. Glutathione is known to neutralize reactive oxygen species [153].

All of the upregulated genes in *ΔyeaC* were involved in motility/chemotaxis-based functions (**Figure 4B**). Significantly upregulated DEGs included genes like *fliF* (Log$_2$ FC=-9.86), *fliO* (Log$_2$ FC=-9.26) and *fliJ* (Log$_2$ FC=-9.26). The upregulation of these genes in *ΔyeaC* signify that the cells prioritize the expression of motility-based genes. They are known to aid the bacterium in adapting to environmental changes by modulating their expression [154]. Moreover, there is often a trade-off between motility and stress resistance. Studies have shown that hypermotile *E. coli* strains, which exhibit increased expression of motility genes, may have reduced expression of stress resistance genes, as was also observed in the downregulated DEGs in the WT. Hence, as opposed to WT, the deletion mutant prioritizes motility and chemotaxis to mitigate oxidative stress.

WT *vs. ΔydgH*

The *ydgH* gene demonstrated an upregulation after the 5h, with a log$_2$FC of approximately 3.7 as compared to 1 hour, indicating a 13-fold increase in expression. Hence, also in this case the 5h time point was chosen for further analysis. A total of 231 DEGs were obtained after filtering (p_{adj}≤0.1), resulting in 111 genes with a positive Log$_2$ FC and 120 genes with a negative Log$_2$ FC.

Top ten positive DEGs (Log$_2$ FC≥1.5) involved six stress response genes, one protein

transport gene, one membrane biosynthesis gene, one fatty acid oxidation gene and one HP-

encoding gene (**Figure 4C**). Significantly upregulated genes were *dsrB* (Log$_2$ FC=3.89),

encoding for a putative stress response protein [155] and *ghoS,* (Log$_2$ FC=3.65), an antitoxin

protein which prevents cell death [156]. Gene *fadM* (Log$_2$ FC=2.4), encoding a thioesterase,

is responsible for breaking down fatty acids into acetyl-CoA units, which can then be utilized

in energy production [157]. Another gene, *tatE* (Log$_2$ FC=2.69) encoding for a transporter, is

involved in transport of folded proteins and maintaining of cellular redox balance [158]. Gene

yoaH, a HP-encoding gene whose function was previously unclear from **Table 1**, might also

play a role in oxidative stress response and should be further investigated. The role of

lipopolysaccharide biosynthesis gene *ais* (Log$_2$ FC= 2.41) was, however, unclear.

In the deletion mutant (*ΔydgH*), the top ten negative DEGs included eight motility/chemotaxis

genes along with one sulphur metabolism and one urate transport gene (**Figure 4C**): *fliF* (Log$_2$

FC=-6.92) and *fliA* (Log$_2$ FC=-5.18) play a crucial role for motility (EcoCyc), *uacT* (Log$_2$ FC=-

6.22) is involved in urate transport, which has already been studied to reduce the effects of

oxidative stress in *E. coli* [159]. The role of gene *cysP* regarding oxygen stress, which is

involved in sulphur metabolism [160] is unclear at the moment and calls for further

investigation.

In WT*,* the upregulation of genes involved in response to stress, i.e. *ghoS* suggest that the

cells prioritize conservation of resources by entering a dormant state, i.e. formation of persister

cells [156]. Due to the absence of *ΔydgH* the cell seems to attempt to compensate for the loss

of oxidoreductive function by utilizing genes related to motility and chemotaxis (**Figure 4C**).

This increased expression could be an adaptive response to maintain survival under

continuous exposure to oxidative stress. Hence, similar to the previous case (WT *vs. ΔyeaC*),

more stress response genes were upregulated in WT while motility/chemotaxis genes were

upregulated in *ΔydgH*.

Figure 4. Comparative differential gene expression analysis in A. wild type (WT) *vs. ΔyhdN*, B. WT *vs. ΔyeaC* and C. WT *vs. ΔydgH* strains. Top ten positively and negatively expressed DEGs are shown. Each row represents a gene, and colour intensity represents the Log$_2$ Fold Change (Log2 FC), with red indicating upregulation in WT and green indicating upregulation in the mutant strains. Functional categories were based on information retrieved from Literature and EcoCyc.

Conclusion and Outlook

The functional characterization of hypothetical protein (HP)-encoding genes remains a critical

challenge, since the rapid accumulation of sequencing data has long surpassed the rate of

possible experimental characterization *in vivo* and *in vitro*. While there have been notable

advancements in methodologies for protein function prediction in the recent past, several

challenges persist, including the heavy reliance on pre-existing data inferred from model

organisms. Unfortunately, even in the best studied model organisms a significant amount of

protein encoding genes could so far not been annotated.

AI-driven approaches, such as machine learning (ML) algorithms, now offer powerful tools to

analyze vast omics datasets. In this study we employed a comprehensive approach,

integrating independent component analyses to decipher transcriptional regulatory networks,

deep learning for structural homolog prediction, and various other bioinformatic tools to

characterize HPs that have no functional information available in public databases. With this

methodology putative/probable functions for at least 64% of 95 hypothetical HPs encoding

genes from *E. coli* K-12 could be extrapolated, facilitating the subsequent functional validation

of these proteins through *in vitro* and *in vivo* experiments in the future. For the remaining 36%

HPs, however, more high-quality datasets including metabolomics and proteomics are

probably needed, as these genes are often seen to be expressed under very specific

environmental conditions from which only one dataset was available. Nevertheless, for most

proteins it is possible to derive at least some characteristics from their general sequence

properties, whether there is evidence of expression, their possible cellular location, details of

homologs, or whether they are predicted or demonstrated to be expressed as part of an

operon and/or regulon. Taking these information into account in databases will aid in the

improvement of using AI tools for annotation in the future.

In this study we furthermore provided an experimental feedback loop for three HP-encoding

genes to verify the *in silico* prediction. Gene *yhdN* ought to play a role in the cells' responding

to heat shock and nutrient limitation, which could be confirmed by comparing the growth

curves and transcriptional responses of wild type and *ΔyhdN* deletion mutant strains. For the

second candidate *yeaC, in silico* predictions suggested a role in aerobic respiration.

Experimental evidence supported this hypothesis by the upregulation of oxidative stress

response and methionine biosynthesis genes, hence it is likely involved in scavenging reactive

oxygen species. In contrast, the significant upregulation of "Motility"/Chemotaxis" genes in the

deletion mutant suggest that during the absence of *yeaC*, bacteria may enhance motility

pathways as an adaptive mechanism under oxidative stress. The third candidate *ydgH*, was

predicted *in silico* to play a role in oxidoreduction, which could also be confirmed by the *in vivo*

experiments showing its role in the protection against oxidative stress. Similar to *yeaC,* in the

mutant an upregulation of motility/chemotaxis genes indicate that in the absence of *ydgH*, the

cell may compensate for reduced oxidoreductive function. However, it is important to

acknowledge that our understanding still remains incomplete since the functional landscape

of proteins is vast and complex. Future studies and more omics datasets are needed to further

refine and expand upon our findings, exploring the broader implications of these proteins

under different environmental conditions.

In conclusion, this study represents a significant step forward in elucidating the functions of

previously uncharacterized HPs with no information in existing knowledge databases.

Leveraging AI-driven annotations and integrating them with experimental laboratory work will

link new functional roles of genes, discover previously unknown cellular and metabolic

processes and maybe even biotechnological potentials. In the future this approach might also

aid in the isolation of new and so far uncultured microbial species, i.e. microbial dark matter,

by harnessing meta-omics datasets and identifying genes critical for growth and adaptation,

offering clues to optimize culture conditions.

Methods

Identification and enumeration of hypothetical proteins (HPs)

For the comprehensive analysis of *Escherichia coli* K-12 (*E. coli*) substrains MG1655 and

BW25113, the complete genome sequences available in the NCBI RefSeq database under

the accession number GCF_000005845.2 and GCF_004355105.2 were used, respectively.

Ribosomal RNA (rRNA), transfer RNA (tRNA), non-coding RNA (ncRNA) were filtered out.

CDS based annotations were used to find hypothetical proteins (HPs). HPs were identified based on keywords like "Putative", "Putative uncharacterized", "Uncharacterized protein" and "DUF (domain of unknown function)-domain containing protein", "UPF" (unknown protein family) as well as proteins having a 'Protein Y…' pattern. Further information was extracted from the extensively curated databases EcoCyc version 26.1 [39], RegulonDB version 11.2 [11], EggNOG v6.0 [12] and UniProt release 2023_04 for proteome UP000000625 [1].

In order for a HP to be classified as a protein with no characterization (target group of this study), it had to fulfill all of the following criteria:

1. The **EcoCyc** protein summary showed the statement 'No information about this'. Additionally, the protein had to be marked as 'Uncharacterized'. Proteins which had the statement but a 'Partial' characterization in EcoCyc were further investigated using UniProt.

2. In **UniProt** the protein lacked any functional information in the 'Function' section and was assigned an annotation score of '1', and a 'protein existence' score of '4' signifying that the protein, although predicted to be expressed, lacked information based on experimental evidence or orthologs in closely related species. For the "Function" section, a score of '1' is the lowest rating in a five-point-scoring system and for the "protein existing" category a score of '4' out of '5', where '5' indicates a higher level of uncertainty regarding the protein's expression.

3. In the **RegulonDB** database it had the label "Weak" evidence for an annotation in the evidence section and no additional information in the 'Note' subsection.

4. In the **EggNOG** database the protein was either classified under the cluster of orthologous groups (COG) with a functional category 'S', where 'S' designates proteins belonging to an unknown function category or it lacked an ortholog which was displayed as 'No orthologs found'.

95 of the 4403 *E. coli* K-12 protein encoding genes fulfilled these criteria. All scripts for the filtering and enumeration of uncharacterized HPs can be found on Figshare

(*Notebooks>HP_curation*). In addition, the here identified HP-encoding genes were compared

with the ones identified by Ghatak et al. in 2019 [8] to check for their current status.

Transcriptomic data curation and processing

RNA-sequencing data from NCBI for analyzing the expression patterns of genes was collected

for *E. coli* K-12 substrains MG1655 and BW25113 using an adapted workflow for *Bacillus*

subtilis [32]. For substrain MG1655 1854 datasets and for substrain BW25113 662 datasets

were downloaded and processed. Raw FASTQ files were obtained using sratoolkit 3.0.1 [161],

and subsequent read trimming was performed using Trim Galore 0.6.5 [162] with the default

settings. Trimmed reads were then subjected to FastQC 0.11.9 [163]. Alignments to the

genomes were done using Bowtie 1.2.3 [164]. The read direction was determined using

RSeQC 3.0.1 [165] before generating read counts using featureCounts [166]. Finally, all

quality control metrics were compiled using MultiQC [167] and the expression compendium

was obtained as log-transformed Transcripts per Million (log-TPM) values. Quality metrics

were established to filter and ensure the selection of high-quality datasets with the following

modifications [168]: Briefly, the datasets were considered to be of high quality if they passed

the fastqc parameters 'per_base_sequence_quality', 'per_sequence_quality_scores',

'per_base_n_content', 'adapter_content' and had at least 500,000 reads mapped to coding

sequences from the reference genome. To preserve biological variability and prevent loss of

information, samples with no replicates were also retained in the dataset. Short non-coding

transcripts (<100 nucleotides) and extremely low-expression transcripts (Fragments Per

Kilobase per Million mapped reads < 10) were removed to reduce noise. 779 datasets for

substrain MG1655 and 135 datasets for substrain BW25113 were then used for further

analysis. All other information on the *E. coli* K-12 genes including their transcription units,

descriptions etc. were sourced from EcoCyc by utilizing the SmartTables feature [39].

Annotations for clusters of orthologous groups (COG) were obtained from EggNOG v6.0 [12]

including Gene Ontology [169]. The transcriptional regulatory network (TRN) comprising

information on genes and their governing regulators (i.e. transcription factors controlling the

expression of genes) were sourced from RegulonDB version 12.0 [11] and EcoCyc [39] as

described by Sastry et al. [32] In order to mitigate any batch effects which might be caused

due to the usage of diverse datasets, a reference condition (wild type *E. coli* K-12 substrain

MG1655 in M9 minimal medium facilitated with glucose and essential micronutrients under

non-stressful conditions) was used [30]. This was done by subtracting the mean expression

of two replicates of the chosen reference condition from the entire gene expression dataset

and calculating the log2-fold-change i.e. relative changes in expression. This normalization

method allowed for the comparison of gene expression differences relative to a common

baseline as previously demonstrated in Sastry et al. [30]. The normalized data was then used

as input for machine learning (*Data processing>Nextflow*).

Independent Component Analysis of Transcriptomic Data

Independent Component Analysis (ICA) developed by McCon et al. -named OptICA- was

applied to prevent both over-decomposition and under-decomposition of transcriptomic

datasets [27]. The application of ICA to the normalized *E. coli* K-12 substrain MG1655 RNAseq

data resulted in two matrices: the first matrix, denoted as the M matrix, encompassing the

robust independent components named iModulons, while the second matrix, termed the A

matrix, comprised the corresponding activities of iModulons over different growth conditions.

Additionally, a PyModulon package was used in conjunction with Jupyter notebooks to

characterize the iModulons [32]. The scripts containing the specific utilization of resources and

curation of iModulons can be found on Figshare (*OptICA* and *iModulon curation*). The

information from substrain MG1655 was then used to infer the iModulons and their member

genes for substrain BW25113, since only for the latter strain single-gene deletion mutants

were available for *in vitro* experimentation. The HP-encoding genes, the regulators governing

them, and their derived functions can be found in **Table 1.**

Possible functions for the 95 HP-encoding genes were further identified through regulators

governing their gene expression, as provided by RegulonDB and EcoCyc and/or by Gene

Ontology (GO) terms derived from co-regulating genes using the PANTHER (Protein ANalysis

THrough Evolutionary Relationships) tool [33]. Here, member genes of an iModulon (including

the gene of interest, i.e. the HP-encoding gene) were used as input to obtain GO terms to see

if any were overrepresented. Statistically significant GO categories with the default false

Discovery Rate (FDR)-adjusted p-value (<0.05) were manually identified for each HP

(*Supplementary Table 1*). These gene-associated functional categories can be found in **Table**

1 and **Figure 2**.

Metadata extraction

Metadata conditions for all datasets were retrieved either manually from the sequence read

archive (SRA), *via* a semi-automated approach [167] or by utilizing data from the Gene

Expression Omnibus Database using the GEOparse tool [170] to obtain experimental

information not available *via* the NCBI's Entrez Programming Utilities (E-utilities) [42]. As a

result, from 779 quality checked datasets for the substrain MG1655, metadata for 602 datasets

could be retrieved with confidence and termed as 'Explicit metadata'. 113 datasets were

termed as 'poor metadata', meaning that information on one or more experimental conditions

(such as media composition, temperature, pH, etc.) was missing or incomplete. Lastly,

datasets with 'unclear metadata' lacked all information. For the substrain BW25113, 128

datasets were 'Explicit' while seven were 'unclear' (*Notebooks>metadata_curation*).

Additionally, metadata condition for the highest expression of the HP- encoding genes against

the reference gene *frr* were extracted from the Tjaden dataset [44] in order to account for the

loss of datasets after quality filtering (also see below). The *frr* gene expression is typically

consistent across different conditions which was also observed when tested as reference

along with other genes like *rpoC*, *rpoD* for the uncharacterized HPs [this study, 171]. Hence,

gene *frr* was chosen as the reference gene serving as a control to validate the relative

expression levels of other genes. The extensive nature of this dataset ensured a broad

representation of gene expression profiles, in addition to the datasets procured for OptICA

analysis (*Notebooks> metadata_curation*).

Co-expression analysis based on public RNA-Seq data

To identify genes with expression patterns similar to those encoding HPs, a Spearman correlation analysis on the Tjaden dataset was performed [172]. This dataset comprises a comprehensive collection of 3.376 *E. coli* strain K-12 RNA-seq datasets, with expression levels quantified in transcripts per million (TPM) [44]. For inferring gene-gene correlations across the transcriptomic datasets for strain *E. coli* K-12, we identified the top ten co-expressed genes to the respective 95 HP-encoding gene and used this information to get an inference on their functions [10] **(Table 1)** (*Notebooks>gene_co-expression*).

Sub-Cellular Localization

For the purpose of determining the subcellular localizations of the 95 HP-encoding genes, BUSCA (Bologna Unified Subcellular Component Annotator) was employed [173]. This tool integrates various machine learning and bioinformatics tools [174–177] to analyze and recognize patterns in protein sequences that are indicative of specific subcellular localizations. The integrative webserver was used to obtain labels such as 'Nucleus', 'Periplasm/cytoplasm', 'Inner/outer membrane' or 'extracellular space'. All 95 HP-encoding genes with their predicted sub-cellular localizations can be found in **Table1**.

Protein-Protein Interactions (PPIs) Analysis

The STRING database (version 12.0) (https://string-db.org) was employed for the identification of PPIs of HP-encoding genes against the proteomic landscape of *E. coli*. The tool incorporates data from diverse origins such as available publications, empirical evidence, predictions derived from co-expression patterns, genomic context analysis, and curated pathway databases. A 'confidence score' for each inferred association was derived, quantitatively expressing the robustness of the predicted interaction [178]. In order to keep only the high-scoring associations, a cut-off of 0.7 was utilized [179]. All 95 HP-encoding genes and their interacting partner proteins are listed in **Table 1**.

Analysis of Gene Context

Shared functional relationships were analyzed using the genomic context of *E. coli* strain K-12 (*Supplementary Figure 1*), particularly the genes located upstream and downstream of the 95 genes of interest. A gene list comprising of gene names was used to create a SmartTable in the EcoCyc database to examine the local genomic neighbourhood and visualize the genes upstream and downstream, including their orientations [36]. (**Table 1)**.

Detection of Remote Protein Homology

In order to identify distantly homologous sequences, HP sequences were used as input for the HHblits tool (version 3.3.0) available in the MPI Bioinformatics Toolkit using a Hidden Markov Model (HMM) iterative sequence search [180]. The HHblits server (https://toolkit.tuebingen.mpg.de/tools/hhblits) takes a single sequence or a multiple sequence alignment as input and iteratively searches through the selected HMM databases [181,182]. HHblits performs sequence search and remote homology detection with higher sensitivity as compared to gold-standard databases like InterPro [180]. Homologous sequences were then searched against the default UniRef30 database [37]. Sequence based remote homologies identified through the HHblits tool are catalogued in **Table 1**.

Structure-Based Homology

For identifying structural homologs, we used the AlphaFold Protein Structure Database (AFDB) (https://cluster.foldseek.com/), a specialized computational database that includes structural models for approximately 215 million protein sequences. This platform employs deep learning algorithms to scan and match protein structures, enabling the discovery of structural similarities, even among proteins with low sequence identity [183]. We analyzed clusters to perform comparisons of HPs with all protein structures in the AFDB. UniProt

accession IDs were used as inputs to identify the cluster to which the HP belongs. The

homology-based function information for each of the 95 HP is listed in **Table 1**. A script to

extract the information can be found on Figshare (*Notebooks> HP_info_curation*).

Bacterial strains and growth conditions

E. coli K-12 substrain BW25113 was sourced from Horizon Discovery Ltd., United States.

Three single-gene knockout mutants from *E. coli* strain K-12 substrain BW25113 *ΔyhdN,*

ΔyeaC, and *ΔydgH* were obtained from the KEIO collection (NBRP, Japan). Glycerol stocks

were stored in an ultra-low temperature freezer (Thermo Fisher, United States) at −80°C until

further processing. Cells were cultured overnight on agar plates from glycerol stocks and

single colonies were picked from the agar plate and then used to Luria Bertani (LB) liquid

medium at 37°C [184].

Media and reagents

E. coli strains were cultured in LB broth (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl;

pH 7.0) at 37°C while shaking at 200 rpm [184]. M9 liquid medium for the pre-culture was

prepared using 42 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.5 mM NaCl, 18.7 mM NH_4Cl, 2 mM

$MgSO_4$, 0.1 mM $CaCl_2$ and 11 mM glucose, supplemented with 0.05% casamino acids to

ensure non-limiting conditions. For nitrogen limited M9 medium, a NH_4Cl concentration of 0.16

mM was used without the addition of casamino acids. For solid medium, 1.5% agar was

added. 30% H_2O_2 was sterile filtered through a 0.22 µm pore size filter to ensure sterility prior

to be used in the oxidative stress experiments. For all experiments with H_2O_2 the LB was

freshly prepared and kept in the dark until use. All chemicals were sourced from Sigma

Chemical Co. (St. Louis, MO, USA).

Measurement of Bacterial Growth Curves

Bacterial cultures were inoculated at a 1:100 dilution unless stated otherwise. Each experiment included triplicates. All optical density values (OD_{600}) were measured *via* a cell density meter (Ultrospec 10, biochrom). OD_{600} was measured every 2h for heat shock in LB and M9 or M9 modified medium and every 1h for oxidative stress experiments. Growth curves were generated for each set of experiments including the wild type and mutant strain in stressed and non-stressed conditions (controls).

Heat stress

A single colony (wild type and Δ*yhdN*) was picked from either LB or M9 agar plates and inoculated into 5 mL LB broth for overnight cultivation at 37°C and shaken with 200 rpm. The overnight culture was then used to inoculate fresh LB medium at a 1:100 dilution, which was then cultured for ~2-3h at 37°C and 200 rpm until an OD_{600} of 0.4 was reached. This was repeated with the M9 liquid medium (preculture), at 37°C to ensure non limiting conditions. For the nitrogen limitation assay, medium was prepared using 0.16mM NH_4Cl. To study the effects of heat shock on bacterial growth, the culture was grown to mid-log phase (OD_{600} 0.4), diluted to OD_{600} 0.04 and then heat shocked for 7 minutes in a shaker-incubator at 50°C. After treatment, the bacterial suspension was incubated for 12 h at 37°C. Samples were measured every 2h for 12h.

Oxidative stress

For studying the effects of oxidative stress, a single colony (wild type, Δ*yeaC* and Δ*ydgH*) was picked from the LB agar plate and inoculated into 5 mL LB for overnight cultivation at 37°C and shaken with 200 rpm. The overnight culture was then used to inoculate fresh LB (dilution of 1:100) and cultured at 37°C and 200 rpm until an OD_{600} of 0.2 was reached. An OD of 0.2 was used since at this stage the cells were actively dividing, making them susceptible to the effects of oxidative stress, allowing for better characterization and analysis of the stress response[137]. H_2O_2 was added at a final concentration of 2.5 mM. After treatment, the bacterial suspension was incubated for 5h at 37°C and the OD_{600} was measured hourly.

RNA Sequencing and data processing

2 ml of cell suspension from the heat shock samples was collected after 2h and 6h of growth.

For the oxidative stress experiments, samples were collected after 1h and 5h of growth. The

medium was discarded, cell pellets were washed with Phosphate Buffer Saline and mixed with

DNA/RNA shield™ (Zymo Research Corp., Freiburg, Germany). All samples were processed

according to the manufacturer's instructions. Samples were preserved at -20°C until RNA

extraction. RNA was extracted using the Direct-zol RNA Miniprep Plus Kit (Zymo Research

Corp., Freiburg, Germany) according to the manufacturer's instructions. Total RNA was eluted

from columns with 100 µL nuclease-free water and quantified using the Qubit RNA Broad

Range Assay Kit (Thermo Fisher Scientific Inc., Waltham, MA, USA). The quality was

assessed by running an RNA-nano chip on an Agilent bioanalyzer (Agilent Technologies,

Waldbronn, Germany). Samples with an RNA integrity score ≥8 were used for further

processing [185]. The rRNA was removed from total RNA preparations using the NEBNext

rRNA Depletion (Bacteria) kit (New England Biolabs, Frankfurt, Germany). Paired-end library

preparation was done using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina

(New England Biolabs, Frankfurt, Germany) following the manufacturer's instructions with an

average insert size of 200 bp. Libraries were run on a NextSeq 550™ (Illumina, San Diego,

CA, United States).

The RNAflow pipeline [186] was implemented using a conda environment. Sequence reads

were checked for quality using FastQC v0.11.9 [187]. Raw reads were processed using fastp

v0.20.0 [188], which employs a sliding window approach for quality trimming and adapter

clipping, with the settings: a -5 to -3 cut for front and tail, a default window of 4 and mean

quality threshold of 20. Trimmed reads shorter than 15 bps were discarded. The remaining

reads were mapped onto the *E. coli* K-12 reference genome (NZ_CP009273.1) using HISAT

v2.1.0 [189]. The generated output was converted into a BAM format using samtools-1.20

[190], followed by the processes of sorting and indexing. A gene transfer format (GTF)

annotation file for *E. coli* substrain BW25113 was used to filter for gene and pseudogene

features. FeatureCounts v2.0.1 [191] was used to quantify the mapped reads based on the

gene annotations. Processed datasets were summarized in a MultiQC v1.9 [167] generated

file. were normalized by the DESeq2 module (v1.28.0) in R (version 4.3.2) using a

variance stabilizing transformation [192].

Comparative gene expression analysis of wild type and isogenic mutant strains

Sequencing reads were obtained from the transcriptomic datasets of wild-type (WT), *ΔyhdN*,

ΔyeaC, and *ΔydgH* mutants, respectively. Non-stressed control and heat shocked samples

from WT and *ΔyhdN* at the 6h timepoint and non-stressed control and oxidative stress samples

from WT, *ΔyeaC,* and *ΔydgH* at the 5h timepoint were used to identify up- or downregulated

genes. DESeq2 [192] was utilized to examine expression differences of genes in the WT and

mutant strains under both control and stress conditions. To focus on the specific effect of

stress on gene expression rather than the difference of WT *vs.* deletion mutants, read counts

under control conditions were subtracted from the read counts under stress condition [193].

Genes with an adjusted P-value (Padj) of 0.1 or less were classified as differentially expressed

to avoid false positives. A cut-off $\log_2$ fold change ($\log_2$FC) of 1.5 was chosen for further

analysis [194]. A $\log_2$FC of 1.5 means that a gene was 2.83 times more expressed in one

strain *vs.* the other, a $\log_2$FC of 2 means that a gene was 4 times more expressed and so on.

Significant positive $\log_2$FC indicated genes that were upregulated in WT relative to the mutant,

while negative values showed an upregulation in the mutant *vs.* WT. To determine the

functional categories of these genes, the EcoCyc database was used [36]. Information of the

differentially expressed genes can be found on Figshare (*DEGs>Table1*).

Data availability

All code and scripts used in this study are available on Figshare. These resources can be

accessed at https://figshare.com/s/0ede175c510cf201e7c2.

Raw sequencing data that supports the findings of this study have been deposited in the Sequence Read Archive with the BioProject ID: PRJNA1209969.

Competing interests

The authors declare that they have no conflict of interest.

Funding

This work was supported by the Karlsruhe Institute of Technology and the Helmholtz Society [POF4; 5207.0004.0012].

Authors' contributions

S.C, H.A. and Z.A did the data analyses, S.C. did the *in vitro* experiments, S.C and A.K.K. wrote the manuscript, A.K.K. provided the funding.

Acknowledgements

The authors acknowledge the support by the state of Baden-Württemberg through bwHPC. We thank Dr. Gunnar Sturm for technical support and insightful discussions and David Thiele for library preparation assistance.

References

1. Consortium TU. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res. 2023;51:D523–31; doi:10.1093/nar/gkac1052

2. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al. InterPro in 2022. Nucleic Acids Res. 2023;51:D418–27; doi:10.1093/nar/gkac993

3. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al. Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49:D412–9; doi:10.1093/nar/gkaa913

4. Sigrist CJA, de Castro E, Cerutti L, Cuche BA, Hulo N, Bridge A, et al. New and continuing developments at PROSITE. Nucleic Acids Res. 2013;41:D344–7; doi:10.1093/nar/gks1067

5. Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in 2020. Nucleic Acids Res. 2021;49:D458–60; doi:10.1093/nar/gkaa937

886   6. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE:
887   the conserved domain database in 2020. Nucleic Acids Res. 2020;48:D265–8;
888   doi:10.1093/nar/gkz991
889

890   7. Durairaj J, Waterhouse AM, Mets T, Brodiazhenko T, Abdullah M, Studer G, et al.
891   Uncovering new families and folds in the natural protein universe. Nature. 2023;622:646–53;
892   doi:10.1038/s41586-023-06622-3
893

894   8. Ghatak S, King ZA, Sastry A, Palsson BO. The y-ome defines the 35% of *Escherichia coli*
895   genes that lack experimental evidence of function. Nucleic Acids Res. 2019;47:2446–54;
896   doi:10.1093/nar/gkz030
897

898   9. Ardern Z, Chakraborty S, Lenk F, Kaster A-K. Elucidating the functional roles of prokaryotic
899   proteins using big data and artificial intelligence. FEMS Microbiol Rev. 2023;47;
900   doi:10.1093/femsre/fuad003
901

902   10. Keseler IM, Gama-Castro S, Mackie A, Billington R, Bonavides-Martínez C, Caspi R, et
903   al. The EcoCyc Database in 2021. Front Microbiol. 2021;12; doi:10.3389/fmicb.2021.711077
904

905   11. Salgado H, Gama-Castro S, Lara P, Mejia-Almonte C, Alarcón-Carranza G, López-Almazo
906   AG, et al. RegulonDB v12.0: a comprehensive resource of transcriptional regulation in *E. coli*
907   K-12. Nucleic Acids Res. 2024;52:D255–64; doi:10.1093/nar/gkad1072
908

909   12. Hernández-Plaza A, Szklarczyk D, Botas J, Cantalapiedra CP, Giner-Lamia J, Mende DR,
910   et al. eggNOG 6.0: enabling comparative genomics across 12 535 organisms. Nucleic Acids
911   Res. 2023;51:D389–94; doi:10.1093/nar/gkac1022
912

913   13. Mills CL, Beuning PJ, Ondrechen MJ. Biochemical functional predictions for protein
914   structures of unknown or uncertain function. Comput Struct Biotechnol J. 2015;13:182–91;
915   doi:10.1016/j.csbj.2015.02.003
916

917   14. Zhang C, Freddolino PL, Zhang Y. COFACTOR: improved protein function prediction by
918   combining structure, sequence and protein–protein interaction information. Nucleic Acids Res.
919   2017;45:W291–9; doi:10.1093/nar/gkx366
920

921   15. Makrodimitris S, van Ham RCHJ, Reinders MJT. Improving protein function prediction
922   using protein sequence and GO-term similarities. Bioinformatics. 2019;35:1116–24. Available
923   from: doi:10.1093/bioinformatics/bty751
924

925   16. Saha S, Chatterjee P, Basu S, Nasipuri M, Plewczynski D. FunPred 3.0: improved protein
926   function prediction using protein interaction network. PeerJ. 2019;7. doi: 10.7717/peerj.6830
927

928   17. Varadi M, Tsenkov M, Velankar S. Challenges in bridging the gap between protein
929   structure prediction and functional interpretation. Proteins: Structure, Function, and
930   Bioinformatics. 2023; doi:10.1002/prot.26614
931

932   18. Vincent AT. Bacterial hypothetical proteins may be of functional interest. Frontiers in
933   Bacteriology. 2024;3; doi:10.3389/fbrio.2024.1334712
934

935   19. Jeffery CJ. Current successes and remaining challenges in protein function prediction.
936   Frontiers in Bioinformatics. 2023;3; doi:10.3389/fbinf.2023.1222182
937

20. Pavlopoulos GA, Baltoumas FA, Liu S, Selvitopi O, Camargo AP, Nayfach S, et al.
Unraveling the functional dark matter through global metagenomics. Nature. 2023;622:594–
602; doi:10.1038/s41586-023-06583-7

21. Escudeiro P, Henry CS, Dias RPM. Functional characterization of prokaryotic dark matter:
the road so far and what lies ahead. Curr Res Microb Sci. 2022;3:100159; doi:
10.1016/j.crmicr.2022.100159

22. da Costa WLO, Araújo CL de A, Dias LM, Pereira LC de S, Alves JTC, Araújo FA, et al.
Functional annotation of hypothetical proteins from the *Exiguobacterium antarcticum* strain B7
reveals proteins involved in adaptation to extreme environments, including high arsenic
resistance. PLoS One. 2018;13:e0198965; doi:10.1371/journal.pone.0198965

23. Grünberger F, Knüppel R, Jüttner M, Fenk M, Borst A, Reichelt R, et al. Exploring
prokaryotic transcription, operon structures, rRNA maturation and modifications using
Nanopore-based native RNA sequencing. bioRxiv. 2020;2019.12.18.880849; doi:
10.1101/2019.12.18.880849

24. Dall'Alba G, Casa PL, Abreu FP de, Notari DL, de Avila e Silva S. A Survey of Biological
Data in a Big Data Perspective. Big Data. 2022;10:279–97; doi:10.1089/big.2020.0383

25. Chen J, Gu Z, Lai L, Pei J. In silico protein function prediction: the rise of machine learning-
based approaches. 2023;3:487–510; doi:10.1515/mr-2023-0038

26. Alteri CJ, Lindner JR, Reiss DJ, Smith SN, Mobley HLT. The broadly conserved regulator
PhoP links pathogen virulence and membrane potential in *Escherichia coli*. Mol Microbiol.
2011;82:145–63; doi:10.1111/j.1365-2958.2011.07804.x

27. Mcconn JL, Lamoureux CR, Poudel S, Palsson BO, Sastry A V. Optimal dimensionality
selection for independent component analysis of transcriptomic data; doi:10.1186/s12859-
021-04497-7
28. Yu J, Li M, Wang J, Hamushan M, Jiang F, Wang B, et al. Identification of *Staphylococcus*
aureus virulence-modulating RNA from transcriptomics data with machine learning. Virulence.
2023;14:2228657; doi:10.1080/21505594.2023.2228657

29. Comon P. Independent component analysis, A new concept? Signal Processing.
1994;36:287–314; doi:10.1016/0165-1684(94)90029-9

30. Sastry A V, Gao Y, Szubin R, Hefner Y, Xu S, Kim D, et al. The *Escherichia coli*
transcriptome mostly consists of independently regulated modules. Nat Commun.
2019;10:5536; doi:10.1038/s41467-019-13483-w

31. Sastry A V, Poudel S, Rychel K, Yoo R, Lamoureux CR, Chauhan S, et al. Title Mining all
publicly available expression data to compute dynamic microbial transcriptional regulatory
networks Authors; doi:10.1101/2021.07.01.450581

32. Rychel K, Sastry A V., Palsson BO. Machine learning uncovers independently regulated
modules in the *Bacillus subtilis* transcriptome. Nat Commun. 2020;11. doi: 10.1038/s41467-
020-20153-9

33. Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou L-P, Mi H. PANTHER:
Making genome-scale phylogenetics accessible to all. Protein Science. 2022;31:8–22;
doi:10.1002/pro.4218
991

34. Savojardo C, Martelli PL, Fariselli P, Profiti G, Casadio R. BUSCA: an integrative web server to predict subcellular localization of proteins. Nucleic Acids Res. 2018;46:W459–66; doi:10.1093/nar/gky320

35. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51:D638–46; doi: 10.1093/nar/gkac1000

36. D KP, Suzanne P, Ron C, Anamika K, Markus K, E MP, et al. The EcoCyc Database (2023). EcoSal Plus. 2023;11:eesp-0002-2023; doi:10.1128/ecosalplus.esp-0002-2023

37. Gabler F, Nam S-Z, Till S, Mirdita M, Steinegger M, Söding J, et al. Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. Curr Protoc Bioinformatics. 2020;72:e108; doi:10.1002/cpbi.108

38. Barrio-Hernandez I, Yeo J, Jänes J, Mirdita M, Gilchrist CLM, Wein T, et al. Clustering predicted structures at the scale of the known protein universe. Nature. 2023;622:637–45; doi:10.1038/s41586-023-06510-w

39. Karp PD, Billington R, Caspi R, Fulcher CA, Latendresse M, Kothari A, et al. The BioCyc collection of microbial genomes and metabolic pathways. Brief Bioinform. 2019;20:1085–93; doi:10.1093/bib/bbx085

40. Choudhary S. pysradb: A Python package to query next-generation sequencing metadata and data from NCBI Sequence Read Archive. F1000Res. 2019;8:532; doi: 10.12688/f1000research.18676.1

41. Gumienny R. GEOparse. Rev.0a257463. 2015. Available from: https://geoparse.readthedocs.io/en/latest/

42. Liu M, Durfee T, Cabrera JE, Zhao K, Jin DJ, Blattner FR. Global transcriptional programs reveal a carbon source foraging strategy by *Escherichia coli*. J Biol Chem. 2005;280:15921–7; doi: 10.1074/jbc.m414050200

43. Tjaden B. *Escherichia coli* transcriptome assembly from a compendium of RNA-seq data sets. RNA Biol. 2023;20:77–84; doi:10.1080/15476286.2023.2189331

44. Rawangkan A, Yosboonruang A, Kiddee A, Siriphap A, Pook-In G, Praphasawat R, et al. Restoring Ampicillin Sensitivity in Multidrug-Resistant *Escherichia coli* Following Treatment in Combination with Coffee Pulp Extracts. J Microbiol Biotechnol. 2023;33:1–10; doi:10.4014/jmb.2304.04051

45. Manat G, El Ghachi M, Auger R, Baouche K, Olatunji S, Kerff F, et al. Membrane Topology and Biochemical Characterization of the *Escherichia coli* BacA Undecaprenyl-Pyrophosphate Phosphatase. PLoS One. 2015;10:e0142870; doi:10.1371/journal.pone.0142870

46. Hajra D, Nair AV, Chakravortty D. An elegant nano-injection machinery for sabotaging the host: Role of Type III secretion system in virulence of different human and animal pathogenic bacteria. Phys Life Rev. 2021;38:25–54; doi:10.1016/j.plrev.2021.05.007

47. Prakash A, Dutta D. Bicyclomycin generates ROS and blocks cell division in *Escherichia coli*. PLoS One. 2024;19:e0293858; doi:10.1371/journal.pone.0293858

48. Ogawa A, Kojima F, Miyake Y, Yoshimura M, Ishijima N, Iyoda S, et al. Regulation of constant cell elongation and Sfm pili synthesis in *Escherichia coli* via two active forms of FimZ orphan response regulator. Genes to Cells. 2022;27:657–74; doi:10.1111/gtc.12982

49. Lamark T, Røkenes TP, McDougall J, Strøm AR. The complex bet promoters of *Escherichia coli*: regulation by oxygen (ArcA), choline (BetI), and osmotic stress. J Bacteriol. 1996;178(6):1655–62; doi:10.1128/jb.178.6.1655-1662.1996.

50. Kambampati R, Lauhon CT. Evidence for the Transfer of Sulfane Sulfur from IscS to ThiI during the in Vitro Biosynthesis of 4-Thiouridine in *Escherichia coli* tRNA*. Journal of Biological Chemistry. 2000;275:10727–30; doi:10.1074/jbc.275.15.10727

51. Laura F-G, Xinyu G, Joy K, Sooyeon S, E BM, Rodolfo G-C, et al. Single-cell analysis reveals that cryptic prophage protease LfgB protects *Escherichia coli* during oxidative stress by cleaving antitoxin MqsA. Microbiol Spectr. 2024;12:e03471-23; doi:10.1128/spectrum.03471-23

52. Zorraquino-Salvo V, Quinones-Soto S, Kim M, Rai N, Tagkopoulos I. Deciphering the genetic and transcriptional basis of cross-stress responses in *Escherichia coli* under complex evolutionary scenarios. bioRxiv. 2014;010595; doi:

53. May T, Okabe S. Enterobactin is required for biofilm development in reduced-genome *Escherichia coli*. Environ Microbiol. 2011;13:3149–62; doi:10.1111/j.1462-2920.2011.02607.x

54. Saier Jr MH, Reddy VS, Tamang DG, Västermark Å. The Transporter Classification Database. Nucleic Acids Res. 2014;42:D251–8; doi:10.1093/nar/gkt1097

55. Muto A, Goto S, Kurita D, Ushida C, Soma A, Himeno H. A leaderless mRNA including tRNA-like sequence encodes a small peptide that regulates the expression of GcvB small RNA in *Escherichia coli*. The Journal of Biochemistry. 2022;171:459–65; doi:10.1093/jb/mvac007

56. Njenga R, Boele J, Öztürk Y, Koch H-G. Coping with stress: How bacteria fine-tune protein synthesis and protein transport. Journal of Biological Chemistry. 2023;299:105163; doi:10.1016/j.jbc.2023.105163

57. Rodríguez-Ochoa JI, Fragoso-Jiménez JC, Flores N, et al. Global transcriptomic response of *Escherichia coli* to p-coumaric acid. Microb Cell Fact. 2022;21:148; doi:10.1186/s12934-022-01874-6.

58. Bak G, Lee J, Suk S, Kim D, Lee J, Kim K, et al. Identification of novel sRNAs involved in biofilm formation, motility, and fimbriae formation in *Escherichia coli*. Sci Rep. 2015;5:15287; doi:10.1038/srep15287

59. Duarte-Velázquez I, Mora J, Ramírez-Prado J, Aguillón-Bárcenas A, Tornero-Gutiérrez F, Cordero-Loreto E, et al. *Escherichia coli* transcription factors of unknown function: sequence features and possible evolutionary relationships. PeerJ. 2022;10:e13772; doi:10.7717/peerj.13772

60. Saier Jr MH, Reddy VS, Tsu B V, Ahmed MS, Li C, Moreno-Hagelsieb G. The Transporter Classification Database (TCDB): recent advances. Nucleic Acids Res. 2016;44:D372–9; doi:10.1093/nar/gkv1103

61. Su C-C, Rutherford DJ, Yu EW. Characterization of the multidrug efflux regulator AcrR from *Escherichia coli*. Biochem Biophys Res Commun. 2007;361:85–90; doi:10.1016/j.bbrc.2007.06.175

62. Börner J, Friedrich T, Bartkuhn M, Klug G. Ribonuclease E strongly impacts bacterial adaptation to different growth conditions. RNA Biol. 2023;20:120–35; doi:10.1080/15476286.2023.2195733

63. A JN, Melissa F, E FS. Disruption of trehalose periplasmic recycling dysregulates cAMP-CRP signaling in *Escherichia coli* during stationary phase. J Bacteriol. 2023;205:e00292-23; doi:10.1128/jb.00292-23

64. Wang M, Fang K, Hong SMC, Kim I, Jang I-S, Hong SH. Medium chain unsaturated fatty acid ethyl esters inhibit persister formation of *Escherichia coli* via antitoxin HipB. Appl Microbiol Biotechnol. 2018;102:8511–24; doi:10.1007/s00253-018-9271-3

65. Muenter MM, Aiken A, Akanji JO, Baig S, Bellou S, Carlson A, et al. The response of *Escherichia coli* to the alkylating agents chloroacetaldehyde and styrene oxide. Mutation Research/Genetic Toxicology and Environmental Mutagenesis. 2019;840:1–10; doi:10.1016/j.mrgentox.2019.02.001

66. Hideji Y, Yasushi M, Shou F, Akiko S, Masami U, Akira W. YqjD Is an Inner Membrane Protein Associated with Stationary-Phase Ribosomes in *Escherichia coli*. J Bacteriol . 2012;194:4178–83; doi:10.1128/jb.00396-12

67. Hemm MR, Paul BJ, Schneider TD, Storz G, Rudd KE. Small membrane proteins found by comparative genomics and ribosome binding site models. Mol Microbiol. 2008;70:1487–501; doi:10.1111/j.1365-2958.2008.06495.x

68. Skjold AC, Ezekiel DH. Analysis of lambda insertions in the fucose utilization region of *Escherichia coli* K-12: use of lambda *fuc* and lambda *argA* transducing bacteriophages to partially order the fucose utilization genes. J Bacteriol. 1982;152:120–5; doi:10.1016/j.mrgentox.2019.02.001

69. Manuel C, Teresa ZM, Blas B, Gonzalo D-R, F JJ, Andrés VJ, et al. Anaerobic Catabolism of Aromatic Compounds: a Genetic and Genomic View. Microbiology and Molecular Biology Reviews. 2009;73:71–133; 10.1128/mmbr.00021-08

70. Arnim W, A KS, Kirsten J. Time-Dependent Proteome Alterations under Osmotic Stress during Aerobic and Anaerobic Growth in *Escherichia coli*. J Bacteriol. 2006;188:7165–75; doi:10.1128/jb.00508-06

71. Teufel R, Mascaraque V, Ismail W, Voss M, Perera J, Eisenreich W, et al. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. Proc Natl Acad Sci U S A. 2010;107(32):14390–5; doi: 10.1073/pnas.1005399107.

72. Tenorio E, Saeki T, Fujita K, Kitakawa M, Baba T, Mori H, et al. Systematic characterization of *Escherichia coli* genes/ORFs affecting biofilm formation. FEMS Microbiol Lett. 2003;225(1):107–14; doi:10.1016/S0378-1097(03)00507-X.

73. Rodríguez-Montelongo L, Volentini SI, Farías RN, Massa EM, Rapisarda VA. The Cu(II)-reductase NADH dehydrogenase-2 of *Escherichia coli* improves the bacterial growth in extreme copper concentrations and increases the resistance to the damage caused by copper and hydroperoxide. Arch Biochem Biophys. 2006;451:1–7; doi:10.1016/j.abb.2006.04.019

74. Harwani D. Regulation of Gene Expression: Cryptic β-glucoside (bgl) Operon of *Escherichia coli* as a Paradigm. Brazilian Journal of Microbiology. 2014; doi:10.1590/S1517-83822014000400003

75. Anne P, K BI, Aurora L-N, F GJ, Valérie de C-L, D HA. FolX and FolM Are Essential for Tetrahydromonapterin Synthesis in *Escherichia coli* and Pseudomonas aeruginosa. J Bacteriol. 2010;192:475–82; doi:10.1128/jb.01198-09

76. Kotaka Y, Hashimoto M, Lee K, Kato J. Mutations identified in engineered *Escherichia coli* with a reduced genome. Front Microbiol. 2023;14; doi:10.3389/fmicb.2023.1189877

77. Doukyu N, Taguchi K. Involvement of catalase and superoxide dismutase in hydrophobic organic solvent tolerance of *Escherichia coli*. AMB Express. 2021;11:97; doi:10.1186/s13568-021-01258-w

78. Pinske C. The Ferredoxin-Like Proteins HydN and YsaA Enhance Redox Dye-Linked Activity of the Formate Dehydrogenase H Component of the Formate Hydrogenlyase Complex. Front Microbiol. 2018;9; doi:10.3389/fmicb.2018.01238

79. Panta PR, Doerrler WT. A link between pH homeostasis and colistin resistance in bacteria. Sci Rep. 2021;11:13230; doi:10.1038/s41598-021-92718-7

80. Vasilyev N, Liu MMJ, Epshtein V, Shamovsky I, Nudler E. General Transcription Factor from *E. coli* with a Distinct Mechanism of Action. bioRxiv. 2023;2023.09.17.558134; doi: 10.1101/2023.09.17.558134

81. Nasreen M, Purushothaman Nair R, Mcewan A, Kappler U. The Peptide Methionine Sulfoxide Reductase (MsrAB) of Haemophilus influenzae Repairs Oxidatively Damaged Outer Membrane and Periplasmic Proteins Involved in Nutrient Acquisition and Virulence. Antioxidants. 2022;11:1557; doi:10.3390/antiox11081557

82. Yuan J, Jin F, Glatter T, Sourjik V. Osmosensing by the bacterial PhoQ/PhoP two-component system. Proceedings of the National Academy of Sciences. 2017;114:E10792–8; doi:10.1073/pnas.1717272114

83. Steinsiek S, Stagge S, Bettenbrock K. Analysis of *Escherichia coli* mutants with a linear respiratory chain. PLoS One. 2014;9(1):e87307; doi:10.1371/journal.pone.0087307.

84. Akinori K, Hiroyuki T, Ryutaro U. Molecular Characterization of the PhoP-PhoQ Two-Component System in *Escherichia coli* K-12: Identification of Extracellular Mg2+-Responsive Promoters. J Bacteriol. 1999;181:5516–20; doi:10.1128/jb.181.17.5516-5520.1999

85. Podlesek Z, Žgur Bertok D. The DNA Damage Inducible SOS Response Is a Key Player in the Generation of Bacterial Persister Cells and Population Wide Tolerance. Front Microbiol .2020;11; doi:10.3389/fmicb.2020.01785

86. Bitoun JP, Wu G, Ding H. *Escherichia coli* FtnA acts as an iron buffer for re-assembly of iron–sulfur clusters in response to hydrogen peroxide stress. BioMetals. 2008;21:693–703; doi:10.1007/s10534-008-9154-7

87. C BD, J TR. Small Multidrug Resistance Protein EmrE Reduces Host pH and Osmotic Tolerance to Metabolic Quaternary Cation Osmoprotectants. J Bacteriol. 2012;194:5941–8; doi:10.1128/jb.00666-12

88. M FE, E EK, N GL, R GE, M MJ, C MA, et al. YfdW and YfdU Are Required for Oxalate-Induced Acid Tolerance in *Escherichia coli* K-12. J Bacteriol. 2013;195:1446–55; doi:10.1128/jb.01936-12

89. Ralph ET, Guest JR, Green J. Altering the anaerobic transcription factor FNR confers a hemolytic phenotype on *Escherichia coli* K12. Proceedings of the National Academy of Sciences. 1998;95:10449–52; doi:10.1073/pnas.95.18.10449

90. Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM, et al. Cryptic prophages help bacteria cope with adverse environments. Nat Commun. 2010;1:147; doi:10.1038/ncomms1146

91. Barabote RD, Rendulic S, Schuster SC, Saier MH. Comprehensive analysis of transport proteins encoded within the genome of *Bdellovibrio bacteriovorus*. Genomics . 2007;90:424–46; doi:10.1016/j.ygeno.2007.06.002

92. Wang J, Ma W, Wang X. Insights into the structure of *Escherichia coli* outer membrane as the target for engineering microbial cell factories. Microb Cell Fact. 2021;20:73; doi:10.1186/s12934-021-01565-8

93. Gurmu D, Lu J, Johnson KA, Nordlund P, Holmgren A, Erlandsen H. The crystal structure of the protein YhaK from *Escherichia coli* reveals a new subclass of redox sensitive enterobacterial bicupins. Proteins: Structure, Function, and Bioinformatics. 2009;74:18–31; doi:10.1002/prot.22128

94. Outten CE, Outten FW, O'Halloran T V. DNA Distortion Mechanism for Transcriptional Activation by ZntR, a Zn(II)-responsive MerR Homologue in *Escherichia coli*. Journal of Biological Chemistry. 1999;274:37517–24; doi:10.1074/jbc.274.53.37517

95. Akbar S, Bhakta S, Sengupta J. Structural insights into the interplay of protein biogenesis factors with the 70S ribosome. Structure. 2021;29:755-767.e4; doi:10.1016/j.str.2021.03.005

96. K VDT, J TL, A CK, L SP, Sima SF. Characterization of the *Escherichia coli* AaeAB Efflux Pump: a Metabolic Relief Valve? J Bacteriol. 2004;186:7196–204; doi:10.1128/jb.186.21.7196-7204.2004

97. Bak G, Lee J, Suk S, Kim D, Young Lee J, Kim K, et al. Identification of novel sRNAs involved in biofilm formation, motility and fimbriae formation in *Escherichia coli*. Sci Rep. 2015;5:15287; doi:10.1038/srep15287

98. P DM, Chi-Fang W, Liang W, J VJ, E BW. DNA Microarray-Based Identification of Genes Controlled by Autoinducer 2-Stimulated Quorum Sensing in *Escherichia coli*. J Bacteriol. 2001;183:5239–47; doi:10.1128/jb.183.18.5239-5247.2001

99. Filenko N, Spiro S, Browning DF, Squire D, Overton TW, Cole J, Constantinidou C. The NsrR regulon of *Escherichia coli* K-12 includes genes encoding the hybrid cluster protein and the periplasmic, respiratory nitrite reductase. J Bacteriol. 2007;189:1203–9; doi:10.1128/jb.00080-07.

100. Khil PP, Camerini-Otero RD. Over 1000 genes are involved in the DNA damage response of *Escherichia coli*. Mol Microbiol. 2002;44:89–105; doi:10.1046/j.1365-2958.2002.02878.x

101. Erol I, Jeong K-C, Baumler DJ, Vykhodets B, Ho Choi S, Kaspar CW. H-NS controls metabolism and stress tolerance in *Escherichia coli* O157:H7 that influence mouse passage. BMC Microbiol. 2006;6:72; doi:10.1186/1471-2180-6-72

102. Zheng L. Catalytic mechanism and function evolvement studies of phosphatases within the haloacid dehalogenase superfamily (HADSF) [dissertation]. Albuquerque (NM): University of New Mexico; 2013. Available from: https://digitalrepository.unm.edu/chem_etds/31.

103. Born TL, Blanchard JS. Structure/function studies on enzymes in the diaminopimelate pathway of bacterial cell wall biosynthesis. Curr Opin Chem Biol. 1999;3:607–13; doi:10.1016/S1367-5931(99)00016-2

104. Hor L, Dobson RCJ, Downton MT, Wagner J, Hutton CA, Perugini MA. Dimerization of Bacterial Diaminopimelate Epimerase Is Essential for Catalysis. J Biol Chem . 2013;288:9238–48

105. Iyer LM, Leipe DD, Koonin E V, Aravind L. Evolutionary history and higher order classification of AAA+ ATPases. J Struct Biol. 2004;146:11–31; doi:

106. Luttringer F, Mulliez E, Dublet B, Lemaire D, Fontecave M. The Zn center of the anaerobic ribonucleotide reductase from *E. coli*. JBIC Journal of Biological Inorganic Chemistry . 2009;14:923–33; doi:10.1007/s00775-009-0505-9

107. Panta PR, Doerrler WT. A link between pH homeostasis and colistin resistance in bacteria. Sci Rep. 2021;11:13230; doi:10.1038/s41598-021-92718-7

108. Dapa T, Fleurier S, Bredeche M-F, Matic I. The SOS and RpoS Regulons Contribute to Bacterial Cell Robustness to Genotoxic Stress by Synergistically Regulating DNA Polymerase Pol II. Genetics. 2017;206:1349–60; doi:10.1534/genetics.116.199471

109. Passariello C, Forleo C, Micheli V, Schippa S, Leone R, Mangani S, et al. Biochemical characterization of the class B acid phosphatase (AphA) of *Escherichia coli* MG1655. Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics. 2006;1764:13–9; doi:10.1016/j.bbapap.2005.08.028

110. Chen SH, Byrne RT, Wood EA, Cox MM. *Escherichia coli rad* gene: a novel function involved in radiation resistance and double-strand break repair. Mol Microbiol. 2015;95:754–68; doi:10.1111/mmi.12885

111. Chuang SE, Blattner FR. Characterization of twenty-six new heat shock genes of *Escherichia coli*. J Bacteriol. 1993;175:5242–52; doi:10.1128/jb.175.16.5242-5252.1993

112. Manganelli R, Gennaro ML. Protecting from Envelope Stress: Variations on the Phage-Shock-Protein Theme. Trends Microbiol. 2017;25:205–16; doi:10.1016/j.tim.2016.10.001

113. Freeman ZN, Dorus S, Waterfield NR. The KdpD/KdpE Two-Component System: Integrating K+ Homeostasis and Virulence. PLoS Pathog. 2013;9:e1003201-; doi:10.1371/journal.ppat.1003201

114. Meadows JA, Wargo MJ. Carnitine in bacterial physiology and metabolism. Microbiology (N Y). 2015;161:1161–74; doi:10.1099/mic.0.000080

115. Kolata P, Efremov RG. Structure of *Escherichia coli* respiratory complex I reconstituted into lipid nanodiscs reveals an uncoupled conformation. Swartz KJ, Carter AP, Kumar V, Letts JA, editors. Elife. 2021;10:e68710; doi:10.7554/eLife.68710

116. Korea C-G, Badouraly R, Prevost M-C, Ghigo J-M, Beloin C. *Escherichia coli* K-12 possesses multiple cryptic but functional chaperone–usher fimbriae with distinct surface specificities. Environ Microbiol. 2010;12:1957–77; doi:10.1111/j.1462-2920.2010.02202.x

117. Fu H, Lee J, Wang T. Heat-shock increases rpoD-dependent β-galactosidase activity in the *Escherichia coli* strains BD792 and B23. 2015;19

118. Gagarinova A, Hosseinnia A, Rahmatbakhsh M, Istace Z, Phanse S, Moutaoufik MT, et al. Auxotrophic and prototrophic conditional genetic networks reveal the rewiring of transcription factors in *Escherichia coli*. Nat Commun. 2022;13:4085; doi:10.1038/s41467-022-31819-x

119. Prusiner S, Davis JN, Stadtman ER. Regulation of glutaminase B in *Escherichia coli*. I. Purification, properties, and cold lability. Journal of Biological Chemistry. 1976;251:3447–56; doi: 10;251(11):3447-56

120. Iacometti C, Marx K, Hönick M, Biletskaia V, Schulz-Mirbach H, Satanowski A, et al. Activating silent glycolysis bypasses *Escherichia coli*; bioRxiv. 2021;2021.11.18.468982; doi: 10.34133/2022/9859643

121. Njenga R, Boele J, Öztürk Y, Koch H-G. Coping with stress: How bacteria fine-tune protein synthesis and protein transport. Journal of Biological Chemistry. 2023;299:105163; doi:10.1016/j.jbc.2023.105163

122. Hajra D, Nair AV, Chakravortty D. An elegant nano-injection machinery for sabotaging the host: Role of Type III secretion system in virulence of different human and animal pathogenic bacteria. Phys Life Rev. 2021;38:25–54; doi:10.1016/j.plrev.2021.05.007

123. Peek J, Christendat D. The shikimate dehydrogenase family: Functional diversity within a conserved structural and mechanistic framework. Arch Biochem Biophys. 2015;566:85–99; doi:10.1016/j.abb.2014.12.006

124. Miwa T, Taguchi H. *Escherichia coli* small heat shock protein IbpA plays a role in regulating the heat shock response by controlling the translation of σ32. Proceedings of the National Academy of Sciences. 2023;120:e2304841120; doi:10.1073/pnas.2304841120

125. Brocklehurst KR, Hobman JL, Lawley B, Blank L, Marshall SJ, Brown NL, et al. ZntR is a Zn(II)-responsive MerR-like transcriptional regulator of zntA in *Escherichia coli*. Mol Microbiol. 1999;31:893–902; doi: 10.1046/j.1365-2958.1999.01229.x

126. T CP. A Galvanizing Story—Protein Stability and Zinc Homeostasis. J Bacteriol. 2007;189:2953–4; doi:10.1128/jb.00173-07

127. Iannuzzi C, Adrover M, Puglisi R, Yan R, Temussi PA, Pastore A. The role of zinc in the stability of the marginally stable IscU scaffold protein. Protein Sci. 2014;23(9):1208-1219. doi:10.1002/pro.2501.

128. Joly N, Engl C, Jovanovic G, Huvet M, Toni T, Sheng X, et al. Managing membrane stress: the phage shock protein (Psp) response, from molecular mechanisms to physiology. FEMS Microbiol Rev. 2010;34:797–827; 10.1111/j.1574-6976.2010.00240.x

129. LeRoux M, Culviner PH, Liu YJ, Littlehale ML, Laub MT. Stress Can Induce Transcription of Toxin-Antitoxin Systems without Activating Toxin. Mol Cell. 2020;79:280-292.e8; doi:10.1016/j.molcel.2020.05.028

130. Flores-Kim J, Darwin AJ. The Phage Shock Protein Response. Annu Rev Microbiol. 2016;70:83–101; doi:10.1146/annurev-micro-102215-095359

131. Walsh P, Bursać D, Law YC, Cyr D, Lithgow T. The J-protein family: modulating protein assembly, disassembly and translocation. EMBO Rep. 2004;5:567-571–571; doi:10.1038/sj.embor.7400172

132. Park DM, Akhtar MdS, Ansari AZ, Landick R, Kiley PJ. The Bacterial Response Regulator ArcA Uses a Diverse Binding Site Architecture to Regulate Carbon Oxidation Globally. PLoS Genet. 2013;9:e1003839; doi:10.1371/journal.pgen.1003839

133. Anand A, Patel A, Chen K, Olson CA, Phaneuf P V, Lamoureux C, et al. Laboratory evolution of synthetic electron transport system variants reveals a larger metabolic respiratory system and its plasticity. Nat Commun. 2022;13:3682; 10.1038/s41467-022-30877-5

134. Henry C, Loiseau L, Vergnes A, Vertommen D, Mérida-Floriano A, Chitteni-Pattu S, et al. Redox controls RecA protein activity via reversible oxidation of its methionine residues. Kowalczykowski SC, Storz G, editors. Elife. 2021;10:e63747; doi:10.7554/eLife.63747

135. Flint DH, Emptage MH, Guest JR. Fumarase A from *Escherichia coli:* purification and characterization as an iron-sulfur cluster containing enzyme. Biochemistry   . 1992;31:10331–7; doi:10.1021/bi00157a022

136. Park S-J, Tseng C-P, Gunsalus RP. Regulation of succinate dehydrogenase (sdhCDAB) operon expression in *Escherichia coli* in response to carbon supply and anaerobiosis: role of ArcA and Fnr. Mol Microbiol. 1995;15:473–82; doi:10.1111/j.1365-2958.1995.tb02261.x

137. Roth M, Jaquet V, Lemeille S, Bonetti E-J, Cambet Y, Francois P, et al. Transcriptomic Analysis of *E. coli* after Exposure to a Sublethal Concentration of Hydrogen Peroxide Revealed a Coordinated Up-Regulation of the Cysteine Biosynthesis Pathway. Antioxidants. 2022;11; doi:10.3390/antiox11040655

138. Friedrich T, Dekovic DK, Burschel S. Assembly of the *Escherichia coli* NADH:ubiquinone oxidoreductase (respiratory complex I). Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2016;1857:214–23; doi:10.1016/j.bbabio.2015.12.004

139. Prasad I, Young B, Schaefler S. Genetic Determination of the Constitutive Biosynthesis of Phospho-β-Glucosidase A in *Escherichia coli* K-12. J Bacteriol. 1973;114:909–15. 10.1128/jb.114.3.909-915.1973

140. Kuznetsova E, Proudfoot M, Gonzalez CF, Brown G, Omelchenko M V, Borozan I, et al. Genome-wide Analysis of Substrate Specificities of the *Escherichia coli* Haloacid Dehalogenase-like Phosphatase Family*. Journal of Biological Chemistry. 2006;281:36149–61; doi:10.1074/jbc.m605449200

141. Sobota JM, Imlay JA. Iron enzyme ribulose-5-phosphate 3-epimerase in *Escherichia coli* is rapidly damaged by hydrogen peroxide but can be protected by manganese. Proceedings of the National Academy of Sciences. 2011;108:5402–7; doi:10.1073/pnas.1100410108

142. Chuanboon K, Na Nakorn P, Pannengpetch S, Laengsri V, Nuchnoi P, Isarankura-Na-Ayudhya C, et al. Proteomics and bioinformatics analysis reveal potential roles of cadmium-binding proteins in cadmium tolerance and accumulation of *Enterobacter cloacae*; PeerJ. 2019;7:e6904; https://pubmed.ncbi.nlm.nih.gov/31534833/

143. Imlay JA, Linn S. Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. J Bacteriol. 1986;166:519–27; doi:10.1128/jb.166.2.519-527.1986

144. McCarthy DJ, Smyth GK. Testing significance relative to a fold-change threshold is a TREAT. Bioinformatics. 2009;25:765–71; doi:10.1093/bioinformatics/btp053

145. Umbarger HE, Brown B. Threonine deamination in *Escherichia coli*. II. Evidence for two
L-threonine deaminases. J Bacteriol. 1957;73(1):105–12; doi:10.1128/jb.73.1.105-112.1957.

146. Julien M, Elisa G, Ronald VE, Enrique TMO, Ladan K, Laleh K, et al. Gene Erosion Can
Lead to Gain-of-Function Alleles That Contribute to Bacterial Fitness. mBio.
2021;12:10.1128/mbio.01129-21; doi:10.1128/mbio.01129-21

147. Greenfield D, McEvoy AL, Shroff H, Crooks GE, Wingreen NS, Betzig E, et al. Self-
Organization of the *Escherichia coli* Chemotaxis Network Imaged with Super-Resolution Light
Microscopy. PLoS Biol. 2009;7:e1000137; doi:10.1371/journal.pbio.1000137

148. Guinote IB, Moreira RN, Barahona S, Freire P, Vicente M, Arraiano CM. Breaking through
the stress barrier: the role of BolA in Gram-negative survival. World J Microbiol Biotechnol.
2014;30:2559–66; 10.1007/s11274-014-1702-4

149. Hemm MR, Paul BJ, Miranda-Ríos J, Zhang A, Soltanzad N, Storz G. Small stress
response proteins in *Escherichia coli*: proteins missed by classical proteomic studies. J
Bacteriol. 2010;192(1):46–58; doi:10.1128/JB.00872-09.

150. J RB, H DR, E PR, L SH, I BP, Aindrila M, et al. Functional Genomic Study of Exogenous
n-Butanol Stress in *Escherichia coli*. Appl Environ Microbiol. 2010;76:1935–45;
doi:10.1128/AEM.02323-09

151. Chen B, Chu T-W, Chiu K, Hong M-C, Wu T-M, Ma J-W, et al. Transcriptomic analysis
elucidates the molecular processes associated with hydrogen peroxide-induced diapause
termination in Artemia-encysted embryos. PLoS One. 2021;16:e0247160;
doi:10.1371/journal.pone.0247160

152. Allison H, Jacquelyn S, Alexandra K, Kathleen C, Matthew Y, Dennis W, et al. The
Bacterial iprA Gene Is Conserved across Enterobacteriaceae, Is Involved in Oxidative Stress
Resistance, and Influences Gene Expression in *Salmonella enterica* Serovar Typhimurium. J
Bacteriol. 2016;198:2166–79; doi:10.1128/jb.00144-16

153. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on metabolism,
oxidative stress, and diseases. Amino Acids. 2017;49:2091–8; doi:10.1007/s00726-017-2494-
2

154. Clausznitzer D, Oleksiuk O, Løvdok L, Sourjik V, Endres RG. Chemotactic Response
and Adaptation Dynamics in *Escherichia coli*. PLoS Comput Biol. 2010;6:e1000784;
doi:10.1371/journal.pcbi.1000784

155. Bouillet S, Hamdallah I, Majdalani N, Tripathi A, Gottesman S. A negative feedback loop
is critical for recovery of RpoS after stress in Escherichia coli. bioRxiv.
2023;2023.11.09.566509; doi:10.1101/2023.11.09.566509.

156. Song S, Wood TK. A primary physiological role of toxin/antitoxin systems is phage
inhibition. Front Microbiol. 2020;11:1895; doi:10.3389/fmicb.2020.01895.

157. chmidt M, Proctor T, Diao R, Freddolino PL. *Escherichia coli* YigI is a conserved
Gammaproteobacterial acyl-CoA thioesterase permitting metabolism of unusual fatty acid
substrates. J Bacteriol. 2022;204:e00014-22; doi:10.1128/JB.00014-22.

158. Jack RL, Sargent F, Berks BC, Sawers G, Palmer T. Constitutive expression of
Escherichia coli tat genes indicates an important role for the twin-arginine translocase during

aerobic and anaerobic growth. J Bacteriol. 2001;183(5):1801–4; doi:10.1128/jb.183.5.1801-1804.2001.

159. Zhao R, Li Z, Sun Y, Ge W, Wang M, Liu H, et al. Engineered *Escherichia coli* Nissle 1917 with urate oxidase and an oxygen-recycling system for hyperuricemia treatment. Gut Microbes. 2022;14:2070391; doi:10.1080/19490976.2022.2070391

160. Hryniewicz M, Sirko A, Pałucha A, Böck A, Hulanicka D. Sulfate and thiosulfate transport in *Escherichia coli* K-12: identification of a gene encoding a novel protein involved in thiosulfate binding. J Bacteriol. 1990;172(6):3358–66; doi:10.1128/jb.172.6.3358-3366.1990.

161. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database resources of the National Center for Biotechnology. 2022; doi:10.1093/nar/gkab1112

162. Krueger F, James F, Ewels P, Afyounian E, Weinstein M, Schuster-Boeckler B, et al. FelixKrueger/TrimGalore: v0.6.10 - add default decompression path. Zenodo; 2023; doi:10.5281/zenodo.7598955

163. FastQC. 2015. Available from: https://qubeshub.org/resources/fastqc

164. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9; doi:10.1038/nmeth.1923

165. Wang L, Wang S, Li W. RSeQC: quality control of RNA-seq experiments. Bioinformatics. 2012;28:2184–5; doi:10.1093/bioinformatics/bts356

166. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30; doi:10.1093/bioinformatics/btt656

167. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics. 2016;32:3047–8. Available from: https://doi.org/10.1093/bioinformatics/btw354

168. Lamoureux CR, Decker KT, Sastry A V, Rychel K, Gao Y, McConn JL, et al. A multi-scale expression and regulation knowledge base for *Escherichia coli*. Nucleic Acids Res. 2023;51:10176–93; doi:10.1093/nar/gkad750

169. Consortium TGO, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al. The Gene Ontology knowledgebase in 2023. Genetics. 2023;224:iyad031; doi:10.1093/genetics/iyad031

170. Gumienny R. GEOparse: Python Library to Access Gene ExpressionOmnibus Database(GEO). Available from: https://pypi.org/project/GEOparse/

171. Kim D, Seo SW, Gao Y, Nam H, Guzman GI, Cho B-K, et al. Systems assessment of transcriptional regulation on central carbon metabolism by Cra and CRP. Nucleic Acids Res . 2018;46:2901–17; doi:10.1093/nar/gky069

172. Spearman Rank Correlation Coefficient. The Concise Encyclopedia of Statistics. New York, NY: Springer New York; 2008. p. 502–5. Available from: https://doi.org/10.1007/978-0-387-32833-1_379

1533 173. Martelli PL, Fariselli P, Casadio R. An ENSEMBLE machine learning approach for the
1534 prediction of all-alpha membrane proteins. Bioinformatics. 2003;19:i205–11. ;
1535 doi:10.1093/bioinformatics/btg1027

1536
1537 174. Pierleoni A, Martelli PL, Fariselli P, Casadio R. BaCelLo: a balanced subcellular
1538 localization predictor. Bioinformatics. 2006;22:e408–16; doi:10.1093/bioinformatics/btl222

1539
1540 175. Pierleoni A, Martelli PL, Casadio R. PredGPI: a GPI-anchor predictor. BMC
1541 Bioinformatics. 2008;9:392; doi:10.1186/1471-2105-9-392

1542
1543 176. Pierleoni A, Martelli PL, Casadio R. MemLoci: predicting subcellular localization of
1544 membrane proteins in eukaryotes. Bioinformatics. 2011;27:1224–30;
1545 doi:10.1093/bioinformatics/btr108

1546
1547 177. Savojardo C, Fariselli P, Casadio R. BETAWARE: a machine-learning tool to detect and
1548 predict transmembrane beta-barrel proteins in prokaryotes. Bioinformatics .
1549 2013;29:504–5; doi:10.1093/bioinformatics/bts728

1550
1551 178. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING
1552 database in 2021: customizable protein–protein networks, and functional characterization of
1553 user-uploaded gene/measurement sets. Nucleic Acids Res. 2021;49:D605–12;
1554 10.1093/nar/gkaa1074

1555
1556 179. Fernando PC, Mabee PM, Zeng E. Integration of anatomy ontology data with protein–
1557 protein interaction networks improves the candidate gene prediction accuracy for anatomical
1558 entities. BMC Bioinformatics. 2020;21:442; doi:10.1186/s12859-020-03773-2

1559
1560 180. Remmert M, Biegert A, Hauser A, Söding J. HHblits: lightning-fast iterative protein
1561 sequence searching by HMM-HMM alignment. Nat Methods. 2012;9:173–5;
1562 doi:10.1038/nmeth.1818

1563
1564 181. Bagheri H, Dyer R, Severin A, Rajan H. Comprehensive Analysis of Non Redundant
1565 Protein Database. 2020; doi:10.21203/rs.3.rs-54568/v1

1566
1567 182. Andreeva A, Kulesha E, Gough J, Murzin AG. The SCOP database in 2020: expanded
1568 classification of representative family and superfamily domains of known protein structures.
1569 Nucleic Acids Res. 2020;48:D376–82. Available from:10.1093/nar/gkz1064

1570
1571 183. Barrio-Hernandez I, Yeo J, Jänes J, Mirdita M, Gilchrist CLM, Wein T, et al. Clustering
1572 predicted structures at the scale of the known protein universe. Nature. 2023;622:637–45;
1573 doi:10.1038/s41586-023-06510-w

1574
1575 184. Tuttle AR, Trahan ND, Son MS. Growth and Maintenance of *Escherichia coli* Laboratory
1576 Strains. Curr Protoc. 2021;1:e20; doi:10.1002/cpz1.20

1577 185. Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, et al. The RIN:
1578 an RNA integrity number for assigning integrity values to RNA measurements. BMC Mol Biol.
1579 2006;7:3; doi:10.1186/1471-2199-7-3

1580
1581 186. Lataretu M, Hölzer M. RNAflow: An Effective and Simple RNA-Seq Differential Gene
1582 Expression Pipeline Using Nextflow. Genes (Basel). 2020;11:1487;
1583 doi:10.3390/genes11121487

1584
1585 187. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
1586 Bioinformatics. 2018;34:i884–90; doi:10.1093/bioinformatics/bty560

1587

1588   188. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
1589   Bioinformatics. 2018;34:i884–90; doi:10.1093/bioinformatics/bty560

1590
1591   189. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
1592   requirements. Nat Methods. 2015;12:357–60; doi:10.1038/nmeth.3317

1593
1594   190. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
1595   Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.;
1596   doi:10.1093/bioinformatics/btp352

1597
1598   191. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
1599   assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30;
1600   doi:10.1093/bioinformatics/btt656

1601
1602   192. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
1603   RNA-seq data with DESeq2. Genome Biol. 2014;15:550; doi:10.1186/s13059-014-0550-8

1604
1605   193. Duda JC, Drenda C, Kästel H, Rahnenführer J, Kappenberg F. Benefit of using interaction
1606   effects for the analysis of high-dimensional time-response or dose-response data for two-
1607   group comparisons. Sci Rep. 2023;13:20804; doi:10.1038/s41598-023-47057-0

1608
1609   194. Liou G-G, Chao Kaberdina A, Wang W-S, Kaberdin VR, Lin-Chao S. Combined
1610   Transcriptomic and Proteomic Profiling of *E. coli* under Microaerobic versus Aerobic
1611   Conditions: The Multifaceted Roles of Noncoding Small RNAs and Oxygen-Dependent
1612   Sensing in Global Gene Expression Control. Int J Mol Sci. 2022;23; doi:10.3390/ijms23052570