**Title**

Deciphering the Proteome of *Escherichia coli* K-12: Integrating Transcriptomics and Machine

Learning to Annotate Hypothetical Proteins

**Author information**

Sagarika Chakraborty[1], Zachary Ardern[1,2], Habibu Aliyu[1] and Anne-Kristin Kaster[1,3]*

[1] Institute for Biological Interfaces 5 (IBG-5), Biotechnology and Microbial Genetics, Karlsruhe

Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-

Leopoldshafen, Germany

[2] Wellcome Trust Sanger Institute, Hinxton, Saffron Walden CB10 1RQ, United Kingdom

[3] Institute for Applied Biosciences (IAB), Karlsruhe Institute of Technology (KIT), Kaiserstraße

12, 76131 Karlsruhe, Germany

* To whom correspondence should be addressed: Email: kaster@kit.edu

**Graphical Abstract**

**Abstract**

Omics technologies have led to the discovery of a vast number of proteins that are expressed but have no functional annotation - so called hypothetical proteins (HPs). Even in the best studied model organism *Escherichia coli* K-12, 2% of the proteome remains uncharacterized, with no sequence homologies available in public databases. This knowledge gap becomes even worse when looking at microbial dark matter. However, knowing the functions of proteins is crucial for elucidating cellular and metabolic processes and harnessing biotechnological potentials. In this study, we employed machine learning (ML) to decipher the transcriptional regulatory network of *E. coli* K-12, deep learning for the prediction of structural homologs as well as other ML and bioinformatic tools, with the goal to assign functions to uncharacterized HPs. We further provide a proof-of-concept for experimental validation of function for three HP-encoding genes (*yhdN*, *yeaC* and *ydgH*) hinted from the *in silico* methods, by analyzing growth patterns of *E. coli* K-12 deletion mutants compared to the wild type, as well as their transcriptional responses to specific conditions. This study demonstrates that the use of Big Omics Data in combination with Artificial Intelligence and experimental controls is a powerful approach to illuminate functional dark matter.

**Keywords**: Functional Dark Matter, Big Omics Data, Artificial Intelligence, Independent Component Analysis (ICA), Functional Annotation of Proteins

## Introduction

Due to the advents in Next Generation Sequencing technologies, the volume of omics data has massively increased over the past years. As of today, gold-standard knowledgebases like the National Center for Biotechnology Information (NCBI)'s Reference Sequences Database (RefSeq) and the Universal Protein Resource Database (UniProt) [1] provide freely accessible sequences and information for over 163 million unique prokaryotic proteins **(Figure 1)**. This rapid accumulation of sequencing data has long surpassed the rate of possible experimental characterization *in vivo* and *in vitro*. Hence, researchers often rely on functional analysis of proteins from model organisms. These findings are then extrapolated *in silico* to closely related sequence homologues. Consequently, the definition of cut-off values plays a crucial role in protein annotation. To manage this challenge, automated annotation pipelines, such as InterPro have become a valuable tool [2]. Here, analysis of protein sequences is provided by classifying them into families and integrating predictive models from multiple protein databases such as e.g. Pfam [3], ProSite [4], SMART [5] and/or CDD [6] to infer functions for experimentally uncharacterized proteins. However, only 83% of all UniProt sequences have so far been functionally annotated by InterPro [7] The remaining 17% of proteins are designated with terms such as 'putative uncharacterized', 'uncharacterized', 'unknown protein families (UPFs)' or proteins bearing 'domains of unknown functions' (DUFs). These proteins are summarized by the term hypothetical proteins (HPs), since they are expressed but could so far not be functionally characterized by classical *in vivo*, *in vitro*, and/or *in silico* methods (8,9). In addition, there is also no information on these proteins available in widely-used knowledge-databases such as BioCyc [10], RegulonDB [11], and EggNOG [12] **(Figure 1).**

While there have been notable advancements in methodologies for protein function prediction in the recent past [13–16] several challenges persist [17,18]: These include different functionalities across homologous proteins, proteins performing different functions in distinct cellular locations, proteins with multiple three-dimensional structures, the lack of conserved proteins across different species to identify shared functional relationships and a heavy

reliance on pre-existing data for inferential annotations [19]. Even in the best studied model

organisms *Escherichia coli* K-12, 31.8% of the genome comprises of protein encoding genes

that lack experimental validation, commonly referred to as 'Putative HPs' [8], and 2.1% that

even lack sequence homologies (this study) (**Figure 1**). The problem of annotating HPs

becomes even larger when looking at microbial dark matter(9, 20).In uncultivated

microorganisms, the proportions of genes with unknown functions can comprise up to 60% in

bacterial and 80% in archaeal genomes [21]. However, elucidating the function of these

proteins is essential for understanding cellular processes, metabolism and evolution, as well

as for harnessing their biotechnological potential [22].

Transcriptional profiling has helped to gain insights on protein functions by monitoring gene

expressions and clustering genes based on their responses to varying environmental

conditions and stimuli [23]. Here, the entire set of RNA transcripts produced by a genome

under specific conditions provides a dynamic representation of the organisms' operational

state. However, standard bioinformatics-based methods for transcriptomic data analysis are

low-throughput and require quite extensive computing power and time due to the high degrees

of complexity and heterogeneity of the data [24]. Today, artificial intelligence (AI)-based

methodologies have made significant advancements over traditional bioinformatics

approaches, offering the capacity to process and analyze data at a scale and speed that were

previously unattainable (24, 25). They are particularly suited for deciphering complex

interactions among genes and transcription factors or proteins that regulate them, as well as

revealing the regulation of gene expression in response to diverse environmental conditions

[27]. Among various techniques for analysis, module-detection methods have proven to

efficiently predict the functions of co-regulated groups of genes from large gene expression

datasets [28]. Independent Component Analysis (ICA), introduced by Comon in 1994 [29],

employs an unsupervised machine learning (ML) approach, categorizing the input data without

any prior information. This technique is especially relevant in the context of transcriptomics,

where the goal is to understand the complex interplay of gene expression signals and to

decipher co-regulated gene sets. ICA therefore allows to reveal the regulatory patterns and

activity levels of genes governed by specific regulators (i.e. transcription factors controlling the

expression of genes) across diverse experimental conditions, so called transcriptional

regulatory networks (TRNs), thereby disclosing the organizational and functional architecture

of genes. Expanding upon this, McConn et al. developed a variant of ICA, termed OptICA,

which can be applied to large TRNs, analyzing interactions between transcription factors and

their target genes. Here, discrete groups of genes are clustered into robust independent

components by avoiding over-clustering of datasets (leading to loss of biological information)

or under-clustering (leading to too few clusters and a loss of output resolution), and therefore

providing an optimal representation of the organism's underlying TRN [27]. This results in

revealing independently regulated gene groups, so called iModulon from transcriptomics

datasets (29, 30). iModulons are therefore the data-driven analogs of so-called regulons - sets

of genes governed by regulators [30]. The OptICA algorithm along with high-quality RNA-seq

data has recently been used to decipher the TRN of *Bacillus subtilis* [32] and *E. coli* [27] to

uncover the detailed responses to environmental conditions and genetic perturbations, e.g.

deletion mutants. However, these studies focused on the identification of regulons that were

not experimentally described before rather than the functional characterization of HPs.

In order to decipher the function of HP-encoding genes in the *E. coli* K-12 proteome, we

adapted the workflow originally designed for *Bacillus subtilis* [32] and analyzed the gene

expression data of the *E. coli* K-12 sub-strains MG1655 and BW25113. Functional

characterization of HP-encoding genes was inferred by identifying co-regulated genes under

the influence of a common regulator from publicly available transcriptomic datasets. This

characterization was then further validated by annotating these genes using specific Gene

Ontology (GO) categories through the PANTHER tool (Protein ANalysis THrough Evolutionary

Relationships) [33]. Additionally, metadata conditions for the transcriptomic datasets were

extracted. Furthermore, other *in silico* methods were employed to determine the sub-cellular

localization [34], protein-protein interactions (PPIs) [35], genomic context [36] and sequence-

based remote homologies [37]. Structural homology information was obtained by using the

AlphaFold Protein Structure Database (AFDB) clusters webserver, which is based on a deep

learning (DL) method [38].

Since AI-predicted functions are usually not experimentally verified, raising the possibility of

untrue annotations [9], we also provide an experimental validation of three candidate HP-

encoding genes (*yhdN*, *yeaC* and *ydgH*). The growth curves and transcriptional responses of

*E. coli* K-12 deletion mutants in comparison to the wild-type strain under specific conditions

were analyzed, showing that the *in silico* predictions were indeed true. Hence, the here

presented pipeline not only facilitates the discovery of previously unidentified regulators, but

also sheds light on our understanding of the function of HPs **(Figure 1).**

**Figure 1. Total number of sequences for all unique prokaryotic and *Escherichia coli* proteins deposited in the National Center for Biotechnology Information (NCBI) as of April 2024 and methodological set-up of this study.** Of the 4403 *E. coli* K-12 protein encoding genes, 1403 (31.8%) encode for unique proteins with functions predicted only *in silico* based on homologous sequences but lacking *in vivo* or *in vitro* experimental evidence (termed "putative hypothetical proteins"). 95 protein encoding genes (2.1%) of *E. coli* K-12 are completely uncharacterized with no sequence homologues according to the four knowledge databases - EcoCyc [39], RegulonDB [11], EggNOG [12] and UniProt (1) (termed "hypothetical proteins"). Transcriptomic datasets from NCBI were filtered and processed using the OptICA approach to generate iModulons [32]. Metadata information was curated in parallel using manual or semi-automated approaches(39–41).Bioinformatics, machine learning and deep learning tools along with the presence of relevant metadata then resulted in potential functions for HP candidates for *in vitro* testing. HPs, hypothetical proteins; ICA, independent component analysis; ML, machine learning.

**Results and Discussion**

**Identification of uncharacterized HPs in *E. coli***

In 2019, Ghatak et al. had identified HP-encoding genes in *E. coli* K-12, however only by employing keywords such as 'possibly', 'predicted' or 'hypothetical' from the EcoCyc database, or annotation scores of two or below in UniProt [8]. This approach did, however, not establish a stringent threshold for accurate identification of completely uncharacterized HP-encoding genes. Hence, a more rigorous approach was used in this study, which included additional keywords for HPs (such as 'Uncharacterized', 'Putative uncharacterized', 'DUF' etc.) and information on the absence of any functional characterization using multiple gold-standard databases (EcoCyc, UniProt , RegulonDB and EggNOG). 158 of the 1600 HP-encoding genes previously listed by Ghatak et al. have now been functionally characterized with experimental evidence in the EcoCyc database. Notably, the HP-encoding gene *ydfX*, curated as one of the HPs in this study, was missing from Ghatak et al.'s list since it was previously categorized as a pseudogene/phantom gene. Its exclusion, however, could not be verified in the current EcoCyc database. We identified 1403 of 4403 genes (31.8%) of the *E. coli* K-12 genome as 'putative' HPs, implying that their functions were inferred based on sequence homologies in the gold-standard databases, but lacking an experimental evidence for function.  A recent study using the EggNOG tool also identified about 30% in a different *E.* strain (O157:H7 strain Sakai) as putative HPs [18].

95 HP-encoding genes (2.1%) could be identified with no sequence homologies in the *E. coli* K-12 genome. These could be further sub-categorized in 'uncharacterized proteins' (53), 'putative uncharacterized' (5), '(DUF)' domain containing proteins (21) and 'UPF' (HPs with unknown protein families, 3). 13 HPs had the characteristic 'Protein Y…' naming scheme. The distribution of those 95 HP-encoding genes within the genome can be found in the Supplementary (*Supplementary Figure 1)*.

**iModulon generation and analysis**

In order to investigate the functions for the 95 uncharacterized HP-encoding genes, publicly

available RNAseq datasets from *E. coli* K-12 substrain MG1655 were downloaded from NCBI

and quality filtered. 779 high-quality datasets from the MG1655 substrain were then used as

input for the OptICA algorithm [27]. For the substrain BW25113, a separate OptICA analysis

was not conducted due to the very small size of the dataset. The iModulons were identified by

utilizing the PyModulon package, specifically employing the *Inferring iModulon Activities*

function [32] for the BW25133 substrain. In total, 131 iModulons were obtained for the entire

*E. coli* K-12 transcriptome. Three single gene iModulons were discarded from the analyses,

since they were considered a result of artificial knockouts or overexpression of single genes

in the dataset. Therefore, a total of 128 iModulons were selected for further analysis.

Functional information on the HP-encoding genes was obtained by mapping the iModulons to

existing experimentally derived regulons (groups of genes regulated by the same transcription

factor), using data from the EcoCyc and RegulonDB databases. Additionally, Gene Ontology

(GO)-based functional annotations for member genes of an iModulon (co-regulated gene sets)

were identified using the PANTHER tool (Protein ANalysis THrough Evolutionary

Relationships) [33]. A total of 51 HP-encoding genes were clustered into iModulons using

OptICA. Of these, 46 genes were assigned putative functions based on information from

regulator proteins and/or Gene Ontology (GO) categories. (**Figure 2**). Two genes, *yfeS* and

*yffL* had information on their regulators (RpoE and NarL, respectively) but no assigned GO

categories, while five genes (*ybaA, yfaP, yjgZ, ymgI and ymgJ*) (white) could not be associated

with a regulator or a GO category. 44 of the HP-encoding genes could not be clustered into

distinct iModulons (grey) (**Figure. 2**).

**Figure 2. Regulator and functional classification for 95 HP-encoding genes in *Escherichia coli K-12*.** 44 HP-encoding genes could not be clustered by the OptICA method [27] (grey), since they did not co-regulate with any other genes. 22 genes were characterized based on information on their regulators as obtained from RegulonDB and/or EcoCyc, as well as by Gene Ontology (GO) categories based on the co-regulating genes using the PANTHER tool (Protein ANalysis THrough Evolutionary Relationships) (orange) [33]. 29 genes co-regulated with other genes, but no information on their regulators could be obtained (blue). 24 out of these 29 genes could be assigned putative functions based on GO annotations of co-regulating genes derived from the PANTHER tool. The striped regions denote genes where the regulator-associated function from EcoCyc/RegulonDB did not match a GO-derived functional category obtained from PANTHER. Highlighted in bold are the three HP-encoding genes which were selected for *in vitro* testing.

In addition to the 95 HP-encoding genes, all other *E. coli K-12* genes were also analysed using this method. 2630 of the 2905 (91%) genes which have experimental evidence of function could also be characterized into iModulons. 1697 (64%) co-regulated with other genes, i.e. they were part of characterized iModulons with information on their regulator on RegulonDB and/or EcoCyc, while 933 of the genes were part of uncharacterized iModulons. For genes with known regulators, only 121 (5%) had either a non-matching functional category when compared with the PANTHER categories or a non-significant (*p*>0.05) functional category, and 95% had a matching PANTHER GO category. For genes which were part of uncharacterized iModulons, with no regulator information, 546 (58%) could be functionally classified using PANTHER. 1061 of the 1403 putative HP-encoding genes (75%), i.e. genes with functional characterisation based on sequence homology information, could be characterized into iModulons. 342 genes (25%) did not co-regulate with other genes. For genes that had co-regulation, 539 (51%) had information on their regulator on RegulonDB

and/or EcoCyc. The other 522 (49%) were part of uncharacterized iModulons. For genes with

known regulators, only 20 genes (4%) had either a non-matching function when compared

with the PANTHER categories or a non-significant ($p > 0.05$) functional category in PANTHER.

For genes with no information on their regulator, 366 (67.9%), could however be characterized

by PANTHER. Hence, about 95% of the iModulon functional categories matched the

PANTHER GO categories, demonstrating the robustness of the method. Moreover, the

inclusion of experimentally characterized genes within uncharacterized iModulons shows

potential for the discovery of new regulons or co-regulated genes.

**Functional analysis of HP-encoding genes using *in silico* tools**

All 95 HP-encoding genes were analysed based on multiple criteria including the generated

iModulons from the OptICA algorithm, functional information based on the regulators of the

co-regulated gene sets obtained *via* RegulonDB/EcoCyc and their GO categories from

PANTHER. Information on metadata conditions pertaining to the highest level of gene

expression relative to the reference gene *frr* (encoding the ribosome recycling factor) [42] and

annotation of the most significant co-expressed gene were also extracted. Additionally, the

sub-cellular localization, protein-protein interactions, genomic context, remote sequence

homology as well as structural homology aided in forming hypothesises for assigning functions

to the respective HPs. 33 of the protein encoding genes (highlighted in green) could be

assigned functions with a high confidence. 28 of the genes were assigned functions based on

lower confidence, since only information from a maximum of two sources could be well-

correlated (highlighted in yellow). A clear inference could not be made for 34 of the genes

(highlighted in red). Information retrieved for each gene and functional inference (if applicable)

is detailed in **Table1**. A pictorial overview of Table 1 based on the functional annotations for

the 95 HPs can be found in *Supplementary Figure 2*.

**Table1. A comprehensive functional interference of HPs based on different *in silico* tools.** Annotations are based on regulator information from RegulonDB/EcoCyc and/or GO category based on co-regulated genes obtained from PANTHER) [33](as shown in Fig. 2). The highest gene expression condition, obtained from the Tjaden dataset [43], with metadata conditions categorized by the type of medium, oxygen availability (normal $O_2$ conditions unless stated otherwise), temperature etc. in comparison to the reference gene 'frr' are described. Functional annotations of co-expressed genes were retrieved from EcoCyc [36]. 'SCL' or sub-cellular localization inferred from the BUSCA tool [34] are abbreviated as either 'NUC' for nuclear 'CYT' for cytoplasmic, 'ECS' for extra cellular space or 'PM' for plasma membrane. Functional information from protein-protein interaction (PPI) analyses was obtained from STRINGdb [35]. Annotations for neighbouring upstream (◄——) and downstream (——►) genes in context of the HP-encoding gene were retrieved from the EcoCyc database. Additionally, remote homology was assessed using the Uniref30 database using the Hhblits tool [37], and structural homology was inferred from the AlphaFold Protein Structure Database [38]. A functional inference can be made for genes supported by multiple lines of evidence when similar functional information was provided by at least two sources. Proteins with well correlated information from at least three sources are highlighted in green (with higher confidence) and proteins with information that can be correlated using at least two sources are highlighted in yellow (with lower confidence). Protein encoding genes highlighted in grey were selected for *in vitro* testing. An unclear inference suggests that, based on the different knowledge bases, the extracted information did not provide enough confidence or could not be well correlated to assign a function (red).

| Gene | Function based on regulator and/or co-regulated genes | Condition with highest gene expression (growth medium) | Annotation of most significant co-expressed gene | SCL | PPI: function of partner protein | Annotation of neighbouring genes | | | Remote homology | Structural homology | Overall functional inference |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1. yabP | Anaerobic respiration | M9, 37°C; 100 µg/ml Ampicillin (inhibits cell wall synthesis) [44] | Growth related protein (yabQ) | CYT | No interaction partner | djlA ◄—— Co-chaperone protein | **yabP** ——► | yabQ Growth related protein | Putative Undecaprenyl pyrophosphate phosphatase (cell wall biosynthesis) [45] | Type III secretion system (bacterial pathogenecity) [46] | Cell wall biosynthesis |
| 2. yagM | Cell adhesion | LB, 37°C; 20 µg/ml Bicyclomycin (produces ROS, blocks cell division) [47] | Transcriptional regulator (fimZ) (involved in fimbriae production) [48] | CYT | DNA-binding protein (YagL) | yagL ◄—— resolvase like prophage | **yagM** ◄—— | ykgV ◄—— Growth related protein | Unknown function | Unknown function | Cell adhesion under antibiotic stress |
| 3. yagK | Glycine betaine bio-synthesis from choline (osmoprotectant) [49] | Na-Phosphate buffer, 37°C; Δthil (thil is responsive to UV) [50] | Putative fimbrial usher protein (yhcD) | NUC | Putative ligase (GtrS) | yagJ ——► Response to X-radiation | **yagK** ◄—— | yagL ◄—— resolvase like prophage | CP4-57 prophage protein (cell growth/biofilm formation/ motility/ transport) [51] | Unknown function | Protection against radiation-induced stress |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *4. yahE* | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (cause cellular stress) [52] | Putative acyl-CoA synthetase (*yahF*) (energy metabolism) | CYT | Putative NAD(P)-binding succinyl-CoA synthase (SucD) (energy metabolism) | *yahD* → Response to UV | **yahE** → | *yahF* → Putative acyl-CoA synthetase | Unknown function | Unknown function | Energy metabolism under cellular stress |
| *5. yahL* | Cell adhesion | LB, 37°C | Membrane protein (*yahC*) | NUC | Unknown function (YahE) | *yahK* → Aldehyde reductase (involved in biofilm formation) [53] | **yahL** → | *xtpA* → Putative transcr. regulator | Unknown function | Unknown function | Cell adhesion |
| *6. yahM* | Glycine betaine bio-synthesis from choline (osmoprotection) [49] | M9, 37°C; 25 µg/ml Kanamycin | Putative amino acid exporter (*yahN)* (Efflux of toxic sunstances) [54] | CYT | Putative replicase (YidX) | *xtpA* → Putative transcr. regulator (confers antibiotic resistance) [55] | **yahM** → | *yahN* ← Putative amino acid exporter | Unknown function | Unknown function | Protection against stress |
| *7. ybaA* | Co-regulated (unknown regulator or function) | LB, 37°C | Putative oxido-reductase (*yohF*) | CYT | Double strand break repair protein (YegP) | *ffs* → Signal recognition particle | **ybaA** → | *pdeB* ← Cyclic di-GMP phospho-diesterase | RNA signal recognition protein [56] | Unknown function | Oxidoreduction/ DNA damage repair |
| *8. ybaP* | Co-regulated (unknown regulator or function) | LB, 37°C | DNA binding transcriptional activator (*AaeR)* (role in biofilm formation) [57] | CYT | Cysteine-tRNA deacylase (YbaK) | *chiX* → Small regulatory RNA (role in biofilm formation) [58] | **ybaP** ← | *ybaQ* → DNA binding transcr. regulator (regulates biofilm based response) [59] | Conjugal transfer protein | Unknown function | Biofilm formation |
| *9. ybdJ* | Not co-regulated | LB, 37°C | Putative transporter (*yeeA*) (aromatic acid exporter) [60] | PM | DUF419 family protein (YbdF) | *ybdF* ← Defense mechanism | **ybdJ** ← | *ybdK* ← Amino acid bio-synthesis | Unknown function | Unknown function | Stress response |

| Gene | Co-regulation | Conditions | Homolog (putative function) | Loc. | Function | Genomic context (left) | | (right) | Neighbor 1 | Neighbor 2 | Predicted role |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 10. ybeR | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (cause cellular stress) [52] | Putative oxamate carbomyl-transferase (fdrA) | CYT | Putative co-chaperone (DjlB) (heat stress response) | ybeQ ← Oxidative stress response | ybeR → | djlB → J-domain type protein | Unknown function | Unknown function | Multi-stress response |
| 11. yceQ | Cell death | LB, 37°C | DNA binding transcriptional repressor (acrR) – involved in efflux of toxic substances [61] | CYT | Endoribonuclease (RNE) – adaptation to different growth conditions [62] | Rne ← Ribosomal nuclease | yceQ | rluC → 23S rRNA pseudo-uridine synthase | Unknown function | Unknown function | Cell death prevention |
| 12. ycgX | Not co-regulated | LB, no O₂, 37°C | Membrane protein (yigF) | CYT | Repressor of blue light-responsive genes (BluR) | ymgL ← Consti-tutively expressed protein in rich medium | ycgX ← | bluR ← DNA binding transcr. regulator | Prophage protein | Unknown function | Unclear |
| 13. ycgY | Stress response in stationary phase | LB, no O₂, 37°C | Outer membrane protein (yaiO) | CYT | Uncharacterized protein (YncH) | ymgE → DNA Recomb-ination | ycgY → | treA ← Trehalase protein (nutrient stress in stationary phase) [63] | Unknown function | Unknown function | Growth-associated response during stationary phase |
| 14. ychS | Not co-regulated | LB, 37°C; 1 ug/mL of Ciprofloxacin+ 500 µM of ET2DA (cause oxidative/ cellular stress) [64] | Putative amino acid de-hydrogenase (ykfG) – role in oxidative stress [65] | CYT | Not found | narI → Nitrate reductase (oxido-reductase activity) (UniProt) | ychS | tpr ← Protamine like protein | Beta-galactosidase | Protein YchS (Unknown function) | Oxidative stress response |

| # Gene | Co-regulated | Conditions | Function notes | Loc | Protein | Left neighbor | Center | Right neighbor | | | Conclusion |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 15. *yciX* | Not co-regulated | M9, 37°C; *yqjD* over-expression (inhibits cell growth) [66] | Membrane-associated regulatory protein (*ymiA)* (in exponential or tationary phase) [67] | CYT | Membrane-associated regulatory protein (YmiA) | *ymiA* → Membrane-associated regulatory protein | **yciX** → | *ymiC* → Stationary phase adaptation protein | Unknown function | Unknown function | Growth-associated response during stationary phase |
| 16. *ydaF* | Fucose metabolism/DNA damage response (Fucose metabolism is disrupted by prophage) [68] | LB, no O₂, 37°C | Prophage protein (*ydaG)* | CYT | Prophage protein (YdaG) | *sieB* → Phage infection exclusion protein | **ydaF** ← | *ydaG* ← Prophage protein | Prophage protein | Unknown function | Prophage protein involved in disruption of fucose metabolism |
| 17. *ydcD* | Phenylacetate catabolic process (prevalant under anaerobic conditions) [69] | LB, no O₂, 37°C | Cytoplasmic trehalase (*treF)* (osmotic stress) [70] | CYT | Putative peri-plasmic protein (YjbT) | *yncH* → | **ydcD** → | *yncO* ← growth-related protein expressed in rich medium) | dCTP de-aminase | Unknown function | Anaerobic respiration |
| 18. *ydcH* | Phenyl acetate catabolic process (prevalant under anaerobic conditions) [71] | LB, no O₂, 37°C | Transcription (*ygeV)* | CYT | Putative structural protein (YaeH) | *opgD* → Glucans synhesis protein | **ydcH** → | *rimL* → Ribosomal transferase | Unknown function | Unknown function | Anaerobic respiration |
| 19. *ydfA* | Not co-regulated | LB, no O₂, 37°C | Putative transcriptional regulator (*ygeV)* | CYT | Qin prophage (constitutively expressed, unknown function) (YdfB) | *dicA* → DNA binding transcr. regulator | **ydfA** → | *ydfB* → Consti-tutively expressed protein | Phage protein | Unknown function | Unclear |
| 20. *ydfC* | Not co-regulated | M9, 37°C; 0.3 M NaCl, + 0.6% n-butanol (causes osmotic stress) [52] | Putative carbamate kinase (*ycbF)* (biofilm formation) [72] | CYT | Qin prophage (constitutively expressed, unknown function) (YdfB) | *ydfB* → Consti-tutively expressed protein | **ydfC** → | *dicB* → Cell division inhibition protein | Putative phage protein | Unknown function | Cellular stress response |

| Gene | Co-regulation | Growth conditions | Function (gene) | Loc. | Neighbor function | Left gene | Center gene | Right gene | Protein | Function | Conclusion |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 21. ydfX | Not co-regulated | LB, 37°C | Putative glycosyl-transferase (yefI) | CYT | Qin prophage (transcriptional regulator) (DicA) | ydfW ← / Response to antibiotics | **ydfX** ← | dicC ← / DNA-binding transcr. regulator | Bacterial toxin protein | Unknown function | Unclear |
| 22. ydgH | Oxidoreduction (Respiratory complex I) (NADH: ubiquinone oxidoreductase, oxidative stress protection) [73] | LB, 37°C; 1ug/ml of Ciprofloxacin (FC=0.38, relative to frr gene) | Beta-glucosidase (bglA) (Bgl operon confers fitness advantage under stress) [74] | ECS | Dihydro-monapterin reductase (FolM) (role in oxidative stress) [75] | pntA ← / Pyrimidine nucleotide hydro-genase – role in oxidative stress [76] | **ydgH** → | ydgI → / Putative Arginine: ornithine antiporter | YdgH/BhsA/McbA-like domain-containing protein (involved in stress response/biofilm formation and patho-genesis) | Unknown function | Oxidative stress response |
| 23. ydiH | Co-regulated (Unknown/ regulator or function) | M9, 37°C; 10% n-hexane (oxidative stress) [77] | Cyclopropane fatty acyl phospholipid synthase (cfa) | CYT | Putative ferrodoxin-like lipoprotein (YdjY) (oxidative stress response) [78] | menI ← / Mena-quinone bio-synthesis | **ydiH** ← | ydiK ← / Putative trans-porter | Unknown function | Unknown function | Oxidative stress response |
| 24. ydjO | Response to acidic pH (upregulation of oxidative stress response genes) [79] | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | Cysteine/ selenocysteine cation transporter (tcyP) [80] | CYT | Cell division modulator (CedA) (induced in oxidative stress) | tcyP → / Cation transporter | **ydjO** ← | cedA ← / Cell division modulator | Unknown function | Unknown function | Oxidoreduction under acidic pH conditions |
| 25. yeaC | Aerobic respiration | LB, 37°C | Methionine sulfoxide reductase (msrB) (oxidative stress) response [81] | CYT | Methionine sulfoxide reductase B (MsrB) | ydjL / Putative Zn-binding dehydro-genase (oxidative stress response) (UniProt) | **yeaC** ← | msrB ← / Meth-ionine sulphoxide reductase | Transcriptional regulator | Unknown function | Oxidative stress response during aerobic conditions |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *26. yebO* | Cellular response to Magnesium ion (during aerobic respiration) [82] | M9, 37°C; double mutant for genes involved in aerobic respiration *(Δndh ΔcyoB)* [83] | PhoQ kinase inhibitor *(mgrB)* | PM | PhoP/PhoQ signaling protein (MgrB) (induced by low Mg2+) [84] | *yobF* ← General stress response protein | *yebO* ← | *mgrB* ← Kinase inhibitor | Unknown function | Unknown function | Response to magnesium ions during aerobic respiration. |
| *27. yebW* | Not co-regulated | LB, 37°C | Phospho-protein phosphatase *(pphA)* | CYT | Lipopolysaccharide biosynthesis protein (WbbK) | *yebV* → Protein binding | *yebW* ← | *pphA* → Phospho-protein phos-phatase | Lipoprotein | DUF1482 family protein | Unclear |
| *28. yecH* | Anaerobic respiration | M9, 37°C; triple mutant of genes involved in aerobic respiration *(ΔndH, ΔcydB, ΔappC)* [83] | DNA gyrase inhibitor *(sbmC)* (involved in SOS response) [85] | CYT | Putative nucleoside tri-phosphatase (SgcQ) | *ftnA* → Ferritin iron storage protein (oxidative stress response) [86] | *yecH* ← | *tyrP* → Tyrosine symporter | Metal-binding protein | YecH family protein | Switching from aerobic to anaerobic respiration under stress |
| *29. yecT* | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (causes osmotic stress) [52] | Multidrug/cho-line/betaine efflux transporter *(emrE)* (osmotic stress response) [87] | CYT | Putative DNA-binding transcriptional regulator (YhfZ) | *yecV* ← Expo-nential phas protein (in rich medium) | *yecT* → | *flhE* ← Flagellar protein | Lysozyme inhibitor | LprI domain-containing protein (lipo-protein) | Osmotic stress response |
| *30. yegJ* | Cellulose biosynthetic process | LB, no O₂, 37°C | Reactive chlorine species responsive protein *(rclC)* | ECS | Putative non-specific DNA-binding protein (YegI) | *yegI* → Ser/Thr Phos-phatase | *yegJ* → | *pphC* Ser/Thr Phosp-hatase | Unknown function | Unknown function | Unclear |
| *31. yegL* | Not co-regulated | LB, no O₂, 37°C | Zinc-finger domain protein *(yehQ)* | CYT | Ser/thr phosphatase realted protein (YegK) | *pphC* Ser/Thr phos-phatase | *yegL* ← | *yegR* → | Homeostasis protein | Mediator of transcription | Unclear |
| *32. yegR* | Phosphorelay signal transduction system / Oxalate catabolic | M9, 37°C; double mutant for genes involved in aerobic respiration *(Δndh ΔcyoB)* [83] | Putative oxido-reductase *(ydeP)* | ECS | Orphan protein (OrgK) | *ogrK* Trans-cription regulation | *yegR* ← | *yegS* Lipid kinase | Lipoprotein | Unknown function | Oxidoreduction |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | process (involved in acid stress resistance) [88] | | | | | | | | |
| 33. yehI | Not co-regulated | LB, no O₂, 37°C | ATP-binding lipopolysaccharide transport protein (*msbA*) | CYT | Putative ATPase (YehL) | *yehF* → **yehI** → *yehK* → / Molybdate transport regulator | Putative regulator | DUF4132 domain-containing protein | Transport protein |
| 34. yehK | Not co-regulated | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | Putative fimbrial adhesin protein (*sfmH*) | CYT | Putative ATPase (YehL) | *yehI* → **yehk** → *orzP* / Small RNA | Unknown function | Unknown function | Unclear |
| 35. yeiS | Not co-regulated | LB, 37°C | Vanomycin responsive protein (*sanA*) | CYT | Cell membrane barrier (SanA) | *sanA* → **yeiS** → *preT* → / Response to Vanomycin / dehydro-genase | Vancomycin high temperature exclusion protein | Unknown function | Maintaining cell membrane integrity against drugs (e.g. Vanomycin) |
| 36. yfbM | Aerobic respiration | M9, no O₂, 37°C | Hemolysin E (*hlyE*) (pore forming protein) [89] | CYT | Unknown function (YfbN) | *yfbL* → **yfbM** → *yfbN* ← / Putative peptidase (proteo-lysis) (EcoCyc) | Unknown function | DUF1877 family protein | Protein lysis |
| 37. yfbN | Response to acidic pH | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | DNA binding transcriptioal regulator (*ydeO*) | CYT | Unknown function, YfbO | *yfbM* → **yfbN** ← *yfbO* → | Unknown function | Unknown function | Unclear |
| 38. yfbO | Not co-regulated | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | Function unknown (*yfbP*) | ECS | Unknown function (YfbP) | *yfbN* ← **yfbO** → *yfbP* → | Immunity protein (cyto-toxicity) | Unknown function | Unclear |
| 39. yfbP | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol | Inner membrane protein (*ynjI*) | CYT | Putative amino-peptidase (YfbL) (Proteolysis) (EcoCyc) | *yfbO* ← **yfbP** → *nuoN* ← / Oxido-reductase | Protein folding and transport | Unknown function | Protein quality control under stress |
| 40. yfdN | Not co-regulated | LB, no O₂, 37°C | Oxidative stress response protein (*yfdO*) | CYT | Prophage protein (phage-related function) (YfdT) | *yfdM* → **yfdN** ← *yfdO* ← / Putative methyl-transferase / Oxidative stress response | Transcriptional activator family protein | Unknown function | Oxidative stress response |

| # / Gene | Co-regulated | Condition | Function (homolog) | Loc | Prophage protein | Left gene | Center gene | Right gene | Protein function | Additional info | Category |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 41. *yfdP* | Not co-regulated | LB, no $O_2$, 37°C | Putative auto-transporter adhesin (*ycgV*) | CYT | Prophage protein (response to oxidative stress) (YfdS) | *yfdO* ← Oxidative stress response | **yfdP** → | *yfdQ* → Phage-related function | Prophage protein | Unknown function | Oxidative stress response |
| 42. *yfeS* | Cellular response to envelope stress | MOPS, 37°C | Putative lipoprotein (*yidX*) | CYT | Putative efflux protein (YdhK) | *yfeK* → Transcr. regulation | **yfeS** → | *cysM* ← Cysteine synthase | Molybdate metabolism protein | Unknown function | Envelope stress response |
| 43. *yffL* | Response to Nitrate/Nitrite (anaerobic respiration) | M9, 37°C | Anaerobic glycerol-3-phosphate dehydrogenase (*glpB*) [89] | CYT | Prophage protein (Integrase) (IntZ) | *intZ* → Putative integrase | **yffL** → | *yffM* → | Prophage protein | CPZ-55 prophage (involved in general stress response) [90] | Anaerobic respiration |
| 44. *yffM* | Not co-regulated | LB, no $O_2$, 37°C | Alpha-mannosidase (*mngB*) | CYT | Prophage (unknown function) (YffN) | *yffL* → | **yffM** → | *yffN* → | Unknown function | Unknown function | Unclear |
| 45. *yffN* | Not co-regulated | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | Tail fiber protein (*stfR*) | CYT | Prophage (unknown function) (YffM) | *yffM* → | **yffN** → | *yffO* → | Unknown function | Unknown function | Unclear |
| 46. *yffP* | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol | Function unknown (*yffO*) | CYT | Prophage (unknown function) (YffO) | *yffO* → | **yffP** → | *yffQ* → | Unknown function | Unknown function | Unclear |
| 47. *yffQ* | Not co-regulated | MOPS, 37°C; 0.4% Glycerol | CRISPR-associated endo-ribonuclease (*cas2*) (defense response to virus) (EcoCyc) | CYT | Prophage (unknown function) (YffP) | *yffP* → | **yffQ** → | *yffR* ← | Unknown function | CPZ-55 prophage (involved in general stress response) [90] | Stress response |
| 48. *yffR* | Not co-regulated | M9, 37°C; triple mutant of genes involved in aerobic respiration (*ΔndH, ΔcydB, ΔappC*) [83] | Bacterial adhesion/biofilm formation (*yhjY*) | CYT | Prophage (unknown function) (YffQ) | *yffQ* → | **yffR** → | *yffS* → DNA bindin | Unknown function | Unknown function | Stress response |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 49. *yfiM* | Cell death | LB, 37°C | Alpha-ketoglutarate: $H^+$ symporter (*kgtP) (*efflux transporter for sugar-phosphate stress) [91] | ECS | Phospholipid biosynthesis protein (PssA) | *pssA* → Phos-phatidyl-serine synthase | **yfiM** → | *kgtP* ← $H^+$ symporter | Lipoprotein | Unknown function | Lipoprotein with a role in mitigating the effects of cellular stress |
| 50. *yfjH* | Not co-regulated | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose | Growth related protei (*htrL*) | CYT | Prophage (transcriptional activation) (AlpA) | *intA* → Integrase protein | **yfjH** ← | *alpA* → Transcr. regulator | Unknown function | Unknown function | Unclear |
| 51. *yfjJ* | Response to acidic pH (upregulation of genes involved in cell adhesion) [92] | MOPS, 37°C; 0.4% Glycerol and 0.5% 2-Deoxy-D-glucose | Putative fimbrial usher protein (*htrE*) – cell adhesion or pilus assembly protein [92] | CYT | Prophage (cytosolic protein) (YfjI) | *yfjI* → Prophage protein | **yfjJ** → | *abpB* ← Putative helicase | Inovirus-type Gp2 protein (viral DNA replication) | Unknown function | Cell adhesion |
| 52. *ygdT* | Not co-regulated | LB, no $O_2$, 37°C | Putative lipoprotein (*yidX)* | CYT | Not found | *rppH* ← RNA phospho-hydrolase | **ygdT** ← | *mutH* → DNA repair protein | Transposase protein | Unknown function | Unclear |
| 53. *yhaL* | Not co-regulated | LB, 37°C; *Δhns* (stress adaptation) | DNA binding transcriptional activator (*fucR)* | CYT | Ser/thr phosphatase-related protein (Yegk) | *yhaK* → bicupin related protein (acid stress/oxidative stress response) [93] | **yhaL** → | *cyuA* ← L-cysteine di-sulphidase | Cytoplasmic protein | Unknown function | Stress response |

| Gene | Co-regulation | Condition | Adjacent gene function | Loc. | Protein function | Left gene | Center gene | Right gene | Col A | Col B | Predicted function |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *54. yhdN* | Response to heat | LB, 30°C (stationary phase) | DNA binding transcriptional activator (*zntR)* (stress responsive transcription factor) [94] | CYT | Transcriptional activator (ZntR) | *zntR* ← DNA binding transcr. regulator | **yhdN** ← | *rplQ* ← 50S ribosome subunit protein (protein bio-genesis) [95] | Conserved protein | DUF1992 domain-containing protein (conserved protein belonging to the DNAJ homolog family, heat shock response) (UniProt) | Regulation of heat shock response, expressed during stationary phase |
| *55. yhfU* | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (osmotic stress) [53] | Aromatic carboxylic acid efflux pump (*aaeA)* (involved in defense mechanisms) [96] | CYT | Putative transport protein (YhfU) | *yhfT* ← Membrane protein | **yhfU** ← | *php* ← Putative hydrolase | Unknown function | Unknown function | Osmotic stress response |
| *56. yhhL* | Not co-regulated | LB, 37°C; *ΔdsrA* (gene involved in stress response, biofilm formation) [97] | 16S rRNA methyl-transferase (*rsmD)* | CYT | 16S rRNA methyl-Transferase (RsmD) | *rsmD* → 16S rRNA methyl transferase | **yhhL** xs | *yhhM* Quorum sensing protein – (stress response, biofilm formation) [98] | rRNA small subunit methyl-transferase protein | Unknown function | Biofilm formation |
| *57. yibI* | Response to nitrosative stress (redox stress) [99] | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol (osmotic stress) [52] | Inner membrane protein (*yibH)* (stress response) [100] | PM | Inner membrane protein (YibH) | *yibH* ← Inner membrane protein | **yibI** ← | *mtlA* → mannitol-specific PTS enzyme | Inner membrane protein | Unknown function | Stress response protein. |
| *58. yicN* | Not co-regulated | LB, 37°C; *Δhns* (stress adaptation) [101] | DNA binding transcriptional regulator (*frlR)* | CYT | Putative Transporter (NepI) | *nepI* ← Purine ribo-nucleoside exporter | **yicN** ← | *adeQ* Adenine trans-porter | Unknown function | Unknown function | Nucleotide transport |

| # Gene | Co-regulation | Conditions | Protein | Loc. | Characterization | Genomic context | | | Left neighbor func. | Right neighbor func. | Conclusion |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 59. yidB | Not co-regulated | LB, 37°C; triple mutant of genes involved in respiration (ΔndH, ΔcydB, ΔappC) [83] | Bitopic inner membrane protein (ypfJ) | CYT | Sugar phosphate phosphatas (YidA) (phosphate metabolism) [102] | yidA ← Sugar phos-phatase | yidB ← | gyrB ← DNA gyrase | Radical SAM domain-containing protein | Unknown function | Unclear |
| 60. yigA | Not co-regulated | LB, 37°C | Diamino-pimelate epimerase (dapF) (cell wall biosynthesis) [103] | CYT | Diamino-pimelate epimerase (DapF) (cell wall structure) [104] | dapF → diamino-pimelate epimerase | yigA → | xerC → Tyrosine recombi-nase | Diamino-pimelate epimerase protein | Not found | Cell wall biosynthesis |
| 61. yigE | Not coregulated | LB, no O$_2$, 37°C | Putative magnesium chelatase (yifB) (chaperonic activity) [105] | CYT | Inner membrane protein (YedI) | uvrD → DNA helicase | yigE ← | ysgD → Consti-tutively expressed protein (in rich medium) | Phospho-diester glycosidase family protein | Anaerobic nucleoside-tri-phosphate reductase-activating protein (DNA synthesis and repair) [106] | Stress response during anaerobic conditions |
| 62. yjbD | Not coregulated | M9, 37°C; 10% n-hexane | Putative methyl-transferase (yafS) | CYT | Cysteine cluster protein (YcgN) | rluF → Pseudo-uridine synthase | yjbD ← | lysC ← Aspartate kinase | Unknown function | Unknown function | Unclear |
| 63. yjbL | Response to acidic pH | LB, no O$_2$, 37°C | Cell envelope stress response protein (yqcG) | CYT | Uncharacterized protein (YjbM) | zur → Transcr. regulator | yjbL → | yjbM → Cell-envelope related protein | Unknown function | Unknown function | Envelope stress response under stress |
| 64. yjbS | Response to acidic pH (upregulation of oxidative stress response genes) [107] | M9, 37°C; 10µg/ml Mitomycin-C (genotoxic agent, ROS producing) [108] | Pathogenicity-related (yqeK) | CYT | Putative oxido-Reductase (YqiJ) ([109] | tyrB → Tyrosine amino-transferase | yjbS ← | aphA → Acid phos-phatase | Unknown function | Unknown function | Oxidative stress response. |
| 65. yjcB | Carbohydrate transport | M9, no phosphate, 37°C; ΔphoB (phosphate transport gene) | Capsule synthesis regulator (rcsB) | CYT | Putative peri-plasmic protein (YpeC) | ssb → ss-DNA binding protein | yjcB ← | pdeC → Cyclic di-GMP Phospho-diesterase | Membrane protein | Not found | Unclear |

| 66. *yjcF* | Response to acidic pH | LB, 37°C | Pathogenicity-related (*ygeH*) | CYT | Ribosomal silencing factor (RsfS) | *yjcE* → Putative transporter | **yjcF** ← | *actP* ← Cation trans-porter | Unknown function | Unknown function | Unclear |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 67. *yjdI* | Oxidoreduction (Respiratory complex I) | LB, 37°C; triple mutant of genes involved in respiration *(ΔndH, ΔcydB, ΔappC)* | Putative N-acetyl-Transferase (*yjdJ*) | CYT | Putative N-acetyl-Transferase (YjdJ) | *dcuS* ← Sensor histidine kinase | **yjdI** → | *yjdJ* → Putative acetyl-trans-ferase | Unknown function | Fer4_19 domain-containing protein (involved in iron-sulphur binding) | Aerobic respiration |
| 68. *yjeJ* | Cell adhesion | LB, 37°C; HS, 50°C for 7' | 6-phospho-gluconate phosphatase (*yieH*) | CYT | Lysylation protein (EpmB) | *yejI* → Lipoprotein (heat shock response) [110] | **yjeJ** ← | *epmB* ← Lysine amino-mutase | Unknown function | Unknown function | Cell adhesion under heat shock response |
| 69. *yjeT* | Not coregulated | LB, 37°C; HS, 50°C for 15' | Lipo-polysaccharide core heptose kinase *(waaP)* | CYT | HflB protease modulator (HflC) (in heat shock response) [111] | *hflC* → ftsH protease regulator (heat shock response) [111] | **yjeT** → | *purA* → Adenylo-suucinate syn-thetase | Unknown function | Unknown function | Heat shock response |
| 70. *yjfI* | Coregulated (unknown regulator or function) | LB, 37°C, reaction buffer (with Lysine) | Inner membrane protein (*yjgN*) | CYT | Membrane chaperone family protein (YjfJ) [112] | *rlmB* → Methyl-transferase | **yjfI** → | *yjfJ* → Transcr. regulator | Cytoplasmic protein | Unknown function | Unclear |
| 71. *yjfK* | Not coregulated | LB, no O$_2$, 37°C | Membrane chaperone family protein (*yjfJ*) | CYT | Inner membrane protein (YjfL) | *yjfJ* → Transcr. regulator | **yjfK** → | *yjfL* → Inner membrane protein | Unknown function | Unknown function | Unclear |
| 72. *yjfZ* | Not coregulated | LB, 37°C | Acyl transferase (*yihG*) | CYT | Putative inner membrane poly-saccharide flippase (YghQ) | *rpiI* → Ribosomal subunit protein | **yjfZ** ← | *ytfB* ← Cell division | Unknown function | Unknown function | Unclear |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| *73. yjgR* | Not coregulated | LB, no O₂, 37°C | NADPH dependent aldehyde reductase (*ahR)* | PM | L-idonate and D-gluconate transporter (IdnT) | *iptG* → / **yjgR** ← / *idnR* ← ; LPS Transport system / DNA binding transcr. regulator | Ornithine amino-transferase | Conjugal transfer protein | Transport |
| *74. yjgZ* | Coregulated (unknown regulator or function) | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol | Putative multidrug efflux pump (*mdtO*) | CYT | Putative receptor protein (YhhM) | *insD6* → / **yjgZ** → / *insG* ← ; Trans-posase / Putative transposase | Transposase | Unknown function | Unclear |
| *75. yjjY* | Cell death | LB, 37°C | DNA binding transcriptional regulator (*arcA)* | CYT | Uncharacterized protein YniD | *arcA* ← / **yjjY** → / *ytjE* → ; DNA binding transcr. regulator / Growth related protein | Unknown function | Unknown function | Unclear |
| *76. ykgL* | Glycine betaine biosynthesis from choline [49] | M9, 37°C; *ΔkdpE*; Tris-maleic acid+115 mM KCl (disrupts potassium uptake under osmotic stress) [113] | 50S ribosomal subunit protein (*ykgM*) | CYT | Carnitinyl-CoA dehydratase (CaiD) – osmoprotectant protein [114] | *matA* ← / **ykgL** → / *ykgO* ← ; DNA binding transcr. regulator / 50S ribosomal subunit protein | Unknown function | Unknown function | Osmotic stress response. |
| *77. ylcH* | Trans-criptional attenuation | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol | Ionic detergents response protein (*yfbK*) | No inf. | Uncharacterized protein (YlcG) | *ybcM* → / **ylcH** → / *ybcN* → ; DNA binding transcr. regulator / DNA base-flipping protein | Unknown function | Unknown function | Unclear |
| *78. ylcI* | Aerobic respiration | LB, 37°C; double mutant of genes involved in respiration (*ΔndH, ΔcyoB*) | Phage related protein (*ybcW*) | CYT | Oxidoreductase (NuoM) – involved in aerobic respiration [115] | *ybcW* → / **ylcI** ← / *nohD* → ; Prophage protein / DNA packing protein | Unknown function | DUF3950 domain-containing protein | Aerobic respiration |
| *79. ymfI* | DNA restriction/ modification system | LB, 37°C; 1 ug/mL of Ciprofloxacin+ 500 µM of ET2DA | Phage related protein (*ymfJ*) | CYT | Uncharacterized protein (*ymfJ*) | *ymfH* ← / **ymfI** → / *ymfJ* ← ; Prophage protein / Prophage protein | Phage protein | Unknown function | Unclear |
| *80. ymgI* | Not co-regulated | M9, no phosphate, 37°C; *ΔphoB* (phosphate transport gene) | Heat shock responsive protein (ymgG) | CYT | Cell division topological specificity protein (MinE) | *ymgG* ← / **ymgI** ← / *ymgJ* → ; Heat shock response protein | Unknown function | Unknown function | Unclear |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *81. ymgJ* | Not co-regulated | M9, 37°C; 0.3 M NaCl + 0.6% n-butanol | Putative fimbrial usher protein (*htrE*) (heat shock response) [116] | CYT | Heat shock response protein (YmgG) | *ymgI* ← | ***ymgJ*** → | *minE* ← Cell division | Transposase protein | Unknown function | Heat shock response |
| *82. ynaJ* | Aerobic respiration | LB, 37°C | ADP-sugar diphosphatase (*nudE*) | CYT | Translation-realted protein (YciU) | *insH4* → Trans-posase | ***ynaJ*** → | *uspE* ← Universal stress protein | Inner membrane protein | Unknown function | Unclear |
| *83. yncH* | Response to acidic pH | LB, 37°C; *rpoD* overexpression (response to heat) [117] | Cell envelope stress response protein (yqcG) | CYT | Glutathione S-transferase homolog (YncG) | *yncG* → Putative Glutata-thione-S transferase | ***yncH*** → | *yncO* ← Consti-tutively expressed protein (in rich medium) | Unknown function | Unknown function | Stress response |
| *84. yneG* | Not co-regulated | M9, no ammonium, 37°C | Glutaminase (*glsB*) Ammonium assimilation [118] | CYT | Putative Glutaminase (GlsB) (amino acid metabolism) [119] | *dgcF* ← Putative Di-guanylate cyclase | ***yneG*** ← | *glsB* ← Glutaminase | Cytoplasmic protein | Unknown function | Nitrogen metabolism during ammonium limitation |
| *85. yneK* | Not co-regulated | MOPS, 37°C; 0.4% Glycerol + 0.5% 2-Deoxy-D-glucose (metabolic stress) [120] | Putative fimbrial usher protein (*htrE*) (heat shock response) [116] | CYT | Putative oxido-reductase (YdhF) | *ptrR* → L-glutamate meta-bolism | ***yneK*** → | *ydeA* → L-arabinose exporter | Unknown function | Unknown function | Stress response |
| *86. ynfO* | Anaerobic respiration | LB, 37°C; 100 µg/ml Ampicillin | Function unknown (*ylcI*) | CYT | Putative (D)-galactarate transporter (GarP) | *nohA* ← Prophage protein | ***ynfO*** → | *ydfO* → Prophage protein | Unknown function | DUF3950 domain-containing protein | Unclear |
| *87. yoaC* | Not co-regulated | LB, 37°C | Lipoprotein (*yfiL*) | CYT | Protein export (MsyB) | *yoaB* → Biofilm based activity | ***yoaC*** → | *yoaH* ← | Unknown function | Unknown function | Unclear |

| | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *88. yoaH* | Not co-regulated | LB, 37°C; *Δhns* (stress adaptation) | DNA-binding transcriptional repressor *(acrR)* | CYT | DUF1889 family protein (YoaC) | *yoaC* → | **yoaH** ← | *pabB* → Amino-deoxy-chorismate synthase | Unknown function | Unknown function | Unclear |
| *89. yodC* | Oxidoreduction (Respiratory complex I) | LB, 37°C | Putative adenosine mono-phosphate *(fic)* | CYT | Ribosome modulation protein (RmF) (response to oxidative stress) [121] | *dgcQ* ← Di-guanylate cyclase | **yodC** ← | *yedI* ← Inner membrane protein | Cytoplasmic protein | Unknown function | Oxidoreduction |
| *90. ypaB* | Not coregulated | M9, 37°C; 0.3 M NaCl +0.6% n-butanol | Putative fimbrial chaperone *(yadV)* | CYT | Not found | *yfaL* ← Putative auto-transporter adhesin | **ypaB** ← | *nrdA* → Ribo-nucleoside reductase | Transposase | Unknown function | Unclear |
| *91. ypeB* | Ribosome structure | M9, 37°C; 50 mM isoprenol | Tetraacyl-disaccharide 4'-kinase *(lpxK)* | CYT | DNA ligase (LigA) | *yfeH* → Putative Na⁺ symporter | **ypeB** ← | *ligA* ← DNA ligase | Cytoplasmic protein | Cyto-plasmic protein | Unclear |
| *92. ypjB* | Not coregulated | LB, no O₂, 37°C | Putative type III secreted effector *(yqeK)* (bacterial patho-genecity) [122] | CYT | Not found | *ypjA* ← auto-transporter | **ypjB** ← | *ypjC* → Patho-genicity | Uncharacterized protein | Unknown function | Pathogenecity |
| *93. yrdB* | Oxidoreduction (Respiratory complex I) | LB, 37°C | Shikimate dehydro-genase *(aroE)* | CYT | Dehydro-shikimate reductase (AroE) - oxidative stress response) [123] | *yrdA* → carbonic anhydrase | **yrdB** ← | *aroE* ← shikimate anhydrase | DUF1488 domain-containing protein | Unknown function | Oxidoreduction |
| *94. yrhD* | Response to acidic pH | LB, no O₂, 37°C | Putative type III secreted effector *(yqeK)* (bacterial patho-genecity) [122] | CYT | Not found | IS1E → Insertion sequence | **yrhD** → | *yrhB* → Putative HS chaperone | Unknown function | Unknown function | Unclear |
| *95. ytiA* | Not co-regulated | LB, no O₂, 37°C | Putative transporter *(yjiJ)* | CYT | Not found | *yjiK* ← Membrane | **ytiA** ← | *yjiL* ← Putative de- | DUF3343 domain-containing | Unknown function | Unclear |

| | | | | | | protein | hydratase | protein | | |
|---|---|---|---|---|---|---|---|---|---|---|

Abbreviations: ATP, Adenosine triphosphate; dCTP, Deoxycytidine triphosphate; ET2DA, Ethyl trans-2-decenoic acid; HS, Heat shock; KCl, Potassium chloride; LB,

Luria Bertani; LPS, Lipopolysaccharide; M9, Minimal medium; MOPS, 3-(N-morpholino) propanesulfonic acid; NADP, Nicotinamide dinucleotide phosphate; PTS,

phosphotransferase system; ROS, Reactive oxygen species; SAM, S-adenosylmethionine; Ser/Thr, Serine/Threonine; Transcr. , Transcription

Genes with a higher confidence in the inference section (green) were analysed for candidates

to be used for *in vitro* testing. Three HP-encoding genes, namely *yhdN*, *yeaC*, and *ydgH*, were

selected for further analysis (highlighted in grey) (**Table 1).**

**Protein encoding gene *yhdN***

Protein YhdN, characterized by the presence of a domain of unknown function (DUF1992),

was predicted to be involved in heat shock response. The gene encoding *yhdN* is regulated

by the RNA polymerase sigma factor RpoH (σ32), which serves as the primary heat shock

transcriptional regulator. Under normal conditions, RpoH levels are kept low due to chaperone-

mediated degradation; however, these levels increase significantly upon heat shock [124].

Gene Ontology (GO) annotation for this cluster is categorized under 'response to heat'. Based

on publicly available transcriptomic data [43], the *yhdN* gene exhibited significant expression

in cells grown to the stationary phase, with a $Log_2FC$ of 5.5 relative to the reference gene *frr*,

signifying a 45 times higher expression of *yhdN* gene. This observation suggests that

the *yhdN* gene is highly expressed during the stationary phase and therefore might have a

role in nutrient stress response. Additionally, *yhdN* expression was upregulated following

exposure to a transient heat shock at 50°C for 15 minutes, with a $Log_2FC$ of 4.8 and 27 times

higher expression, suggesting that the gene might also play a role in heat shock response.

Co-expression analysis using Spearman correlation of the gene resulted in a high correlation

with the gene *zntR* (0.93) encoding an HTH (helix-turn-helix)-type transcriptional regulator

[125] which is involved in protein stability and prevention of aggregates [126,127]. Genes *yhdN*

and *zntR* are also part of the same transcription unit (EcoCyc database). Other genes with

high correlations to *yhdN* were *relB* (0.84)*, pspB* (0.84) *and pspC* (0.84), all of which are

involved in stress response [128] (*Supplementary Figure 3A.*). Gene *relB* encodes for an

antitoxin protein involved in regulation of growth [129] while genes *pspB* and *pspC* encode for

phage shock proteins that are known to be induced by infection with bacteriophages or

ethanol, osmotic and/or nutrient stress as well as heat shock conditions [130]. Protein-protein

interaction (PPI) analysis showed highest interaction confidence with ZntR. The local genomic

context analysis showed genes *zntR* and *rplQ* (a ribosomal subunit protein) as neighbouring

genes. Protein YhdN showed remote sequence homology to a protein cluster comprising of

uncharacterized conserved proteins and was structurally homologous to conserved proteins

with similarity to J-domains of chaperones, regulating the activity of heat-shock proteins [131].

To elucidate the potential function of *yhdN*, we hypothesized that its absence would affect

bacterial growth under heat shock conditions and possibly the cells stationary phase under

nutrient limitation. This hypothesis was tested by comparing the growth patterns of the

*E. coli* K-12 substrain BW25113 wild type and its isogenic *ΔyhdN* knockout mutant under a

transient heat shock condition (50°C) and recovery at 37°C in LB and nutrient-limited M9

medium.

**Protein encoding gene *yeaC***

Protein YeaC is characterized by the presence of a DUF1315 domain. Gene *yeaC* is predicted

to be governed by the regulator ArcA, which regulates a group of proteins involved in redox

homeostasis and aerobic respiration [132]. The Gene Ontology (GO) annotation for this cluster

was predicted as 'aerobic respiration'. Analysis of gene *yeaC* in publicly available

transcriptomic datasets showed slightly higher expression levels against gene *frr* in LB

medium at 37°C under normal $O_2$ conditions ($\log_2$ FC 1.9) (**Table1**) and in knock-out mutants

for genes encoding type II NADH:quinone oxidoreductase (*ndh*), Cytochrome bd-I ubiquinol

oxidase (*cydB*) and Cytochrome bd-II ubiquinol oxidase (*appC*), involved in the Tricarboxylic

acid cycle or oxidative phosphorylation [133] ($\log_2$ FC 1.7). Based on the co-expression

analysis of public data, the highest correlation was observed with a gene encoding for a

peptide methionine sulfoxide reductase (*msrB),* which is an oxidoreductase and known to be

highly expressed during oxidative stress to maintain the cell-redox homeostasis [134]

(*Supplementary Figure3B.*). Other highly correlated genes involved succinate dehydrogenase

cytochrome b556 subunit (*sdhC*), succinate dehydrogenase hydrophobic membrane anchor

subunit (*sdhD)* and Fumarate hydratase class I, aerobic (*fumA),* which are all involved in the

tricarboxylic acid or TCA cycle [135,136] (*Supplementary Figure 3B.*). PPI analysis also

showed highest interaction confidence with MsrB. Additionally, it neighbours the *msrB* gene

and a putative zinc-binding dehydrogenase, *ydjL* which is also involved in oxidoreduction

(UniProt) (**Table 1**). Protein YeaC shows remote sequence homology to other proteins

involved in transcriptional regulation. Structural homology revealed similarities to a group of

uncharacterized proteins with unknown functions. We hypothesized that *yeaC* affects the cell-

redox homeostasis and cellular growth under oxidative stress. This hypothesis was tested by

comparing the growth patterns of the *E. coli* K-12 substrain BW25113 wild type and its isogenic

*ΔyeaC* knockout mutant with a sub-lethal concentration of $H_2O_2$ (2.5 mM) [137].

**Protein encoding gene *ydgH***

Protein YdgH has a DUF1471 domain of unknown function. It is predicted to be part of an

unknown cluster, with no information on its regulator. The GO annotation for this cluster,

however, is categorized under oxidoreduction (complex I). The energy-converting

NADH:ubiquinone oxidoreductase respiratory complex I, is the main entry point for electrons

from NADH into the respiratory chains in bacteria [138]. Based on the public datasets, we

could not obtain a gene-specific condition where gene *ydgH* had higher expression compared

to gene *frr.* Co-expression analysis of *ydgH* from the publicly available RNAseq datasets

revealed highest correlation with a 6-phospho-beta-glucosidase (*bglA*) encoding hydrolase,

involved in the hydrolysis of phosphorylated beta-glucosides [139]. Other highly correlated

genes were phosphoglycolate phosphatase *(gph)* [140] and ribulose-phosphate 3-epimerase

*(rpe)* [141], involved in oxidative stress response (*Supplementary Figure 3C.*). PPI analysis

showed highest interaction confidence with YjfY*,* a putative HP, with a possible role in stress

resistance (UniProt). Genomic context provided information on a neighbouring upstream gene

involved in NAD(P)$^+$ transhydrogenase activity, annotated as NAD(P) transhydrogenase,

*pntA*, which is known to play a role in oxidative stress [142]. The protein showed remote

sequence homology to a YdgH/BhsA/McbA-like domain involved in stress response/biofilm

formation and pathogenesis [143]. The structural similarity was only to uncharacterized

proteins with unknown function. Following a similar hypothesis to that proposed for *yeaC*, we

evaluated the growth patterns of the wild type and its isogenic *ΔydgH* knockout mutant under

oxidative stress.

### *In vitro* analyses of candidate proteins

Effect of heat shock on growth

To determine the effect of a transient heat shock at 50°C (7 minutes), the wild type and the

mutant *ΔyhdN E. coli* BW25113 strains were grown in LB medium. The *yhdN* mutant was seen

to have a growth defect of ~18% after 6h of growth based on the OD values obtained (**Figure**

**3A**). It was observed that after 6h of growth, the mutant could not resume normal growth

compared to the wild type. Additionally, the effects of a transient heat shock in Nitrogen-limited

M9 medium showed a significant growth defect of ~50% (**Figure 3B**). Growth curves for WT

and *ΔyhdN* in different temperatures can be found in the supplementary (*Supplementary*

*Figure 4A.*). Growth defects were also observed in the *yhdN* mutant compared to the wild type

when both the strains were grown continuously at 50°C. Since growth at 50°C is however

severely inhibited and cell started dying after 3h (*Supplementary Figure 4B.*), it was difficult to

say whether this was solely based on the absence of the protein YhdN in the mutant cells.

Effect of sublethal $H_2O_2$ concentration on growth

To determine the effect of sublethal $H_2O_2$ concentration, the wild type (WT), *ΔydgH* and *ΔyeaC*

*E. coli* BW25113 strains were grown in LB medium until an OD of 0.2 and then exposed to an

$H_2O_2$ concentration of 2.5 mM, which is known to causes DNA damage and oxidative stress

[143] (**Figure 3C & D**). To study the responses due to oxidative stress, the absorbance values

($OD_{600}$) after each hour over a 5h time period were measured. $H_2O_2$ was added to the cell

culture during the exponential phase of growth, which is characterized by rapid cell division

and metabolic activity and where the quorum-sensing signalling, i.e. process which allows the

bacteria to communicate and adjust gene expression according to cell density, is minimal.

Under non-stress conditions, the WT and mutant strains (*ΔydgH* and *ΔyeaC*) exhibited

comparable growth, with no significant differences observed.  A growth lag was observed in

the mutants (*ΔydgH* and *ΔyeaC*) as compared to wild type when the medium was

supplemented with 2.5 mM $H_2O_2$. A growth defect of ~25% was observed in the mutant when

WT was compared to *ΔyeaC* at the 1h timepoint. This was reduced to ~7% after 5h of growth

underscoring the probability of the gene being expressed at this time point to counterfeit the

oxidative stress. Similarly, a growth defect of ~36% was observed in *ΔydgH* as compared to

WT at 1h, which decreased to ~2% after 5h, again probably due to stress adaptation and

expression of gene *ydgH*.

372
373
**Figure 3. Growth curves of *E.* coli K-12 BW25113 wild type (WT) and isogenic deletion mutants.**
**(A, B)** Effect of transient heat shock on WT and *ΔyhdN* cells. WT and the respective mutant strain were
exposed to a transient heat shock at 50°C for 7 minutes, (at $OD_{600}$ nm = 0.04) and grown for 12 h in **A.**
LB medium and **B.** in Nitrogen-limited M9 medium. **(C, D)** Bacterial growth of WT and respective
mutants at 37°C exposed to sub-lethal concentration of 2.5 mM $H_2O_2$, added during the exponential
phase ($OD_{600}$ = 0.2) and grown for 5h. **C.** WT and *ΔyeaC* and D. WT and *ΔydgH* cells. WT are indicated
by filled circles (●) and mutants by filled triangles (▲). Green lines represent controls, orange lines stress
conditions. Average of three independent readings taken for each specified condition. Error bars on the
graph indicate standard deviation from the mean.

**Differential gene expression analysis**

For a comparative analysis of differentially expressed genes (DEGs) and their functions in WT

*vs.* the mutant strains, all up- and down-regulated genes were analysed (*Figshare > DEGs >*

*Table1*). For positively regulated DEGs, representing genes upregulated in the WT strain, a

Log$_2$ fold change (Log$_2$FC) threshold of ≥1.5 was applied, corresponding to a minimum 2.8-

fold increase in expression levels. This cut-off ensured the inclusion of genes with significant

upregulation while minimizing noise [144]. Conversely, for negatively regulated DEGs,

corresponding to genes upregulated in the mutant strains, a Log2FC threshold of ≤-1.5 was

used.

WT *vs. ΔyhdN*

Analysis of the WT strain at the 2h and 6h time points showed that *yhdN* was only significantly

expressed after 6h, hence this time point was chosen. In total, 484 DEGs ($p_{adj}$≤0.1) were

identified, (181) 213 genes with a positive Log$_2$ FC and 271 with a negative Log$_2$FC of 1.5.

Within the positive DEGs, an analysis of the top 10 genes, resulted in one gene involved in

protein folding, five in stress response and four in motility/chemotaxis (**Figure 4A**).

Significantly upregulated genes in the WT strain were *tdcB* (Log$_2$ FC=7.37) and *pyrB* (Log$_2$

FC=6.14). *tdcB*, encoding for threonine dehydratase is known for being expressed during

nutrient deprivation (EcoCyc) and gene *pyrB* is involved in pyrimidine metabolism and

associated with motility under stress [145]. Gene *tnaA* (Log$_2$ FC=6.02), which encodes for

tryptophanase, has been recently studied for its role in protein folding and dissolution of protein

aggregates [146]. Motility and chemotaxis genes such as *fliL* (Log$_2$ FC=6.24)*, flgD* (Log$_2$

FC=5.99) are also known to be involved in response to environmental stresses [147].

In the deletion mutant (*ΔyhdN*), the top ten negative DEGs included two genes involved in

heat shock, six in some type of stress response and interestingly two other were

uncharacterized hypothetical proteins. *bolA* (Log$_2$ FC=-5.53)*,* a DNA binding transcriptional

regulator, plays a role in cellular stress response including heat stress [148] and *mgtS* (Log$_2$

FC=-5.01), a small inner membrane protein encoding gene, is known to be expressed upon

heat shock [149] (**Figure 4A**). Other upregulated genes in *ΔyhdN* involved *glgS* (Log$_2$ FC=-

5.34), encoding for a surface composition regulator for biofilms and *ymgA* (Log$_2$ FC=-4.44)

encoding a putative two-component system connector protein, known to play an important role

in biofilm formation (EcoCyc). Upregulation of these genes in the deletion mutant suggests a

compensatory mechanism to mitigate stress. Additionally, a HP-encoding gene, *yodC* was

also upregulated (Log$_2$ FC=-6.39) in *ΔyhdN*. Interestingly, this gene was predicted to play a

role in oxidoreduction (**Table1)**. The results of this transcriptomic experiment could further aid

in forming a hypothesis of this gene for *in vitro* testing. Another HP-encoding gene *yffR* (Log$_2$

FC=-4.85)*,* was predicted to play a role in stress response (this study, Table 1) which aligns

with the fact that a majority of stress-response genes were upregulated in the WT, when gene

*yhdN* is present.

All observations highlight WT strain's prioritization of growth and motility, possibly facilitating

better environmental sensing and resource acquisition under stress while the absence of *yhdN*

gene seems to upregulate other heat shock and stress response genes which reorient

metabolic processes towards alternative survival strategies such as biofilm formation. Hence,

the protein encoding gene *yhdN* is predicted to be a stress response gene involved in the

regulation of heat shock or nutrient limitation response, possibly also playing a broader role in

general stress response.

WT *vs. ΔyeaC*

In WT strain, the *yeaC* gene exhibited a significant upregulation after the 5h, with a log$_2$ FC of

3 relative to the 1h time point, where its expression was undetectable. This log$_2$FC

corresponds to an 8-fold increase in expression. Hence, the 5h time point was chosen for the

analysis. A total of 156 DEGs for WT *vs. ΔyeaC* were obtained after filtering ($p_{adj}$≤0.1). In WT

*vs. yeaC*, 87 genes had a positive Log$_2$ FC while 69 had a negative Log$_2$ FC.

Analysis of the top ten upregulated DEGs in the WT strain resulted in three oxidative stress

response genes, two other stress response genes, two methionine biosynthesis genes and

three genes involved in bacterial secretion system and transposition (EcoCyc). The

upregulation of the latter three genes in WT was difficult to explain, since there was no

conclusive literature available (**Figure 4B**). Significantly upregulated DEGs involved *ymcE*

(Log$_2$ FC=6.81) which provides tolerance to n-butanol [145], and is also known to provide

protection against oxidative stress [150], *prpE* (Log$_2$ FC=4.75) which responds to hydrogen

peroxide [151] and *iprA* (Log$_2$ FC=4.34), which encodes for an inhibitor of hydrogen peroxide

[152]. *metF* (Log$_2$ FC=4.49), a methionine biosynthesis gene, is also known to provide

protection against oxidative stress: Methionine is a precursor in the synthesis of cysteine,

which is a component of glutathione—a major intracellular antioxidant. Glutathione is known

to neutralize reactive oxygen species [153].

All of the upregulated genes in *ΔyeaC* were involved in motility/chemotaxis-based functions

(**Figure 4B**). Significantly upregulated DEGs included genes like *fliF* (Log$_2$ FC=-9.86), *fliO*

(Log$_2$ FC=-9.26) and *fliJ* (Log$_2$ FC=-9.26). The upregulation of these genes in *ΔyeaC* signify

that the cells prioritize the expression of motility-based genes. They are known to aid the

bacterium in adapting to environmental changes by modulating their expression [154].

Moreover, there is often a trade-off between motility and stress resistance. Studies have

shown that hypermotile *E. coli* strains, which exhibit increased expression of motility genes,

may have reduced expression of stress resistance genes, as was also observed in the

downregulated DEGs in the WT. Hence, as opposed to WT, the deletion mutant prioritizes

motility and chemotaxis to mitigate oxidative stress.

WT *vs. ΔydgH*

The *ydgH* gene demonstrated an upregulation after the 5h, with a log$_2$FC of approximately 3.7

as compared to 1 hour, indicating a 13-fold increase in expression. Hence, also in this case

the 5h time point was chosen for further analysis. A total of 231 DEGs were obtained after

filtering ($p_{adj}$≤0.1), resulting in 111 genes with a positive Log$_2$ FC and 120 genes with a negative

Log$_2$ FC.

Top ten positive DEGs (Log$_2$ FC≥1.5) involved six stress response genes, one protein

transport gene, one membrane biosynthesis gene, one fatty acid oxidation gene and one HP-

encoding gene (**Figure 4C**). Significantly upregulated genes were *dsrB* (Log$_2$ FC=3.89),

encoding for a putative stress response protein [155] and *ghoS,* (Log$_2$ FC=3.65), an antitoxin

protein which prevents cell death [156]. Gene *fadM* (Log$_2$ FC=2.4), encoding a thioesterase,

is responsible for breaking down fatty acids into acetyl-CoA units, which can then be utilized

in energy production [157]. Another gene, *tatE* (Log$_2$ FC=2.69) encoding for a transporter, is

involved in transport of folded proteins and maintaining of cellular redox balance [158]. Gene

*yoaH,* a HP-encoding gene whose function was previously unclear from **Table 1**, might also

play a role in oxidative stress response and should be further investigated. The role of

lipopolysaccharide biosynthesis gene *ais* (Log$_2$ FC= 2.41) was, however, unclear.

In the deletion mutant (*ΔydgH*), the top ten negative DEGs included eight motility/chemotaxis

genes along with one sulphur metabolism and one urate transport gene (**Figure 4C**): *fliF* (Log$_2$

FC=-6.92) and *fliA* (Log$_2$ FC=-5.18) play a crucial role for motility (EcoCyc), *uacT* (Log$_2$ FC=-

6.22) is involved in urate transport, which has already been studied to reduce the effects of

oxidative stress in *E. coli* [159]. The role of gene *cysP* regarding oxygen stress, which is

involved in sulphur metabolism [160] is unclear at the moment and calls for further

investigation.

In WT*,* the upregulation of genes involved in response to stress, i.e. *ghoS* suggest that the

cells prioritize conservation of resources by entering a dormant state, i.e. formation of persister

cells [156]. Due to the absence of *ΔydgH* the cell seems to attempt to compensate for the loss

of oxidoreductive function by utilizing genes related to motility and chemotaxis (**Figure 4C**).

This increased expression could be an adaptive response to maintain survival under

continuous exposure to oxidative stress. Hence, similar to the previous case (WT *vs. ΔyeaC*),

more stress response genes were upregulated in WT while motility/chemotaxis genes were

upregulated in *ΔydgH*.

**Figure 4. Comparative differential gene expression analysis in A. wild type (WT) *vs. ΔyhdN*, B. WT *vs. ΔyeaC* and C. WT *vs. ΔydgH* strains.** Top ten positively and negatively expressed DEGs are shown. Each row represents a gene, and colour intensity represents the Log$_2$ Fold Change (Log2 FC), with red indicating upregulation in WT and green indicating upregulation in the mutant strains. Functional categories were based on information retrieved from Literature and EcoCyc.

## Conclusion and Outlook

The functional characterization of hypothetical protein (HP)-encoding genes remains a critical challenge, since the rapid accumulation of sequencing data has long surpassed the rate of possible experimental characterization *in vivo* and *in vitro*. While there have been notable advancements in methodologies for protein function prediction in the recent past, several challenges persist, including the heavy reliance on pre-existing data inferred from model organisms. Unfortunately, even in the best studied model organisms a significant amount of protein encoding genes could so far not been annotated.

AI-driven approaches, such as machine learning (ML) algorithms, now offer powerful tools to

analyze vast omics datasets. In this study we employed a comprehensive approach,

integrating independent component analyses to decipher transcriptional regulatory networks,

deep learning for structural homolog prediction, and various other bioinformatic tools to

characterize HPs that have no functional information available in public databases. With this

methodology putative/probable functions for at least 64% of 95 hypothetical HPs encoding

genes from *E. coli* K-12 could be extrapolated, facilitating the subsequent functional validation

of these proteins through *in vitro* and *in vivo* experiments in the future. For the remaining 36%

HPs, however, more high-quality datasets including metabolomics and proteomics are

probably needed, as these genes are often seen to be expressed under very specific

environmental conditions from which only one dataset was available. Nevertheless, for most

proteins it is possible to derive at least some characteristics from their general sequence

properties, whether there is evidence of expression, their possible cellular location, details of

homologs, or whether they are predicted or demonstrated to be expressed as part of an

operon and/or regulon. Taking these information into account in databases will aid in the

improvement of using AI tools for annotation in the future.

In this study we furthermore provided an experimental feedback loop for three HP-encoding

genes to verify the *in silico* prediction. Gene *yhdN* ought to play a role in the cells' responding

to heat shock and nutrient limitation, which could be confirmed by comparing the growth

curves and transcriptional responses of wild type and *ΔyhdN* deletion mutant strains. For the

second candidate *yeaC, in silico* predictions suggested a role in aerobic respiration.

Experimental evidence supported this hypothesis by the upregulation of oxidative stress

response and methionine biosynthesis genes, hence it is likely involved in scavenging reactive

oxygen species. In contrast, the significant upregulation of "Motility"/Chemotaxis" genes in the

deletion mutant suggest that during the absence of *yeaC*, bacteria may enhance motility

pathways as an adaptive mechanism under oxidative stress. The third candidate *ydgH*, was

predicted *in silico* to play a role in oxidoreduction, which could also be confirmed by the *in vivo*

experiments showing its role in the protection against oxidative stress. Similar to *yeaC,* in the mutant an upregulation of motility/chemotaxis genes indicate that in the absence of *ydgH*, the cell may compensate for reduced oxidoreductive function. However, it is important to acknowledge that our understanding still remains incomplete since the functional landscape of proteins is vast and complex. Future studies and more omics datasets are needed to further refine and expand upon our findings, exploring the broader implications of these proteins under different environmental conditions.

In conclusion, this study represents a significant step forward in elucidating the functions of previously uncharacterized HPs with no information in existing knowledge databases. Leveraging AI-driven annotations and integrating them with experimental laboratory work will link new functional roles of genes, discover previously unknown cellular and metabolic processes and maybe even biotechnological potentials. In the future this approach might also aid in the isolation of new and so far uncultured microbial species, i.e. microbial dark matter, by harnessing meta-omics datasets and identifying genes critical for growth and adaptation, offering clues to optimize culture conditions.

**Methods**

**Identification and enumeration of hypothetical proteins (HPs)**

For the comprehensive analysis of *Escherichia coli* K-12 (*E. coli*) substrains MG1655 and BW25113, the complete genome sequences available in the NCBI RefSeq database under the accession number GCF_000005845.2 and GCF_004355105.2 were used, respectively. Ribosomal RNA (rRNA), transfer RNA (tRNA), non-coding RNA (ncRNA) were filtered out. CDS based annotations were used to find hypothetical proteins (HPs). HPs were identified based on keywords like "Putative", "Putative uncharacterized", "Uncharacterized protein" and "DUF (domain of unknown function)-domain containing protein", "UPF" (unknown protein family) as well as proteins having a 'Protein Y…' pattern. Further information was extracted

from the extensively curated databases EcoCyc version 26.1 [39], RegulonDB version 11.2

[11], EggNOG v6.0 [12] and UniProt release 2023_04 for proteome UP000000625 [1].

In order for a HP to be classified as a protein with no characterization (target group of this

study), it had to fulfill all of the following criteria:

1. The **EcoCyc** protein summary showed the statement 'No information about this'.

Additionally, the protein had to be marked as 'Uncharacterized'. Proteins which had

the statement but a 'Partial' characterization in EcoCyc were further investigated using

UniProt.

2. In **UniProt** the protein lacked any functional information in the 'Function' section and

was assigned an annotation score of '1', and a 'protein existence' score of '4' signifying

that the protein, although predicted to be expressed, lacked information based on

experimental evidence or orthologs in closely related species. For the "Function"

section, a score of '1' is the lowest rating in a five-point-scoring system and for the

"protein existing" category a score of '4' out of '5', where '5' indicates a higher level of

uncertainty regarding the protein's expression.

3. In the **RegulonDB** database it had the label "Weak" evidence for an annotation in the

evidence section and no additional information in the 'Note' subsection.

4. In the **EggNOG** database the protein was either classified under the cluster of

orthologous groups (COG) with a functional category 'S', where 'S' designates proteins

belonging to an unknown function category or it lacked an ortholog which was

displayed as 'No orthologs found'.

95 of the 4403 *E. coli* K-12 protein encoding genes fulfilled these criteria. All scripts for the

filtering and enumeration of uncharacterized HPs can be found on Figshare

(*Notebooks>HP_curation*). In addition, the here identified HP-encoding genes were compared

with the ones identified by Ghatak et al. in 2019 [8] to check for their current status.

**Transcriptomic data curation and processing**

RNA-sequencing data from NCBI for analyzing the expression patterns of genes was collected

for *E. coli* K-12 substrains MG1655 and BW25113 using an adapted workflow for *Bacillus*

*subtilis* [32]. For substrain MG1655 1854 datasets and for substrain BW25113 662 datasets

were downloaded and processed. Raw FASTQ files were obtained using sratoolkit 3.0.1 [161],

and subsequent read trimming was performed using Trim Galore 0.6.5 [162] with the default

settings. Trimmed reads were then subjected to FastQC 0.11.9 [163]. Alignments to the

genomes were done using Bowtie 1.2.3 [164]. The read direction was determined using

RSeQC 3.0.1 [165] before generating read counts using featureCounts [166]. Finally, all

quality control metrics were compiled using MultiQC [167] and the expression compendium

was obtained as log-transformed Transcripts per Million (log-TPM) values. Quality metrics

were established to filter and ensure the selection of high-quality datasets with the following

modifications [168]: Briefly, the datasets were considered to be of high quality if they passed

the  fastqc  parameters  'per_base_sequence_quality',  'per_sequence_quality_scores',

'per_base_n_content', 'adapter_content' and had at least 500,000 reads mapped to coding

sequences from the reference genome. To preserve biological variability and prevent loss of

information, samples with no replicates were also retained in the dataset. Short non-coding

transcripts (<100 nucleotides) and extremely low-expression transcripts (Fragments Per

Kilobase per Million mapped reads < 10) were removed to reduce noise. 779 datasets for

substrain MG1655 and 135 datasets for substrain BW25113 were then used for further

analysis. All other information on the *E. coli* K-12 genes including their transcription units,

descriptions etc. were sourced from EcoCyc by utilizing the SmartTables feature [39].

Annotations for clusters of orthologous groups (COG) were obtained from EggNOG v6.0 [12]

including Gene Ontology [169]. The transcriptional regulatory network (TRN) comprising

information on genes and their governing regulators (i.e. transcription factors controlling the

expression of genes) were sourced from RegulonDB version 12.0 [11] and EcoCyc [39] as

described by Sastry et al. [32] In order to mitigate any batch effects which might be caused

due to the usage of diverse datasets, a reference condition (wild type *E. coli* K-12 substrain

MG1655 in M9 minimal medium facilitated with glucose and essential micronutrients under

non-stressful conditions) was used [30]. This was done by subtracting the mean expression

of two replicates of the chosen reference condition from the entire gene expression dataset

and calculating the log2-fold-change i.e. relative changes in expression. This normalization

method allowed for the comparison of gene expression differences relative to a common

baseline as previously demonstrated in Sastry et al. [30]. The normalized data was then used

as input for machine learning (*Data processing>Nextflow*).

**Independent Component Analysis of Transcriptomic Data**

Independent Component Analysis (ICA) developed by McCon et al. -named OptICA- was

applied to prevent both over-decomposition and under-decomposition of transcriptomic

datasets [27]. The application of ICA to the normalized *E. coli* K-12 substrain MG1655 RNAseq

data resulted in two matrices: the first matrix, denoted as the M matrix, encompassing the

robust independent components named iModulons, while the second matrix, termed the A

matrix, comprised the corresponding activities of iModulons over different growth conditions.

Additionally, a PyModulon package was used in conjunction with Jupyter notebooks to

characterize the iModulons [32]. The scripts containing the specific utilization of resources and

curation of iModulons can be found on Figshare (*OptICA* and *iModulon curation*). The

information from substrain MG1655 was then used to infer the iModulons and their member

genes for substrain BW25113, since only for the latter strain single-gene deletion mutants

were available for *in vitro* experimentation. The HP-encoding genes, the regulators governing

them, and their derived functions can be found in **Table 1.**

Possible functions for the 95 HP-encoding genes were further identified through regulators

governing their gene expression, as provided by RegulonDB and EcoCyc and/or by Gene

Ontology (GO) terms derived from co-regulating genes using the PANTHER (Protein ANalysis

THrough Evolutionary Relationships) tool [33]. Here, member genes of an iModulon (including

the gene of interest, i.e. the HP-encoding gene) were used as input to obtain GO terms to see

if any were overrepresented. Statistically significant GO categories with the default false

Discovery Rate (FDR)-adjusted p-value (<0.05) were manually identified for each HP

(*Supplementary Table 1*). These gene-associated functional categories can be found in **Table**

**1** and **Figure 2**.

**Metadata extraction**

Metadata conditions for all datasets were retrieved either manually from the sequence read

archive (SRA), *via* a semi-automated approach [167] or by utilizing data from the Gene

Expression Omnibus Database using the GEOparse tool [170] to obtain experimental

information not available *via* the NCBI's Entrez Programming Utilities (E-utilities) [42]. As a

result, from 779 quality checked datasets for the substrain MG1655, metadata for 602 datasets

could be retrieved with confidence and termed as 'Explicit metadata'. 113 datasets were

termed as 'poor metadata', meaning that information on one or more experimental conditions

(such as media composition, temperature, pH, etc.) was missing or incomplete. Lastly,

datasets with 'unclear metadata' lacked all information. For the substrain BW25113, 128

datasets were 'Explicit' while seven were 'unclear' (*Notebooks>metadata_curation*).

Additionally, metadata condition for the highest expression of the HP- encoding genes against

the reference gene *frr* were extracted from the Tjaden dataset [44] in order to account for the

loss of datasets after quality filtering (also see below). The *frr* gene expression is typically

consistent across different conditions which was also observed when tested as reference

along with other genes like *rpoC*, *rpoD* for the uncharacterized HPs [this study, 171]. Hence,

gene *frr* was chosen as the reference gene serving as a control to validate the relative

expression levels of other genes. The extensive nature of this dataset ensured a broad

representation of gene expression profiles, in addition to the datasets procured for OptICA

analysis (*Notebooks> metadata_curation*).

**Co-expression analysis based on public RNA-Seq data**

To identify genes with expression patterns similar to those encoding HPs, a Spearman

correlation analysis on the Tjaden dataset was performed [172]. This dataset comprises a

comprehensive collection of 3.376 *E. coli* strain K-12 RNA-seq datasets, with expression

levels quantified in transcripts per million (TPM) [44]. For inferring gene-gene correlations

across the transcriptomic datasets for strain *E. coli* K-12, we identified the top ten co-

expressed genes to the respective 95 HP-encoding gene and used this information to get an

inference on their functions [10] **(Table 1)** (*Notebooks>gene_co-expression*).

**Sub-Cellular Localization**

For the purpose of determining the subcellular localizations of the 95 HP-encoding genes,

BUSCA (Bologna Unified Subcellular Component Annotator) was employed [173]. This tool

integrates various machine learning and bioinformatics tools [174–177] to analyze and

recognize patterns in protein sequences that are indicative of specific subcellular localizations.

The integrative webserver was used to obtain labels such as 'Nucleus', 'Periplasm/cytoplasm',

'Inner/outer membrane' or 'extracellular space'. All 95 HP-encoding genes with their predicted

sub-cellular localizations can be found in **Table1**.

**Protein-Protein Interactions (PPIs) Analysis**

The STRING database (version 12.0) (https://string-db.org) was employed for the

identification of PPIs of HP-encoding genes against the proteomic landscape of *E. coli*. The

tool incorporates data from diverse origins such as available publications, empirical evidence,

predictions derived from co-expression patterns, genomic context analysis, and curated

pathway databases. A 'confidence score' for each inferred association was derived,

quantitatively expressing the robustness of the predicted interaction [178]. In order to keep

only the high-scoring associations, a cut-off of 0.7 was utilized [179]. All 95 HP-encoding

genes and their interacting partner proteins are listed in **Table 1**.

**Analysis of Gene Context**

Shared functional relationships were analyzed using the genomic context of

*E. coli* strain K-12 (*Supplementary Figure 1*), particularly the genes located upstream and

downstream of the 95 genes of interest. A gene list comprising of gene names was used to

create a SmartTable in the EcoCyc database to examine the local genomic neighbourhood

and visualize the genes upstream and downstream, including their orientations [36]. (**Table**

**1)**.

**Detection of Remote Protein Homology**

In order to identify distantly homologous sequences, HP sequences were used as input for

the HHblits tool (version 3.3.0) available in the MPI Bioinformatics Toolkit using a Hidden

Markov Model (HMM) iterative sequence search [180]. The HHblits server

(https://toolkit.tuebingen.mpg.de/tools/hhblits) takes a single sequence or a multiple sequence

alignment as input and iteratively searches through the selected HMM databases [181,182].

HHblits performs sequence search and remote homology detection with higher sensitivity as

compared to gold-standard databases like InterPro [180]. Homologous sequences were then

searched against the default UniRef30 database [37]. Sequence based remote homologies

identified through the HHblits tool are catalogued in **Table 1**.

**Structure-Based Homology**

For identifying structural homologs, we used the AlphaFold Protein Structure Database

(AFDB) (https://cluster.foldseek.com/), a specialized computational database that includes

structural models for approximately 215 million protein sequences. This platform employs

deep learning algorithms to scan and match protein structures, enabling the discovery of

structural similarities, even among proteins with low sequence identity [183]. We analyzed

clusters to perform comparisons of HPs with all protein structures in the AFDB. UniProt

accession IDs were used as inputs to identify the cluster to which the HP belongs. The

homology-based function information for each of the 95 HP is listed in **Table 1**. A script to

extract the information can be found on Figshare (*Notebooks> HP_info_curation*).

**Bacterial strains and growth conditions**

*E. coli* K-12 substrain BW25113 was sourced from Horizon Discovery Ltd., United States.

Three single-gene knockout mutants from *E. coli* strain K-12 substrain BW25113 *ΔyhdN,*

*ΔyeaC*, and *ΔydgH* were obtained from the KEIO collection (NBRP, Japan). Glycerol stocks

were stored in an ultra-low temperature freezer (Thermo Fisher, United States) at −80°C until

further processing. Cells were cultured overnight on agar plates from glycerol stocks and

single colonies were picked from the agar plate and then used to Luria Bertani (LB) liquid

medium at 37°C [184].

**Media and reagents**

*E. coli* strains were cultured in LB broth (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl;

pH 7.0) at 37°C while shaking at 200 rpm [184]. M9 liquid medium for the pre-culture was

prepared using 42 mM $Na_2HPO_4$, 22 mM $KH_2PO_4$, 8.5 mM NaCl, 18.7 mM $NH_4Cl$, 2 mM

$MgSO_4$, 0.1 mM $CaCl_2$ and 11 mM glucose, supplemented with 0.05% casamino acids to

ensure non-limiting conditions. For nitrogen limited M9 medium, a $NH_4Cl$ concentration of 0.16

mM was used without the addition of casamino acids. For solid medium, 1.5% agar was

added. 30% $H_2O_2$ was sterile filtered through a 0.22 µm pore size filter to ensure sterility prior

to be used in the oxidative stress experiments. For all experiments with $H_2O_2$ the LB was

freshly prepared and kept in the dark until use. All chemicals were sourced from Sigma

Chemical Co. (St. Louis, MO, USA).

**Measurement of Bacterial Growth Curves**

Bacterial cultures were inoculated at a 1:100 dilution unless stated otherwise. Each

experiment included triplicates. All optical density values ($OD_{600}$) were measured *via* a cell

density meter (Ultrospec 10, biochrom). $OD_{600}$ was measured every 2h for heat shock in LB

and M9 or M9 modified medium and every 1h for oxidative stress experiments. Growth curves

were generated for each set of experiments including the wild type and mutant strain in

stressed and non-stressed conditions (controls).

Heat stress

A single colony (wild type and Δ*yhdN*) was picked from either LB or M9 agar plates and

inoculated into 5 mL LB broth for overnight cultivation at 37°C and shaken with 200 rpm. The

overnight culture was then used to inoculate fresh LB medium at a 1:100 dilution, which was

then cultured for ~2-3h at 37°C and 200 rpm until an $OD_{600}$ of 0.4 was reached. This was

repeated with the M9 liquid medium (preculture), at 37°C to ensure non limiting conditions.

For the nitrogen limitation assay, medium was prepared using 0.16mM $NH_4Cl$. To study the

effects of heat shock on bacterial growth, the culture was grown to mid-log phase ($OD_{600}$ 0.4),

diluted to $OD_{600}$ 0.04 and then heat shocked for 7 minutes in a shaker-incubator at 50°C. After

treatment, the bacterial suspension was incubated for 12 h at 37°C. Samples were measured

every 2h for 12h.

Oxidative stress

For studying the effects of oxidative stress, a single colony (wild type, Δ*yeaC* and Δ*ydgH*) was

picked from the LB agar plate and inoculated into 5 mL LB for overnight cultivation at 37°C

and shaken with 200 rpm. The overnight culture was then used to inoculate fresh LB (dilution

of 1:100) and cultured at 37°C and 200 rpm until an $OD_{600}$ of 0.2 was reached. An OD of 0.2

was used since at this stage the cells were actively dividing, making them susceptible to the

effects of oxidative stress, allowing for better characterization and analysis of the stress

response[137]. $H_2O_2$ was added at a final concentration of 2.5 mM. After treatment, the

bacterial suspension was incubated for 5h at 37°C and the $OD_{600}$ was measured hourly.

**RNA Sequencing and data processing**

2 ml of cell suspension from the heat shock samples was collected after 2h and 6h of growth.

For the oxidative stress experiments, samples were collected after 1h and 5h of growth. The

medium was discarded, cell pellets were washed with Phosphate Buffer Saline and mixed with

DNA/RNA shield™ (Zymo Research Corp., Freiburg, Germany). All samples were processed

according to the manufacturer's instructions. Samples were preserved at -20°C until RNA

extraction. RNA was extracted using the Direct-zol RNA Miniprep Plus Kit (Zymo Research

Corp., Freiburg, Germany) according to the manufacturer's instructions. Total RNA was eluted

from columns with 100 μL nuclease-free water and quantified using the Qubit RNA Broad

Range Assay Kit (Thermo Fisher Scientific Inc., Waltham, MA, USA). The quality was

assessed by running an RNA-nano chip on an Agilent bioanalyzer (Agilent Technologies,

Waldbronn, Germany). Samples with an RNA integrity score ≥8 were used for further

processing [185]. The rRNA was removed from total RNA preparations using the NEBNext

rRNA Depletion (Bacteria) kit (New England Biolabs, Frankfurt, Germany). Paired-end library

preparation was done using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina

(New England Biolabs, Frankfurt, Germany) following the manufacturer's instructions with an

average insert size of 200 bp. Libraries were run on a NextSeq 550™ (Illumina, San Diego,

CA, United States).

The RNAflow pipeline [186] was implemented using a conda environment. Sequence reads

were checked for quality using FastQC v0.11.9 [187]. Raw reads were processed using fastp

v0.20.0 [188], which employs a sliding window approach for quality trimming and adapter

clipping, with the settings: a -5 to -3 cut for front and tail, a default window of 4 and mean

quality threshold of 20. Trimmed reads shorter than 15 bps were discarded. The remaining

reads were mapped onto the *E. coli* K-12 reference genome (NZ_CP009273.1) using HISAT

v2.1.0 [189]. The generated output was converted into a BAM format using samtools-1.20

[190], followed by the processes of sorting and indexing. A gene transfer format (GTF)

annotation file for *E. coli* substrain BW25113 was used to filter for gene and pseudogene

features. FeatureCounts v2.0.1 [191] was used to quantify the mapped reads based on the

gene annotations. Processed datasets were summarized in a MultiQC v1.9 [167] generated

file and were normalized by the DESeq2 module (v1.28.0) in R (version 4.3.2) using a

variance stabilizing transformation [192].

**Comparative gene expression analysis of wild type and isogenic mutant strains**

Sequencing reads were obtained from the transcriptomic datasets of wild-type (WT), *ΔyhdN*,

*ΔyeaC,* and *ΔydgH* mutants, respectively. Non-stressed control and heat shocked samples

from WT and *ΔyhdN* at the 6h timepoint and non-stressed control and oxidative stress samples

from WT, *ΔyeaC,* and *ΔydgH* at the 5h timepoint were used to identify up- or downregulated

genes. DESeq2 [192] was utilized to examine expression differences of genes in the WT and

mutant strains under both control and stress conditions. To focus on the specific effect of

stress on gene expression rather than the difference of WT *vs.* deletion mutants, read counts

under control conditions were subtracted from the read counts under stress condition [193].

Genes with an adjusted P-value (Padj) of 0.1 or less were classified as differentially expressed

to avoid false positives. A cut-off $\log_2$ fold change ($\log_2$FC) of 1.5 was chosen for further

analysis [194]. A $\log_2$FC of 1.5 means that a gene was 2.83 times more expressed in one

strain *vs.* the other, a $\log_2$FC of 2 means that a gene was 4 times more expressed and so on.

Significant positive $\log_2$FC indicated genes that were upregulated in WT relative to the mutant,

while negative values showed an upregulation in the mutant *vs*. WT. To determine the

functional categories of these genes, the EcoCyc database was used [36]. Information of the

differentially expressed genes can be found on Figshare (*DEGs>Table1*).

**Data availability**

All code and scripts used in this study are available on Figshare. These resources can be

accessed at https://figshare.com/s/0ede175c510cf201e7c2.

Raw sequencing data that supports the findings of this study have been deposited in the

Sequence Read Archive with the BioProject ID: PRJNA1209969.

**Competing interests**

The authors declare no competing interests.

**Funding**

This work was supported by the Karlsruhe Institute of Technology and the Helmholtz Society

[POF4; 5207.0004.0012].

**Authors' contributions**

S.C, H.A. and Z.A did the data analyses, S.C. did the *in vitro* experiments, S.C and A.K.K.

wrote the manuscript, A.K.K. provided the funding.

**Acknowledgements**

The authors acknowledge the support by the state of Baden-Württemberg through bwHPC.

We thank Dr. Gunnar Sturm for technical support and insightful discussions and David Thiele

for library preparation assistance.

**References**

1. Consortium TU. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res.
2023;51:D523–31; doi:10.1093/nar/gkac1052
865
866 2. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al. InterPro
867 in 2022. Nucleic Acids Res. 2023;51:D418–27; doi:10.1093/nar/gkac993
868
869 3. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.
870 Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49:D412–9;
871 doi:10.1093/nar/gkaa913
872
873 4. Sigrist CJA, de Castro E, Cerutti L, Cuche BA, Hulo N, Bridge A, et al. New and continuing
874 developments at PROSITE. Nucleic Acids Res. 2013;41:D344–7; doi:10.1093/nar/gks1067
875
876 5. Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in
877 2020. Nucleic Acids Res. 2021;49:D458–60; doi:10.1093/nar/gkaa937
878
879 6. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE:
880 the conserved domain database in 2020. Nucleic Acids Res. 2020;48:D265–8;
881 doi:10.1093/nar/gkz991
882

7. Durairaj J, Waterhouse AM, Mets T, Brodiazhenko T, Abdullah M, Studer G, et al.
Uncovering new families and folds in the natural protein universe. Nature. 2023;622:646–53;
doi:10.1038/s41586-023-06622-3

8. Ghatak S, King ZA, Sastry A, Palsson BO. The y-ome defines the 35% of *Escherichia coli*
genes that lack experimental evidence of function. Nucleic Acids Res. 2019;47:2446–54;
doi:10.1093/nar/gkz030

9. Ardern Z, Chakraborty S, Lenk F, Kaster A-K. Elucidating the functional roles of prokaryotic
proteins using big data and artificial intelligence. FEMS Microbiol Rev. 2023;47;
doi:10.1093/femsre/fuad003

10. Keseler IM, Gama-Castro S, Mackie A, Billington R, Bonavides-Martínez C, Caspi R, et
al. The EcoCyc Database in 2021. Front Microbiol. 2021;12; doi:10.3389/fmicb.2021.711077

11. Salgado H, Gama-Castro S, Lara P, Mejia-Almonte C, Alarcón-Carranza G, López-Almazo
AG, et al. RegulonDB v12.0: a comprehensive resource of transcriptional regulation in *E. coli*
K-12. Nucleic Acids Res. 2024;52:D255–64; doi:10.1093/nar/gkad1072

12. Hernández-Plaza A, Szklarczyk D, Botas J, Cantalapiedra CP, Giner-Lamia J, Mende DR,
et al. eggNOG 6.0: enabling comparative genomics across 12 535 organisms. Nucleic Acids
Res. 2023;51:D389–94; doi:10.1093/nar/gkac1022

13. Mills CL, Beuning PJ, Ondrechen MJ. Biochemical functional predictions for protein
structures of unknown or uncertain function. Comput Struct Biotechnol J. 2015;13:182–91;
doi:10.1016/j.csbj.2015.02.003

14. Zhang C, Freddolino PL, Zhang Y. COFACTOR: improved protein function prediction by
combining structure, sequence and protein–protein interaction information. Nucleic Acids Res.
2017;45:W291–9; doi:10.1093/nar/gkx366

15. Makrodimitris S, van Ham RCHJ, Reinders MJT. Improving protein function prediction
using protein sequence and GO-term similarities. Bioinformatics. 2019;35:1116–24. Available
from: doi:10.1093/bioinformatics/bty751

16. Saha S, Chatterjee P, Basu S, Nasipuri M, Plewczynski D. FunPred 3.0: improved protein
function prediction using protein interaction network. PeerJ. 2019;7. doi: 10.7717/peerj.6830

17. Varadi M, Tsenkov M, Velankar S. Challenges in bridging the gap between protein
structure prediction and functional interpretation. Proteins: Structure, Function, and
Bioinformatics. 2023; doi:10.1002/prot.26614

18. Vincent AT. Bacterial hypothetical proteins may be of functional interest. Frontiers in
Bacteriology. 2024;3; doi:10.3389/fbrio.2024.1334712

19. Jeffery CJ. Current successes and remaining challenges in protein function prediction.
Frontiers in Bioinformatics. 2023;3; doi:10.3389/fbinf.2023.1222182

20. Pavlopoulos GA, Baltoumas FA, Liu S, Selvitopi O, Camargo AP, Nayfach S, et al.
Unraveling the functional dark matter through global metagenomics. Nature. 2023;622:594–
602; doi:10.1038/s41586-023-06583-7

21. Escudeiro P, Henry CS, Dias RPM. Functional characterization of prokaryotic dark matter:
the road so far and what lies ahead. Curr Res Microb Sci. 2022;3:100159; doi:
10.1016/j.crmicr.2022.100159

938
939 22. da Costa WLO, Araújo CL de A, Dias LM, Pereira LC de S, Alves JTC, Araújo FA, et al.
940 Functional annotation of hypothetical proteins from the *Exiguobacterium antarcticum* strain B7
941 reveals proteins involved in adaptation to extreme environments, including high arsenic
942 resistance. PLoS One. 2018;13:e0198965; doi:10.1371/journal.pone.0198965

943
944 23. Grünberger F, Knüppel R, Jüttner M, Fenk M, Borst A, Reichelt R, et al. Exploring
945 prokaryotic transcription, operon structures, rRNA maturation and modifications using
946 Nanopore-based native RNA sequencing. bioRxiv. 2020;2019.12.18.880849; doi:
947 10.1101/2019.12.18.880849

948
949 24. Dall'Alba G, Casa PL, Abreu FP de, Notari DL, de Avila e Silva S. A Survey of Biological
950 Data in a Big Data Perspective. Big Data. 2022;10:279–97; doi:10.1089/big.2020.0383

951
952 25. Chen J, Gu Z, Lai L, Pei J. In silico protein function prediction: the rise of machine learning-
953 based approaches. 2023;3:487–510; doi:10.1515/mr-2023-0038

954
955 26. Alteri CJ, Lindner JR, Reiss DJ, Smith SN, Mobley HLT. The broadly conserved regulator
956 PhoP links pathogen virulence and membrane potential in *Escherichia coli*. Mol Microbiol.
957 2011;82:145–63; doi:10.1111/j.1365-2958.2011.07804.x

958
27. Mcconn JL, Lamoureux CR, Poudel S, Palsson BO, Sastry A V. Optimal dimensionality
selection for independent component analysis of transcriptomic data; doi:10.1186/s12859-
021-04497-7
28. Yu J, Li M, Wang J, Hamushan M, Jiang F, Wang B, et al. Identification of *Staphylococcus*
*aureus* virulence-modulating RNA from transcriptomics data with machine learning. Virulence.
2023;14:2228657; doi:10.1080/21505594.2023.2228657

965
966 29. Comon P. Independent component analysis, A new concept? Signal Processing.
967 1994;36:287–314; doi:10.1016/0165-1684(94)90029-9

968
969 30. Sastry A V, Gao Y, Szubin R, Hefner Y, Xu S, Kim D, et al. The *Escherichia coli*
970 transcriptome mostly consists of independently regulated modules. Nat Commun.
971 2019;10:5536; doi:10.1038/s41467-019-13483-w

972
973 31. Sastry A V, Poudel S, Rychel K, Yoo R, Lamoureux CR, Chauhan S, et al. Title Mining all
974 publicly available expression data to compute dynamic microbial transcriptional regulatory
975 networks Authors; doi:10.1101/2021.07.01.450581

976
977 32. Rychel K, Sastry A V., Palsson BO. Machine learning uncovers independently regulated
978 modules in the *Bacillus subtilis* transcriptome. Nat Commun. 2020;11. doi: 10.1038/s41467-
979 020-20153-9

980
981 33. Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou L-P, Mi H. PANTHER:
982 Making genome-scale phylogenetics accessible to all. Protein Science. 2022;31:8–22;
983 doi:10.1002/pro.4218

984
985 34. Savojardo C, Martelli PL, Fariselli P, Profiti G, Casadio R. BUSCA: an integrative web
986 server to predict subcellular localization of proteins. Nucleic Acids Res. 2018;46:W459–66;
987 doi:10.1093/nar/gky320

988
989 35. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING
990 database in 2023: protein–protein association networks and functional enrichment analyses
991 for any sequenced genome of interest. Nucleic Acids Res. 2023;51:D638–46; doi:
992 10.1093/nar/gkac1000

993
994 36. D KP, Suzanne P, Ron C, Anamika K, Markus K, E MP, et al. The EcoCyc Database
995 (2023). EcoSal Plus. 2023;11:eesp-0002-2023; doi:10.1128/ecosalplus.esp-0002-2023
996
997 37. Gabler F, Nam S-Z, Till S, Mirdita M, Steinegger M, Söding J, et al. Protein Sequence
998 Analysis Using the MPI Bioinformatics Toolkit. Curr Protoc Bioinformatics. 2020;72:e108;
999 doi:10.1002/cpbi.108
1000
1001 38. Barrio-Hernandez I, Yeo J, Jänes J, Mirdita M, Gilchrist CLM, Wein T, et al. Clustering
1002 predicted structures at the scale of the known protein universe. Nature. 2023;622:637–45;
1003 doi:10.1038/s41586-023-06510-w
1004
1005 39. Karp PD, Billington R, Caspi R, Fulcher CA, Latendresse M, Kothari A, et al. The BioCyc
1006 collection of microbial genomes and metabolic pathways. Brief Bioinform. 2019;20:1085–93;
1007 doi:10.1093/bib/bbx085
1008
1009 40. Choudhary S. pysradb: A Python package to query next-generation sequencing metadata
1010 and data from NCBI Sequence Read Archive. F1000Res. 2019;8:532; doi:
1011 10.12688/f1000research.18676.1
1012
1013 41. Gumienny R. GEOparse. Rev.0a257463. 2015. Available from:
1014 https://geoparse.readthedocs.io/en/latest/
1015
1016 42. Liu M, Durfee T, Cabrera JE, Zhao K, Jin DJ, Blattner FR. Global transcriptional programs
1017 reveal a carbon source foraging strategy by *Escherichia coli*. J Biol Chem. 2005;280:15921–
1018 7; doi: 10.1074/jbc.m414050200
1019
1020 43. Tjaden B. *Escherichia coli* transcriptome assembly from a compendium of RNA-seq data
1021 sets. RNA Biol. 2023;20:77–84; doi:10.1080/15476286.2023.2189331
1022
1023 44. Rawangkan A, Yosboonruang A, Kiddee A, Siriphap A, Pook-In G, Praphasawat R, et al.
1024 Restoring Ampicillin Sensitivity in Multidrug-Resistant *Escherichia coli* Following Treatment in
1025 Combination with Coffee Pulp Extracts. J Microbiol Biotechnol. 2023;33:1–10;
1026 doi:10.4014/jmb.2304.04051
1027
1028 45. Manat G, El Ghachi M, Auger R, Baouche K, Olatunji S, Kerff F, et al. Membrane Topology
1029 and Biochemical Characterization of the *Escherichia coli* BacA Undecaprenyl-Pyrophosphate
1030 Phosphatase. PLoS One. 2015;10:e0142870; doi:10.1371/journal.pone.0142870
1031
1032 46. Hajra D, Nair AV, Chakravortty D. An elegant nano-injection machinery for sabotaging the
1033 host: Role of Type III secretion system in virulence of different human and animal pathogenic
1034 bacteria. Phys Life Rev. 2021;38:25–54; doi:10.1016/j.plrev.2021.05.007
1035
1036 47. Prakash A, Dutta D. Bicyclomycin generates ROS and blocks cell division in *Escherichia
1037 coli*. PLoS One. 2024;19:e0293858; doi:10.1371/journal.pone.0293858
1038
1039 48. Ogawa A, Kojima F, Miyake Y, Yoshimura M, Ishijima N, Iyoda S, et al. Regulation of
1040 constant cell elongation and Sfm pili synthesis in *Escherichia coli* via two active forms of FimZ
1041 orphan response regulator. Genes to Cells. 2022;27:657–74; doi:10.1111/gtc.12982
1042
1043 49. Lamark T, Røkenes TP, McDougall J, Strøm AR. The complex bet promoters of
1044 *Escherichia coli*: regulation by oxygen (ArcA), choline (BetI), and osmotic stress. J Bacteriol.
1045 1996;178(6):1655–62; doi:10.1128/jb.178.6.1655-1662.1996.
1046

50. Kambampati R, Lauhon CT. Evidence for the Transfer of Sulfane Sulfur from IscS to ThiI during the in Vitro Biosynthesis of 4-Thiouridine in *Escherichia coli* tRNA*. Journal of Biological Chemistry. 2000;275:10727–30; doi:10.1074/jbc.275.15.10727

51. Laura F-G, Xinyu G, Joy K, Sooyeon S, E BM, Rodolfo G-C, et al. Single-cell analysis reveals that cryptic prophage protease LfgB protects *Escherichia coli* during oxidative stress by cleaving antitoxin MqsA. Microbiol Spectr. 2024;12:e03471-23; doi:10.1128/spectrum.03471-23

52. Zorraquino-Salvo V, Quinones-Soto S, Kim M, Rai N, Tagkopoulos I. Deciphering the genetic and transcriptional basis of cross-stress responses in *Escherichia coli* under complex evolutionary scenarios. bioRxiv. 2014;010595; doi:

53. May T, Okabe S. Enterobactin is required for biofilm development in reduced-genome *Escherichia coli*. Environ Microbiol. 2011;13:3149–62; doi:10.1111/j.1462-2920.2011.02607.x

54. Saier Jr MH, Reddy VS, Tamang DG, Västermark Å. The Transporter Classification Database. Nucleic Acids Res. 2014;42:D251–8; doi:10.1093/nar/gkt1097

55. Muto A, Goto S, Kurita D, Ushida C, Soma A, Himeno H. A leaderless mRNA including tRNA-like sequence encodes a small peptide that regulates the expression of GcvB small RNA in *Escherichia coli*. The Journal of Biochemistry. 2022;171:459–65; doi:10.1093/jb/mvac007

56. Njenga R, Boele J, Öztürk Y, Koch H-G. Coping with stress: How bacteria fine-tune protein synthesis and protein transport. Journal of Biological Chemistry. 2023;299:105163; doi: 10.1016/j.jbc.2023.105163

57. Rodríguez-Ochoa JI, Fragoso-Jiménez JC, Flores N, et al. Global transcriptomic response of *Escherichia coli* to p-coumaric acid. Microb Cell Fact. 2022;21:148; doi:10.1186/s12934-022-01874-6.

58. Bak G, Lee J, Suk S, Kim D, Lee J, Kim K, et al. Identification of novel sRNAs involved in biofilm formation, motility, and fimbriae formation in *Escherichia coli*. Sci Rep. 2015;5:15287; doi:10.1038/srep15287

59. Duarte-Velázquez I, Mora J, Ramírez-Prado J, Aguillón-Bárcenas A, Tornero-Gutiérrez F, Cordero-Loreto E, et al. *Escherichia coli* transcription factors of unknown function: sequence features and possible evolutionary relationships. PeerJ. 2022;10:e13772; doi:10.7717/peerj.13772

60. Saier Jr MH, Reddy VS, Tsu B V, Ahmed MS, Li C, Moreno-Hagelsieb G. The Transporter Classification Database (TCDB): recent advances. Nucleic Acids Res. 2016;44:D372–9; doi:10.1093/nar/gkv1103

61. Su C-C, Rutherford DJ, Yu EW. Characterization of the multidrug efflux regulator AcrR from *Escherichia coli*. Biochem Biophys Res Commun. 2007;361:85–90; doi:10.1016/j.bbrc.2007.06.175

62. Börner J, Friedrich T, Bartkuhn M, Klug G. Ribonuclease E strongly impacts bacterial adaptation to different growth conditions. RNA Biol. 2023;20:120–35; doi:10.1080/15476286.2023.2195733

63. A JN, Melissa F, E FS. Disruption of trehalose periplasmic recycling dysregulates cAMP-CRP signaling in *Escherichia coli* during stationary phase. J Bacteriol. 2023;205:e00292-23; doi:10.1128/jb.00292-23

64. Wang M, Fang K, Hong SMC, Kim I, Jang I-S, Hong SH. Medium chain unsaturated fatty acid ethyl esters inhibit persister formation of *Escherichia coli* via antitoxin HipB. Appl Microbiol Biotechnol. 2018;102:8511–24; doi:10.1007/s00253-018-9271-3

65. Muenter MM, Aiken A, Akanji JO, Baig S, Bellou S, Carlson A, et al. The response of *Escherichia coli* to the alkylating agents chloroacetaldehyde and styrene oxide. Mutation Research/Genetic Toxicology and Environmental Mutagenesis. 2019;840:1–10; doi:10.1016/j.mrgentox.2019.02.001

66. Hideji Y, Yasushi M, Shou F, Akiko S, Masami U, Akira W. YqjD Is an Inner Membrane Protein Associated with Stationary-Phase Ribosomes in *Escherichia coli*. J Bacteriol . 2012;194:4178–83; doi:10.1128/jb.00396-12

67. Hemm MR, Paul BJ, Schneider TD, Storz G, Rudd KE. Small membrane proteins found by comparative genomics and ribosome binding site models. Mol Microbiol. 2008;70:1487–501; doi:10.1111/j.1365-2958.2008.06495.x

68. Skjold AC, Ezekiel DH. Analysis of lambda insertions in the fucose utilization region of *Escherichia coli* K-12: use of lambda *fuc* and lambda *argA* transducing bacteriophages to partially order the fucose utilization genes. J Bacteriol. 1982;152:120–5; doi:10.1016/j.mrgentox.2019.02.001

69. Manuel C, Teresa ZM, Blas B, Gonzalo D-R, F JJ, Andrés VJ, et al. Anaerobic Catabolism of Aromatic Compounds: a Genetic and Genomic View. Microbiology and Molecular Biology Reviews. 2009;73:71–133; 10.1128/mmbr.00021-08

70. Arnim W, A KS, Kirsten J. Time-Dependent Proteome Alterations under Osmotic Stress during Aerobic and Anaerobic Growth in *Escherichia coli*. J Bacteriol. 2006;188:7165–75; doi:10.1128/jb.00508-06

71. Teufel R, Mascaraque V, Ismail W, Voss M, Perera J, Eisenreich W, et al. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. Proc Natl Acad Sci U S A. 2010;107(32):14390–5; doi: 10.1073/pnas.1005399107.

72. Tenorio E, Saeki T, Fujita K, Kitakawa M, Baba T, Mori H, et al. Systematic characterization of *Escherichia coli* genes/ORFs affecting biofilm formation. FEMS Microbiol Lett. 2003;225(1):107–14; doi:10.1016/S0378-1097(03)00507-X.

73. Rodríguez-Montelongo L, Volentini SI, Farías RN, Massa EM, Rapisarda VA. The Cu(II)-reductase NADH dehydrogenase-2 of *Escherichia coli* improves the bacterial growth in extreme copper concentrations and increases the resistance to the damage caused by copper and hydroperoxide. Arch Biochem Biophys. 2006;451:1–7; doi:10.1016/j.abb.2006.04.019

74. Harwani D. Regulation of Gene Expression: Cryptic β-glucoside (bgl) Operon of *Escherichia coli* as a Paradigm. Brazilian Journal of Microbiology. 2014; doi:10.1590/S1517-83822014000400003

75. Anne P, K BI, Aurora L-N, F GJ, Valérie de C-L, D HA. FolX and FolM Are Essential for Tetrahydromonapterin Synthesis in *Escherichia coli* and Pseudomonas aeruginosa. J Bacteriol. 2010;192:475–82; doi:10.1128/jb.01198-09

76. Kotaka Y, Hashimoto M, Lee K, Kato J. Mutations identified in engineered *Escherichia coli* with a reduced genome. Front Microbiol. 2023;14; doi:10.3389/fmicb.2023.1189877

77. Doukyu N, Taguchi K. Involvement of catalase and superoxide dismutase in hydrophobic organic solvent tolerance of *Escherichia coli*. AMB Express. 2021;11:97; doi:10.1186/s13568-021-01258-w

78. Pinske C. The Ferredoxin-Like Proteins HydN and YsaA Enhance Redox Dye-Linked Activity of the Formate Dehydrogenase H Component of the Formate Hydrogenlyase Complex. Front Microbiol. 2018;9; doi:10.3389/fmicb.2018.01238

79. Panta PR, Doerrler WT. A link between pH homeostasis and colistin resistance in bacteria. Sci Rep. 2021;11:13230; doi:10.1038/s41598-021-92718-7

80. Vasilyev N, Liu MMJ, Epshtein V, Shamovsky I, Nudler E. General Transcription Factor from *E. coli* with a Distinct Mechanism of Action. bioRxiv. 2023;2023.09.17.558134; doi: 10.1101/2023.09.17.558134

81. Nasreen M, Purushothaman Nair R, Mcewan A, Kappler U. The Peptide Methionine Sulfoxide Reductase (MsrAB) of Haemophilus influenzae Repairs Oxidatively Damaged Outer Membrane and Periplasmic Proteins Involved in Nutrient Acquisition and Virulence. Antioxidants. 2022;11:1557; doi:10.3390/antiox11081557

82. Yuan J, Jin F, Glatter T, Sourjik V. Osmosensing by the bacterial PhoQ/PhoP two-component system. Proceedings of the National Academy of Sciences. 2017;114:E10792–8; doi:10.1073/pnas.1717272114

83. Steinsiek S, Stagge S, Bettenbrock K. Analysis of *Escherichia coli* mutants with a linear respiratory chain. PLoS One. 2014;9(1):e87307; doi:10.1371/journal.pone.0087307.

84. Akinori K, Hiroyuki T, Ryutaro U. Molecular Characterization of the PhoP-PhoQ Two-Component System in *Escherichia coli* K-12: Identification of Extracellular Mg2+-Responsive Promoters. J Bacteriol. 1999;181:5516–20; doi:10.1128/jb.181.17.5516-5520.1999

85. Podlesek Z, Žgur Bertok D. The DNA Damage Inducible SOS Response Is a Key Player in the Generation of Bacterial Persister Cells and Population Wide Tolerance. Front Microbiol .2020;11; doi:10.3389/fmicb.2020.01785

86. Bitoun JP, Wu G, Ding H. *Escherichia coli* FtnA acts as an iron buffer for re-assembly of iron–sulfur clusters in response to hydrogen peroxide stress. BioMetals. 2008;21:693–703; doi:10.1007/s10534-008-9154-7

87. C BD, J TR. Small Multidrug Resistance Protein EmrE Reduces Host pH and Osmotic Tolerance to Metabolic Quaternary Cation Osmoprotectants. J Bacteriol. 2012;194:5941–8; doi:10.1128/jb.00666-12

88. M FE, E EK, N GL, R GE, M MJ, C MA, et al. YfdW and YfdU Are Required for Oxalate-Induced Acid Tolerance in *Escherichia coli* K-12. J Bacteriol. 2013;195:1446–55; doi:10.1128/jb.01936-12

89. Ralph ET, Guest JR, Green J. Altering the anaerobic transcription factor FNR confers a hemolytic phenotype on *Escherichia coli* K12. Proceedings of the National Academy of Sciences. 1998;95:10449–52; doi:10.1073/pnas.95.18.10449

90. Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM, et al. Cryptic prophages help bacteria cope with adverse environments. Nat Commun. 2010;1:147; doi:10.1038/ncomms1146

91. Barabote RD, Rendulic S, Schuster SC, Saier MH. Comprehensive analysis of transport
proteins encoded within the genome of *Bdellovibrio bacteriovorus.* Genomics .
2007;90:424–46; doi:10.1016/j.ygeno.2007.06.002

92. Wang J, Ma W, Wang X. Insights into the structure of *Escherichia coli* outer membrane as
the target for engineering microbial cell factories. Microb Cell Fact. 2021;20:73;
doi:10.1186/s12934-021-01565-8
93. Gurmu D, Lu J, Johnson KA, Nordlund P, Holmgren A, Erlandsen H. The crystal structure
of the protein YhaK from *Escherichia coli* reveals a new subclass of redox sensitive
enterobacterial bicupins. Proteins: Structure, Function, and Bioinformatics. 2009;74:18–31;
doi:10.1002/prot.22128

94. Outten CE, Outten FW, O'Halloran T V. DNA Distortion Mechanism for Transcriptional
Activation by ZntR, a Zn(II)-responsive MerR Homologue in *Escherichia coli*. Journal of
Biological Chemistry. 1999;274:37517–24; doi:10.1074/jbc.274.53.37517

95. Akbar S, Bhakta S, Sengupta J. Structural insights into the interplay of protein biogenesis
factors with the 70S ribosome. Structure. 2021;29:755-767.e4; doi:10.1016/j.str.2021.03.005

96. K VDT, J TL, A CK, L SP, Sima SF. Characterization of the *Escherichia coli* AaeAB Efflux
Pump: a Metabolic Relief Valve? J Bacteriol. 2004;186:7196–204;
doi:10.1128/jb.186.21.7196-7204.2004

97. Bak G, Lee J, Suk S, Kim D, Young Lee J, Kim K, et al. Identification of novel sRNAs
involved in biofilm formation, motility and fimbriae formation in *Escherichia coli*. Sci Rep.
2015;5:15287; doi:10.1038/srep15287

98. P DM, Chi-Fang W, Liang W, J VJ, E BW. DNA Microarray-Based Identification of Genes
Controlled by Autoinducer 2-Stimulated Quorum Sensing in *Escherichia coli*. J Bacteriol.
2001;183:5239–47; doi:10.1128/jb.183.18.5239-5247.2001

99. Filenko N, Spiro S, Browning DF, Squire D, Overton TW, Cole J, Constantinidou C. The
NsrR regulon of *Escherichia coli* K-12 includes genes encoding the hybrid cluster protein and
the periplasmic, respiratory nitrite reductase. J Bacteriol. 2007;189:1203–9;
doi:10.1128/jb.00080-07.

100. Khil PP, Camerini-Otero RD. Over 1000 genes are involved in the DNA damage response
of *Escherichia coli*. Mol Microbiol. 2002;44:89–105; doi:10.1046/j.1365-2958.2002.02878.x

101. Erol I, Jeong K-C, Baumler DJ, Vykhodets B, Ho Choi S, Kaspar CW. H-NS controls
metabolism and stress tolerance in *Escherichia coli* O157:H7 that influence mouse passage.
BMC Microbiol. 2006;6:72; doi:10.1186/1471-2180-6-72

102. Zheng L. Catalytic mechanism and function evolvement studies of phosphatases within
the haloacid dehalogenase superfamily (HADSF) [dissertation]. Albuquerque (NM): University
of New Mexico; 2013. Available from: https://digitalrepository.unm.edu/chem_etds/31.

103. Born TL, Blanchard JS. Structure/function studies on enzymes in the diaminopimelate
pathway of bacterial cell wall biosynthesis. Curr Opin Chem Biol. 1999;3:607–13;
doi:10.1016/S1367-5931(99)00016-2

104. Hor L, Dobson RCJ, Downton MT, Wagner J, Hutton CA, Perugini MA. Dimerization of
Bacterial Diaminopimelate Epimerase Is Essential for Catalysis. J Biol Chem .
2013;288:9238–48

105. Iyer LM, Leipe DD, Koonin E V, Aravind L. Evolutionary history and higher order
classification of AAA+ ATPases. J Struct Biol. 2004;146:11–31; doi:

106. Luttringer F, Mulliez E, Dublet B, Lemaire D, Fontecave M. The Zn center of the anaerobic
ribonucleotide reductase from *E. coli*. JBIC Journal of Biological Inorganic Chemistry .
2009;14:923–33; doi:10.1007/s00775-009-0505-9

107. Panta PR, Doerrler WT. A link between pH homeostasis and colistin resistance in
bacteria. Sci Rep. 2021;11:13230; doi:10.1038/s41598-021-92718-7

108. Dapa T, Fleurier S, Bredeche M-F, Matic I. The SOS and RpoS Regulons Contribute to
Bacterial Cell Robustness to Genotoxic Stress by Synergistically Regulating DNA Polymerase
Pol II. Genetics. 2017;206:1349–60; doi:10.1534/genetics.116.199471

109. Passariello C, Forleo C, Micheli V, Schippa S, Leone R, Mangani S, et al. Biochemical
characterization of the class B acid phosphatase (AphA) of *Escherichia coli* MG1655.
Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics. 2006;1764:13–9;
doi:10.1016/j.bbapap.2005.08.028

110. Chen SH, Byrne RT, Wood EA, Cox MM. *Escherichia coli rad* gene: a novel function
involved in radiation resistance and double-strand break repair. Mol Microbiol. 2015;95:754–
68; doi:10.1111/mmi.12885

111. Chuang SE, Blattner FR. Characterization of twenty-six new heat shock genes of
*Escherichia coli*. J Bacteriol. 1993;175:5242–52; doi:10.1128/jb.175.16.5242-5252.1993

112. Manganelli R, Gennaro ML. Protecting from Envelope Stress: Variations on the Phage-
Shock-Protein Theme. Trends Microbiol. 2017;25:205–16; doi:10.1016/j.tim.2016.10.001

113. Freeman ZN, Dorus S, Waterfield NR. The KdpD/KdpE Two-Component System:
Integrating K+ Homeostasis and Virulence. PLoS Pathog. 2013;9:e1003201-;
doi:10.1371/journal.ppat.1003201

114. Meadows JA, Wargo MJ. Carnitine in bacterial physiology and metabolism. Microbiology
(N Y). 2015;161:1161–74; doi:10.1099/mic.0.000080

115. Kolata P, Efremov RG. Structure of *Escherichia coli* respiratory complex I reconstituted
into lipid nanodiscs reveals an uncoupled conformation. Swartz KJ, Carter AP, Kumar V, Letts
JA, editors. Elife. 2021;10:e68710; doi:10.7554/eLife.68710

116. Korea C-G, Badouraly R, Prevost M-C, Ghigo J-M, Beloin C. *Escherichia coli* K-12
possesses multiple cryptic but functional chaperone–usher fimbriae with distinct surface
specificities. Environ Microbiol. 2010;12:1957–77; doi:10.1111/j.1462-2920.2010.02202.x

117. Fu H, Lee J, Wang T. Heat-shock increases rpoD-dependent β-galactosidase activity in
the *Escherichia coli* strains BD792 and B23. 2015;19

118. Gagarinova A, Hosseinnia A, Rahmatbakhsh M, Istace Z, Phanse S, Moutaoufik MT, et
al. Auxotrophic and prototrophic conditional genetic networks reveal the rewiring of
transcription factors in *Escherichia coli*. Nat Commun. 2022;13:4085; doi:10.1038/s41467-
022-31819-x

119. Prusiner S, Davis JN, Stadtman ER. Regulation of glutaminase B in *Escherichia coli*. I.
Purification, properties, and cold lability. Journal of Biological Chemistry. 1976;251:3447–56;
doi: 10;251(11):3447-56

120. Iacometti C, Marx K, Hönick M, Biletskaia V, Schulz-Mirbach H, Satanowski A, et al.
Activating silent glycolysis bypasses *Escherichia coli*; bioRxiv. 2021;2021.11.18.468982; doi:
10.34133/2022/9859643

121. Njenga R, Boele J, Öztürk Y, Koch H-G. Coping with stress: How bacteria fine-tune
protein synthesis and protein transport. Journal of Biological Chemistry. 2023;299:105163;
doi:10.1016/j.jbc.2023.105163

122. Hajra D, Nair AV, Chakravortty D. An elegant nano-injection machinery for sabotaging
the host: Role of Type III secretion system in virulence of different human and animal
pathogenic bacteria. Phys Life Rev. 2021;38:25–54; doi:10.1016/j.plrev.2021.05.007

123. Peek J, Christendat D. The shikimate dehydrogenase family: Functional diversity within
a conserved structural and mechanistic framework. Arch Biochem Biophys. 2015;566:85–99;
doi:10.1016/j.abb.2014.12.006

124. Miwa T, Taguchi H. *Escherichia coli* small heat shock protein IbpA plays a role in
regulating the heat shock response by controlling the translation of σ32. Proceedings of the
National Academy of Sciences. 2023;120:e2304841120; doi:10.1073/pnas.2304841120

125. Brocklehurst KR, Hobman JL, Lawley B, Blank L, Marshall SJ, Brown NL, et al. ZntR is
a Zn(II)-responsive MerR-like transcriptional regulator of zntA in *Escherichia coli*. Mol
Microbiol. 1999;31:893–902; doi: 10.1046/j.1365-2958.1999.01229.x

126. T CP. A Galvanizing Story—Protein Stability and Zinc Homeostasis. J Bacteriol.
2007;189:2953–4; doi:10.1128/jb.00173-07

127. Iannuzzi C, Adrover M, Puglisi R, Yan R, Temussi PA, Pastore A. The role of zinc in the
stability of the marginally stable IscU scaffold protein. Protein Sci. 2014;23(9):1208-1219.
doi:10.1002/pro.2501.

128. Joly N, Engl C, Jovanovic G, Huvet M, Toni T, Sheng X, et al. Managing membrane
stress: the phage shock protein (Psp) response, from molecular mechanisms to physiology.
FEMS Microbiol Rev. 2010;34:797–827; 10.1111/j.1574-6976.2010.00240.x

129. LeRoux M, Culviner PH, Liu YJ, Littlehale ML, Laub MT. Stress Can Induce Transcription
of Toxin-Antitoxin Systems without Activating Toxin. Mol Cell. 2020;79:280-292.e8;
doi:10.1016/j.molcel.2020.05.028

130. Flores-Kim J, Darwin AJ. The Phage Shock Protein Response. Annu Rev Microbiol.
2016;70:83–101; doi:10.1146/annurev-micro-102215-095359

131. Walsh P, Bursać D, Law YC, Cyr D, Lithgow T. The J-protein family: modulating protein
assembly, disassembly and translocation. EMBO Rep. 2004;5:567-571–571;
doi:10.1038/sj.embor.7400172

132. Park DM, Akhtar MdS, Ansari AZ, Landick R, Kiley PJ. The Bacterial Response Regulator
ArcA Uses a Diverse Binding Site Architecture to Regulate Carbon Oxidation Globally. PLoS
Genet. 2013;9:e1003839; doi:10.1371/journal.pgen.1003839

133. Anand A, Patel A, Chen K, Olson CA, Phaneuf P V, Lamoureux C, et al. Laboratory
evolution of synthetic electron transport system variants reveals a larger metabolic respiratory
system and its plasticity. Nat Commun. 2022;13:3682; 10.1038/s41467-022-30877-5

134. Henry C, Loiseau L, Vergnes A, Vertommen D, Mérida-Floriano A, Chitteni-Pattu S, et
al. Redox controls RecA protein activity via reversible oxidation of its methionine residues.
Kowalczykowski SC, Storz G, editors. Elife. 2021;10:e63747; doi:10.7554/eLife.63747
1376
1377 135. Flint DH, Emptage MH, Guest JR. Fumarase A from *Escherichia coli:* purification and
1378 characterization as an iron-sulfur cluster containing enzyme. Biochemistry .
1379 1992;31:10331–7; doi:10.1021/bi00157a022
1380
1381 136. Park S-J, Tseng C-P, Gunsalus RP. Regulation of succinate dehydrogenase (sdhCDAB)
1382 operon expression in *Escherichia coli* in response to carbon supply and anaerobiosis: role of
1383 ArcA and Fnr. Mol Microbiol. 1995;15:473–82; doi:10.1111/j.1365-2958.1995.tb02261.x
1384
1385 137. Roth M, Jaquet V, Lemeille S, Bonetti E-J, Cambet Y, Francois P, et al. Transcriptomic
1386 Analysis of *E. coli* after Exposure to a Sublethal Concentration of Hydrogen Peroxide
1387 Revealed a Coordinated Up-Regulation of the Cysteine Biosynthesis Pathway. Antioxidants.
1388 2022;11; doi:10.3390/antiox11040655
1389
1390 138. Friedrich T, Dekovic DK, Burschel S. Assembly of the *Escherichia coli* NADH:ubiquinone
1391 oxidoreductase (respiratory complex I). Biochimica et Biophysica Acta (BBA) - Bioenergetics.
1392 2016;1857:214–23; doi:10.1016/j.bbabio.2015.12.004
1393
1394 139. Prasad I, Young B, Schaefler S. Genetic Determination of the Constitutive Biosynthesis
1395 of Phospho-β-Glucosidase A in *Escherichia coli* K-12. J Bacteriol. 1973;114:909–15.
1396 10.1128/jb.114.3.909-915.1973
1397
1398 140. Kuznetsova E, Proudfoot M, Gonzalez CF, Brown G, Omelchenko M V, Borozan I, et al.
1399 Genome-wide Analysis of Substrate Specificities of the *Escherichia coli* Haloacid
1400 Dehalogenase-like Phosphatase Family*. Journal of Biological Chemistry. 2006;281:36149–
1401 61; doi:10.1074/jbc.m605449200
1402
1403 141. Sobota JM, Imlay JA. Iron enzyme ribulose-5-phosphate 3-epimerase in *Escherichia coli*
1404 is rapidly damaged by hydrogen peroxide but can be protected by manganese. Proceedings
1405 of the National Academy of Sciences. 2011;108:5402–7; doi:10.1073/pnas.1100410108
1406
1407 142. Chuanboon K, Na Nakorn P, Pannengpetch S, Laengsri V, Nuchnoi P, Isarankura-Na-
1408 Ayudhya C, et al. Proteomics and bioinformatics analysis reveal potential roles of cadmium-
1409 binding proteins in cadmium tolerance and accumulation of *Enterobacter cloacae*; PeerJ.
1410 2019;7:e6904; https://pubmed.ncbi.nlm.nih.gov/31534833/
1411
1412 143. Imlay JA, Linn S. Bimodal pattern of killing of DNA-repair-defective or anoxically grown
1413 *Escherichia coli* by hydrogen peroxide. J Bacteriol. 1986;166:519–27;
1414 doi:10.1128/jb.166.2.519-527.1986
1415
1416 144. McCarthy DJ, Smyth GK. Testing significance relative to a fold-change threshold is a
1417 TREAT. Bioinformatics. 2009;25:765–71; doi:10.1093/bioinformatics/btp053
1418
1419 145. Umbarger HE, Brown B. Threonine deamination in *Escherichia coli*. II. Evidence for two
1420 L-threonine deaminases. J Bacteriol. 1957;73(1):105–12; doi:10.1128/jb.73.1.105-112.1957.
1421
1422 146. Julien M, Elisa G, Ronald VE, Enrique TMO, Ladan K, Laleh K, et al. Gene Erosion Can
1423 Lead to Gain-of-Function Alleles That Contribute to Bacterial Fitness. mBio.
1424 2021;12:10.1128/mbio.01129-21; doi:10.1128/mbio.01129-21
1425

1426 147. Greenfield D, McEvoy AL, Shroff H, Crooks GE, Wingreen NS, Betzig E, et al. Self-
1427 Organization of the *Escherichia coli* Chemotaxis Network Imaged with Super-Resolution Light
1428 Microscopy. PLoS Biol. 2009;7:e1000137; doi:10.1371/journal.pbio.1000137
1429
1430 148. Guinote IB, Moreira RN, Barahona S, Freire P, Vicente M, Arraiano CM. Breaking through
1431 the stress barrier: the role of BolA in Gram-negative survival. World J Microbiol Biotechnol.
1432 2014;30:2559–66; 10.1007/s11274-014-1702-4
1433
1434 149. Hemm MR, Paul BJ, Miranda-Ríos J, Zhang A, Soltanzad N, Storz G. Small stress
1435 response proteins in *Escherichia coli*: proteins missed by classical proteomic studies. J
1436 Bacteriol. 2010;192(1):46–58; doi:10.1128/JB.00872-09.
1437
1438 150. J RB, H DR, E PR, L SH, I BP, Aindrila M, et al. Functional Genomic Study of Exogenous
1439 n-Butanol Stress in *Escherichia coli*. Appl Environ Microbiol. 2010;76:1935–45;
1440 doi:10.1128/AEM.02323-09
1441
1442 151. Chen B, Chu T-W, Chiu K, Hong M-C, Wu T-M, Ma J-W, et al. Transcriptomic analysis
1443 elucidates the molecular processes associated with hydrogen peroxide-induced diapause
1444 termination in Artemia-encysted embryos. PLoS One. 2021;16:e0247160;
1445 doi:10.1371/journal.pone.0247160
1446
1447 152. Allison H, Jacquelyn S, Alexandra K, Kathleen C, Matthew Y, Dennis W, et al. The
1448 Bacterial iprA Gene Is Conserved across Enterobacteriaceae, Is Involved in Oxidative Stress
1449 Resistance, and Influences Gene Expression in *Salmonella enterica* Serovar Typhimurium. J
1450 Bacteriol. 2016;198:2166–79; doi:10.1128/jb.00144-16
1451
1452 153. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on metabolism,
1453 oxidative stress, and diseases. Amino Acids. 2017;49:2091–8; doi:10.1007/s00726-017-2494-
1454 2
1455
1456 154. Clausznitzer D, Oleksiuk O, Løvdok L, Sourjik V, Endres RG. Chemotactic Response
1457 and Adaptation Dynamics in *Escherichia coli*. PLoS Comput Biol. 2010;6:e1000784;
1458 doi:10.1371/journal.pcbi.1000784
1459
1460 155. Bouillet S, Hamdallah I, Majdalani N, Tripathi A, Gottesman S. A negative feedback loop
1461 is critical for recovery of RpoS after stress in Escherichia coli. bioRxiv.
1462 2023;2023.11.09.566509; doi:10.1101/2023.11.09.566509.
1463
1464 156. Song S, Wood TK. A primary physiological role of toxin/antitoxin systems is phage
1465 inhibition. Front Microbiol. 2020;11:1895; doi:10.3389/fmicb.2020.01895.
1466
1467 157. Schmidt M, Proctor T, Diao R, Freddolino PL. *Escherichia coli* YigI is a conserved
1468 Gammaproteobacterial acyl-CoA thioesterase permitting metabolism of unusual fatty acid
1469 substrates. J Bacteriol. 2022;204:e00014-22; doi:10.1128/JB.00014-22.
1470
1471 158. Jack RL, Sargent F, Berks BC, Sawers G, Palmer T. Constitutive expression of
1472 *Escherichia coli* tat genes indicates an important role for the twin-arginine translocase during
1473 aerobic and anaerobic growth. J Bacteriol. 2001;183(5):1801–4; doi:10.1128/jb.183.5.1801-
1474 1804.2001.
1475
1476 159. Zhao R, Li Z, Sun Y, Ge W, Wang M, Liu H, et al. Engineered *Escherichia coli* Nissle
1477 1917 with urate oxidase and an oxygen-recycling system for hyperuricemia treatment. Gut
1478 Microbes. 2022;14:2070391; doi:10.1080/19490976.2022.2070391
1479

160. Hryniewicz M, Sirko A, Pałucha A, Böck A, Hulanicka D. Sulfate and thiosulfate transport
in *Escherichia coli* K-12: identification of a gene encoding a novel protein involved in thiosulfate
binding. J Bacteriol. 1990;172(6):3358–66; doi:10.1128/jb.172.6.3358-3366.1990.

161. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database
resources of the National Center for Biotechnology. 2022; doi:10.1093/nar/gkab1112

162. Krueger F, James F, Ewels P, Afyounian E, Weinstein M, Schuster-Boeckler B, et al.
FelixKrueger/TrimGalore: v0.6.10 - add default decompression path. Zenodo; 2023;
doi:10.5281/zenodo.7598955

163. FastQC. 2015. Available from: https://qubeshub.org/resources/fastqc

164. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9:357–9; doi:10.1038/nmeth.1923

165. Wang L, Wang S, Li W. RSeQC: quality control of RNA-seq experiments. Bioinformatics.
2012;28:2184–5; doi:10.1093/bioinformatics/bts356

166. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30;
doi:10.1093/bioinformatics/btt656

167. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics. 2016;32:3047–8. Available from:
https://doi.org/10.1093/bioinformatics/btw354

168. Lamoureux CR, Decker KT, Sastry A V, Rychel K, Gao Y, McConn JL, et al. A multi-scale
expression and regulation knowledge base for *Escherichia coli*. Nucleic Acids Res.
2023;51:10176–93; doi:10.1093/nar/gkad750

169. Consortium TGO, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al. The
Gene Ontology knowledgebase in 2023. Genetics. 2023;224:iyad031;
doi:10.1093/genetics/iyad031

170. Gumienny R. GEOparse: Python Library to Access Gene ExpressionOmnibus
Database(GEO). Available from: https://pypi.org/project/GEOparse/

171. Kim D, Seo SW, Gao Y, Nam H, Guzman GI, Cho B-K, et al. Systems assessment of
transcriptional regulation on central carbon metabolism by Cra and CRP. Nucleic Acids Res .
2018;46:2901–17; doi:10.1093/nar/gky069

172. Spearman Rank Correlation Coefficient. The Concise Encyclopedia of Statistics. New
York, NY: Springer New York; 2008. p. 502–5. Available from: https://doi.org/10.1007/978-0-
387-32833-1_379

173. Martelli PL, Fariselli P, Casadio R. An ENSEMBLE machine learning approach for the
prediction of all-alpha membrane proteins. Bioinformatics. 2003;19:i205–11. ;
doi:10.1093/bioinformatics/btg1027

174. Pierleoni A, Martelli PL, Fariselli P, Casadio R. BaCelLo: a balanced subcellular
localization predictor. Bioinformatics. 2006;22:e408–16; doi:10.1093/bioinformatics/btl222

175. Pierleoni A, Martelli PL, Casadio R. PredGPI: a GPI-anchor predictor. BMC
Bioinformatics. 2008;9:392; doi:10.1186/1471-2105-9-392

176. Pierleoni A, Martelli PL, Casadio R. MemLoci: predicting subcellular localization of membrane proteins in eukaryotes. Bioinformatics. 2011;27:1224–30; doi:10.1093/bioinformatics/btr108

177. Savojardo C, Fariselli P, Casadio R. BETAWARE: a machine-learning tool to detect and predict transmembrane beta-barrel proteins in prokaryotes. Bioinformatics . 2013;29:504–5; doi:10.1093/bioinformatics/bts728

178. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING database in 2021: customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Res. 2021;49:D605–12; 10.1093/nar/gkaa1074

179. Fernando PC, Mabee PM, Zeng E. Integration of anatomy ontology data with protein–protein interaction networks improves the candidate gene prediction accuracy for anatomical entities. BMC Bioinformatics. 2020;21:442; doi:10.1186/s12859-020-03773-2

180. Remmert M, Biegert A, Hauser A, Söding J. HHblits: lightning-fast iterative protein sequence searching by HMM-HMM alignment. Nat Methods. 2012;9:173–5; doi:10.1038/nmeth.1818

181. Bagheri H, Dyer R, Severin A, Rajan H. Comprehensive Analysis of Non Redundant Protein Database. 2020; doi:10.21203/rs.3.rs-54568/v1

182. Andreeva A, Kulesha E, Gough J, Murzin AG. The SCOP database in 2020: expanded classification of representative family and superfamily domains of known protein structures. Nucleic Acids Res. 2020;48:D376–82. Available from:10.1093/nar/gkz1064

183. Barrio-Hernandez I, Yeo J, Jänes J, Mirdita M, Gilchrist CLM, Wein T, et al. Clustering predicted structures at the scale of the known protein universe. Nature. 2023;622:637–45; doi:10.1038/s41586-023-06510-w

184. Tuttle AR, Trahan ND, Son MS. Growth and Maintenance of *Escherichia coli* Laboratory Strains. Curr Protoc. 2021;1:e20; doi:10.1002/cpz1.20

185. Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, et al. The RIN: an RNA integrity number for assigning integrity values to RNA measurements. BMC Mol Biol. 2006;7:3; doi:10.1186/1471-2199-7-3

186. Lataretu M, Hölzer M. RNAflow: An Effective and Simple RNA-Seq Differential Gene Expression Pipeline Using Nextflow. Genes (Basel). 2020;11:1487; doi:10.3390/genes11121487

187. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34:i884–90; doi:10.1093/bioinformatics/bty560

188. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34:i884–90; doi:10.1093/bioinformatics/bty560

189. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. Nat Methods. 2015;12:357–60; doi:10.1038/nmeth.3317

190. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.;
doi:10.1093/bioinformatics/btp352

191. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30;
doi:10.1093/bioinformatics/btt656

192. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15:550; doi:10.1186/s13059-014-0550-8

193. Duda JC, Drenda C, Kästel H, Rahnenführer J, Kappenberg F. Benefit of using interaction
effects for the analysis of high-dimensional time-response or dose-response data for two-
group comparisons. Sci Rep. 2023;13:20804; doi:10.1038/s41598-023-47057-0

194. Liou G-G, Chao Kaberdina A, Wang W-S, Kaberdin VR, Lin-Chao S. Combined
Transcriptomic and Proteomic Profiling of *E. coli* under Microaerobic versus Aerobic
Conditions: The Multifaceted Roles of Noncoding Small RNAs and Oxygen-Dependent
Sensing in Global Gene Expression Control. Int J Mol Sci. 2022;23; doi:10.3390/ijms23052570

63