Investigating the role of urban vegetation alongside other

environmental variables in shaping *Aedes albopictus*

presence and abundance in Montpellier, France

Colombine Bartholomée[1*], Paul Taconet[1], Mathilde Mercat[1], Coralie Grail[1], Emilie Bouhsira[2],

Florence Fournet[1] and Nicolas Moiroux[1*]

[1] MIVEGEC, Univ. Montpellier, CNRS, IRD, Montpellier, France

[2] InTheres, Univ. Toulouse, INRAE, ENVT, Toulouse, France

Corresponding authors

E-mail: colombine.bartholomee@ird.fr (CB), nicolas.moiroux@ird.fr (NM)

Abstract

Urban greening helps address urbanization challenges, but it may also favor mosquito species, vectors of pathogens causing human diseases. This study examines the relationship between urban vegetation and the presence and abundance of *Aedes albopictus* in Montpellier, the second greenest city of France, while accounting for meteorology, microclimate, air quality, human socio-demography, and landscape.

From May to October 2023, adult mosquitoes were collected monthly in urban parks, residential areas, and the highly impervious city center using BG-Pro traps with odor and CO_2 attractants. Microclimate data (air temperature and relative humidity) were recorded on-site at each trap location using Hygrobutton data-loggers. Vegetation, land cover, meteorological, air quality, and human demographic data were gathered from open-access databases. *Ae. albopictus* presence and abundance were analysed according to environmental variables taken at different time lags and spatial distances using a two-stage modeling approach: bivariate analyses using generalized linear mixed models were conducted to select variables for inclusion in a multivariate random forest model, aiming to identify the factors that best explain *Ae. albopictus* presence and abundance.

While urban vegetation had a limited effect on *Ae. albopictus* presence, the average patch size, and the percentage of area covered by low vegetation were among the most important predictors of abundance. The main predictors for presence were minimum hourly temperature (24h-48h before sampling), minimum atmospheric pressure during sampling, and the weekly cumulated rainfall recorded six weeks before sampling. The most important predictors of abundance were the average patch size of low vegetation, the maximum

46 hourly temperatures during sampling, and the length of roads. To our knowledge, this is the

47 first study examining urban vegetation's influence on *Ae. albopictus* in France, offering

48 insights for urban planning and suggesting further research on vegetation's role in mosquito-

49 borne disease transmission, particularly in the context of increasing dengue incidence in

50 Europe.

Introduction

According to Hussain and Imitiaz [1], urbanisation is a complex, dynamic, and evolving social

process that drives the emergence and growth of cities. It occurs unevenly across time and

space, with disparities between continents and countries. Cities are mosaics of

heterogeneous patches of diverse ecosystems with multiple interfaces [2]. Urbanisation can

improve public health by enhancing living standards for citizens [3]. However, it also results

in habitat loss and fragmentation, leading to a decline in biodiversity, as observed in China

for birds species diversity [4]. Urbanisation also exacerbates air pollution [5], contributing to

human health issues [6]. Moreover, rapid urbanisation often accentuates social disparities,

as access to green spaces, healthcare, and healthy living conditions is not equally distributed

across urban populations. These inequalities, together with reduced opportunities for

contact with nature, increase vulnerability to stress and mental health disorders [7]. Finally,

the expansion of asphalt-covered areas that absorb energy during the day and release it as

heat at night amplifies the urban heat island (UHI) effect [8], worsening its impact on human

health [9].

Given these multiple pressures, strategies that simultaneously address environmental and

social challenges are increasingly needed. Nature-Based Solutions (NBS), such as urban

greening, are promoted to achieve the 11th Sustainable Development Goal: "to make cities

and human settlements inclusive, safe, resilient, and sustainable" [10]. NBS also support the

12th target of the COP 15 Biodiversity Plan for Life on Earth, aiming to "enhance green

spaces and urban planning for human well-being and biodiversity" [11], and align with the

sustainability goals of the European Green Deal [12]. Urban greening involves deliberately

integrating vegetation into urban areas (13), encompassing initiatives such as soil de-

imperviousness, developing green and blue infrastructures, conserving urban green spaces

(UGS), and promoting urban agriculture. UGS provide cultural and regular ecosystem

services [13]. They help mitigate the UHI effect [14,15], and contribute to improving mental

health [16]. Effective management of UGS can help sequester carbon dioxide [17] through

tree plantations and enhance animal biodiversity by increasing the presence of indigenous

vegetation [18].

Urban greening may also have negative side effects. It can increase pollen allergies [19] and

attract agricultural pests like tree moths [20]. UGS increase human-wildlife interactions,

raising the risk of zoonotic [21] and arthropod-borne diseases [22], as shown for the Lyme

disease pathogen (*Borrelia burgdorferi*) transmitted by ticks in urban parks across Europe

[23]. In Europe, while numerous studies have explored the relationship between UGS and

tick populations, the relationship between urban greening and mosquitoes is less

documented [24], with only one study on *Culex pipiens* in the Netherlands [25] and two

studies on the tiger mosquito *Aedes albopictus* in Italy [26,27]. As shown in the scoping

review by Mercat *et al* (2025) [24], the effects of urban green infrastructures (UGIs) on

mosquito populations are highly dependent on the vector system studied and the

environmental context. Urban greening is a dynamic process, and its environmental changes

may take time to manifest [28]. While it can create favorable conditions for vectors and

increase human-vector interactions, it may also promote the establishment of predators

that could regulate vector populations.

Ae. albopictus is an invasive mosquito in Europe. It is native to Southeast Asia and a vector of

several infectious agents, including the dengue virus (DENV), chikungunya virus (CHIKV), and

Zika virus. The species adapted to anthropogenic environments [29], including urban areas,

using natural and artificial breeding sites. It can survive extreme conditions through egg

diapause [30]. Its abundance is influenced by environmental factors such as weather [31],

microclimate [32], and landscape features [33]. Urban vegetation may favour *Ae. albopictus*

by creating cooler and more humid microclimates during warm season [34], facilitating

access to sugar meals, providing natural resting sites [35], and attracting blood sources

(humans and animals). In France, since its first detection in 2004 [36], *Ae. albopictus* spreads

to 78 of 96 metropolitan France's departments by 2024, including all departments in the

Occitanie region, where it has been linked to several autochthonous arboviral disease

clusters since 2014.

The Occitanie region is actively pursuing urban greening projects in its two largest cities,

Toulouse and Montpellier, including plans to plant 150 000 trees. As Occitanie is a hotspot

for both arboviral diseases and urban greening, it is crucial to assess the impact of these

greening initiatives on *Ae. albopictus* population dynamics. With the goal to guide future

research to improve the design and planning of urban green spaces, this study aims to

evaluate the influence of urban vegetation in Montpellier in 2023 on the presence and

abundance of *Ae. albopictus* females, using an explanatory approach taking into account

land cover, microclimate, meteorology, air quality, and socio-demographics.

Material and methods

Mosquito sampling and study areas

Study areas

Montpellier is a city of 300 000 inhabitants located in the Occitanie region in southern France (Fig 1). The city has a Mediterranean climate with an average annual temperature of 15°C and an average annual precipitation of 739 mm [37]. The city features more than 10% of vegetated areas within densely urbanized areas. It ranks among the top three greenest cities in metropolitan France [38]. The study was conducted in nine areas belonging to three different environments: two urban parks (PRK), three residential neighborhoods (RES), and four highly impervious (i.e., areas of the city center dominated by sealed surfaces such as concrete and asphalt) areas (IMP) (Fig 1). Urban parks (Botanical Garden (PRK-BOT), and Park of Aiguelongue (PRK-AGL)) and residential neighborhoods (Lemasson (RES-LEM), Aiguerelles (RES-AGR), and Soulas (RES-SOUL)) were vegetated areas. The highly impervious areas included two areas with some vegetation (Saint Charles University (IMP-SCU) and Acapulco Hotel (IMP-ACA)) and two with little to no vegetation (Buisson Bertrand Institute (IMP-BBI) and Diderot School (IMP-DID)) (Fig 1). Detailed descriptions of study areas are provided in S1 Table, and images of study areas in S1 Fig. The minimum distance between study areas was 115 m. Although *Aedes* mosquitoes can disperse from 75 m [39] to 290 m [40] in urban areas over 8 days, we treated the study areas as independent units due to impervious roads between them, that are known to limit mosquito dispersal [41,42].

Fig 1: **Study areas**. Fig 1A and 1B show the location of Montpellier in France (map from Map data © OpenStreetMap contributors, licensed under ODbL) and the land cover map of Montpellier with the study areas, respectively. Sampling environments are shown in the legend of Fig 1B, delineating urban parks, residential areas, and highly impervious areas (boundaries from Montpellier Open Data). Land cover categories are shown in the legend of

Fig 1B. Fig 1C, 1D, and 1E show a garden of a residential area, the southern part of the

Botanical Garden (PRK-BOT), the Bouisson Bertrand Institute (IMP-BBI) and the Diderot

School (IMP-DID) respectively.

Mosquito sampling

Adult mosquitoes were captured using BG-Pro traps (BioGents, Regensburg, Germany)

modified according to L'Ambert *et al* [43], each equipped with a BG lure—synthetic human

odor mimic—and a daily replenished carbon dioxide source made from yeast, sugar, and

water (S1 Fig). Two BG-Pro traps were installed in each residential area (RES-LEM, RES-AGR,

RES-SOUL), and in the park of Aiguelongue (PRK-AGL). Due to their smaller size (S1 Table),

only one BG-Pro trap was deployed in each of the highly impervious areas (IMP-ACA, IMP-

BBI, IMP-DID and IMP-SCU). Because of its larger size (S1 Table), three BG-Pro traps were

deployed in the Botanical Garden (PRK-BOT).

Traps were placed in shaded, wind-sheltered sites at a height of one meter. Therefore, 15

sites were sampled: five in the two urban parks, six in the three residential areas, and four in

highly impervious areas. Every trap was separated by a minimum distance of 50 m to

prevent interferences [44]. However, due to accessibility constraints, the two traps in RES-

SOUL were placed 22 m apart.

One 48-hour trapping session was conducted per site per month from May to October 2023.

During each session, mosquitoes were retrieved each morning, resulting in two collections

per session. Mosquitoes were transported at 4°C to the laboratory (MIVEGEC laboratory, IRD

regional delegation, Montpellier), and stored at -20°C. Mosquito individuals were then

counted, sorted by sex, and morphologically identified using the multicriteria key

MosKeyTool [45]. These data are available on GBIF [46].

Environmental data collection and preparation

To assess the impact of urban vegetation on the presence and abundance of *Ae. albopictus*,

it was essential to consider all factors that may influence its population dynamics. For this

task, we adapted the conceptual model of human-*Anopheles* contact by Moiroux (2012) [47]

to the case of *Ae. albopictus*. Based on a literature review, we documented and

hypothesised known or potential predictors and their pathways to mosquito abundance

[48]. This model guided the selection of environmental data to be collected as described

below.

Fine-scale vegetation and land cover data

A harmonised land-cover map of Montpellier was generated at 0.5 m resolution by

combining three complementary datasets (see Supplementary S2 Table): (i) the fine-scale

vegetation dataset of Montpellier (Montpellier Méditerranée Métropole, 2019), which

distinguishes vegetation patches using deep learning and photo-interpretation of Pléiades

satellite imagery and a LiDAR-derived elevation model; (ii) the French national building

database (Base de données nationale des bâtiments, BDNB); and (iii) the Copernicus Urban

Atlas. Following rasterisation and integration of these sources, five land-cover categories

were retained (Fig 1B): buildings, roads, low vegetation (< 3 m height), high vegetation (> 3

m height), and "others" (including railways, pathways, industrial areas, and sports facilities).

For each trap location, spatial metrics were computed within circular buffer zones of 20 m,

50 m, 100 m, and 250 m radii. For each land cover category, we calculated the percentage

cover (proportion of buffer area occupied by the category), total edge length (sum of the

perimeters of all patches of the category), and average patch size (mean area of individual

patches). For vegetation categories (low vegetation < 3 m height, high vegetation > 3 m

height), we also extracted the number of patches. The landscape diversity (Shannon index)

was calculated for all buffer zones. This index quantifies diversity by considering both the

number of land cover categories and their relative proportions, with higher values indicating

more heterogeneous landscapes (see Supplementary S2 Table for the formula). Buffer sizes

were chosen according to previous works investigating the effect of urban vegetation [27,49]

on the dispersal of *Ae. albopictus* [40]. Supplementary S3 Table provides the land cover

variable values for each trap and buffer radii.

Microclimatic data

We continuously collected hourly temperature and relative humidity near each trap using

Hygro Button data loggers from May to October 2023. The minimum (lowest recorded

value), maximum (highest recorded value), and mean temperatures, as well as relative

humidity, were recorded for 24-hour periods starting from 48 hours prior to the beginning of

each sampling session and continuing until the end of the session (S2 Table).

Meteorological and air quality data

Hourly temperature and rainfall data were obtained from the Observatoire Départemental

de l'Eau et de l'Environnement (ODEE) station located at Château d'Ô in the north-west of

Montpellier (S2 Table). Relative humidity and wind speed data were obtained from the

Frejorgues airport station (belonging to the Meteo France national meteorology agency

network) at a 3-hour resolution (S2 Table). Growing Degree Days (GDD), a known predictor

of mosquito development, were calculated from daily mean temperatures above 11°C, as

this was shown to be the annual mean temperature limit for the establishment of *Ae.*

albopictus populations [50]. Temperatures above this level are conducive to mosquito

development. For each sampling event, weather data were aggregated at a weekly

resolution for the sampling week (W0) and for all possible lag periods between W−6 and W0.

These lag periods ranged in duration from one to seven consecutive weeks (e.g., W−2 only,

W−4 to W−2, W−6 to W0). Rainfall and GDD were summed, while temperature, humidity,

and wind speed were averaged over each lagged window. Precipitation data from the

Château d'Ô station were also aggregated over 24-hour periods prior to and during each

sampling session, following the same temporal aggregation approach used for microclimate

data from the data loggers, because precipitation may influence mosquito activity and

therefore sampling.

As atmospheric pressure can influence insects' flight and activity [51–53], we evaluate the

effect of daily atmospheric pressure on *Ae. albopictus*. For this purpose, the minimum

(lowest recorded value), maximum (highest recorded value), and mean atmospheric

pressure, measured every three hours at the Frejorgues airport station, were extracted per

24-hour periods before and during sampling sessions. The daily mean atmospheric pressure

difference between two consecutive days was calculated per 24-hour periods before and

during sampling sessions. S3 Fig depicts the temporal evolution of these meteorological

variables (temperature, rainfall, wind speed, relative humidity, and atmospheric pressure).

As air pollution can influence insect behaviour and abundance [54–56], we evaluate the

effect of air quality data on *Ae. albopictus*. For each trap location, we collected daily

concentrations of nitrogen monoxide (NO), nitrogen dioxide (NO2), particulate matter (of

diameter less than 10 (PM10) and 2.5 μm (PM2.5)), and ozone (O3) from the closest station

of the Atmo Occitanie air quality observatory (S2 Table). As for weather data, air quality data

were aggregated at a weekly resolution for the sampling week (W0) and for all possible lag

periods between W–6 and W0, by averaging the daily concentrations of each pollutant over

these windows.

Social and demographic data

Socioeconomic factors such as low living standards [57], aging or degrading infrastructure

[58], and population densities [59] can influence mosquito presence. Number of buildings of

different ages and number of low-income households (2019) were obtained from the Filosofi

database of the French Institut National de la Statistique et des Études Économiques (INSEE),

provided as a 200 m resolution grid. For each buffer area, values of intersecting grid cells

were summed. Human population data were obtained from the Montpellier fine-scale

population dataset of 2020 (Supplementary S2 Table), in which IRIS-level population counts

(~2000 inhabitants per unit) were redistributed across land plots in proportion to living area

and assigned to the centroid of significant buildings. Population counts were then

aggregated within the same four buffer zones (20, 50, 100, and 250 m radii).

Assessment of vector control interventions

No municipal adulticide or larvicide treatments were implemented in the study areas during

the sampling period. No individual larval habitat removal by residents was observed.

However, potential actions by residents near some sampling sites could not be

systematically assessed, and their effects on mosquito abundance cannot be excluded.

Statistical analyses of the presence and abundance of female *Ae.* *albopictus*

To study the relationships between the presence and the abundance of female *Ae.* *albopictus* and environmental conditions, we analysed the data using a hurdle modeling approach, i.e., considering the data responding to two distinguished processes: one causing zero vs. non-zero (« presence » models) and one causing positive counts of females (« abundance » models). This approach is justified because determinants of mosquito presence are expected to be different than abundance ones [60].

For both presence and abundance analysis, we adopt a two-stage analysis method as described by Taconet *et al* [61]. The first step involves bivariate analyses using Generalized Linear Mixed Models (GLMMs), followed by the second step, a multivariate analysis employing an interpretable machine learning method based on random forest models [62]. The bivariate analyses aimed to improve our understanding of (i) the time-lagged effects of meteorological and air quality variables, (ii) the impact of atmospheric pressure and microclimate on the tiger mosquito, (iii) land cover-scale effects, (iv) effects of socio-economic and demographic variables, and (iv) to identify and select key variables for multivariate analysis. The multivariate analysis aimed to identify the most impactful determinants of the presence and abundance of *Ae. albopictus* and the shape of their association while considering potentially complex (e.g. non-linear) associations and interactions between variables, with the overall aim of assessing more precisely the impact of urban vegetation. The modeling workflow is summarised in S2 Fig and in S1 Text.

Bivariate analysis using GLMMs

Every explanatory variable was tested using a GLMM with crossed random intercepts for trap location, with sites nested within the sampling area, and for the sampling session. We fitted binomial and zero-truncated negative binomial models to presence/absence and positive counts data, respectively. Marginal R^2, representing the proportion of variance explained by the fixed effects excluding the contribution of random intercepts [63], was calculated and used for interpretation. Time-lagged effects of meteorological and air quality variables were interpreted using cross-correlation maps (CCMs) [64] of marginal R^2 adjusted for the direction (+/-) of the relationship.

Multivariate analysis using random forests

The results of the bivariate analyses were used to select the variables to be included in the multivariate analysis. First, we excluded variables poorly correlated with the response variables ($p > 0.2$). For each meteorological, microclimatic, air quality, land cover, and demographic variable, we retained the time lag or radii with the higher marginal R^2. Multicollinearity was addressed using Pearson's correlation coefficient with a threshold of 0.7. Then, based on empirical knowledge, an initial selection of correlated variables was made. Other collinear covariates were removed until the Variance Inflation Factor (VIF) of the remaining variables was below three [65] (S2 Fig and S1 Text). Binary classification random forests (RFs) and regression classification RFs were generated for the presence and abundance models, respectively. To improve model efficiency and reduce the number of explanatory variables, recursive feature elimination was performed for each model [66]. To account for spatial and temporal autocorrelation, sampling variables (trap location, sampling day, and sampling session) were included. For each set of hyperparameters and to assess the

predictive power of each model, both a spatial and temporal cross-validation framework

[67] was used, taking into account the potential spatial and temporal autocorrelation in the

data. Spatial cross-validation (leave-area-out) involved training the model recursively on

data from eight out of the nine sampled areas, testing it with data from the remaining area,

and finally averaging the performance metric for the nine areas [61]. The temporal cross-

validation (leave-sampling-session-out) involved training the model recursively on data from

six out of the seven sampling sessions and testing it with data from the remaining session.

The performance metrics selected were the Area Under the Curve (AUC) for the presence

models [68] and the Mean Absolute Error (MAE) for the abundance models. The model

results were interpreted using Variable Importance Plots (VIPs) [62] and Partial Dependence

Plots (PDPs) [69]. Residual spatial autocorrelation was assessed for each model.

Softwares used

The R version 4.2.3 programming language [70] was used. The land cover map creation and

landscape metrics calculation involved the use of the 'sf' [71], 'raster' [72], 'fasterize' [73],

and 'landscapemetrics' [74] packages. GLMMs were fitted using the 'glmmTMB' package

[75]. Marginal R^2 was calculated using the 'performance' package [76]. The 'correlation'

package was used to evaluate multicollinearity [77]. Multivariate analyses were performed

using the 'caret' [78] and 'ranger' [79] packages. The 'CAST' package [80] was used for the

cross-validation and the interpretation plots were performed using the 'iml' [81], 'pdp' [82],

'MLmetrics' [83], 'precrec' [84] and 'patchwork' [85] packages. The package 'spatialRF' [86]

was used to assess spatial autocorrelation. All scripts are available on Software Heritage

[87].

Results

Entomological data

A total of 2344 mosquitoes were collected, predominantly *Ae. albopictus* (85.66%, N=2008),

with smaller proportions of *Culex pipiens* (13.91%, N=326), *Culiseta longiareolata* (0.39%,

N=9), and *Culiseta annulata* (0.04%, N=1). Among 2008 collected *Ae. albopictus*, there were

1106 females (55.08%) and 902 males (44.92%). The mean density of female *Ae. albopictus*

was 3.83 mosquitoes/trap/24h (SE = 1.97) in urban parks, 3.31 mosquitoes/trap/24h (SE =

1.65) in residential areas, and 1.99 mosquitoes/trap/24h (SE = 1.08) in highly impervious

areas (Table 1). Mean mosquito densities (mosquitoes/trap/24 h), stratified by sampling

environment (or sampling areas), are presented in Table 1 (or S4 Table). The temporal

distribution of *Ae. albopictus* (S4 Fig) exhibited a bimodal pattern, with an initial emergence

in May, a primary peak in late June, and a secondary, smaller peak in late August.

Table 1: Mosquito density/trap/24h by species and sex in the three different sampling

environments.

Environment	Mosquito female /trap/24h (± SE)	Mosquito male /trap/24h (± SE)	*Ae. albopictus* female /trap/24h (± SE)	*Ae. albopictus* male /trap/24h (± SE)	*Cx. pipiens* female /trap/24h (± SE)	*Cx. pipiens* male /trap/24h (± SE)	*Cs. longiareolata* female /trap/24h (± SE)	*Cs. longiareolata* male /trap/24h (± SE)	*Cs. annulata* male /trap/24h (± SE)
Urban parks	6.46 (± 2.80)	3.59 (± 2.89)	3.83 (± 1.97)	1.83 (± 1.67)	1.48 (± 0.79)	0.49 (± 0.38)	0.02 (± 0.02)	0.02 (± 0.02)	0.01 (± 0.01)
Residential areas	3.97 (± 1.64)	1.58 (± 1.14)	3.31 (± 1.65)	0.93 (± 0.83)	0.30 (± 0.16)	0.07 (± 0.05)	0.01 (± 0.01)	0.01 (± 0.01)	0
Highly impervi	2.91 (± 1.37)	1.19 (± 0.87)	1.99 (± 1.08)	0.72 (± 0.69)	0.26 (± 0.18)	0.16 (± 0.12)	0.01 (± 0.01)	0.01 (± 0.01)	0

ous areas								

SE: Standard error

Bivariate analysis

Vegetation and other land cover variables

Fig 2 shows the vegetation and other land cover variables correlated with the probability of

presence or abundance of female *Ae. albopictus*. Two variables related to low vegetation

were significantly and positively associated with the probability of presence within a 20 m

radius buffer around the traps: the total length (in m) of low vegetation edges (OR =1.007, R^2

=0.042, p =0.042, CI 95% [1.001; 1.014]) and the average patch size (in m^2) of low vegetation

(OR =1.01, R^2 =0.07, p =0.006, CI 95% [1.003; 1.017]). The percentage of area occupied by

buildings within a 50 m radius buffer around the traps was negatively correlated with the

probability of presence (OR =0.959, R^2 =0.047, p =0.048, CI 95% [0.920; 0.999]). Within a 20

m radius buffer around the trap, the total length edge of roads and the average size of a

patch of roads were negatively correlated with the probability of presence (Fig 2).

Regarding abundance, positive associations were observed with all vegetation-related

variables except for the average patch size of low vegetation within a 50 m radius buffer that

showed a negative and significant association (RR =0.996, R^2 =0.098, p =0.0001, CI 95%

[0.994; 0.998]). The most remarkable positive association was for the total edge length of

low vegetation within a 100 m buffer (RR =1.0008, R^2 =0.190, p =0.007, CI 95% [1.0002;

1.0014]). Variables representing the building land cover category consistently showed

negative relationships with female *Ae. albopictus* abundance, especially for the percentage

area variable (across all buffer radii) and for all variables within the 100m buffer radius. RR

for the percentage of building areas in the 100m buffer was 0.945 (R^2 =0.240, p =0.0007, CI

95% [0.914; 0.976]) (Fig 2).

Fig 2: Bivariate relationships between *Ae. albopictus* and land cover variables for varied

buffer sizes. Relationships are analysed based on *Ae. albopictus* presence or abundance and

land cover variables for different buffer sizes around traps. The adjusted marginal R^2 reflects

the variance explained by the explanatory variable, adjusted for correlation direction. Boxes

are colored if the p-value was < 0.2 (No asterisk: p-value $\in$ [0.05; 0.2], *: p-value $\in$ [0.01,

0.05], **: p-value $\in$ [0.001; 0.01]; ***: p-value $\in$ [0; 0.001]). Box color depends on the

direction of the relationship (blue: negative, red: positive). Color intensity varies according to

the marginal R^2 value.

Microclimatic, meteorological, and air quality variables

Fig 3 shows the relationship between presence or abundance and microclimatic or

meteorological variables recorded from 48h before the start of sampling and during

sampling. Local relative humidity (RH) variable showed positive associations with presence

when recorded before sampling (with the strongest association for maximum hourly relative

humidity 24-48h before sampling; OR = 1.194, R^2 =0.201, p = 0.002, CI 95% [1.070; 1.332])

and negative association when recorded during sampling (with the strongest association for

minimum RH; OR =0.924, R^2 =0.134, p = 0.001, CI 95% [0.880;0.969]). Significant associations

of RH data with abundance were negative whatever the period considered.

Fig 3: Bivariate relationships between *Ae. albopictus* and microclimatic and meteorological

variables 48h before and during sampling. Relationships are analysed based on *Ae.*

albopictus presence or abundance and variables recorded 48h before the start and during

sampling sessions. Microclimatic variables include minimum hourly relative humidity

(rhmin), maximum hourly relative humidity (rhmax), mean hourly relative humidity

(rhmean), minimum hourly temperature (tmin), maximum hourly temperature (tmax) and

mean hourly temperature (tmean). Meteorological variables include daily cumulated rainfall,

minimum atmospheric pressure (Patmin), maximum atmospheric pressure (Patmax), mean

atmospheric pressure (Patmean), and the difference of mean atmospheric pressure between

two consecutive days (Patdiff). The adjusted marginal R^2 reflects the variance explained by

the explanatory variable, adjusted for correlation direction. Boxes are colored if the p-value

is < 0.2 (No asterisk: p-value $\in$ [0.05; 0.2], *: p-value $\in$ [0.01, 0.05], **: p-value $\in$ [0.001;

0.01]; ***: p-value $\in$ [0; 0.001]). Box color depends on the direction of the relationship

(blue: negative, red: positive). Color intensity varies according to the marginal R^2 value.

bef=before

Local temperatures during and before sampling consitently showed positive correlations

with presence, particularly for mean and maximum hourly temperatures recorded during

sampling (tmean: OR =1.592, R^2 =0.408, p =0.0002, CI 95% [1.251; 2.026]; tmax: OR =1.497,

R^2 =0.402, p =0.0001, CI 95% [1.226; 1.828]). Abundance was also positively associated with

tmean (RR =1.143, R^2 =0.197, p =0.0002, CI 95% [1.064; 1.228]) and tmax (RR =1.057, R^2

=0.061, p =0.07, CI 95% [0.996; 1.122]) during sampling, but with lower R^2.

Daily total precipitation was negatively associated with presence during sampling (OR

=0.863, R^2 =0.137, p =0.003, CI 95% [0.783; 0.951]). *Ae. albopictus* presence was positively

associated with all variables describing atmospheric pressure recorded during the sampling

hours, but negatively with lagged data. The strongest effect size was for the difference in

mean daily atmospheric pressure (Patdiff) between the sampling day and the previous day

(OR = 1.008, R^2 = 0.194, p = 0.008, CI 95% [1.002; 1.014]). Average (patmean) and maximum

(patmax) atmospheric pressure during the 24-48h period preceding sampling showed

negative association with abundance (RR =0.997, R^2 =0.243, p =0.018, CI 95% [0.994; 0.999]

and RR =0.998, R^2 =0.177, p =0.0001, CI 95% [0.996; 0.999], respectively).

Local temperatures during and before sampling consitently showed positive correlations

with female presence, particularly for mean and maximum hourly temperatures recorded

during sampling (tmean: OR =1.662, R^2 =0.439, p =0.0001, CI 95% [1.298; 2.129]; tmax: OR

=1.553, R^2 =0.424, p =0.0001, CI 95% [1.262; 1.991]). Abundance was also positively

associated with tmean (RR =1.134, R^2 =0.197, p =0.0001, CI 95% [1.055; 1.218]) and tmax (RR

=1.052, R^2 =0.050, p =0.098, CI 95% [0.991; 1.117]) during sampling, but with lower R^2.

Daily total precipitation was negatively associated with presence during sampling (OR

=0.859, R^2 =0.133, p =0.003, CI 95% [0.777; 0.949]). *Ae. albopictus* presence was positively

associated with all variables describing atmospheric pressure recorded during the sampling

hours, but negatively with lagged data. The strongest effect size was for the difference in

mean daily atmospheric pressure (Patdiff) between the sampling day and the previous day

(OR = 1.008, R^2 = 0.217, p = 0.003, CI 95% [1.003; 1.015]). Average (patmean) and maximum

(patmax) atmospheric pressure during the 24-48h period preceding sampling showed

negative association with abundance (RR =0.997, R^2 =0.217, p =0.049, CI 95% [0.994; 0.999]

and RR =0.998, R^2 =0.183, p =0.0003, CI 95% [0.996; 0.999], respectively).

Fig 4 shows the effect size and significance (p < 0.2) of weeks-lagged meteorological

variables on the presence and abundance of female *Ae. albopictus*. The relationship

between weekly cumulated rainfall (CUMRF) and both the probability of presence and

abundance was positive, with a statistically significant association for abundance and the

highest effect size observed the 6th week before sampling (for presence: OR = 1.430, R^2 =

0.334, p = 0.168, CI 95% [0.860; 2.375]; for abundance: RR = 1.075, R^2 = 0.127, p = 0.04, CI

95% [1.003; 1.152]). The probability of presence was positively and significantly associated

with minimum (TMIN), average (TMEAN), and maximum (TMAX) temperatures. The highest

effect size was observed for TMIN one week before sampling (OR =1.620, R^2 =0.372, p

=0.001, CI 95% [1.209; 2.172]. For abundance, relationships with TMIN, TMEAN, and TMAX

were also positive and significant for time interval 0 to 2 weeks before sampling, with the

highest effect size observed for TMAX (RR = 1.161, R^2 =0.192, p = 0.0006, CI 95% [1.067;

1.267]). Both responses were positively and significantly correlated with the weekly growing

degree days (GDD). The highest effect on presence and abundance was observed for GDD

during the week preceding sampling (OR = 1.085, R^2 = 0.357, p = 0.002, CI 95% [1.031;

1.141]) and for time interval 0 to 2 weeks before sampling (RR = 1.011, R^2 = 0.189, p =

0.0009, CI 95% [1.004; 1.017]), respectively. Both responses were negatively and significantly

correlated with relative humidity (RH), with the highest effect observed for RH recorded 6

weeks before sampling (OR = 0.792, R^2 = 0.472, p = 0.006, CI 95% [0.670; 0.937] for

presence; RR = 0.948, R^2 = 0.183, p = 0.007, CI 95% [0.912; 0.986] for abundance). The

relationship between wind speed (WS) and abundance was negative, with the highest effect

observed for time interval 0 to 5 weeks before collection (RR =0.343, R^2 = 0.086, p =0.157, CI

95% [0.078; 1 .510]).

Fig 4: Cross correlation maps showing the lagged effect of meteorological variables on *Ae.*

***albopictus* presence or abundance**. Lagged meteorological variables include weekly

cumulated daily rainfall (CUMRF), mean daily temperature (TMEAN), daily minimal

temperature (TMIN), daily maximal temperature (TMAX), the weekly Growing Degree Days

(GDD), mean daily relative humidity (RH), and mean daily wind speed (WS). Time lags are

expressed in weeks prior to sampling. The adjusted marginal R^2 reflects the variance

explained by the explanatory variable, adjusted for correlation direction. Red-bordered

squares highlight the time lag with the highest marginal R^2, with the interval indicated in the

top left corner. Black-bordered squares denote marginal R^2 values close to the highest

(within 10%). Gray squares represent correlations with p-values > 0.2.

Daily nitrogen dioxide (NO2) concentrations were positively associated with the probability

of presence of female *Ae. albopictus*, with the highest effect size observed for the sampling

week (OR = 1.109, R^2 = 0.126, p = 0.163, CI 95% [0.959; 1.282]; S5 Fig). Particulate matter

concentrations (PM2.5 and PM10) were also positively correlated with the probability of

presence and with the abundance. However, PM2.5 and PM10 concentrations were strongly

correlated with all time-lagged temperature metrics (Pearson coefficient > 0.8).

Tropospheric ozone (O3) concentration (three weeks before sampling) was negatively

correlated with the probability of presence and positively (one week before sampling) with

abundance. Daily O3 concentrations were also highly correlated with weekly cumulated

rainfall, wind speed, and daily relative humidity (Pearson coefficient > 0.8).

Socio-demographic variables

Human population density in a 250 m radius buffer around traps was positively correlated

with the *Ae. albopictus* probability of presence (OR=1.0013, R2=0.056, p=0.029, CI 95%

[1.0001; 1.0026]; S6 Fig).

511 Multivariate analysis

After bivariate analyses, multicollinearity assessments, and recursive feature elimination, ten

and nine variables, in addition to sampling design variables, were kept for the multivariate

random forest models of presence and abundance, respectively (S5 Table shows reason for

exclusion). Fig 5 and Fig 6 display the multivariate RF model interpretation plots for presence

and abundance, respectively. Fig 5A and Fig 6A illustrate the contribution of each variable in

explaining the response variable (VIPs). Fig 5B and Fig 6B depict the relationship between

the response variable and the explanatory variables (PDPs). Models of presence and

abundance did not show any residual spatial autocorrelation.

Fig 5: **Interpretation plots for the multivariate random forest model of *Ae. albopictus***

presence. The left plot (Fig 5A) is the variable importance plot (VIP). Rights plots (Fig 5B) are

smoothed partial dependence plots (PDPs) with Confidence Intervals for each selected

variable in the models. The y-axis in PDPs represents the probability of catching at least one

mosquito. The range of values in the x-axis represents the range of values available in the

data for the considered variable. The rugs above the x-axis represent the actual values

available in the data for the variable. The black line represents the unsmoothed PDP, while

the green line shows the smoothed version. The shaded gray area corresponds to the 95%

confidence intervals, calculated using the standard deviation and adjusted for local data

density. b/w: between. bef.: before. tmin: minimum hourly temperature. rhmean: hourly

average relative humidity. Patmin: minimum daily atmospheric pressure. Patdiff: difference

of mean daily atmospheric pressure. CUMRF: Cumulated Rainfall.

Fig 6: Interpretation plots for the multivariate random forest model of *Ae. albopictus*

abundance. The left plot (Fig 6A) is the variable importance plot (VIP). Rights plots (Fig 6B)

are smoothed partial dependence plots (PDPs) with Confidence Intervals for each selected

variable in the models. The y-axis in PDPs represents the number of *Ae. albopictus* females

caught/trap/24h. The range of values in the x-axis represents the range of values available in

the data for the considered variable. The rugs above the x-axis represent the actual values

available in the data for the variable. The black line represents the unsmoothed PDP, while

the green line shows the smoothed version. The shaded gray area corresponds to the 95%

confidence intervals, calculated using the standard deviation and adjusted for local data

density. b/w: between. bef.: before. tmax: maximum hourly temperature. GDD: Growing

Degree Day. WS: Wind Speed

Presence model

The most contributing variables to presence probability were a combination of real-time and

time-lagged microclimatic and meteorological variables (Fig 5A), with only a minimal

contribution of vegetation variables. The first contributing variable was the local minimum

hourly temperature recorded 24–48 hours before sampling. The probability of presence

showed a slight increase between 11.9°C and 16°C, rose sharply from 16°C to 25°C, and

plateaued beyond 25°C (Fig 5B). Minimum atmospheric pressure during sampling was the

second most influential variable, with a unimodal relationship: presence probability

increased up to 1012.91 HPa before declining. Cumulated rainfall 6 weeks before sampling

emerged as the third key variable, displaying a linear positive association with presence

probability, from 0 mm to 15 mm. The fourth major factor was the mean daily atmospheric

pressure difference between the sampling and previous day. Presence probabilities

decreased for descreasing pressures and were stable for stable or increasing pressures.

Secondary contributors included mean local hourly relative humidity 24–48 hours before

sampling, daily NO_2 concentrations, and vegetation metrics within a 20 m buffer around

traps (average patch size of low vegetation, number of high vegetation patches, and total

length of low vegetation edges). These variables slightly increased presence probability,

whereas the total length of edges of roads in a 20 m radius buffer weakly reduced it. The

model demonstrated strong predictive accuracy (AUC = 0.89; S7A Fig). Cross-validation with

the leave-area-out method produced consistent predictions, except in the Diderot School

area (IMP-DID). In contrast, leave-sampling-session-out cross-validation exhibited greater

variability between sampling areas (S7B Fig).

Abundance model

The most contributing variables to *Ae. albopictus* abundance comprised a combination of

microclimatic, weeks-lagged meteorological, vegetation and other land cover variables. The

first contributing variable was the average size of low-vegetation patches within a 50 m

radius buffer around traps (Fig 6A). Abundance increased markedly for areas between 0 and

120 m², plateaued up to 237 m², and subsequently decreased slightly (Fig 6B). The second

most important variable was the local maximum hourly temperature during sampling. A

strong positive correlation with abundance was observed between 24°C and 38.7°C,

followed by a negative correlation beyond 40.6°C. The third key variable was the total length

of road edges within a 250 m buffer around traps, with a strong negative correlation

between 8230 m and 11 000 m. The fourth contributor was the percentage of low

vegetation within a 250 m buffer, which showed a positive effect between 10% and 23%,

beyond which it had no observable effect. The fifth and sixth contributors were the

cumulated rainfall in the sixth week before sampling and the weekly GDD between the

sampling day and two weeks prior, both of which were positively associated with mosquito

abundance. The average wind speed between the day of sampling and five weeks before had

no effect on abundance between 3.7 m/s and 4.3 m/s, but showed a negative correlation

beyond this range. Finally, the last two contributors were the average patch size of roads

within a 250 m buffer and the maximum hourly relative humidity during the 24 hours

preceding sampling, both of which were slightly negatively correlated with mosquito

abundance.

The abundance model demonstrated good overall accuracy, except for high abundance

cases (>11 mosquitoes, Mean Absolute Error (MAE)=8.00, S8A Fig). Spatial cross-validation

retained overall abundance trends, except in RES-LEM and PRK-AIG, and showed an

underestimation of high abundance and an overestimation of low abundance. Temporal

cross-validation, using leave-sampling-session-out, did not preserve abundance trends but

captured relative differences in *Ae. albopictus* abundance between areas (S8B Fig).

Discussion

This study examined, for the first time in France, the impact of urban vegetation on the

presence and abundance of *Ae. albopictus*, showing a minimal influence on the presence and

a positive association with the abundance. The analytical methodology employed in this

study provides a nuanced evaluation of how different environmental factors contribute to

mosquito presence and abundance. It highlights the existence of non-linear relationships,

offering valuable insights for identifying effective vector control measures.

Influence of urban vegetation and other land cover variables

This study highlights that urban vegetation within a 20 m buffer radius around traps had a

minimal weakly positive influence on the probability of the presence of *Ae. albopictus*.

Regarding the abundance, urban vegetation demonstrated a positive effect and emerged as

one of the four most influential predictors. This positive effect of urban vegetation on the

abundance of *Ae. albopictus* has also been observed in other countries, including Italy [26],

Brazil [88], Singapore [89], and China [33]. In our study, the first vegetation metric

influencing abundance was the average size of low-vegetation patches within a 50 m buffer

radius. These findings are consistent with those of Cianci *et al* (2015) [26], who reported a

positive relationship between vegetation cover within a 50 m radius buffer around traps and

Ae. albopictus eggs abundance in Italy. Manica *et al* (2016) observed that, in Roma (Italy),

vegetation cover within a 20 m radius around traps positively influenced *Ae. albopictus*

female abundance [27]. They hypothesised that the tiger mosquito prefers "small green

islands". Our findings refine this hypothesis by demonstrating that *Ae. albopictus* abundance

significantly increases with the average size of low-vegetation patches, particularly within

the 10 to 120 m² range. Beyond 120 m², the effect stabilises. This nuance provides additional

detail to the assumption proposed by Manica *et al*. This hypothesis, along with the identified

size range, requires further research to confirm its generalisability across different urban

contexts and mosquito populations.

The percentage of low vegetation cover within a 250 m radius around traps moderately

increased abundance. These results are consistent with the study of Sun *et al* (2021), which

demonstrated that, in Singapore, *Ae. albopictus* abundance increased with managed

vegetation within a 300 m radius buffer around traps [89]. However, Manica *et al* (2016)

found a negative relationship between vegetation cover within a 300 m radius buffer and

mosquito abundance [27]. These differences can be attributed to variations in city

architecture and study design. In our study, the percentage of low vegetation cover ranged

from 10% to 23.5%, whereas in Manica *et al*, it ranged from 30% to 55%. With different

ranges of vegetation cover, differing study designs, and contrasting urban environments,

direct comparisons are challenging.

The other two land cover variables, the third and eighth contributors to abundance, are road

metrics within a 250 m radius buffer around traps. The first was the total length of road

edges, which was negatively correlated with abundance. These findings aligned with the

study by Wang *et al* (2023) in Shanghai (China) which reported a negative correlation

between road length and *Ae. albopictus* eggs density [33]. This supports our initial

assumption that roads act as barriers to mosquito movement. This hypothesis was also

consistent with Regilme *et al* (2021), who observed genetic differentiation in *Ae. aegypti*

populations located north and south of a major road in Shanghai [42]. The second road-

related variable was the average size of road patches. Roads may represent unfavorable

habitats for host-seeking *Ae. albopictus* which could explain their limited abundance in areas

dominated by large roads.

Influence of microclimatic and meteorological variables

647   The relationship between atmospheric pressure and the presence of *Ae. albopictus* in the

648   field has, to our knowledge, been investigated for the first time in this study. The minimum

649   atmospheric pressure during the 24-hour sampling period and the difference in daily mean

atmospheric pressure between the sampling day and the previous day were the second and

fourth main variables contributing to presence. Atmospheric pressure can influence insects'

flight [51] and locomotor activity [52], as shown in *Drosophila* and beetles. Haufe (1954)

demonstrated that atmospheric pressure affects the flight activity of *Ae. aegyp*ti, especially

the change in atmospheric pressure after six hours of acclimatisation [53]. A study by Knop

et al (2023) in Switzerland showed that the traffic rate of high-flying insects (number of

insects per hour) had a positive correlation with air pressure [90]. Our results suggest an

effect of the atmospheric pressure metrics on the activity and behaviour of the tiger

mosquito. Further studies under controlled laboratory and field conditions are needed to

investigate these findings in more detail.

Local hourly temperatures emerged as the primary determinant of *Ae. albopictus* presence

and the second most influential predictor of its abundance. Evans *et al* (2019) found an

exponential relationship between the minimum temperature during the 7 days preceding

sampling and the abundance of females, as well as an unimodal relationship between the

maximum temperature and the abundance of females, with an optimal daily maximum

temperature around 35°C [32]. Our results suggest that increasing daily maximum

temperature increases the activity of *Ae. albopictus* up to 38°C.

The third variable influencing the presence of *Ae. albopictus* was the weekly cumulated

rainfall recorded six weeks before sampling, which also ranked as the fifth most important

predictor of abundance. While other studies have identified a positive relationship between

Ae. albopictus dynamics and rainfall accumulated over several weeks before sampling, none

have specifically examined a six-week delay. Roiz *et al* (2015) observed in Montpellier that

strong rainfall, particularly that accumulated each week over the four weeks before

sampling, positively influenced abundance [31]. Similarly, Manica *et al* (2016) reported a

positive relationship between the cumulated rainfall one month before sampling and mosquito abundance [27]. The positive effect of rainfall in our study may be explained by the filling of larval habitats, benefiting previous mosquito generations that influenced the sampled population. This hypothesis requires confirmation through additional temporal replicates to validate the observed trend.

The sixth most influential variable of abundance was the weekly GDD during the two weeks preceding sampling. Based on minimum, maximum, and average temperatures, the GDD is widely used as a key predictor for many insect species [91]. For *Ae. albopictus*, the GDD is calculated using a baseline temperature of 11°C, promoting mosquito development when temperatures exceed this value [50]. This baseline corresponds to the average annual temperature required for the spread of *Ae. albopictus* [92].

One of the last variables influencing abundance is the mean daily wind speed during the five weeks preceding sampling. Le Goff *et al* (2024) observed a similar negative relationship between monthly average daily wind speed and *Ae. albopictus* eggs abundance on the Réunion Island [93]. Adeleke *et al* (2022) reported that in regions where the tiger mosquito is present, wind speeds exceeding 4 m·s^{-1} reduce the likelihood of occurrence [94]. The threshold identified in our study (4.3 m·s^{-1}) can be considered comparable to the findings of Adeleke *et al*, although direct comparisons between the studies are challenging due to differences in spatial and temporal scales.

Limitations, perspectives and conclusion

In order to increase the resilience of Urban Green Infrastructures (UGIs) [95] and to adapt the UGIs management to promote biodiversity [96], since 2017, studies have highlighted the need to consider UGIs as socio-ecological ecosystems. This reflects the complexity of urban

greening, which must account for land cover, socio-demographic variables, and urban architecture, as demonstrated in the systematic review by Menconi *et al* (2021) [97]. The challenge of studying the relationship between urban greening and the risk of vector-borne diseases is one of the key conclusions of the scoping review by Mercat *et al* (2025) [24]. The difficulty stems from the need to integrate multiple variables essential to the analysis of this socio-ecosystem, the time required to observe its effects [28], and the challenge of assessing vector-borne disease risk using standardised protocols.

Due to limited financial and human resources, this study was limited to specific sampling areas within vegetated environments representative of the city's UGS and non-vegetated environments. A randomised sampling approach would have better captured the diversity of land cover categories. The predicted abundance of *Ae. albopictus* in the RES-SOUL residential area was therefore certainly underestimated. While the abundance model effectively captures temporal variation, it does not account well for spatial variation. This suggests that key factors such as environmental carrying capacity, defined as the 'equilibrium density of larvae in a system' [98], and the availability of resting sites may have been overlooked (35). This is also related to the complexity of describing urban vegetation using remote sensing [24]. In our study, we employed a fine-scale characterisation of urban vegetation generated through deep learning and photo interpretation, alongside a surface modelling metric derived from lidar data. However, this approach does not capture the diversity of plant species or genera, which can vary markedly between the Botanical Garden, the Park of Aiguelongue, and residential areas. Finally, as described in the Materials and Methods, no municipal vector control interventions were observed in the sampling areas. Nevertheless, potential actions by residents in houses close to the sampled sites could not be assessed.

The objectives of this study were to explore the relationships between urban vegetation and

other environmental factors with the presence and abundance of *Ae. albopictus*, to generate

hypotheses using an exploratory approach. The realisation of a predictive model is another

objective, which requires a better understanding of the ecological and biological

mechanisms [99]. The results of the study are based on only one sampling year. To draw

more robust conclusions, additional years of sampling would be needed to assess inter-

annual variation. In addition, this study has focused solely on one aspect of the vector risk,

the presence and abundance of the vector, without considering its capacity to transmit

pathogens, or human exposure and vulnerability. The vector's capacity depends on the

abundance of females, the daily survival rate, and the extrinsic incubation period of

pathogens [100]. It would be relevant to investigate the effect of urban vegetation on these

elements. Together with these investigations, a socio-ecological approach is essential to gain

a comprehensive understanding of vector-borne disease risks in urban green spaces.

Developing social surveys to assess citizens' perceptions, alongside employing co-

construction methods [101] with policymakers, urban designers, and ecologists, could help

formulate effective recommendations to mitigate this risk. One example could be the

development of green wind corridors, as implemented in Stuttgart [102] and Lisbon [103],

which can reduce urban heat islands and increase wind speed, limiting mosquito abundance.

Another possible implication for urban planning would be to promote the establishment of

small vegetation patches located at least 50 m away from residential areas. These patches

could be separated by permeable soil or arranged as discontinuous corridors composed of

diverse vegetation types (e.g., grass, scattered trees). Vegetation management and good

gardening practices could further limit mosquito proliferation, for instance by emptying

potential breeding sites at least once a week, avoiding plants that naturally serve as larval

habitats (e.g., bromeliads [104]), and reducing features such as bamboo nodes [29].

Maintaining less dense shrubs could help reduce the number of available resting sites [35].

Such practices would allow urban green spaces to retain their ecological and recreational

functions while reducing mosquito abundance. However, these suggestions are currently

limited to the local context of Montpellier and would require additional sampling, including

in other locations, to ensure more robust and generalizable recommendations.

Urban greening appears to be a promising solution to address the environmental and health

challenges associated with increasing urbanisation. It offers numerous health and

environmental benefits, but can also contribute to an increased risk of vector-borne

diseases. This study, the first in France to investigate the effect of urban vegetation on *Ae.*

albopictus, provides a nuanced assessment of how urban vegetation influences both the

presence and abundance of mosquitoes. Our results indicate that urban vegetation

moderately increases abundance, but has no effect on presence. Urban greening is a

dynamic and complex phenomenon that requires a social-ecological approach in order to be

studied. Further research is needed to better elucidate the role of urban vegetation in the

ecology of the tiger mosquito and its influence on different aspects of risk.

Acknowledgements

We would also like to thank Gilbert Le Goff, Nil Rahola, Christophe Paupy for the help to the

design sampling and Mathilde Mercat, Coralie Grail and all the other field and laboratory

staff for their strong commitment to the V2MOC project. We would like to thank the

managers and staff of the Botanical Garden who facilitated our access, the managers of the

green spaces of the city of Montpellier for access to the Parc de l'Aiguelongue, as well as the

managers of the Insituts Diderot, Bouisson Bertrand, the Hotel Acapulco and the Saint

Charles site of the University of Montpellier III, as well as the individuals who gave us access

to their gardens. We thank all the persons who worked on the open access databases used

in this work.

References

1.   Manzoor Hussain, Imitiaz I. Urbanization concepts, dimension and factors. Int J Recent Sci Res.
2018;

2.   Breuste J, Niemelä J, Snep RPH. Applying landscape ecological principles in urban
environments. Landsc Ecol. 2008 Dec;23(10):1139–42.

3.   Zhang Z, Zhao M, Zhang Y, Feng Y. How does urbanization affect public health? New evidence
from 175 countries worldwide. Front Public Health. 2023 Jan 6;10:1096964.

4.   Xu X, Xie Y, Qi K, Luo Z, Wang X. Detecting the response of bird communities and biodiversity
to habitat loss and fragmentation due to urbanization. Sci Total Environ. 2018 May;624:1561–
76.

5.   Qi F, Abu-Rumman A, Al Shraah A, Muda I, Huerta-Soto R, Hai Yen TT, et al. Moving a step
closer towards environmental sustainability in Asian countries: focusing on real income,
urbanization, transport infrastructure, and research and development. Econ Res-Ekon
Istraživanja. 2023 Jul 10;36(2):2111317.

6.   Moore M, Gould P, Keary BS. Global urbanization and impact on health. Int J Hyg Environ
Health. 2003;206(4–5):269–78.

7.   Ventriglio A, Torales J, Castaldelli-Maia JM, De Berardis D, Bhugra D. Urbanization and
emerging mental health issues. CNS Spectr. 2021 Feb;26(1):43–50.

8.   Almusaed A. The urban heat island phenomenon upon urban components. In: biophilic and
bioclimatic architecture [Internet]. London: Springer London; 2011 [cited 2024 Sep 18]. p.
139–50. Available from: http://link.springer.com/10.1007/978-1-84996-534-7_10

9.   Heaviside C, Macintyre H, Vardoulakis S. The urban heat island: implications for health in a
changing environment. Curr Environ Health Rep. 2017 Sep;4(3):296–305.

10. United Nations (UN). Goal 11: Make cities and human settlements inclusive, safe, resilient and
sustainable. New York: UN. [Internet]. [cited 2024 Nov 19]. Available from:
https://sdgs.un.org/goals/goal11

11.   Unit B. Convention on Biological Diversity [Internet]. Secretariat of the Convention on
802   Biological Diversity; 2022 [cited 2024 Oct 2]. Available from:
803   https://www.cbd.int/gbf/targets/12

804   12.   The European Green Deal: A growth strategy that protects the climate [Internet]. 2022 [cited
805   2024 Oct 2]. Available from: https://ec.europa.eu/stories/european-green-deal

806   13.   Chang J, Qu Z, Xu R, Pan K, Xu B, Min Y, et al. Assessing the ecosystem services provided by
807   urban green spaces along urban center-edge gradients. Sci Rep. 2017 Sep 11;7(1):11226.

808   14.   Liu HL, Shen YS. The impact of green space changes on air pollution and microclimates: a case
809   study of the Taipei Metropolitan Area. Sustainability. 2014 Dec 3;6(12):8827–55.

810   15.   Gunawardena KR, Wells MJ, Kershaw T. Utilising green and blue spaces to mitigate urban heat
811   island intensity. Sci Total Environ. 2017 Apr;584–585:1040–55.

812   16.   Callaghan A, McCombe G, Harrold A, McMeel C, Mills G, Moore-Cherry N, et al. The impact of
813   green spaces on mental health in urban settings: a scoping review. J Ment Health. 2021 Mar
814   4;30(2):179–93.

815   17.   Zhang Y, Meng W, Yun H, Xu W, Hu B, He M, et al. Is urban green space a carbon sink or
816   source? - A case study of China based on LCA method. Environ Impact Assess Rev. 2022
817   May;94:106766.

818   18.   Threlfall CG, Mata L, Mackie JA, Hahs AK, Stork NE, Williams NSG, et al. Increasing biodiversity
819   in urban green spaces through simple vegetation interventions. Beggs J, editor. J Appl Ecol.
820   2017 Dec;54(6):1874–83.

821   19.   Cariñanos P, Casares-Porcel M. Urban green zones and related pollen allergy: A review. Some
822   guidelines for designing spaces with low allergy impact. Landsc Urban Plan. 2011
823   Jun;101(3):205–14.

824   20.   Ciceoi R, Gutue C, Gutue M, Roşca I. Current status of pests associated with urban vegetation
825   in bucharest area. 2017.

826   21.   Lõhmus M, Balbus J. Making green infrastructure healthier infrastructure. Infect Ecol
827   Epidemiol. 2015 Jan;5(1):30082.

828   22.   Fournet F, Simard F, Fontenille D. Green cities and vector-borne diseases: emerging concerns
829   and opportunities. Eurosurveillance [Internet]. 2024 Mar 7 [cited 2024 Aug 19];29(10).
830   Available from: https://www.eurosurveillance.org/content/10.2807/1560-
831   7917.ES.2024.29.10.2300548

832   23.   Heylen D, Lasters R, Adriaensen F, Fonville M, Sprong H, Matthysen E. Ticks and tick-borne
833   diseases in the city: Role of landscape connectivity and green space characteristics in a
834   metropolitan area. Sci Total Environ. 2019 Jun;670:941–9.

835   24.   Mercat M, Bartholomée C, Fournet F, Alcover Amengual M, Bourquia M, Bouhsira E, et al.
836   Green cities and the risk for vector-borne disease transmission: a scoping review. 2025.
837   bioRxiv 2025.04.01.646559; doi: https://doi.org/10.1101/2025.04.01.646559

25. Krol L, Langezaal M, Budidarma L, Wassenaar D, Didaskalou EA, Trimbos K, et al. Distribution
of *Culex pipiens* life stages across urban green and grey spaces in Leiden, The Netherlands.
Parasit Vectors. 2024 Jan 29;17(1):37.

26. Cianci D, Hartemink N, Zeimes CB, Vanwambeke SO, Ienco A, Caputo B. High resolution spatial
analysis of habitat preference of *Aedes albopictus* (Diptera: Culicidae) in an urban
environment. J Med Entomol. 2015 May 1;52(3):329–35.

27. Manica M, Filipponi F, D'Alessandro A, Screti A, Neteler M, Rosà R, et al. Spatial and temporal
hot spots of *Aedes albopictus* abundance inside and outside a South European Metropolitan
Area. McCall PJ, editor. PLoS Negl Trop Dis. 2016 Jun 22;10(6):e0004758.

28. Mata L, Hahs AK, Palma E, Backstrom A, Johnston N, King T, et al. Large positive ecological
changes of small urban greening actions. Ecol Solut Evid. 2023 Jul;4(3):e12259.

29. Paupy C, Delatte H, Bagny L, Corbel V, Fontenille D. *Aedes albopictus,* an arbovirus vector:
From the darkness to the light. Microbes Infect. 2009 Dec;11(14–15):1177–85.

30. Bonizzoni M, Gasperi G, Chen X, James AA. The invasive mosquito species *Aedes albopictus*:
current knowledge and future perspectives. Trends Parasitol. 2013 Sep;29(9):460–8.

31. Roiz D, Boussès P, Simard F, Paupy C, Fontenille D. Autochthonous Chikungunya transmission
and extreme climate events in Southern France. Carvalho MS, editor. PLoS Negl Trop Dis. 2015
Jun 16;9(6):e0003854.

32. Evans MV, Hintz CW, Jones L, Shiau J, Solano N, Drake JM, et al. Microclimate and larval
habitat density predict adult *Aedes albopictus* abundance in urban areas. Am J Trop Med Hyg.
2019 Aug 7;101(2):362–70.

33. Wang F, Zhu Y, Zhang H, Fan J, Leng P, Zhou J, et al. Spatial and temporal analyses of the
influences of meteorological and environmental factors on *Aedes albopictus* (Diptera:
Culicidae) population dynamics during the peak abundance period at a city scale. Acta Trop.
2023 Sep;245:106964.

34. Murdock CC, Evans MV, McClanahan TD, Miazgowicz KL, Tesla B. Fine-scale variation in
microclimate across an urban landscape shapes variation in mosquito population dynamics
and the potential of Aedes albopictus to transmit arboviral disease. Scarpino SV, editor. PLoS
Negl Trop Dis. 2017 May 30;11(5):e0005640.

35. Samson DM, Qualls WA, Roque D, Naranjo DP, Alimi T, Arheart KL, et al. Resting and Energy
Reserves of *Aedes albopictus* collected in common landscaping vegetation in St. Augustine,
Florida. J Am Mosq Control Assoc. 2013 Sep;29(3):231–6.

36. Chabasse D. *Aedes albopictus,* le "moustique tigre", vecteur de la dengue, du chikungunya et
du Zika, s'implante durablement en France métropolitaine. Rev Francoph Lab. 2023
Nov;2023(556):51–9.

37. Climate Data. Climat Montpellier: Pluviométrie et Température moyenne Montpellier,
diagramme ombrothermique pour Montpellier [Internet]. 2020 [cited 2024 Oct 2]. Available
from: https://fr.climate-data.org/europe/france/languedoc-roussillon/montpellier-1126/

38. Kermap. Nos villes vertes : la carte de France de la nature en ville [Internet]. 2023 [cited 2024
Oct 2]. Available from: https://www.nosvillesvertes.fr/Accueil

39.   Russell RC, Webb CE, Williams CR, Ritchie SA. Mark–release–recapture study to measure
dispersal of the mosquito *Aedes aegypti* in Cairns, Queensland, Australia. Med Vet Entomol.
2005 Dec;19(4):451–7.

40.   Marini F, Caputo B, Pombi M, Tarsitani G, Della Torre A. Study of *Aedes albopictus* dispersal in
Rome, Italy, using sticky traps in mark-release-recapture experiments. Med Vet Entomol. 2010
Dec;24(4):361–8.

41.   Hemme RR, Thomas CL, Chadee DD, Severson DW. Influence of urban landscapes on
population dynamics in a short-distance migrant mosquito: evidence for the dengue vector
Aedes aegypti. Gubler DJ, editor. PLoS Negl Trop Dis. 2010 Mar 16;4(3):e634.

42.   Regilme MAF, Carvajal TM, Honnen A, Amalin DM, Watanabe K. The influence of roads on the
fine-scale population genetic structure of the dengue vector *Aedes aegypti* (Linnaeus).
Bonizzoni M, editor. PLoS Negl Trop Dis. 2021 Feb 26;15(2):e0009139.

43.   L'Ambert G, Gendrot M, Briolant S, Nguyen A, Pages S, Bosio L, et al. Analysis of trapped
mosquito excreta as a noninvasive method to reveal biodiversity and arbovirus circulation.
Mol Ecol Resour. 2023 Feb;23(2):410–23.

44.   Le Goff G, Damiens D, Payet L, Ruttee AH, Jean F, Lebon C, et al. Enhancement of the BG-
sentinel trap with varying number of mice for field sampling of male and female *Aedes
albopictus* mosquitoes. Parasit Vectors. 2016 Dec;9(1):514.

45.   Gunay F, Picard M, Robert V. MosKeyTool, an interactive identification key for mosquitoes of
Euro-Mediterranean. [Internet]. 2017. Available from: www.medilabsecure.com/moskeytool.

46.   Bartholomée C, Taconet P, Mercat M, Sutter C, Grail C, Garcia-Marin C, et al. Monthly
mosquito sampling in Montpellier (Occitanie), France, 2023 [Internet]. IRD - Institute of
Research   for   Development;   2024   [cited   2025   Mar   6].   Available   from:
https://www.gbif.org/dataset/8e52f35a-2522-4865-8361-5c249310a7cf   /   doi:
https://doi.org/10.15468/4qafbu

47.   Moiroux N. 'Chapitre 5◻: Modélisation spatio-temporelle du contact homme-vecteur'. In:
Modélisation du risque d'exposition aux moustiques vecteurs de Plasmodium spp dans un
contexte de lutte anti-vectorielle" [Internet]. Université Montpellier-II Sciences et Techniques
du Languedoc; 2012. p. 76. Available from: https://theses.hal.science/tel-00812118

48.   Bartholomée C, Moiroux N. Conceptual model of *Aedes albopictus* in urban environments
[Internet].   Zenodo;   2025   [cited   2025   Mar   6].   Available   from:
https://zenodo.org/doi/10.5281/zenodo.14981760

49.   Arduino M de B, Mucci LF, Santos LM dos, Soares MF de S. Importance of microenvironment
to arbovirus vector distribution in an urban area, São Paulo, Brazil. Rev Soc Bras Med Trop.
2020;53:e20190504.

50.   Neteler M, Roiz D, Rocchini D, Castellani C, Rizzoli A. Terra and Aqua satellites track tiger
mosquito invasion: modelling the potential distribution of *Aedes albopictus* in north-eastern
Italy. Int J Health Geogr. 2011;10(1):49.

51.   Chadwick LE, Williams CM. The effects of atmospheric pressure and composition on the flight
of drosophila. Biol Bull. 1949 Oct;97(2):115–37.

52.   Pellegrino AC, Peñaflor MFGV, Nardi C, Bezner-Kerr W, Guglielmo CG, Bento JMS, et al.
919   Weather forecasting by insects: modified sexual behaviour in response to atmospheric
920   pressure changes. Lazzari CR, editor. PLoS ONE. 2013 Oct 2;8(10):e75004.

921   53.   Haufe WO. The effects of atmospheric pressure on the flight responses of *Aëdes aegypti* (L.).
922   Bull Entomol Res. 1954 Sep;45(3):507–26.

923   54.   Duque L, Steffan-Dewenter I. Air pollution: a threat to insect pollination. Front Ecol Environ.
924   2024 Apr;22(3):e2701.

925   55.   Ryalls JMW, Langford B, Mullinger NJ, Bromfield LM, Nemitz E, Pfrang C, et al. Anthropogenic
926   air pollutants reduce insect-mediated pollination services. Environ Pollut. 2022
927   Mar;297:118847.

928   56.   Ryalls JMW, Bishop J, Mofikoya AO, Bromfield LM, Nakagawa S, Girling RD. Air pollution
929   disproportionately impairs beneficial invertebrates: a meta-analysis. Nat Commun. 2024 Jul
930   11;15(1):5447.

931   57.   Zellweger RM, Cano J, Mangeas M, Taglioni F, Mercier A, Despinoy M, et al. Socioeconomic
932   and environmental determinants of dengue transmission in an urban setting: An ecological
933   study in Nouméa, New Caledonia. Messer WB, editor. PLoS Negl Trop Dis. 2017 Apr
934   3;11(4):e0005471.

935   58.   Little E, Biehler D, Leisnham PT, Jordan R, Wilson S, LaDeau SL. Socio-Ecological mechanisms
936   supporting high densities of *Aedes albopictus* (Diptera: Culicidae) in Baltimore, MD. J Med
937   Entomol. 2017 Sep 1;54(5):1183–92.

938   59.   Rodrigues M de M, Marques GRAM, Serpa LLN, Arduino M de B, Voltolini JC, Barbosa GL, et al.
939   Density of *Aedes aegypti* and *Aedes albopictus* and its association with number of residents
940   and meteorological variables in the home environment of dengue endemic area, São Paulo,
941   Brazil. Parasit Vectors. 2015 Dec;8(1):115.

942   60.   Moiroux N, Djènontin A, Bio-Bangana AS, Chandre F, Corbel V, Guis H. Spatio-temporal
943   analysis of abundances of three malaria vector species in southern Benin using zero-truncated
944   models. Parasit Vectors. 2014;7(1):103.

945   61.   Taconet P, Porciani A, Soma DD, Mouline K, Simard F, Koffi AA, et al. Data-driven and
946   interpretable machine-learning modeling to explore the fine-scale environmental
947   determinants of malaria vectors biting rates in rural Burkina Faso. Parasit Vectors. 2021 Jun
948   29;14(1):345.

949   62.   Breiman L. Random Forest. Mach Learn. 2001;45(1):5–32.

950   63.   Nakagawa S, Schielzeth H. A general and simple method for obtaining R^2 from generalized
951   linear mixed-effects models. O'Hara RB, editor. Methods Ecol Evol. 2013 Feb;4(2):133–42.

952   64.   Curriero FC, Shone SM, Glass GE. Cross Correlation Maps: A tool for visualizing and modeling
953   time lagged associations. Vector-Borne Zoonotic Dis. 2005 Sep;5(3):267–75.

954   65.   Zuur AF, Ieno EN, Elphick CS. A protocol for data exploration to avoid common statistical
955   problems: Data exploration. Methods Ecol Evol. 2010 Mar;1(1):3–14.

66. Speiser JL, Miller ME, Tooze J, Ip E. A comparison of random forest variable selection methods
for classification prediction modeling. Expert Syst Appl. 2019 Nov;134:93–101.

67. Meyer H, Reudenbach C, Hengl T, Katurji M, Nauss T. Improving performance of spatio-
temporal machine learning models using forward feature selection and target-oriented
validation. Environ Model Softw. 2018 Mar;101:1–9.

68. Bradley AP. The use of the area under the ROC curve in the evaluation of machine learning
algorithms. Pattern Recognit. 1997 Jul;30(7):1145–59.

69. Friedman JH. Greedy function approximation: A gradient boosting machine. Ann Stat
[Internet]. 2001 Oct 1 [cited 2024 Aug 29];29(5). Available from:
https://projecteuclid.org/journals/annals-of-statistics/volume-29/issue-5/Greedy-function-
approximation-A-gradient-boosting-machine/10.1214/aos/1013203451.full

70. R Core Team. R: A language and environment for statistical. R Foundation for Statistical
Computing, Vienna, Austria. [Internet]. 2021. Available from: https://www.R-project.org/.

71. Pebesma E. Simple Features for R: Standardized Support for Spatial Vector Data. R J.
2018;10(1):439.

72. Hijmans RJ. raster: Geographic Data Analysis and Modeling [Internet]. 2010 [cited 2024 Aug
29]. p. 3.6-26. Available from: https://CRAN.R-project.org/package=raster

73. Ross N. fasterize: Fast Polygon to Raster Conversion [Internet]. 2018 [cited 2024 Aug 29]. p.
1.0.5. Available from: https://CRAN.R-project.org/package=fasterize

74. Hesselbarth MHK, Sciaini M, Nowosad J, Hanss S. landscapemetrics: Landscape Metrics for
Categorical Map Patterns [Internet]. 2018 [cited 2024 Aug 29]. p. 2.1.4. Available from:
https://CRAN.R-project.org/package=landscapemetrics

75. Brooks M E, Kristensen K, Benthem K J ,van, Magnusson A, Berg C W, Nielsen A, et al.
glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear
Mixed Modeling. R J. 2017;9(2):378.

76. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. performance: An R Package for
Assessment, Comparison and Testing of Statistical Models. J Open Source Softw. 2021 Apr
21;6(60):3139.

77. Makowski D, Ben-Shachar M, Patil I, Lüdecke D. Methods and Algorithms for Correlation
Analysis in R. J Open Source Softw. 2020 Jul 16;5(51):2306.

78. Kuhn M. Building Predictive Models in R Using the caret Package. J Stat Softw [Internet]. 2008
[cited 2024 Jun 6];28(5). Available from: http://www.jstatsoft.org/v28/i05/

79. Wright MN, Ziegler A. ranger: A Fast Implementation of random forests for high dimensional
Data in C++ and R. J Stat Softw [Internet]. 2017 [cited 2024 Jun 6];77(1). Available from:
http://www.jstatsoft.org/v77/i01/

80. Meyer H, Ludwig M, Milà C, Linnenbrink J, Schumacher F. The CAST package for training and
assessment of spatial prediction models in R [Internet]. arXiv; 2024 [cited 2025 Jan 8].
Available from: https://arxiv.org/abs/2404.06978

81.   Molnar C. iml: An R package for Interpretable Machine Learning. J Open Source Softw. 2018
Jun 27;3(26):786.

82.   Greenwell B M. pdp: An R Package for Constructing Partial Dependence Plots. R J.
2017;9(1):421.

83.   Yan Y. MLmetrics: Machine Learning Evaluation Metrics [Internet]. 2015 [cited 2025 Jan 8]. p.
1.1.3. Available from: https://CRAN.R-project.org/package=MLmetrics

84.   Saito T, Rehmsmeier M. Precrec: fast and accurate precision–recall and ROC curve calculations
in R. Wren J, editor. Bioinformatics. 2017 Jan 1;33(1):145–7.

85.   Pedersen TL. patchwork: The Composer of Plots [Internet]. 2019 [cited 2025 Jan 8]. p. 1.3.0.
Available from: https://CRAN.R-project.org/package=patchwork

86.   Benito B. BlasBenito/spatialRF: spatialRF: Easy Spatial Regression with Random Forest
[Internet]. Zenodo; 2021 [cited 2025 Jan 8]. Available from:
https://zenodo.org/record/4745208

87.   Taconet P, Bartholomée C. R scripts to reproduce the analysis of the publication 'Investigating
the role of urban vegetation alongside other environmental variables in shaping Aedes
albopictus presence and abundance in Montpellier, France' [Internet]. [cited 2025 Mar 11].
Available   from:
https://archive.softwareheritage.org/swh:1:dir:a1c83c4be076b5645d1834265219d4f2e6e5b5
44;origin=https://github.com/ptaconet/modeling_vector_mtp;visit=swh:1:snp:741ed0f1af295
a2a2ef477b310cda3f0faa3f1a0;anchor=swh:1:rev:11de6e23243904d7fa7a21d9d6ade29486d
06349

88.   Honório NA, Castro MG, Barros FSM de, Magalhães M de AFM, Sabroza PC. The spatial
distribution of *Aedes aegypti* and *Aedes albopictus* in a transition zone, Rio de Janeiro, Brazil.
Cad Saúde Pública. 2009 Jun;25(6):1203–14.

89.   Sun H, Dickens BL, Richards D, Ong J, Rajarethinam J, Hassim MEE, et al. Spatio-temporal
analysis of the main dengue vector populations in Singapore. Parasit Vectors. 2021
Dec;14(1):41.

90.   Knop E, Grimm ML, Korner-Nievergelt F, Schmid B, Liechti F. Patterns of high-flying insect
abundance are shaped by landscape type and abiotic conditions. Sci Rep. 2023 Sep
13;13(1):15114.

91.   Keefe HE, Kharouba HM. Growing degree-days do not explain moth species' distributions at
broad scales. Ecosphere. 2024 Jul;15(7):e4885.

92.   Reinhold J, Lazzari C, Lahondère C. Effects of the environmental temperature on *Aedes aegypti*
and *Aedes albopictus* Mosquitoes: A Review. Insects. 2018 Nov 6;9(4):158.

93.   Le Goff G, Damiens D, Ruttee AH, Jean F, Payet L, Lebon C, et al. Spatial and temporal
characterization of *Aedes albopictus* oviposition activity in candidate urban settings for sterile
insect technique testing in La Reunion Island. Infect Dis Poverty. 2024 Oct 25;13(1):78.

94.   Adeleke ED, Shittu RA, Beierkuhnlein C, Thomas SM. High wind speed prevents the
establishment of the disease vector mosquito *Aedes albopictus* in its climatic niche in Europe.
Front Environ Sci. 2022 Jun 8;10:846243.

95.   Afriyanie D, Akbar R, Suroso DSA. Socio-ecological resilience for urban green space allocation.
1035   IOP Conf Ser Earth Environ Sci. 2018 Apr;145:012120.

1036   96.   Paudel S, States SL. Urban green spaces and sustainability: Exploring the ecosystem services
1037   and disservices of grassy lawns versus floral meadows. Urban For Urban Green. 2023
1038   Jun;84:127932.

1039   97.   Menconi ME, Palazzoni L, Grohmann D. Core themes for an urban green systems thinker: A
1040   review of complexity management in provisioning cultural ecosystem services. Urban For
1041   Urban Green. 2021 Nov;65:127355.

1042   98.   Smith DL, Perkins TA, Tusting LS, Scott TW, Lindsay SW. Mosquito Population Regulation and
1043   Larval Source Management in Heterogeneous Environments. Vasilakis N, editor. PLoS ONE.
1044   2013 Aug 7;8(8):e71247.

1045   99.   Tredennick AT, Hooker G, Ellner SP, Adler PB. A practical guide to selecting models for
1046   exploration, inference, and prediction in ecology. Ecology. 2021 Jun;102(6):e03336.

1047   100.   Tran A, Biteau-Coroller F, Guis H. Modelisation des maladies vectorielles. 2005;

1048   101.   Foudriat M. La co-construction [Internet]. 2éd. Presses de l'EHESP; 2019 [cited 2025 Feb 27].
1049   Available from: http://www.cairn.info/la-co-construction--9782810908257.htm

1050   102.   Kapp R. City of Stuttgart and Green and Blue Space Adaptation for Urban Areas and Eco Towns
1051   (GRaBS) [Internet]. 2022 [cited 2024 Oct 11]. Available from: https://climate-
1052   adapt.eea.europa.eu/fr/metadata/case-studies/stuttgart-combating-the-heat-island-effect-
1053   and-poor-air-quality-with-green-ventilation-corridors

1054   103.   Conseil des Architectes d'Europe. Corridor vert de Lisbonne [Internet]. [cited 2024 Oct 11].
1055   Available   from: https://www.ace-cae.eu/fr/activites/projets-europeens/projets-
1056   termines/connectarch/20-projets-architecturaux-contre-le-changement-climatique/corridor-
1057   vert-de-lisbonne/

1058   104.   Wilke ABB, Vasquez C, Mauriello PJ, Beier JC. Ornamental bromeliads of Miami-Dade County,
1059   Florida are important breeding sites for Aedes aegypti (Diptera: Culicidae). Parasit Vectors.
1060   2018 Dec;11(1):283.

1061

Supporting informations

1063

1064   **S1 Table: Table with sampling areas characterization.**

1065

1066   **S2 Table: Summary of the different variables used, with their processing, acquisition**

1067   **methods or source database and temporal/spatial scale.**

1068

1069 **S3 Table: Description of the landscape metrics for each landscape class for each trap, by**

1070 **buffer.**

1071

1072 **S4 Table: Mosquito density/trap/24h by species and sex in the nine different sampling**

1073 **areas.** The mosquito density represents the marginal mean of the number of mosquitoes

1074 caught per trap per 24 hours, calculated based on the sampling environment, with the

1075 standard error specified (SE).

1076

1077 **S5 Table: Variables Selected After Bivariate Analyses, with Justifications for Exclusion from**

1078 **Multivariate Analysis** (Correlation, VIF, Recursive Feature Elimination)

1079

1080 **S1 Fig: BG-PRO trap on the field and pictures of studied areas.** S1A Fig represents a BG-Pro

1081 on the field, S1B Fig the Park of Aiguelongue, S1C the Botanical Garden, S1D The Bouisson

1082 Bertrand Institute and 1E the Acapulco Hostel.

1083

1084 **S2 Fig: Analysis workflow: from the data acquisition to the multivariate analysis**

1085

1086 **S3 Fig: Time series of weekly cumulated precipitations (1A), mean (TMEAN), minimum**

1087 **(TMIN) and maximum (TMAX) temperature, (1B), growing degree-days (1B), mean relative**

1088 **humidity (1C), mean wind speed (1D) and mean (Patmean), minimum (Patmin) and**

1089 **maximum atmospheric pressure (Patmax), during the studing period.** Vertical dotted red

1090 lines indicate monthly mosquito sampling sessions.

1091

1092 **S4 Fig: Number of *Ae. albopictus* per trap per 24h during the study period, by sampling**

1093 **environment.** Boxes represent medians and interquartile ranges, whiskers indicate minimum

1094 and maximum values, and points show the abundance for each sampling session and day, for

1095 each trap, within the three environments.

1096

S5 Fig: Cross correlation maps showing the lagged effect of air-quality variables on *Ae.*

***albopictus* presence or abundance**. Lagged air-quality variables include daily concentration

of nitrogen monoxide (NO), of nitrogen dioxide (NO2), of particulate matter (PM10, PM2.5)

and of ozone (O3). Time lags are expressed in weeks prior to sampling. The adjusted

marginal R^2 reflects the variance explained by the explanatory variable, adjusted for

correlation direction. Red-bordered squares highlight the time lag with the highest marginal

R^2, with the interval indicated in the top left corner. Black-bordered squares denote marginal

R^2 values close to the highest (within 10%). Gray squares represent correlations with p-

values > 0.2.

1106

S6 Fig: Bivariate relationships between *Ae. albopictus* presence or abundance and the

sampling environment and the socio-demographic variables. The adjusted marginal R^2

reflects the variance explained by the explanatory variable, adjusted for correlation

direction. Boxes are colored if the p-value was < 0.2 (No asterisk: p-value $\in$ [0.05; 0.2], *: p-

value $\in$ [0.01, 0.05], **: p-value $\in$ [0.001; 0.01]; ***: p-value $\in$ [0; 0.001]). Box color

depends on the direction of the relationship (blue: negative, red: positive). Color intensity

varies according to the marginal R^2 value.

1114

S7 Fig: Evaluation of presence model. S7A Fig is the Receiver Operating Characteristics (ROC) Curve for the presence model. It illustrates the model's sensitivity (its ability to predict the presence of *Ae. albopictus*) against 1 - Specificity (its ability to detect the absence of *Ae. albopictus*). The Area Under the Curve (AUC) represents the model's ability to discriminate between the presence and absence of the mosquito, with values ranging from 0 to 1. The closer the AUC is to 1, the better the model's predictive performance. S7A Fig shows that the model demonstrated good predictive accuracy, with an AUC of 0.89. S7B Fig compares the observed (in green) versus predicted presence probabilities for each out-of-sample leave-area (yellow) and for each out-of-sample leave-sampling-session (in blue) . The y-axis represents the probability of *Ae. albopictus* presence, while the x-axis corresponds to the sampling session. Figure 7B indicates that the model accurately predicted the spatiotemporal trends of presence/absence, although it tended to overestimate presence.

S8 Fig: Evaluation of abundance model. S8A Fig is a violin plot showing the distribution of residuals for the abundance model. The black dots indicate the median values. The Mean Absolute Error (MAE) and the number of observations (n) are displayed in a square above the plot. This figure demonstrates that the model tends to overestimate low counts of *Aedes albopictus* (fewer than three caught in 24 hours per trap), accurately estimates counts between 4 and 20, and underestimates higher counts (more than 20). S8B Fig compares the observed versus predicted numbers of *Ae. albopictus* caught per site per sampling session (in green), versus predicted numbers for each out-of-sample leave-area (yellow) and for each out-of-sample leave-sampling-session (in blue). The y-axis represents the mean number of d *Ae. albopictus* females caught per trap per area per sampling session, while the x-axis represents the sampling session. This figure confirms the observations made in S8A Fig.

1139

1140   **S1 Text: Detailed description of bivariate and multivariate analysis.**

A. Location of Montpellier

C. Garden of residential area

D. Botanical Garden

E. One impervous area

B. Studied areas

Sampling environments

- Urban parks (5 traps)
- Highly impervious center (4 traps)
- Residential areas (6 traps)

Land cover categories

- Low vegetation (< 3 m)
- High vegetation (> 3 m)
- Buildings
- Roads
- Others

A. Variable importance plot

Average patch size of low vegetation
in a 50m radius buffer around traps (at 50m)

Maximum hourly temperature (tmax)
during sampling

The total length of roads edge
in a 250m radius buffer around traps (at 250m)

Percentage of cover of low vegetation
in a 250m radius buffer around traps (at 250m)

Weekly cumulated rainfaill (CUMRF)
during the 6th week bef. sampling

Weekly Growing Degree Day (GDD)
b/w the sampling and two weeks bef.

Mean daily Wind Speed (WS)
b/w the sampling and five weeks bef.

Average patch size of roads
in a 250m radius buffer around traps (at 250m)

Traps location

Day and session of sampling

Maximum hourly relative humidity (RHMAX)
during the 24 h before

Variable topics

- Real−time and 48h−lagged micro−climatic
- Weeks−lagged meteorological
- Sampling variables
- Land cover − roads and building
- Land cover − vegetation

B. Partial Dependence Plots